

8. REFERENCES

- *ACGIH. 1989. TLVs Threshold limit values and biological exposure indices for 1989-1990. American Conference of Governmental Industrial Hygienists. Cincinnati, OH.
- *Alderson M, Rattan N. 1980. Mortality of workers on the isopropyl alcohol plant and two MEK dewaxing plants. *Br J Ind Med* 37:85-89.
- *Allen N, Mendell J, Billmaier D, et al. 1975. Toxic polyneuropathy due to methyl-n-butyl-ketone. *Arch Neurol* 32:209.
- Almaguer D, Kramkowski R, Orris P. 2984. Health hazard evaluation report no. HETA-83-296-1491. Report to Hazard Evaluations and Technical Assistance Branch, NIOSH, U.S. Department of Health and Human Services by Johnson Controls, Incorporated, Watertown, WI.
- *Altenkirch H, Mager J, Stoltenburg G, et al. 1977. Toxic polyneuropathies after sniffing a glue thinner. *J Neurology* 214:137-152.
- *Altenkirch H, Stoltenburg G, Wagner H. 1978a. Experimental studies on hydrocarbon neuropathies induced by methyl-ethyl-ketone. *J Neurol* 219:159-170.
- *Altenkirch H, Stoltenburg-Didinger G, Wagner H. 1978b. Experimental data on the neurotoxicity of methyl-ethyl-ketone. *Experientia* 35:503-504.
- *Altenkirch H, Wagner H, Stoltenburg G, et al. 1982a. Nervous system responses of rats to subchronic inhalation of N-hexane and N-hexane plus methylethyl-ketone mixtures. *J Neurol Sci* 57:209-220.
- *Altenkirch H, Wagner H, Stoltenburg-Didinger G, et al. 1982b. Potentiation of hexacarbon-neurotoxicity by methyl-ethyl-ketone (MEK) and other substances: Clinical and experimental aspects. *Neurobehav Toxicol Teratol* 4:623-627.
- *Amoore J, Hautala E. 1983. Odor as an aid to chemical safety: Odor thresholds compared with threshold limit values and volatilities for 214 industrial chemicals in air and water dilution. *J Appl Toxicol* 3:272-290.
- *Anbar M, Neta P. 1967. A compilation of specific bimolecular rate constant for the reactions of hydrated electrons, hydrogen atoms and hydroxyl radical with inorganic and organic compounds in aqueous solution. *Int J Appl Radiation Isotopes* 18:593-523.
- *Anderson C, Sundberg K, Groth O. 1986. Animal model for assessment of skin irritancy. *Contact Dermatitis* 15:143-151.

*Cited in text

8. REFERENCES

- Apol A. 1982. Health hazard evaluation report no. HETA-81-105-831. Report to National Institute for Occupational Safety and Health, Cincinnati, OH. Division of Surveillance, Hazard Evaluations and Field Studies by Labels West, Inc., Resmond, WA.
- *Atkinson R. 1985. Kinetics and mechanisms of the gas-phase reactions of hydroxyl radical with organic compounds under atmospheric conditions. Chem Rev 85:69-201.
- Atkinson R, Darnall KR, Lloyd AC, et al. 1979. Kinetics and mechanisms of the reactions of the hydroxyl radical with organic compounds in the gas phase. Adv Photochem 11:375-488.
- Atkinson R, Aschmann SM, Carter WPL, et al. 1982. Rate constants for the gas-phase reaction of OH radicals with a series of ketones at 299 K. Int J Chem Kinet 14:839-847.
- *Babeu L, Vaishnav DD. 1987. Prediction of biodegradability for selected organic chemicals. J Ind Microb 2:107-115.
- Bang K. 1984. Health effects of common organic solvents in the workplace. Health Hazards Occup Environ 7:15-29.
- *Barnes D, Bellen J, DeRosa C, et al. 1988. Reference dose (RfD): description and use in health risk assessments. Volume I, Appendix A: Integrated risk information system supportive documentation. Washington, DC; US Environmental Protection Agency, Office of Health and Environmental Assessment. EPA/600/8-860-32a.
- *Basler A. 1986. Aneuploidy-inducing chemicals in yeast evaluated by the micronucleus test. Mutat Res 174:11-13.
- Battista JR, Connelly JP. 1989. VOC contamination at selected Wisconsin landfills--sampling results and policy implications. Madison, WI: Wisconsin Department of Natural Resources. PUBL-SW-094-89.
- *Berg E. 1971. Retrobulbar Neuritis. Ann Ophthalmol 3:1351-1353.
- Bernard A, De Russis R, Normand J, et al. 1989. Evaluation of the subacute nephrotoxicity of cyclohexane and other industrial solvents in the female Sprague-Dawley rat. Toxicol Lett 45:271-280.
- *Bertsch W, Anderson E, Holzer G. 1975. Trace analysis of organic volatiles in water by gas chromatography-mass spectrometry with glass capillary columns. J Chromatog 112:701-718.

8. REFERENCES

- *Billmaier D, Yee H, Allen N, et al. 1974. Peripheral neuropathy in a coated fabrics plant. *J Occup Med* 16:665-671.
- Binaschi S, Gazzaniga G, Crovato E. 1976. Behavioral toxicology in the evaluation of the effects of solvent mixtures. *Adverse Eff Environ Chem Psych Drug* 2:91-98.
- Botta D, Castellani PL, Mantica E. 1984. Ground water pollution solvents and their microbial degradation products. *Comm Eur Comm* 8518:261-275.
- *Brady JF, Li D, Ishizaki H, et al. 1989. Induction of cytochromes P450IIE1 and P450IIB1 by secondary ketones and the role of P450IIE1 in chloroform metabolism. *Toxicol Appl Pharmacol* 100:342-349.
- *Bridie AL, Wolff CJM, Winter M. 1979. BOD and COD of some petrochemicals. *Water Res* 13:627-630. . .
- *Bronstein AC, Currance PL. 1988. Emergency care for hazardous materials exposure. Washington, DC: The CV Mosby Company, 93-94.
- *Brown E, Hewitt W. 1984. Dose-response relationships in ketone-induced potentiation of chloroform hepato- and nephrotoxicity. *Toxicol Appl Pharmacol* 76:437-453.
- *Brown W, Setzer J, Dick R, et al. 1986. Body burden profiles of single and mixed exposure to solvents. In: Third joint U.S.-Finnish Science Symposium. Frankfurt, KY: The changing nature of work and workforce, 111-114.
- *Brown W, Setzer J, Dick R, et al. 1987. Body burden profiles of single and mixed solvent exposures. *J Occup Med* 29:877-883.
- *Browning E. 1965. Ketones. In: Toxicity and metabolism of industrial solvents. New York, NY: Elsevier Publishing Co, 412-462.
- *Brugnone F, Perbellini L, Apostoli P, et al. 1983. Environmental and biological monitoring of occupational methylethyl ketone exposure. *Dev Sci Pratt Toxicol* 571-574.
- Buttery RG, Ling LC, Guadagni DG. 1969. Food volatiles: Volatiles of aldehydes, ketones, and esters in dilute water solution. *J Agric Food Chem* 17:385-389.
- *Cammaerts MC, Inwood MR, Morgan ED, et al. 1978. Comparative study of the pheromones emitted by workers of the ants *Myrmica rubra* and *Mvrmica scabrinodis*. *J Insect Physiol* 24: 207-214.

8. REFERENCES

- *Carpenter C, Smyth H, Pozzani U. 1949. The assay of acute vapor toxicity, and the grading and interpretation of results on 96 chemical compounds. *J Ind Hyg Toxicol* 31:343-346.
- *CAS . 1989. Chemical Abstracts Service, Chemical Abstracts Registry File [Database]. September 19, 1989.
- *Cavender FL, Casey HW. 1981. 90-Day vapor inhalation toxicity study of methyl ethyl ketone in albino rats. *Toxigenics, Inc., Decatur, IL.*
- *Cavender F, Casey H, Salem H, et al. 1983. A go-day vapor inhalation toxicity study of methyl ethyl ketone. *Fund Appl Toxicol* 3:264-270.
- *Chemical Marketing Reporter. 1987. Chemical profile: Methyl ethyl ketone. August 24, 1987.
- *Chemline. 1989. National Library of Medicine Chemline Database. September 19, 1989.
- *Chou WL, Speece RE, Siddiqi RH. 1979. Acclimation and degradation of petrochemical wastewater components by methane fermentation. *Biotechnol Bioeng symp* 8:391-414.
- *CLPSD. 1988. Contract Laboratory Program Statistical Database. August 16, 1988.
- *Coleman WE, Lingg RD, Melton RG, et al. 1976. The occurrence of volatile organics in five drinking water supplies using gas chromatography/ mass spectrometry. In: Keith L, ed. *Analysis and identification of organic substances in water*. Chapter 21. Ann Arbor, MI: Ann Arbor Science, 305-327.
- *Conkle JP, Camp BJ, Welch BE. 1975. Trace composition of human respiratory gas. *Arch Environ Health* 30:290-295.
- *Corwin JF. 1969. Volatile oxygen-containing organic compounds in sea water: Determination. *Bull Mar Sci* 19:504-559.
- Couri D, Hetland L, Abdel-Rahman M, et al. 1977. The influence of inhaled ketone solvent vapors on hepatic microsomal biotransformation activities. *Toxicol Appl Pharmacol* 41:285-289.
- *Couri D, Abdel-Rahman M, Hetland L. 1978. Biotransformation of n-hexane and methyl n-butyl ketone in guinea pigs and mice. *Am Ind Hyg Assoc J* 39:295-300.
- *Cox RA, Derwent RG, Williams MR. 1980. Atmospheric photooxidation reactions. Rates, reactivity, and mechanism for reaction of organic compounds with hydroxyl radicals. *Environ Sci Technol* 14:57-61.

8. REFERENCES

- *Cox RA, Patrick KF, Chant SA. 1981. Mechanism of atmospheric photooxidation of organic compounds. Reactions of alkoxy radicals in oxidation of N-butane and simple ketones. *Environ Sci Technol* 15:587-592.
- *Cunningham J, Sharkawi M, Plaa GL. 1989. Pharmacological and metabolic interactions between ethanol and methyl D-butyl ketone, methyl isobutyl ketone, methyl ethyl ketone, or acetone in mice. *Fundam Appl Pharmacol* 13:102-109.
- *Darnall KR, Lloyd AC, Winer AM, et al. 1976. Reactivity scales for atmospheric hydrocarbons based on reaction with hydroxyl radicals. *Environ Sci Technol* 10:692-696.
- *Davis T, Baker R. 1975. Eye irritation tests with four samples of methyl ethyl ketone in albino rabbits. Report to Celanese Corporation, New York, NY by Industrial Bio-Test Laboratories Inc., Northbrook, IL. OTS Submission 206023. [Peer reviewed]
- *Deacon M, Pilny M, John J, et al. 1981. Embryotoxicity and fetotoxicity of inhaled methylethyl ketone in rats. *Toxicol Appl Pharmacol* 59:620-622.
- Dearden JC. 1985. Partitioning and lipophilicity in quantitative structureactivity. *Relationships Environ Health Perspec* ;61:203-228.
- *De Ceaurriz J, Desiles J, Bonnet P, et al. 1983. Concentration-dependent behavioral changes in mice following short-term inhalation exposure to various industrial solvents. *Toxicol Appl Pharmacol* 67:383-389.
- *Decker DW, Clark CS, Elia VJ, et al. 1983. Worker exposure to organic vapors at a liquid chemical waste incinerator. *Am Ind Hyg Assoc J* 44:296-300.
- *Delfino JJ, Miles CJ. 1985. Aerobic and anaerobic degradation of organic contaminants in Florida groundwater. *Soil Crop Sci Sot FL Proc* 44:9-14.
- *Del Rosario R, De Lumen BO, 'Habu T, et al. 1984. Comparison of headspace volatiles from winged beans and soybeans. *J Agric Food Chem* 32:1011-1015.
- *Deveaux M, Huvenne JP. 1987. Identification of solvents of abuse using gas chromatography/Fourier transform infrared spectrometry after headspace sampling. *Chromatographia* 23:626-630.
- *Dick R, Setzer J, Wait R, et al. 1984. Effects of acute exposure of toluene and methyl ethyl ketone on psychomotor performance. *Int Arch Occup Environ Health* 54:91-109.
- *Dick R, Brown W, Setzer J, et al. 1988. Effects of short duration exposures to acetone and methyl ethyl ketone. *Toxicol Lett* 43:31-49.

8. REFERENCES

*Dick R, Setzer J, Taylor B, et al. 1989. Neurobehavioral effects of short duration exposures to acetone and methyl ethyl ketone. *Br J Ind Med* 46:111-121.

Dietz F. 1979. The role of 2-butanol and 2-butanone metabolism in the potentiation of carbon-tetrachloride induced hepatotoxicity. *Dissertation Abstracts International* 41/01-B.

*Dietz F, Traiger G. 1979. Potentiation of CCl₄ of hepatotoxicity in rats by a metabolite of 2-butanone: 2,3-butanediol. *Toxicology* 14:209-215.

*Dietz F, Rodriguez-Giayola M, Traiger G, et al. 1981. Pharmacokinetics of 2-butanol and its metabolites in the rat. *J Pharmacokinet Biopharm* 9:533-576.

*Dilling WL, Bredeweg CJ, Tefertiller NB. 1976. Organic photochemistry. XIII. Simulated atmospheric photodecomposition rates of methylene chloride, 1,1,1-trichloroethane, trichloroethylene, tetrachloroethylene, and other compounds. *Environ Sci Technol* 10:351-356.

*DiVincenzo G, Kaplan C, Dedinas J. 1976. Characterization of the metabolites of methyl n-butyl ketone, methyl iso-butyl ketone, and methyl ethyl ketone in guinea pig serum and their clearance. *Toxicol Appl Pharm* 36:511-522.

*Doty R, Deems D, Frye R, et al. 1988. Olfactory sensitivity, nasal resistance, and autonomic function in patients with multiple chemical sensitivities. *Arch Otolaryngol Head Neck Surg* 114:1422-1427.

Douglas R, Coe J. 1987. The relative sensitivity of the human eye and lung to irritant gases. *Annals Occup Hyg* 31:2265-267.

*Dumont JP, Adda J. 1978. Occurrence of sesquiterpenes in mountain cheese volatiles. *J Agric Food Chem* 26:364-367.

*Dunovant VS, Clark CS, Que Hee SS, et al. 1986. Volatile organics in the wastewater and airspaces of three wastewater plants. *J Water Pollut Contr Fed* 58:886-895.

*Dyksen JE, Hess AF III. 1982. Alternatives for controlling organics in ground water supplies. *J Am Water Works Assoc* 74:394-403.

Eastman Kodak. 1976a. Electromyographic examination of dogs treated with ketone solvents. Eastman Kodak Company, Rochester, NY. OTS Submission U 206514. [Peer reviewed]

8. REFERENCES

- Eastman Kodak. 1976b. Acute intraperitoneal toxicity of methyl ethyl ketone, methyl n-butyl ketone and methyl iso-butyl ketone to rats and guinea pigs.
- Eastman Kodak. 1977. Chronic intraperitoneal administration of MnBK, MEK, MiBK and combinations of these ketones to rats. Eastman Kodak Company, Rochester, NY. OTS Submission 0206514. [Peer reviewed]
- *Eastman Kodak. 1978. Chronic skin application of MnBK, MEK, MiBK, MEK/MiBK and MEK/MnBK. Eastman Kodak Company, Rochester, NY. OTS Submission 0206514. [Peer reviewed]
- *Edney EO, Corse EW. 1986. Validation of OH radical reaction rate constant test protocol. Washington, DC: U.S. Environmental Protection Agency. NTIS PB86-166758/AS.
- *Edney EO, Kleindienst TE, Corse EW. 1986. Room temperature rate constants for the reaction of OH with selected chlorinated and oxygenated hydrocarbons. Int J Chem Kinet 18:1355-1371.
- *Eisenreich SJ, Looney BB, Thornton JD. 1981. Airborne organic contaminants in the Great Lakes ecosystem. Env Sci Tech 15:30-38.
- *EPA. 1974. New Orleans area water supply study: Draft analytical report by the Lower Mississippi River Facility, Slidell, LA. Dallas, TX: U.S. Environmental Protection Agency.
- *EPA. 1975. Preliminary assessment of suspected carcinogens in drinking water: Interim report to Congress, June, 1975. Washington, DC: U.S. Environmental Protection Agency.
- EPA. 1984. Health effects assessment for methyl ethyl ketone. Cincinnati OH: U.S. Environmental Protection Agency, Environmental Criteria and Assessment Office. ECAO-CIN-H003.
- *EPA. 1985a. Notification requirements; reportable quantity adjustments; final rule and proposed rule. Federal Register 50:13479.
- EPA. 1985b. Health and environmental effects profile for methyl ethyl ketone. Cincinnati OH: U.S. Environmental Protection Agency, Office of Health Effect Assessment, Environmental Criteria and Assessment Office. ECAO-CW-P143.
- *EPA. 1988. Contract Laboratory Program statement of work for organic analysis. Washington, DC: U.S. Environmental Protection Agency, D-1 to D-38.
- *EPA. 1989a. Interim methods for development of inhalation reference doses. U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. Washington, DC. EPA 600/8-88-066F.

8. REFERENCES

- *EPA. 1989b. Coordinated list of chemicals: 2-Butanone, CAS i/78933. Washington, DC: U.S. Environmental Protection Agency.
- *EPA. 1989c. Updated health effects assessment for methyl ethyl ketone. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Health Effects Assessment, Environmental Criteria and Assessment Office. ECAO-CIN-H003a.
- *Fisher GS, Legendre MG, Lovgren NV, et al. 1979. Volatile constituents of southernpea seed [*Vigna unguiculata* (L) Walp.]. *J Agric Food Chem* 27:7-11.
- *Francis AJ, Iden CR, Nine BJ, et al. 1980. Characterization of organics in leachates from low-level radioactive waste disposal sites. *Nuclear Tech* 50:158-163.
- *FSTRAC. 1988. Summary of state and federal drinking water standards and guidelines. Prepared by Chemical Communication Subcommittee Federal-State Toxicology and Regulatory Alliance Committee (FSTRAC).
- *Garman JR, Freund T, Lawless EW. 1987. Testing for groundwater contamination at hazardous waste sites. *J Chromatogr Sci* 25:328-337.
- *Gaudy AF Jr, Turner BG, Pusztaszeri S. 1963. Biological treatment of volatile waste components. *J Water Pollut Contr Fed* 35:75-93.
- *Geller I, Gause E, Kaplan H, et al. 1979. Effects of acetone, methyl ethyl ketone and methyl isobutyl ketone on a match-to-sample task in the baboon. *Pharmacol Biochem Behav* 11:401-406.
- Gerhold RM, Malaney GW. 1966. Structural determinants in the oxidation of aliphatic compounds by activated sludge. *J Water Pollut Contr Fed* 38:562-579.
- *Ghittori S, Imbriani M, Pezzagno G, et al. 1987. The urinary concentration of solvents as a biological indicator of exposure: Proposal for the biological equivalent exposure limit for nine solvents. *Am Ind Hyg Assoc J* 48:786-790.
- Goldstein M. 1981. Effect of methyl-n-butyl ketone and other ketone homologues on mammalian cells in culture. *Govt Reports Announcements* 6: Index 23.
- *Gordon DT, Morgan ME. 1979. Principal volatile compounds in feed flavored milk. *J Dairy Sci* 55:905-912.
- Gould JP, Ramsey RE, Giabbai M, et al. 1983. Formation of volatile haloorganic compounds in the chlorination of municipal landfill leachates. In: *Water chlorination environmental impact and health effects* 4:525-539.

8. REFERENCES

- *Grey TC, Shrimpton DH, 1967. Volatile components of raw breast muscle. *Br Poultry Sci* 8:23-33.
- *Grosjean D. 1982. Formaldehyde and other carbonyls in Los Angeles ambient air. *Environ Sci Technol* 16:254-262.
- *Grosjean D, Wright B. 1983. Carbonyls in urban fog, ice fog, cloudwater and rainwater. *Atmos Environ* 17:2093-2096.
- *Gusten H, Klasinc L, Marie D. 1984. Prediction of the abiotic degradability of organic compounds in the troposphere. *J Atmos Chem* 2:83-94.
- *Halvorsen F, Ohneck R. 1985. MEK groundwater decontamination. In: *Hazardous wastes and environmental emergencies*. Silver Springs, MD: Hazardous Materials Control Research Institute, 193-195.
- *Hampton CV, Pierson WR, Harvey TM. 1982. Hydrocarbon gases emitted from vehicles on the road. I. A Qualitative gas chromatography/mass spectrometry survey. *Environ Sci Technol* 16:287-298.
- *Hansch C, Leo AJ. 1950. *Medchem Project*. Issue No. 26. Claremont, CA: Pomona College.
- Harada K, Nagano M, Ohmori S, et al. 1989. Combined effects of methyl B-butyl ketone and methyl ethyl ketone on activities of microsomal drug metabolizing enzymes in rats. *Jpn J Ind Health* 31:156-157.
- Harkov R, Katz R, Bozzelli J. 1981. Toxic and carcinogenic air pollutants in New Jersey-volatile organic substances. In: McGovern J, ed. *International technical conference: Toxic air contamination*. Pittsburgh, PA: Air Pollution Control Association, 104-119.
- *Haskell Laboratory. 1971, Federal hazardous substances act tests - rabbit eye irritation. Report by Haskell Laboratory to E. I. duPont de Nemours and Co., Inc., Wilmington, DE. OTS Submission 215038. (Peer reviewed)
- Hawkes C, Cavanagh J, Fox A. 1989. Motoneuron disease: A disorder secondary to solvent exposure? *Lancet* 1:73-76.
- *Hawthorne SB, Sievers RE. 1984. Emission of organic air pollutants from shale oil wastewaters. *Environ Sci Technol* 18:483-490.
- Hawthorne SE, Sievers RE, Barkley RM. 1985. Organic emissions from shale oil wastewaters and their implications for air quality. *Environ Sci Tech* 19:922-927.

8. REFERENCES

- *Hazleton Laboratories. 1963a. Primary skin irritation - rabbits. Report to Esso Research and Engineering Company by Hazleton Laboratories, Inc., Falls Church, VA. OTS Submission 206264. [Peer reviewed]
- *Hazleton Laboratories. 1963b. Acute eye irritation - rabbits. Report to Esso Research and Engineering Company by Hazleton Laboratories, Inc., Falls Church, VA. OTS Submission 206264. [Peer reviewed]
- Hewitt W, Brown E. 1984. Nephrotoxic interactions between ketonic solvents and halogenated aliphatic chemicals. *Fundam Appl Toxicol* 4:902-908.
- Hewitt W, Miyajima H, Cote M, et al. 1980. Modification of haloalkane-induced hepatotoxicity by exogenous ketones and metabolic ketosis. *Fed Proc* 39:3118-3123.
- *Hewitt W, Brown E, Plaa G. 1983. Relationship between the carbon skeleton length of ketonic solvents and potentiation of chloroform-induced hepatotoxicity in rats. *Toxicol Lett* 16:297-304.
- *Hewitt L, Ayotte P, Plaa G. 1986. Modifications in rat hepatobiliary function following treatment with acetone, 2-butanone, 2-hexanone, mirex or chlordecone and subsequently exposed to chloroform. *Toxicol Appl Pharmacol* 83:465-473.
- *Hewitt L, Valiquetta C, Plaa G. 1987. The role of biotransformation/detoxication in acetone-, 2-butanone-, and 2-hexanone-potentiated chloroform-induced hepatotoxicity. *Can J Physiol Pharmacol* 65:2313-2318.
- Hewitt LA, Palmason C, Masson S, et al. 1990. Evidence for the involvement of organelles in the mechanism of ketone-potentiated chloroform-induced hepatotoxicity. *Liver* 10:35-48.
- *Higgins CE, Griest WH, Olerich, G. 1983. Application of Tenax trapping to analysis of gas phase organic compounds in ultra-low tar cigarette smoke. *J Assoc Off Anal Chem* 66:1074-1083.
- Hites RA, Lopez-Avila V. 1980. Sedimentary accumulations of industrial organic compounds discharged into a river system. In: Baker RA, ed. *Contaminants and sediments. Vol. 1: Fate and transport case studies, modeling, toxicity.* Ann Arbor, MI: Ann Arbor Science.
- *Hodgkin JH, Galbraith MN, Chong YK. 1982. Combustion products from burning polyethylene. *J Macromol Sci Chem* 17:35-43.
- *HSDB. 1989. Hazardous Substances Data Bank. National Library of Medicine, National Toxicology Information Program, Bethesda, MD. September 11, 1989.

8. REFERENCES

- Ingram A. 1979. Interaction of benzo(a)pyrene and a hyperplastic agent in epidermal nuclear enlargement in the mouse. A dose-response study. *Chem Biol Interact* 26:103-113.
- Ingram A, Grass P. 1985. Nuclear enlargement-- an early change produced in mouse epidermis by carcinogenic chemicals applied topically in the presence of a promoter. *J Appl Toxicol* 5:53-60.
- IRIS. 1989. Integrated Risk Information System. U.S. Environmental Protection Agency, Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH. September 1, 1989.
- *Isidorov VA, Zenkevich IG, Ioffe BV. 1985. Volatile organic compounds in the atmosphere of forests. *Atmos Environ* 19:1-8.
- Iwata M, Takeuchi Y, Hisanaga N, et al. 1983. Changes of n-hexane metabolites in urine of rats exposed to various concentrations of n-hexane and to its mixture with toluene or MEK. *Int Arch Occup Environ Health* 53:1-8.
- Iwata M, Takeuchi Y, Hisanaga N, et al. 1984. Changes of n-hexane neurotoxicity and its urinary metabolites by long-term co-exposure with MEK or toluene. *Int Arch Occup Environ Health* 54:273-281.
- *Jarke FH, Dravnieks A, Gordon SM. 1981. Organic contaminants in indoor air and their relation to outdoor contaminants. *ASHRAE Trans* 87:153-166.
- John J, Wroblewski D, Schwetz B. 1984. Teratogenicity of experimental and occupational exposure to industrial chemicals. *Issues Rev Teratol* 8:267-324.
- *Jonsson A, Persson KA, Grigoriadis V. 1985. Measurements of some low molecular-weight oxygenated, aromatic and chlorinated hydrocarbons in ambient air and in vehicle emissions. *Environ Int* 11:383-392.
- *Jungclaus GA, Lopez-Avila V, Hites RA. 1978. Organic compounds in an industrial waste water: A case study of their environmental impact. *Environ Sci Technol* 12:88-96.
- *Katz G. 1985. Chemical and biological interactions affecting neurotoxicity. In: O'Donoghue, ed. *Neurotoxicity of industrial and commercial chemicals*, Vol. I. Boca Raton, FL: CRC Press, Inc., 149-158.
- *Keen AR, Walker NJ, Perberdy MF. 1974. The formation of 2-butanone and Z-butanol in cheddar cheese. *J Dairy Res* 41:249-257.
- *Kennah H II, Hignet S, Laux P, et al. 1989. An objective procedure for quantitating eye irritation based upon changes of corneal thickness. *Fund Appl Toxicol* 12:258-268.

8. REFERENCES

- *Kezic S, Monster AC. 1988. Determination of methyl ethyl ketone and its metabolites in urine using capillary gas chromatography. *J Chromatogr* 428:175-280.
- *Kimura E, Ebert D, Dodge P. 1971. Acute toxicity and limits of solvent residue for sixteen organic, solvents. *Toxicol Appl Pharmacol* 19:699-704.
- *King P, Morris J, Pollard J. 1985. Glue sniffing neuropathy. *Aust N Z J Med* 15:293-299.
- *Kinlin TE, Muralidhara R, Pittet AO. 1972. Volatile components in roasted filberts. *J Agric Food Chem* 20:1021-1028.
- *Klimisch H. 1988. The inhalation hazard test; principle and method. *Arch Toxicol* 61:411-416.
- *Kolb DK. 1988. Teratogenic chemicals in undergraduate general chemistry laboratories. *Stud Environ Sci* 31:2;47-255.
- *Kollig HP, Ellington JJ, Hamrick KJ, et al. 1987. Hydrolysis rate constants, partition coefficients, and their water solubilities for 129 chemicals. A summary of fate constants provided for the concentration-based listing program, 1987. Prepublication. Athens, GA: U.S. Environmental Research Laboratory, Office of Research and Development, Environmental Protection Agency.
- *Kool HJ, van Kreijl CF, Zoeteman BCJ. 1982. Toxicology assessment of organic compounds in drinking water. *Crit Rev Env Contr* 12:307-357.
- *Kopelman P, Kalfayan P. 1983. Severe metabolic acidosis after ingestion of butanone. *Br Med J [Clin Res]* 286:21-22.
- *Kopfler FC, Melton RG, Mullaney JL, et al. 1977. Human exposure to water pollutants. *Adv Environ Sci Technol* 8:4;19-433.
- Krasavage WJ, O' Donoghue JL, DiVincenzo GD. 1982. Ketones. In: Clayton GD, Clayton FE, ed. *Patty's industrial hygiene and toxicology*. Vol. 2C, 3rd ed. New York, NY: John Wiley and Sons, 4728-4733.
- Krill RM, Sonzogni WC. 1986. Chemical monitoring of Wisconsin's groundwater. *J Am Water Works Assoc* 78:70-75.
- *Krotoszynski NK, Bruneau GM, O'Neill HJ. 1979. Measurement of chemical inhalation exposure in urban population in the presence of endogenous effluents. *J Anal Toxicol* 3:225-234.

8. REFERENCES

- *Kupferschmid LL, Perkins JL. 1986. Organic solvent recycling plant exposure levels. *Appl Ind Hyg* 1:122-124.
- *LaBelle C, Brieger H. 1955. The vapor toxicity of a composite solvent and its principal components. *Arch Ind Health* 12:623-627.
- *LaRegina J, Bozzelli JW, Harkov R, et al. 1986. Volatility organic compounds at hazardous waste sites and a sanitary landfill in New Jersey. An up-to-date review of the present situation. *Environ Prog* 5:18-27.
- Lauwerys R. 1983. Industrial chemical exposure: guidelines for biological monitoring. Davis, CA: Biomedical Publications.
- *Lee S, Frederick L. 1981. Health hazard evaluation report no. HHE80-103-827. Report to National Institute for Occupational Safety and Health, Cincinnati, OH. Division of Surveillance, Hazard Evaluations and Field Studies by Joel and Aronoff, Ridgefield, New Jersey.
- *Lee S, Murphy D. 1982. Health hazard evaluation report no. HETA-82-030-1184. Report to Hazard Evaluations and Technical Assistance Branch, NIOSH, Cincinnati, Ohio by Medalist-Gladiator Athletic Products Company, Leesburg, Florida.
- *Lee S, Parkinson D. 1982. Health hazard evaluation report no. HETA 80-168-1204. Report to National Inst for Occupational Safety and Health, Cincinnati, OH by Rola-Esmark Company, Du Bois, Pennsylvania.
- *Li G, Yin S, Watanabe T, et al. 1986. Benzene-specific increase in leukocyte alkaline phosphatase activity in rats exposed to vapors of various organic solvents. *J Toxicol Environ Health* 19:581-589.
- *Liebich HM, Bertsch W, Zlatkis A, et al. 1975. Volatile organic components in the Skylab 4 spacecraft atmosphere. *Aviat Space Environ Med* 46:1002-1007.
- *Liira J, Riihimaki V, Pfaffli P. 1988a. Kinetics of methyl ethyl ketone in man: absorption, distribution and elimination in inhalation exposure. *Int Arch Occup Environ Health* 60:195-200.
- *Liira J, Riihimaki V, Engstrom K, et al. 1988b. Coexposure of man to m-xylene and methyl ethyl ketone: kinetics and metabolism. *Stand J Work, Environ Health* 14:322-327.
- *Liira J, Riihimaki V, Engstrom K. 1990. Effects of ethanol on the kinetics of methyl ethyl ketone in man. *British Journal of Industrial Medicine* 47:325-330.

8. REFERENCES

- *Liira J, Elovaara E, Raunio H, et al. 1991. Metabolic interaction and disposition of methyl ethyl ketone and m-xylene in rats at single and repeated inhalation exposures. *Xenobiotica* 21:53-63.
- Lissi EA, Abuin E, Envina MV. 1973. Photochemistry of butanone and methyl butanone. *J Photochem* 2:377-392.
- *Lovegren NV, Fisher GS, Legendre MG, et al. 1979. Volatile constituents of dried legumes. *J Agric Food Chem* 27:851-853.
- *Lowry L. 1987. The biological exposure index: Its use in assessing chemical exposures in the workplace. *Toxicology* 47:55-69.
- Lundberg I, Ekdahl M, Kronevi T, et al. 1986. Relative hepatotoxicity of some industrial solvents after intraperitoneal injection or inhalation exposure in rats. *Environ Res* 40:411-420.
- *Lyman WJ, Reehl WF, Rosenblatt DH. 1982. Handbook of chemical property estimation methods. New York, NY: McGraw Hill Book Co. Chapters 4, 7, 15.
- Maizlish N, Langolf G, Whitehead L, et al. 1985. Behavioral evaluation of workers exposed to mixtures of organic solvents. *Br J Ind Med* 42:579-590.
- *MAORS. 1989. Massachusetts drinking water standards and guidelines. Boston, MA: Office of Research and Standards.
- *Mast T, Dill J, Evanoff J, et al. 1989. Inhalation developmental toxicology studies: teratology study of methyl ethyl ketone in mice. Report by Pacific Northwest Laboratory, Richland, WA to National Institute of Environmental Health Science, National Toxicology Program. PNL-6833 DE89 009563. [Peer reviewed]
- *Mayer VW, Goin CJ. 1987. Effects of chemical combinations on the induction of aneuploidy in *Saccharomyces cerevisiae*. *Mutat Res* 187:21-30.
- Medinilla J, Espigares M. 1988. Contamination by organic solvents in auto paint shops. *Ann Occup Hyg* 32:509-513.
- Meyera V. 1983. Chemicals which cause birth defects - teratogens: A special concern of research chemists. *Sci Total Environ* 32:1-12.
- *Mill T. 1982. Hydrolysis and oxidation processes in the environment. *Environ Toxicol Chem.* 1:135-141.
- *Mill T, Hendry DG, Richardson H. 1980. Free-radical oxidants in natural waters. *Science* 207:886-887.

8. REFERENCES

- *Misumi J, Nagano M. 1985. Experimental study on the enhancement of the neurotoxicity of methyl n-butylketone by non-neurotoxic aliphatic monoketones. *Br J Ind Med* 42:155-161.
- Miyamoto Y. 1986. Eye and respiratory irritants in jet engine exhaust. *Aviation Space and Environ Med* 57:1104-1108.
- *Miyasaka M, Kumai M, Koizumi A, et al. 1982 . Biological monitoring of occupational exposure to methyl ethyl ketone by means of urinalysis for methyl ethyl ketone itself. *Int Arch Occup Environ Health* 50:131-137.
- Moody P, Hartle R. 1982. Health hazard evaluation report no. HETA-81-174-962. Report to National Inst for Occupational Safety and Health, Cincinnati, OH. Div of Surveillance, Hazard Evaluations and Field Studies by Sport Craft, Inc., Perry, Florida.
- Morimoto Y, Seki T, Sugibayashi K, et al. 1988. Basic studies on controlled transdermal delivery of nifedipine hydrochloride using ethylene-vinyl acetate and ethylene-vinyl alcohol copolymer membranes. *Chem Pharm Bull* 36:2633-2641.
- *NAS/NRC. 1989. Biologic markers in reproductive toxicology. National Academy of Sciences/National Research Council. Washington, DC: National Academy Press, 15-35.
- *NATICH. 1988. National Toxics Information Clearinghouse. Database report of state, local and EPA air toxics activities. Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency, Research Triangle Park, NC.
- *Neier W, Strehlke G. 1985. 2-Butanone. In: Ullman's encyclopedia of industrial chemistry. Deerfield Beach, FL: UCH, 475-481.
- *Nelson K, Ege J, Ross M, et.al. 1943. Sensory response to certain industrial solvent vapours. *J Ind Hyg Toxicol* 25:282.
- *NIOSH. 1978. Criteria for a recommended standard...occupational exposure to ketones. Cincinnati, OH; U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health. DHHS (NIOSH) publication no. 78-173.
- *NIOSH. 1984. NIOSH manual of analytical methods. 3rd ed. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health, 2500-1 to 2500-3.
- *NIOSH. 1989. National occupational exposure survey (NOES). Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control, National Institute for Occupational Safety and Health.

8. REFERENCES

- Noma E, Berglund B, Berglund U, et al. 1988. Joint representation of physical locations and volatile organic compounds in indoor air from a healthy and a sick building. *Atmos Environ* 22:451-460.
- *O'Donoghue J, Krasavage W, DiVincenzo G, et al. 1982. Commercial-grade methyl heptyl ketone (S-methyl-2-octanone) neurotoxicity: Contribution of 5-nonanone. *Toxicol Appl Pharmacol* 62:307-316.
- *O'Donoghue J, Krasavage W, DiVincenzo G, et al. 1984. Further studies on ketone neurotoxicity and interactions. *Toxicol Appl Pharmacol* 72:201-209.
- O'Donoghue J, Haworth S, Curren R, et al. 1988. Mutagenicity studies on ketone solvents: Methyl ethyl ketone, methyl isobutyl ketone, and isophorone. *Mutat Res* 206:149-161.
- *Ogawa I, Fritz JS. 1985. Determination of low concentrations of lowmolecular-weight aldehydes and ketones in aqueous samples. *J Chromatogr* 329:81-89.
- *OHM/TADS. 1989. Oil and Hazardous Materials Technical Assistance Data System. U.S. Environmental Protection Agency, National Institute of Health, Washington, DC. September 11, 1989.
- *Osborne JS, Adamek S, Hobbs ME. 1956. Some components of gas phase of cigarette smoke. *Anal Chem* 28:211-215.
- *OSHA. 1989. Air contaminants. Final rule. U.S. Department of Labor, Occupational Safety and Health Administration. *Federal Register* 54:2926.
- *Panson R, Winek C. 1980. Aspiration toxicity of ketones. *Clin Toxicol* 17:271-317.
- *Papa AJ, Sherman PD Jr. 1981. Ketones. In: Kirk-Othmer encyclopedia of chemical technology. Vol. 13, 3rd ed. New York, NY: John Wiley and Sons, Inc., 894-941.
- *Patty F, Schrenk H, Yant W. 1935. Acute response of guinea pigs to vapours of some new commercial organic compounds. *US Treas Publ Health Rep* 50:1217.
- *Pellizzari ED. 1982. Analysis for organic vapor emissions near industrial and chemical waste disposal sites. *Environ Sci Technol* 16:781-785.
- *Pellizzari ED, Castillo NP, Willis S, et al. 1979. Identification of organic components in aqueous effluents from energy-related processes. *ASTM Spec Tech Publ STP* 686: 256-274.

8. REFERENCES

- *Pellizzari ED, Hartwell TD, Harris HSB, et al. 1982. Purgeable organic compounds in mother's milk. Bull Environ Contam Toxicol 28:322-328.
- *Perbellini L, Brugnone F, Mezzo P, et al. 1984. Methyl ethyl ketone exposure in industrial workers. Uptake and kinetics. Int Arch Occup Environ Health 54:73-81.
- *Phillips WE, Jr, Perry JJ. 1974. Metabolism of butane and 2-butanone by *Mycobacterium vaccae*. J Bacteriol 120:987-989.
- *Price KS, Waggy GT, Conway RA. 1974. Brine shrimp bioassay and seawater BOD of petrochemicals. J Water Pollut Contr Fed 46:63-77.
- *Ralston W, Hilderbrand R, Uddin D, et al. 1985. Potentiation of 2,5-hexanedione neurotoxicity by methyl ethyl ketone. Toxicol Appl Pharmacol 81:319-327.
- Rathbun, RE, Tai DY. 1982. Volatilization of ketones. Water air soil pollution 17:281-293.
- *Rathbun RE, Tai DY. 1987. Vapor pressures and gas-film coefficients for ketones. Chemosphere 16:69-78.
- *Raunio H, Liira J, Elovaara E, et al. 1990. Cytochrome P450 isozyme induction by methyl ethyl ketone and m-xylene in rat liver. Toxicol Appl Pharmacol 103:175-179.
- Remington M, Basu D. Review and validation of health effects of assessment for methyl ethyl ketone. Prepared for U.S. EPA, ECAO/Cincinnati OH under contract 68-03-3112.
- *Riddick JA, Bunger WB, Sakano TK. 1986. Organic solvents: Physical properties and methods of purification. Techniques of chemistry. 4th ed. New York, NY: Wiley-Interscience, Vol 2, 338-339.
- *Riihimaki V. 1986. Metabolism and excretion of organic solvents. In: Riihimaki V, Ulfvarson U, ed. Progress in clinical and biological research, Vol. 220. New York, NY: Alan R. Liss, Inc., 61-72.
- *Robertson P, White EL, Bus JS. 1989. Effects of methyl ethyl ketone pretreatment on hepatic mixed-function oxidase activity and on in vivo metabolism of n-hexane. Xenobiotica 19:721-729.
- *Roy WR, Griffin RA. 1985. Mobility of organic solvents in water-saturated soil materials. Environ Geol Water Sci 7:241-247.
- *Sabel GV, Clark TP. 1984. Volatile organic compounds as indicators of municipal solid waste leachate contamination. Waste Manag Res 2:119-130.

8. REFERENCES

- *Saida K, Mendell J, Weiss H. 1976. Peripheral nerve changes induced by methyl n-butyl ketone and potentiation by methyl ethyl ketone. *J Neuropath Exp Neurol* 35:207-205.
- *Sakata M, Kikuchi J, Masanobu H, et al. 1989. Disposition of acetone, methyl ethyl ketone and cyclohexanone in acute poisoning. *Clin Toxicol* 27:67-77.
- *SANSS . 1989. Structure and Nomenclature Search System. September 1, 1989.
- Sato A, Nakajima T. 1979. Partition coefficients of some aromatic hydrocarbons and ketones in water, blood and oil. *Br J Ind Med* 36:231-234.
- *Sawhney BL, Kozloski RP. 1984. Organic pollutants in leachates from landfill sites. *J Environ Qual* 13:349-352.
- *Sax NI, Lewis RJ. 1987. *Hawley's condensed chemical dictionary*. 11th ed. New York, NY: Van Nostrand Reinhold Co., 769.
- *Scheiman MA, Saunders RA, Saalfeld FE. 1974. Organic contaminants in the District of Columbia water supply. *Biomed Mass Spectrom* 4:209-211.
- *Schmidt R, Schnoy N, Altenkirch H, et al. 1984. Ultrastructural alteration of intrapulmonary nerves after exposure to organic solvents. A contribution to 'sniffers disease'. *Respiration* 46:362-369.
- *Schwetz B, Leong B, Gehring P. 1974. Embryo- and fetotoxicity of inhaled carbon tetrachloride, 1,1-dichloroethane and methyl ethyl ketone in rats. *Toxicol Appl Pharmacol* 28:452-464.
- Seixas N, Ordin D. 1985. Health hazard evaluation report no. HETA-85-268-1641. Report to Hazard Evaluations and Technical Assistance Branch, U.S. Department of Health and Human Services, Cincinnati, Ohio by Products Research Chemical Corp., Gloucester City, New Jersey.
- *Seizinger DE, Dimitriades B. 1972. Oxygenates in exhaust from simple hydrocarbon fuels. *J Air Pollut Contr Assoc* 22:47-51.
- Seki T, Sugibayashi K, Morimoto Y. 1987. Effect of solvents on the permeation of nicardipine hydrochloride through the hairless rat skin. *Chem Pharm Bull* 35:3054-3057.
- *Shah JJ, Heyerdahl EK. 1988. National ambient volatile organic compounds (VOCS) database update. EPA 600/3-88-010.
- Shah JJ, Singh HB. 1988. Distribution of volatile organic chemicals in outdoor and indoor air. *Environ Sci Technol* 22:1381-1388.

8. REFERENCES

- *Smoyer JC, Shaffer DE, Dewitt IL. 1971. A program to sample and analyze air pollution in the vicinity of a chemical reclamation plant. *Inst Environ Sci* 339-345.
- *Smyth H, Carpenter C, Weil C, et al. 1962. Range-finding toxicity data: List VI. *Am Ind Hyg Assoc J* 23:95-107.
- *Snider JR, Dawson GA. 1985. Tropospheric light alcohols, carbonyls, and acetonitrile concentrations in the southwestern United States and Henry's law data. *J Geophys Res D Atmos* 90:3797-3805.
- *Sosebee JBJR, Geiszler PC, Winegardner DL, et al. 1983. Contamination of groundwater samples with poly(vinyl chloride) adhesives and poly(vinyl chloride) primer from monitor wells. *ASTM Spec Tech Publ* 805(Hazar Ind Solid Waste Test):38-50.
- *Sosulski F, Mahmoud RM. 1979. Effects of protein supplements on carbonyl compounds and flavor in bread. *Cereal Chem* 56:533-536.
- *Speece RE. 1983. Anaerobic biotechnology for industrial wastewater treatment. *Environ Sci Technol* 17:416A-427A.
- Spencer P, Schaumburg H. 1976. Feline nervous system response to chronic intoxication with commercial grades of methyl n-butyl ketone, methyl isobutyl ketone, and methyl ethyl ketone. *Toxicol Appl Pharmacol* 37:301-311
- *SRI. 1989. Directory of chemical producers. Menlo Park, CA: Stanford Research Institute International, 785.
- *State of Kentucky. 1986. New or modified sources emitting toxic air pollutants. Department for Environmental Protection. 401 KAR 63:022.
- *Stillmeadow, Inc. 1978. Rat acute oral toxicity. Report to Shell Oil Company, Houston, TX by Stillmeadow, Inc., Houston TX. OTS Submission 206206. [Peer reviewed]
- *Stoltenburg-Didinger G, Altenkirch H, Wagner M. 1990. Neurotoxicity of organic solvent mixtures: embryotoxicity and fetotoxicity. *Neurotoxicol Teratol* 12:585-589.
- *Stonebraker RD, Smith AJ Jr. 1980. Containment and treatment of a mixed chemical discharge from "The Valley of the Drums" near Louisville, Kentucky. In: National Conference: Control of hazardous materials spills. Nashville, TN, 1-10.

8. REFERENCES

- *Stutz DR, Janusz SJ. 1988. Hazardous materials injuries: a handbook for pre-hospital care, 2nd ed. Beltsville MD: Bradford Communications Corporation, 312-313.
- *Swarm RL, Laskowski DA, McCall PJ, et al. 1983. A rapid method for the estimation of the environmental parameters octanol/water partition coefficient, soil sorption constant, water to air ratio and water solubility. Residue Rev 85:17-28.
- *Takeoka GR, Flath RA, Guntert M, et al. 1988. Nectarine volatiles: Vacuum steam distillation versus headspace sampling. J Agric Food Chem 36:553-560.
- *Takeuchi Y, Ono Y, Hisanaga N, et al. 1983. An experimental study of the combined effects of n-hexane and methyl ethyl ketone. Br J Ind Med 40:199-203.
- *Tanii H, Tsuji H, Hashimoto K. 1986. Structure-toxicity relationship of monoketones. Toxicol Lett (AMST) 30:13-18.
- *Tewari YB, Miller MM, Wasik SP, et al. 1982. Aqueous solubility and octanol/water partition coefficient of organic compounds at 25°C. J Chem Eng Data 27:451-454.
- Tham R, Bunnfors I, Eriksson B, et al. 1984. Vestibulo-ocular disturbances in rats exposed to organic solvents. Acta Pharmacol Toxicol 54:58-63.
- *Tharr D, Murphy D, Mortimer V. 1982. Health hazard evaluation report no. HETA-81-455-1229. Report to Hazard Evaluations and Technical Assistance Branch, NIOSH, Cincinnati, Ohio by Red Wing Shoe Company, Red Wing, Minnesota.
- *Thorpe E. 1982. Toxicity studies with chemical solvents: short-term in vitro tests for genotoxic activity with methyl ethyl ketone. Report to Shell Oil Company, Houston, TX by Sittington Research Centre, Sittington, Kent, UK. OTS Submission 206206. [Peer reviewed]
- *Tichenor BA. 1987. Organic emission measurements via small chamber testing. U.S. Environmental Protection Agency, Washington, DC. EPA 600/D-87-187.
- *Tichenor BA, Mason MA. 1988. Organic emissions from consumer products and building materials to the indoor environment. J Air Pollut Contr Assoc 38:264-268.
- *Toftgard R, Nilsen O, Gustafsson J-A. 1981. Changes in rat liver microsomal cytochrome P-450 and enzymatic activities after the inhalation of n-hexane, xylene, methyl ethyl ketone and methylchloroform for four weeks. Stand J Work, Environ Health 7:31-37.

8. REFERENCES

- *Tolos W, Setzer J, MacKenzie B, et al. 1987. Biological monitoring of experimental human inhalation exposures of methyl ethyl ketone and toluene. In: Ho M, Dillon H, ed., Biological monitoring of exposure to chemicals, organic compounds. New York, NY: John Wiley and Sons, 133-142.
- *Traiger GJ, Bruckner JV, Jiang W-D, et al. 1989. Effect of 2-butanol and 2-butanone on rat hepatic ultrastructure and drug metabolizing enzyme activity. *J Toxicol Environ Health* 28:235-248.
- *TRI. 1989. Toxic Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- Truong KN, Blackburn JW. 1984. The stripping of organic chemicals in biological treatment processes. *Environ Prog* 3:143-152.
- *Tsuchiya Y, 1987. Volatile organic compounds in indoor air, New Orleans, LA: American Chemical Society, Division of Environmental Chemistry, Vol. 27, 183-185.
- *Tsukamoto S, Chiba S, Muto T, et al. 1985a. Study on the metabolism of volatile hydrocarbons in mice-propane, n-butane, and iso-butane. *J Toxicol Sci* 10:323-332.
- Tsukamoto S, Chiba S, Muto T, et al. 1985b. Studies on the metabolism of volatile hydrocarbons in propane gas (LPG) inhalation. Detection of the metabolites. *Nippon Hoigaku Zasshi* 39:124-130.
- *Urano K, Kato 2. 1986. Evaluation of biodegradation ranks of priority organic compounds. *J Hazar Mater* 13:147-159.
- *USITC. 1987. United States International Trade Commission. Synthetic organic chemicals: U.S. Production and Sales, 1986. USITC Publication 2009. Washington, DC: U.S. Government Printing Office, 210, 226, 238-239.
- *USITC. 1988. United States International Trade Commission. Synthetic organic chemicals: U.S. Production and Sales, 1987. USITC Publication 2118. Washington, DC: U.S. Government Printing Office, 15-5, 15-22, 15-36, 15-37.
- *USITC. 1989. United States International Trade Commission. Synthetic organic chemicals: U.S. Production and Sales, 1988. USITC Publication 2219. Washington, DC: U.S. Government Printing Office, 15-5, 15-23, 15-36 to 15-38.
- *Vaishnav DC, Boethling RS, Babeu L. 1987. Quantitative structurebiodegradability relationships for alcohols, ketones and alicyclic compounds. *Chemosphere* 16:695-703.

8. REFERENCES

- Vaishnav DD, Lopas DM. 1985. Relationship between lipophilicity and biodegradation inhibition of selected industrial chemicals. *Dev Ind Microbiol* 26:557-565.
- *Vallat J, Leboutet M, Loubet A, et al. 1981. N-hexane-induced and methylethylketone-induced polyneuropathy: Abnormal accumulation of glycogen in unmyelinated axons. *Acta Neuropathol* 55:275-280.
- *Van Doorn JE, De Cock J, Kezic S, et al. 1989. Determination of methyl ethyl ketone in human urine after derivatization with onitrophenolhydrazine, using solid-phase extraction and reversed-phase high-performance liquid chromatography and ultraviolet detection. *J Chromatogr* 489:419-424.
- *Varigos G, Nurse D. 1986. Contact urticaria from methyl ethyl ketone. *Contact Dermatitis* 15:259-260.
- VDEC. 1990. Vermont primary groundwater quality standards. Waterbury, VT: Vermont Department of Environmental Conservation.
- Veronesi B. 1984. An ultrastructural study of methyl ethyl ketone's effect on cultured nerve tissues. *Neurotoxicology* 5:31-44.
- *Veronesi B, Lington A, Spencer P. 1984. A tissue culture model of methyl ethyl ketone's potentiation of n-hexane neurotoxicity. *Neurotoxicology* 5:43-52.
- *View Database. 1989. Agency for Toxic Substances and Disease Registry (ATSDR), Office of External Affairs, Exposure and Disease Registry Branch, Atlanta, GA. September 25, 1989.
- *Wahlberg J. 1984. Edema-inducing effects of solvents following topical administration. *Derm Beruf Umwelt* 32:91-94.
- *Wallace LA, Pellizzari E, Hartwell T, et al. 1984. Personal exposure to volatile organic compounds. 1. Direct measurements in breathing-zone air, drinking water, food, and exhaled breath. *Environ Res* 35:293-319.
- *Wallington TJ, Kurylo MJ. 1987. Flash photolysis resonance fluorescence investigation of the gas-phase reactions of hydroxyl radicals with a series of aliphatic ketones over the temperature range 240-440 K. *J Phys Chem* 91:5050-5054.
- *Wallington TJ, Dagaut P, Kurylo MJ. 1988. Correlation between gas-phase and solution-phase reactivities of hydroxyl radicals towards saturated organic compounds. *J Phys Chem* 92:5024-5028.
- *Wang TC, Bricker JL. 1979. 2-Butanone and tetrahydrofuran contamination in the water supply. *Bull Environ Contam Toxicol* 23:620-623.

8. REFERENCES

- *Weast RC, Astle MJ, Beyer WH. 1988. CRC handbook of chemistry and physics. 69th ed. Boca Raton, FL: CRC Press, Inc., C-170.
- *Wen C, Tsai S, Weiss N, et al. 1985. Long-term mortality study of oil refinery workers. IV. Exposure to the lubricating-dewaxing process. J Natl Cancer Inst 74:11-18.
- *Whelan JK, Tarafa ME, Hunt JM. 1982. Volatile C₁-C₈ organic compounds in macroalgae. Nature 299:50-52.
- *Whitehead LW, Ball GL, Fine LJ, et al. 1984. Solvent vapor exposures in both spray painting and spray glowing, and associated operations. Am Ind Hyg Assoc J 45:767-772.
- *Whittaker SG, Zimmermann FK, Dicus B, et al. 1990. Detection of induced chromosome loss in *Saccharomyces cerevisiae* - an interlaboratory assessment of 12 chemicals. Mutat Res 241:225-242.
- Yang R. 1986. The toxicology of methyl ethyl ketone. Residue Rev 97: 121-143.
- *Young RHF, Ryckman DW, Buzzell JC Jr. 1968. An improved tool for measuring biodegradability. J Water Pollut Contr Fed. 40:354-368.
- *Zimmermann FK, Scheel I, Resnick MA. 1989. Induction of chromosome loss by mixtures of organic solvents including neurotoxins. Mutat Res 224:287-303.

