

4. CHEMICAL AND PHYSICAL INFORMATION

4.1 CHEMICAL IDENTITY


Information regarding the chemical identity of carbon tetrachloride is located in Table 4-1.

4.2 PHYSICAL AND CHEMICAL PROPERTIES

Information regarding the physical and chemical properties of carbon tetrachloride is located in Table 4-2.

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-1. Chemical Identity of Carbon Tetrachloride

Characteristic	Information	Reference
Chemical name	Carbon tetrachloride	IARC 1979
Synonym(s)	Carbona; carbon chloride; carbon tet; methane tetrachloride; perchloromethane; tetrachloromethane; benzinoform	HSDB 2004
Registered trade name(s)	Benzinoform; Fasciolin; Flukoids; Freon 10; Halon 104; Tetraform; Tetrasol	IARC 1979
Chemical formula	CCl_4	IARC 1979
Chemical structure		IARC 1979
Identification numbers:		
CAS registry	56-23-5	NLM 1988
NIOSH RTECS	FG4900000	HSDB 2004
EPA hazardous waste	U211; D019	HSDB 2004
OHM/TADS	7216634	HSDB 2004
DOT/UN/NA/IMCO shipping	UN1846; IMCO 6.1	HSDB 2004
HSDB	53	HSDB 2004
NCI	No data	

CAS = Chemical Abstracts Service; DOT/UN/NA/IMCO = Department of Transportation/United Nations/North America/International Maritime Dangerous Goods Code; EPA = Environmental Protection Agency; HSDB = Hazardous Substances Data Bank; NCI = National Cancer Institute; NIOSH = National Institute for Occupational Safety and Health; OHM/TADS = Oil and Hazardous Materials/Technical Assistance Data System; RTECS = Registry of Toxic Effects of Chemical Substances

4. CHEMICAL AND PHYSICAL INFORMATION

Table 4-2. Physical and Chemical Properties of Carbon Tetrachloride

Property	Information	Reference
Molecular weight	153.82	Lide 1993
Color	Colorless	Verschueren 1983
Physical state	Liquid	Verschueren 1983
Melting point	-23 °C	Lide 1992
Boiling point	76.5 °C	Lide 1992
Density	1.594 g/mL	Lide 1992
Odor	Aromatic, sweet	HSDB 2004
Odor threshold:		
Water	0.52 mg/L	IRIS 2004
Air	10–71,000 mg/m ³ 96 ppm (600 mg/m ³) 60–1,500 mg/m ³	Verschueren 1983 Amoore and Hautala 1983 Ruth 1986
Solubility:		
Water at 20 °C	800 mg/L	Verschueren 1983
Organic solvent(s)	Miscible	HSDB 2004
Partition coefficients:		
Log K _{ow}	2.64	EPA 1984
Log K _{oc}	2.04	Kenaga 1980
Vapor pressure at 20 °C	90 mmHg	Verschueren 1983
Henry's law constant:		
at 25 °C	2.94x10 ⁻² atm·m ³ /mol	Yaws et al. 1991
at 24.8 °C	3.04x10 ⁻² atm·m ³ /mol	HSDB 2004
at 20 °C	2.04x10 ⁻² atm·m ³ /mol	Tse et al. 1992
at 30 °C	3.37x10 ⁻² atm·m ³ /mol	Tse et al. 1992
Autoignition temperature	Nonflammable	HSDB 2004
Flashpoint	Nonflammable	HSDB 2004
Flammability limits	Nonflammable	HSDB 2004
Conversion factors		
ppm (v/v) to mg/m ³ in air (25 °C)	1 ppm=6.39 mg/m ³	HSDB 2004
mg/m ³ to ppm (v/v) in air (25 °C)	1 mg/m ³ =0.16 ppm	Verschueren 1983
Explosive limits	No data	