

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET ALL CAUSES												
BOTH SEXES												
1959	938.6	2,748.7	106.9	47.7	107.5	147.8	293.9	743.3	1,711.7	3,757.3	8,584.9	19,417.5
1958	950.8	2,814.5	111.7	46.6	107.5	147.4	298.9	753.5	1,739.8	3,836.0	8,794.4	19,795.3
1957	958.6	2,797.5	111.8	48.8	115.6	152.8	307.1	764.9	1,780.6	3,902.7	8,807.5	19,789.8
1956	935.1	2,823.1	110.2	47.4	115.5	149.1	300.9	746.8	1,744.0	3,809.3	8,785.6	19,229.7
1955	930.4	2,848.5	113.4	48.8	115.6	149.6	306.8	756.3	1,729.7	3,811.1	8,794.6	18,983.3
1954	919.0	2,928.2	118.3	50.4	113.8	153.2	309.8	776.2	1,737.4	3,785.1	8,603.5	18,157.5
1953	959.0	3,076.2	130.0	54.8	123.8	162.7	332.0	817.2	1,838.5	3,939.0	9,063.5	19,187.3
1952	961.4	3,207.7	141.1	59.3	130.1	172.7	343.9	833.9	1,859.9	3,940.7	9,054.6	19,056.6
1951	966.7	3,234.4	136.9	59.3	128.0	176.6	353.6	847.2	1,881.2	4,005.5	9,235.9	19,758.8
1950	963.8	3,299.2	139.4	60.1	128.1	178.7	358.7	853.9	1,901.0	4,104.3	9,331.1	20,196.9
MALE												
1959	1,084.4	3,105.8	116.2	57.8	154.0	189.4	367.8	972.5	2,275.1	4,787.6	9,912.1	20,543.4
1958	1,096.7	3,163.9	119.3	55.7	154.6	188.0	373.1	977.1	2,298.1	4,852.4	10,135.3	20,833.4
1957	1,108.1	3,156.7	119.7	58.5	165.2	193.5	381.1	987.6	2,346.7	4,916.9	10,098.4	20,726.1
1956	1,082.5	3,201.3	118.8	56.8	167.3	191.6	370.7	964.1	2,288.7	4,762.2	10,078.0	20,045.3
1955	1,076.4	3,213.8	123.1	59.0	165.8	191.8	377.0	973.9	2,256.9	4,734.6	10,044.1	19,588.4
1954	1,065.3	3,308.3	128.4	60.2	163.7	196.2	377.2	987.1	2,254.2	4,673.8	9,800.6	18,741.1
1953	1,112.0	3,478.9	141.7	65.9	176.4	206.4	406.3	1,038.1	2,377.8	4,814.2	10,275.9	19,915.9
1952	1,111.9	3,613.2	152.0	69.6	180.6	215.6	417.0	1,054.5	2,386.4	4,786.6	10,228.7	19,673.6
1951	1,114.8	3,665.2	147.6	71.0	173.2	218.2	427.8	1,061.8	2,386.2	4,862.0	10,350.5	20,818.0
1950	1,106.1	3,728.0	151.7	70.9	167.9	216.5	428.8	1,067.1	2,395.3	4,931.4	10,426.0	21,636.0
FEMALE												
1959	797.1	2,379.9	97.3	37.2	62.2	107.8	223.3	520.7	1,182.3	2,856.5	7,551.0	18,681.4
1958	809.0	2,452.5	103.8	37.2	61.8	108.4	227.9	535.9	1,212.7	2,940.2	7,738.9	19,103.3
1957	813.2	2,425.2	103.6	38.8	68.0	113.8	236.2	547.6	1,243.2	3,000.7	7,781.6	19,151.0
1956	791.6	2,431.7	101.3	37.7	66.1	108.4	233.9	534.1	1,223.7	2,956.1	7,750.5	18,662.3
1955	788.2	2,469.4	103.3	38.2	67.9	109.4	239.2	543.0	1,222.6	2,978.0	7,785.5	18,554.8
1954	776.8	2,534.7	107.8	40.2	67.5	112.3	244.8	569.0	1,236.7	2,979.1	7,625.9	17,740.0
1953	810.4	2,659.0	118.0	43.3	75.3	121.4	260.3	599.7	1,312.3	3,139.9	8,064.8	18,663.4
1952	815.0	2,786.8	129.8	48.6	83.4	132.1	273.2	615.8	1,341.0	3,164.0	8,081.8	18,614.4
1951	821.9	2,789.0	125.9	47.2	85.2	137.3	281.6	634.3	1,379.0	3,213.4	8,296.7	19,005.6
1950	823.5	2,854.6	126.7	48.9	89.1	142.7	290.3	641.5	1,404.8	3,333.2	8,399.6	19,194.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET ALL CAUSES												
WHITE, BOTH SEXES												
1959	931.6	2,404.1	93.3	45.0	99.7	125.0	255.6	681.2	1,616.2	3,686.4	8,681.7	19,928.9
1958	941.3	2,451.9	97.9	44.4	99.8	123.7	261.5	685.9	1,638.3	3,757.8	8,888.8	20,327.3
1957	947.8	2,459.6	98.2	45.9	106.7	128.3	267.2	693.4	1,673.2	3,831.0	8,910.1	20,308.7
1956	926.3	2,501.6	97.4	44.8	107.0	125.9	261.4	678.5	1,642.2	3,752.5	8,894.0	19,778.8
1955	921.8	2,530.4	100.1	46.4	106.6	126.9	267.7	685.1	1,636.2	3,758.3	8,905.4	19,532.4
1954	908.4	2,610.4	105.1	47.8	103.4	129.1	269.2	699.8	1,645.5	3,739.8	8,708.5	18,617.3
1953	944.8	2,750.9	116.0	51.9	111.5	136.6	286.7	733.0	1,737.0	3,884.3	9,160.9	19,627.4
1952	944.8	2,859.4	125.3	56.3	116.0	144.3	296.0	747.9	1,756.1	3,889.2	9,150.1	19,478.2
1951	950.0	2,928.9	121.4	55.5	113.0	148.4	304.8	759.3	1,774.8	3,955.1	9,331.9	20,219.3
1950	945.7	2,992.6	124.1	56.4	111.7	148.3	307.6	765.0	1,799.6	4,023.1	9,416.5	20,678.6
WHITE, MALE												
1959	1,078.4	2,741.5	101.2	55.1	145.5	163.4	328.7	914.7	2,195.2	4,731.1	10,047.6	21,085.9
1958	1,088.7	2,773.0	105.8	53.4	145.9	162.2	336.6	916.1	2,212.8	4,788.2	10,267.8	21,345.6
1957	1,098.7	2,799.5	105.5	55.3	156.0	166.0	341.9	922.6	2,251.1	4,863.7	10,241.8	21,162.7
1956	1,075.6	2,849.7	105.7	54.0	157.5	166.0	333.0	903.7	2,201.0	4,726.7	10,229.6	20,508.6
1955	1,069.6	2,877.9	109.6	56.6	156.1	167.3	338.5	910.1	2,175.2	4,698.7	10,199.2	20,063.6
1954	1,055.9	2,969.9	114.6	57.4	152.3	169.9	338.3	917.6	2,170.7	4,645.8	9,936.3	19,081.2
1953	1,098.3	3,142.2	126.4	62.7	163.3	177.6	361.4	958.8	2,282.4	4,771.5	10,407.6	20,234.0
1952	1,095.6	3,233.4	135.3	66.5	166.1	185.4	369.6	975.0	2,289.4	4,751.2	10,335.2	19,990.9
1951	1,099.3	3,329.4	131.2	66.8	158.5	188.7	381.5	980.5	2,289.4	4,827.2	10,476.2	21,192.8
1950	1,089.5	3,400.5	135.5	67.2	152.4	185.3	380.9	984.5	2,304.4	4,864.9	10,526.3	22,116.3
WHITE, FEMALE												
1959	788.5	2,053.3	85.0	34.5	54.6	87.5	185.4	453.8	1,073.0	2,775.1	7,628.1	19,180.6
1958	797.6	2,117.2	89.7	35.0	54.7	86.2	189.2	461.5	1,097.3	2,851.6	7,814.5	19,654.2
1957	800.6	2,105.1	90.5	36.1	59.1	91.7	195.1	469.1	1,125.5	2,915.6	7,859.6	19,727.6
1956	780.4	2,137.5	88.8	35.2	58.5	87.1	192.2	457.8	1,109.6	2,882.3	7,836.4	19,271.0
1955	777.4	2,167.5	90.3	35.8	59.2	87.9	199.1	464.3	1,119.4	2,911.4	7,876.1	19,156.1
1954	764.6	2,235.6	95.2	37.9	57.6	90.0	202.3	485.5	1,137.8	2,918.8	7,716.0	18,284.5
1953	795.1	2,341.0	105.1	40.7	63.2	97.4	214.2	510.4	1,205.9	3,075.5	8,148.3	19,189.2
1952	797.5	2,468.3	114.8	45.8	69.4	105.2	224.3	523.4	1,232.0	3,100.3	8,177.2	19,109.7
1951	803.3	2,512.1	111.2	43.7	69.5	110.1	229.9	539.9	1,265.1	3,151.8	8,382.0	19,524.4
1950	803.3	2,566.8	112.2	45.1	71.5	112.8	235.8	546.4	1,293.8	3,242.8	8,481.5	19,679.5

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET ALL CAUSES												
ALL OTHER, BOTH SEXES												
1959	994.9	4,744.6	189.6	65.9	164.8	325.6	627.1	1,330.8	2,712.4	4,610.2	7,247.5	13,175.0
1958	1,027.1	4,983.4	196.4	62.1	163.2	335.6	624.5	1,397.1	2,824.3	4,780.3	7,480.8	13,395.5
1957	1,046.1	4,814.3	197.5	69.3	178.3	351.3	654.8	1,448.5	2,963.6	4,760.8	7,394.6	13,253.1
1956	1,007.4	4,758.2	192.2	65.8	175.6	339.1	643.2	1,400.4	2,896.2	4,488.0	7,247.8	12,604.9
1955	1,001.0	4,804.0	198.9	66.0	179.1	338.0	644.7	1,439.5	2,812.7	4,443.0	7,236.3	12,270.7
1954	1,006.9	4,938.0	204.1	68.6	188.5	354.5	660.4	1,509.5	2,826.8	4,327.4	7,111.1	12,183.6
1953	1,077.9	5,194.9	222.9	75.3	213.7	382.2	724.6	1,623.2	3,077.0	4,593.4	7,637.2	13,778.8
1952	1,101.8	5,517.0	247.9	80.3	233.2	412.4	761.1	1,654.3	3,159.7	4,556.6	7,653.5	13,851.0
1951	1,109.0	5,209.0	248.1	86.1	239.6	416.5	779.4	1,685.3	3,235.9	4,597.8	7,814.7	14,183.0
1950	1,119.4	5,368.4	250.8	86.0	251.3	440.2	805.3	1,706.1	3,126.6	5,205.0	8,039.7	14,473.6
ALL OTHER, MALE												
1959	1,133.2	5,255.3	209.0	76.9	216.9	405.1	719.5	1,529.1	3,098.5	5,450.5	8,180.6	14,444.8
1958	1,162.4	5,555.4	204.2	72.0	219.1	405.6	699.2	1,568.8	3,200.2	5,608.5	8,438.8	15,611.1
1957	1,186.0	5,337.6	210.9	81.4	232.6	429.9	730.4	1,615.3	3,382.3	5,556.7	8,255.6	15,503.8
1956	1,140.4	5,369.5	205.0	76.9	239.0	413.8	704.3	1,547.1	3,265.8	5,180.1	8,116.9	14,448.0
1955	1,133.3	5,348.3	212.2	76.4	237.0	405.9	716.1	1,589.0	3,191.2	5,159.4	8,018.4	13,766.7
1954	1,144.7	5,500.8	220.3	80.8	248.9	428.3	720.2	1,656.2	3,227.1	5,006.0	7,995.0	14,381.8
1953	1,228.1	5,708.5	244.7	89.2	276.8	460.3	803.6	1,798.4	3,511.5	5,304.1	8,501.8	15,857.1
1952	1,251.6	6,205.4	266.9	92.0	291.8	482.5	839.1	1,814.4	3,575.0	5,220.0	8,781.8	15,660.0
1951	1,248.2	5,889.1	266.6	101.3	287.0	480.5	842.6	1,834.8	3,593.9	5,272.8	8,721.7	16,142.1
1950	1,251.1	5,991.6	271.2	97.1	289.9	496.2	860.7	1,857.3	3,480.8	5,794.9	9,029.6	16,022.1
ALL OTHER, FEMALE												
1959	864.2	4,221.5	170.3	54.8	115.9	257.2	543.5	1,144.0	2,341.1	3,848.7	6,413.0	12,230.8
1958	899.1	4,413.4	188.5	52.1	111.7	275.3	555.8	1,235.9	2,469.1	4,014.8	6,610.5	12,173.7
1957	913.9	4,292.7	184.1	57.1	128.8	283.3	585.2	1,289.2	2,555.5	4,034.0	6,593.1	12,029.7
1956	881.6	4,151.5	179.5	54.6	118.3	274.4	587.3	1,257.7	2,533.3	3,831.9	6,429.0	11,325.0
1955	875.9	4,282.1	185.7	55.4	127.3	279.0	577.9	1,294.5	2,437.5	3,783.2	6,440.9	11,214.7
1954	877.1	4,379.6	188.0	56.3	136.0	290.5	604.8	1,366.5	2,426.5	3,700.5	6,219.8	11,053.1
1953	936.4	4,683.4	201.0	61.3	159.7	314.2	651.1	1,451.3	2,631.0	3,918.4	6,759.5	12,371.0
1952	960.3	4,837.8	228.8	68.6	182.6	350.9	688.7	1,496.2	2,733.0	3,934.2	6,583.6	12,603.4
1951	976.9	4,535.1	229.4	70.9	197.6	360.2	721.1	1,534.8	2,863.2	3,959.8	6,948.6	12,853.6
1950	993.5	4,749.0	230.3	75.0	216.4	390.4	754.0	1,554.9	2,763.0	4,610.7	7,064.7	13,366.8

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS, ALL FORMS (001-019)												
BOTH SEXES												
1959	6.5	1.2	0.8	0.2	0.7	2.8	5.3	10.3	16.7	24.7	33.0	38.0
1958	7.1	1.5	0.9	0.2	0.9	3.4	6.2	11.4	18.3	26.6	34.5	35.0
1957	7.8	1.9	0.9	0.2	1.2	3.9	7.1	12.7	19.3	28.9	36.6	39.1
1956	8.4	2.5	1.5	0.2	1.2	4.2	7.8	13.5	20.7	30.2	39.5	42.2
1955	9.1	2.6	1.7	0.3	1.6	4.8	8.9	15.0	21.6	32.0	39.7	39.4
1954	10.2	4.0	1.9	0.3	2.2	5.8	10.3	16.6	23.6	34.6	43.3	41.0
1953	12.4	4.1	3.1	0.6	3.4	7.7	12.5	20.0	29.3	39.2	47.7	45.7
1952	15.8	6.5	4.2	0.9	5.9	11.5	17.3	25.4	35.1	46.1	53.3	43.7
1951	20.1	7.1	5.2	1.4	8.8	16.9	22.7	32.7	43.9	52.8	58.9	54.2
1950	22.5	8.5	6.3	1.8	11.3	19.1	26.1	35.9	47.4	58.2	63.2	47.7
MALE												
1959	9.5	1.6	0.8	0.1	0.5	2.9	6.3	15.4	27.9	42.4	53.3	61.4
1958	10.5	1.5	1.0	0.2	0.8	3.1	7.7	17.9	30.6	45.0	54.2	54.2
1957	11.5	1.8	1.0	0.2	1.1	3.8	8.9	19.3	32.8	48.9	59.0	60.8
1956	12.1	2.5	1.4	0.2	0.9	3.8	9.8	20.6	34.8	49.7	62.2	62.1
1955	13.0	3.2	1.8	0.3	1.3	4.9	10.7	23.3	36.0	52.1	58.9	58.8
1954	14.5	4.3	1.9	0.3	1.7	5.6	12.8	25.3	39.7	56.1	66.0	62.5
1953	17.4	4.2	3.3	0.6	3.0	7.7	16.1	31.0	47.8	62.7	69.7	64.4
1952	21.9	6.3	4.4	0.9	5.1	11.5	22.0	39.6	57.9	72.3	76.1	63.0
1951	27.3	7.2	5.3	1.2	7.5	16.8	29.3	50.5	71.0	81.0	84.9	87.1
1950	30.1	8.6	6.4	1.6	9.0	18.7	33.5	55.7	76.3	90.5	88.7	68.4
FEMALE												
1959	3.6	*	0.9	0.2	0.8	2.7	4.4	5.2	6.2	9.3	17.2	22.7
1958	3.9	1.5	0.8	0.2	1.0	3.6	4.7	5.1	6.6	10.3	18.9	21.9
1957	4.3	2.0	0.9	0.2	1.2	4.0	5.4	6.2	6.6	11.0	18.7	24.3
1956	4.8	2.4	1.5	0.2	1.5	4.5	5.9	6.5	7.3	12.6	21.3	28.3
1955	5.3	1.9	1.7	0.3	1.9	4.8	7.1	6.9	7.7	13.8	24.3	25.6
1954	6.0	3.7	2.0	0.3	2.7	6.0	7.9	8.0	8.0	15.1	24.7	25.6
1953	7.4	4.0	2.9	0.6	3.8	7.8	9.0	9.3	11.2	17.7	29.7	32.3
1952	9.9	6.7	4.0	1.0	6.7	11.5	12.8	11.4	12.7	22.0	34.4	29.9
1951	13.1	7.0	5.2	1.6	10.0	17.0	16.3	15.0	16.9	26.7	37.2	30.8
1950	15.1	8.5	6.3	2.0	13.6	19.5	18.9	16.2	18.5	28.1	41.5	33.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS, ALL FORMS (001-019)												
WHITE, BOTH SEXES												
1959	5.5	0.7	0.5	0.1	0.4	1.4	3.6	8.1	14.2	22.7	31.3	35.7
1958	6.0	1.0	0.6	0.1	0.4	1.8	4.2	9.0	15.5	23.9	32.9	32.1
1957	6.6	1.0	0.6	0.1	0.6	2.2	5.2	10.1	16.5	26.5	34.6	37.5
1956	7.0	1.5	0.9	0.2	0.5	2.2	5.7	10.4	17.7	28.0	38.4	39.3
1955	7.6	1.6	1.1	0.2	0.7	2.7	6.3	11.8	18.8	29.6	37.9	37.0
1954	8.4	2.8	1.2	0.2	1.1	3.3	7.4	13.0	20.4	32.3	41.3	40.4
1953	10.1	2.3	2.1	0.4	1.6	4.4	8.9	15.6	25.2	36.1	46.2	44.9
1952	12.7	3.6	2.8	0.6	2.6	6.8	12.2	20.0	30.8	43.3	51.8	41.5
1951	16.2	5.0	3.6	0.9	4.5	10.8	16.8	26.0	38.2	49.4	57.0	52.2
1950	17.9	5.4	4.6	1.1	5.5	11.7	19.1	28.4	41.8	54.2	61.4	46.4
WHITE, MALE												
1959	8.3	*	0.5	*	0.3	1.3	4.4	12.2	24.4	39.5	50.5	57.4
1958	8.9	*	0.7	0.2	0.4	1.7	5.3	14.2	26.4	40.6	51.6	48.4
1957	10.0	*	0.7	0.1	0.6	2.0	6.5	15.5	28.5	45.2	56.7	56.9
1956	10.5	1.5	0.9	0.2	0.4	1.9	7.2	16.2	30.3	46.7	60.4	55.6
1955	11.2	1.8	1.1	0.2	0.6	2.8	7.6	18.9	31.7	48.9	56.1	54.8
1954	12.4	3.1	1.0	0.2	1.0	3.4	9.2	20.2	34.7	52.7	63.0	61.0
1953	14.6	2.6	2.3	0.4	1.5	4.2	11.4	24.7	41.8	58.2	67.8	62.7
1952	18.1	3.4	2.9	0.6	2.3	6.6	15.3	31.8	51.1	68.3	73.8	57.7
1951	22.8	4.9	3.6	0.8	3.8	10.5	21.9	40.9	62.8	76.4	81.5	83.5
1950	25.0	5.1	4.4	1.1	4.3	11.4	25.4	45.4	68.6	84.8	85.7	67.4
WHITE, FEMALE												
1959	2.8	*	0.5	*	0.4	1.4	2.9	4.1	4.7	8.0	16.4	21.6
1958	3.0	*	0.4	*	0.4	1.9	3.1	4.0	5.3	9.2	18.3	21.3
1957	3.3	*	0.5	*	0.5	2.4	3.8	4.7	5.2	10.0	17.2	24.3
1956	3.7	1.4	0.8	*	0.6	2.4	4.3	4.8	5.8	11.3	20.9	27.9
1955	4.1	1.3	1.1	0.2	0.7	2.7	5.1	5.0	6.3	12.3	23.4	24.3
1954	4.6	2.5	1.3	0.2	1.2	3.3	5.6	5.9	6.6	13.9	23.8	25.7
1953	5.6	2.1	1.9	0.3	1.7	4.6	6.4	6.7	9.1	16.0	28.7	32.1
1952	7.4	3.8	2.6	0.5	3.0	6.9	9.1	8.3	10.9	20.4	33.8	29.8
1951	9.8	5.1	3.5	0.9	5.2	11.0	11.8	11.1	13.8	24.5	36.7	29.8
1950	10.8	5.7	4.9	1.2	6.6	12.0	13.0	11.5	15.0	25.9	41.0	31.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS, ALL FORMS (001-019)												
ALL OTHER, BOTH SEXES												
1959	14.5	4.0	2.7	0.7	2.8	14.0	20.3	30.7	42.8	49.0	57.7	66.2
1958	16.6	4.5	3.0	0.6	4.6	15.9	23.7	34.0	48.0	58.8	56.5	69.7
1957	17.6	7.5	2.8	0.6	5.4	17.4	24.1	37.4	50.1	56.8	65.0	57.8
1956	19.5	8.4	5.4	0.7	6.0	20.6	26.2	43.0	54.3	55.9	55.2	77.0
1955	21.5	8.6	5.9	1.1	8.7	22.3	30.7	45.2	54.4	59.9	66.0	69.0
1954	24.6	11.8	7.0	1.2	10.4	26.4	35.8	50.9	61.7	62.6	71.7	47.3
1953	31.4	15.6	10.0	2.1	16.6	35.3	43.5	62.0	78.5	75.7	69.7	55.8
1952	42.1	25.7	13.7	3.7	30.2	51.5	62.0	77.2	89.0	79.3	74.3	71.4
1951	53.1	20.6	17.2	5.0	40.3	69.3	73.9	96.6	115.7	93.6	88.0	78.7
1950	62.3	29.3	18.7	6.6	55.3	83.0	87.1	107.8	115.4	112.1	89.5	62.5
ALL OTHER, MALE												
1959	19.5	*	2.5	*	1.7	15.4	23.8	46.4	64.8	76.5	88.9	106.9
1958	22.9	*	2.7	*	3.8	15.4	29.6	52.8	76.1	97.2	88.5	129.6
1957	24.1	7.0	2.8	*	4.7	18.4	29.8	55.4	79.4	93.1	88.9	107.7
1956	25.9	8.8	5.1	*	4.1	20.2	32.8	63.7	84.9	85.4	85.4	140.0
1955	28.4	11.9	6.8	1.1	6.9	23.4	38.2	66.1	84.9	89.5	95.2	108.3
1954	32.0	12.1	7.6	*	7.0	25.5	44.7	74.1	98.2	97.4	105.0	*
1953	41.1	15.0	10.3	1.8	14.4	38.1	57.6	91.2	119.1	115.1	94.7	*
1952	54.0	26.1	14.6	2.9	26.8	54.8	81.4	113.4	140.5	119.7	106.4	130.0
1951	66.6	22.3	17.8	3.9	36.2	72.7	95.3	141.9	172.2	136.1	133.0	131.6
1950	74.7	32.1	21.2	5.4	46.3	84.2	107.1	153.8	168.4	164.5	129.8	*
ALL OTHER, FEMALE												
1959	9.8	*	2.8	0.9	3.9	12.8	17.1	15.9	21.6	24.0	29.8	*
1958	10.7	*	3.3	*	5.2	16.3	18.2	16.2	21.0	23.8	27.5	*
1957	11.6	8.0	2.8	*	6.0	16.5	18.9	20.1	21.5	23.6	42.8	*
1956	13.5	8.0	5.7	*	7.7	20.9	20.1	23.0	24.2	28.6	26.8	*
1955	15.0	*	5.1	1.1	10.2	21.4	23.8	24.9	24.1	32.6	37.9	*
1954	17.6	11.6	6.4	1.5	13.3	27.2	27.6	28.3	25.3	30.4	39.7	*
1953	22.2	16.2	9.8	2.5	18.4	33.0	30.5	33.3	37.1	38.8	45.5	*
1952	30.8	25.3	12.8	4.5	33.1	48.6	44.1	41.5	36.0	41.5	44.0	*
1951	40.3	18.9	16.6	6.0	44.0	66.3	54.2	51.6	56.9	53.5	45.0	*
1950	50.6	26.5	16.2	7.7	63.5	82.0	68.6	61.8	60.9	59.2	49.8	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)												
BOTH SEXES												
1959	6.0	*	0.3	*	0.5	2.5	5.0	9.7	15.8	23.4	31.4	36.0
1958	6.6	0.6	0.3	0.1	0.8	3.0	5.7	10.9	17.3	25.2	32.9	33.3
1957	7.3	0.9	0.3	0.1	1.0	3.5	6.7	11.9	18.3	27.4	34.8	37.9
1956	7.8	1.1	0.5	0.1	1.0	3.8	7.3	12.7	19.5	28.4	37.6	41.2
1955	8.3	1.0	0.6	0.1	1.4	4.4	8.1	14.0	20.2	30.3	37.7	38.2
1954	9.3	1.9	0.7	0.1	1.9	5.3	9.6	15.6	22.0	32.6	40.8	39.9
1953	11.3	1.4	1.0	0.2	2.9	7.0	11.5	18.8	27.7	37.0	45.8	43.3
1952	14.4	2.8	1.3	0.4	5.1	10.6	16.1	23.9	33.4	43.7	50.6	41.6
1951	18.5	3.2	1.9	0.7	7.8	15.7	21.2	31.0	41.6	49.9	55.9	51.3
1950	20.6	3.4	2.1	0.9	10.0	17.7	24.7	34.1	44.9	55.2	60.3	44.5
MALE												
1959	8.9	*	*	*	0.3	2.5	6.0	14.8	26.7	40.4	51.3	58.9
1958	9.8	*	0.3	*	0.7	2.8	7.2	17.3	29.1	43.1	52.1	51.2
1957	10.8	*	0.3	*	0.9	3.3	8.4	18.4	31.3	46.7	56.1	59.3
1956	11.3	1.1	0.5	*	0.7	3.4	9.2	19.6	33.0	47.2	59.9	60.2
1955	12.1	1.3	0.5	*	1.1	4.5	9.8	22.0	34.2	49.8	55.9	58.2
1954	13.4	2.1	0.7	*	1.5	5.1	12.0	23.9	37.5	53.0	62.6	61.2
1953	16.1	1.6	1.2	0.2	2.5	6.8	15.0	29.4	45.7	59.5	66.9	61.2
1952	20.3	3.1	1.3	0.3	4.3	10.6	20.6	37.7	55.5	69.2	72.6	60.1
1951	25.4	3.3	1.9	0.5	6.5	15.6	27.5	48.3	68.3	77.1	80.6	81.6
1950	27.9	3.6	2.2	0.6	7.7	17.2	31.8	53.2	72.8	86.5	85.4	65.0
FEMALE												
1959	3.2	*	0.3	*	0.6	2.5	4.1	4.8	5.5	8.5	15.9	21.0
1958	3.5	*	*	*	0.9	3.2	4.4	4.7	6.1	9.4	17.8	21.3
1957	3.8	1.0	0.3	*	1.0	3.7	5.0	5.7	6.0	10.3	17.8	23.3
1956	4.3	1.2	0.5	*	1.3	4.1	5.5	6.0	6.5	11.6	19.8	27.9
1955	4.7	*	0.7	*	1.7	4.4	6.5	6.1	6.7	12.7	23.0	24.1
1954	5.3	1.7	0.8	0.1	2.3	5.5	7.2	7.3	7.1	14.0	23.1	24.7
1953	6.5	1.3	0.8	0.3	3.3	7.1	8.1	8.4	10.1	16.4	28.5	30.3
1952	8.7	2.6	1.4	0.5	5.9	10.6	11.9	10.3	11.6	20.3	32.4	28.3
1951	11.7	3.1	1.8	0.9	9.0	15.8	15.2	13.8	15.0	24.7	35.2	29.7
1950	13.5	3.3	2.0	1.2	12.3	18.3	17.7	15.1	17.0	26.0	38.8	30.3

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)												
WHITE, BOTH SEXES												
1959	5.2	*	*	*	0.3	1.2	3.4	7.8	13.6	21.6	29.9	33.7
1958	5.6	*	0.2	*	0.4	1.7	3.9	8.7	14.9	22.8	31.5	30.7
1957	6.2	*	0.2	*	0.4	2.1	4.9	9.6	15.9	25.3	33.1	36.5
1956	6.6	0.8	0.3	*	0.4	2.0	5.4	9.9	16.8	26.5	36.6	38.5
1955	7.1	0.7	0.3	*	0.5	2.6	5.9	11.3	17.7	28.3	36.0	36.0
1954	7.8	1.3	0.4	0.1	0.9	3.1	7.0	12.3	19.2	30.5	39.2	39.3
1953	9.4	0.9	0.7	0.1	1.4	4.1	8.3	14.8	24.1	34.2	44.4	42.9
1952	11.8	1.6	0.8	0.2	2.3	6.4	11.5	19.1	29.5	41.2	49.4	39.8
1951	15.1	2.4	1.2	0.3	4.0	10.1	15.9	24.8	36.5	46.8	54.0	49.6
1950	16.6	2.1	1.4	0.5	4.8	11.0	18.3	27.2	39.9	51.4	58.7	43.6
WHITE, MALE												
1959	7.9	*	*	*	0.2	1.2	4.2	11.8	23.6	37.8	48.9	54.9
1958	8.5	*	*	*	0.3	1.6	4.9	13.9	25.4	39.1	49.7	45.6
1957	9.5	*	*	*	0.5	1.9	6.2	14.9	27.5	43.4	54.4	55.9
1956	9.9	*	0.3	*	0.3	1.8	6.8	15.5	29.0	44.5	58.3	54.0
1955	10.5	*	*	*	0.4	2.6	7.1	18.1	30.3	47.1	53.5	54.1
1954	11.6	1.4	0.3	*	0.9	3.1	8.8	19.2	32.9	49.8	59.8	59.6
1953	13.8	*	0.9	*	1.3	3.8	10.8	23.7	40.3	55.5	65.3	60.4
1952	17.1	1.6	0.8	0.2	2.0	6.2	14.5	30.7	49.4	65.6	70.6	55.7
1951	21.4	2.4	1.2	0.3	3.3	9.8	20.7	39.4	60.9	73.0	77.5	78.9
1950	23.5	2.1	1.2	0.4	3.6	10.6	24.3	43.8	65.9	81.0	82.7	64.2
WHITE, FEMALE												
1959	2.5	*	*	*	0.3	1.3	2.8	3.9	4.3	7.5	15.2	20.0
1958	2.8	*	*	*	0.4	1.7	3.0	3.7	4.9	8.5	17.3	20.9
1957	3.1	*	*	*	0.4	2.3	3.6	4.4	4.8	9.3	16.4	23.2
1956	3.4	*	*	*	0.5	2.3	4.1	4.5	5.3	10.4	19.5	27.6
1955	3.7	*	0.4	*	0.6	2.5	4.8	4.6	5.6	11.3	22.1	23.1
1954	4.1	1.3	0.5	*	1.0	3.1	5.2	5.5	5.8	12.9	22.6	24.7
1953	5.1	*	0.5	*	1.4	4.4	5.9	6.1	8.3	14.8	27.5	30.2
1952	6.6	1.7	0.8	0.2	2.6	6.5	8.6	7.6	9.9	18.8	32.0	28.4
1951	8.8	2.4	1.1	0.4	4.7	10.4	11.2	10.3	12.3	22.7	34.6	28.6
1950	9.7	2.1	1.5	0.5	6.0	11.4	12.4	10.8	13.8	24.0	38.6	29.3

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)												
ALL OTHER, BOTH SEXES												
1959	12.8	*	1.0	*	2.1	12.5	18.8	28.0	38.3	44.9	52.5	63.2
1958	14.7	*	1.0	*	3.7	14.0	21.6	31.5	42.8	53.3	52.4	63.6
1957	15.6	*	1.0	*	4.8	15.2	22.0	34.7	45.0	52.0	58.2	54.7
1956	17.1	*	1.8	*	5.1	18.2	23.6	39.3	49.1	51.3	51.5	73.8
1955	18.6	*	2.0	*	7.4	19.9	27.3	40.0	48.7	54.4	61.7	65.5
1954	21.6	5.4	2.8	0.6	9.0	23.4	32.0	46.9	56.1	57.7	65.2	47.3
1953	27.2	5.3	3.3	1.1	14.4	30.9	39.3	56.8	71.0	69.8	65.8	48.1
1952	36.7	11.0	4.9	2.0	26.2	46.4	56.6	69.8	82.1	73.3	68.6	63.3
1951	47.2	8.8	6.8	3.2	35.6	63.3	67.6	89.9	106.4	86.6	83.4	72.3
1950	55.4	12.3	7.1	3.9	49.2	75.5	80.9	100.0	105.9	106.1	83.1	55.8
ALL OTHER, MALE												
1959	17.5	*	*	*	*	13.3	22.0	43.1	59.3	71.0	81.9	103.4
1958	20.7	*	*	*	3.2	13.5	27.2	50.1	67.6	89.7	82.7	118.5
1957	21.5	*	*	*	4.2	15.8	27.6	52.1	72.2	85.2	79.3	100.0
1956	23.3	*	*	*	3.4	17.9	30.1	59.5	78.4	78.7	80.0	136.0
1955	24.9	*	*	*	5.6	20.5	34.0	60.2	78.5	82.3	88.0	108.3
1954	28.6	*	3.1	*	5.9	22.3	40.5	69.2	90.0	90.6	100.0	*
1953	36.3	*	3.6	*	11.9	33.7	52.5	83.7	109.3	106.4	88.6	*
1952	47.7	13.5	4.4	1.3	21.9	49.2	74.8	104.5	130.0	111.3	100.0	115.0
1951	59.6	9.5	7.0	2.1	30.9	66.3	87.7	133.6	159.6	125.4	124.5	115.8
1950	66.6	13.3	9.1	2.2	39.8	75.9	99.8	143.4	154.7	157.2	123.9	*
ALL OTHER, FEMALE												
1959	8.3	*	*	*	2.8	11.8	15.8	13.9	18.2	21.3	26.1	*
1958	9.0	*	*	*	4.3	14.5	16.5	14.0	19.1	20.0	24.8	*
1957	10.0	*	*	*	5.3	14.8	16.8	18.1	18.5	21.7	38.6	*
1956	11.2	*	*	*	6.5	18.4	17.6	19.8	20.4	25.9	24.6	*
1955	12.6	*	2.0	*	9.0	19.4	21.0	20.4	19.1	28.8	36.4	*
1954	15.0	*	2.5	*	11.6	24.5	24.2	25.1	22.2	27.3	31.7	*
1953	18.7	*	3.0	1.4	16.5	28.4	27.1	30.3	32.1	35.5	43.8	*
1952	26.3	*	5.4	2.6	29.8	43.8	39.7	35.5	32.8	37.5	38.8	*
1951	35.4	*	6.7	4.3	39.8	60.6	49.1	46.5	51.1	50.0	44.1	*
1950	44.8	11.3	5.1	5.6	57.7	75.2	63.4	56.6	55.8	54.7	43.0	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS, OTHER FORMS (010-019)												
BOTH SEXES												
1959	0.5	0.8	0.6	0.1	0.2	0.3	0.3	0.5	0.9	1.3	1.6	*
1958	0.5	0.9	0.6	0.1	0.2	0.3	0.4	0.5	1.0	1.4	1.6	*
1957	0.6	1.0	0.6	0.1	0.2	0.4	0.5	0.7	1.0	1.5	1.8	*
1956	0.6	1.3	1.0	0.1	0.2	0.4	0.5	0.8	1.2	1.7	1.9	*
1955	0.8	1.5	1.2	0.2	0.3	0.4	0.7	1.0	1.4	1.7	2.1	*
1954	0.8	2.1	1.2	0.2	0.3	0.5	0.7	1.0	1.6	2.1	2.4	*
1953	1.1	2.6	2.1	0.4	0.5	0.8	1.0	1.2	1.6	2.2	1.9	*
1952	1.4	3.7	2.9	0.5	0.8	0.9	1.2	1.5	1.7	2.4	2.6	*
1951	1.7	3.8	3.4	0.7	1.0	1.2	1.4	1.7	2.3	2.9	3.0	*
1950	1.9	5.1	4.3	0.9	1.3	1.4	1.4	1.8	2.5	3.0	2.9	*
MALE												
1959	0.6	1.1	0.6	*	*	0.4	0.4	0.7	1.2	2.0	2.0	*
1958	0.6	*	0.7	0.1	*	0.3	0.6	0.6	1.6	1.9	2.2	*
1957	0.7	1.0	0.7	0.1	0.2	0.4	0.5	0.9	1.5	2.2	2.9	*
1956	0.8	1.4	1.0	*	*	0.4	0.6	1.0	1.7	2.5	2.4	*
1955	0.9	1.9	1.3	0.2	0.3	0.5	0.9	1.3	1.8	2.2	3.0	*
1954	1.0	2.2	1.2	0.2	0.3	0.6	0.8	1.3	2.3	3.2	3.4	*
1953	1.3	2.6	2.1	0.4	0.5	0.8	1.1	1.6	2.1	3.2	2.8	*
1952	1.6	3.2	3.1	0.6	0.8	0.9	1.4	1.9	2.4	3.1	3.4	*
1951	1.9	3.8	3.4	0.7	1.0	1.3	1.8	2.1	2.7	4.0	4.3	*
1950	2.2	5.0	4.2	1.0	1.3	1.5	1.7	2.5	3.6	4.1	3.3	*
FEMALE												
1959	0.4	*	0.6	0.2	0.2	0.2	0.2	0.4	0.7	0.7	1.3	*
1958	0.4	*	0.6	0.1	*	0.3	0.3	0.5	0.5	0.9	1.1	*
1957	0.4	1.0	0.6	*	0.2	0.3	0.4	0.5	0.6	0.8	1.0	*
1956	0.5	1.2	1.0	0.1	0.2	0.4	0.4	0.6	0.8	1.0	1.5	*
1955	0.6	1.1	1.0	0.2	0.3	0.3	0.6	0.7	1.0	1.1	1.3	*
1954	0.7	2.1	1.2	0.2	0.4	0.5	0.6	0.7	0.9	1.1	1.6	*
1953	0.9	2.7	2.1	0.3	0.5	0.7	0.9	0.8	1.1	1.3	1.2	*
1952	1.1	4.1	2.6	0.5	0.8	0.9	0.9	1.1	1.1	1.7	2.0	*
1951	1.4	3.9	3.4	0.7	1.0	1.2	1.1	1.2	1.8	2.0	2.0	*
1950	1.6	5.2	4.3	0.8	1.2	1.3	1.1	1.1	1.5	2.1	2.7	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS, OTHER FORMS (010-019)												
WHITE, BOTH SEXES												
1959	0.3	*	0.4	0.1	*	0.1	0.2	0.3	0.6	1.1	1.4	*
1958	0.3	*	0.4	0.1	*	0.1	0.2	0.3	0.6	1.0	1.4	*
1957	0.4	*	0.4	0.1	0.1	0.1	0.3	0.5	0.6	1.2	1.5	*
1956	0.4	0.7	0.6	0.1	*	0.2	0.3	0.5	0.9	1.5	1.8	*
1955	0.5	0.8	0.7	0.2	0.1	0.2	0.4	0.6	1.1	1.4	1.9	*
1954	0.6	1.5	0.7	0.1	0.2	0.2	0.4	0.7	1.2	1.9	2.1	*
1953	0.7	1.5	1.4	0.3	0.3	0.3	0.6	0.8	1.1	1.9	1.8	*
1952	0.9	2.0	2.0	0.4	0.4	0.4	0.7	0.9	1.3	2.1	2.4	*
1951	1.2	2.6	2.4	0.5	0.5	0.6	0.9	1.1	1.7	2.6	2.9	*
1950	1.3	3.3	3.3	0.7	0.6	0.7	0.9	1.2	2.0	2.8	2.7	*
WHITE, MALE												
1959	0.4	*	0.4	*	*	*	0.2	0.4	0.8	1.7	1.6	*
1958	0.4	*	0.5	0.1	*	*	0.4	0.4	0.9	1.5	1.9	*
1957	0.5	*	0.5	*	*	*	0.3	0.7	0.9	1.7	2.4	*
1956	0.6	*	0.6	*	*	*	0.3	0.7	1.3	2.2	2.1	*
1955	0.6	*	0.8	0.2	*	*	0.5	0.8	1.4	1.8	2.6	*
1954	0.8	1.7	0.7	*	*	0.3	0.5	0.9	1.7	2.8	3.3	*
1953	0.9	1.5	1.4	0.3	0.2	0.4	0.7	1.0	1.5	2.7	2.5	*
1952	1.0	1.8	2.1	0.4	0.3	0.4	0.8	1.1	1.7	2.7	3.2	*
1951	1.3	2.5	2.4	0.5	0.4	0.7	1.1	1.5	1.9	3.4	4.0	*
1950	1.5	3.0	3.2	0.7	0.6	0.7	1.1	1.7	2.7	3.8	3.1	*
WHITE, FEMALE												
1959	0.3	*	0.4	*	*	*	*	0.2	0.5	0.6	1.2	*
1958	0.2	*	0.3	*	*	*	*	0.3	0.4	0.7	1.0	*
1957	0.3	*	0.4	*	*	*	0.2	0.4	0.4	0.7	*	*
1956	0.3	*	0.6	*	*	*	0.2	0.3	0.5	0.9	1.5	*
1955	0.4	*	0.7	*	*	*	0.3	0.4	0.7	0.9	1.3	*
1954	0.4	*	0.8	*	*	0.2	0.3	0.4	0.7	1.0	1.2	*
1953	0.6	1.4	1.4	0.2	0.3	0.2	0.6	0.6	0.8	1.2	1.2	*
1952	0.7	2.1	1.9	0.3	0.4	0.4	0.5	0.6	1.0	1.5	1.8	*
1951	1.0	2.8	2.4	0.6	0.5	0.6	0.6	0.8	1.5	1.8	2.1	*
1950	1.1	3.7	3.3	0.7	0.6	0.6	0.7	0.7	1.2	1.9	2.4	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
TUBERCULOSIS, OTHER FORMS (010-019)												
ALL OTHER, BOTH SEXES												
1959	1.7	*	1.7	0.4	0.7	1.6	1.5	2.6	4.4	4.1	*	*
1958	2.0	*	2.0	*	0.8	1.9	2.1	2.5	5.2	5.5	*	*
1957	2.0	4.2	1.8	*	*	2.2	2.2	2.7	5.0	4.8	*	*
1956	2.4	5.1	3.6	*	0.9	2.4	2.5	3.7	5.2	4.7	*	*
1955	3.0	5.7	4.0	0.7	1.3	2.4	3.5	5.2	5.7	5.4	*	*
1954	3.0	6.4	4.2	0.6	1.4	3.0	3.8	4.0	5.6	4.9	*	*
1953	4.1	10.2	6.7	1.0	2.2	4.5	4.2	5.2	7.5	5.9	*	*
1952	5.4	14.8	8.9	1.7	4.0	5.1	5.4	7.4	6.9	6.1	*	*
1951	5.9	11.8	10.4	1.8	4.7	6.0	6.3	6.7	9.3	7.0	*	*
1950	6.9	17.0	11.6	2.6	6.1	7.5	6.2	7.8	9.5	5.9	*	*
ALL OTHER, MALE												
1959	2.0	*	1.8	*	*	2.2	1.8	3.3	5.5	5.6	*	*
1958	2.3	*	1.8	*	*	1.9	2.4	2.7	8.5	7.5	*	*
1957	2.5	*	1.9	*	*	2.7	2.2	3.3	7.1	7.9	*	*
1956	2.6	*	3.3	*	*	2.3	2.6	4.3	6.5	6.7	*	*
1955	3.5	*	4.9	*	*	2.9	4.1	5.9	6.4	7.2	*	*
1954	3.3	*	4.5	*	*	3.2	4.2	4.9	8.2	6.8	*	*
1953	4.8	9.4	6.7	*	2.5	4.4	5.1	7.5	9.8	8.7	*	*
1952	6.4	12.6	10.3	1.5	4.9	5.6	6.6	8.9	10.5	8.4	*	*
1951	7.0	12.7	10.8	1.8	5.3	6.4	7.6	8.3	12.6	10.7	*	*
1950	8.1	18.8	12.0	3.2	6.5	8.3	7.4	10.4	13.7	7.3	*	*
ALL OTHER, FEMALE												
1959	1.4	*	*	*	*	*	*	2.0	3.4	*	*	*
1958	1.7	*	2.3	*	*	1.8	1.7	2.3	*	*	*	*
1957	1.5	*	*	*	*	1.7	2.1	*	*	*	*	*
1956	2.2	*	3.9	*	*	2.5	2.4	3.1	3.8	*	*	*
1955	2.4	*	3.0	*	*	2.0	2.9	4.4	5.0	*	*	*
1954	2.7	*	3.9	*	1.7	2.8	3.5	3.2	*	*	*	*
1953	3.5	11.1	6.7	*	1.9	4.6	3.4	3.0	5.1	*	*	*
1952	4.5	16.9	7.4	1.8	3.3	4.8	4.3	6.0	*	*	*	*
1951	5.0	10.8	10.0	1.7	4.2	5.7	5.1	5.1	5.7	*	*	*
1950	5.8	15.2	11.1	2.1	5.8	6.7	5.2	5.1	5.2	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SYPHILIS AND ITS SEQUELAE (020-029)												
BOTH SEXES												
1959	1.7	0.5	*	*	*	0.2	0.5	2.2	6.0	9.2	9.6	8.7
1958	2.0	0.6	*	*	*	0.2	0.6	2.7	6.9	10.2	10.9	10.9
1957	2.2	0.7	*	*	*	0.2	0.8	3.4	7.6	11.2	11.6	11.0
1956	2.3	0.7	*	*	*	0.2	0.8	3.7	8.1	11.4	10.7	10.4
1955	2.3	0.9	*	*	*	0.2	1.0	4.0	7.9	11.2	10.7	7.3
1954	3.0	1.2	*	*	0.1	0.4	1.3	4.8	10.1	14.8	14.9	12.1
1953	3.3	1.6	*	*	0.1	0.3	1.7	5.8	11.4	15.5	14.7	12.2
1952	3.7	2.7	*	*	0.2	0.4	2.0	6.8	12.4	16.6	15.2	14.1
1951	4.1	3.9	*	*	0.2	0.5	2.3	7.8	14.0	17.5	16.1	11.0
1950	5.0	6.4	*	*	0.3	0.7	3.3	9.3	15.8	22.0	22.7	16.6
MALE												
1959	2.5	*	*	*	*	0.2	0.8	3.0	9.3	14.7	13.2	13.8
1958	2.9	*	*	*	*	0.3	0.8	3.7	11.0	16.1	16.6	15.4
1957	3.3	*	*	*	*	0.2	1.1	4.9	11.9	18.0	17.5	14.8
1956	3.4	*	*	*	*	0.2	1.1	5.5	12.8	18.2	15.9	15.0
1955	3.5	1.0	*	*	*	0.3	1.5	5.8	12.6	17.9	16.6	9.1
1954	4.4	1.3	*	*	*	0.4	1.7	6.8	16.1	23.8	22.5	15.1
1953	5.0	1.8	*	*	*	0.3	2.3	8.8	18.1	24.7	22.2	15.6
1952	5.5	3.1	*	*	*	0.4	2.6	10.2	19.9	26.7	23.6	20.5
1951	6.2	4.0	*	*	0.2	0.6	3.1	11.9	22.5	27.9	24.2	12.1
1950	7.4	7.1	*	*	0.3	0.8	4.3	13.8	25.1	35.1	33.8	20.7
FEMALE												
1959	1.0	*	*	*	*	*	0.3	1.5	2.8	4.3	6.7	5.3
1958	1.1	*	*	*	*	*	0.5	1.7	3.1	5.0	6.4	7.9
1957	1.2	*	*	*	*	0.2	0.6	1.9	3.6	5.2	6.9	8.3
1956	1.2	*	*	*	*	0.2	0.6	1.9	3.6	5.3	6.6	7.2
1955	1.2	*	*	*	*	0.2	0.6	2.2	3.4	5.2	5.9	6.0
1954	1.6	*	*	*	*	0.3	0.9	2.8	4.3	6.6	8.6	9.9
1953	1.7	1.3	*	*	*	0.3	1.2	2.9	4.9	7.2	8.5	9.7
1952	1.8	2.3	*	*	0.2	0.3	1.4	3.4	5.0	7.3	8.3	9.4
1951	2.1	3.8	*	*	*	0.4	1.6	3.8	5.6	7.9	9.3	10.3
1950	2.7	5.6	*	*	0.2	0.6	2.4	4.8	6.6	9.9	13.2	13.8

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SYPHILIS AND ITS SEQUELAE (020-029)												
WHITE, BOTH SEXES												
1959	1.3	*	*	*	*	*	0.3	1.2	4.2	7.8	8.4	7.8
1958	1.5	*	*	*	*	0.1	0.2	1.4	4.8	8.2	9.1	9.1
1957	1.7	*	*	*	*	0.1	0.4	1.9	5.5	9.1	9.8	9.1
1956	1.7	*	*	*	*	0.1	0.3	2.2	5.7	9.5	9.4	8.8
1955	1.7	*	*	*	*	0.1	0.5	2.2	5.5	9.0	9.1	5.8
1954	2.3	*	*	*	*	0.1	0.6	2.6	7.3	12.7	13.9	10.4
1953	2.4	*	*	*	*	0.1	0.7	3.2	8.3	13.2	13.0	11.1
1952	2.7	1.1	*	*	0.1	0.1	0.9	3.9	9.2	14.2	13.4	13.1
1951	3.0	1.4	*	*	0.1	0.2	1.1	4.6	10.5	14.9	14.6	9.8
1950	3.7	2.7	*	*	0.2	0.3	1.5	5.3	12.1	19.4	20.7	15.6
WHITE, MALE												
1959	2.0	*	*	*	*	*	0.4	1.7	6.9	12.9	11.5	11.7
1958	2.2	*	*	*	*	*	0.3	2.0	7.7	13.2	13.7	12.0
1957	2.5	*	*	*	*	*	0.4	2.8	8.9	14.8	14.6	12.5
1956	2.6	*	*	*	*	*	0.4	3.4	9.4	15.4	13.4	13.9
1955	2.5	*	*	*	*	*	0.7	3.2	8.9	14.5	14.3	7.5
1954	3.4	*	*	*	*	*	0.8	3.7	11.8	20.7	21.2	11.3
1953	3.7	*	*	*	*	*	1.0	5.0	13.4	21.4	19.3	14.2
1952	4.2	*	*	*	*	0.2	1.1	6.1	15.0	23.4	21.2	18.2
1951	4.6	*	*	*	*	0.2	1.5	7.2	17.4	24.1	21.9	11.4
1950	5.6	3.1	*	*	0.2	0.3	1.9	8.0	19.6	31.3	30.9	18.8
WHITE, FEMALE												
1959	0.7	*	*	*	*	*	*	0.7	1.7	3.4	6.0	5.4
1958	0.8	*	*	*	*	*	*	0.8	1.9	3.9	5.6	7.1
1957	0.9	*	*	*	*	*	0.3	1.0	2.3	4.1	5.9	6.8
1956	0.9	*	*	*	*	*	0.2	1.1	2.1	4.2	6.1	5.3
1955	0.8	*	*	*	*	*	0.3	1.2	2.3	4.0	4.9	*
1954	1.1	*	*	*	*	*	0.3	1.5	2.9	5.4	8.0	9.7
1953	1.2	*	*	*	*	*	0.5	1.4	3.3	5.7	8.0	8.9
1952	1.3	*	*	*	*	*	0.6	1.8	3.5	5.8	7.1	9.4
1951	1.4	1.5	*	*	*	0.2	0.7	2.0	3.7	6.5	8.6	8.7
1950	1.9	2.3	*	*	*	0.2	1.1	2.7	4.7	8.3	12.0	13.4

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SYPHILIS AND ITS SEQUELAE (020-029)												
ALL OTHER, BOTH SEXES												
1959	4.9	*	*	*	*	*	2.9	11.9	24.4	24.8	25.6	*
1958	6.4	3.5	*	*	*	0.8	4.1	15.0	30.1	34.4	36.0	33.3
1957	6.9	*	*	*	*	1.0	4.8	17.8	30.8	35.8	37.1	32.8
1956	7.1	*	*	*	*	1.0	5.3	17.4	35.8	34.4	30.2	*
1955	7.9	4.8	*	*	*	1.2	6.1	21.5	35.7	38.3	33.6	*
1954	9.2	6.2	*	*	*	2.3	7.6	25.8	43.3	39.8	28.7	*
1953	10.9	9.0	*	*	*	1.9	10.6	31.0	49.3	43.5	39.3	*
1952	11.7	13.4	*	*	*	2.2	11.6	33.6	51.7	45.4	40.7	*
1951	13.4	20.1	*	*	0.8	3.0	13.6	38.7	58.6	48.7	38.2	*
1950	16.1	31.3	*	*	0.8	4.6	19.1	47.1	60.3	58.3	53.1	*
ALL OTHER, MALE												
1959	6.8	*	*	*	*	*	4.2	15.4	34.5	36.1	34.7	*
1958	9.1	*	*	*	*	*	5.1	20.1	45.2	50.6	54.0	*
1957	9.8	*	*	*	*	*	6.7	25.2	43.8	55.4	54.1	*
1956	10.2	*	*	*	*	*	7.0	25.3	51.3	51.5	47.7	*
1955	11.6	*	*	*	*	*	9.1	31.0	55.1	58.0	47.2	*
1954	13.3	8.5	*	*	*	2.5	9.3	36.9	65.6	60.5	39.5	*
1953	15.7	10.7	*	*	*	1.8	13.8	44.9	73.9	63.4	62.3	*
1952	17.1	16.2	*	*	*	2.1	15.8	49.6	79.2	66.6	55.5	*
1951	19.3	21.8	*	*	*	4.4	18.2	56.1	85.8	73.7	56.6	*
1950	23.2	35.1	*	*	*	5.9	25.4	69.1	90.3	84.5	74.3	*
ALL OTHER, FEMALE												
1959	3.1	*	*	*	*	*	1.7	8.6	14.7	14.6	17.4	*
1958	3.8	*	*	*	*	*	3.1	10.1	15.5	19.5	19.6	*
1957	4.1	*	*	*	*	*	3.2	10.7	18.2	17.8	21.4	*
1956	4.2	*	*	*	*	*	3.8	9.8	20.6	18.5	*	*
1955	4.3	*	*	*	*	*	3.2	12.3	16.5	20.1	20.5	*
1954	5.4	*	*	*	*	2.1	6.0	14.9	21.0	20.7	18.3	*
1953	6.3	*	*	*	*	2.1	7.6	17.4	24.4	24.9	17.4	*
1952	6.6	10.7	*	*	*	2.2	7.7	17.8	23.6	25.5	26.7	*
1951	7.7	18.5	*	*	*	1.8	9.3	21.5	30.2	25.1	20.7	*
1950	9.4	27.5	*	*	*	3.4	13.3	25.1	29.6	31.9	32.2	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DYSENTERY, ALL FORMS (045-048)												
BOTH SEXES												
1959	0.2	4.2	0.3	*	*	*	0.1	*	0.2	0.3	0.8	*
1958	0.2	4.6	0.3	*	*	*	*	0.1	0.1	0.4	0.7	*
1957	0.3	5.4	0.4	*	*	*	*	*	0.2	0.4	0.7	*
1956	0.3	5.6	0.3	0.1	*	*	*	0.1	0.3	0.5	1.2	*
1955	0.3	7.2	0.4	0.1	*	*	*	*	0.2	0.5	1.0	3.4
1954	0.4	8.3	0.5	*	*	*	0.1	0.2	0.3	0.6	1.5	3.2
1953	0.6	13.8	0.8	0.1	*	*	*	0.2	0.2	0.6	1.5	3.8
1952	0.6	15.2	0.8	0.1	*	*	0.1	0.2	0.4	0.6	1.5	4.7
1951	0.7	17.1	1.1	0.1	*	*	0.1	0.2	0.3	0.8	2.1	3.2
1950	0.6	16.6	0.9	0.1	*	0.1	0.1	0.2	0.4	0.6	1.3	3.5
MALE												
1959	0.2	4.3	0.3	*	*	*	*	*	*	*	1.0	*
1958	0.3	5.7	0.4	*	*	*	*	*	*	0.6	*	*
1957	0.3	5.3	0.3	*	*	*	*	*	*	0.5	*	*
1956	0.3	5.8	0.3	*	*	*	*	*	0.4	0.6	1.2	*
1955	0.4	7.2	0.5	*	*	*	*	*	*	0.6	*	*
1954	0.4	8.4	0.6	*	*	*	*	0.2	0.3	0.9	1.6	*
1953	0.6	15.1	0.8	*	*	*	*	0.3	0.3	0.7	1.6	*
1952	0.7	16.3	0.9	*	*	*	*	*	0.3	0.7	1.8	*
1951	0.7	18.9	1.0	*	*	*	*	0.3	0.4	0.8	2.1	*
1950	0.7	19.1	0.9	*	*	*	0.2	0.2	0.5	0.7	*	*
FEMALE												
1959	0.2	4.1	0.3	*	*	*	*	*	*	*	*	*
1958	0.2	3.5	0.3	*	*	*	*	*	*	*	*	*
1957	0.2	5.5	0.4	*	*	*	*	*	*	*	*	*
1956	0.3	5.5	*	*	*	*	*	*	*	*	1.2	*
1955	0.3	7.2	0.4	*	*	*	*	*	*	0.4	1.1	*
1954	0.4	8.1	0.4	*	*	*	*	*	0.3	*	1.4	*
1953	0.5	12.5	0.7	*	*	*	*	*	*	0.5	1.4	*
1952	0.6	14.0	0.8	*	*	*	*	*	0.4	0.5	1.3	5.2
1951	0.6	15.2	1.2	*	*	*	*	*	0.3	0.8	2.0	*
1950	0.5	13.9	0.8	*	*	*	*	*	*	0.5	1.3	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DYSENTERY, ALL FORMS (045-048)												
WHITE, BOTH SEXES												
1959	0.2	2.7	0.2	*	*	*	*	*	*	*	0.7	*
1958	0.2	3.2	0.2	*	*	*	*	*	*	0.3	0.6	*
1957	0.2	3.7	0.3	*	*	*	*	*	*	0.2	0.7	*
1956	0.2	3.8	0.2	0.1	*	*	*	*	*	0.4	1.1	*
1955	0.2	5.2	0.3	0.1	*	*	*	*	*	0.3	0.9	3.1
1954	0.3	5.3	0.3	*	*	*	*	*	0.2	0.4	1.4	*
1953	0.4	10.3	0.6	0.1	*	*	*	0.1	0.2	0.5	1.2	3.6
1952	0.5	11.1	0.5	0.1	*	*	*	0.1	0.3	0.5	1.4	4.0
1951	0.6	14.6	0.9	*	*	*	*	0.2	0.3	0.6	1.8	*
1950	0.5	14.6	0.7	0.1	*	*	0.1	*	0.3	0.5	1.1	*
WHITE, MALE												
1959	0.2	3.0	*	*	*	*	*	*	*	*	*	*
1958	0.2	3.8	*	*	*	*	*	*	*	*	*	*
1957	0.2	3.1	*	*	*	*	*	*	*	*	*	*
1956	0.2	3.8	*	*	*	*	*	*	*	*	*	*
1955	0.3	4.9	0.4	*	*	*	*	*	*	*	*	*
1954	0.3	5.5	0.4	*	*	*	*	*	*	0.6	1.4	*
1953	0.5	10.8	0.6	*	*	*	*	0.2	*	0.6	1.3	*
1952	0.5	11.9	0.6	*	*	*	*	*	*	0.6	1.5	*
1951	0.6	15.9	0.8	*	*	*	*	*	0.3	0.7	2.0	*
1950	0.6	17.0	0.8	*	*	*	*	*	0.4	0.5	*	*
WHITE, FEMALE												
1959	0.1	2.5	0.3	*	*	*	*	*	*	*	*	*
1958	0.1	2.6	*	*	*	*	*	*	*	*	*	*
1957	0.2	4.3	*	*	*	*	*	*	*	*	*	*
1956	0.2	3.8	*	*	*	*	*	*	*	*	1.2	*
1955	0.2	5.6	*	*	*	*	*	*	*	*	*	*
1954	0.3	5.1	*	*	*	*	*	*	*	*	1.5	*
1953	0.4	9.8	0.6	*	*	*	*	*	*	*	1.2	*
1952	0.4	10.1	0.5	*	*	*	*	*	0.4	0.5	1.2	*
1951	0.5	13.3	1.0	*	*	*	*	*	*	0.6	1.7	*
1950	0.4	12.2	0.7	*	*	*	*	*	*	*	1.2	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DYSENTERY, ALL FORMS (045-048)												
ALL OTHER, BOTH SEXES												
1959	0.7	12.6	*	*	*	*	*	*	*	*	*	*
1958	0.8	13.1	1.0	*	*	*	*	*	*	*	*	*
1957	0.8	15.5	*	*	*	*	*	*	*	*	*	*
1956	0.9	16.8	*	*	*	*	*	*	1.7	*	*	*
1955	1.1	19.4	1.3	*	*	*	*	*	*	2.6	*	*
1954	1.5	26.9	1.9	*	*	*	*	*	*	3.1	*	*
1953	1.7	36.9	1.9	*	*	*	*	*	*	*	*	*
1952	1.9	42.3	2.8	*	*	*	*	*	*	*	*	*
1951	1.7	33.0	2.7	*	*	*	*	*	*	*	*	*
1950	1.5	29.5	2.0	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	0.8	11.7	*	*	*	*	*	*	*	*	*	*
1958	1.0	17.3	*	*	*	*	*	*	*	*	*	*
1957	1.0	18.8	*	*	*	*	*	*	*	*	*	*
1956	1.1	18.4	*	*	*	*	*	*	*	*	*	*
1955	1.2	21.8	*	*	*	*	*	*	*	*	*	*
1954	1.7	27.4	*	*	*	*	*	*	*	*	*	*
1953	2.0	44.0	2.2	*	*	*	*	*	*	*	*	*
1952	2.1	45.9	2.8	*	*	*	*	*	*	*	*	*
1951	1.8	39.1	2.9	*	*	*	*	*	*	*	*	*
1950	1.8	33.6	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	0.7	13.6	*	*	*	*	*	*	*	*	*	*
1958	0.5	9.0	*	*	*	*	*	*	*	*	*	*
1957	0.7	12.2	*	*	*	*	*	*	*	*	*	*
1956	0.8	15.3	*	*	*	*	*	*	*	*	*	*
1955	1.0	17.1	*	*	*	*	*	*	*	*	*	*
1954	1.3	26.4	*	*	*	*	*	*	*	*	*	*
1953	1.4	29.8	*	*	*	*	*	*	*	*	*	*
1952	1.6	38.7	2.8	*	*	*	*	*	*	*	*	*
1951	1.5	27.0	2.6	*	*	*	*	*	*	*	*	*
1950	1.1	25.5	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)												
BOTH SEXES												
1959	0.1	*	*	0.1	*	*	*	*	*	*	*	*
1958	0.1	*	0.2	*	*	*	*	*	*	*	*	*
1957	0.1	0.6	0.2	0.1	*	*	0.1	0.1	0.2	*	*	*
1956	0.1	0.7	0.2	0.1	*	0.1	0.1	0.1	*	0.2	0.6	*
1955	0.1	0.7	0.3	0.1	*	*	0.1	0.1	*	*	*	*
1954	0.2	0.7	0.2	0.1	0.2	0.1	0.1	*	*	*	*	*
1953	0.2	1.0	0.3	0.2	0.1	0.1	0.1	0.2	*	0.3	*	*
1952	0.2	1.1	0.4	0.2	0.1	0.1	0.1	0.1	0.3	0.3	0.6	*
1951	0.2	1.2	0.4	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.7	*
1950	0.2	0.9	0.5	0.3	0.1	*	0.1	0.1	0.2	0.3	0.7	*
MALE												
1959	0.1	*	*	*	*	*	*	*	*	*	*	*
1958	0.1	*	0.3	*	*	*	*	*	*	*	*	*
1957	0.1	*	*	*	*	*	*	*	*	*	*	*
1956	0.1	*	*	0.1	*	*	*	*	*	*	*	*
1955	0.2	*	0.3	*	*	*	*	*	*	*	*	*
1954	0.2	*	*	0.1	*	*	0.2	*	*	*	*	*
1953	0.2	1.3	*	*	0.2	*	*	*	*	0.5	*	*
1952	0.2	1.2	0.5	0.2	*	*	*	*	0.3	*	*	*
1951	0.2	1.5	0.4	0.2	*	*	0.2	*	*	*	*	*
1950	0.2	1.2	0.6	0.3	*	*	0.2	*	*	*	*	*
FEMALE												
1959	0.1	*	*	*	*	*	*	*	*	*	*	*
1958	0.1	*	*	*	*	*	*	*	*	*	*	*
1957	0.1	*	*	*	*	*	*	*	*	*	*	*
1956	0.1	*	*	*	*	*	*	*	*	*	*	*
1955	0.1	*	*	*	*	*	*	*	*	*	*	*
1954	0.2	*	*	0.2	*	*	*	*	*	*	*	*
1953	0.2	*	0.4	0.2	*	*	*	*	*	*	*	*
1952	0.2	*	0.4	0.2	*	*	*	*	*	*	*	*
1951	0.2	*	0.4	*	*	0.2	*	*	*	*	*	*
1950	0.2	*	0.5	0.3	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)												
WHITE, BOTH SEXES												
1959	0.1	*	*	0.1	*	*	*	*	*	*	*	*
1958	0.1	*	0.2	*	*	*	*	*	*	*	*	*
1957	0.1	*	0.2	0.1	*	*	0.1	*	0.2	*	*	*
1956	0.1	*	0.2	0.1	*	*	*	*	*	0.2	0.6	*
1955	0.1	0.7	0.3	0.1	*	*	*	*	*	*	*	*
1954	0.2	*	0.2	0.2	0.1	*	0.1	*	*	*	*	*
1953	0.2	0.9	0.3	0.2	0.1	0.1	*	0.2	*	0.3	*	*
1952	0.2	1.0	0.4	0.2	0.1	0.1	*	*	0.3	0.3	0.6	*
1951	0.2	1.2	0.4	0.1	0.1	0.1	0.1	0.2	0.2	*	0.7	*
1950	0.2	0.8	0.5	0.3	0.1	*	0.1	*	*	0.3	0.7	*
WHITE, MALE												
1959	0.1	*	*	*	*	*	*	*	*	*	*	*
1958	0.1	*	*	*	*	*	*	*	*	*	*	*
1957	0.1	*	*	*	*	*	*	*	*	*	*	*
1956	0.1	*	*	*	*	*	*	*	*	*	*	*
1955	0.1	*	*	*	*	*	*	*	*	*	*	*
1954	0.2	*	*	*	*	*	*	*	*	*	*	*
1953	0.2	*	*	*	*	*	*	*	*	*	*	*
1952	0.2	*	0.4	*	*	*	*	*	0.3	*	*	*
1951	0.2	1.6	0.3	*	*	*	*	*	*	*	*	*
1950	0.2	*	0.6	0.3	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	0.1	*	*	*	*	*	*	*	*	*	*	*
1958	0.1	*	*	*	*	*	*	*	*	*	*	*
1957	0.1	*	*	*	*	*	*	*	*	*	*	*
1956	0.1	*	*	*	*	*	*	*	*	*	*	*
1955	0.1	*	*	*	*	*	*	*	*	*	*	*
1954	0.1	*	*	0.2	*	*	*	*	*	*	*	*
1953	0.2	*	0.4	0.2	*	*	*	*	*	*	*	*
1952	0.2	*	0.4	0.2	*	*	*	*	*	*	*	*
1951	0.2	*	0.4	*	*	*	*	*	*	*	*	*
1950	0.2	*	0.4	0.3	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)												
ALL OTHER, BOTH SEXES												
1959	0.1	*	*	*	*	*	*	*	*	*	*	*
1958	0.1	*	*	*	*	*	*	*	*	*	*	*
1957	0.2	*	*	*	*	*	*	*	*	*	*	*
1956	0.2	*	*	*	*	*	*	*	*	*	*	*
1955	0.2	*	*	*	*	*	*	*	*	*	*	*
1954	0.3	*	*	*	*	*	*	*	*	*	*	*
1953	0.2	*	*	*	*	*	*	*	*	*	*	*
1952	0.3	*	*	*	*	*	*	*	*	*	*	*
1951	0.3	*	*	*	*	*	*	*	*	*	*	*
1950	0.3	*	*	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	*	*	*	*	*	*	*	*	*	*	*	*
1958	*	*	*	*	*	*	*	*	*	*	*	*
1957	0.3	*	*	*	*	*	*	*	*	*	*	*
1956	0.3	*	*	*	*	*	*	*	*	*	*	*
1955	0.2	*	*	*	*	*	*	*	*	*	*	*
1954	*	*	*	*	*	*	*	*	*	*	*	*
1953	*	*	*	*	*	*	*	*	*	*	*	*
1952	0.3	*	*	*	*	*	*	*	*	*	*	*
1951	0.4	*	*	*	*	*	*	*	*	*	*	*
1950	0.3	*	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	*	*	*	*	*	*	*	*	*	*	*	*
1958	*	*	*	*	*	*	*	*	*	*	*	*
1957	*	*	*	*	*	*	*	*	*	*	*	*
1956	*	*	*	*	*	*	*	*	*	*	*	*
1955	*	*	*	*	*	*	*	*	*	*	*	*
1954	0.3	*	*	*	*	*	*	*	*	*	*	*
1953	*	*	*	*	*	*	*	*	*	*	*	*
1952	0.2	*	*	*	*	*	*	*	*	*	*	*
1951	0.3	*	*	*	*	*	*	*	*	*	*	*
1950	0.4	*	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SEPTICEMIA (053)												
BOTH SEXES												
1959	1.0	10.4	0.9	0.2	0.2	0.2	0.4	0.7	1.5	2.7	4.6	7.9
1958	1.0	10.7	0.9	0.1	0.2	0.2	0.4	0.6	1.4	2.4	4.7	8.3
1957	0.8	10.8	0.8	0.2	0.2	0.1	0.2	0.5	1.0	1.8	3.4	3.0
1956	0.7	8.6	1.0	0.1	0.1	0.2	0.2	0.3	0.8	1.6	2.6	4.3
1955	0.6	8.3	0.6	0.2	0.1	0.1	0.2	0.4	0.7	1.2	1.9	3.3
1954	0.6	8.0	0.7	0.2	0.1	0.2	0.2	0.4	0.5	1.1	1.7	3.3
1953	0.5	6.6	0.8	0.2	0.1	0.1	0.2	0.4	0.6	0.8	1.5	*
1952	0.5	7.2	0.7	0.2	0.1	0.1	0.2	0.3	0.5	0.6	1.3	*
1951	0.4	5.8	0.7	0.1	0.1	0.1	0.1	0.3	0.6	0.8	1.1	*
1950	0.4	5.7	0.6	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.5	*
MALE												
1959	1.1	11.4	1.0	0.2	*	0.2	0.3	0.8	2.0	3.4	5.6	9.6
1958	1.1	12.1	1.0	*	*	0.2	0.3	0.7	1.8	3.2	6.1	10.2
1957	1.0	12.3	0.8	0.2	0.2	*	0.3	0.7	1.3	2.6	4.7	*
1956	0.8	10.3	0.9	0.1	*	0.2	0.3	0.4	1.1	1.8	3.5	*
1955	0.7	9.7	0.7	0.2	*	*	*	0.4	0.9	1.6	2.6	*
1954	0.6	8.7	0.9	0.2	*	*	0.2	0.5	0.7	1.3	1.9	*
1953	0.6	7.8	0.9	0.2	*	*	*	0.4	0.9	1.2	1.6	*
1952	0.5	7.8	0.8	*	*	*	*	0.3	0.7	0.9	1.9	*
1951	0.5	6.4	0.8	0.2	*	*	*	0.3	0.5	1.2	1.3	*
1950	0.4	6.2	0.7	*	*	*	0.2	0.4	0.6	0.7	2.0	*
FEMALE												
1959	0.9	9.3	0.7	0.2	*	0.3	0.4	0.7	1.0	2.2	3.9	6.8
1958	0.8	9.3	0.8	0.2	0.3	0.3	0.4	0.5	1.0	1.7	3.6	7.0
1957	0.6	9.2	0.7	0.1	*	0.2	0.2	0.4	0.7	1.1	2.4	*
1956	0.6	6.8	1.0	*	*	*	0.2	0.3	0.5	1.5	1.9	4.3
1955	0.5	6.8	0.6	0.2	*	*	0.2	0.3	0.5	0.8	1.4	*
1954	0.5	7.4	0.6	0.2	*	0.2	0.2	0.3	0.4	1.0	1.6	*
1953	0.4	5.3	0.7	*	*	*	*	0.3	0.3	0.5	1.4	*
1952	0.4	6.6	0.6	0.2	*	0.2	0.2	0.3	*	*	*	*
1951	0.4	5.2	0.6	*	*	0.2	*	0.3	0.6	0.5	*	*
1950	0.4	5.2	0.5	*	*	*	0.2	*	*	0.6	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SEPTICEMIA (053)												
WHITE, BOTH SEXES												
1959	0.9	8.2	0.7	0.2	0.1	0.2	0.3	0.6	1.3	2.7	4.6	7.6
1958	0.9	8.9	0.8	0.1	0.2	0.2	0.3	0.5	1.3	2.3	4.6	8.3
1957	0.7	9.2	0.7	0.2	0.2	0.1	0.2	0.5	0.9	1.7	3.3	3.1
1956	0.6	7.4	0.9	0.1	0.1	0.2	0.2	0.3	0.7	1.7	2.5	4.2
1955	0.5	7.3	0.6	0.2	0.1	*	0.2	0.3	0.6	1.1	1.9	3.4
1954	0.5	7.0	0.7	0.2	0.1	0.1	0.2	0.3	0.5	1.1	1.7	3.6
1953	0.4	6.0	0.7	0.1	*	0.1	0.1	0.3	0.5	0.8	1.5	*
1952	0.4	6.3	0.7	0.1	*	0.1	0.2	0.3	0.4	0.6	1.3	*
1951	0.4	5.3	0.7	0.2	*	0.1	*	0.3	0.5	0.8	1.1	*
1950	0.4	5.7	0.6	0.1	*	*	0.2	0.3	0.4	0.6	1.4	*
WHITE, MALE												
1959	1.0	9.1	0.9	0.2	*	*	0.3	0.7	1.8	3.3	5.3	8.3
1958	1.0	10.2	0.9	*	*	*	0.3	0.6	1.6	3.0	6.0	10.1
1957	0.9	10.4	0.7	0.2	*	*	0.3	0.6	1.2	2.5	4.4	*
1956	0.7	9.2	0.9	*	*	0.2	0.2	0.3	1.0	1.9	3.4	*
1955	0.6	9.0	0.7	0.2	*	*	*	0.4	0.8	1.4	2.6	*
1954	0.6	7.8	0.9	0.2	*	*	*	0.4	0.5	1.2	1.8	*
1953	0.5	6.9	0.9	*	*	*	*	0.3	0.8	1.2	1.6	*
1952	0.5	6.9	0.8	*	*	*	*	0.3	0.7	0.9	1.8	*
1951	0.4	5.8	0.8	*	*	*	*	0.3	0.5	1.1	*	*
1950	0.4	6.2	0.7	*	*	*	*	0.4	0.6	0.6	1.8	*
WHITE, FEMALE												
1959	0.8	7.3	0.6	0.1	*	*	0.3	0.5	0.9	2.1	4.0	7.1
1958	0.8	7.4	0.7	0.2	0.2	0.3	0.3	0.5	0.9	1.7	3.5	7.1
1957	0.6	8.0	0.7	*	*	*	*	0.3	0.6	1.1	2.4	*
1956	0.5	5.5	1.0	*	*	*	*	*	0.4	1.5	1.9	*
1955	0.4	5.5	0.6	*	*	*	*	0.3	0.4	0.8	1.4	*
1954	0.4	6.2	0.5	*	*	*	0.2	0.2	0.4	1.0	1.6	*
1953	0.4	5.0	0.6	*	*	*	*	*	*	0.5	1.4	*
1952	0.3	5.6	0.6	0.2	*	*	*	*	*	*	*	*
1951	0.3	4.8	0.6	*	*	*	*	*	0.5	0.5	*	*
1950	0.3	5.2	0.5	*	*	*	*	*	*	0.5	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SEPTICEMIA (053)												
ALL OTHER, BOTH SEXES												
1959	1.9	23.0	1.5	*	*	*	0.9	2.3	2.7	3.6	*	*
1958	1.7	21.8	1.3	*	*	*	1.0	1.3	2.6	3.5	*	*
1957	1.4	20.2	1.1	*	*	*	*	1.2	2.3	3.0	*	*
1956	1.2	15.8	1.3	*	*	*	*	*	1.8	*	*	*
1955	1.0	14.5	*	*	*	*	*	*	*	*	*	*
1954	1.0	14.7	*	*	*	*	*	*	*	*	*	*
1953	0.9	10.7	*	*	*	*	*	1.4	*	*	*	*
1952	0.8	13.4	*	*	*	*	*	*	*	*	*	*
1951	0.8	9.0	*	*	*	*	*	*	*	*	*	*
1950	0.7	5.4	*	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	2.1	25.3	*	*	*	*	*	2.4	3.7	*	*	*
1958	1.9	23.5	*	*	*	*	*	*	*	5.4	*	*
1957	1.7	23.7	*	*	*	*	*	*	*	*	*	*
1956	1.2	16.9	*	*	*	*	*	*	*	*	*	*
1955	1.0	14.6	*	*	*	*	*	*	*	*	*	*
1954	1.0	14.5	*	*	*	*	*	*	*	*	*	*
1953	1.2	14.1	*	*	*	*	*	*	*	*	*	*
1952	0.8	13.5	*	*	*	*	*	*	*	*	*	*
1951	0.8	10.5	*	*	*	*	*	*	*	*	*	*
1950	0.6	*	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	1.7	20.5	*	*	*	*	*	2.2	*	*	*	*
1958	1.5	20.0	*	*	*	*	*	*	*	*	*	*
1957	1.2	16.7	*	*	*	*	*	*	*	*	*	*
1956	1.1	14.6	*	*	*	*	*	*	*	*	*	*
1955	1.0	14.4	*	*	*	*	*	*	*	*	*	*
1954	0.9	14.8	*	*	*	*	*	*	*	*	*	*
1953	0.6	*	*	*	*	*	*	*	*	*	*	*
1952	0.8	13.3	*	*	*	*	*	*	*	*	*	*
1951	0.8	*	*	*	*	*	*	*	*	*	*	*
1950	0.8	*	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DIPHTHERIA (055)												
BOTH SEXES												
1959	0.0	*	0.2	0.1	*	*	*	*	*	*	*	*
1958	0.0	*	0.2	0.1	*	*	*	*	*	*	*	*
1957	0.0	*	0.3	0.1	*	*	*	*	*	*	*	*
1956	0.1	*	0.3	0.1	*	*	*	*	*	*	*	*
1955	0.1	*	0.4	0.2	*	*	*	*	*	*	*	*
1954	0.1	*	0.5	0.2	*	*	*	*	*	*	*	*
1953	0.1	*	0.5	0.2	*	*	*	*	*	*	*	*
1952	0.1	*	0.7	0.2	*	*	*	*	*	*	*	*
1951	0.2	*	1.1	0.3	*	*	*	*	*	*	*	*
1950	0.3	0.7	1.6	0.5	*	*	*	*	*	*	*	*
MALE												
1959	0.0	*	*	*	*	*	*	*	*	*	*	*
1958	0.0	*	*	*	*	*	*	*	*	*	*	*
1957	0.0	*	0.3	*	*	*	*	*	*	*	*	*
1956	0.1	*	0.3	0.1	*	*	*	*	*	*	*	*
1955	0.1	*	0.4	0.2	*	*	*	*	*	*	*	*
1954	0.1	*	0.6	0.2	*	*	*	*	*	*	*	*
1953	0.1	*	0.5	0.2	*	*	*	*	*	*	*	*
1952	0.2	*	0.7	0.2	*	*	*	*	*	*	*	*
1951	0.2	*	1.3	0.3	*	*	*	*	*	*	*	*
1950	0.3	*	1.6	0.5	*	*	*	*	*	*	*	*
FEMALE												
1959	0.0	*	*	*	*	*	*	*	*	*	*	*
1958	0.0	*	*	*	*	*	*	*	*	*	*	*
1957	0.1	*	0.3	*	*	*	*	*	*	*	*	*
1956	0.0	*	*	*	*	*	*	*	*	*	*	*
1955	0.1	*	0.4	0.2	*	*	*	*	*	*	*	*
1954	0.1	*	0.4	0.2	*	*	*	*	*	*	*	*
1953	0.1	*	0.4	0.2	*	*	*	*	*	*	*	*
1952	0.1	*	0.7	0.2	*	*	*	*	*	*	*	*
1951	0.2	*	0.9	0.3	*	*	*	*	*	*	*	*
1950	0.3	*	1.6	0.5	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DIPHTHERIA (055)												
WHITE, BOTH SEXES												
1959	0.0	*	0.2	*	*	*	*	*	*	*	*	*
1958	0.0	*	*	0.1	*	*	*	*	*	*	*	*
1957	0.0	*	0.2	*	*	*	*	*	*	*	*	*
1956	0.0	*	*	0.1	*	*	*	*	*	*	*	*
1955	0.1	*	0.3	0.1	*	*	*	*	*	*	*	*
1954	0.1	*	0.2	0.1	*	*	*	*	*	*	*	*
1953	0.1	*	0.3	0.1	*	*	*	*	*	*	*	*
1952	0.1	*	0.5	0.2	*	*	*	*	*	*	*	*
1951	0.2	*	0.8	0.3	*	*	*	*	*	*	*	*
1950	0.2	*	1.4	0.4	*	*	*	*	*	*	*	*
WHITE, MALE												
1959	0.0	*	*	*	*	*	*	*	*	*	*	*
1958	0.0	*	*	*	*	*	*	*	*	*	*	*
1957	0.0	*	*	*	*	*	*	*	*	*	*	*
1956	0.0	*	*	*	*	*	*	*	*	*	*	*
1955	0.1	*	*	*	*	*	*	*	*	*	*	*
1954	0.1	*	*	*	*	*	*	*	*	*	*	*
1953	0.1	*	*	*	*	*	*	*	*	*	*	*
1952	0.1	*	0.5	*	*	*	*	*	*	*	*	*
1951	0.2	*	0.9	0.3	*	*	*	*	*	*	*	*
1950	0.3	*	1.4	0.4	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	0.0	*	*	*	*	*	*	*	*	*	*	*
1958	0.0	*	*	*	*	*	*	*	*	*	*	*
1957	0.0	*	*	*	*	*	*	*	*	*	*	*
1956	0.0	*	*	*	*	*	*	*	*	*	*	*
1955	0.0	*	*	*	*	*	*	*	*	*	*	*
1954	0.1	*	*	*	*	*	*	*	*	*	*	*
1953	0.1	*	*	*	*	*	*	*	*	*	*	*
1952	0.1	*	0.6	0.2	*	*	*	*	*	*	*	*
1951	0.1	*	0.8	0.3	*	*	*	*	*	*	*	*
1950	0.2	*	1.4	0.4	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DIPHTHERIA (055)												
ALL OTHER, BOTH SEXES												
1959	0.1	*	*	*	*	*	*	*	*	*	*	*
1958	0.1	*	*	*	*	*	*	*	*	*	*	*
1957	0.2	*	0.9	*	*	*	*	*	*	*	*	*
1956	0.2	*	1.1	*	*	*	*	*	*	*	*	*
1955	0.3	*	1.4	0.8	*	*	*	*	*	*	*	*
1954	0.3	*	2.0	*	*	*	*	*	*	*	*	*
1953	0.3	*	1.5	*	*	*	*	*	*	*	*	*
1952	0.3	*	1.8	*	*	*	*	*	*	*	*	*
1951	0.5	*	2.7	*	*	*	*	*	*	*	*	*
1950	0.6	*	3.4	0.9	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	*	*	*	*	*	*	*	*	*	*	*	*
1958	*	*	*	*	*	*	*	*	*	*	*	*
1957	*	*	*	*	*	*	*	*	*	*	*	*
1956	0.3	*	*	*	*	*	*	*	*	*	*	*
1955	0.3	*	*	*	*	*	*	*	*	*	*	*
1954	0.4	*	2.6	*	*	*	*	*	*	*	*	*
1953	0.4	*	*	*	*	*	*	*	*	*	*	*
1952	0.4	*	*	*	*	*	*	*	*	*	*	*
1951	0.6	*	3.9	*	*	*	*	*	*	*	*	*
1950	0.6	*	3.3	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	*	*	*	*	*	*	*	*	*	*	*	*
1958	*	*	*	*	*	*	*	*	*	*	*	*
1957	*	*	*	*	*	*	*	*	*	*	*	*
1956	*	*	*	*	*	*	*	*	*	*	*	*
1955	0.3	*	*	*	*	*	*	*	*	*	*	*
1954	0.3	*	*	*	*	*	*	*	*	*	*	*
1953	0.3	*	*	*	*	*	*	*	*	*	*	*
1952	0.3	*	*	*	*	*	*	*	*	*	*	*
1951	0.3	*	*	*	*	*	*	*	*	*	*	*
1950	0.5	*	3.6	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
WHOOPING COUGH (056)												
BOTH SEXES												
1959	0.2	4.5	0.5	*	*	*	*	*	*	*	*	*
1958	0.1	3.2	0.3	*	*	*	*	*	*	*	*	*
1957	0.1	3.1	0.3	*	*	*	*	*	*	*	*	*
1956	0.2	4.5	0.5	*	*	*	*	*	*	*	*	*
1955	0.3	8.9	0.8	*	*	*	*	*	*	*	*	*
1954	0.2	7.2	0.7	*	*	*	*	*	*	*	*	*
1953	0.2	5.2	0.5	*	*	*	*	*	*	*	*	*
1952	0.3	7.7	0.9	*	*	*	*	*	*	*	*	*
1951	0.6	19.7	1.9	0.1	*	*	*	*	*	*	*	*
1950	0.7	23.7	2.5	0.2	*	*	*	*	*	*	*	*
MALE												
1959	0.1	3.4	0.4	*	*	*	*	*	*	*	*	*
1958	0.1	2.9	0.3	*	*	*	*	*	*	*	*	*
1957	0.1	2.6	0.3	*	*	*	*	*	*	*	*	*
1956	0.2	4.3	0.5	*	*	*	*	*	*	*	*	*
1955	0.3	8.1	0.6	*	*	*	*	*	*	*	*	*
1954	0.2	6.0	0.7	*	*	*	*	*	*	*	*	*
1953	0.2	4.7	0.4	*	*	*	*	*	*	*	*	*
1952	0.3	7.5	0.9	*	*	*	*	*	*	*	*	*
1951	0.6	21.0	1.4	*	*	*	*	*	*	*	*	*
1950	0.7	21.8	1.9	*	*	*	*	*	*	*	*	*
FEMALE												
1959	0.2	5.6	0.6	*	*	*	*	*	*	*	*	*
1958	0.1	3.5	0.3	*	*	*	*	*	*	*	*	*
1957	0.1	3.6	0.3	*	*	*	*	*	*	*	*	*
1956	0.2	4.7	0.6	*	*	*	*	*	*	*	*	*
1955	0.3	9.8	1.1	*	*	*	*	*	*	*	*	*
1954	0.3	8.5	0.7	*	*	*	*	*	*	*	*	*
1953	0.2	5.8	0.6	*	*	*	*	*	*	*	*	*
1952	0.3	7.9	0.9	*	*	*	*	*	*	*	*	*
1951	0.6	18.4	2.3	*	*	*	*	*	*	*	*	*
1950	0.8	25.6	3.2	0.2	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
WHOOPING COUGH (056)												
WHITE, BOTH SEXES												
1959	0.1	2.2	0.3	*	*	*	*	*	*	*	*	*
1958	0.1	1.9	0.2	*	*	*	*	*	*	*	*	*
1957	0.1	1.8	0.2	*	*	*	*	*	*	*	*	*
1956	0.1	2.0	0.2	*	*	*	*	*	*	*	*	*
1955	0.2	4.7	0.5	*	*	*	*	*	*	*	*	*
1954	0.2	5.1	0.5	*	*	*	*	*	*	*	*	*
1953	0.1	3.4	0.4	*	*	*	*	*	*	*	*	*
1952	0.1	4.2	0.6	*	*	*	*	*	*	*	*	*
1951	0.3	11.1	0.9	*	*	*	*	*	*	*	*	*
1950	0.5	17.4	1.7	0.1	*	*	*	*	*	*	*	*
WHITE, MALE												
1959	0.1	1.7	*	*	*	*	*	*	*	*	*	*
1958	0.1	1.6	*	*	*	*	*	*	*	*	*	*
1957	0.1	1.5	*	*	*	*	*	*	*	*	*	*
1956	0.1	1.9	*	*	*	*	*	*	*	*	*	*
1955	0.1	4.1	0.3	*	*	*	*	*	*	*	*	*
1954	0.1	4.0	0.5	*	*	*	*	*	*	*	*	*
1953	0.1	2.9	*	*	*	*	*	*	*	*	*	*
1952	0.1	3.9	0.6	*	*	*	*	*	*	*	*	*
1951	0.3	12.1	0.6	*	*	*	*	*	*	*	*	*
1950	0.5	15.9	1.2	*	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	0.1	2.8	0.4	*	*	*	*	*	*	*	*	*
1958	0.1	2.2	*	*	*	*	*	*	*	*	*	*
1957	0.1	2.0	*	*	*	*	*	*	*	*	*	*
1956	0.1	2.0	*	*	*	*	*	*	*	*	*	*
1955	0.2	5.4	0.7	*	*	*	*	*	*	*	*	*
1954	0.2	6.3	0.4	*	*	*	*	*	*	*	*	*
1953	0.1	3.9	0.4	*	*	*	*	*	*	*	*	*
1952	0.2	4.5	0.6	*	*	*	*	*	*	*	*	*
1951	0.3	10.1	1.2	*	*	*	*	*	*	*	*	*
1950	0.6	18.9	2.4	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
WHOOPING COUGH (056)												
ALL OTHER, BOTH SEXES												
1959	0.7	17.5	1.8	*	*	*	*	*	*	*	*	*
1958	0.4	11.1	1.0	*	*	*	*	*	*	*	*	*
1957	0.5	10.8	1.4	*	*	*	*	*	*	*	*	*
1956	0.9	19.6	2.7	*	*	*	*	*	*	*	*	*
1955	1.4	34.7	2.9	*	*	*	*	*	*	*	*	*
1954	0.9	20.3	2.5	*	*	*	*	*	*	*	*	*
1953	0.6	17.3	1.4	*	*	*	*	*	*	*	*	*
1952	1.2	30.9	2.9	*	*	*	*	*	*	*	*	*
1951	3.1	75.3	8.6	*	*	*	*	*	*	*	*	*
1950	2.6	66.2	8.1	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	0.6	13.7	*	*	*	*	*	*	*	*	*	*
1958	0.5	10.7	*	*	*	*	*	*	*	*	*	*
1957	0.5	9.1	*	*	*	*	*	*	*	*	*	*
1956	0.9	19.1	2.5	*	*	*	*	*	*	*	*	*
1955	1.3	33.3	2.5	*	*	*	*	*	*	*	*	*
1954	0.8	18.5	2.4	*	*	*	*	*	*	*	*	*
1953	0.6	17.1	*	*	*	*	*	*	*	*	*	*
1952	1.2	32.0	2.7	*	*	*	*	*	*	*	*	*
1951	3.2	80.0	7.2	*	*	*	*	*	*	*	*	*
1950	2.5	63.2	7.1	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	0.9	21.2	2.1	*	*	*	*	*	*	*	*	*
1958	0.4	11.4	*	*	*	*	*	*	*	*	*	*
1957	0.6	12.5	*	*	*	*	*	*	*	*	*	*
1956	0.9	20.1	2.8	*	*	*	*	*	*	*	*	*
1955	1.4	36.1	3.3	*	*	*	*	*	*	*	*	*
1954	0.9	22.0	2.5	*	*	*	*	*	*	*	*	*
1953	0.7	17.4	*	*	*	*	*	*	*	*	*	*
1952	1.2	29.8	3.1	*	*	*	*	*	*	*	*	*
1951	3.0	70.7	10.0	*	*	*	*	*	*	*	*	*
1950	2.8	69.2	9.2	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MENINGOCOCCAL INFECTIONS (057)												
BOTH SEXES												
1959	0.4	5.0	1.5	0.3	0.2	0.1	*	0.1	0.2	0.2	*	*
1958	0.4	6.0	1.7	0.2	0.1	*	*	0.2	0.2	0.3	*	*
1957	0.5	6.1	1.8	0.3	0.2	*	*	0.1	0.2	0.3	*	*
1956	0.5	6.6	1.7	0.3	0.2	*	0.1	0.1	0.2	*	*	*
1955	0.6	6.8	2.2	0.3	0.3	0.1	0.1	0.2	0.2	0.2	*	*
1954	0.6	8.3	2.4	0.5	0.3	*	0.1	0.2	0.2	0.3	*	*
1953	0.8	11.2	3.2	0.7	0.4	0.1	0.1	0.3	0.3	0.3	*	*
1952	0.9	11.6	3.7	0.8	0.4	0.1	0.1	0.2	0.3	0.3	*	*
1951	0.7	9.2	2.8	0.6	0.4	0.1	0.2	0.3	0.2	0.4	*	*
1950	0.6	8.0	2.6	0.5	0.3	0.1	0.2	0.3	0.3	0.4	*	*
MALE												
1959	0.4	5.5	1.7	0.3	0.2	*	*	*	*	*	*	*
1958	0.4	6.3	1.7	0.2	*	*	*	*	*	*	*	*
1957	0.5	7.0	2.1	0.3	0.2	*	*	*	*	*	*	*
1956	0.5	7.5	2.0	0.3	0.2	*	*	*	*	*	*	*
1955	0.6	7.8	2.4	0.4	0.4	*	*	0.2	0.3	*	*	*
1954	0.7	9.2	2.7	0.6	0.5	*	*	*	*	*	*	*
1953	1.0	13.1	3.6	0.8	0.5	*	*	0.3	0.4	*	*	*
1952	1.0	12.1	4.2	0.8	0.5	*	*	0.2	*	*	*	*
1951	0.8	10.7	2.9	0.5	0.6	*	0.2	0.4	*	*	*	*
1950	0.7	8.7	3.1	0.5	0.3	*	0.2	0.3	0.3	*	*	*
FEMALE												
1959	0.3	4.5	1.3	0.2	*	*	*	*	*	*	*	*
1958	0.4	5.6	1.7	0.2	*	*	*	*	*	*	*	*
1957	0.4	5.2	1.5	0.3	*	*	*	*	*	*	*	*
1956	0.4	5.6	1.4	0.3	0.2	*	*	*	*	*	*	*
1955	0.5	5.8	2.1	0.3	*	*	*	*	*	*	*	*
1954	0.5	7.4	2.2	0.5	*	*	*	*	*	*	*	*
1953	0.7	9.1	2.8	0.6	0.4	*	*	0.2	0.3	*	*	*
1952	0.8	11.0	3.2	0.8	0.4	*	*	*	*	*	*	*
1951	0.7	7.6	2.7	0.6	0.3	*	*	0.2	*	*	*	*
1950	0.5	7.3	2.2	0.5	0.2	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MENINGOCOCCAL INFECTIONS (057)												
WHITE, BOTH SEXES												
1959	0.4	5.0	1.4	0.2	0.2	*	*	*	0.2	*	*	*
1958	0.4	5.8	1.7	0.2	0.1	*	*	0.1	0.2	0.3	*	*
1957	0.4	6.4	1.8	0.3	0.2	*	*	*	*	0.3	*	*
1956	0.4	6.5	1.8	0.3	0.2	*	*	*	0.2	*	*	*
1955	0.5	6.8	2.2	0.3	0.2	0.1	*	0.2	0.2	*	*	*
1954	0.6	8.3	2.5	0.5	0.2	*	0.1	0.2	0.2	0.3	*	*
1953	0.8	11.3	3.2	0.6	0.4	0.1	0.1	0.3	0.3	0.3	*	*
1952	0.9	11.8	3.7	0.7	0.4	0.1	0.1	0.2	0.3	0.3	*	*
1951	0.7	8.9	2.9	0.5	0.4	0.1	0.1	0.2	0.2	0.4	*	*
1950	0.6	8.0	2.7	0.5	0.3	0.1	0.2	0.2	0.2	0.3	*	*
WHITE, MALE												
1959	0.4	5.8	1.7	0.2	0.2	*	*	*	*	*	*	*
1958	0.4	5.9	1.7	0.2	*	*	*	*	*	*	*	*
1957	0.5	7.4	2.2	0.2	0.2	*	*	*	*	*	*	*
1956	0.5	7.2	2.1	0.3	*	*	*	*	*	*	*	*
1955	0.6	7.8	2.4	0.4	0.3	*	*	0.2	*	*	*	*
1954	0.7	9.3	2.7	0.5	0.4	*	*	*	*	*	*	*
1953	0.9	13.5	3.7	0.7	0.4	*	*	0.3	0.4	*	*	*
1952	0.9	12.7	4.2	0.7	0.5	*	*	*	*	*	*	*
1951	0.8	10.6	2.9	0.5	0.5	*	*	0.3	*	*	*	*
1950	0.7	8.5	3.2	0.5	0.3	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	0.3	4.2	1.2	0.2	*	*	*	*	*	*	*	*
1958	0.4	5.8	1.7	0.2	*	*	*	*	*	*	*	*
1957	0.4	5.3	1.5	0.3	*	*	*	*	*	*	*	*
1956	0.4	5.7	1.4	0.2	*	*	*	*	*	*	*	*
1955	0.4	5.8	2.0	0.3	*	*	*	*	*	*	*	*
1954	0.5	7.3	2.2	0.4	*	*	*	*	*	*	*	*
1953	0.6	9.0	2.7	0.5	0.3	*	*	0.2	*	*	*	*
1952	0.8	11.0	3.2	0.8	0.3	*	*	*	*	*	*	*
1951	0.6	7.1	2.8	0.6	0.3	*	*	*	*	*	*	*
1950	0.5	7.4	2.3	0.5	0.2	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MENINGOCOCCAL INFECTIONS (057)												
ALL OTHER, BOTH SEXES												
1959	0.6	4.8	1.7	0.5	*	*	*	*	*	*	*	*
1958	0.7	6.9	1.9	*	*	*	*	*	*	*	*	*
1957	0.6	4.5	1.7	0.5	*	*	*	*	*	*	*	*
1956	0.7	7.3	1.4	*	*	*	*	*	*	*	*	*
1955	0.9	6.9	2.7	*	*	*	*	*	*	*	*	*
1954	0.9	8.2	2.2	1.0	*	*	*	*	*	*	*	*
1953	1.3	10.2	3.3	1.4	1.0	*	*	*	*	*	*	*
1952	1.2	9.8	3.6	1.2	*	*	*	*	*	*	*	*
1951	1.1	11.1	2.8	1.0	0.8	*	*	*	*	*	*	*
1950	0.8	8.4	2.0	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	0.7	*	1.9	*	*	*	*	*	*	*	*	*
1958	0.7	9.0	1.8	*	*	*	*	*	*	*	*	*
1957	0.7	*	*	*	*	*	*	*	*	*	*	*
1956	0.8	9.2	*	*	*	*	*	*	*	*	*	*
1955	0.9	8.0	2.9	*	*	*	*	*	*	*	*	*
1954	1.1	8.5	2.4	1.1	*	*	*	*	*	*	*	*
1953	1.4	10.7	2.8	1.7	*	*	*	*	*	*	*	*
1952	1.3	*	4.0	1.3	*	*	*	*	*	*	*	*
1951	1.4	11.8	3.0	*	*	*	*	*	*	*	*	*
1950	1.1	9.9	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	0.6	*	*	*	*	*	*	*	*	*	*	*
1958	0.6	*	1.9	*	*	*	*	*	*	*	*	*
1957	0.6	*	1.9	*	*	*	*	*	*	*	*	*
1956	0.6	*	*	*	*	*	*	*	*	*	*	*
1955	0.8	*	2.4	*	*	*	*	*	*	*	*	*
1954	0.7	8.0	*	*	*	*	*	*	*	*	*	*
1953	1.1	9.8	3.7	*	*	*	*	*	*	*	*	*
1952	1.2	11.1	3.1	*	*	*	*	*	*	*	*	*
1951	0.9	10.4	2.6	*	*	*	*	*	*	*	*	*
1950	0.6	*	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ACUTE POLIOMYELITIS (080)												
BOTH SEXES												
1959	0.3	*	0.4	0.3	0.4	0.5	0.2	*	*	*	*	*
1958	0.1	*	0.3	0.1	0.2	0.3	0.1	*	*	*	*	*
1957	0.1	*	0.3	0.1	0.2	0.2	*	*	*	*	*	*
1956	0.3	0.8	0.7	0.4	0.6	0.6	0.2	*	*	*	*	*
1955	0.6	1.0	0.8	0.9	0.9	1.1	0.4	0.2	*	*	*	*
1954	0.8	1.9	1.3	1.4	1.1	1.4	0.4	0.2	*	*	*	*
1953	0.9	1.5	1.4	1.6	1.2	1.4	0.6	0.1	*	*	*	*
1952	2.0	3.4	3.4	3.7	2.8	3.0	0.9	0.2	0.2	*	*	*
1951	1.0	2.0	1.9	2.0	1.4	1.2	0.4	0.1	*	*	*	*
1950	1.3	2.6	2.0	2.5	1.8	1.7	0.6	0.2	*	*	*	*
MALE												
1959	0.4	*	0.6	0.4	0.5	0.7	0.2	*	*	*	*	*
1958	0.2	*	0.3	0.2	0.3	0.4	*	*	*	*	*	*
1957	0.2	*	0.3	0.2	0.2	0.2	*	*	*	*	*	*
1956	0.4	*	0.8	0.4	0.5	0.7	0.2	*	*	*	*	*
1955	0.7	1.3	0.8	1.0	0.9	1.3	0.6	*	*	*	*	*
1954	1.0	2.0	1.5	1.5	1.3	1.7	0.6	0.2	*	*	*	*
1953	1.1	1.7	1.5	1.9	1.5	1.6	0.7	0.2	*	*	*	*
1952	2.4	3.8	3.9	4.3	3.4	3.6	1.2	0.3	*	*	*	*
1951	1.2	2.1	2.2	2.5	1.7	1.4	0.5	*	*	*	*	*
1950	1.5	3.1	2.2	2.9	2.1	2.0	0.7	0.3	*	*	*	*
FEMALE												
1959	0.2	*	0.3	0.2	0.3	0.2	*	*	*	*	*	*
1958	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1957	0.1	*	*	0.1	*	*	*	*	*	*	*	*
1956	0.3	*	0.5	0.3	0.7	0.5	*	*	*	*	*	*
1955	0.6	*	0.8	0.9	0.9	0.9	0.3	*	*	*	*	*
1954	0.7	1.7	1.1	1.2	0.9	1.1	0.3	*	*	*	*	*
1953	0.7	1.2	1.3	1.2	0.9	1.3	0.5	*	*	*	*	*
1952	1.6	3.0	2.8	3.1	2.4	2.4	0.7	*	*	*	*	*
1951	0.8	2.0	1.7	1.5	1.2	1.0	0.2	*	*	*	*	*
1950	1.0	2.0	1.7	2.0	1.4	1.4	0.5	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ACUTE POLIOMYELITIS (080)												
WHITE, BOTH SEXES												
1959	0.2	*	0.3	0.3	0.3	0.5	0.2	*	*	*	*	*
1958	0.1	*	0.2	0.1	0.2	0.3	*	*	*	*	*	*
1957	0.1	*	0.2	0.1	0.2	0.2	*	*	*	*	*	*
1956	0.3	0.6	0.5	0.3	0.6	0.6	0.2	*	*	*	*	*
1955	0.7	1.1	0.9	1.0	1.0	1.2	0.5	0.2	*	*	*	*
1954	0.9	1.9	1.4	1.5	1.2	1.5	0.5	0.2	*	*	*	*
1953	1.0	1.3	1.4	1.7	1.3	1.6	0.6	0.2	*	*	*	*
1952	2.2	3.6	3.5	4.0	3.1	3.3	1.0	0.3	0.2	*	*	*
1951	1.1	1.9	1.9	2.1	1.6	1.4	0.4	*	*	*	*	*
1950	1.3	2.7	2.1	2.7	1.9	1.8	0.6	0.2	*	*	*	*
WHITE, MALE												
1959	0.3	*	0.4	0.4	0.4	0.8	0.2	*	*	*	*	*
1958	0.2	*	*	0.2	0.3	0.4	*	*	*	*	*	*
1957	0.1	*	*	*	0.2	0.2	*	*	*	*	*	*
1956	0.4	*	0.7	0.3	0.5	0.8	0.2	*	*	*	*	*
1955	0.8	1.2	0.8	1.1	1.0	1.4	0.6	*	*	*	*	*
1954	1.1	1.9	1.6	1.6	1.4	1.8	0.6	0.3	*	*	*	*
1953	1.2	1.4	1.4	2.1	1.6	1.7	0.7	0.2	*	*	*	*
1952	2.6	3.7	4.1	4.7	3.7	3.9	1.3	0.3	*	*	*	*
1951	1.3	1.9	2.2	2.7	1.8	1.6	0.6	*	*	*	*	*
1950	1.6	3.2	2.3	3.3	2.3	2.1	0.8	0.3	*	*	*	*
WHITE, FEMALE												
1959	0.1	*	*	0.2	0.2	0.3	*	*	*	*	*	*
1958	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1957	0.1	*	*	*	*	*	*	*	*	*	*	*
1956	0.3	*	0.4	0.3	0.7	0.5	*	*	*	*	*	*
1955	0.6	*	0.9	0.9	1.0	1.0	0.3	*	*	*	*	*
1954	0.7	1.8	1.1	1.3	1.0	1.2	0.3	*	*	*	*	*
1953	0.8	*	1.3	1.3	1.0	1.4	0.5	*	*	*	*	*
1952	1.8	3.5	2.9	3.4	2.6	2.6	0.8	*	*	*	*	*
1951	0.8	2.0	1.6	1.6	1.4	1.1	0.2	*	*	*	*	*
1950	1.1	2.1	1.8	2.1	1.5	1.5	0.5	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ACUTE POLIOMYELITIS (080)												
ALL OTHER, BOTH SEXES												
1959	0.4	*	1.1	0.4	*	*	*	*	*	*	*	*
1958	0.2	*	*	*	*	*	*	*	*	*	*	*
1957	0.2	*	*	*	*	*	*	*	*	*	*	*
1956	0.5	*	1.3	0.6	*	*	*	*	*	*	*	*
1955	0.2	*	*	*	*	*	*	*	*	*	*	*
1954	0.4	*	*	0.6	*	*	*	*	*	*	*	*
1953	0.5	*	1.6	0.6	*	*	*	*	*	*	*	*
1952	0.8	*	2.5	1.1	0.9	*	*	*	*	*	*	*
1951	0.6	*	2.1	1.0	*	*	*	*	*	*	*	*
1950	0.6	*	1.6	0.8	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	0.6	*	*	*	*	*	*	*	*	*	*	*
1958	0.3	*	*	*	*	*	*	*	*	*	*	*
1957	0.2	*	*	*	*	*	*	*	*	*	*	*
1956	0.5	*	*	*	*	*	*	*	*	*	*	*
1955	0.3	*	*	*	*	*	*	*	*	*	*	*
1954	0.5	*	*	*	*	*	*	*	*	*	*	*
1953	0.6	*	*	*	*	*	*	*	*	*	*	*
1952	1.0	*	2.9	1.5	*	*	*	*	*	*	*	*
1951	0.8	*	2.3	*	*	*	*	*	*	*	*	*
1950	0.6	*	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	0.2	*	*	*	*	*	*	*	*	*	*	*
1958	*	*	*	*	*	*	*	*	*	*	*	*
1957	*	*	*	*	*	*	*	*	*	*	*	*
1956	0.4	*	*	*	*	*	*	*	*	*	*	*
1955	0.2	*	*	*	*	*	*	*	*	*	*	*
1954	0.3	*	*	*	*	*	*	*	*	*	*	*
1953	0.4	*	*	*	*	*	*	*	*	*	*	*
1952	0.6	*	*	*	*	*	*	*	*	*	*	*
1951	0.5	*	*	*	*	*	*	*	*	*	*	*
1950	0.6	*	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MEASLES (085)												
BOTH SEXES												
1959	0.2	1.8	1.1	0.3	*	*	*	*	*	*	*	*
1958	0.3	3.3	1.8	0.3	*	*	*	*	*	*	*	*
1957	0.2	1.9	1.1	0.3	*	*	*	*	*	*	*	*
1956	0.3	3.2	1.6	0.4	*	*	*	*	*	*	*	*
1955	0.2	2.2	1.0	0.3	*	*	*	*	*	*	*	*
1954	0.3	3.0	1.7	0.4	*	*	*	*	*	*	*	*
1953	0.3	3.5	1.4	0.3	*	*	*	*	*	*	*	*
1952	0.4	4.2	1.9	0.6	0.1	*	*	*	*	*	*	*
1951	0.4	4.9	2.2	0.6	0.1	*	*	*	*	*	*	*
1950	0.3	3.6	1.4	0.4	0.1	*	*	*	*	*	*	*
MALE												
1959	0.2	1.5	1.1	0.3	*	*	*	*	*	*	*	*
1958	0.3	3.5	1.8	0.3	*	*	*	*	*	*	*	*
1957	0.2	1.9	1.1	0.4	*	*	*	*	*	*	*	*
1956	0.3	2.9	1.8	0.3	*	*	*	*	*	*	*	*
1955	0.2	2.4	1.1	0.2	*	*	*	*	*	*	*	*
1954	0.3	2.7	1.6	0.4	*	*	*	*	*	*	*	*
1953	0.3	3.6	1.3	0.3	*	*	*	*	*	*	*	*
1952	0.4	4.5	1.7	0.5	*	*	*	*	*	*	*	*
1951	0.5	5.8	2.1	0.6	*	*	*	*	*	*	*	*
1950	0.3	3.3	1.5	0.4	0.2	*	*	*	*	*	*	*
FEMALE												
1959	0.2	2.0	1.1	0.3	*	*	*	*	*	*	*	*
1958	0.3	3.1	1.8	0.3	*	*	*	*	*	*	*	*
1957	0.2	1.9	1.2	0.3	*	*	*	*	*	*	*	*
1956	0.3	3.6	1.5	0.5	*	*	*	*	*	*	*	*
1955	0.2	1.9	1.0	0.3	*	*	*	*	*	*	*	*
1954	0.3	3.3	1.7	0.5	*	*	*	*	*	*	*	*
1953	0.3	3.4	1.6	0.4	*	*	*	*	*	*	*	*
1952	0.4	3.8	2.1	0.6	*	*	*	*	*	*	*	*
1951	0.4	3.9	2.3	0.6	*	*	*	*	*	*	*	*
1950	0.3	3.9	1.3	0.4	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MEASLES (085)												
WHITE, BOTH SEXES												
1959	0.2	1.5	0.9	0.3	*	*	*	*	*	*	*	*
1958	0.3	2.3	1.6	0.3	*	*	*	*	*	*	*	*
1957	0.2	1.4	1.0	0.3	*	*	*	*	*	*	*	*
1956	0.3	3.0	1.3	0.4	*	*	*	*	*	*	*	*
1955	0.2	1.6	0.9	0.3	*	*	*	*	*	*	*	*
1954	0.3	2.1	1.5	0.4	*	*	*	*	*	*	*	*
1953	0.3	3.1	1.4	0.4	*	*	*	*	*	*	*	*
1952	0.3	3.4	1.5	0.5	*	*	*	*	*	*	*	*
1951	0.4	4.1	2.0	0.6	*	*	*	*	*	*	*	*
1950	0.3	3.1	1.2	0.4	0.1	*	*	*	*	*	*	*
WHITE, MALE												
1959	0.2	1.2	1.0	0.2	*	*	*	*	*	*	*	*
1958	0.3	2.4	1.7	0.3	*	*	*	*	*	*	*	*
1957	0.2	1.5	0.9	0.4	*	*	*	*	*	*	*	*
1956	0.3	2.7	1.5	0.3	*	*	*	*	*	*	*	*
1955	0.2	1.8	0.9	0.2	*	*	*	*	*	*	*	*
1954	0.3	2.1	1.4	0.4	*	*	*	*	*	*	*	*
1953	0.3	3.4	1.3	0.3	*	*	*	*	*	*	*	*
1952	0.3	4.0	1.4	0.5	*	*	*	*	*	*	*	*
1951	0.4	5.0	2.0	0.6	*	*	*	*	*	*	*	*
1950	0.3	3.0	1.3	0.4	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	0.2	1.8	0.9	0.3	*	*	*	*	*	*	*	*
1958	0.3	2.2	1.6	0.3	*	*	*	*	*	*	*	*
1957	0.2	1.3	1.1	0.3	*	*	*	*	*	*	*	*
1956	0.3	3.3	1.2	0.5	*	*	*	*	*	*	*	*
1955	0.2	1.3	0.8	0.3	*	*	*	*	*	*	*	*
1954	0.3	2.2	1.6	0.5	*	*	*	*	*	*	*	*
1953	0.3	2.9	1.4	0.4	*	*	*	*	*	*	*	*
1952	0.3	2.8	1.7	0.6	*	*	*	*	*	*	*	*
1951	0.4	3.1	2.0	0.5	*	*	*	*	*	*	*	*
1950	0.3	3.2	1.2	0.4	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MEASLES (085)												
ALL OTHER, BOTH SEXES												
1959	0.5	3.3	2.3	0.4	*	*	*	*	*	*	*	*
1958	0.7	8.8	2.8	*	*	*	*	*	*	*	*	*
1957	0.5	5.0	2.0	*	*	*	*	*	*	*	*	*
1956	0.6	4.4	3.6	*	*	*	*	*	*	*	*	*
1955	0.5	5.7	2.2	*	*	*	*	*	*	*	*	*
1954	0.7	8.4	2.8	*	*	*	*	*	*	*	*	*
1953	0.5	5.8	2.0	*	*	*	*	*	*	*	*	*
1952	0.9	8.9	4.3	0.8	*	*	*	*	*	*	*	*
1951	0.9	10.2	3.4	0.8	*	*	*	*	*	*	*	*
1950	0.6	6.9	2.6	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	0.4	*	2.2	*	*	*	*	*	*	*	*	*
1958	0.7	9.7	2.9	*	*	*	*	*	*	*	*	*
1957	0.5	*	2.3	*	*	*	*	*	*	*	*	*
1956	0.7	*	3.7	*	*	*	*	*	*	*	*	*
1955	0.5	*	2.1	*	*	*	*	*	*	*	*	*
1954	0.7	*	3.1	*	*	*	*	*	*	*	*	*
1953	0.4	*	*	*	*	*	*	*	*	*	*	*
1952	0.8	*	3.5	*	*	*	*	*	*	*	*	*
1951	0.8	11.4	2.9	*	*	*	*	*	*	*	*	*
1950	0.5	*	2.8	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	0.5	*	2.3	*	*	*	*	*	*	*	*	*
1958	0.6	7.9	2.6	*	*	*	*	*	*	*	*	*
1957	0.4	*	*	*	*	*	*	*	*	*	*	*
1956	0.6	*	3.5	*	*	*	*	*	*	*	*	*
1955	0.5	*	2.3	*	*	*	*	*	*	*	*	*
1954	0.7	10.0	2.6	*	*	*	*	*	*	*	*	*
1953	0.5	*	2.4	*	*	*	*	*	*	*	*	*
1952	1.0	10.2	5.1	*	*	*	*	*	*	*	*	*
1951	0.9	9.0	4.0	*	*	*	*	*	*	*	*	*
1950	0.6	*	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER INFECTIVE AND PARASITIC DISEASES												
BOTH SEXES												
1959	3.1	21.7	3.0	1.2	1.0	1.5	1.7	2.9	4.6	7.0	11.3	20.6
1958	3.0	20.9	3.0	1.0	1.1	1.3	1.8	2.8	4.4	6.6	11.4	20.6
1957	2.8	20.4	2.6	1.0	1.0	1.3	1.5	2.9	3.9	5.7	9.9	15.4
1956	2.6	19.1	2.8	0.9	0.9	1.3	1.6	2.4	3.7	5.5	9.5	17.9
1955	2.5	17.3	2.4	1.0	1.0	1.1	1.6	2.4	3.5	5.4	8.7	19.7
1954	2.6	18.7	2.7	1.0	1.1	1.3	1.8	2.6	3.4	4.9	8.1	17.4
1953	2.6	18.2	3.0	1.3	1.0	1.4	1.7	2.4	3.6	5.3	8.5	16.8
1952	2.8	20.0	3.3	1.4	1.1	1.3	1.9	2.7	3.7	4.7	9.1	18.3
1951	2.7	18.4	3.1	1.3	1.0	1.3	1.8	2.7	3.6	5.2	8.1	18.2
1950	2.5	17.3	3.0	1.1	1.0	1.3	1.8	2.6	3.3	4.8	8.2	16.5
MALE												
1959	3.4	24.1	2.9	1.2	1.0	1.4	1.8	3.1	5.5	8.8	13.5	19.4
1958	3.3	23.8	3.1	1.0	1.0	1.1	1.8	3.2	5.5	8.1	13.7	20.9
1957	3.1	23.0	2.7	1.1	1.2	1.1	1.7	3.2	4.8	7.3	12.1	15.7
1956	2.9	21.6	2.9	1.0	0.9	1.4	1.8	2.7	4.8	6.5	11.2	17.4
1955	2.8	19.5	2.4	1.1	1.1	1.1	1.8	2.8	4.2	6.4	9.8	17.3
1954	2.9	20.6	3.1	1.1	1.2	1.3	1.9	3.1	4.0	5.7	8.2	18.8
1953	3.1	20.9	3.3	1.4	1.1	1.4	1.9	2.9	4.6	6.6	10.4	17.0
1952	3.1	21.5	3.6	1.5	1.1	1.1	1.8	3.1	4.6	5.7	11.0	21.2
1951	2.9	20.0	3.4	1.3	1.0	1.1	2.0	3.0	4.2	6.2	9.8	18.4
1950	2.8	18.8	3.4	1.2	1.1	1.2	1.9	3.2	4.3	5.9	9.7	16.9
FEMALE												
1959	2.8	19.3	3.1	1.1	1.0	1.5	1.6	2.7	3.8	5.4	9.7	21.4
1958	2.8	18.0	2.8	1.0	1.2	1.6	1.8	2.5	3.4	5.2	9.6	20.3
1957	2.4	17.8	2.6	0.9	0.9	1.5	1.3	2.6	3.1	4.3	8.0	15.2
1956	2.4	16.5	2.7	0.8	1.0	1.2	1.5	2.2	2.8	4.7	8.1	18.3
1955	2.3	15.1	2.4	1.0	0.9	1.2	1.5	2.1	2.9	4.5	7.8	21.4
1954	2.3	16.7	2.4	0.8	1.0	1.4	1.8	2.2	2.7	4.2	8.0	16.5
1953	2.3	15.4	2.7	1.1	1.0	1.3	1.5	2.0	2.6	4.1	7.0	16.7
1952	2.5	18.5	3.1	1.3	1.1	1.4	1.9	2.3	2.7	3.9	7.6	16.3
1951	2.4	16.8	2.8	1.2	1.0	1.4	1.5	2.3	3.0	4.4	6.6	18.1
1950	2.2	15.8	2.6	0.9	0.9	1.3	1.7	2.1	2.3	3.8	6.9	16.2

* Figure does not meet standard of reliability or precision

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and ICD Nos. 030-039, 041, 042, 044, 049, 052-054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER INFECTIVE AND PARASITIC DISEASES												
WHITE, BOTH SEXES												
1959	2.7	16.1	2.5	1.1	0.8	1.2	1.3	2.4	4.2	6.8	11.2	20.1
1958	2.7	16.4	2.7	0.9	0.9	1.2	1.5	2.4	4.1	6.4	11.3	21.3
1957	2.5	16.8	2.3	1.0	0.9	1.0	1.3	2.5	3.5	5.6	9.9	15.8
1956	2.3	15.2	2.5	0.8	0.8	1.0	1.3	2.2	3.3	5.4	9.6	18.6
1955	2.2	13.8	2.1	0.9	0.8	0.9	1.3	2.1	3.3	5.2	8.8	20.2
1954	2.3	15.3	2.5	0.9	0.8	1.0	1.5	2.3	2.9	4.7	8.1	17.9
1953	2.3	14.7	2.7	1.1	0.8	1.0	1.4	2.1	3.2	5.2	8.6	16.9
1952	2.5	16.5	2.9	1.2	0.9	1.1	1.6	2.3	3.3	4.6	9.1	19.2
1951	2.3	14.7	2.7	1.1	0.8	1.0	1.4	2.3	3.2	5.2	8.2	19.3
1950	2.2	14.6	2.7	0.9	0.8	1.0	1.5	2.2	3.0	4.5	7.9	17.3
WHITE, MALE												
1959	3.0	18.0	2.5	1.1	0.8	1.2	1.4	2.6	5.0	8.6	13.2	18.4
1958	3.0	18.9	2.9	0.9	0.8	1.0	1.5	2.8	5.1	7.8	13.5	21.5
1957	2.8	18.9	2.4	1.1	1.1	0.8	1.5	2.8	4.3	7.3	12.2	15.8
1956	2.6	17.8	2.5	0.8	0.7	1.2	1.4	2.4	4.1	6.3	11.2	17.9
1955	2.5	16.1	2.1	1.0	0.8	0.9	1.5	2.4	3.8	6.1	10.0	18.0
1954	2.5	17.0	2.9	1.0	0.9	1.0	1.6	2.8	3.4	5.4	8.1	19.1
1953	2.7	16.7	2.9	1.2	0.9	1.1	1.6	2.5	4.1	6.4	10.6	16.8
1952	2.7	17.9	3.1	1.3	0.9	0.9	1.7	2.6	4.1	5.6	10.8	21.3
1951	2.6	16.1	2.9	1.2	0.8	0.9	1.6	2.6	3.7	6.1	9.9	19.4
1950	2.5	15.7	3.1	1.1	0.8	0.9	1.7	2.8	4.0	5.6	9.2	18.3
WHITE, FEMALE												
1959	2.5	14.2	2.5	1.0	0.8	1.1	1.2	2.2	3.5	5.3	9.7	21.2
1958	2.5	13.7	2.5	1.0	1.0	1.3	1.6	2.1	3.2	5.1	9.5	21.1
1957	2.2	14.7	2.1	0.9	0.7	1.2	1.1	2.2	2.7	4.1	8.2	15.8
1956	2.1	12.4	2.5	0.7	0.8	0.9	1.2	1.9	2.6	4.6	8.3	19.1
1955	2.0	11.3	2.1	0.9	0.8	0.9	1.2	1.8	2.7	4.3	7.9	21.7
1954	2.1	13.6	2.1	0.8	0.8	1.1	1.4	1.9	2.5	4.1	8.0	17.0
1953	2.0	12.5	2.4	1.0	0.8	0.9	1.2	1.6	2.3	4.0	7.0	17.0
1952	2.2	15.1	2.6	1.1	0.9	1.2	1.5	1.9	2.4	3.7	7.7	17.6
1951	2.1	13.2	2.5	1.1	0.8	1.2	1.2	2.0	2.7	4.3	6.7	19.3
1950	1.8	13.4	2.3	0.8	0.7	1.0	1.3	1.7	2.1	3.4	6.8	16.6

* Figure does not meet standard of reliability or precision

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and ICD Nos. 030-039, 041, 042, 044, 049, 052-054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER INFECTIVE AND PARASITIC DISEASES												
ALL OTHER, BOTH SEXES												
1959	6.1	54.1	5.8	1.7	2.3	3.6	4.8	7.3	8.9	8.9	12.8	*
1958	5.3	48.4	4.6	1.4	2.0	2.8	4.2	6.5	7.5	9.0	13.0	*
1957	5.0	41.9	5.1	1.1	2.0	3.3	3.6	6.7	8.7	7.2	8.6	*
1956	5.1	42.7	5.0	1.8	2.2	3.5	4.1	5.3	8.5	7.1	7.8	*
1955	4.8	39.2	4.7	1.9	2.1	3.0	4.3	5.4	6.4	8.1	*	*
1954	5.2	40.0	4.2	1.7	3.0	3.8	4.7	5.6	8.2	7.5	8.2	*
1953	5.3	41.4	5.2	2.3	2.6	4.1	3.9	6.1	8.2	6.7	*	*
1952	5.3	43.2	6.6	2.4	2.4	2.8	4.1	6.7	8.7	6.2	9.3	*
1951	5.4	42.5	6.0	2.2	2.6	3.2	5.1	6.4	8.7	5.8	*	*
1950	5.2	35.7	5.3	2.0	2.8	3.8	4.4	6.5	6.6	9.3	12.3	*
ALL OTHER, MALE												
1959	6.7	60.0	5.1	1.9	2.8	3.1	5.0	7.8	11.0	11.6	16.7	*
1958	5.7	53.6	4.5	1.7	1.9	2.2	4.2	6.4	9.3	11.6	15.1	*
1957	5.6	47.7	5.1	1.4	2.2	3.1	3.8	7.3	10.1	8.4	*	*
1956	5.5	44.9	5.4	1.8	1.7	3.3	4.6	4.8	11.9	8.1	*	*
1955	5.2	41.0	4.9	1.8	2.8	2.8	4.1	6.3	7.8	9.9	*	*
1954	6.0	44.0	4.3	2.1	3.6	4.0	4.9	6.6	11.3	9.4	*	*
1953	6.0	48.7	5.6	2.9	3.1	4.0	3.9	6.6	9.4	8.4	*	*
1952	5.7	45.9	7.0	2.5	2.3	3.0	2.9	7.4	10.9	6.9	*	*
1951	6.0	45.9	7.0	2.3	2.7	3.1	5.6	7.6	10.7	6.4	*	*
1950	5.5	40.0	5.7	2.0	3.3	3.7	3.7	7.3	8.3	10.1	*	*
ALL OTHER, FEMALE												
1959	5.6	48.0	6.5	1.6	1.9	4.1	4.7	6.7	6.9	6.4	*	*
1958	4.9	43.1	4.8	1.1	2.0	3.3	4.2	6.6	5.8	6.6	*	*
1957	4.5	36.1	5.1	*	1.8	3.5	3.4	6.0	7.3	6.0	*	*
1956	4.7	40.5	4.5	1.7	2.7	3.8	3.6	5.6	5.1	6.2	*	*
1955	4.5	37.6	4.5	2.0	1.5	3.2	4.4	4.4	5.0	6.4	*	*
1954	4.4	36.0	4.0	1.2	2.4	3.6	4.5	4.6	5.0	5.8	*	*
1953	4.6	34.0	4.8	1.6	2.2	4.1	3.9	5.7	6.9	*	*	*
1952	5.0	40.4	6.1	2.3	2.4	2.7	5.1	6.0	6.5	5.6	*	*
1951	4.8	39.2	4.9	2.1	2.6	3.3	4.6	5.2	6.6	*	*	*
1950	4.8	31.4	5.0	2.0	2.4	4.0	5.1	5.7	4.8	8.4	*	*

* Figure does not meet standard of reliability or precision

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and ICD Nos. 030-039, 041, 042, 044, 049, 052-.054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
MALIGNANT NEOPLASMS (ICD NOS. 140-205)												
BOTH SEXES												
1959	147.3	6.8	10.6	7.1	8.5	19.9	59.4	174.3	394.4	706.2	1,101.8	1,447.3
1958	146.8	6.5	10.9	6.8	8.8	19.4	60.2	173.5	392.4	707.6	1,112.8	1,443.1
1957	148.6	7.0	11.0	6.8	8.7	20.1	58.9	176.1	399.3	717.3	1,140.2	1,489.7
1956	147.8	7.9	11.1	7.2	8.6	20.1	60.8	173.9	399.5	706.3	1,161.5	1,483.7
1955	146.5	7.7	11.1	7.0	9.2	19.3	61.3	174.6	392.3	711.5	1,146.4	1,465.3
1954	145.6	8.4	11.3	6.7	9.1	19.7	61.5	176.3	390.3	708.3	1,151.8	1,447.7
1953	144.8	8.5	10.9	6.6	8.9	19.5	62.1	176.7	396.1	699.8	1,154.5	1,456.0
1952	143.4	8.5	11.6	6.5	9.4	20.3	61.4	175.8	394.3	697.4	1,152.9	1,455.4
1951	140.6	8.2	11.8	6.5	9.3	19.7	61.7	175.9	388.7	686.5	1,133.4	1,452.4
1950	139.8	8.7	11.7	6.7	8.6	20.0	62.7	175.1	390.7	698.8	1,153.3	1,451.0
MALE												
1959	159.9	7.2	11.4	8.2	10.4	18.4	48.5	168.6	453.1	872.8	1,356.4	1,762.8
1958	157.8	6.9	11.7	7.4	10.7	18.2	48.2	165.7	446.4	862.5	1,358.6	1,704.9
1957	160.0	7.0	12.3	7.7	10.5	18.1	46.6	165.7	453.6	879.3	1,384.5	1,753.4
1956	158.5	8.4	11.5	7.8	10.4	18.1	47.5	163.6	451.9	849.3	1,416.6	1,767.9
1955	155.8	8.4	12.2	8.3	11.2	17.4	45.9	163.8	437.2	851.0	1,373.9	1,694.0
1954	154.3	9.4	12.0	7.4	11.0	18.1	47.1	162.3	433.0	839.4	1,371.6	1,688.8
1953	152.4	8.6	11.7	7.6	10.8	17.4	48.0	162.9	437.2	815.0	1,358.9	1,677.9
1952	149.5	8.3	12.2	7.1	11.2	17.6	45.9	158.9	428.3	809.1	1,349.2	1,665.6
1951	145.1	8.7	13.1	7.1	11.4	17.4	44.9	158.3	415.8	788.5	1,307.3	1,689.1
1950	142.9	9.7	12.5	7.4	9.7	17.7	45.6	156.2	413.1	791.5	1,332.6	1,668.3
FEMALE												
1959	135.1	6.4	9.8	5.9	6.6	21.3	69.9	179.7	339.4	560.5	903.2	1,241.1
1958	136.1	6.0	10.1	6.2	7.0	20.7	71.7	181.0	341.3	571.2	919.4	1,268.6
1957	137.5	7.0	9.8	5.8	7.0	22.1	70.6	186.2	347.7	573.2	945.9	1,309.7
1956	137.5	7.3	10.6	6.4	7.0	21.9	73.6	184.1	349.4	578.2	957.3	1,286.0
1955	137.4	6.9	10.0	5.7	7.3	21.0	76.1	185.1	349.1	585.7	962.7	1,303.3
1954	137.2	7.4	10.5	6.0	7.2	21.2	75.3	190.2	348.9	589.4	972.5	1,275.3
1953	137.4	8.4	10.0	5.6	7.2	21.5	75.7	190.2	356.0	594.6	986.3	1,296.5
1952	137.5	8.7	10.8	5.8	7.7	22.8	76.5	192.6	360.8	594.8	990.2	1,304.7
1951	136.1	7.6	10.4	5.8	7.4	21.9	77.9	193.4	361.8	592.1	987.0	1,284.2
1950	136.8	7.6	10.8	6.0	7.6	22.2	79.3	194.0	368.2	612.3	1,000.7	1,299.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
MALIGNANT NEOPLASMS (ICD NOS. 140-205)												
WHITE, BOTH SEXES												
1959	150.9	6.9	11.3	7.4	8.4	19.2	56.6	168.1	385.6	706.9	1,121.2	1,488.9
1958	150.4	6.8	11.5	7.0	8.9	18.6	57.5	166.6	383.8	709.3	1,134.9	1,493.3
1957	152.3	7.5	11.8	7.0	8.8	19.7	56.4	168.8	390.2	719.7	1,163.5	1,541.1
1956	151.5	7.9	11.7	7.4	8.7	19.5	57.7	166.8	390.5	711.7	1,183.5	1,535.7
1955	150.4	8.0	11.8	7.3	9.3	18.7	58.2	167.2	385.4	718.6	1,173.1	1,519.0
1954	149.3	8.7	12.0	7.0	8.9	19.1	58.4	168.7	382.5	716.3	1,177.6	1,500.1
1953	148.7	8.5	11.5	6.9	9.0	18.9	59.4	169.3	390.5	707.9	1,183.1	1,511.6
1952	147.3	8.4	12.2	6.6	9.5	19.8	58.4	168.4	388.8	707.2	1,180.5	1,512.1
1951	144.5	8.6	12.3	6.8	9.4	19.1	58.8	169.0	382.7	697.0	1,163.9	1,510.5
1950	143.5	8.7	12.2	7.0	8.7	19.3	59.7	168.3	386.0	704.2	1,182.6	1,505.9
WHITE, MALE												
1959	163.4	7.2	12.1	8.6	10.4	18.2	46.4	162.4	443.5	870.7	1,379.1	1,808.6
1958	161.6	7.4	12.5	7.6	10.9	17.8	46.5	159.4	438.3	862.5	1,387.5	1,769.0
1957	163.8	7.3	12.9	8.0	10.7	18.2	45.1	158.6	443.2	881.1	1,416.1	1,806.8
1956	162.4	8.5	12.1	8.1	10.5	18.0	46.2	157.6	442.0	855.3	1,445.0	1,823.8
1955	160.0	8.7	13.1	8.6	11.3	17.5	44.3	157.8	429.1	858.8	1,409.1	1,746.9
1954	158.2	9.5	12.7	7.8	10.8	18.1	45.4	155.6	424.7	846.0	1,403.9	1,740.8
1953	156.7	8.5	12.4	7.9	10.9	17.6	46.9	156.8	431.0	822.4	1,398.0	1,735.1
1952	153.8	8.1	12.8	7.2	11.4	17.8	44.4	153.0	422.2	820.6	1,383.1	1,722.9
1951	149.5	9.2	13.8	7.6	11.5	17.3	44.2	152.8	409.5	800.2	1,346.4	1,748.9
1950	147.2	9.6	13.1	7.6	9.9	17.7	44.5	150.8	409.4	798.7	1,367.6	1,732.7
WHITE, FEMALE												
1959	138.7	6.5	10.4	6.2	6.4	20.3	66.3	173.7	331.2	563.9	922.0	1,282.1
1958	139.4	6.1	10.6	6.4	7.0	19.4	68.1	173.6	332.6	574.5	938.1	1,311.1
1957	141.0	7.6	10.6	6.1	7.0	21.2	67.2	178.7	339.8	576.7	964.5	1,360.4
1956	140.7	7.3	11.2	6.7	7.0	20.9	68.9	175.8	341.5	583.4	976.4	1,335.3
1955	141.0	7.2	10.6	6.0	7.4	19.9	71.8	176.5	343.6	592.3	985.1	1,357.6
1954	140.7	7.9	11.2	6.3	7.2	20.0	70.9	181.5	341.8	598.8	994.8	1,327.5
1953	140.9	8.6	10.6	5.9	7.2	20.2	71.5	181.7	351.1	603.5	1,008.6	1,350.1
1952	141.0	8.7	11.5	6.0	7.7	21.7	72.0	183.6	355.8	603.4	1,014.1	1,360.5
1951	139.5	8.0	10.8	6.1	7.4	20.8	73.0	185.0	356.1	601.9	1,012.4	1,340.4
1950	139.9	7.8	11.3	6.3	7.5	20.9	74.5	185.8	362.5	616.5	1,026.6	1,348.3

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
MALIGNANT NEOPLASMS (ICD NOS. 140-205)												
ALL OTHER, BOTH SEXES												
1959	118.8	6.7	6.6	4.8	8.9	24.8	84.5	232.7	487.1	698.8	834.1	939.7
1958	118.0	4.7	7.2	5.6	8.0	25.9	83.5	239.1	483.2	688.4	805.5	839.4
1957	119.0	4.0	6.5	5.1	8.1	23.5	80.6	245.8	499.9	688.3	815.4	848.4
1956	118.3	7.5	7.4	5.6	8.2	24.7	87.4	242.8	500.8	641.9	850.0	855.7
1955	114.0	6.1	6.5	5.0	8.3	23.7	87.6	244.6	472.0	627.5	767.6	808.6
1954	114.9	6.4	6.6	4.8	9.8	24.7	88.3	250.2	482.7	612.4	781.6	778.2
1953	111.4	8.1	6.6	4.4	8.5	24.2	85.9	247.0	464.7	602.9	735.5	773.1
1952	110.6	9.4	7.4	5.4	8.6	24.2	88.2	246.5	464.1	580.3	748.7	755.1
1951	107.5	5.2	7.9	4.0	8.6	24.9	86.7	242.2	466.0	559.7	681.6	748.9
1950	108.1	8.6	7.7	4.9	8.2	26.1	88.9	240.4	447.5	625.5	710.6	799.4
ALL OTHER, MALE												
1959	131.7	7.3	7.1	5.4	9.8	19.9	66.9	229.0	551.7	897.0	1,067.4	1,248.3
1958	126.4	*	7.0	5.8	8.8	20.9	63.2	227.1	532.3	862.0	988.5	1,018.5
1957	129.2	*	8.4	5.9	8.7	17.4	60.0	234.9	565.7	860.4	978.5	1,115.4
1956	125.4	7.7	7.8	6.1	9.8	19.2	59.6	221.8	562.4	778.4	1,048.5	1,092.0
1955	119.9	*	6.7	5.9	10.0	17.0	60.5	221.9	530.0	759.1	913.6	1,045.8
1954	121.5	8.9	7.1	5.1	12.6	17.8	62.2	226.6	530.6	761.1	942.0	1,022.7
1953	115.2	9.0	7.3	5.1	10.1	15.5	57.9	221.6	510.9	725.9	832.5	947.6
1952	112.5	9.9	8.2	6.1	9.5	16.1	59.1	214.4	503.6	676.4	889.1	940.0
1951	107.2	*	8.0	3.6	10.1	18.1	51.5	209.7	495.3	650.2	782.1	942.1
1950	106.1	10.4	8.2	5.7	7.7	17.9	56.0	207.4	457.1	697.8	844.4	916.0
ALL OTHER, FEMALE												
1959	106.7	*	6.1	4.3	8.1	29.1	100.5	236.3	425.1	519.2	625.5	710.3
1958	110.1	*	7.5	5.4	7.4	30.3	102.0	250.8	437.2	528.7	639.2	734.2
1957	109.3	*	4.6	4.3	7.5	28.7	99.4	256.3	435.8	531.1	663.4	683.8
1956	111.5	7.3	7.0	5.1	6.8	29.6	113.1	262.7	440.4	513.6	663.0	691.7
1955	108.4	*	6.3	4.2	6.8	29.5	112.6	266.7	414.4	506.4	623.5	641.2
1954	108.7	*	6.1	4.5	7.4	30.7	112.6	273.2	434.8	475.1	623.8	634.4
1953	107.9	*	6.0	3.6	7.2	31.9	111.9	271.9	417.1	486.7	638.0	654.8
1952	108.7	8.9	6.6	4.8	7.9	31.4	115.1	278.2	423.4	490.2	615.5	627.6
1951	107.8	*	7.9	4.4	7.3	31.0	119.3	274.1	435.5	474.3	585.6	617.9
1950	110.1	*	7.1	4.0	8.6	33.5	119.3	273.3	437.7	552.7	578.9	716.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)												
BOTH SEXES												
1959	3.4	*	*	*	0.1	0.2	1.1	5.1	10.0	15.6	25.0	34.6
1958	3.3	*	*	*	0.1	0.2	1.1	4.5	10.1	14.9	25.1	37.8
1957	3.4	*	*	*	0.1	0.2	1.0	4.7	9.5	16.5	27.4	40.6
1956	3.3	*	*	0.1	0.1	0.2	1.0	4.3	9.8	15.5	27.2	38.5
1955	3.4	*	*	*	0.1	0.2	1.1	4.2	9.5	16.4	26.8	40.0
1954	3.3	*	*	*	0.1	0.2	1.1	4.2	9.4	16.9	26.8	41.2
1953	3.2	*	*	*	*	0.2	1.0	3.9	9.3	15.8	27.9	42.0
1952	3.2	*	*	*	0.1	0.2	1.0	3.9	9.3	16.3	27.6	41.6
1951	3.2	*	*	*	0.1	0.2	1.0	3.8	9.3	16.3	28.2	39.1
1950	3.4	*	*	*	0.1	0.2	1.1	4.0	9.7	17.9	32.5	43.5
MALE												
1959	5.4	*	*	*	0.2	0.4	1.5	8.1	16.8	27.0	41.0	61.7
1958	5.2	*	*	*	*	0.3	1.5	7.0	16.9	25.4	41.2	64.2
1957	5.5	*	*	*	*	0.3	1.5	7.4	16.0	28.9	45.5	69.1
1956	5.3	*	*	*	*	0.3	1.4	6.9	16.5	26.5	46.3	67.3
1955	5.4	*	*	*	*	0.3	1.6	6.6	16.1	28.1	46.0	68.6
1954	5.3	*	*	*	*	0.3	1.5	6.6	15.8	28.5	44.9	70.4
1953	5.1	*	*	*	*	0.2	1.4	6.0	15.7	26.9	46.5	70.9
1952	5.1	*	*	*	0.2	0.3	1.3	6.2	16.0	26.5	45.9	67.0
1951	5.1	*	*	*	*	0.2	1.5	5.9	15.6	27.2	47.8	64.5
1950	5.4	*	*	*	*	0.2	1.6	6.3	16.2	30.2	54.8	75.2
FEMALE												
1959	1.5	*	*	*	*	*	0.7	2.1	3.6	5.6	12.5	16.9
1958	1.5	*	*	*	*	0.2	0.7	2.1	3.7	5.7	12.4	20.2
1957	1.4	*	*	*	*	*	0.5	2.0	3.3	5.4	13.0	21.1
1956	1.4	*	*	*	*	0.2	0.6	1.8	3.4	5.6	11.9	18.5
1955	1.4	*	*	*	*	0.2	0.7	1.9	3.1	5.9	11.3	19.8
1954	1.4	*	*	*	*	*	0.7	1.9	3.2	6.3	12.0	20.2
1953	1.4	*	*	*	*	*	0.5	1.8	3.1	5.6	12.6	21.1
1952	1.4	*	*	*	*	0.2	0.7	1.7	2.8	6.8	12.5	23.4
1951	1.4	*	*	*	*	*	0.6	1.8	3.1	6.3	11.7	21.1
1950	1.4	*	*	*	*	0.2	0.5	1.8	3.3	6.4	13.6	21.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)												
WHITE, BOTH SEXES												
1959	3.5	*	*	*	0.1	0.2	0.9	4.8	9.9	15.9	25.7	35.8
1958	3.4	*	*	*	*	0.2	1.0	4.4	10.0	15.2	25.7	39.5
1957	3.5	*	*	*	0.1	0.2	0.9	4.4	9.3	16.7	28.1	41.9
1956	3.4	*	*	*	*	0.2	0.9	4.1	9.6	15.7	28.2	40.5
1955	3.5	*	*	*	*	0.2	1.1	4.1	9.5	16.7	27.8	41.5
1954	3.5	*	*	*	*	0.2	0.9	4.1	9.2	17.4	27.5	43.6
1953	3.3	*	*	*	*	0.2	0.9	3.7	9.2	16.2	28.8	44.6
1952	3.3	*	*	*	*	0.2	0.9	3.7	9.3	16.6	28.3	44.3
1951	3.3	*	*	*	0.1	0.1	0.9	3.5	9.1	16.8	29.5	41.3
1950	3.5	*	*	*	0.1	0.2	1.0	3.8	9.5	18.3	33.8	44.9
WHITE, MALE												
1959	5.6	*	*	*	*	0.4	1.3	7.7	16.7	27.6	42.4	64.4
1958	5.3	*	*	*	*	0.2	1.4	6.8	16.7	26.0	42.2	67.1
1957	5.6	*	*	*	*	0.2	1.4	7.0	15.6	29.2	46.9	73.0
1956	5.5	*	*	*	*	0.3	1.3	6.6	16.3	27.1	48.5	70.9
1955	5.6	*	*	*	*	0.3	1.5	6.4	16.1	28.9	48.1	71.4
1954	5.5	*	*	*	*	0.3	1.3	6.4	15.5	29.5	46.5	74.8
1953	5.3	*	*	*	*	0.2	1.3	5.8	15.6	27.7	48.2	75.0
1952	5.3	*	*	*	*	0.3	1.2	6.0	15.9	27.4	47.4	71.5
1951	5.2	*	*	*	*	*	1.3	5.6	15.4	28.2	50.2	67.5
1950	5.7	*	*	*	*	0.2	1.5	6.1	15.9	31.1	57.4	78.4
WHITE, FEMALE												
1959	1.5	*	*	*	*	*	0.6	2.1	3.5	5.6	12.9	17.3
1958	1.6	*	*	*	*	*	0.6	2.0	3.6	5.7	12.7	21.3
1957	1.5	*	*	*	*	*	0.5	1.9	3.2	5.5	13.3	20.8
1956	1.4	*	*	*	*	*	0.5	1.6	3.3	5.6	12.2	19.4
1955	1.4	*	*	*	*	*	0.7	1.8	3.2	5.8	11.5	20.2
1954	1.5	*	*	*	*	*	0.6	1.8	3.1	6.4	12.3	21.1
1953	1.4	*	*	*	*	*	0.5	1.6	3.0	5.7	13.0	22.6
1952	1.4	*	*	*	*	*	0.6	1.5	2.7	6.6	12.7	24.7
1951	1.4	*	*	*	*	*	0.5	1.5	2.9	6.3	12.3	22.6
1950	1.4	*	*	*	*	0.2	0.5	1.6	3.1	6.4	13.8	21.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)												
ALL OTHER, BOTH SEXES												
1959	2.7	*	*	*	*	*	2.3	7.1	11.1	12.2	14.8	*
1958	2.5	*	*	*	*	*	1.9	5.8	11.3	11.5	16.8	*
1957	2.8	*	*	*	*	*	1.7	6.9	12.4	14.0	17.1	*
1956	2.6	*	*	*	*	*	1.4	7.0	11.8	12.4	13.1	*
1955	2.4	*	*	*	*	*	1.8	5.7	9.6	13.4	12.5	*
1954	2.3	*	*	*	*	*	2.1	5.0	12.1	10.2	16.0	*
1953	2.2	*	*	*	*	*	1.5	5.6	10.4	10.4	14.1	*
1952	2.4	*	*	*	*	*	1.9	5.7	10.1	12.4	16.8	*
1951	2.4	*	*	*	*	*	2.1	6.5	11.9	10.5	9.2	*
1950	2.4	*	*	*	*	*	1.6	6.3	12.3	12.4	13.8	*
ALL OTHER, MALE												
1959	4.2	*	*	*	*	*	3.2	11.9	18.1	19.4	24.3	*
1958	3.8	*	*	*	*	*	2.4	8.5	18.3	18.1	27.3	*
1957	4.5	*	*	*	*	*	2.8	11.4	20.5	24.5	26.7	*
1956	3.9	*	*	*	*	*	1.8	10.4	18.8	19.7	17.7	*
1955	3.5	*	*	*	*	*	2.6	8.1	16.5	19.1	18.4	*
1954	3.6	*	*	*	*	*	2.9	7.7	19.7	16.2	24.4	*
1953	3.3	*	*	*	*	*	1.9	7.8	15.9	17.7	22.8	*
1952	3.6	*	*	*	*	*	2.5	8.6	16.3	15.8	25.5	*
1951	3.6	*	*	*	*	*	3.1	9.1	18.3	15.3	*	*
1950	3.5	*	*	*	*	*	2.4	8.4	18.9	18.1	*	*
ALL OTHER, FEMALE												
1959	1.2	*	*	*	*	*	*	2.5	4.3	5.7	*	*
1958	1.3	*	*	*	*	*	*	3.3	4.6	5.4	*	*
1957	1.1	*	*	*	*	*	*	2.6	4.6	*	*	*
1956	1.4	*	*	*	*	*	*	3.8	4.9	5.7	*	*
1955	1.3	*	*	*	*	*	*	3.4	*	8.1	*	*
1954	1.2	*	*	*	*	*	*	2.5	4.5	*	*	*
1953	1.1	*	*	*	*	*	*	3.5	4.7	*	*	*
1952	1.3	*	*	*	*	*	*	2.9	*	9.2	*	*
1951	1.3	*	*	*	*	*	*	3.9	5.1	6.1	*	*
1950	1.4	*	*	*	*	*	*	4.2	5.6	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)												
BOTH SEXES												
1959	50.7	*	0.3	0.1	0.6	2.9	12.1	46.4	128.9	271.0	476.3	633.3
1958	50.9	0.6	0.3	0.1	0.6	2.8	12.1	46.2	129.1	277.1	481.4	633.4
1957	51.7	0.6	0.2	0.1	0.5	2.9	12.3	46.2	132.3	283.9	492.0	658.8
1956	52.4	0.8	0.3	0.1	0.7	2.9	13.2	47.7	135.3	284.2	512.9	649.7
1955	52.3	1.1	0.2	0.2	0.7	2.8	12.6	47.3	133.8	291.3	510.7	654.9
1954	53.0	1.1	0.2	0.2	0.7	3.0	13.6	50.0	136.8	296.0	516.3	655.7
1953	53.4	0.6	0.3	0.1	0.7	2.9	13.3	50.4	141.8	296.3	530.8	652.7
1952	53.5	0.9	0.4	0.1	0.7	3.1	13.6	51.1	143.4	298.3	537.1	654.7
1951	53.4	*	0.3	0.2	0.8	3.4	14.0	52.3	145.4	303.1	525.5	663.0
1950	54.2	*	0.4	0.2	0.8	3.4	14.3	53.4	150.0	313.5	554.8	670.7
MALE												
1959	56.4	*	0.3	0.2	0.6	3.1	12.9	53.4	155.0	332.0	555.8	691.0
1958	56.4	*	0.3	0.1	0.7	3.1	12.9	53.1	154.0	336.1	561.9	659.6
1957	57.6	*	0.3	0.2	0.6	2.9	13.2	52.4	157.9	346.9	570.2	705.3
1956	58.6	1.1	0.3	*	0.9	2.9	14.0	54.4	161.0	343.5	599.6	700.0
1955	58.4	1.7	0.3	0.2	0.8	2.9	12.7	54.2	158.9	353.4	589.0	685.8
1954	59.2	1.5	*	0.2	0.8	3.0	14.2	56.4	162.4	357.0	590.6	678.9
1953	59.6	*	0.4	0.2	0.8	3.1	14.1	57.1	169.0	353.6	604.1	678.5
1952	59.6	*	0.4	*	0.8	3.0	14.4	57.1	169.1	357.7	606.9	672.2
1951	59.2	*	0.4	0.3	1.0	3.2	14.2	57.8	170.4	360.4	589.4	710.2
1950	60.1	*	0.4	0.2	0.9	3.7	14.9	59.9	174.8	369.0	623.6	706.4
FEMALE												
1959	45.2	*	*	*	0.5	2.8	11.4	39.6	104.4	217.6	414.3	595.6
1958	45.5	*	0.3	*	0.5	2.4	11.3	39.5	105.6	225.0	418.1	615.9
1957	46.0	*	*	*	0.5	2.8	11.5	40.1	107.9	227.8	429.8	627.1
1956	46.5	*	*	0.1	0.4	2.9	12.4	41.2	110.6	231.1	443.4	614.7
1955	46.3	*	*	*	0.5	2.7	12.4	40.6	109.7	235.2	447.5	633.0
1954	47.1	*	*	0.1	0.6	2.9	13.0	43.8	112.0	240.8	455.6	639.1
1953	47.3	*	*	*	0.5	2.8	12.6	43.8	115.3	244.0	470.5	634.1
1952	47.5	*	*	*	0.7	3.2	13.0	45.3	118.1	243.9	479.2	642.3
1951	47.6	*	*	*	0.7	3.5	13.8	46.9	120.6	250.1	471.6	629.4
1950	48.4	*	0.4	*	0.6	3.1	13.8	47.0	125.1	261.8	496.3	645.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)												
WHITE, BOTH SEXES												
1959	52.0	*	0.3	0.1	0.5	2.6	11.3	43.8	124.2	270.7	485.9	654.3
1958	52.1	0.6	0.3	0.1	0.5	2.6	11.5	43.1	124.2	277.0	491.8	658.2
1957	53.0	0.7	0.3	0.1	0.5	2.7	11.6	43.2	127.6	284.9	502.9	681.0
1956	53.8	0.7	0.2	0.1	0.6	2.7	12.3	44.9	130.5	286.8	523.3	672.8
1955	53.7	1.0	0.2	0.1	0.6	2.6	11.8	44.2	129.6	293.3	524.4	681.5
1954	54.4	1.2	0.2	0.2	0.6	2.8	12.8	46.6	132.9	298.6	527.1	679.1
1953	55.0	*	0.3	0.1	0.6	2.8	12.5	47.3	138.3	299.7	544.9	679.8
1952	55.0	0.9	0.4	0.1	0.7	3.0	12.8	47.6	140.0	301.9	550.8	681.8
1951	55.0	*	0.3	0.2	0.8	3.2	13.1	49.4	142.1	307.6	540.0	692.1
1950	55.8	*	0.3	0.2	0.7	3.1	13.5	49.9	146.8	316.1	569.5	696.5
WHITE, MALE												
1959	57.1	*	0.3	0.1	0.5	2.8	11.7	49.7	148.3	329.6	566.3	709.5
1958	57.3	*	0.4	*	0.7	2.9	12.1	49.3	147.9	333.6	575.2	686.1
1957	58.5	*	*	0.2	0.5	2.6	12.4	48.2	151.2	346.8	582.3	722.2
1956	59.4	*	0.3	*	0.8	2.6	12.9	50.6	154.1	344.8	610.7	720.2
1955	59.3	1.6	*	0.2	0.8	2.7	11.6	50.1	152.4	353.8	605.0	709.2
1954	60.0	1.6	*	0.2	0.7	2.8	13.1	51.8	156.9	357.2	601.9	695.4
1953	60.8	*	0.3	0.2	0.8	3.0	13.3	53.1	164.0	355.2	620.1	697.8
1952	60.7	*	0.4	*	0.8	2.9	13.3	52.5	164.0	360.1	621.2	692.1
1951	60.6	*	0.4	0.3	0.9	3.0	13.5	54.0	165.6	364.8	605.8	735.9
1950	61.3	*	0.3	0.2	0.9	3.4	14.0	55.7	170.7	370.4	639.4	730.2
WHITE, FEMALE												
1959	47.0	*	*	*	0.5	2.5	10.8	38.1	101.6	219.3	423.9	618.7
1958	47.1	*	0.3	*	0.4	2.3	10.9	37.1	101.9	227.2	426.9	639.7
1957	47.8	*	*	*	0.4	2.7	10.9	38.2	105.2	230.0	440.2	653.0
1956	48.4	*	*	0.2	0.4	2.7	11.6	39.3	108.1	234.9	454.1	639.9
1955	48.3	*	*	*	0.5	2.5	11.9	38.3	107.7	238.9	460.2	661.9
1954	49.0	*	*	*	0.5	2.8	12.5	41.5	109.7	245.5	466.6	667.4
1953	49.2	*	*	*	0.5	2.6	11.8	41.6	113.3	249.1	483.9	666.8
1952	49.5	*	*	*	0.6	3.1	12.4	42.8	116.5	248.6	493.1	674.4
1951	49.5	*	*	*	0.6	3.3	12.8	44.8	118.9	254.9	485.4	660.8
1950	50.3	*	0.4	*	0.6	2.9	13.0	44.1	123.0	265.8	510.7	673.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)												
ALL OTHER, BOTH SEXES												
1959	40.7	*	*	*	1.1	5.3	19.8	70.8	177.9	275.4	343.6	376.5
1958	40.8	*	*	*	1.0	4.2	17.3	75.7	181.2	278.2	336.3	334.8
1957	40.9	*	*	*	0.9	4.4	18.7	75.0	183.8	271.9	340.4	382.8
1956	41.0	*	*	*	1.2	4.7	21.3	74.9	188.7	253.7	364.9	370.5
1955	40.2	*	*	*	0.8	4.5	19.5	77.6	182.8	266.5	316.4	329.3
1954	41.4	*	*	*	1.2	4.6	20.7	82.9	183.1	265.2	361.5	356.4
1953	39.9	*	*	*	1.0	4.0	19.9	80.4	185.2	256.2	324.4	319.2
1952	40.5	*	*	*	1.3	4.0	20.9	84.8	186.2	256.1	335.0	320.4
1951	39.4	*	*	*	1.3	5.2	21.6	80.1	187.9	248.0	310.6	310.6
1950	41.0	*	*	*	0.9	5.8	21.7	87.4	188.4	278.6	332.0	364.0
ALL OTHER, MALE												
1959	50.8	*	*	*	*	6.1	23.3	89.2	224.5	361.1	422.2	482.8
1958	49.4	*	*	*	*	5.4	19.9	90.1	218.7	364.9	391.4	374.1
1957	50.7	*	*	*	*	5.4	20.8	92.8	231.2	349.1	414.1	503.8
1956	51.2	*	*	*	2.1	5.5	23.5	91.1	238.6	328.6	455.4	456.0
1955	50.2	*	*	*	*	4.8	22.3	93.1	232.6	348.3	379.2	400.0
1954	52.1	*	*	*	*	5.1	24.0	101.2	226.7	354.3	440.3	468.2
1953	49.4	*	*	*	*	3.6	20.9	96.0	228.9	333.4	388.6	433.3
1952	50.4	*	*	*	*	4.2	23.9	100.7	231.5	330.4	411.8	420.0
1951	47.6	*	*	*	*	5.6	20.8	93.3	231.2	307.6	369.8	389.5
1950	49.6	*	*	*	*	6.4	23.4	100.3	224.1	350.7	403.4	428.5
ALL OTHER, FEMALE												
1959	31.1	*	*	*	*	4.5	16.6	53.5	133.0	197.7	273.3	297.4
1958	32.7	*	*	*	*	3.2	14.9	62.3	145.4	198.6	286.3	315.8
1957	31.7	*	*	*	*	3.6	16.8	58.1	137.7	201.4	271.7	308.1
1956	31.5	*	*	*	*	4.0	19.2	59.2	139.7	184.0	279.7	311.1
1955	30.7	*	*	*	*	4.2	16.8	62.6	133.5	191.1	254.5	279.4
1954	31.4	*	*	*	*	4.1	17.7	65.0	139.4	182.9	284.1	290.6
1953	31.0	*	*	*	*	4.3	18.9	65.2	140.5	183.5	261.2	241.9
1952	31.1	*	*	*	*	3.9	18.2	69.1	139.7	186.3	262.1	251.7
1951	31.6	*	*	*	*	4.9	22.3	67.0	142.9	191.6	254.1	257.1
1950	32.7	*	*	*	*	5.2	20.1	74.5	151.8	206.1	261.6	317.8

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)												
BOTH SEXES												
1959	21.2	*	*	*	0.1	1.1	6.9	30.4	79.5	111.4	96.4	77.4
1958	20.4	*	*	0.1	0.2	1.1	6.9	30.0	76.0	106.7	94.7	71.5
1957	19.9	*	*	*	0.2	1.1	6.3	29.0	75.2	103.3	93.0	72.8
1956	19.2	*	*	0.1	0.2	1.0	6.3	28.3	73.1	99.3	90.5	66.5
1955	18.2	*	*	0.1	0.3	1.0	5.9	27.4	68.5	92.9	88.2	65.8
1954	17.1	*	*	*	0.3	1.0	5.8	26.0	64.8	87.7	81.1	63.2
1953	16.7	*	*	*	0.3	0.9	5.9	25.9	64.9	82.0	78.7	63.0
1952	15.6	*	*	*	0.3	0.9	5.6	24.0	60.7	78.2	74.5	60.9
1951	14.7	*	*	0.1	0.3	0.8	5.2	23.9	56.7	73.1	69.6	60.4
1950	14.1	*	*	0.1	0.2	0.9	5.1	22.9	54.9	69.9	69.3	64.0
MALE												
1959	36.7	*	*	*	0.2	1.6	10.9	51.9	146.1	208.4	171.2	137.5
1958	35.3	*	*	*	0.3	1.7	10.6	51.6	140.0	197.6	164.5	112.8
1957	34.4	*	*	*	0.3	1.5	9.9	50.1	138.3	189.8	160.8	117.8
1956	33.1	*	*	*	0.3	1.5	10.0	49.2	132.7	180.9	154.8	109.2
1955	31.2	*	*	*	0.4	1.4	9.3	48.0	123.9	166.5	150.1	100.6
1954	29.3	*	*	*	0.5	1.4	9.3	45.1	117.1	155.9	131.1	97.7
1953	28.4	*	*	*	0.4	1.4	9.3	45.0	116.4	143.4	128.1	94.8
1952	26.2	*	*	*	0.4	1.3	8.8	41.3	106.8	133.6	117.1	93.8
1951	24.4	*	*	*	0.4	1.1	8.0	40.9	98.8	125.4	105.8	85.5
1950	23.1	*	*	*	0.3	1.3	8.1	39.3	94.2	116.3	105.1	95.4
FEMALE												
1959	6.1	*	*	*	*	0.7	3.2	9.5	16.9	26.5	38.0	38.1
1958	6.0	*	*	*	*	0.6	3.3	8.9	15.6	26.6	39.9	44.0
1957	5.8	*	*	*	*	0.6	2.9	8.4	15.2	26.5	39.1	42.1
1956	5.7	*	*	*	*	0.6	2.8	7.8	16.1	26.2	39.0	36.8
1955	5.6	*	*	*	0.2	0.6	2.7	7.2	15.1	26.6	38.3	41.2
1954	5.3	*	*	*	*	0.5	2.4	7.3	14.2	25.7	40.4	38.6
1953	5.3	*	*	*	*	0.6	2.6	7.0	14.7	25.9	38.3	40.0
1952	5.4	*	*	*	0.2	0.5	2.4	6.9	15.2	27.4	39.1	37.3
1951	5.2	*	*	*	0.3	0.5	2.4	7.2	14.7	24.7	39.2	42.5
1950	5.2	*	*	*	0.2	0.6	2.3	6.7	15.4	26.7	38.8	42.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)												
WHITE, BOTH SEXES												
1959	21.9	*	*	*	0.1	1.1	6.5	29.6	79.5	113.4	98.0	79.0
1958	21.2	*	*	0.1	0.2	1.1	6.4	29.3	76.4	108.5	97.3	73.8
1957	20.6	*	*	*	0.2	1.0	6.2	28.2	75.1	105.4	95.6	75.3
1956	20.0	*	*	0.1	0.2	1.0	6.1	27.8	73.1	101.8	93.0	69.3
1955	18.9	*	*	*	0.3	1.0	5.7	26.8	68.9	95.6	90.6	68.5
1954	17.8	*	*	*	0.3	0.9	5.6	25.5	65.3	90.2	83.4	64.9
1953	17.3	*	*	*	0.3	0.9	5.7	25.4	65.2	84.5	81.0	64.6
1952	16.3	*	*	*	0.3	0.9	5.3	23.7	61.2	80.5	76.8	62.8
1951	15.3	*	*	0.1	0.3	0.8	5.1	23.7	56.8	75.0	71.7	63.1
1950	14.7	*	*	0.1	0.2	0.9	5.1	22.8	55.7	71.6	71.6	68.0
WHITE, MALE												
1959	37.9	*	*	*	*	1.6	10.0	50.8	146.5	213.1	174.8	139.9
1958	36.5	*	*	*	0.3	1.6	9.8	50.3	141.3	201.6	169.6	116.1
1957	35.6	*	*	*	0.3	1.5	9.7	48.8	138.5	194.3	165.9	119.9
1956	34.4	*	*	*	0.4	1.4	9.7	48.3	133.2	185.9	159.7	113.9
1955	32.5	*	*	*	0.3	1.4	8.9	47.2	125.3	172.1	155.2	105.1
1954	30.4	*	*	*	0.4	1.3	9.0	44.4	118.0	160.8	135.0	99.6
1953	29.5	*	*	*	0.4	1.3	9.0	44.4	117.3	148.0	132.0	97.4
1952	27.2	*	*	*	0.4	1.2	8.4	40.9	108.1	137.8	121.0	95.3
1951	25.3	*	*	*	0.4	1.1	8.0	40.7	99.4	128.9	109.4	89.5
1950	24.1	*	*	*	0.3	1.2	7.9	39.1	95.9	119.4	109.1	102.7
WHITE, FEMALE												
1959	6.3	*	*	*	*	0.6	3.1	9.1	16.7	26.5	38.7	39.7
1958	6.2	*	*	*	*	0.6	3.1	8.7	15.3	26.7	41.0	45.8
1957	6.0	*	*	*	*	0.5	2.8	8.2	15.1	26.5	40.3	44.9
1956	5.9	*	*	*	*	0.6	2.6	7.6	15.9	26.7	40.1	38.2
1955	5.7	*	*	*	*	0.6	2.6	6.8	14.8	26.7	39.1	42.7
1954	5.6	*	*	*	*	0.5	2.3	6.9	14.4	26.2	41.9	39.9
1953	5.5	*	*	*	*	0.5	2.5	6.7	14.6	26.6	39.7	41.0
1952	5.6	*	*	*	*	0.5	2.2	6.8	15.1	27.9	40.5	39.5
1951	5.4	*	*	*	0.3	0.5	2.3	6.9	14.5	25.5	40.5	44.3
1950	5.4	*	*	*	*	0.5	2.2	6.5	15.5	27.2	40.0	44.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)												
ALL OTHER, BOTH SEXES												
1959	15.7	*	*	*	*	1.4	10.9	37.5	79.4	86.9	73.4	57.4
1958	14.7	*	*	*	*	1.5	10.6	36.6	71.7	84.7	59.6	43.9
1957	14.2	*	*	*	*	1.2	7.4	36.0	75.5	78.7	56.4	42.2
1956	13.3	*	*	*	*	1.4	8.7	33.0	72.2	69.1	56.0	32.8
1955	12.3	*	*	*	*	1.4	7.9	33.2	63.6	61.3	53.5	*
1954	11.5	*	*	*	*	1.5	7.7	31.0	59.1	57.3	48.0	41.8
1953	11.2	*	*	*	*	1.4	7.5	30.4	60.7	52.1	45.3	42.3
1952	10.3	*	*	*	*	1.2	7.8	26.5	54.7	51.2	40.7	*
1951	9.7	*	*	*	*	0.9	5.3	26.0	54.9	49.2	37.8	*
1950	8.8	*	*	*	*	1.6	5.9	24.6	44.7	47.8	33.9	*
ALL OTHER, MALE												
1959	27.0	*	*	*	*	2.2	18.4	63.0	141.0	154.0	125.0	110.3
1958	25.4	*	*	*	*	2.7	17.4	64.0	126.5	150.1	99.3	77.8
1957	24.6	*	*	*	*	*	11.6	63.1	136.0	136.4	94.8	92.3
1956	23.0	*	*	*	*	2.1	13.4	57.8	126.5	121.8	91.5	*
1955	20.6	*	*	*	*	2.2	12.9	56.3	108.0	100.6	83.2	*
1954	20.0	*	*	*	*	2.4	12.4	51.5	106.5	97.4	79.8	*
1953	19.2	*	*	*	*	1.7	11.8	51.6	105.0	88.7	75.4	*
1952	17.4	*	*	*	*	1.9	12.5	45.0	92.2	83.3	64.5	*
1951	16.0	*	*	*	*	*	7.6	42.3	91.1	84.1	56.6	*
1950	14.5	*	*	*	*	2.1	9.4	40.6	74.4	74.8	48.6	*
ALL OTHER, FEMALE												
1959	5.0	*	*	*	*	*	4.1	13.5	20.1	26.1	27.3	*
1958	4.5	*	*	*	*	*	4.5	10.7	18.9	24.9	23.5	*
1957	4.3	*	*	*	*	*	3.5	10.2	16.5	26.0	20.7	*
1956	4.1	*	*	*	*	*	4.3	9.1	18.9	20.5	22.5	*
1955	4.5	*	*	*	*	*	3.3	10.9	19.6	25.2	25.0	*
1954	3.6	*	*	*	*	*	3.4	11.1	11.6	20.2	17.5	*
1953	3.7	*	*	*	*	*	3.5	9.6	15.8	17.9	16.5	*
1952	3.6	*	*	*	*	*	3.4	8.2	16.1	21.0	18.1	*
1951	3.7	*	*	*	*	*	3.2	9.8	17.2	16.2	19.8	*
1950	3.4	*	*	*	*	*	2.6	8.7	14.1	20.7	19.5	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF BREAST (170)												
BOTH SEXES												
1959	13.1	*	*	*	*	2.2	10.0	24.8	37.0	48.2	71.5	117.3
1958	13.1	*	*	*	*	1.9	10.6	24.9	36.1	48.6	73.4	111.7
1957	13.3	*	*	*	*	2.1	10.0	25.5	36.6	50.1	77.0	120.1
1956	13.3	*	*	*	*	1.9	10.8	24.8	36.8	49.9	76.8	119.2
1955	13.4	*	*	*	*	2.0	10.9	25.2	36.6	49.5	79.2	118.0
1954	13.0	*	*	*	*	2.1	10.2	24.5	35.6	49.5	76.4	115.0
1953	13.0	*	*	*	0.1	2.0	10.6	24.5	35.3	49.6	78.2	117.2
1952	12.8	*	*	*	*	2.1	10.6	23.8	36.5	48.3	74.7	109.6
1951	12.5	*	*	*	*	1.9	10.0	23.2	35.2	48.2	79.0	116.2
1950	12.6	*	*	*	*	1.9	10.6	23.7	35.3	50.1	76.8	116.7
MALE												
1959	0.2	*	*	*	*	*	*	0.3	0.6	1.2	2.0	*
1958	0.3	*	*	*	*	*	*	0.2	0.8	1.2	2.1	*
1957	0.3	*	*	*	*	*	*	0.3	0.8	1.9	2.7	*
1956	0.3	*	*	*	*	*	*	0.3	0.7	1.1	2.5	*
1955	0.3	*	*	*	*	*	*	0.3	0.9	1.5	2.2	*
1954	0.3	*	*	*	*	*	*	0.3	0.7	1.5	2.6	*
1953	0.3	*	*	*	*	*	*	0.4	0.8	1.4	2.2	*
1952	0.3	*	*	*	*	*	*	0.3	0.8	1.5	3.3	*
1951	0.3	*	*	*	*	*	*	0.4	0.9	1.6	1.8	*
1950	0.3	*	*	*	*	*	*	0.4	0.8	1.9	2.7	*
FEMALE												
1959	25.5	*	*	*	*	4.3	19.5	48.5	71.1	89.2	125.9	191.3
1958	25.6	*	*	*	*	3.8	20.7	48.9	69.5	90.4	129.5	182.8
1957	26.0	*	*	*	*	4.1	19.6	50.0	70.6	93.0	136.0	199.8
1956	26.0	*	*	*	*	3.8	21.1	48.8	71.2	93.6	136.3	199.1
1955	26.1	*	*	*	*	3.8	21.3	49.6	71.0	92.8	141.3	198.2
1954	25.3	*	*	*	*	4.1	19.9	48.2	69.4	93.2	136.4	193.9
1953	25.3	*	*	*	0.2	3.9	20.6	48.3	69.0	93.6	140.5	198.8
1952	24.9	*	*	*	*	4.0	20.7	47.0	71.8	91.2	133.9	184.8
1951	24.5	*	*	*	*	3.6	19.7	45.8	69.4	91.1	143.6	196.7
1950	24.7	*	*	*	*	3.8	20.8	46.9	69.9	95.0	139.8	195.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF BREAST (170)												
WHITE, BOTH SEXES												
1959	13.6	*	*	*	*	2.1	9.9	24.9	37.2	49.3	73.7	121.4
1958	13.6	*	*	*	*	1.8	10.4	25.1	36.4	49.9	75.6	115.9
1957	13.9	*	*	*	*	2.0	9.8	25.5	37.0	51.3	79.1	125.5
1956	13.8	*	*	*	*	1.9	10.6	24.7	37.1	51.1	79.0	122.7
1955	13.9	*	*	*	*	1.8	10.7	25.4	37.1	50.6	81.5	124.1
1954	13.4	*	*	*	*	1.9	9.9	24.6	36.0	50.8	78.8	119.7
1953	13.5	*	*	*	*	1.8	10.5	24.6	36.0	51.1	80.6	121.0
1952	13.3	*	*	*	*	2.0	10.4	24.0	36.9	49.7	76.5	114.7
1951	13.0	*	*	*	*	1.8	9.9	23.1	35.6	49.6	81.7	120.4
1950	13.1	*	*	*	*	1.9	10.6	23.8	35.8	50.9	79.2	121.8
WHITE, MALE												
1959	0.2	*	*	*	*	*	*	0.3	0.6	1.2	1.9	*
1958	0.3	*	*	*	*	*	*	*	0.8	1.1	2.2	*
1957	0.3	*	*	*	*	*	*	0.3	0.8	1.9	2.7	*
1956	0.2	*	*	*	*	*	*	0.3	0.7	1.1	2.4	*
1955	0.3	*	*	*	*	*	*	0.3	1.0	1.5	2.3	*
1954	0.3	*	*	*	*	*	*	0.3	0.6	1.5	2.5	*
1953	0.3	*	*	*	*	*	*	0.4	0.9	1.3	2.3	*
1952	0.3	*	*	*	*	*	*	0.3	0.7	1.5	3.3	*
1951	0.3	*	*	*	*	*	*	0.3	1.0	1.7	1.9	*
1950	0.3	*	*	*	*	*	*	0.4	0.8	1.9	2.8	*
WHITE, FEMALE												
1959	26.6	*	*	*	*	4.1	19.3	48.8	71.5	91.1	129.3	197.8
1958	26.7	*	*	*	*	3.6	20.3	49.4	69.9	92.7	132.9	188.9
1957	27.1	*	*	*	*	4.0	19.3	50.1	71.3	95.1	139.1	208.8
1956	27.0	*	*	*	*	3.7	20.7	48.6	71.7	95.9	139.8	205.5
1955	27.2	*	*	*	*	3.6	21.0	50.0	71.8	94.9	144.8	208.9
1954	26.3	*	*	*	*	3.7	19.3	48.4	70.0	95.4	140.2	202.8
1953	26.4	*	*	*	*	3.6	20.5	48.5	70.2	96.4	144.1	205.7
1952	26.0	*	*	*	*	3.9	20.4	47.6	72.4	93.8	136.6	193.8
1951	25.5	*	*	*	*	3.5	19.4	45.6	69.9	93.7	148.0	203.9
1950	25.7	*	*	*	*	3.7	20.8	47.1	70.9	96.3	143.6	204.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF BREAST (170)												
ALL OTHER, BOTH SEXES												
1959	8.9	*	*	*	*	3.0	11.5	23.7	34.6	34.9	41.3	66.2
1958	8.7	*	*	*	*	2.8	12.4	22.7	33.3	33.2	42.1	62.1
1957	9.0	*	*	*	*	2.5	11.7	25.5	32.8	36.1	47.5	53.1
1956	9.3	*	*	*	*	2.2	13.2	26.3	33.3	35.0	44.4	77.0
1955	8.9	*	*	*	*	3.0	12.7	23.4	30.7	36.2	46.5	43.1
1954	9.0	*	*	*	*	3.6	13.1	23.7	31.8	34.8	42.2	54.5
1953	8.5	*	*	*	*	3.3	11.5	23.7	27.6	32.3	44.0	71.2
1952	8.5	*	*	*	*	2.7	12.3	21.4	32.3	31.4	48.7	46.9
1951	8.4	*	*	*	*	2.4	11.4	24.2	30.8	30.9	38.7	66.0
1950	8.3	*	*	*	*	2.5	10.9	22.8	29.6	38.9	39.8	55.8
ALL OTHER, MALE												
1959	0.3	*	*	*	*	*	*	*	*	*	*	*
1958	0.2	*	*	*	*	*	*	*	*	*	*	*
1957	0.3	*	*	*	*	*	*	*	*	*	*	*
1956	0.3	*	*	*	*	*	*	*	*	*	*	*
1955	*	*	*	*	*	*	*	*	*	*	*	*
1954	0.4	*	*	*	*	*	*	*	*	*	*	*
1953	0.3	*	*	*	*	*	*	*	*	*	*	*
1952	0.3	*	*	*	*	*	*	*	*	*	*	*
1951	0.3	*	*	*	*	*	*	*	*	*	*	*
1950	0.3	*	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	17.0	*	*	*	*	5.5	21.7	45.8	66.9	65.7	75.8	107.7
1958	16.8	*	*	*	*	5.3	23.6	43.7	64.9	61.9	79.1	105.3
1957	17.2	*	*	*	*	4.6	22.3	49.5	63.1	66.7	89.7	89.2
1956	17.7	*	*	*	*	4.1	25.3	50.9	65.1	65.4	82.6	122.2
1955	17.1	*	*	*	*	5.5	24.5	45.8	60.7	67.9	89.4	67.6
1954	17.2	*	*	*	*	6.7	24.9	46.1	62.0	65.6	77.0	84.4
1953	16.3	*	*	*	*	6.3	21.6	46.7	55.0	60.2	84.3	116.1
1952	16.2	*	*	*	*	5.0	23.5	41.9	64.0	58.8	91.4	75.9
1951	16.1	*	*	*	*	4.5	21.8	47.1	62.0	59.5	74.8	110.7
1950	16.0	*	*	*	*	4.7	20.8	45.3	58.1	76.4	78.1	91.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)												
BOTH SEXES												
1959	21.6	*	*	0.1	0.9	3.5	11.1	24.7	49.8	100.1	189.0	265.8
1958	22.1	*	0.1	0.1	0.9	3.8	11.3	25.9	51.5	102.2	195.2	271.2
1957	22.6	*	0.2	0.1	0.9	4.0	11.2	27.6	52.5	103.9	199.0	274.7
1956	22.6	*	*	0.1	0.9	4.1	11.6	27.0	52.4	102.1	206.3	281.4
1955	22.8	*	0.1	0.1	1.0	4.0	12.5	28.2	53.7	104.6	202.3	272.9
1954	22.9	*	*	0.1	1.1	3.8	12.5	28.7	54.2	104.2	209.8	268.7
1953	22.9	*	*	0.1	0.9	4.0	13.0	29.4	54.3	105.7	205.8	267.4
1952	22.7	*	0.2	0.1	1.0	4.1	12.8	29.7	54.4	104.8	204.0	274.5
1951	22.5	*	*	0.1	1.1	4.0	13.5	30.6	54.3	103.3	202.7	252.9
1950	23.1	*	0.2	0.2	1.0	4.3	14.2	31.9	56.7	107.1	207.9	262.4
MALE												
1959	17.3	*	*	*	0.9	1.6	1.9	3.8	23.9	102.0	278.5	467.9
1958	17.6	*	*	*	1.0	1.8	1.9	4.1	24.8	104.3	286.9	470.1
1957	18.2	*	*	*	0.8	1.7	1.7	4.2	25.4	108.4	299.8	479.2
1956	18.1	*	*	*	0.8	2.0	1.7	4.0	25.6	105.3	309.1	488.4
1955	17.9	*	*	*	0.9	1.9	1.7	4.5	26.6	106.4	296.2	465.7
1954	17.9	*	*	*	1.1	2.0	1.7	4.2	26.6	106.1	305.5	464.1
1953	17.3	*	*	*	0.9	1.8	1.8	4.1	24.7	105.7	296.9	451.9
1952	17.1	*	*	*	0.9	1.7	1.6	4.6	26.4	103.5	290.2	464.5
1951	16.5	*	*	*	1.0	2.1	1.8	4.8	25.4	100.6	286.5	421.9
1950	16.4	*	*	*	0.8	2.0	1.7	4.8	26.2	103.9	291.4	436.2
FEMALE												
1959	25.8	*	*	0.2	0.9	5.2	19.8	45.1	74.2	98.4	119.2	133.7
1958	26.5	*	*	0.2	0.9	5.7	20.3	47.2	76.7	100.3	123.0	138.6
1957	26.9	*	*	0.2	1.0	6.2	20.4	50.4	78.2	99.8	118.9	135.2
1956	26.9	*	*	0.1	1.0	6.2	21.1	49.5	78.1	99.3	124.1	137.4
1955	27.7	*	*	0.2	1.1	5.9	22.9	51.5	79.7	103.0	126.5	136.3
1954	27.8	*	*	0.2	1.0	5.5	23.0	52.8	80.9	102.5	132.0	128.9
1953	28.4	*	*	*	0.9	6.1	23.8	54.3	83.2	105.8	131.0	134.8
1952	28.2	*	*	*	1.0	6.4	23.6	54.5	82.0	106.0	132.6	138.3
1951	28.4	*	*	*	1.1	5.8	24.8	56.1	83.1	105.7	132.3	132.8
1950	29.6	*	*	0.2	1.2	6.4	26.3	59.0	87.4	110.1	136.9	141.4

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)												
WHITE, BOTH SEXES												
1959	21.1	*	*	0.1	0.9	3.2	9.9	22.5	46.0	95.3	187.0	266.5
1958	21.7	*	*	0.1	0.9	3.4	10.1	23.5	47.6	97.9	194.1	272.5
1957	22.1	*	0.2	0.1	0.9	3.7	10.0	24.9	48.6	99.0	198.6	279.2
1956	22.0	*	*	0.1	0.9	3.8	10.2	24.4	47.9	98.0	204.9	287.4
1955	22.4	*	0.2	0.1	1.0	3.6	10.9	25.3	50.0	101.7	201.8	277.2
1954	22.5	*	*	0.1	1.1	3.5	11.2	25.8	50.0	101.4	210.4	274.4
1953	22.5	*	*	0.1	0.9	3.6	11.5	26.4	50.9	102.2	206.7	273.1
1952	22.3	*	0.2	0.1	1.0	3.7	11.3	26.4	51.0	102.6	204.9	278.7
1951	22.1	*	*	0.1	1.0	3.6	11.9	27.6	51.0	101.2	203.9	257.6
1950	22.7	*	*	0.2	0.9	3.8	12.4	29.0	53.4	104.5	209.2	266.5
WHITE, MALE												
1959	17.0	*	*	*	0.9	1.8	1.9	2.9	20.5	94.3	275.9	473.3
1958	17.5	*	*	*	1.1	2.0	1.9	3.5	21.3	98.2	286.6	477.5
1957	18.0	*	*	*	0.9	1.9	1.6	3.3	22.0	101.1	301.1	489.4
1956	18.1	*	*	*	0.9	2.1	1.7	3.3	21.6	100.3	310.2	500.7
1955	17.9	*	*	*	1.0	2.0	1.7	3.6	23.2	102.3	297.1	475.5
1954	17.9	*	*	*	1.2	2.1	1.7	3.3	22.6	101.8	308.3	477.3
1953	17.4	*	*	*	1.0	1.9	1.8	3.3	21.6	101.5	302.1	464.6
1952	17.2	*	*	*	1.0	1.8	1.6	3.6	23.4	101.7	292.4	476.3
1951	16.6	*	*	*	1.0	2.2	1.8	3.8	22.6	98.0	291.1	430.8
1950	16.6	*	*	*	0.9	2.1	1.7	3.7	23.7	101.7	294.5	447.8
WHITE, FEMALE												
1959	25.2	*	*	0.1	0.9	4.5	17.6	41.6	69.8	96.1	118.3	132.7
1958	25.8	*	*	0.2	0.8	4.8	18.0	42.9	72.4	97.7	122.1	137.0
1957	26.1	*	*	0.2	0.9	5.5	18.1	45.9	73.8	97.1	117.9	136.1
1956	25.9	*	*	0.1	0.9	5.4	18.4	45.1	73.1	95.9	121.5	138.9
1955	26.9	*	*	*	0.9	5.1	19.9	46.7	75.8	101.2	125.7	136.6
1954	27.0	*	*	*	0.9	4.7	20.3	48.0	76.4	101.1	131.6	128.8
1953	27.4	*	*	*	0.8	5.2	21.0	49.1	79.4	102.9	129.2	134.8
1952	27.3	*	*	*	1.0	5.4	20.7	49.0	78.0	103.5	133.1	136.6
1951	27.6	*	*	*	1.1	4.9	21.7	51.3	79.2	104.0	131.5	134.0
1950	28.6	*	*	0.2	1.0	5.5	22.9	54.3	83.1	107.1	137.3	140.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)												
ALL OTHER, BOTH SEXES												
1959	25.2	*	*	*	0.9	5.8	21.3	45.9	90.3	157.5	217.0	257.4
1958	25.6	*	*	*	0.8	6.8	21.8	49.6	93.6	153.3	209.9	254.5
1957	26.3	*	*	*	1.1	6.5	22.0	53.4	94.7	163.0	206.1	217.2
1956	26.8	*	*	*	1.0	6.9	23.6	52.4	103.4	151.2	226.9	209.8
1955	26.2	*	*	*	1.1	6.9	25.9	56.1	95.8	138.8	210.5	220.7
1954	26.2	*	*	*	1.1	6.6	24.4	56.5	104.0	137.8	202.0	194.5
1953	26.3	*	*	*	0.9	7.3	25.5	58.1	95.9	147.5	192.7	198.1
1952	26.1	*	*	*	1.0	7.9	25.8	61.1	97.5	130.9	190.7	222.4
1951	25.6	*	*	*	1.1	7.4	27.7	58.6	96.4	128.1	184.3	195.7
1950	26.5	*	*	*	1.4	7.9	29.4	60.0	97.0	142.6	188.8	214.4
ALL OTHER, MALE												
1959	19.5	*	*	*	*	*	2.1	12.2	59.3	192.4	311.8	406.9
1958	18.3	*	*	*	*	*	1.9	9.3	62.2	176.2	290.6	400.0
1957	19.2	*	*	*	*	*	2.2	12.5	61.9	195.8	283.7	357.7
1956	18.4	*	*	*	*	*	*	11.3	70.5	163.3	294.6	340.0
1955	17.7	*	*	*	*	*	2.5	12.7	65.5	155.2	284.0	345.8
1954	17.8	*	*	*	*	*	*	13.7	72.8	157.4	268.1	295.5
1953	16.2	*	*	*	*	*	1.9	11.4	62.0	154.7	227.2	290.5
1952	15.7	*	*	*	*	*	1.9	14.4	63.1	125.7	260.9	315.0
1951	15.3	*	*	*	*	*	1.9	14.6	60.4	130.6	225.5	310.5
1950	14.8	*	*	*	*	*	1.9	15.4	56.1	132.5	247.8	300.0
ALL OTHER, FEMALE												
1959	30.5	*	*	*	1.4	10.5	38.7	77.7	120.1	125.9	132.3	146.2
1958	32.5	*	*	*	1.4	12.3	40.0	87.8	124.2	132.0	136.6	157.9
1957	33.0	*	*	*	2.0	11.9	40.1	92.4	126.7	133.0	133.8	124.3
1956	34.7	*	*	*	1.7	12.4	44.0	92.0	135.7	139.3	163.0	119.4
1955	34.2	*	*	*	1.9	12.3	47.6	98.1	125.8	123.7	139.4	132.4
1954	34.2	*	*	*	1.9	11.8	45.6	98.3	135.3	119.7	138.1	131.3
1953	35.9	*	*	*	*	13.2	47.4	103.9	130.0	140.4	158.7	135.5
1952	35.9	*	*	*	1.5	14.1	48.0	107.3	132.8	135.9	124.1	158.6
1951	35.5	*	*	*	1.5	13.0	51.6	102.1	133.9	125.7	145.0	117.9
1950	37.6	*	*	*	2.4	14.5	54.9	104.6	139.1	152.8	130.8	153.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)												
BOTH SEXES												
1959	7.1	*	1.2	0.3	*	0.3	1.3	6.0	17.5	37.7	66.5	87.9
1958	7.0	*	1.1	0.3	0.1	0.3	1.4	6.1	16.7	38.1	65.9	88.7
1957	7.1	0.6	1.1	0.3	0.1	0.3	1.5	6.3	18.2	37.6	67.1	91.3
1956	7.0	0.6	1.1	0.3	*	0.3	1.4	6.0	18.2	37.2	67.1	90.2
1955	7.0	*	1.1	0.3	*	0.3	1.3	6.0	18.0	38.3	67.6	86.2
1954	6.9	*	1.1	0.3	*	0.4	1.5	6.2	18.0	37.6	67.7	83.4
1953	6.9	0.6	1.0	0.3	0.1	0.3	1.6	6.4	18.6	37.8	64.0	84.5
1952	6.9	*	1.1	0.3	0.1	0.3	1.6	6.3	18.8	37.8	65.4	82.0
1951	6.6	0.8	1.2	0.3	0.1	0.3	1.4	6.5	18.3	35.9	63.3	88.3
1950	6.7	0.9	1.2	0.3	0.1	0.3	1.6	6.2	19.1	38.0	64.2	75.1
MALE												
1959	9.6	*	1.3	0.3	*	0.3	1.7	8.6	25.8	55.5	97.0	126.8
1958	9.4	*	1.2	0.3	*	0.4	1.9	8.4	24.5	55.8	94.5	120.3
1957	9.6	*	1.1	0.3	*	0.3	1.9	8.8	26.5	55.2	93.3	123.1
1956	9.4	*	1.1	0.3	*	0.3	1.7	8.4	26.9	54.1	93.8	122.3
1955	9.3	*	1.2	0.3	*	0.4	1.7	8.4	25.9	55.3	91.6	111.0
1954	9.4	*	1.3	0.2	*	0.5	2.0	8.5	26.5	53.2	93.2	116.4
1953	9.2	*	1.1	0.3	*	0.3	2.0	8.6	27.6	53.1	86.4	116.3
1952	9.2	*	0.9	0.2	*	0.4	2.0	9.0	27.2	53.0	89.2	111.0
1951	8.8	*	1.3	0.3	*	0.4	1.7	9.1	26.1	49.3	86.3	119.9
1950	8.8	*	1.2	0.2	*	0.3	1.9	8.7	26.9	52.1	84.8	115.3
FEMALE												
1959	4.6	*	1.1	0.3	*	0.3	1.0	3.5	9.8	22.1	42.6	62.4
1958	4.6	*	1.0	0.4	*	0.2	1.0	3.9	9.4	22.4	43.4	67.6
1957	4.8	*	1.1	0.3	*	0.3	1.1	3.8	10.4	21.9	46.2	69.6
1956	4.6	*	1.1	0.4	*	0.2	1.1	3.7	9.9	21.9	45.7	67.9
1955	4.7	*	1.0	0.4	*	0.3	1.0	3.6	10.4	23.0	48.2	68.6
1954	4.6	*	1.0	0.3	*	0.2	1.0	3.8	9.8	23.5	46.9	59.8
1953	4.6	*	0.9	0.4	*	0.3	1.2	4.3	9.8	23.8	45.6	61.7
1952	4.6	*	1.2	0.3	*	0.3	1.1	3.7	10.5	23.8	45.6	61.2
1951	4.5	*	1.1	0.3	*	0.3	1.0	3.9	10.6	23.6	44.0	65.8
1950	4.5	*	1.2	0.4	*	0.2	1.2	3.8	11.2	24.8	46.6	47.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)												
WHITE, BOTH SEXES												
1959	7.4	*	1.2	0.3	*	0.3	1.2	5.8	17.5	38.4	68.9	92.4
1958	7.3	*	1.1	0.3	*	0.2	1.4	5.7	16.7	38.9	68.4	93.8
1957	7.4	0.6	1.1	0.3	*	0.3	1.4	6.0	18.1	38.4	69.5	96.0
1956	7.3	0.6	1.1	0.3	*	0.2	1.3	5.8	18.2	38.2	69.3	94.8
1955	7.4	*	1.1	0.3	*	0.3	1.3	5.9	18.1	39.7	70.3	90.1
1954	7.3	*	1.2	0.3	*	0.3	1.4	5.9	18.0	38.7	70.3	87.0
1953	7.2	0.7	1.0	0.3	0.1	0.3	1.5	6.1	18.7	38.7	66.4	89.4
1952	7.2	*	1.1	0.3	0.1	0.3	1.5	6.2	18.8	38.9	67.6	87.3
1951	6.9	0.9	1.2	0.3	0.1	0.3	1.2	6.2	18.4	36.9	66.0	92.8
1950	7.0	0.9	1.2	0.3	0.1	0.2	1.4	6.1	19.3	38.7	66.7	78.0
WHITE, MALE												
1959	10.1	*	1.3	0.3	*	0.3	1.6	8.5	26.1	56.8	101.2	134.4
1958	9.9	*	1.2	0.3	*	0.3	1.8	8.0	24.8	57.5	98.4	127.8
1957	10.0	*	1.1	0.3	*	0.3	1.8	8.5	26.7	56.8	97.6	130.2
1956	9.9	*	1.1	0.3	*	0.2	1.6	8.3	27.2	56.1	97.1	128.1
1955	9.9	*	1.2	0.3	*	0.3	1.6	8.4	26.1	57.8	95.9	117.0
1954	9.8	*	1.3	0.2	*	0.5	2.0	8.3	26.6	55.1	97.9	120.9
1953	9.7	*	1.1	0.3	*	0.2	2.0	8.5	28.0	54.6	90.6	123.5
1952	9.7	*	0.9	0.3	*	0.3	1.9	9.0	27.4	55.0	92.7	119.0
1951	9.3	*	1.4	0.3	*	0.3	1.6	9.0	26.5	50.7	90.5	124.9
1950	9.3	*	1.2	0.2	*	0.3	1.8	8.7	27.6	53.6	88.1	120.1
WHITE, FEMALE												
1959	4.8	*	1.1	0.3	*	0.3	0.8	3.2	9.4	22.4	43.9	65.3
1958	4.8	*	1.0	0.4	*	*	0.9	3.4	9.1	22.6	45.0	71.3
1957	4.9	*	1.1	0.3	*	0.3	1.0	3.6	9.9	22.2	47.4	72.6
1956	4.8	*	1.1	0.3	*	*	1.0	3.3	9.6	22.3	47.2	71.7
1955	5.0	*	1.1	0.4	*	0.3	0.9	3.4	10.4	23.4	49.9	71.1
1954	4.8	*	1.0	0.3	*	0.2	0.9	3.5	9.6	23.8	48.1	62.6
1953	4.8	*	0.9	0.4	*	0.3	1.1	3.8	9.6	24.3	46.9	64.7
1952	4.8	*	1.3	0.3	*	0.3	1.0	3.4	10.3	24.3	47.0	64.5
1951	4.7	*	1.1	0.3	*	0.2	0.9	3.5	10.5	24.2	45.7	69.9
1950	4.7	*	1.1	0.5	*	*	1.0	3.5	11.0	24.8	48.7	48.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)												
ALL OTHER, BOTH SEXES												
1959	4.5	*	1.1	*	*	*	2.4	7.8	18.5	28.7	33.1	32.4
1958	4.5	*	1.0	*	*	*	2.3	10.0	16.6	28.1	31.8	*
1957	4.5	*	*	*	*	*	2.3	8.8	19.9	27.3	32.9	34.4
1956	4.3	*	1.1	*	*	*	2.5	8.2	17.9	24.0	36.2	34.4
1955	3.8	*	*	*	*	*	1.9	6.6	16.8	22.3	28.9	37.9
1954	4.3	*	*	*	*	*	2.2	9.0	18.5	24.4	29.5	38.2
1953	4.2	*	*	*	*	*	2.4	9.4	17.6	26.7	28.6	*
1952	4.1	*	*	*	*	*	2.4	7.6	19.0	24.3	32.7	*
1951	3.9	*	*	*	*	*	2.4	8.7	16.4	23.5	23.5	*
1950	4.0	*	1.3	*	*	*	2.8	7.4	16.3	28.5	26.1	*
ALL OTHER, MALE												
1959	5.6	*	*	*	*	*	2.4	9.3	23.3	40.7	43.8	*
1958	5.6	*	*	*	*	*	2.6	12.1	21.4	37.0	44.6	*
1957	5.7	*	*	*	*	*	2.7	11.4	24.5	36.4	38.5	*
1956	5.3	*	*	*	*	*	2.4	9.3	23.0	31.3	50.8	*
1955	4.7	*	*	*	*	*	2.4	8.0	23.2	26.2	35.2	*
1954	5.3	*	*	*	*	*	2.6	10.8	24.6	30.4	31.1	*
1953	5.1	*	*	*	*	*	2.4	10.2	22.0	35.8	30.7	*
1952	5.1	*	*	*	*	*	2.7	8.7	25.6	30.1	41.8	*
1951	4.9	*	*	*	*	*	2.7	10.1	20.9	31.8	30.2	*
1950	4.8	*	*	*	*	*	2.7	8.9	18.4	33.0	39.6	*
ALL OTHER, FEMALE												
1959	3.4	*	*	*	*	*	2.3	6.3	13.8	17.8	23.6	*
1958	3.4	*	*	*	*	*	2.0	8.0	12.0	20.0	20.3	*
1957	3.5	*	*	*	*	*	1.9	6.3	15.3	19.0	27.6	*
1956	3.4	*	*	*	*	*	2.7	7.0	12.9	17.3	22.5	*
1955	2.9	*	*	*	*	*	*	5.3	10.5	18.6	22.7	*
1954	3.3	*	*	*	*	*	1.8	7.2	12.4	18.9	27.8	*
1953	3.4	*	*	*	*	*	2.3	8.7	13.1	18.2	26.4	*
1952	3.1	*	*	*	*	*	2.0	6.6	12.2	18.8	24.1	*
1951	3.0	*	*	*	*	*	2.1	7.4	11.7	15.6	*	*
1950	3.3	*	*	*	*	*	2.8	6.0	14.1	23.8	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)												
BOTH SEXES												
1959	16.2	2.3	3.0	2.5	2.7	4.2	9.1	21.3	40.6	64.5	99.7	162.5
1958	16.1	2.2	3.1	2.3	2.8	4.2	8.8	20.6	40.9	64.6	102.6	160.2
1957	16.7	2.0	3.2	2.2	2.8	4.5	8.6	21.6	43.0	65.9	109.7	169.2
1956	16.3	2.6	3.3	2.4	2.7	4.2	8.6	20.8	41.3	64.3	107.2	177.3
1955	16.4	2.4	3.2	2.5	2.8	3.9	9.0	21.4	42.0	66.2	104.8	169.0
1954	16.5	2.4	3.3	2.3	2.9	4.1	9.2	21.7	42.2	65.7	108.5	166.3
1953	16.3	2.6	3.3	2.3	2.8	4.0	9.3	21.7	41.8	65.5	107.5	179.3
1952	16.6	2.5	3.2	2.2	3.0	4.3	9.0	22.1	42.0	67.1	112.9	186.9
1951	16.1	2.7	3.4	2.1	2.9	4.0	9.2	21.5	41.2	64.7	111.7	184.4
1950	14.7	2.5	3.2	2.3	2.8	3.8	8.6	19.3	37.9	60.3	97.7	181.3
MALE												
1959	17.7	2.2	3.1	2.7	3.4	4.8	10.1	23.8	47.0	74.3	114.8	181.7
1958	17.4	2.7	3.1	2.5	3.3	4.8	9.5	23.0	46.4	74.1	113.2	173.3
1957	18.0	2.1	3.6	2.5	3.3	5.0	9.0	24.0	48.9	77.5	118.9	179.8
1956	17.6	2.8	3.3	2.6	3.3	4.6	9.5	22.7	48.4	70.9	119.5	194.8
1955	17.8	2.7	3.3	2.7	3.5	4.3	9.2	23.9	48.2	75.0	117.0	181.1
1954	17.9	2.5	3.6	2.6	3.5	4.5	9.3	23.1	48.1	75.5	121.7	185.2
1953	17.6	2.8	3.5	2.3	3.3	4.3	10.0	24.1	46.7	72.6	118.7	196.9
1952	17.7	3.0	3.4	2.4	3.6	4.5	9.1	23.0	47.1	75.5	126.2	197.8
1951	17.0	2.9	3.7	2.3	3.5	4.3	9.0	22.4	44.5	70.9	121.5	220.3
1950	15.6	2.7	3.4	2.4	3.2	4.0	8.8	20.5	41.6	66.7	109.0	189.6
FEMALE												
1959	14.8	2.3	2.8	2.2	2.1	3.6	8.1	18.8	34.7	56.0	88.0	149.9
1958	15.0	1.7	3.1	2.1	2.2	3.6	8.1	18.3	35.7	56.2	94.2	151.6
1957	15.4	1.9	2.9	1.8	2.3	3.9	8.3	19.2	37.4	55.6	102.4	161.9
1956	15.1	2.5	3.2	2.3	2.3	3.8	7.8	19.0	34.6	58.3	97.3	165.1
1955	15.1	2.1	3.0	2.3	2.1	3.5	8.9	19.0	36.0	58.2	94.9	160.4
1954	15.1	2.2	3.1	2.0	2.4	3.7	9.1	20.3	36.4	56.8	97.7	152.7
1953	15.1	2.4	3.2	2.2	2.3	3.7	8.6	19.3	37.0	59.0	98.3	166.7
1952	15.5	2.0	3.0	2.0	2.4	4.0	9.0	21.2	37.0	59.5	101.9	179.0
1951	15.3	2.5	3.0	1.9	2.3	3.8	9.4	20.6	37.8	58.9	103.5	158.9
1950	13.8	2.3	3.0	2.1	2.4	3.5	8.4	18.1	34.3	54.4	88.1	175.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)												
WHITE, BOTH SEXES												
1959	16.7	2.3	3.2	2.5	2.7	4.3	9.1	20.9	40.1	64.6	101.4	167.8
1958	16.6	2.3	3.2	2.3	2.8	4.2	8.8	20.3	40.5	64.8	104.3	167.6
1957	17.2	2.2	3.3	2.2	2.8	4.6	8.6	21.1	42.5	66.2	111.8	177.0
1956	16.8	2.8	3.4	2.5	2.8	4.2	8.6	20.4	41.1	64.6	109.4	184.6
1955	16.9	2.5	3.3	2.6	2.9	4.0	9.0	20.9	41.5	66.6	107.1	174.3
1954	17.0	2.5	3.5	2.4	3.0	4.1	9.1	21.2	41.6	66.6	112.0	174.8
1953	16.9	2.7	3.6	2.4	2.8	4.1	9.1	21.4	41.6	66.2	110.4	186.7
1952	17.2	2.6	3.3	2.2	3.1	4.4	9.0	21.7	42.0	68.5	116.2	195.5
1951	16.6	2.9	3.4	2.2	3.0	4.1	9.1	20.9	40.9	65.8	115.2	192.3
1950	15.2	2.6	3.3	2.3	2.8	3.9	8.5	19.3	37.9	60.8	100.1	190.0
WHITE, MALE												
1959	18.3	2.0	3.4	2.7	3.4	5.0	10.2	23.8	46.7	74.5	117.0	187.7
1958	17.9	2.8	3.2	2.5	3.3	4.8	9.6	22.9	46.3	74.7	115.1	183.9
1957	18.6	2.3	3.5	2.5	3.4	5.2	9.0	23.7	48.9	78.3	122.0	189.1
1956	18.2	3.0	3.5	2.7	3.2	4.7	9.6	22.6	48.6	71.6	121.8	202.0
1955	18.3	2.8	3.5	2.8	3.6	4.5	9.3	23.7	47.8	75.6	119.8	183.7
1954	18.5	2.6	3.8	2.7	3.5	4.7	9.2	23.1	48.1	76.7	126.5	195.7
1953	18.2	2.9	3.7	2.4	3.3	4.5	10.1	23.9	46.4	73.3	123.2	206.0
1952	18.4	3.2	3.5	2.4	3.7	4.7	9.2	23.2	47.3	77.1	131.1	208.3
1951	17.6	3.0	3.8	2.4	3.7	4.4	9.2	22.0	44.6	72.3	125.8	230.4
1950	16.1	2.8	3.5	2.4	3.2	4.2	8.9	20.6	41.8	67.4	112.2	199.4
WHITE, FEMALE												
1959	15.2	2.5	3.0	2.3	2.1	3.7	8.0	18.2	33.9	56.0	89.5	155.0
1958	15.4	1.8	3.2	2.1	2.3	3.6	8.0	17.9	35.1	56.1	96.0	156.9
1957	15.8	2.1	3.1	1.9	2.3	4.0	8.2	18.6	36.4	55.6	103.7	168.7
1956	15.5	2.5	3.3	2.4	2.3	3.7	7.6	18.3	34.0	58.3	99.6	172.6
1955	15.5	2.2	3.1	2.4	2.2	3.5	8.6	18.1	35.4	58.6	96.9	167.7
1954	15.5	2.4	3.1	2.0	2.5	3.6	8.9	19.4	35.2	57.5	100.3	159.8
1953	15.7	2.6	3.4	2.3	2.4	3.8	8.2	18.9	37.0	59.8	99.9	172.8
1952	15.9	1.9	3.1	2.1	2.5	4.0	8.7	20.3	36.7	60.7	104.0	186.4
1951	15.7	2.7	3.0	2.0	2.3	3.9	9.0	19.9	37.2	59.9	106.4	165.1
1950	14.2	2.4	3.1	2.2	2.5	3.6	8.1	17.9	33.9	54.6	90.0	183.5

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)												
ALL OTHER, BOTH SEXES												
1959	12.4	*	1.8	1.8	2.9	3.3	8.8	24.3	46.4	63.7	76.1	97.1
1958	12.3	*	2.6	2.1	2.4	4.1	9.1	23.1	45.6	62.8	78.4	71.2
1957	12.7	*	2.6	1.8	2.3	3.6	9.2	25.9	49.0	61.8	80.7	73.4
1956	12.2	*	2.5	1.9	2.7	4.0	8.9	24.8	43.7	60.5	75.4	88.5
1955	12.6	*	2.2	1.8	2.4	3.5	9.8	26.9	47.7	60.5	72.3	103.4
1954	12.2	*	2.4	1.8	2.9	3.5	10.4	26.1	49.8	54.4	57.8	58.2
1953	11.7	*	2.0	1.7	2.7	3.0	10.5	24.2	43.9	56.7	66.2	88.5
1952	11.4	*	2.5	2.0	2.1	3.6	9.5	25.7	42.1	50.4	64.2	79.6
1951	11.5	*	3.0	1.4	2.2	3.2	9.9	26.6	44.5	50.5	59.9	89.4
1950	10.6	*	2.9	2.1	2.7	3.1	9.5	19.9	39.1	54.5	60.5	78.2
ALL OTHER, MALE												
1959	13.4	*	*	2.1	3.8	3.6	9.2	24.4	49.8	73.0	86.8	113.8
1958	13.0	*	2.5	2.3	3.0	4.4	9.0	24.2	48.5	67.7	89.9	*
1957	13.3	*	3.7	2.2	2.4	4.0	9.3	26.8	49.0	68.9	79.3	*
1956	12.6	*	2.1	2.0	3.7	3.7	8.4	23.3	46.8	62.5	90.0	108.0
1955	13.3	*	*	1.9	2.8	3.3	8.4	26.4	52.7	68.2	80.0	150.0
1954	12.2	*	2.3	1.9	3.9	3.0	10.1	22.9	48.2	62.2	58.8	*
1953	12.6	*	2.3	1.8	3.4	3.0	9.4	26.0	49.8	64.5	57.9	*
1952	11.2	*	2.8	2.4	2.9	3.1	7.4	21.4	44.0	56.4	59.1	*
1951	11.5	*	3.4	*	2.4	3.3	7.5	26.4	43.0	54.4	63.2	*
1950	10.9	*	3.0	2.6	3.3	2.8	8.0	19.4	39.6	57.7	64.4	*
ALL OTHER, FEMALE												
1959	11.5	*	1.9	1.6	2.1	3.0	8.5	24.2	43.1	55.4	66.5	84.6
1958	11.7	*	2.8	1.9	1.8	3.9	9.2	22.0	42.8	58.1	68.0	84.2
1957	12.1	*	*	1.5	2.1	3.3	9.1	25.1	48.9	55.4	82.1	78.4
1956	11.8	*	2.8	1.7	1.7	4.3	9.4	26.1	40.7	58.3	61.6	75.0
1955	11.9	*	2.5	1.7	2.0	3.6	11.0	27.3	42.8	53.4	64.4	70.6
1954	12.2	*	2.5	1.7	2.0	3.9	10.8	29.2	51.4	47.2	56.3	65.6
1953	10.9	*	*	1.6	2.1	2.9	11.5	22.5	37.9	49.3	73.6	93.5
1952	11.6	*	2.2	1.7	1.5	4.1	11.4	30.0	40.1	44.8	69.0	89.7
1951	11.6	*	2.7	1.7	1.9	3.1	12.0	26.7	46.0	46.8	56.8	85.7
1950	10.3	*	2.7	1.5	2.1	3.4	10.8	20.4	38.5	51.2	56.6	80.4

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
LEUKEMIA AND ALEUKEMIA (204)												
BOTH SEXES												
1959	7.0	2.9	5.2	3.2	2.0	2.2	3.3	6.1	12.8	27.4	42.2	44.3
1958	6.9	2.3	5.4	3.2	2.3	2.0	3.4	6.0	12.9	26.6	42.6	45.8
1957	6.9	2.7	5.6	3.1	2.2	2.3	3.4	6.3	12.9	26.2	41.6	38.9
1956	6.8	2.2	5.6	3.1	2.1	2.4	3.3	6.0	13.7	25.5	41.0	38.1
1955	6.6	2.7	5.4	3.0	2.2	2.2	3.4	6.0	12.9	25.1	37.6	39.1
1954	6.5	3.2	5.6	3.0	2.0	2.2	3.1	6.4	12.3	25.1	37.4	33.5
1953	6.3	3.6	5.3	2.8	2.0	2.1	3.3	6.1	12.6	23.5	35.6	34.0
1952	6.3	3.7	5.9	2.9	2.2	2.1	3.3	6.4	12.8	23.6	33.2	28.7
1951	6.1	3.2	6.0	2.9	2.2	2.3	3.4	6.1	12.8	21.5	30.8	30.5
1950	5.9	4.0	5.9	2.8	1.9	2.4	3.4	5.8	12.1	21.7	29.7	24.3
MALE												
1959	8.2	3.2	5.6	3.8	2.5	2.4	3.8	7.0	15.8	35.6	53.9	59.2
1958	8.1	2.3	5.9	3.4	2.8	2.3	3.9	6.8	15.5	34.4	56.3	71.8
1957	8.0	2.8	6.0	3.3	2.6	2.7	3.6	7.2	16.0	33.8	53.2	49.0
1956	7.9	1.7	5.8	3.5	2.5	2.8	3.6	6.8	16.2	33.0	52.3	51.7
1955	7.8	2.6	5.8	3.6	2.8	2.6	3.9	6.9	15.4	31.9	47.5	53.1
1954	7.5	3.4	5.7	3.2	2.4	2.6	3.4	7.1	14.8	31.7	48.9	48.4
1953	7.4	3.2	5.6	3.3	2.7	2.5	3.7	7.0	15.0	29.8	44.2	45.7
1952	7.3	3.1	6.4	3.2	2.8	2.3	3.6	6.9	15.1	30.1	42.1	35.5
1951	7.2	3.5	6.6	3.0	2.8	2.6	3.5	7.2	15.4	28.1	39.3	41.4
1950	6.8	4.7	6.3	3.1	2.1	2.6	3.5	6.5	14.1	27.9	36.4	33.4
FEMALE												
1959	5.7	2.5	4.8	2.6	1.5	2.0	2.8	5.2	9.9	20.2	33.1	34.6
1958	5.7	2.4	4.9	3.0	1.8	1.7	2.8	5.2	10.4	19.8	31.7	28.5
1957	5.8	2.6	5.1	2.9	1.8	1.9	3.2	5.3	9.9	19.4	32.4	32.0
1956	5.7	2.7	5.3	2.8	1.7	2.0	3.0	5.3	11.3	18.8	32.0	28.7
1955	5.4	2.9	5.0	2.3	1.6	1.9	3.0	5.0	10.6	18.9	29.5	29.2
1954	5.4	2.9	5.5	2.8	1.7	1.9	2.8	5.6	9.8	19.2	28.0	22.8
1953	5.2	4.1	5.0	2.3	1.4	1.8	2.9	5.2	10.3	17.8	28.5	25.6
1952	5.4	4.2	5.4	2.6	1.7	2.0	3.0	5.9	10.6	17.7	25.9	23.9
1951	5.0	2.9	5.3	2.7	1.6	2.0	3.3	5.1	10.2	15.5	23.7	22.8
1950	5.0	3.4	5.4	2.5	1.6	2.1	3.3	5.1	10.1	15.8	24.0	17.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
LEUKEMIA AND ALEUKEMIA (204)												
WHITE, BOTH SEXES												
1959	7.3	2.8	5.6	3.5	2.0	2.2	3.2	6.1	12.9	28.4	44.1	46.9
1958	7.3	2.4	5.8	3.4	2.3	1.9	3.4	6.0	13.1	27.6	44.7	48.0
1957	7.2	2.9	6.1	3.3	2.2	2.4	3.5	6.3	13.0	27.1	43.4	41.3
1956	7.1	2.1	6.0	3.3	2.1	2.4	3.3	6.0	13.9	26.4	42.8	40.1
1955	7.0	2.8	5.9	3.2	2.3	2.3	3.5	6.0	13.2	26.2	39.1	41.2
1954	6.8	3.1	6.1	3.2	2.0	2.3	3.1	6.4	12.5	26.1	39.2	35.0
1953	6.6	3.6	5.7	3.0	2.1	2.2	3.3	6.1	12.9	24.7	37.4	35.5
1952	6.6	3.5	6.3	3.1	2.4	2.2	3.3	6.5	13.1	24.7	34.8	30.1
1951	6.4	3.4	6.5	3.1	2.2	2.3	3.4	6.2	13.0	22.6	32.3	32.2
1950	6.2	4.0	6.4	3.0	1.9	2.5	3.4	6.0	12.4	22.6	31.2	26.1
WHITE, MALE												
1959	8.6	3.3	5.9	4.2	2.6	2.4	3.9	7.0	16.1	36.7	56.1	62.9
1958	8.6	2.4	6.4	3.7	2.8	2.1	4.0	6.9	15.8	35.6	59.2	75.9
1957	8.4	3.0	6.5	3.5	2.7	2.7	3.6	7.2	16.1	34.9	55.7	51.8
1956	8.3	1.7	6.1	3.7	2.6	2.8	3.6	6.8	16.4	34.2	54.6	53.6
1955	8.2	2.6	6.4	3.8	2.9	2.7	3.9	7.0	15.7	33.2	49.7	56.1
1954	7.9	3.2	6.2	3.4	2.4	2.5	3.4	7.1	15.0	32.7	51.5	49.6
1953	7.7	3.1	6.0	3.5	2.8	2.5	3.7	7.1	15.4	31.1	46.5	47.4
1952	7.7	2.8	6.8	3.4	3.0	2.4	3.7	7.1	15.4	31.3	44.1	37.2
1951	7.6	3.7	7.2	3.3	2.8	2.6	3.5	7.3	15.6	29.4	41.3	42.6
1950	7.1	4.6	6.8	3.4	2.2	2.7	3.5	6.7	14.4	29.1	38.4	36.2
WHITE, FEMALE												
1959	6.0	2.3	5.2	2.8	1.5	2.0	2.6	5.2	9.9	21.2	34.8	36.5
1958	6.0	2.4	5.3	3.1	1.8	1.7	2.8	5.1	10.5	20.4	33.4	29.5
1957	6.1	2.8	5.6	3.1	1.8	2.0	3.3	5.3	10.0	20.2	33.7	34.1
1956	6.0	2.5	5.9	2.9	1.7	2.0	2.9	5.3	11.5	19.5	33.5	30.6
1955	5.7	3.0	5.4	2.4	1.8	1.9	3.0	4.9	10.8	19.9	30.7	30.6
1954	5.8	3.0	6.0	3.0	1.7	2.0	2.8	5.6	10.0	20.1	29.3	24.4
1953	5.5	4.1	5.4	2.5	1.4	1.8	2.9	5.1	10.5	18.8	30.0	27.0
1952	5.6	4.2	5.8	2.7	1.8	2.0	2.9	5.8	10.9	18.6	27.1	25.0
1951	5.3	3.1	5.7	2.9	1.6	2.0	3.4	5.1	10.4	16.3	24.8	24.7
1950	5.3	3.4	5.9	2.7	1.6	2.2	3.3	5.2	10.3	16.5	25.2	19.1

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
LEUKEMIA AND ALEUKEMIA (204)												
ALL OTHER, BOTH SEXES												
1959	4.1	*	2.9	1.6	1.9	2.2	3.5	5.8	11.5	14.9	17.0	*
1958	4.1	*	2.5	2.0	2.0	2.9	3.4	5.9	10.6	15.8	12.3	*
1957	3.9	*	2.4	1.8	1.8	1.9	3.2	6.1	11.5	15.1	17.1	*
1956	4.1	*	2.8	1.9	1.6	2.4	3.3	6.4	11.8	14.2	15.7	*
1955	3.5	*	2.2	1.6	1.7	1.5	3.1	6.2	9.7	11.0	15.6	*
1954	3.8	*	2.6	1.5	2.0	2.1	3.1	6.1	9.9	13.4	11.5	*
1953	3.5	*	2.8	1.4	1.8	1.9	3.4	6.3	9.1	9.6	9.0	*
1952	3.6	4.7	3.0	1.7	1.4	2.0	3.3	5.7	9.1	11.3	10.2	*
1951	3.3	*	2.4	1.4	1.8	2.4	3.0	5.2	9.8	9.4	9.2	*
1950	3.1	*	2.4	1.6	1.4	1.6	3.4	4.7	9.0	9.1	*	*
ALL OTHER, MALE												
1959	4.6	*	3.2	1.7	1.8	2.5	2.7	6.3	12.3	22.5	26.4	*
1958	4.6	*	2.5	1.8	2.4	3.9	3.2	6.0	12.4	19.9	19.4	*
1957	4.9	*	2.9	2.1	2.3	2.4	3.7	7.1	14.6	21.6	21.5	*
1956	4.9	*	3.6	2.1	1.8	3.1	3.0	6.9	14.3	18.9	22.3	*
1955	4.3	*	2.3	2.1	2.7	1.8	3.4	6.4	12.0	15.7	19.2	*
1954	4.8	*	2.9	1.7	2.8	3.1	3.2	7.0	12.5	19.6	*	*
1953	4.1	*	2.8	1.9	2.6	2.1	3.8	5.9	10.4	13.7	*	*
1952	4.0	*	3.6	2.0	*	2.0	2.7	5.0	12.0	15.5	*	*
1951	4.0	*	*	1.5	2.4	2.6	3.8	5.6	12.6	12.8	*	*
1950	3.6	*	2.8	1.6	*	1.9	3.6	5.0	11.0	12.2	*	*
ALL OTHER, FEMALE												
1959	3.6	*	2.7	1.5	2.0	2.0	4.1	5.2	10.7	8.0	*	*
1958	3.6	*	2.5	2.1	1.7	2.0	3.5	5.8	8.8	12.0	*	*
1957	3.0	*	*	1.5	*	1.4	2.7	5.2	8.5	9.2	*	*
1956	3.4	*	2.1	1.7	*	1.8	3.6	6.0	9.4	9.9	*	*
1955	2.8	*	2.0	1.1	*	*	2.9	6.0	7.6	6.6	*	*
1954	2.8	*	2.3	1.4	*	*	3.0	5.2	7.2	7.6	*	*
1953	3.0	*	2.7	*	*	1.8	3.0	6.7	7.9	5.7	*	*
1952	3.2	*	2.4	1.5	*	2.0	3.8	6.4	6.1	7.3	*	*
1951	2.7	*	2.7	1.3	*	2.1	2.3	4.8	6.8	6.1	*	*
1950	2.6	*	*	1.6	*	*	3.2	4.4	6.9	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)												
BOTH SEXES												
1959	7.1	0.5	0.8	0.7	1.9	3.2	4.5	9.6	18.3	30.4	35.1	24.3
1958	6.9	0.6	0.7	0.7	1.8	3.1	4.6	9.3	19.0	28.9	32.0	22.8
1957	7.0	0.8	0.6	0.9	1.8	2.8	4.4	9.1	19.1	30.0	33.5	23.2
1956	6.8	1.2	0.7	0.9	1.8	3.0	4.6	8.9	18.9	28.4	32.5	22.7
1955	6.5	0.7	1.0	0.8	1.9	2.8	4.4	8.7	17.3	27.3	29.3	19.4
1954	6.3	0.9	0.7	0.8	1.8	3.0	4.4	8.7	17.1	25.6	27.8	20.9
1953	6.1	0.6	0.7	0.8	2.0	3.1	4.2	8.4	17.4	23.6	25.9	15.9
1952	5.8	0.7	0.8	0.8	1.8	3.1	4.0	8.4	16.3	22.9	23.5	16.5
1951	5.5	0.8	0.7	0.8	1.7	2.8	4.0	8.0	15.5	20.5	22.6	17.5
1950	5.2	*	0.7	0.8	1.6	2.8	3.8	7.7	14.9	20.2	20.4	13.2
MALE												
1959	8.4	*	0.9	1.1	2.5	4.0	5.7	11.7	22.0	36.6	42.2	33.2
1958	8.1	*	0.9	0.9	2.3	3.8	5.9	11.5	23.4	33.6	38.0	27.6
1957	8.4	*	0.9	1.2	2.5	3.7	5.7	11.4	23.7	36.9	40.2	26.7
1956	8.2	1.4	0.8	1.1	2.4	3.7	5.6	10.9	23.8	34.0	38.7	30.0
1955	7.8	*	1.3	1.2	2.5	3.5	5.7	10.9	21.3	33.0	34.4	23.3
1954	7.5	*	0.8	1.0	2.5	3.7	5.5	10.9	21.0	30.0	32.9	23.0
1953	7.5	*	0.8	1.2	2.4	4.0	5.5	10.6	21.3	28.5	31.8	19.0
1952	7.1	*	0.9	1.0	2.3	4.0	5.0	10.4	19.9	27.7	28.4	19.0
1951	6.7	*	0.8	1.0	2.4	3.5	5.1	9.9	18.7	25.1	28.9	22.3
1950	6.4	*	0.9	1.1	2.0	3.6	4.9	9.7	18.2	23.5	25.0	13.5
FEMALE												
1959	5.8	*	0.6	0.4	1.2	2.4	3.4	7.5	14.8	25.0	29.6	18.4
1958	5.7	*	0.6	0.5	1.3	2.4	3.4	7.1	14.8	24.8	27.2	19.6
1957	5.5	*	0.3	0.6	1.2	2.0	3.1	6.9	14.8	23.8	28.1	20.9
1956	5.5	*	0.5	0.6	1.3	2.3	3.6	6.9	14.2	23.4	27.5	17.7
1955	5.2	*	0.6	0.4	1.4	2.2	3.2	6.5	13.5	22.1	25.2	16.7
1954	5.1	*	0.5	0.5	1.1	2.3	3.4	6.5	13.2	21.5	23.6	19.3
1953	4.8	*	0.5	0.4	1.6	2.2	3.0	6.3	13.5	19.1	21.1	13.7
1952	4.7	*	0.8	0.5	1.4	2.2	3.1	6.4	12.8	18.5	19.5	14.7
1951	4.3	*	0.6	0.5	1.1	2.2	3.0	6.1	12.3	16.3	17.3	14.2
1950	4.1	*	0.6	0.4	1.2	2.1	2.7	5.7	11.5	17.1	16.6	12.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)												
WHITE, BOTH SEXES												
1959	7.3	0.6	0.8	0.8	1.9	3.2	4.6	9.6	18.4	30.9	36.4	24.7
1958	7.2	0.7	0.8	0.7	1.9	3.2	4.6	9.2	19.0	29.6	33.0	23.9
1957	7.2	0.9	0.6	0.9	1.9	2.9	4.4	9.2	19.0	30.8	34.6	24.1
1956	7.1	1.2	0.7	0.9	1.9	3.1	4.6	8.8	19.0	29.0	33.5	23.5
1955	6.7	0.7	1.0	0.8	2.0	2.9	4.3	8.7	17.5	28.1	30.5	20.6
1954	6.5	1.0	0.7	0.8	1.8	3.1	4.4	8.5	17.3	26.4	28.8	21.8
1953	6.4	*	0.8	0.9	2.0	3.1	4.3	8.4	17.6	24.6	26.9	16.9
1952	6.1	0.7	0.8	0.7	1.8	3.2	4.0	8.5	16.6	23.8	24.5	16.9
1951	5.8	0.9	0.7	0.8	1.8	2.9	4.1	8.2	15.7	21.4	23.6	18.8
1950	5.5	*	0.8	0.8	1.7	2.8	3.8	7.8	15.2	20.8	21.2	14.1
WHITE, MALE												
1959	8.6	*	1.0	1.1	2.6	4.0	5.7	11.6	22.0	36.9	43.6	33.1
1958	8.4	*	0.9	1.0	2.4	3.9	5.8	11.4	23.4	34.2	39.1	29.1
1957	8.7	*	0.9	1.3	2.6	3.8	5.6	11.6	23.5	37.9	41.9	28.0
1956	8.4	1.5	0.8	1.2	2.5	3.8	5.6	10.9	23.8	34.3	39.9	30.8
1955	8.1	*	1.4	1.2	2.5	3.5	5.7	11.0	21.5	33.7	35.9	24.5
1954	7.8	*	0.9	1.1	2.4	3.8	5.5	10.8	21.3	30.7	33.9	23.4
1953	7.8	*	0.9	1.3	2.4	4.0	5.6	10.4	21.8	29.6	32.9	19.8
1952	7.3	*	0.9	0.9	2.3	4.1	4.9	10.5	20.1	28.6	29.9	18.6
1951	7.0	*	0.8	1.1	2.4	3.5	5.3	10.2	18.9	26.2	30.4	24.1
1950	6.6	*	0.9	1.2	2.1	3.6	5.0	9.8	18.7	24.0	25.7	14.7
WHITE, FEMALE												
1959	6.1	*	0.7	0.4	1.2	2.4	3.5	7.5	15.0	25.6	30.9	19.2
1958	6.0	*	0.6	0.5	1.4	2.5	3.5	7.1	14.8	25.5	28.2	20.5
1957	5.8	*	0.3	0.6	1.3	2.0	3.2	6.9	14.9	24.5	28.9	21.4
1956	5.8	*	0.5	0.6	1.3	2.5	3.6	6.7	14.3	24.2	28.5	18.4
1955	5.4	*	0.6	0.4	1.5	2.3	3.1	6.4	13.7	23.0	26.2	17.8
1954	5.3	*	0.6	0.5	1.2	2.4	3.3	6.3	13.4	22.6	24.7	20.6
1953	5.0	*	0.6	0.5	1.7	2.2	3.1	6.4	13.6	20.0	22.0	14.8
1952	4.9	*	0.7	0.5	1.4	2.3	3.1	6.5	13.1	19.3	20.0	15.6
1951	4.5	*	0.5	0.5	1.2	2.3	3.0	6.2	12.5	17.1	17.8	15.1
1950	4.3	*	0.6	0.5	1.3	2.1	2.7	5.7	11.7	17.7	17.3	13.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)												
ALL OTHER, BOTH SEXES												
1959	4.8	*	*	0.5	1.6	2.9	4.1	9.8	17.6	24.5	17.7	*
1958	4.7	*	*	0.6	1.1	2.3	4.7	9.8	19.3	21.0	18.2	*
1957	4.6	*	*	0.7	1.2	2.4	4.4	8.1	20.4	20.3	17.1	*
1956	4.7	*	*	0.8	1.3	1.8	4.5	9.8	18.0	21.7	17.5	*
1955	4.3	*	*	0.7	1.5	2.3	5.0	8.8	15.2	17.6	11.3	*
1954	4.1	*	*	*	1.7	2.1	4.5	9.9	14.5	14.9	13.1	*
1953	3.8	*	*	0.6	1.5	2.8	3.8	8.8	14.3	11.5	11.1	*
1952	3.7	*	*	0.8	1.6	1.9	4.3	8.0	13.2	12.4	9.7	*
1951	3.2	*	*	*	1.4	2.4	3.4	6.2	13.5	9.7	*	*
1950	3.4	*	*	0.7	1.2	2.7	3.6	7.2	11.2	13.1	*	*
ALL OTHER, MALE												
1959	6.2	*	*	*	1.9	4.0	5.4	12.4	22.1	32.8	24.3	*
1958	5.9	*	*	*	*	2.6	6.7	12.4	23.5	26.6	24.5	*
1957	5.8	*	*	*	1.9	2.8	6.7	9.3	26.5	25.1	17.8	*
1956	6.0	*	*	*	*	3.0	5.4	11.1	23.2	30.7	22.3	*
1955	5.5	*	*	*	2.4	3.4	5.8	10.4	19.1	24.0	*	*
1954	5.3	*	*	*	3.1	2.9	5.0	11.3	18.1	22.2	19.3	*
1953	5.0	*	*	*	2.0	3.8	5.2	12.5	16.3	15.1	*	*
1952	4.8	*	*	*	2.2	2.5	5.2	10.1	17.6	17.0	*	*
1951	4.0	*	*	*	2.0	3.2	3.9	7.1	17.0	12.8	*	*
1950	4.2	*	*	*	*	3.3	4.3	9.0	12.8	17.0	*	*
ALL OTHER, FEMALE												
1959	3.5	*	*	*	*	2.0	3.0	7.5	13.2	16.9	*	*
1958	3.6	*	*	*	*	2.0	2.8	7.2	15.4	15.8	*	*
1957	3.4	*	*	*	*	2.0	2.3	7.0	14.4	15.9	16.6	*
1956	3.5	*	*	*	*	*	3.7	8.6	13.0	13.3	*	*
1955	3.1	*	*	*	*	*	4.2	7.2	11.3	11.7	*	*
1954	2.9	*	*	*	*	1.5	4.1	8.5	10.9	8.1	*	*
1953	2.6	*	*	*	*	1.9	2.5	5.2	12.2	8.1	*	*
1952	2.7	*	*	*	*	*	3.4	5.9	8.6	8.1	*	*
1951	2.5	*	*	*	*	1.7	2.8	5.4	9.9	6.6	*	*
1950	2.6	*	*	*	*	2.1	3.0	5.4	9.5	9.1	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)												
BOTH SEXES												
1959	2.8	3.1	0.8	0.7	0.8	1.2	2.4	4.6	6.0	8.4	10.9	20.4
1958	2.9	3.3	0.8	0.6	0.8	1.4	2.7	4.5	6.1	8.5	11.1	19.5
1957	3.0	3.3	0.8	0.6	0.8	1.4	2.8	4.7	6.4	8.4	12.3	21.5
1956	2.9	4.1	0.9	0.6	0.9	1.4	2.7	4.7	6.4	7.9	11.7	18.9
1955	3.1	4.3	1.0	0.7	1.0	1.5	3.1	5.0	6.4	8.7	12.8	19.8
1954	3.2	4.6	1.0	0.7	0.8	1.6	3.1	5.3	6.6	7.9	12.8	18.0
1953	3.3	4.5	1.0	0.8	0.8	1.7	3.5	5.6	6.8	8.0	13.4	20.0
1952	3.3	4.4	1.1	0.7	1.0	1.7	3.9	5.4	6.9	7.7	11.5	21.1
1951	3.4	4.5	1.0	0.7	0.9	1.8	3.8	5.9	7.0	8.2	11.9	23.4
1950	3.6	4.4	1.1	0.8	0.9	1.7	4.1	5.9	7.9	9.2	13.5	21.8
MALE												
1959	2.5	3.0	0.9	0.8	0.7	1.0	1.6	3.5	6.1	9.1	10.6	15.8
1958	2.6	2.5	0.9	0.6	0.8	1.1	2.0	4.1	6.5	8.8	11.3	16.6
1957	2.7	3.8	0.8	0.7	0.9	1.1	2.0	4.2	6.4	9.0	11.8	18.4
1956	2.6	4.6	0.8	0.7	0.8	1.0	1.7	3.9	6.5	8.4	11.2	15.3
1955	2.7	4.5	0.9	0.6	0.9	1.1	1.9	4.1	6.5	8.9	13.0	17.0
1954	2.7	4.8	1.1	0.7	1.0	1.3	2.0	4.0	6.6	8.2	11.9	12.8
1953	2.9	5.0	1.1	0.8	0.8	1.3	2.0	4.0	7.1	8.6	13.2	19.7
1952	2.7	4.7	1.1	0.7	0.9	1.3	2.3	4.1	6.3	7.1	10.2	19.0
1951	2.8	4.1	1.0	0.7	0.9	1.2	2.2	4.1	6.7	8.4	11.5	14.1
1950	2.8	4.6	1.2	0.8	0.9	1.1	2.2	4.0	7.2	9.1	11.5	15.6
FEMALE												
1959	3.1	3.2	0.6	0.6	0.8	1.5	3.0	5.6	5.9	7.9	11.2	23.4
1958	3.1	4.2	0.8	0.7	0.7	1.7	3.3	4.8	5.7	8.2	11.0	21.5
1957	3.2	2.9	0.8	0.6	0.8	1.7	3.5	5.3	6.3	7.8	12.8	23.7
1956	3.2	3.6	0.9	0.6	1.0	1.7	3.8	5.4	6.4	7.4	12.1	21.5
1955	3.5	4.2	1.1	0.8	1.0	1.9	4.2	5.9	6.3	8.5	12.6	21.8
1954	3.6	4.4	1.0	0.7	0.7	2.0	4.3	6.6	6.5	7.7	13.6	21.6
1953	3.8	3.9	1.0	0.9	0.9	2.1	5.1	7.2	6.6	7.4	13.6	20.1
1952	3.9	4.0	1.1	0.7	1.0	2.0	5.5	6.8	7.5	8.2	12.5	22.6
1951	4.0	4.8	0.9	0.7	0.9	2.4	5.3	7.6	7.2	8.1	12.2	30.0
1950	4.3	4.1	1.1	0.7	0.9	2.3	5.9	7.7	8.6	9.3	15.1	26.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)												
WHITE, BOTH SEXES												
1959	2.7	3.1	0.8	0.7	0.7	1.1	2.0	4.2	5.8	8.3	10.9	20.4
1958	2.8	3.4	0.8	0.6	0.7	1.2	2.3	4.2	5.8	8.4	11.1	19.5
1957	2.9	3.4	0.8	0.6	0.7	1.3	2.4	4.3	6.0	8.4	12.5	22.0
1956	2.8	4.0	0.8	0.6	0.8	1.2	2.3	4.3	6.2	7.7	11.7	19.2
1955	3.0	3.9	1.0	0.7	0.9	1.2	2.5	4.6	6.2	8.5	12.6	20.3
1954	3.0	4.4	1.0	0.7	0.8	1.4	2.7	4.8	6.3	7.9	12.7	18.1
1953	3.2	4.4	0.9	0.8	0.8	1.5	2.9	5.1	6.7	8.0	13.6	20.7
1952	3.1	4.3	1.1	0.7	0.9	1.4	3.2	4.8	6.6	7.6	11.6	21.7
1951	3.2	4.7	1.0	0.6	0.8	1.5	3.1	5.4	6.7	8.3	11.9	24.3
1950	3.4	4.5	1.1	0.8	0.9	1.5	3.3	5.4	7.6	8.9	13.8	22.7
WHITE, MALE												
1959	2.5	3.2	0.9	0.8	0.7	1.0	1.6	3.3	5.9	9.0	10.8	15.0
1958	2.6	2.5	0.8	0.5	0.9	1.0	1.9	4.0	6.3	8.8	11.2	17.4
1957	2.7	3.7	0.8	0.7	0.8	1.1	2.0	4.1	6.1	9.1	12.1	19.0
1956	2.6	4.6	0.8	0.7	0.7	1.1	1.6	3.9	6.3	8.3	11.5	15.2
1955	2.7	4.4	0.9	0.6	0.9	1.0	1.9	4.0	6.5	8.7	12.9	18.4
1954	2.7	4.7	1.0	0.7	0.9	1.2	1.9	3.8	6.5	8.3	11.8	12.4
1953	2.9	4.9	0.9	0.8	0.8	1.3	1.8	3.8	7.2	8.6	13.3	21.3
1952	2.7	4.6	1.1	0.7	0.8	1.3	2.2	3.9	6.3	7.3	10.4	19.4
1951	2.8	4.3	1.0	0.6	0.8	1.1	2.1	4.0	6.6	8.7	11.7	14.3
1950	2.9	4.7	1.1	0.9	0.8	1.1	2.2	3.9	7.1	9.1	12.2	15.6
WHITE, FEMALE												
1959	2.9	3.1	0.6	0.6	0.7	1.3	2.3	5.0	5.6	7.7	11.0	23.8
1958	2.9	4.2	0.8	0.7	0.6	1.4	2.6	4.4	5.3	8.0	11.1	20.9
1957	3.0	3.0	0.7	0.6	0.7	1.4	2.8	4.5	5.9	7.7	12.9	24.1
1956	3.0	3.5	0.9	0.5	0.9	1.3	3.0	4.6	6.1	7.3	11.9	21.9
1955	3.2	3.5	1.1	0.7	1.0	1.5	3.1	5.1	5.9	8.3	12.4	21.7
1954	3.3	4.2	0.9	0.7	0.7	1.6	3.4	5.8	6.2	7.5	13.4	22.1
1953	3.5	3.8	0.9	0.9	0.8	1.7	4.0	6.4	6.2	7.4	13.7	20.2
1952	3.4	3.9	1.1	0.7	0.9	1.5	4.1	5.7	7.0	7.8	12.5	23.3
1951	3.6	5.0	1.0	0.6	0.9	1.9	4.1	6.8	6.7	8.0	12.0	31.3
1950	3.9	4.2	1.1	0.7	0.9	1.8	4.5	6.9	8.1	8.8	15.2	27.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)												
ALL OTHER, BOTH SEXES												
1959	3.6	*	*	0.9	1.1	2.2	5.8	8.4	8.4	9.6	11.1	*
1958	3.7	*	1.0	0.7	1.2	2.6	6.3	7.2	9.2	9.6	10.6	*
1957	4.0	*	*	0.9	1.4	2.8	6.5	8.9	10.7	8.4	9.6	*
1956	3.9	4.4	1.1	0.6	1.3	3.0	6.4	8.3	8.8	9.2	11.9	*
1955	4.5	6.9	1.1	0.9	1.2	3.5	8.3	9.4	8.9	11.1	14.8	*
1954	4.5	5.6	1.4	1.0	1.0	3.3	7.3	10.7	9.6	8.6	15.2	*
1953	4.6	5.3	1.9	0.9	1.1	3.4	8.9	10.3	8.8	8.0	11.1	*
1952	5.2	4.9	1.2	0.8	1.7	3.9	10.8	11.6	10.7	8.8	10.2	*
1951	4.8	*	*	1.0	1.6	4.1	9.5	10.2	10.8	7.0	12.0	*
1950	5.1	*	*	*	1.2	4.0	10.5	10.6	11.1	13.1	*	*
ALL OTHER, MALE												
1959	2.6	*	*	*	*	*	2.3	5.3	7.5	9.6	*	*
1958	2.7	*	*	*	*	*	3.4	5.3	8.3	8.8	*	*
1957	2.9	*	*	*	1.7	*	2.9	5.2	10.1	7.7	*	*
1956	2.5	*	*	*	*	*	2.4	3.9	8.1	9.4	*	*
1955	2.8	*	*	*	*	*	2.3	5.0	6.8	11.3	*	*
1954	3.1	*	*	*	*	1.7	2.8	6.6	8.4	7.1	*	*
1953	2.9	*	*	*	*	*	3.2	6.2	5.9	8.7	*	*
1952	2.9	*	*	*	1.7	*	3.2	6.4	7.1	*	*	*
1951	2.8	*	*	*	1.9	1.9	3.2	5.0	7.5	*	*	*
1950	2.6	*	*	*	*	*	2.1	5.6	7.5	9.4	*	*
ALL OTHER, FEMALE												
1959	4.5	*	*	0.9	*	2.8	9.0	11.4	9.2	9.6	13.7	*
1958	4.5	*	*	*	1.7	3.5	9.0	9.0	10.1	10.4	*	*
1957	5.0	*	*	1.0	*	3.9	9.8	12.4	11.2	9.2	*	*
1956	5.2	*	*	*	1.7	4.8	10.2	12.6	9.6	8.9	15.9	*
1955	6.2	8.4	*	*	1.5	5.2	13.8	13.5	11.0	10.9	15.2	*
1954	5.8	*	*	*	*	4.7	11.5	14.6	10.8	10.0	16.7	*
1953	6.1	*	*	*	*	5.1	14.1	14.3	11.6	7.3	*	*
1952	7.4	*	*	*	1.7	5.9	17.8	16.6	14.3	12.6	*	*
1951	6.7	*	*	*	*	6.1	15.3	15.3	14.2	8.7	*	*
1950	7.5	*	*	*	*	6.3	18.4	15.6	14.7	16.8	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ASTHMA (241)												
BOTH SEXES												
1959	2.8	0.8	0.4	0.2	0.3	0.8	1.6	3.6	7.6	12.9	17.0	17.3
1958	2.9	0.8	0.4	0.2	0.3	0.7	1.4	3.5	7.9	13.5	18.6	21.3
1957	3.9	1.3	0.6	0.2	0.4	0.8	1.7	4.2	10.5	19.4	28.2	27.8
1956	3.6	1.6	0.6	0.1	0.4	0.6	1.2	3.6	9.4	19.0	25.4	29.4
1955	3.6	1.3	0.6	0.2	0.4	0.7	1.5	3.7	9.8	18.5	25.6	31.7
1954	3.8	1.7	0.6	0.1	0.4	0.7	1.6	4.1	10.3	19.3	27.5	30.7
1953	4.3	2.3	0.6	0.2	0.3	0.7	1.7	4.4	12.0	22.3	30.2	34.0
1952	4.5	2.7	0.7	0.2	0.4	0.8	1.8	4.9	12.7	23.2	30.3	35.6
1951	4.5	1.9	0.7	0.1	0.4	0.8	1.7	5.2	12.7	22.9	32.9	43.7
1950	2.9	2.3	0.8	0.2	0.3	0.7	1.4	3.9	8.1	12.6	18.1	25.1
MALE												
1959	3.8	1.1	0.3	0.2	0.2	0.6	1.4	4.4	11.6	20.3	26.1	23.1
1958	4.0	1.0	0.4	0.2	0.2	0.6	1.2	4.2	12.1	22.2	29.9	28.2
1957	5.4	1.5	0.6	0.2	0.3	0.6	1.6	5.5	16.3	31.1	42.5	34.7
1956	4.9	2.0	0.6	*	0.3	0.4	1.1	4.7	14.5	30.9	36.6	31.2
1955	4.9	1.3	0.6	0.2	0.3	0.5	1.4	4.7	14.7	29.4	37.3	42.5
1954	5.1	2.0	0.7	*	0.3	0.6	1.5	5.1	15.5	29.8	38.0	39.5
1953	5.7	2.5	0.5	0.2	0.2	0.5	1.9	5.4	18.0	33.5	43.0	43.9
1952	6.0	3.2	0.8	*	0.3	0.5	1.7	6.5	18.9	35.3	43.0	39.2
1951	6.0	2.2	0.8	0.2	0.4	0.6	1.5	6.6	19.4	34.0	44.8	50.8
1950	3.6	2.6	0.9	0.2	0.2	0.4	1.4	4.9	11.2	18.2	24.8	31.2
FEMALE												
1959	1.9	*	0.5	0.2	0.4	0.9	1.7	2.7	3.9	6.4	9.9	13.4
1958	1.9	*	0.4	0.2	0.5	0.8	1.5	2.9	4.0	5.8	9.8	16.7
1957	2.5	1.1	0.5	0.2	0.5	0.9	1.7	3.0	5.1	9.1	16.7	23.1
1956	2.3	1.1	0.5	0.1	0.4	0.8	1.3	2.6	4.6	8.4	16.5	28.1
1955	2.4	1.4	0.6	0.2	0.5	0.9	1.5	2.6	5.1	8.7	16.2	24.1
1954	2.5	1.4	0.6	*	0.5	0.8	1.6	3.0	5.2	9.8	19.0	24.5
1953	2.8	2.0	0.6	*	0.4	0.9	1.6	3.4	6.1	12.1	19.7	26.9
1952	3.0	2.1	0.5	0.2	0.5	1.1	1.9	3.4	6.7	12.0	19.7	33.1
1951	3.0	1.7	0.7	*	0.4	1.0	1.8	3.7	6.1	12.7	22.9	38.6
1950	2.1	2.1	0.7	*	0.5	1.0	1.4	3.0	5.0	7.5	12.4	20.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ASTHMA (241)												
WHITE, BOTH SEXES												
1959	2.8	0.7	0.3	0.1	0.2	0.5	1.2	3.3	7.4	13.3	17.4	17.6
1958	2.9	*	0.3	0.1	0.3	0.5	1.0	3.2	7.9	13.6	19.3	22.4
1957	3.9	1.1	0.5	0.1	0.3	0.5	1.3	3.8	10.2	19.7	29.0	28.9
1956	3.6	1.3	0.5	0.1	0.3	0.5	1.0	3.3	9.3	19.4	26.1	31.1
1955	3.7	1.2	0.5	0.1	0.3	0.5	1.2	3.5	9.8	19.1	26.6	32.3
1954	3.9	1.3	0.6	0.1	0.4	0.6	1.4	3.9	10.3	19.9	28.1	32.1
1953	4.4	1.9	0.5	0.2	0.3	0.6	1.6	4.2	11.9	23.2	31.3	36.0
1952	4.5	2.6	0.6	0.2	0.4	0.6	1.6	4.6	12.7	23.7	31.1	35.5
1951	4.6	1.9	0.6	0.1	0.3	0.6	1.5	4.9	12.7	23.7	34.1	45.2
1950	2.9	1.9	0.7	0.1	0.3	0.6	1.2	3.9	8.1	12.9	18.5	25.6
WHITE, MALE												
1959	3.8	*	*	0.1	*	0.4	1.0	4.0	11.5	20.9	27.3	24.2
1958	4.0	*	0.3	0.1	*	0.4	1.0	3.9	12.1	22.4	31.4	29.7
1957	5.4	1.2	0.4	0.2	0.2	0.4	1.3	4.9	16.1	31.6	43.6	36.7
1956	5.0	1.7	0.5	*	0.3	0.3	0.9	4.3	14.4	31.8	38.0	33.1
1955	5.1	1.2	0.5	*	*	0.3	1.2	4.6	14.8	30.3	38.7	42.5
1954	5.3	1.5	0.6	*	0.3	0.5	1.3	4.9	15.6	30.9	39.0	41.8
1953	5.9	2.1	0.5	0.2	*	0.4	1.7	5.2	18.1	34.9	44.9	46.3
1952	6.2	3.0	0.7	*	*	0.4	1.5	6.2	19.1	36.4	44.7	41.1
1951	6.2	2.2	0.7	*	0.3	0.5	1.4	6.4	19.8	35.4	46.6	54.0
1950	3.7	2.1	0.8	*	*	0.4	1.1	4.8	11.4	18.6	25.3	32.1
WHITE, FEMALE												
1959	1.8	*	0.4	*	0.2	0.6	1.3	2.5	3.5	6.6	9.8	13.3
1958	1.7	*	0.3	*	0.4	0.5	1.0	2.5	3.9	5.8	9.9	17.6
1957	2.4	*	0.5	*	0.3	0.6	1.4	2.6	4.7	9.1	17.5	23.6
1956	2.2	*	0.4	*	0.4	0.6	1.1	2.3	4.5	8.3	16.8	29.7
1955	2.4	1.3	0.6	*	0.4	0.7	1.2	2.5	5.0	9.0	16.9	25.1
1954	2.5	*	0.5	*	0.4	0.7	1.5	2.8	5.1	10.0	19.4	25.2
1953	2.8	1.7	0.5	*	0.4	0.7	1.4	3.2	5.8	12.5	20.3	28.6
1952	2.9	2.2	0.5	0.2	0.5	0.9	1.6	3.1	6.4	12.1	19.9	31.5
1951	3.0	1.6	0.6	*	0.3	0.8	1.6	3.4	5.8	13.0	23.8	38.9
1950	2.1	1.7	0.6	*	0.4	0.8	1.2	2.9	4.8	7.7	12.7	21.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ASTHMA (241)												
ALL OTHER, BOTH SEXES												
1959	3.4	*	1.0	0.6	1.1	2.7	5.1	6.3	10.2	8.9	11.1	*
1958	3.3	*	*	0.5	0.9	2.2	4.7	7.3	8.7	12.9	8.9	*
1957	4.3	*	1.3	0.5	1.2	2.7	4.6	8.6	14.2	16.4	16.4	*
1956	3.3	*	1.1	*	0.8	1.7	3.0	7.2	10.8	14.3	15.3	*
1955	3.1	*	*	*	1.2	2.3	3.6	5.2	10.1	12.1	12.1	*
1954	3.2	4.4	1.1	*	0.9	1.8	3.3	6.0	10.8	11.9	18.4	*
1953	3.5	4.9	1.2	*	0.8	2.0	3.1	6.8	13.4	12.2	14.1	*
1952	3.9	*	1.2	*	0.9	2.2	3.8	7.7	12.8	16.6	18.1	*
1951	3.5	*	1.6	*	0.8	1.9	3.1	7.5	12.0	13.5	14.7	*
1950	2.7	4.9	1.5	*	*	1.8	3.3	4.3	7.6	8.9	12.3	*
ALL OTHER, MALE												
1959	3.8	*	*	*	*	2.7	4.6	8.0	12.3	13.6	*	*
1958	3.6	*	*	*	*	1.8	3.9	7.1	11.8	19.6	*	*
1957	5.2	*	1.9	*	*	1.8	4.7	10.6	18.8	24.5	28.1	*
1956	4.2	*	*	*	*	1.8	2.9	9.0	15.8	19.7	19.2	*
1955	3.8	*	*	*	*	1.8	3.0	5.9	13.9	19.3	18.4	*
1954	3.9	*	*	*	*	1.7	3.6	7.1	14.3	17.0	24.4	*
1953	4.0	*	*	*	*	*	3.2	7.6	16.7	16.9	*	*
1952	4.5	*	*	*	*	2.0	3.2	9.5	15.5	22.7	20.0	*
1951	3.9	*	*	*	*	*	3.1	8.5	14.2	17.7	20.8	*
1950	3.0	*	*	*	*	*	3.4	5.3	8.5	12.9	*	*
ALL OTHER, FEMALE												
1959	3.0	*	*	*	1.6	2.7	5.4	4.7	8.1	4.6	*	*
1958	3.0	*	*	*	*	2.5	5.5	7.4	5.6	6.8	*	*
1957	3.4	*	*	*	1.5	3.5	4.6	6.7	9.7	8.9	*	*
1956	2.6	*	*	*	*	1.7	3.1	5.4	5.9	9.4	*	*
1955	2.5	*	*	*	*	2.7	4.1	4.4	6.4	5.3	*	*
1954	2.6	*	*	*	*	1.8	3.0	4.9	7.2	7.1	*	*
1953	2.9	*	*	*	*	2.4	3.0	6.0	10.1	7.9	*	*
1952	3.4	*	*	*	*	2.4	4.3	6.0	10.0	10.9	*	*
1951	3.2	*	*	*	*	2.3	3.2	6.6	9.7	9.5	*	*
1950	2.3	*	*	*	*	2.3	3.2	3.4	6.7	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
DIABETES (260)												
BOTH SEXES												
1959	15.9	*	0.2	0.4	0.9	2.6	3.8	11.4	37.5	89.9	154.5	158.8
1958	15.9	*	0.3	0.3	0.9	2.5	4.1	11.2	37.0	91.6	151.7	169.9
1957	16.0	*	0.3	0.4	1.2	2.6	4.5	11.2	38.0	92.5	149.8	158.1
1956	15.7	*	0.2	0.3	1.0	2.4	3.8	11.0	37.5	92.1	151.7	162.6
1955	15.5	0.6	0.3	0.4	1.0	2.4	3.9	10.9	37.3	91.7	150.6	163.1
1954	15.6	*	0.3	0.4	1.0	2.2	3.9	11.3	38.3	92.3	153.9	155.3
1953	16.3	*	0.3	0.4	1.2	2.4	4.1	11.7	41.7	96.2	161.7	156.4
1952	16.4	0.8	0.5	0.6	1.3	2.3	4.0	12.3	40.8	99.7	161.5	152.8
1951	16.3	*	0.4	0.5	1.2	2.2	3.9	12.4	42.7	99.8	161.2	151.1
1950	16.2	0.7	0.3	0.6	1.1	2.2	4.2	12.4	42.1	101.2	166.7	150.3
MALE												
1959	12.9	*	*	0.3	0.8	3.0	4.2	11.4	31.0	71.7	133.5	153.5
1958	12.8	*	0.4	0.3	0.9	2.9	4.8	10.6	30.1	73.1	129.0	163.4
1957	12.6	*	0.3	0.4	1.0	3.0	5.0	10.4	29.3	71.9	127.8	145.1
1956	12.5	*	*	0.2	0.8	2.7	4.2	9.9	30.6	71.5	129.8	150.5
1955	12.4	*	0.3	0.4	0.9	2.6	4.4	9.9	29.8	70.7	131.2	143.4
1954	12.4	*	0.3	0.4	1.0	2.5	4.5	10.3	29.4	70.0	131.0	143.1
1953	12.7	*	0.3	0.3	1.1	2.8	4.5	9.9	30.6	71.9	136.1	151.6
1952	13.0	*	0.5	0.4	1.1	2.3	4.3	10.7	30.7	75.8	140.9	149.5
1951	12.5	*	0.4	0.3	1.0	2.2	4.0	10.8	32.1	74.6	129.4	138.3
1950	12.5	*	*	0.5	0.9	2.1	4.3	10.2	30.3	76.3	141.9	140.6
FEMALE												
1959	18.8	*	0.3	0.5	1.0	2.2	3.5	11.5	43.6	105.8	170.9	162.2
1958	18.8	*	0.3	0.4	0.9	2.1	3.6	11.9	43.5	107.8	169.6	174.2
1957	19.2	*	0.3	0.6	1.3	2.3	4.1	12.0	46.2	110.8	167.2	167.0
1956	18.7	*	*	0.4	1.2	2.0	3.5	12.1	44.0	110.5	169.3	171.1
1955	18.5	*	*	0.4	1.1	2.2	3.3	11.9	44.5	110.5	166.3	177.1
1954	18.7	*	0.3	0.5	1.1	1.9	3.4	12.3	46.9	112.5	172.5	164.0
1953	19.8	*	*	0.5	1.2	2.0	3.7	13.4	52.5	118.3	182.6	160.0
1952	19.7	*	0.4	0.7	1.4	2.2	3.7	13.8	50.7	121.7	178.6	155.1
1951	20.0	*	0.5	0.7	1.4	2.2	3.9	14.1	53.3	122.9	187.9	160.3
1950	19.9	*	0.4	0.7	1.4	2.2	4.1	14.5	54.1	124.4	187.9	157.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
DIABETES (260)												
WHITE, BOTH SEXES												
1959	15.8	*	0.2	0.4	0.8	2.4	3.3	9.4	33.9	88.6	156.3	162.5
1958	15.8	*	0.3	0.3	0.8	2.3	3.6	9.4	34.1	90.4	153.9	173.6
1957	16.0	*	0.3	0.4	1.0	2.5	3.9	9.2	35.3	92.3	153.1	163.4
1956	15.8	*	0.2	0.3	0.9	2.2	3.4	9.4	34.7	92.2	155.5	169.0
1955	15.7	*	0.3	0.4	1.0	2.3	3.3	9.2	35.2	91.9	154.1	169.7
1954	15.8	*	0.3	0.4	0.9	2.1	3.4	9.4	36.2	93.1	158.6	159.6
1953	16.5	*	0.3	0.4	1.1	2.2	3.4	9.7	39.8	97.3	165.6	160.3
1952	16.6	0.8	0.5	0.5	1.2	2.1	3.4	10.6	39.2	101.0	165.8	157.7
1951	16.6	*	0.4	0.5	1.1	2.1	3.2	10.6	41.4	101.2	165.8	155.9
1950	16.4	0.8	0.3	0.5	1.0	2.0	3.3	10.5	41.0	102.0	171.3	154.9
WHITE, MALE												
1959	13.0	*	*	0.3	0.8	2.8	4.1	10.2	29.2	70.9	135.8	161.7
1958	13.0	*	0.4	0.2	0.8	2.8	4.4	9.8	29.3	72.9	132.1	165.2
1957	12.8	*	0.3	0.3	0.9	2.8	4.7	9.6	28.4	71.7	132.0	148.6
1956	12.8	*	*	0.2	0.7	2.6	4.0	9.4	29.3	72.0	133.9	157.9
1955	12.8	*	0.3	0.3	0.9	2.5	4.1	9.4	29.1	71.7	134.5	150.3
1954	12.7	*	*	0.3	0.9	2.5	4.3	9.6	28.6	70.9	135.4	147.2
1953	13.0	*	0.4	0.3	1.1	2.6	4.2	9.3	30.0	73.5	139.4	155.2
1952	13.3	*	0.6	0.4	1.0	2.2	4.1	10.2	30.2	76.8	144.7	154.5
1951	12.8	*	0.4	0.3	0.9	2.2	3.7	10.1	31.4	75.5	133.6	141.8
1950	12.8	*	*	0.5	0.8	2.0	3.9	9.4	30.0	77.0	145.7	147.6
WHITE, FEMALE												
1959	18.5	*	*	0.4	0.9	2.0	2.5	8.7	38.2	104.1	172.3	163.1
1958	18.5	*	0.3	0.3	0.8	1.8	2.8	9.0	38.6	105.7	170.9	179.1
1957	19.0	*	*	0.5	1.2	2.1	3.1	8.9	41.8	110.5	169.6	173.5
1956	18.6	*	*	0.3	1.0	1.8	2.8	9.3	39.7	110.3	172.6	176.7
1955	18.5	*	*	0.4	1.1	2.1	2.5	9.0	41.1	110.1	169.8	183.4
1954	18.8	*	*	0.5	1.0	1.8	2.6	9.2	43.6	113.2	177.2	168.4
1953	19.9	*	*	0.6	1.2	1.8	2.7	10.0	49.5	119.1	186.8	163.9
1952	19.9	*	0.4	0.6	1.3	2.1	2.8	11.0	48.0	123.2	183.0	159.9
1951	20.3	*	0.5	0.7	1.3	2.0	2.8	11.2	51.3	124.9	192.6	166.0
1950	20.0	*	*	0.6	1.2	1.9	2.8	11.5	52.0	125.2	192.7	159.9

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
DIABETES (260)												
ALL OTHER, BOTH SEXES												
1959	16.8	*	*	*	1.1	3.7	8.6	30.3	75.2	105.4	129.5	113.2
1958	16.3	*	*	0.6	1.6	3.9	8.9	29.0	68.0	106.5	121.2	125.8
1957	15.8	*	*	0.6	2.0	4.0	10.5	29.8	67.5	95.3	102.9	92.2
1956	14.7	*	*	*	1.9	3.6	7.8	27.0	69.1	90.8	97.8	85.2
1955	14.3	*	*	*	1.5	3.1	8.9	27.2	61.4	88.3	101.2	82.8
1954	14.1	*	*	0.6	1.6	3.0	8.4	29.5	62.6	82.5	86.5	100.0
1953	14.7	*	*	*	1.7	3.8	9.5	30.8	63.8	82.0	104.7	109.6
1952	14.1	*	*	0.9	2.1	3.2	8.6	28.3	60.9	84.2	98.7	91.8
1951	14.1	*	*	0.7	2.2	3.2	9.9	29.6	59.8	82.6	93.5	93.6
1950	14.4	*	*	0.7	2.2	3.6	11.5	30.5	56.4	90.3	98.4	96.0
ALL OTHER, MALE												
1959	12.1	*	*	*	*	4.1	5.7	22.4	48.6	81.6	104.9	*
1958	11.2	*	*	*	1.5	3.8	7.8	18.4	38.4	76.0	89.2	148.1
1957	10.8	*	*	*	2.0	4.5	7.5	17.4	38.6	74.1	72.6	103.8
1956	10.1	*	*	*	*	3.4	6.1	15.0	44.5	66.0	76.2	*
1955	9.7	*	*	*	*	3.4	7.3	14.2	37.8	59.1	88.0	*
1954	9.8	*	*	*	*	2.9	6.4	17.0	38.4	59.9	72.3	90.9
1953	9.8	*	*	*	*	4.1	6.9	16.0	37.6	52.9	91.2	104.8
1952	9.9	*	*	*	2.0	3.4	5.4	15.9	37.1	64.2	88.2	*
1951	10.2	*	*	*	2.3	2.6	6.5	17.6	40.4	64.8	72.6	*
1950	10.0	*	*	*	1.7	3.0	8.3	18.1	33.3	67.1	88.2	*
ALL OTHER, FEMALE												
1959	21.2	*	*	*	1.5	3.4	11.3	37.7	100.7	127.0	151.6	151.3
1958	21.1	*	*	*	1.7	3.9	9.8	39.1	96.8	134.1	150.3	113.2
1957	20.4	*	*	*	2.0	3.5	13.3	41.7	95.6	114.7	131.0	86.5
1956	19.1	*	*	*	2.2	3.8	9.4	38.5	93.2	113.1	118.1	102.8
1955	18.5	*	*	*	1.5	2.9	10.4	39.7	84.9	115.3	112.9	100.0
1954	18.1	*	*	*	1.5	3.0	10.3	41.8	86.9	103.4	99.2	109.4
1953	19.4	*	*	*	1.9	3.5	12.0	45.4	90.2	108.9	116.5	112.9
1952	18.1	*	*	1.3	2.2	3.1	11.5	40.4	85.5	103.1	108.6	96.6
1951	17.9	*	*	*	2.2	3.7	13.1	41.5	79.9	99.4	113.5	92.9
1950	18.7	*	*	*	2.6	4.2	14.5	43.0	80.2	113.5	108.4	122.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)												
BOTH SEXES												
1959	1.3	20.5	2.5	0.4	0.3	0.3	0.5	0.6	1.2	1.6	1.6	*
1958	1.3	21.7	2.8	0.4	0.3	0.3	0.3	0.8	1.2	1.5	1.5	2.4
1957	1.2	19.2	2.4	0.5	0.3	0.2	0.4	0.7	1.1	1.3	1.6	*
1956	1.2	19.5	2.4	0.5	0.2	0.2	0.4	0.7	1.1	1.4	1.5	*
1955	1.1	18.0	2.5	0.4	0.3	0.3	0.5	0.7	1.0	1.3	1.7	*
1954	1.1	17.7	2.4	0.5	0.2	0.3	0.4	0.7	1.0	1.3	2.1	*
1953	1.3	19.5	3.1	0.5	0.3	0.3	0.6	0.9	1.2	1.4	1.6	*
1952	1.2	18.9	2.7	0.6	0.3	0.3	0.6	0.9	1.0	1.2	1.5	*
1951	1.2	17.1	2.8	0.6	0.4	0.4	0.6	0.8	1.1	1.4	1.4	*
1950	1.2	18.2	2.8	0.6	0.4	0.4	0.5	0.8	1.1	1.4	1.4	*
MALE												
1959	1.5	23.9	3.1	0.5	0.4	0.3	0.6	0.7	1.3	2.1	2.0	*
1958	1.5	24.5	3.2	0.5	0.3	0.4	0.4	0.9	1.3	2.1	1.9	*
1957	1.4	21.9	2.6	0.5	0.4	0.2	0.4	0.8	1.5	1.6	2.0	*
1956	1.4	22.6	2.8	0.4	0.3	0.2	0.4	0.9	1.4	1.9	1.9	*
1955	1.4	21.4	2.6	0.5	0.4	0.2	0.5	0.8	1.3	1.6	2.0	*
1954	1.3	19.7	2.8	0.5	0.2	0.3	0.5	0.9	1.2	1.6	2.6	*
1953	1.6	21.7	3.6	0.7	0.3	0.4	0.7	1.1	1.5	1.9	2.1	*
1952	1.4	20.6	3.1	0.6	0.3	0.4	0.7	1.2	1.1	1.3	1.7	*
1951	1.5	19.4	3.5	0.8	0.5	0.5	0.8	1.1	1.4	2.0	1.7	*
1950	1.5	20.2	3.2	0.7	0.5	0.4	0.7	1.1	1.5	1.8	1.7	*
FEMALE												
1959	1.0	17.1	1.8	0.4	0.2	0.3	0.5	0.5	1.0	1.2	1.4	*
1958	1.1	18.8	2.4	0.3	0.2	0.2	0.3	0.6	1.1	0.9	1.2	*
1957	1.0	16.4	2.2	0.5	0.2	0.3	0.3	0.6	0.7	1.1	1.4	*
1956	0.9	16.3	2.0	0.5	*	0.2	0.3	0.5	0.7	0.9	1.1	*
1955	0.9	14.5	2.3	0.4	*	0.3	0.4	0.5	0.7	1.0	1.3	*
1954	0.9	15.6	1.9	0.4	0.2	0.3	0.3	0.5	0.7	1.1	1.7	*
1953	1.0	17.2	2.6	0.4	0.2	0.3	0.4	0.7	0.8	1.0	1.2	*
1952	1.1	17.1	2.4	0.5	0.3	0.2	0.5	0.6	0.8	1.2	1.3	*
1951	0.9	14.7	2.2	0.4	0.3	0.3	0.4	0.6	0.7	0.9	1.2	*
1950	1.0	16.2	2.4	0.5	0.3	0.3	0.4	0.6	0.7	0.9	1.1	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)												
WHITE, BOTH SEXES												
1959	1.0	16.4	2.1	0.4	0.2	0.2	0.4	0.5	0.9	1.5	1.5	*
1958	1.1	17.3	2.4	0.4	0.3	0.2	0.2	0.6	0.9	1.3	1.5	*
1957	0.9	14.9	2.1	0.4	0.2	0.2	0.3	0.5	0.8	1.2	1.4	*
1956	0.9	15.2	2.0	0.4	0.2	0.2	0.2	0.5	0.8	1.2	1.4	*
1955	0.9	14.2	2.2	0.4	0.2	0.2	0.3	0.5	0.8	1.1	1.4	*
1954	0.9	13.7	2.0	0.4	0.2	0.2	0.3	0.5	0.8	1.1	1.8	*
1953	1.1	15.8	2.8	0.5	0.2	0.3	0.4	0.7	1.0	1.3	1.5	*
1952	1.0	15.2	2.3	0.5	0.2	0.2	0.5	0.7	0.9	1.1	1.3	*
1951	1.0	13.3	2.4	0.5	0.2	0.3	0.5	0.7	0.9	1.2	1.3	*
1950	1.0	14.0	2.4	0.5	0.3	0.3	0.4	0.7	0.9	1.3	1.2	*
WHITE, MALE												
1959	1.2	18.5	2.6	0.5	0.3	0.2	0.5	0.5	1.1	1.9	1.8	*
1958	1.3	19.9	2.8	0.4	0.3	0.3	0.3	0.7	1.1	1.8	2.0	*
1957	1.1	16.7	2.3	0.4	0.3	*	0.4	0.6	1.2	1.6	1.8	*
1956	1.1	18.4	2.4	0.4	0.2	0.2	0.3	0.8	1.0	1.8	1.8	*
1955	1.1	16.9	2.5	0.4	0.3	0.2	0.4	0.6	1.1	1.3	1.9	*
1954	1.1	15.5	2.3	0.5	*	0.2	0.4	0.7	1.0	1.4	2.2	*
1953	1.3	17.5	3.2	0.6	0.3	0.3	0.5	0.9	1.4	1.8	2.0	*
1952	1.2	17.3	2.6	0.5	0.3	0.3	0.5	0.9	1.0	1.1	1.6	*
1951	1.3	15.4	3.0	0.7	0.3	0.4	0.6	0.9	1.2	1.8	1.6	*
1950	1.2	16.2	2.6	0.6	0.4	0.3	0.5	0.9	1.2	1.7	1.4	*
WHITE, FEMALE												
1959	0.8	14.2	1.6	0.3	*	0.2	0.3	0.4	0.8	1.1	1.3	*
1958	0.9	14.7	2.1	0.3	0.2	*	0.2	0.5	0.8	0.9	1.1	*
1957	0.8	13.0	1.8	0.4	*	*	0.2	0.4	0.5	1.0	1.2	*
1956	0.7	11.9	1.7	0.5	*	*	0.2	0.3	0.6	0.7	1.0	*
1955	0.7	11.4	1.9	0.3	*	0.2	0.3	0.3	0.4	0.9	1.1	*
1954	0.7	11.9	1.6	0.3	*	0.2	0.2	0.4	0.5	0.9	1.5	*
1953	0.8	14.0	2.4	0.3	*	0.2	0.2	0.5	0.6	0.9	1.2	*
1952	0.8	13.1	2.0	0.5	0.2	*	0.4	0.5	0.7	1.0	1.1	*
1951	0.7	11.1	1.8	0.3	*	*	0.3	0.4	0.6	0.7	*	*
1950	0.8	11.7	2.1	0.4	0.3	0.3	0.3	0.5	0.6	0.8	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)												
ALL OTHER, BOTH SEXES												
1959	3.2	44.4	4.6	0.7	0.9	1.0	1.8	1.8	3.5	3.5	*	*
1958	3.3	47.7	5.3	0.6	*	1.1	1.2	2.5	3.9	3.5	*	*
1957	3.2	44.9	4.4	0.9	0.9	0.8	1.2	2.8	3.9	*	*	*
1956	3.1	45.4	4.6	0.6	*	*	1.6	2.5	4.2	3.4	*	*
1955	3.0	41.5	4.1	0.7	*	*	1.6	2.7	4.0	3.7	*	*
1954	3.2	42.8	5.1	1.1	*	1.0	1.2	2.4	3.2	3.8	*	*
1953	3.1	43.7	5.3	0.9	*	1.0	2.1	2.7	3.2	*	*	*
1952	3.2	43.2	5.9	0.8	0.9	1.5	1.8	2.7	2.4	3.0	*	*
1951	3.2	41.6	6.0	0.7	1.2	1.4	2.0	2.3	3.2	3.7	*	*
1950	3.3	46.5	6.2	1.1	0.8	1.2	1.9	2.7	3.4	*	*	*
ALL OTHER, MALE												
1959	3.8	55.3	6.4	*	*	*	*	2.1	3.4	*	*	*
1958	3.7	52.6	6.1	*	*	*	*	2.9	3.6	5.7	*	*
1957	3.6	53.7	4.4	*	*	*	*	3.1	5.2	*	*	*
1956	3.6	48.5	5.5	*	*	*	1.9	2.3	6.2	*	*	*
1955	3.4	50.2	3.6	*	*	*	1.9	3.2	3.5	*	*	*
1954	3.6	46.8	6.3	1.2	*	*	*	2.4	*	*	*	*
1953	3.7	49.6	6.4	*	*	*	2.8	3.1	*	*	*	*
1952	3.6	43.2	6.8	*	*	1.7	2.0	3.8	*	*	*	*
1951	3.8	45.9	6.7	*	*	*	2.7	2.5	*	*	*	*
1950	3.9	47.4	7.6	*	*	*	2.6	3.4	5.0	*	*	*
ALL OTHER, FEMALE												
1959	2.6	33.4	2.9	*	*	*	2.0	*	3.7	*	*	*
1958	2.8	42.8	4.5	*	*	*	*	2.2	4.3	*	*	*
1957	2.8	36.1	4.4	*	*	*	*	2.5	*	*	*	*
1956	2.7	42.3	3.7	*	*	*	*	2.8	*	*	*	*
1955	2.6	33.1	4.7	*	*	*	*	2.2	4.5	*	*	*
1954	2.8	38.8	3.9	*	*	*	*	2.3	*	*	*	*
1953	2.7	37.9	4.1	*	*	*	*	2.4	*	*	*	*
1952	2.8	43.1	5.1	*	*	*	1.7	*	*	*	*	*
1951	2.7	37.4	5.3	*	*	*	*	*	*	*	*	*
1950	2.7	45.6	4.7	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)												
BOTH SEXES												
1959	515.9	13.1	2.9	2.6	8.6	26.8	99.7	346.1	939.0	2,358.7	6,086.1	14,736.9
1958	523.5	13.7	2.9	2.6	9.0	27.5	102.4	354.2	963.0	2,420.5	6,242.7	14,994.4
1957	523.5	14.5	3.2	3.1	10.5	27.8	105.5	357.5	983.1	2,454.2	6,219.8	14,859.9
1956	510.5	15.0	3.1	3.2	10.1	27.5	103.0	351.7	964.7	2,407.6	6,200.2	14,424.8
1955	506.0	14.9	3.1	3.4	10.8	27.8	105.7	355.5	958.3	2,403.3	6,219.2	14,222.4
1954	495.2	16.3	3.2	3.9	11.4	29.1	107.3	370.5	970.0	2,385.0	6,036.2	13,485.7
1953	514.8	17.7	3.5	4.4	12.3	31.3	117.1	392.8	1,029.3	2,496.5	6,330.5	14,012.0
1952	511.8	17.3	3.8	4.8	13.2	31.9	120.0	401.9	1,046.0	2,494.3	6,348.4	13,926.1
1951	513.2	15.9	3.4	5.0	13.3	32.1	121.7	406.2	1,054.5	2,538.1	6,469.8	14,333.4
1950	510.8	11.4	3.8	5.5	13.5	32.4	122.4	411.5	1,068.3	2,609.0	6,522.9	14,625.6
MALE												
1959	584.0	14.5	3.2	2.6	9.0	31.7	139.0	500.7	1,299.7	2,986.4	6,822.9	15,085.6
1958	592.0	15.1	3.0	2.7	9.8	32.0	141.6	502.2	1,320.2	3,041.8	6,994.1	15,293.3
1957	593.4	16.5	3.3	3.1	11.4	32.5	144.6	504.1	1,345.2	3,060.8	6,932.8	15,102.1
1956	580.3	16.4	3.2	3.1	10.7	32.6	140.2	496.3	1,314.2	2,991.1	6,917.8	14,584.1
1955	576.2	16.2	3.4	3.4	11.5	32.8	143.5	497.8	1,300.5	2,972.7	6,923.1	14,334.9
1954	565.3	18.2	3.5	3.8	12.2	33.7	142.6	510.1	1,303.2	2,929.4	6,713.1	13,510.5
1953	587.9	20.5	3.5	4.2	13.1	35.9	153.3	535.1	1,374.7	3,040.5	7,011.8	14,149.8
1952	583.8	20.4	4.2	4.6	14.5	36.0	154.7	542.2	1,386.1	3,014.5	7,014.6	14,061.5
1951	584.1	18.1	3.7	4.9	13.5	36.1	155.2	539.7	1,379.8	3,072.2	7,121.8	14,771.9
1950	580.0	13.9	4.1	5.1	13.8	35.9	154.2	545.4	1,388.1	3,129.4	7,163.8	15,402.3
FEMALE												
1959	449.8	11.6	2.6	2.6	8.2	22.2	62.2	195.9	600.1	1,809.8	5,512.6	14,508.8
1958	456.9	12.2	2.7	2.6	8.2	23.2	64.8	210.1	625.7	1,872.9	5,651.4	14,795.2
1957	455.5	12.4	3.0	3.1	9.5	23.2	68.1	214.4	639.3	1,914.6	5,653.2	14,694.7
1956	442.6	13.6	3.0	3.3	9.6	22.6	67.2	210.3	630.9	1,885.2	5,625.5	14,314.0
1955	437.6	13.6	2.8	3.4	10.0	23.1	69.2	216.0	629.3	1,889.7	5,650.8	14,142.8
1954	427.0	14.3	2.9	3.9	10.7	24.8	73.3	233.3	647.2	1,891.3	5,482.9	13,468.0
1953	443.9	14.9	3.4	4.6	11.5	26.9	82.1	252.7	692.2	1,999.9	5,768.7	13,912.9
1952	441.8	14.0	3.4	5.0	11.9	28.0	86.4	263.2	710.8	2,016.8	5,796.3	13,829.1
1951	443.8	13.7	3.2	5.0	13.1	28.3	89.1	273.7	731.1	2,044.0	5,919.9	14,021.7
1950	442.6	8.9	3.5	6.0	13.3	29.2	91.4	278.1	747.3	2,123.9	5,977.6	14,084.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)												
WHITE, BOTH SEXES												
1959	523.7	11.1	2.3	2.3	7.4	21.5	84.4	312.0	878.2	2,306.6	6,162.5	15,162.5
1958	529.8	11.6	2.4	2.3	7.9	21.7	87.1	315.8	896.8	2,364.1	6,318.3	15,444.0
1957	528.8	12.6	2.5	2.7	9.0	21.8	87.9	318.0	913.2	2,401.9	6,302.4	15,295.2
1956	516.3	12.5	2.5	2.9	8.7	22.1	86.7	312.9	898.0	2,362.6	6,284.8	14,873.9
1955	511.8	12.5	2.6	3.0	9.3	22.7	88.7	315.3	895.6	2,362.7	6,303.9	14,674.0
1954	499.8	13.8	2.7	3.4	9.8	23.5	89.6	327.3	909.5	2,349.2	6,117.5	13,872.3
1953	518.1	15.1	3.0	3.9	10.4	25.1	97.8	345.3	963.0	2,457.9	6,410.1	14,406.4
1952	513.8	14.7	3.5	4.2	11.2	25.6	100.4	353.3	975.8	2,453.9	6,424.5	14,289.3
1951	515.2	14.6	3.2	4.3	11.4	26.0	101.2	357.7	985.0	2,499.5	6,545.7	14,741.1
1950	512.3	10.0	3.4	5.0	11.6	26.4	101.7	362.4	1,001.2	2,552.9	6,593.2	15,053.1
WHITE, MALE												
1959	595.9	12.9	2.5	2.4	8.1	26.9	125.9	475.5	1,257.8	2,955.3	6,934.1	15,530.4
1958	602.7	13.2	2.6	2.4	9.0	27.2	130.0	474.4	1,273.9	3,005.6	7,099.4	15,727.2
1957	603.2	14.2	2.6	2.7	10.2	27.3	130.4	475.5	1,293.2	3,032.0	7,050.5	15,469.1
1956	590.8	13.8	2.6	2.9	9.5	28.0	128.1	468.2	1,267.0	2,968.5	7,036.5	14,965.2
1955	586.3	14.3	2.8	3.1	10.3	28.8	129.5	468.1	1,254.8	2,952.3	7,042.1	14,731.3
1954	574.3	15.6	3.0	3.4	10.9	29.2	128.9	477.9	1,257.2	2,915.8	6,819.1	13,807.1
1953	595.0	18.0	2.9	3.9	11.6	30.5	138.3	497.3	1,321.5	3,019.4	7,116.5	14,451.9
1952	589.8	17.6	3.9	4.0	12.8	31.2	139.9	505.6	1,329.4	2,991.3	7,103.7	14,342.3
1951	590.6	16.3	3.3	4.4	12.1	31.2	139.7	502.7	1,326.0	3,052.9	7,222.0	15,115.6
1950	585.9	12.7	3.7	4.7	12.2	31.0	138.6	508.1	1,337.3	3,091.8	7,251.5	15,840.2
WHITE, FEMALE												
1959	453.4	9.2	2.1	2.2	6.7	16.3	44.5	152.9	522.0	1,740.6	5,567.8	14,924.6
1958	458.8	10.0	2.2	2.2	6.9	16.3	45.8	161.0	541.7	1,800.0	5,709.8	15,256.7
1957	456.3	10.9	2.4	2.6	7.8	16.5	46.9	164.0	553.2	1,843.4	5,711.7	15,176.8
1956	443.6	11.1	2.3	2.9	8.0	16.4	46.7	160.8	546.4	1,821.4	5,689.5	14,810.4
1955	438.9	10.6	2.3	3.0	8.3	16.8	49.2	165.3	551.3	1,831.9	5,717.3	14,633.5
1954	427.2	11.9	2.5	3.4	8.7	18.1	51.6	179.0	573.4	1,835.7	5,549.8	13,919.1
1953	443.1	12.1	3.1	3.9	9.2	20.1	58.5	195.5	613.8	1,945.9	5,836.3	14,373.6
1952	439.6	11.7	3.0	4.4	9.6	20.4	61.8	202.8	628.5	1,962.1	5,866.9	14,251.1
1951	441.2	12.9	3.1	4.3	10.7	21.0	63.6	213.8	647.2	1,989.9	5,984.3	14,473.8
1950	439.4	7.2	3.2	5.3	11.0	22.0	65.5	217.3	664.5	2,053.3	6,038.5	14,506.2

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)												
ALL OTHER, BOTH SEXES												
1959	453.6	24.8	6.3	4.8	17.5	68.4	233.5	668.4	1,577.1	2,984.6	5,031.1	9,541.2
1958	472.5	26.3	5.7	4.6	16.7	74.1	235.7	719.9	1,670.0	3,100.6	5,190.8	9,586.4
1957	479.8	25.9	7.1	6.3	21.2	75.9	259.1	734.8	1,752.1	3,080.0	5,079.6	9,404.7
1956	463.0	30.4	7.4	5.4	20.4	71.7	243.8	723.2	1,718.9	2,945.7	5,000.4	9,006.6
1955	458.2	29.9	6.5	5.9	21.4	70.2	251.6	741.6	1,684.7	2,888.7	5,026.2	8,701.7
1954	457.0	32.1	6.4	7.1	23.1	75.8	260.3	785.7	1,687.3	2,813.8	4,878.7	8,496.4
1953	487.5	34.8	6.4	7.9	26.0	82.9	284.2	847.5	1,838.3	2,959.4	5,165.4	9,165.4
1952	495.2	34.0	6.2	8.9	27.7	84.8	291.1	865.2	1,924.1	2,978.5	5,230.5	9,442.9
1951	496.1	24.4	5.1	9.7	27.4	84.1	300.4	868.8	1,939.6	2,991.4	5,347.9	9,397.9
1950	498.0	21.2	6.4	9.2	28.0	84.1	303.1	882.6	1,878.8	3,370.3	5,460.3	9,545.6
ALL OTHER, MALE												
1959	488.0	24.3	7.2	4.5	15.8	71.8	256.4	743.8	1,732.0	3,350.8	5,403.5	10,086.2
1958	504.8	27.0	6.0	4.6	15.8	72.5	245.7	771.1	1,809.9	3,468.7	5,645.3	10,781.5
1957	512.5	30.7	7.6	6.1	20.8	77.2	270.8	780.2	1,908.8	3,409.0	5,420.7	10,711.5
1956	492.9	32.0	7.3	5.0	19.9	72.2	247.6	767.3	1,839.6	3,257.7	5,382.3	9,980.0
1955	491.4	28.7	7.2	5.7	20.9	67.6	266.4	784.3	1,822.7	3,214.4	5,368.0	9,479.2
1954	490.0	35.1	7.1	7.1	21.6	73.7	263.8	819.7	1,839.1	3,090.6	5,303.4	9,709.1
1953	527.3	36.8	7.4	6.9	24.5	83.7	286.4	897.1	2,007.0	3,280.8	5,600.0	10,295.2
1952	532.0	39.2	6.5	9.0	27.2	78.6	286.4	891.1	2,081.5	3,297.6	5,802.7	10,510.0
1951	527.7	29.5	6.6	9.1	24.1	79.8	294.9	891.2	2,050.9	3,299.7	5,813.2	10,484.2
1950	528.4	22.2	7.0	7.9	26.3	79.0	294.4	902.6	1,994.8	3,618.2	5,942.4	10,285.0
ALL OTHER, FEMALE												
1959	421.0	25.2	5.4	5.1	19.1	65.6	212.7	597.3	1,428.0	2,652.9	4,698.1	9,135.9
1958	442.1	25.5	5.4	4.7	17.4	75.5	226.3	672.2	1,539.6	2,758.1	4,777.8	8,989.5
1957	448.8	21.2	6.7	6.6	21.6	74.7	248.4	691.5	1,599.4	2,779.5	4,762.1	8,740.5
1956	434.6	28.8	7.6	5.7	20.9	71.3	240.5	679.8	1,600.3	2,645.4	4,640.6	8,330.6
1955	426.8	31.2	5.8	6.2	21.8	72.5	237.6	700.1	1,547.9	2,588.8	4,664.4	8,152.9
1954	426.0	29.2	5.7	7.0	24.4	77.6	257.2	752.4	1,535.5	2,558.0	4,438.9	7,928.1
1953	450.0	32.8	5.4	8.9	27.2	82.2	282.1	798.8	1,664.0	2,651.8	4,713.2	8,400.0
1952	460.4	28.9	6.0	8.8	28.2	90.3	295.4	839.6	1,762.5	2,679.0	4,687.9	8,706.9
1951	466.1	19.4	3.5	10.3	30.2	87.8	305.6	845.5	1,823.8	2,700.0	4,903.6	8,660.7
1950	468.8	20.1	5.7	10.5	29.6	88.6	311.1	862.5	1,759.8	3,120.5	4,985.4	9,017.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)												
BOTH SEXES												
1959	508.9	12.3	2.7	2.1	7.1	24.0	95.5	338.6	926.1	2,334.3	6,027.7	14,596.8
1958	515.5	13.0	2.6	2.1	7.1	24.6	97.7	345.8	948.6	2,391.0	6,174.3	14,826.0
1957	514.6	14.0	2.9	2.4	8.3	24.3	100.1	348.6	966.5	2,421.6	6,146.3	14,678.9
1956	501.4	14.2	2.7	2.5	8.2	24.2	97.8	342.6	947.9	2,373.4	6,117.6	14,228.1
1955	496.3	14.4	2.7	2.7	8.5	24.7	100.3	345.9	940.6	2,366.5	6,128.7	14,021.5
1954	484.6	15.4	2.6	3.1	9.2	25.5	101.9	359.6	949.9	2,344.3	5,934.7	13,271.1
1953	503.0	17.1	2.8	3.5	9.8	27.4	110.8	380.6	1,007.3	2,450.1	6,217.5	13,751.5
1952	498.5	16.4	3.1	3.9	10.9	28.0	113.0	387.7	1,020.4	2,442.1	6,214.3	13,639.6
1951	498.5	15.2	2.6	4.0	10.7	28.0	114.4	391.5	1,026.0	2,479.0	6,318.8	13,977.1
1950	494.4	10.7	2.8	4.6	11.0	27.9	114.1	395.2	1,037.4	2,539.2	6,348.6	14,214.6
MALE												
1959	576.3	13.6	3.0	2.2	7.3	28.2	134.0	491.6	1,284.8	2,958.1	6,753.8	14,918.9
1958	583.2	14.3	2.8	2.1	7.6	28.7	135.9	492.6	1,303.6	3,007.5	6,913.4	15,093.9
1957	583.8	16.0	3.0	2.4	8.8	28.2	138.5	493.9	1,326.8	3,022.9	6,850.7	14,894.7
1956	570.4	15.4	2.8	2.5	8.5	28.5	134.5	486.0	1,295.3	2,952.9	6,823.1	14,348.0
1955	565.8	15.2	2.9	2.7	8.9	29.1	137.3	487.1	1,281.0	2,930.9	6,817.6	14,107.2
1954	553.8	17.3	2.9	3.0	9.6	29.6	136.5	498.3	1,281.1	2,882.3	6,593.8	13,264.8
1953	575.0	19.7	2.8	3.3	10.4	31.5	146.5	521.4	1,350.8	2,988.1	6,878.8	13,846.4
1952	569.2	19.5	3.3	3.6	11.8	31.5	147.3	526.0	1,357.6	2,954.9	6,856.7	13,731.9
1951	568.3	17.0	2.7	3.9	10.8	31.6	147.5	523.2	1,348.3	3,005.6	6,949.6	14,381.3
1950	562.5	12.8	3.0	4.1	10.8	31.0	145.5	527.8	1,353.9	3,051.7	6,964.4	14,934.9
FEMALE												
1959	443.5	11.0	2.4	2.0	7.0	19.9	58.7	190.0	589.1	1,788.9	5,462.6	14,386.2
1958	449.7	11.6	2.5	2.1	6.7	20.6	61.2	202.9	613.5	1,847.8	5,592.5	14,647.5
1957	447.4	11.9	2.8	2.3	7.8	20.6	63.3	206.8	624.5	1,886.8	5,586.4	14,531.8
1956	434.2	13.0	2.7	2.6	8.0	20.0	62.5	202.2	616.1	1,854.6	5,552.5	14,144.7
1955	428.7	13.5	2.4	2.7	8.3	20.4	64.7	207.5	613.2	1,857.4	5,572.4	13,960.8
1954	417.3	13.4	2.3	3.2	8.9	21.5	68.5	223.4	629.1	1,856.4	5,395.8	13,275.5
1953	432.9	14.5	2.8	3.7	9.3	23.4	76.3	241.8	672.1	1,959.0	5,672.0	13,683.3
1952	429.7	13.3	2.9	4.1	10.0	24.7	79.8	250.9	688.0	1,971.3	5,682.1	13,573.5
1951	430.2	13.3	2.5	4.1	10.7	24.7	82.2	260.8	705.6	1,991.9	5,786.8	13,689.7
1950	427.3	8.5	2.6	5.0	11.1	25.0	83.6	263.1	719.7	2,061.4	5,824.7	13,712.9

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)												
WHITE, BOTH SEXES												
1959	517.6	10.4	2.2	1.8	6.0	19.0	81.1	306.3	867.7	2,286.3	6,108.3	15,029.4
1958	522.8	10.9	2.2	1.9	6.1	19.1	83.5	309.3	885.8	2,338.6	6,254.6	15,283.6
1957	521.0	12.1	2.3	2.0	6.9	18.9	83.8	311.1	899.7	2,373.6	6,234.2	15,123.3
1956	508.2	11.7	2.1	2.3	6.9	19.3	82.8	306.1	884.4	2,333.1	6,207.4	14,684.8
1955	503.1	11.9	2.2	2.3	7.1	20.0	84.7	307.9	881.2	2,330.1	6,219.1	14,480.8
1954	490.3	13.1	2.1	2.7	7.7	20.4	85.4	318.8	892.9	2,313.7	6,022.2	13,667.1
1953	507.5	14.5	2.4	3.0	8.1	21.8	92.9	335.8	944.8	2,417.2	6,305.2	14,155.4
1952	502.1	13.9	2.8	3.3	9.0	22.3	95.1	342.6	955.3	2,408.4	6,301.0	14,021.8
1951	502.1	14.0	2.4	3.3	9.0	22.6	95.6	346.4	961.5	2,448.2	6,404.5	14,403.9
1950	497.6	9.6	2.5	4.0	9.1	22.6	95.4	349.7	975.9	2,492.6	6,430.7	14,655.8
WHITE, MALE												
1959	589.0	12.0	2.4	2.0	6.4	23.7	121.8	468.1	1,245.2	2,931.8	6,869.7	15,374.8
1958	594.8	12.5	2.3	1.9	6.9	24.1	125.2	466.8	1,260.6	2,975.8	7,023.9	15,541.5
1957	594.6	13.7	2.4	2.0	7.7	23.5	125.4	466.9	1,277.9	2,998.3	6,974.7	15,277.5
1956	581.9	13.0	2.3	2.3	7.3	24.3	123.3	460.0	1,251.5	2,935.5	6,948.2	14,742.4
1955	576.8	13.2	2.4	2.4	7.6	25.4	124.5	459.3	1,238.4	2,915.1	6,943.3	14,510.2
1954	563.9	14.9	2.4	2.6	8.4	25.5	123.9	468.3	1,238.4	2,875.3	6,709.7	13,577.0
1953	583.3	17.2	2.3	2.9	8.9	26.6	132.5	486.1	1,301.4	2,973.8	6,993.9	14,160.1
1952	576.8	16.7	3.0	3.0	10.2	27.1	133.9	492.7	1,306.0	2,939.3	6,960.1	14,034.0
1951	576.4	15.5	2.4	3.3	9.4	27.1	133.3	489.2	1,299.2	2,995.1	7,062.7	14,753.2
1950	570.1	12.1	2.6	3.7	9.3	26.7	131.2	493.4	1,308.7	3,025.0	7,066.8	15,395.6
WHITE, FEMALE												
1959	448.0	8.7	2.0	1.7	5.5	14.5	42.0	148.8	513.4	1,723.1	5,521.5	14,806.0
1958	452.5	9.3	2.0	1.8	5.4	14.3	43.3	155.8	532.7	1,778.2	5,655.3	15,113.2
1957	449.2	10.5	2.2	1.9	6.2	14.4	43.6	158.8	541.3	1,819.7	5,649.6	15,018.4
1956	436.2	10.4	2.0	2.2	6.5	14.5	43.6	155.3	534.6	1,795.0	5,620.7	14,644.7
1955	431.1	10.5	2.0	2.3	6.7	14.7	46.2	159.2	538.7	1,803.3	5,643.6	14,460.0
1954	418.5	11.1	1.8	2.8	7.0	15.6	48.1	171.8	559.0	1,804.8	5,465.9	13,731.8
1953	433.5	11.7	2.5	3.1	7.4	17.3	54.5	187.6	597.6	1,909.9	5,745.8	14,152.0
1952	429.1	10.9	2.5	3.5	7.9	17.8	57.2	194.2	610.8	1,922.4	5,759.9	14,013.1
1951	429.2	12.4	2.4	3.3	8.6	18.3	58.9	204.7	626.9	1,944.6	5,858.0	14,154.5
1950	425.9	7.1	2.3	4.4	9.0	18.7	60.2	206.6	642.4	1,999.1	5,894.7	14,141.8

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)												
ALL OTHER, BOTH SEXES												
1959	439.9	23.3	6.1	4.1	15.5	62.4	220.5	644.2	1,538.6	2,911.8	4,913.8	9,316.2
1958	457.2	25.0	5.2	4.0	14.5	67.7	221.9	692.9	1,620.6	3,023.9	5,056.5	9,321.2
1957	462.9	25.2	6.6	5.5	18.2	68.1	242.1	706.7	1,702.2	2,997.0	4,929.6	9,117.2
1956	445.4	28.9	6.8	4.5	17.6	64.2	228.0	691.4	1,666.5	2,855.9	4,843.7	8,718.0
1955	440.6	29.5	5.7	5.2	18.5	63.4	235.4	711.3	1,628.8	2,802.5	4,854.3	8,406.9
1954	437.4	30.1	5.6	6.2	20.1	67.6	244.7	751.7	1,626.2	2,710.9	4,685.2	8,169.1
1953	465.2	34.3	5.4	7.1	22.4	74.0	265.6	809.0	1,769.4	2,843.7	4,932.5	8,788.5
1952	468.4	33.6	5.4	7.9	24.4	76.0	269.1	817.8	1,834.9	2,846.2	4,943.4	8,920.4
1951	468.0	22.9	4.0	8.7	23.7	74.4	277.7	821.5	1,848.2	2,840.1	5,051.6	8,810.6
1950	466.8	17.7	5.2	8.4	24.9	73.6	278.0	831.5	1,781.4	3,171.3	5,107.2	8,971.8
ALL OTHER, MALE												
1959	473.4	22.7	7.0	3.9	13.8	65.1	243.0	717.8	1,692.7	3,266.4	5,272.9	9,793.1
1958	488.4	25.3	5.4	3.7	13.2	67.4	231.6	743.4	1,758.3	3,379.8	5,499.3	10,414.8
1957	495.1	30.0	6.7	5.3	17.3	68.4	254.6	754.4	1,856.3	3,320.6	5,257.8	10,315.4
1956	474.5	30.1	6.2	3.9	16.8	65.3	233.7	736.5	1,783.2	3,158.2	5,205.4	9,584.0
1955	473.5	28.0	6.3	4.9	18.1	61.2	250.3	755.7	1,768.6	3,118.0	5,176.0	9,170.8
1954	468.5	32.7	6.1	6.0	18.2	65.8	248.4	787.1	1,778.3	2,965.1	5,052.1	9,263.6
1953	504.2	36.3	6.3	6.3	21.9	75.0	270.2	860.4	1,937.4	3,148.3	5,328.9	9,842.9
1952	503.7	38.3	5.5	7.9	23.5	70.3	266.5	844.1	1,990.6	3,148.1	5,451.8	9,910.0
1951	498.6	26.8	5.1	8.0	21.1	71.4	274.9	846.2	1,960.6	3,130.3	5,461.3	9,742.1
1950	496.3	17.8	5.6	7.3	23.0	70.3	273.6	856.7	1,894.4	3,398.4	5,537.1	9,551.1
ALL OTHER, FEMALE												
1959	408.3	23.8	5.1	4.3	17.0	60.2	200.2	574.7	1,390.5	2,590.4	4,592.5	8,961.5
1958	427.6	24.8	5.1	4.4	15.6	68.1	212.9	645.7	1,492.4	2,692.7	4,654.2	8,789.5
1957	432.4	20.5	6.4	5.6	19.0	67.8	230.7	661.0	1,552.1	2,701.4	4,624.1	8,521.6
1956	417.8	27.7	7.4	5.0	18.2	63.2	222.9	647.3	1,551.9	2,564.9	4,502.9	8,116.7
1955	409.5	31.2	5.2	5.5	18.9	65.3	221.4	668.3	1,490.2	2,512.0	4,512.9	7,867.6
1954	408.1	27.6	5.1	6.4	21.7	69.2	241.3	717.1	1,474.0	2,476.1	4,301.6	7,671.9
1953	428.4	32.3	4.6	8.0	22.9	73.1	261.3	758.5	1,595.9	2,552.0	4,518.2	8,074.2
1952	435.2	28.9	5.4	8.0	25.1	81.0	271.5	791.9	1,674.9	2,563.0	4,461.2	8,237.9
1951	438.9	18.9	2.9	9.4	26.0	77.0	280.2	795.9	1,731.2	2,565.9	4,660.4	8,178.6
1950	438.6	17.7	4.9	9.6	26.6	76.6	282.0	806.2	1,665.4	2,942.5	4,683.8	8,557.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
CEREBROVASCULAR DISEASES (ICD 330-334)												
BOTH SEXES												
1959	108.4	4.1	1.0	0.7	1.7	4.8	15.0	49.3	150.8	478.4	1,508.1	3,640.9
1958	110.1	4.6	1.0	0.7	1.8	4.9	15.5	51.7	156.5	495.0	1,539.3	3,676.9
1957	110.2	4.9	1.1	0.8	2.0	4.9	16.6	55.2	161.8	508.2	1,525.4	3,581.9
1956	106.3	4.6	1.1	0.7	1.9	4.8	16.2	54.6	161.1	497.6	1,489.8	3,389.3
1955	106.0	5.6	0.9	0.7	1.9	5.0	16.6	56.2	163.2	506.0	1,497.6	3,290.6
1954	104.1	5.1	0.9	0.7	1.9	4.6	17.9	61.2	172.0	503.8	1,460.0	3,057.3
1953	107.3	5.8	1.0	0.6	1.7	5.1	19.3	67.1	183.8	531.3	1,503.1	3,051.5
1952	106.8	5.3	1.1	0.7	2.0	5.0	19.1	69.4	189.2	534.8	1,500.4	3,060.1
1951	106.7	4.9	0.9	0.6	1.8	4.8	19.5	70.6	192.3	550.9	1,516.4	2,997.1
1950	104.0	5.1	0.9	0.5	1.6	4.2	18.7	70.4	194.2	554.7	1,499.6	2,990.1
MALE												
1959	104.7	5.1	1.0	0.8	1.8	4.5	14.9	52.3	166.6	534.1	1,566.7	3,558.0
1958	106.6	5.1	1.2	0.8	2.0	4.7	14.8	53.7	174.3	552.4	1,601.0	3,529.1
1957	107.7	5.7	1.1	0.8	2.2	5.0	16.3	57.0	179.9	561.8	1,594.1	3,474.5
1956	103.7	5.2	1.0	0.7	1.9	4.8	15.7	55.9	176.8	548.1	1,547.5	3,257.8
1955	104.0	6.0	1.1	0.8	1.8	5.1	16.2	56.3	180.2	554.7	1,553.1	3,185.5
1954	102.2	5.4	1.0	0.8	1.9	4.5	17.5	60.6	185.8	550.9	1,514.9	2,887.2
1953	105.5	7.0	1.0	0.7	2.0	5.1	18.5	65.8	199.0	573.2	1,552.7	2,918.3
1952	105.4	6.9	1.2	0.8	2.1	4.9	17.9	68.1	204.2	572.8	1,559.1	2,965.9
1951	104.9	5.9	1.0	0.6	1.7	4.7	18.6	68.3	204.7	593.7	1,546.0	2,925.8
1950	102.5	6.4	1.1	0.5	1.8	4.2	17.5	67.9	205.2	589.6	1,543.6	3,048.6
FEMALE												
1959	112.1	3.1	1.1	0.5	1.7	5.0	15.1	46.3	136.0	429.7	1,462.9	3,695.0
1958	113.4	4.1	0.8	0.7	1.5	5.1	16.2	49.8	139.6	444.4	1,490.7	3,775.4
1957	112.6	4.1	1.1	0.7	1.8	4.8	16.9	53.5	144.6	460.5	1,470.9	3,655.3
1956	108.8	4.0	1.2	0.7	1.8	4.7	16.6	53.3	146.2	452.4	1,443.6	3,480.9
1955	107.9	5.3	0.7	0.6	1.9	4.8	17.0	56.1	147.0	462.1	1,452.8	3,365.0
1954	106.0	4.7	0.8	0.7	1.9	4.6	18.3	61.7	158.6	461.2	1,414.6	3,179.1
1953	109.0	4.7	0.9	0.6	1.5	5.0	20.1	68.3	169.0	493.1	1,461.6	3,147.3
1952	108.2	3.7	1.1	0.7	1.8	5.1	20.3	70.8	174.4	499.9	1,451.7	3,127.6
1951	108.5	3.9	0.7	0.7	1.8	4.8	20.4	72.9	180.0	511.1	1,490.8	3,047.8
1950	105.6	3.7	0.7	0.4	1.5	4.3	19.9	72.9	183.1	522.1	1,462.2	2,949.4

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
CEREBROVASCULAR DISEASES (ICD 330-334)												
WHITE, BOTH SEXES												
1959	106.9	3.7	0.9	0.6	1.5	3.8	10.5	37.4	125.0	448.1	1,514.9	3,730.8
1958	108.4	4.0	0.9	0.7	1.6	3.6	11.1	39.3	129.9	463.8	1,546.8	3,781.0
1957	108.4	4.3	1.0	0.7	1.8	3.5	11.6	42.2	135.4	479.3	1,537.4	3,677.3
1956	104.6	3.8	0.9	0.7	1.7	3.5	11.7	41.0	135.3	472.6	1,502.5	3,477.7
1955	104.3	4.5	0.9	0.7	1.6	3.8	11.6	42.8	137.8	481.1	1,511.4	3,384.5
1954	102.2	4.4	0.8	0.7	1.7	3.5	12.7	46.1	147.6	481.0	1,471.8	3,136.7
1953	105.4	4.9	0.9	0.6	1.5	3.7	14.0	51.2	159.1	509.9	1,516.5	3,133.3
1952	104.7	4.3	1.1	0.7	1.7	3.9	13.8	53.7	163.8	512.8	1,512.5	3,126.1
1951	104.6	4.7	0.8	0.6	1.4	3.6	14.3	54.9	167.6	529.8	1,526.4	3,075.0
1950	101.9	4.4	0.9	0.4	1.4	3.1	13.3	54.4	169.6	532.5	1,510.2	3,062.7
WHITE, MALE												
1959	103.1	4.4	0.9	0.8	1.7	3.7	10.9	41.4	142.4	505.9	1,579.6	3,649.1
1958	104.7	4.6	1.1	0.7	1.9	3.7	11.3	42.7	148.5	522.1	1,611.2	3,616.1
1957	105.9	4.7	1.0	0.8	2.0	4.0	12.2	45.4	155.0	535.0	1,609.8	3,560.8
1956	102.2	4.5	0.8	0.7	1.8	3.7	12.2	44.3	153.5	524.3	1,563.3	3,336.1
1955	102.3	5.0	1.0	0.8	1.6	4.2	11.6	45.3	156.4	531.4	1,569.9	3,262.6
1954	100.4	4.8	0.9	0.7	1.7	3.6	13.2	47.6	163.1	530.6	1,530.1	2,941.5
1953	103.6	6.1	0.9	0.7	1.8	4.0	14.2	51.2	175.7	552.7	1,570.4	2,982.5
1952	103.4	5.7	1.2	0.7	1.8	4.2	13.7	54.3	180.2	553.1	1,572.0	3,015.0
1951	103.0	5.6	0.9	0.6	1.4	3.8	14.5	54.2	182.5	575.7	1,558.9	2,984.8
1950	100.5	5.9	1.1	0.5	1.6	3.4	13.1	53.7	182.2	569.7	1,556.3	3,127.1
WHITE, FEMALE												
1959	110.6	2.9	1.0	0.5	1.4	3.8	10.1	33.4	108.7	397.6	1,465.5	3,783.7
1958	111.9	3.5	0.7	0.7	1.3	3.6	10.8	36.1	112.4	412.4	1,496.6	3,890.0
1957	110.9	3.8	1.0	0.6	1.5	3.0	11.1	39.1	116.9	430.0	1,479.8	3,756.7
1956	107.0	3.0	1.0	0.7	1.6	3.3	11.3	37.7	117.9	426.4	1,454.3	3,576.3
1955	106.2	4.1	0.7	0.6	1.6	3.5	11.6	40.5	119.9	435.8	1,465.5	3,470.8
1954	103.9	3.9	0.7	0.6	1.8	3.3	12.2	44.7	132.6	436.1	1,424.1	3,276.8
1953	107.1	3.6	0.9	0.5	1.3	3.4	13.9	51.2	143.0	470.9	1,472.8	3,242.3
1952	105.9	2.7	1.0	0.6	1.5	3.6	13.9	53.0	147.7	476.0	1,463.6	3,206.0
1951	106.2	3.7	0.8	0.7	1.5	3.5	14.0	55.7	152.9	487.6	1,499.4	3,139.5
1950	103.3	2.9	0.6	0.4	1.2	2.9	13.6	55.0	156.9	498.1	1,471.3	3,017.9

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
CEREBROVASCULAR DISEASES (ICD 330-334)												
ALL OTHER, BOTH SEXES												
1959	120.4	6.8	1.6	0.9	3.2	12.7	54.1	162.1	421.7	841.4	1,413.1	2,542.6
1958	123.8	8.1	1.8	1.1	3.0	15.1	54.1	169.9	440.6	870.5	1,434.9	2,424.2
1957	124.3	8.7	1.8	1.1	3.7	16.3	60.3	179.5	451.6	854.5	1,362.9	2,381.3
1956	119.8	9.5	2.2	0.5	3.1	15.0	54.7	184.7	453.5	796.4	1,310.4	2,323.0
1955	119.9	12.6	1.4	0.8	3.8	14.4	59.6	184.6	458.4	804.4	1,305.1	2,143.1
1954	120.1	9.6	1.6	1.2	3.5	13.9	62.8	205.6	461.1	777.2	1,294.3	2,027.3
1953	123.6	12.2	1.5	0.8	3.4	16.5	65.2	218.6	485.3	787.4	1,306.0	2,046.2
1952	125.2	12.5	1.3	1.0	4.1	14.4	65.0	220.1	506.4	797.5	1,323.0	2,244.9
1951	124.4	6.8	*	0.8	4.2	14.5	65.1	220.0	507.4	801.0	1,369.1	2,053.2
1950	122.3	9.6	*	0.7	3.6	13.7	65.9	224.3	491.6	854.6	1,339.6	2,127.9
ALL OTHER, MALE												
1959	117.3	9.3	*	1.1	2.6	11.2	50.3	156.9	417.2	864.4	1,401.4	2,534.5
1958	122.5	8.3	2.4	1.2	2.9	13.1	46.0	160.7	447.1	909.0	1,470.5	2,640.7
1957	121.9	11.5	*	1.0	3.6	13.8	52.9	169.0	449.8	879.4	1,391.9	2,442.3
1956	116.5	9.6	2.0	*	2.9	14.8	47.2	167.7	435.7	828.6	1,343.8	2,312.0
1955	118.0	12.3	*	*	3.7	13.4	56.7	163.4	451.0	830.9	1,333.6	2,241.7
1954	117.3	9.3	*	1.4	3.7	12.3	54.9	186.1	450.2	791.5	1,313.4	2,190.9
1953	122.0	12.8	*	*	3.6	14.3	56.5	205.4	477.4	813.1	1,314.0	2,100.0
1952	122.9	14.9	*	*	4.5	11.3	55.3	200.5	497.7	806.9	1,384.5	2,345.0
1951	121.1	*	*	*	4.5	13.1	54.8	202.6	482.1	806.4	1,384.0	2,189.5
1950	119.5	10.4	*	*	3.3	11.9	57.4	204.2	479.3	847.8	1,365.7	2,132.0
ALL OTHER, FEMALE												
1959	123.3	*	*	*	3.7	14.0	57.5	166.9	426.0	820.6	1,423.6	2,548.7
1958	124.9	7.9	*	1.0	3.1	16.7	61.6	178.8	435.7	833.4	1,402.6	2,334.2
1957	126.6	*	2.0	1.2	3.7	18.5	67.0	189.5	453.3	831.8	1,335.9	2,402.7
1956	123.0	9.5	2.4	*	3.3	15.3	61.7	201.0	471.0	763.0	1,279.0	2,330.6
1955	121.8	12.9	*	*	3.9	15.3	62.2	205.2	465.8	779.9	1,268.2	2,073.5
1954	122.8	10.0	*	*	3.2	15.3	70.1	224.7	472.0	764.0	1,265.9	1,978.1
1953	125.1	11.5	*	*	3.1	18.5	73.3	231.5	492.3	761.2	1,287.6	2,009.7
1952	127.3	10.2	*	*	3.8	17.1	74.0	239.5	515.3	788.8	1,264.7	2,175.9
1951	127.6	*	*	*	3.9	15.8	74.6	237.1	533.9	796.0	1,355.0	1,960.7
1950	124.9	*	*	*	4.0	15.2	73.7	244.3	504.2	861.4	1,314.0	2,125.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
DISEASES OF HEART (ICD 400-402, 410-413)												
BOTH SEXES												
1959	363.2	7.1	1.5	1.3	4.3	16.5	73.8	272.6	728.7	1,721.1	4,030.7	9,166.8
1958	367.7	7.4	1.4	1.3	4.4	16.8	75.3	276.4	746.8	1,754.6	4,138.3	9,300.0
1957	369.4	7.7	1.6	1.5	5.4	16.9	77.3	278.2	761.8	1,783.1	4,155.4	9,314.3
1956	361.0	8.1	1.5	1.6	5.5	16.9	75.5	272.8	746.4	1,750.5	4,158.7	9,079.8
1955	356.5	7.4	1.6	1.8	5.6	17.2	77.4	275.3	738.9	1,739.7	4,144.3	8,917.2
1954	348.3	8.7	1.5	2.2	6.3	17.9	77.6	283.1	740.1	1,725.0	4,016.6	8,472.6
1953	361.3	9.5	1.6	2.7	6.9	19.4	84.4	297.1	783.7	1,794.6	4,213.3	8,789.1
1952	357.6	9.2	1.7	3.0	7.8	20.0	86.7	301.8	790.7	1,783.9	4,201.6	8,661.8
1951	357.0	8.2	1.5	3.2	7.7	20.4	87.5	304.6	793.1	1,798.8	4,258.7	8,973.5
1950	356.8	4.1	1.6	3.9	8.2	20.9	88.3	309.2	804.3	1,857.2	4,311.0	9,152.5
MALE												
1959	432.2	7.3	1.8	1.3	4.7	21.1	111.5	418.5	1,056.7	2,248.9	4,633.1	9,540.6
1958	436.6	7.8	1.4	1.3	4.9	21.5	113.7	417.5	1,069.2	2,275.9	4,741.3	9,655.2
1957	439.1	8.7	1.6	1.4	5.9	21.0	115.3	418.0	1,089.7	2,294.6	4,735.7	9,596.4
1956	430.4	8.4	1.5	1.6	6.0	21.5	112.0	411.4	1,063.9	2,246.0	4,745.7	9,288.4
1955	425.8	7.5	1.7	1.8	6.1	21.8	114.3	412.1	1,050.9	2,224.8	4,713.5	9,044.7
1954	417.8	9.9	1.6	2.1	6.9	22.4	112.3	418.8	1,047.1	2,189.3	4,562.6	8,637.2
1953	433.2	10.5	1.6	2.5	7.6	23.8	120.4	435.1	1,100.3	2,263.7	4,760.6	9,008.7
1952	428.1	10.4	1.9	2.8	8.9	24.1	121.8	437.8	1,101.3	2,235.7	4,733.5	8,817.9
1951	427.2	9.2	1.5	3.1	8.2	24.3	121.4	435.7	1,093.2	2,256.6	4,808.1	9,375.4
1950	424.7	4.7	1.7	3.5	8.3	24.4	120.4	441.2	1,100.5	2,310.2	4,825.8	9,661.4
FEMALE												
1959	296.1	6.8	1.2	1.3	3.9	12.0	37.7	131.0	420.4	1,259.6	3,561.5	8,922.5
1958	300.7	6.9	1.4	1.3	3.9	12.3	38.5	139.2	442.4	1,295.2	3,663.6	9,063.2
1957	301.7	6.6	1.6	1.5	4.9	13.0	40.8	141.9	450.5	1,328.3	3,694.3	9,121.9
1956	293.5	7.9	1.4	1.6	4.9	12.4	40.3	137.2	443.1	1,306.9	3,688.6	8,934.7
1955	289.0	7.2	1.6	1.8	5.1	12.8	41.9	141.1	438.9	1,302.2	3,684.7	8,826.9
1954	280.8	7.5	1.3	2.3	5.7	13.7	44.1	149.8	442.8	1,303.8	3,570.8	8,354.8
1953	291.5	8.4	1.6	2.9	6.4	15.2	49.8	161.2	474.8	1,366.2	3,762.3	8,631.3
1952	288.9	8.0	1.6	3.2	6.8	16.1	52.6	167.5	484.6	1,369.1	3,760.9	8,549.9
1951	288.3	7.2	1.5	3.3	7.3	16.7	54.7	174.5	494.7	1,375.4	3,796.1	8,687.8
1950	289.7	3.4	1.6	4.3	8.2	17.6	57.0	177.8	507.0	1,434.9	3,873.0	8,798.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
DISEASES OF HEART (ICD 400-402, 410-413)												
WHITE, BOTH SEXES												
1959	373.1	5.6	1.0	1.0	3.5	13.5	66.3	255.6	701.2	1,708.8	4,100.4	9,446.5
1958	376.6	6.0	1.2	1.1	3.7	13.7	67.8	256.5	715.2	1,739.6	4,208.4	9,593.7
1957	377.5	6.8	1.2	1.1	4.4	13.8	68.2	257.1	726.2	1,769.6	4,227.1	9,605.9
1956	369.4	6.8	1.0	1.4	4.5	14.1	67.1	253.1	713.3	1,740.3	4,231.2	9,384.6
1955	364.9	6.2	1.2	1.5	4.7	14.5	69.1	253.8	709.5	1,733.6	4,216.3	9,222.3
1954	356.0	7.3	1.2	1.8	5.2	14.9	68.5	260.9	711.6	1,722.1	4,088.4	8,738.7
1953	367.9	8.2	1.3	2.2	5.6	16.1	74.1	272.0	750.8	1,788.8	4,283.3	9,044.9
1952	363.3	7.9	1.4	2.4	6.5	16.5	76.2	275.8	755.8	1,776.7	4,270.3	8,916.4
1951	362.6	7.5	1.3	2.5	6.5	17.0	76.3	278.6	758.3	1,793.8	4,327.7	9,249.6
1950	362.0	3.8	1.3	3.4	6.8	17.5	77.1	283.0	772.0	1,838.3	4,375.2	9,441.7
WHITE, MALE												
1959	446.5	6.5	1.2	1.1	4.1	18.4	105.9	409.3	1,045.8	2,257.1	4,731.0	9,835.6
1958	450.0	6.6	1.2	1.1	4.4	18.8	108.7	406.4	1,056.2	2,279.9	4,836.5	9,959.5
1957	451.5	7.6	1.2	1.1	5.0	18.2	108.5	406.1	1,070.5	2,302.2	4,836.0	9,841.2
1956	443.3	7.0	1.2	1.4	5.1	19.2	106.5	400.0	1,048.1	2,257.0	4,847.1	9,542.1
1955	438.5	6.7	1.2	1.5	5.3	19.7	108.4	398.8	1,036.2	2,238.4	4,813.9	9,316.0
1954	429.8	8.4	1.3	1.8	6.1	19.9	106.1	405.3	1,031.2	2,206.9	4,657.1	8,854.3
1953	443.7	9.4	1.2	2.1	6.5	20.7	112.9	418.2	1,079.3	2,276.2	4,852.6	9,206.3
1952	437.9	9.1	1.5	2.2	7.7	21.3	114.6	421.4	1,078.7	2,243.8	4,818.4	9,019.0
1951	437.1	8.3	1.3	2.6	7.3	21.5	113.2	419.2	1,071.3	2,268.9	4,901.4	9,620.7
1950	434.2	4.6	1.3	3.1	7.1	21.5	112.6	424.1	1,082.6	2,309.4	4,908.0	9,952.3
WHITE, FEMALE												
1959	301.7	4.8	0.9	1.0	3.0	8.8	28.3	106.0	377.8	1,230.6	3,614.0	9,194.8
1958	305.0	5.4	1.1	1.0	3.1	8.7	28.5	110.2	393.9	1,264.4	3,719.1	9,351.9
1957	305.2	5.8	1.1	1.1	3.8	9.6	29.4	111.5	400.0	1,297.5	3,746.6	9,445.7
1956	297.2	6.6	0.9	1.3	4.0	9.2	29.0	109.1	394.2	1,278.7	3,743.5	9,275.1
1955	293.0	5.6	1.2	1.5	4.1	9.5	31.2	111.4	396.2	1,279.0	3,740.9	9,155.9
1954	284.1	6.2	1.0	1.9	4.3	10.1	32.1	118.9	402.9	1,282.7	3,628.6	8,655.7
1953	293.9	6.9	1.4	2.3	4.8	11.8	36.6	127.8	431.0	1,344.4	3,820.9	8,928.3
1952	290.5	6.6	1.3	2.7	5.3	12.0	38.8	132.0	438.6	1,349.2	3,820.3	8,842.6
1951	289.5	6.6	1.4	2.5	5.7	12.7	40.3	139.0	448.1	1,356.3	3,851.5	8,984.6
1950	290.5	2.9	1.4	3.7	6.6	13.6	42.4	142.4	460.7	1,401.6	3,926.2	9,086.9

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
DISEASES OF HEART (ICD 400-402, 410-413)												
ALL OTHER, BOTH SEXES												
1959	283.9	15.3	4.2	3.0	9.7	39.3	138.4	433.7	1,016.8	1,869.7	3,068.2	5,752.9
1958	295.9	15.5	2.9	2.8	9.5	41.4	140.2	466.7	1,084.5	1,937.7	3,162.3	5,766.7
1957	303.9	13.0	4.4	4.0	12.1	41.7	156.1	479.7	1,153.8	1,945.0	3,161.4	5,670.3
1956	292.6	16.1	4.2	3.6	11.9	39.5	148.0	462.0	1,121.4	1,873.3	3,130.2	5,401.6
1955	287.2	14.5	4.3	4.2	12.2	39.7	149.0	481.3	1,079.9	1,813.0	3,125.8	5,187.9
1954	284.4	17.7	3.6	4.5	14.3	43.3	156.0	496.3	1,077.8	1,759.6	2,989.3	5,052.7
1953	306.4	17.9	3.6	6.1	16.5	46.8	173.7	537.5	1,185.1	1,864.0	3,187.2	5,646.2
1952	309.2	17.9	3.6	6.7	17.5	49.7	177.4	549.9	1,228.3	1,869.2	3,194.2	5,518.4
1951	309.1	12.7	2.9	7.7	16.9	49.6	185.5	552.8	1,237.0	1,852.5	3,238.7	5,631.9
1950	311.8	6.2	3.7	7.6	18.8	50.5	186.4	561.0	1,195.8	2,113.8	3,340.2	5,716.2
ALL OTHER, MALE												
1959	317.2	12.0	5.2	2.7	9.7	43.8	161.8	506.7	1,167.7	2,153.0	3,382.6	6,224.1
1958	326.5	15.2	2.5	2.4	9.1	44.2	158.4	525.3	1,206.0	2,228.7	3,522.3	6,451.9
1957	336.2	15.0	4.4	3.8	11.9	44.8	175.8	532.9	1,297.6	2,211.1	3,445.9	6,669.2
1956	322.9	16.5	3.4	3.6	12.3	42.0	160.5	521.5	1,240.3	2,116.2	3,434.6	6,224.0
1955	319.4	12.6	4.5	3.9	12.0	40.2	166.6	540.0	1,219.8	2,063.5	3,401.6	5,720.8
1954	316.6	19.8	4.0	4.3	12.9	44.7	167.1	548.2	1,232.6	1,981.0	3,305.0	5,854.5
1953	343.6	17.9	4.3	5.5	15.9	51.5	186.4	596.6	1,348.7	2,110.2	3,521.1	6,485.7
1952	344.7	18.9	4.0	6.7	17.8	49.4	185.5	594.1	1,379.5	2,146.0	3,580.0	6,275.0
1951	341.0	15.0	3.3	6.7	15.2	49.4	194.7	591.4	1,367.7	2,112.2	3,571.7	6,315.8
1950	342.0	*	4.3	6.3	17.9	50.2	191.4	604.5	1,314.6	2,321.0	3,679.8	6,262.1
ALL OTHER, FEMALE												
1959	252.4	18.5	3.1	3.4	9.7	35.4	117.2	365.0	871.7	1,613.0	2,787.0	5,402.6
1958	266.9	15.9	3.3	3.1	9.8	39.0	123.3	411.7	970.1	1,668.2	2,835.3	5,431.6
1957	273.5	11.1	4.4	4.2	12.3	39.0	138.0	428.9	1,013.8	1,701.9	2,896.6	5,121.6
1956	263.8	15.7	4.9	3.6	11.6	37.4	136.5	404.2	1,004.6	1,641.5	2,843.5	4,830.6
1955	256.8	16.3	4.1	4.5	12.3	39.2	132.5	424.3	941.2	1,582.2	2,840.9	4,811.8
1954	254.0	15.6	3.2	4.7	15.6	42.0	145.6	445.6	922.9	1,555.1	2,667.5	4,659.4
1953	271.4	17.9	2.9	6.7	17.1	42.7	161.9	479.5	1,017.1	1,629.5	2,846.3	5,077.4
1952	275.5	16.9	3.3	6.7	17.3	50.0	169.9	506.2	1,072.9	1,609.5	2,828.4	4,996.6
1951	278.9	10.4	2.6	8.7	18.4	49.8	177.0	513.8	1,101.0	1,606.9	2,920.7	5,167.9
1950	283.0	*	3.1	8.9	19.6	50.9	181.7	517.6	1,073.9	1,905.1	3,005.7	5,326.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)												
BOTH SEXES												
1959	10.4	*	0.2	0.8	2.2	4.9	10.2	19.7	28.6	31.0	37.0	56.8
1958	10.8	*	0.2	0.8	2.2	5.2	10.5	20.8	29.2	32.1	42.2	62.9
1957	11.8	*	0.2	0.8	2.9	5.6	11.4	22.4	31.6	35.7	45.6	68.6
1956	12.0	*	0.3	1.1	3.1	5.7	11.3	22.2	31.0	35.3	49.0	80.3
1955	12.0	*	0.3	1.2	3.1	5.9	12.0	21.8	30.4	35.5	48.3	79.3
1954	12.1	*	0.4	1.6	3.6	6.3	11.9	21.8	29.8	34.8	50.4	79.0
1953	13.3	0.7	0.5	1.9	4.0	6.9	13.5	24.1	31.4	37.5	56.6	87.4
1952	13.7	*	0.5	2.2	4.4	7.1	14.6	23.1	31.9	37.9	62.5	98.9
1951	14.1	0.6	0.5	2.5	4.2	7.3	14.9	24.5	31.7	39.4	65.6	109.7
1950	14.8	*	0.5	2.9	4.6	7.8	15.5	24.1	32.8	42.3	74.0	136.4
MALE												
1959	10.1	*	0.3	0.7	2.2	4.8	10.5	20.1	28.9	31.5	34.7	53.5
1958	10.6	*	*	0.7	2.3	5.4	11.1	20.4	29.4	32.3	40.6	61.6
1957	11.7	*	0.3	0.7	2.8	5.4	11.9	22.5	33.1	35.9	45.8	62.9
1956	12.0	*	0.3	1.0	3.1	5.8	11.7	22.7	32.4	36.4	49.3	78.9
1955	12.1	*	0.4	1.1	3.1	5.8	12.7	22.0	32.1	37.1	48.9	70.8
1954	12.1	*	0.5	1.4	3.6	6.3	12.0	21.7	30.5	36.8	51.4	77.3
1953	13.4	*	0.5	1.7	4.2	6.9	13.4	24.3	33.6	40.4	57.7	80.6
1952	14.0	*	0.4	1.9	5.0	7.1	15.2	24.0	33.8	40.0	65.8	95.2
1951	14.3	*	0.6	2.3	4.3	7.5	15.3	24.6	33.2	42.1	66.0	111.7
1950	15.1	*	0.4	2.6	4.4	8.2	15.6	24.3	35.0	46.4	78.8	130.9
FEMALE												
1959	10.6	*	*	0.9	2.3	5.0	9.9	19.3	28.4	30.5	38.8	58.9
1958	11.1	*	0.3	0.8	2.1	5.0	9.9	21.2	29.0	31.8	43.4	63.8
1957	12.0	*	*	0.9	3.0	5.7	10.8	22.4	30.1	35.5	45.4	72.5
1956	12.0	*	0.3	1.1	3.1	5.5	11.0	21.7	29.6	34.4	48.8	81.3
1955	12.0	*	0.3	1.3	3.2	6.0	11.3	21.7	28.8	34.0	47.8	85.3
1954	12.2	*	0.3	1.7	3.5	6.2	11.7	22.0	29.0	33.0	49.6	80.2
1953	13.2	*	0.5	2.2	3.8	6.9	13.7	24.0	29.3	34.9	55.7	92.3
1952	13.4	*	0.5	2.5	3.9	7.1	14.0	22.2	29.9	36.0	59.7	101.6
1951	14.0	*	0.5	2.7	4.1	7.2	14.6	24.3	30.3	36.9	65.2	108.3
1950	14.5	*	0.6	3.3	4.8	7.4	15.3	23.8	30.5	38.6	69.8	140.3

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)												
WHITE, BOTH SEXES												
1959	10.7	*	*	0.6	1.9	4.5	10.2	20.1	29.6	31.9	37.5	57.0
1958	11.2	*	0.2	0.6	1.9	4.8	10.3	21.2	30.0	33.1	43.1	63.6
1957	12.1	*	0.2	0.6	2.5	5.1	11.2	22.7	32.2	36.9	46.3	68.8
1956	12.2	*	0.2	0.8	2.6	5.4	11.1	22.6	31.5	36.1	49.7	82.5
1955	12.3	*	0.2	0.9	2.7	5.7	11.8	22.0	31.0	36.0	49.2	80.0
1954	12.3	*	0.3	1.3	3.1	5.8	11.7	22.1	30.4	35.5	50.2	79.9
1953	13.5	*	0.4	1.6	3.5	6.6	13.4	24.4	31.8	37.6	57.1	85.3
1952	13.7	*	0.4	1.8	3.9	6.8	14.7	23.0	31.8	37.6	62.1	98.7
1951	14.2	*	0.4	2.0	3.8	7.0	15.0	24.7	31.9	39.4	65.4	106.2
1950	14.8	*	0.3	2.5	4.1	7.6	15.4	24.0	32.6	41.8	72.7	134.7
WHITE, MALE												
1959	10.3	*	*	0.5	1.8	4.5	10.5	20.4	29.8	32.2	35.1	52.5
1958	10.9	*	*	0.5	2.1	5.1	11.0	20.8	30.1	33.2	41.1	62.3
1957	11.9	*	*	0.5	2.6	5.0	11.7	22.7	33.7	36.9	46.7	62.1
1956	12.2	*	*	0.7	2.7	5.6	11.6	23.1	32.9	36.7	49.6	78.8
1955	12.3	*	*	0.9	2.7	5.8	12.6	22.1	32.7	37.7	49.8	72.1
1954	12.2	*	0.4	1.2	3.3	6.0	11.8	21.8	31.2	37.5	51.2	75.5
1953	13.6	*	0.4	1.5	3.7	6.7	13.4	24.7	34.0	40.4	58.1	78.0
1952	14.1	*	0.3	1.5	4.6	7.0	15.4	23.9	33.9	39.6	65.2	92.1
1951	14.4	*	0.5	1.9	4.1	7.3	15.3	24.9	33.4	42.2	65.2	108.4
1950	15.0	*	*	2.2	4.0	8.0	15.6	24.3	34.6	45.8	76.6	131.6
WHITE, FEMALE												
1959	11.0	*	*	0.6	2.0	4.5	9.9	19.9	29.5	31.7	39.4	59.9
1958	11.4	*	*	0.6	1.8	4.4	9.6	21.6	30.0	33.1	44.6	64.4
1957	12.3	*	*	0.6	2.5	5.2	10.7	22.7	30.7	36.8	46.0	73.3
1956	12.3	*	*	0.9	2.5	5.1	10.6	22.2	30.1	35.6	49.8	85.0
1955	12.2	*	*	1.0	2.7	5.7	11.0	21.9	29.5	34.5	48.6	85.5
1954	12.3	*	*	1.4	2.9	5.6	11.6	22.4	29.5	33.6	49.4	83.0
1953	13.3	*	0.4	1.7	3.3	6.5	13.3	24.0	29.6	35.1	56.3	90.6
1952	13.4	*	0.4	2.0	3.3	6.6	14.0	22.2	29.7	35.7	59.6	103.4
1951	14.1	*	0.4	2.1	3.5	6.8	14.7	24.5	30.5	36.9	65.6	104.5
1950	14.5	*	0.4	2.8	4.2	7.2	15.2	23.8	30.6	38.0	69.3	137.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)												
ALL OTHER, BOTH SEXES												
1959	7.9	*	*	2.3	4.7	7.5	10.5	15.9	18.2	19.4	30.2	54.4
1958	8.3	*	*	2.0	4.2	8.3	12.1	17.1	20.1	19.4	30.1	54.5
1957	9.7	*	*	2.4	5.6	9.4	13.1	19.8	25.2	21.7	35.4	65.6
1956	9.9	*	*	2.9	6.3	8.1	13.4	17.9	24.9	26.5	38.8	54.1
1955	10.1	*	1.1	3.0	6.2	7.6	13.3	20.2	23.6	28.9	35.9	70.7
1954	10.8	*	*	3.6	7.0	10.2	13.5	19.6	22.6	27.0	53.3	67.3
1953	12.3	*	1.4	4.5	7.7	9.4	15.2	21.9	26.7	36.5	49.1	113.5
1952	13.4	*	1.2	5.1	8.0	10.0	14.2	23.4	32.8	42.2	68.1	102.0
1951	13.1	*	1.3	6.2	6.9	10.0	14.4	22.1	29.8	39.1	67.7	153.2
1950	15.0	*	1.7	6.1	8.7	10.1	16.1	24.2	35.1	50.3	93.4	156.3
ALL OTHER, MALE												
1959	8.0	*	*	1.8	4.5	6.9	10.6	17.3	19.7	23.7	29.9	*
1958	8.1	*	*	1.6	3.6	7.7	12.0	16.6	21.7	22.0	33.8	*
1957	9.7	*	*	2.0	4.6	8.5	13.8	21.1	26.8	23.2	34.1	*
1956	10.0	*	*	2.8	5.9	7.7	11.9	18.9	26.3	32.9	44.6	80.0
1955	9.9	*	*	2.5	6.0	6.1	13.3	20.6	26.0	30.7	36.0	*
1954	10.9	*	*	3.1	6.0	9.3	14.6	21.1	22.6	29.0	52.9	100.0
1953	11.9	*	*	3.6	7.5	9.1	13.5	20.7	27.8	40.4	52.6	114.3
1952	13.4	*	*	4.7	8.3	8.0	14.1	24.9	33.3	44.5	74.5	135.0
1951	12.9	*	*	5.4	5.4	8.9	14.6	21.6	31.8	41.3	76.4	152.6
1950	15.3	*	*	4.9	8.1	10.4	16.0	24.4	40.4	53.6	110.0	123.2
ALL OTHER, FEMALE												
1959	7.7	*	*	2.7	4.8	8.1	10.3	14.5	16.9	15.6	30.4	*
1958	8.4	*	*	2.3	4.7	8.9	12.2	17.7	18.6	16.9	26.8	55.3
1957	9.8	*	*	2.7	6.5	10.2	12.5	18.6	23.7	20.2	36.6	62.2
1956	9.7	*	*	3.0	6.7	8.5	14.7	16.9	23.6	20.5	33.3	*
1955	10.2	*	*	3.5	6.3	9.0	13.3	19.8	21.1	27.2	35.6	82.4
1954	10.7	*	*	4.1	7.8	10.9	12.5	18.0	22.6	25.2	53.2	*
1953	12.6	*	*	5.4	7.8	9.6	16.7	23.2	25.5	32.8	45.5	112.9
1952	13.3	*	*	5.6	7.7	11.8	14.3	21.9	32.2	40.1	62.1	79.3
1951	13.3	*	*	7.0	8.2	10.9	14.3	22.6	27.7	37.0	59.5	153.6
1950	14.6	*	*	7.2	9.3	9.8	16.3	24.0	29.6	47.0	77.1	180.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)												
BOTH SEXES												
1959	268.6	*	*	0.1	0.8	7.2	49.9	206.9	571.9	1,332.3	2,912.5	6,119.6
1958	266.2	*	*	0.1	0.7	7.4	50.3	205.7	574.7	1,329.6	2,920.7	6,048.4
1957	265.6	1.2	*	0.1	0.9	7.0	50.5	202.7	580.1	1,340.1	2,932.4	6,054.4
1956	255.5	1.6	*	0.1	0.9	6.9	49.3	197.4	563.7	1,297.9	2,868.6	5,732.9
1955	247.0	1.3	0.1	0.1	1.0	7.0	48.7	196.9	549.7	1,265.8	2,780.4	5,467.7
1954	235.7	2.1	0.1	0.1	0.9	6.7	47.8	198.3	540.9	1,225.2	2,624.3	5,003.4
1953	236.1	1.7	0.2	0.2	1.1	7.1	49.4	200.4	556.2	1,239.6	2,645.2	4,940.5
1952	226.2	1.4	*	0.2	1.1	7.1	48.5	200.1	550.6	1,189.5	2,532.0	4,649.1
1951	219.6	1.1	*	0.1	1.3	6.7	48.0	196.6	540.3	1,175.0	2,479.9	4,562.8
1950	213.0	1.2	*	0.1	1.1	6.4	46.7	196.0	534.4	1,180.9	2,423.9	4,424.0
MALE												
1959	339.7	*	*	*	1.1	11.7	85.6	343.7	875.5	1,820.5	3,499.2	6,606.5
1958	337.9	*	*	*	1.0	11.6	85.9	339.5	876.0	1,810.1	3,513.7	6,570.6
1957	336.4	1.7	*	0.2	1.1	11.0	86.0	335.0	881.6	1,804.5	3,490.2	6,473.6
1956	325.4	1.6	*	*	1.3	11.0	84.2	327.1	856.0	1,745.2	3,414.2	6,118.3
1955	316.2	1.2	*	*	1.3	11.1	82.5	324.6	834.3	1,700.6	3,311.6	5,777.0
1954	304.5	2.4	*	0.2	1.1	10.7	80.6	325.6	818.2	1,642.8	3,130.6	5,292.8
1953	305.9	2.0	*	0.2	1.2	11.0	83.3	327.2	835.5	1,650.0	3,155.3	5,296.2
1952	293.6	1.7	*	0.2	1.3	11.0	81.2	324.6	825.1	1,574.0	3,001.4	4,946.9
1951	286.7	1.3	*	0.2	1.5	10.3	80.0	318.6	804.7	1,564.0	2,957.7	4,981.6
1950	277.8	1.6	*	*	1.3	9.7	77.2	316.6	792.9	1,560.7	2,873.5	4,907.8
FEMALE												
1959	199.5	*	*	*	0.5	2.9	15.9	74.2	286.5	905.4	2,455.1	5,801.3
1958	196.5	*	*	*	0.5	3.4	16.3	75.5	290.2	906.1	2,453.9	5,700.2
1957	196.8	*	*	*	0.7	3.2	16.4	73.7	293.9	927.1	2,489.1	5,768.4
1956	187.5	1.6	*	0.1	0.6	3.0	15.8	70.6	284.6	897.4	2,431.5	5,464.7
1955	179.6	1.3	*	*	0.7	3.1	16.0	71.7	275.9	873.6	2,351.4	5,248.6
1954	168.8	1.8	*	*	0.7	3.0	16.2	73.2	272.2	846.3	2,211.4	4,796.5
1953	168.3	1.4	*	*	0.9	3.4	16.7	75.6	283.5	864.9	2,225.6	4,684.8
1952	160.5	*	*	0.2	0.9	3.4	16.8	77.1	280.0	836.5	2,143.0	4,435.7
1951	153.9	*	*	*	1.1	3.3	17.0	75.7	277.3	815.5	2,078.2	4,265.0
1950	149.1	*	*	*	0.8	3.4	16.9	76.0	275.0	826.8	2,041.4	4,087.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)												
WHITE, BOTH SEXES												
1959	283.0	*	*	0.1	0.6	6.5	48.2	204.1	571.3	1,355.9	2,999.2	6,370.5
1958	280.2	*	*	0.1	0.6	6.5	48.9	202.2	574.2	1,353.5	3,009.3	6,304.4
1957	279.3	1.0	*	0.1	0.7	6.1	48.3	198.8	577.8	1,366.8	3,022.6	6,301.4
1956	269.1	1.3	*	0.1	0.8	6.2	47.5	194.8	563.0	1,325.3	2,959.4	5,986.1
1955	260.4	0.9	*	0.1	0.8	6.2	47.3	193.7	551.1	1,297.2	2,868.3	5,715.1
1954	248.6	1.8	*	0.1	0.7	6.1	46.1	195.9	543.1	1,259.0	2,710.7	5,220.4
1953	248.4	1.4	0.2	0.1	0.8	6.2	47.7	196.5	558.0	1,272.2	2,730.8	5,152.4
1952	238.4	1.3	*	0.1	0.9	6.2	46.8	197.7	553.0	1,224.9	2,616.9	4,848.6
1951	231.5	1.0	*	0.1	1.1	6.0	46.0	194.5	543.2	1,211.2	2,564.7	4,773.8
1950	224.6	1.2	*	0.1	0.9	5.7	45.1	194.6	540.4	1,208.9	2,502.7	4,634.5
WHITE, MALE												
1959	358.8	*	*	*	1.0	10.9	85.2	348.3	890.8	1,865.7	3,617.2	6,883.1
1958	356.8	*	*	*	0.9	10.8	86.5	343.6	892.3	1,855.3	3,633.3	6,862.0
1957	354.7	1.5	*	0.1	0.9	10.2	85.6	339.0	894.9	1,853.5	3,610.6	6,712.2
1956	344.0	1.3	*	*	1.2	10.5	84.3	332.2	871.1	1,795.5	3,538.4	6,364.9
1955	334.5	*	*	*	1.1	10.7	83.0	328.9	850.4	1,753.5	3,433.8	6,024.8
1954	322.0	2.2	*	0.2	1.0	10.2	81.1	331.0	833.7	1,696.7	3,244.3	5,495.0
1953	322.9	1.8	*	0.2	1.1	10.2	83.3	330.9	850.7	1,704.1	3,268.6	5,486.6
1952	310.4	1.6	*	*	1.2	10.3	81.6	330.2	840.9	1,628.0	3,113.8	5,136.4
1951	303.3	*	*	*	1.3	9.8	79.8	324.2	821.2	1,619.9	3,074.0	5,189.0
1950	294.0	1.6	*	*	1.1	9.0	77.5	323.1	812.9	1,608.2	2,978.7	5,139.8
WHITE, FEMALE												
1959	209.2	*	*	*	0.3	2.2	12.6	63.7	271.6	911.2	2,522.2	6,038.9
1958	205.6	*	*	*	0.3	2.2	12.7	64.2	274.4	912.3	2,523.2	5,935.8
1957	205.6	*	*	*	0.5	2.1	12.4	61.8	277.3	935.4	2,559.1	6,021.9
1956	196.0	1.3	*	*	0.5	2.0	11.8	60.2	269.3	905.2	2,500.8	5,722.6
1955	188.1	*	*	*	0.4	1.9	12.7	61.1	264.0	886.3	2,417.9	5,495.7
1954	177.0	1.4	*	*	0.5	2.1	12.2	62.9	262.2	862.3	2,279.8	5,023.4
1953	175.8	*	*	*	0.6	2.3	13.1	64.0	273.0	878.5	2,293.9	4,911.1
1952	168.0	*	*	*	0.7	2.4	12.9	66.7	270.2	856.0	2,208.9	4,641.8
1951	161.0	*	*	*	0.8	2.4	13.0	65.8	267.7	834.9	2,142.0	4,477.4
1950	156.0	*	*	*	0.7	2.6	13.2	66.6	267.5	838.9	2,101.6	4,283.4

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)												
ALL OTHER, BOTH SEXES												
1959	154.0	*	*	*	1.7	13.0	65.2	233.4	577.8	1,050.9	1,713.4	3,057.4
1958	153.3	*	*	*	1.5	15.0	62.8	239.0	579.9	1,043.9	1,687.7	2,968.2
1957	154.6	*	*	*	2.1	13.9	69.6	239.9	606.5	1,020.5	1,673.6	2,995.3
1956	144.4	3.7	*	*	1.7	12.7	65.5	222.5	572.4	970.2	1,580.2	2,677.0
1955	136.2	*	*	*	2.6	13.7	60.5	227.6	533.0	890.3	1,528.1	2,443.1
1954	128.0	4.2	*	*	2.0	12.3	62.9	221.4	515.2	819.6	1,379.1	2,249.1
1953	132.5	*	*	*	3.1	14.9	64.2	238.2	534.6	848.9	1,392.3	2,336.5
1952	123.3	*	*	*	2.2	14.5	63.7	222.9	521.2	765.3	1,286.3	2,185.7
1951	118.2	*	*	*	2.8	12.7	65.3	217.1	504.0	737.9	1,225.3	2,008.5
1950	113.4	*	*	*	2.3	12.5	60.7	209.9	461.6	800.5	1,232.9	1,922.5
ALL OTHER, MALE												
1959	185.9	*	*	*	1.6	17.9	89.1	298.9	715.9	1,290.7	1,991.7	3,496.6
1958	183.1	*	*	*	1.7	18.2	80.7	300.0	703.5	1,277.8	1,982.7	3,403.7
1957	184.6	*	*	*	2.3	17.3	89.8	296.6	736.7	1,232.7	1,942.2	3,619.2
1956	171.3	*	*	*	2.1	15.3	82.9	277.6	687.1	1,152.6	1,808.5	3,140.0
1955	162.9	*	*	*	2.5	14.9	77.7	283.1	651.8	1,076.0	1,715.2	2,741.7
1954	155.7	*	*	*	2.0	15.0	76.6	272.9	638.5	1,002.8	1,616.8	2,700.0
1953	160.6	*	*	*	2.4	18.2	82.5	291.4	652.4	1,006.1	1,628.1	2,866.7
1952	149.6	*	*	*	2.1	17.5	78.0	270.5	635.0	940.6	1,473.6	2,550.0
1951	143.1	*	*	*	2.5	15.2	81.7	264.7	602.4	904.0	1,387.7	2,394.7
1950	136.6	*	*	*	3.1	15.2	74.3	254.2	554.1	943.1	1,407.3	2,196.3
ALL OTHER, FEMALE												
1959	123.8	*	*	*	1.8	8.8	43.6	171.6	445.0	833.6	1,464.6	2,730.8
1958	125.2	*	*	*	*	12.2	46.4	181.5	462.0	828.5	1,419.6	2,736.8
1957	126.2	*	*	*	1.9	11.0	51.0	185.8	479.6	826.7	1,423.4	2,637.8
1956	118.8	*	*	*	*	10.5	49.5	169.2	459.8	798.3	1,365.2	2,355.6
1955	110.9	*	*	*	2.7	12.6	44.5	173.8	415.1	719.3	1,339.4	2,232.4
1954	101.9	*	*	*	2.1	9.9	50.1	171.2	391.9	650.4	1,143.7	2,009.4
1953	106.1	*	*	*	3.6	12.1	47.2	185.9	414.3	700.0	1,158.7	1,977.4
1952	98.5	*	*	*	2.3	11.8	50.4	175.8	404.3	600.8	1,108.6	1,934.5
1951	94.6	*	*	*	3.0	10.4	50.2	169.5	401.6	580.9	1,070.3	1,746.4
1950	91.2	*	*	*	1.6	10.2	48.1	165.7	366.7	656.8	1,061.1	1,726.8

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)												
BOTH SEXES												
1959	31.8	1.1	0.2	0.1	0.3	1.0	2.6	9.6	32.8	116.9	466.9	1,596.9
1958	33.8	1.3	0.3	0.1	0.3	1.0	3.0	10.6	36.5	127.6	499.0	1,690.1
1957	36.8	1.6	0.4	0.2	0.5	1.1	3.3	12.5	41.7	144.7	537.7	1,802.5
1956	37.7	1.7	0.4	0.1	0.5	1.1	3.5	12.8	42.7	149.9	566.8	1,858.8
1955	39.9	1.9	0.4	0.1	0.5	1.2	4.2	14.6	45.2	158.7	614.5	1,957.6
1954	41.0	1.6	0.3	0.2	0.5	1.2	4.5	16.0	48.8	169.5	631.9	1,979.1
1953	46.3	1.7	0.3	0.2	0.6	1.6	5.1	18.1	57.3	195.4	725.1	2,250.7
1952	49.3	2.3	0.4	0.3	0.7	1.6	6.1	19.6	62.9	212.9	785.7	2,346.6
1951	52.6	1.4	0.3	0.2	0.7	1.9	6.4	21.9	69.5	229.8	846.7	2,604.2
1950	56.5	...	0.3	0.3	0.9	1.9	7.4	24.8	78.4	256.7	918.2	2,849.2
MALE												
1959	31.9	*	0.3	0.1	0.4	1.2	3.1	12.1	42.2	139.8	506.6	1,596.9
1958	33.8	1.4	0.3	0.1	0.4	1.3	3.8	13.3	45.1	151.5	537.4	1,652.9
1957	37.4	1.5	0.4	0.2	0.6	1.2	4.1	15.6	52.8	172.4	584.7	1,771.8
1956	38.6	1.8	0.4	0.2	0.5	1.4	4.4	16.3	52.8	179.1	623.3	1,800.3
1955	40.9	1.8	0.4	0.2	0.5	1.4	5.1	18.1	56.2	189.5	666.6	1,902.2
1954	42.6	1.8	0.4	0.2	0.7	1.5	5.3	19.3	61.2	201.5	693.9	1,928.6
1953	48.3	2.0	0.3	0.2	0.7	1.7	6.1	23.2	72.3	230.4	780.7	2,223.9
1952	51.6	2.4	0.5	0.3	0.8	1.7	7.1	23.8	78.2	254.0	858.9	2,289.0
1951	55.5	1.5	*	0.2	0.9	2.1	7.3	26.5	86.2	275.5	919.6	2,646.1
1950	59.4	...	0.3	0.3	0.8	1.9	8.4	31.1	97.3	303.1	986.2	2,901.3
FEMALE												
1959	31.7	1.2	*	*	0.2	0.8	2.1	7.1	23.9	96.9	436.1	1,596.9
1958	33.9	1.3	0.3	*	0.3	0.8	2.3	8.0	28.5	106.6	468.9	1,714.9
1957	36.2	1.7	0.5	0.2	0.3	1.0	2.6	9.5	31.2	120.0	500.4	1,823.5
1956	36.8	1.6	0.4	*	0.4	0.9	2.6	9.4	33.1	123.8	521.6	1,899.6
1955	39.0	2.0	0.3	*	0.4	1.1	3.2	11.1	34.7	131.0	572.5	1,996.9
1954	39.5	1.4	0.3	0.2	0.4	1.0	3.7	12.7	36.7	140.5	581.2	2,015.3
1953	44.5	1.4	0.4	0.2	0.4	1.4	4.1	13.0	42.6	163.3	679.1	2,269.9
1952	46.9	2.3	0.3	0.3	0.6	1.6	5.1	15.4	47.9	175.2	725.0	2,387.9
1951	49.7	1.3	0.3	0.2	0.6	1.7	5.5	17.2	53.0	187.6	785.1	2,574.4
1950	53.7	...	*	0.3	1.0	1.9	6.4	18.5	59.5	213.4	860.4	2,812.9

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)												
WHITE, BOTH SEXES												
1959	32.0	0.8	0.2	0.1	0.2	0.6	1.8	7.5	27.5	109.1	469.7	1,639.4
1958	33.8	1.2	0.2	0.1	0.3	0.7	2.1	8.1	30.0	118.7	500.8	1,732.0
1957	36.7	1.3	0.3	0.1	0.4	0.7	2.3	9.4	34.8	134.6	540.5	1,860.4
1956	37.7	1.6	0.2	0.1	0.4	0.7	2.4	9.6	35.9	141.2	569.0	1,917.5
1955	40.1	1.7	0.3	0.1	0.4	0.8	2.9	10.9	38.4	150.6	619.8	2,025.2
1954	41.0	1.4	0.3	0.1	0.5	0.8	3.1	11.9	41.6	160.9	636.8	2,037.8
1953	46.4	1.6	0.3	0.2	0.4	1.1	3.6	13.8	49.4	186.1	731.7	2,322.7
1952	49.2	2.1	0.3	0.2	0.5	1.1	4.3	14.9	54.0	203.7	792.1	2,420.5
1951	52.6	1.2	0.3	0.2	0.5	1.3	4.6	16.5	59.5	220.5	854.0	2,692.3
1950	56.5	...	0.2	0.3	0.7	1.3	5.3	19.4	68.5	245.4	926.1	2,945.0
WHITE, MALE												
1959	31.8	*	*	0.1	0.3	0.8	2.4	10.2	36.7	131.3	510.1	1,638.0
1958	33.4	1.2	*	0.1	0.3	1.0	2.9	10.8	38.4	141.7	539.3	1,688.3
1957	37.0	1.2	0.3	*	0.5	0.8	3.1	12.2	45.3	162.2	590.3	1,818.6
1956	38.4	1.7	*	0.1	0.4	1.0	3.4	13.0	46.1	170.2	625.7	1,845.4
1955	40.8	1.6	0.3	0.2	0.5	1.0	3.9	14.8	49.1	181.1	671.2	1,955.4
1954	42.4	1.7	0.3	*	0.6	1.1	4.1	15.5	53.3	193.1	697.9	1,974.8
1953	47.9	1.9	*	0.2	0.6	1.2	4.7	18.4	63.2	220.5	785.8	2,284.7
1952	51.3	2.1	0.4	0.3	0.6	1.3	5.5	19.3	68.9	244.1	864.1	2,347.4
1951	55.1	*	*	0.2	0.6	1.5	5.6	20.9	75.6	265.4	926.9	2,725.3
1950	59.2	...	*	0.2	0.7	1.3	6.7	25.4	87.4	292.6	994.6	2,993.2
WHITE, FEMALE												
1959	32.1	*	*	*	*	0.4	1.3	4.9	18.8	89.7	438.6	1,640.3
1958	34.1	1.2	*	*	0.3	0.4	1.4	5.5	22.2	98.5	470.9	1,760.9
1957	36.4	1.4	0.4	*	*	0.6	1.5	6.6	24.9	110.1	501.1	1,888.8
1956	37.1	1.5	*	*	0.3	0.5	1.5	6.4	26.2	115.3	524.0	1,967.7
1955	39.4	1.8	*	*	0.3	0.6	2.0	7.1	28.2	123.2	579.1	2,074.7
1954	39.7	*	*	0.2	0.3	0.6	2.2	8.3	30.3	131.8	587.2	2,083.0
1953	45.0	*	*	*	0.3	1.0	2.5	9.2	36.0	154.7	687.8	2,350.1
1952	47.2	2.1	0.3	0.2	0.4	0.9	3.2	10.4	39.4	166.8	733.1	2,473.0
1951	50.0	*	*	*	0.5	1.1	3.6	12.2	43.6	179.2	793.6	2,668.7
1950	53.9	...	*	0.3	0.8	1.4	4.0	13.4	49.6	201.6	868.5	2,911.6

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)												
ALL OTHER, BOTH SEXES												
1959	30.2	*	*	*	*	4.3	9.3	29.0	88.9	210.4	428.2	1,077.9
1958	34.0	*	*	*	*	3.8	10.9	34.6	106.1	234.6	474.0	1,186.4
1957	37.2	*	1.1	*	1.2	3.9	12.1	42.7	117.9	265.9	501.1	1,079.7
1956	37.5	*	1.7	*	1.1	4.3	12.7	43.3	118.9	254.0	536.6	1,150.8
1955	38.4	*	*	*	0.8	4.7	14.7	50.0	123.8	255.6	540.6	1,131.0
1954	40.9	*	*	*	1.1	4.7	16.1	55.4	133.5	272.3	563.9	1,223.6
1953	45.5	*	*	*	1.6	5.9	18.2	59.1	153.2	306.2	627.8	1,365.4
1952	49.5	*	*	*	2.2	6.3	21.5	64.8	174.5	322.8	691.6	1,434.7
1951	53.0	*	*	*	2.2	6.9	22.2	72.5	196.8	340.4	737.8	1,538.3
1950	56.3	...	*	*	2.5	7.3	25.6	77.1	197.9	409.7	798.7	1,710.4
ALL OTHER, MALE												
1959	32.2	*	*	*	*	5.2	9.7	30.9	99.2	239.4	461.8	1,134.5
1958	36.5	*	*	*	*	4.1	12.0	38.2	116.1	265.9	512.9	1,300.0
1957	40.1	*	*	*	*	4.6	12.8	48.6	134.1	293.1	513.3	1,211.5
1956	40.5	*	*	*	*	5.0	13.9	48.3	126.7	283.6	592.3	1,256.0
1955	41.8	*	*	*	*	4.7	16.2	50.7	136.9	288.4	606.4	1,250.0
1954	44.6	*	*	*	*	5.2	16.3	56.1	153.2	301.1	641.2	1,336.4
1953	51.4	*	*	*	*	6.9	18.2	69.1	180.7	346.8	712.3	1,447.6
1952	54.3	*	*	*	2.3	5.3	21.6	66.5	192.4	371.9	789.1	1,550.0
1951	58.9	*	*	*	2.9	7.2	22.8	79.6	217.4	394.2	828.3	1,657.9
1950	61.0	...	*	*	2.0	7.8	24.1	86.4	215.8	439.0	870.2	1,826.7
ALL OTHER, FEMALE												
1959	28.2	*	*	*	*	3.6	9.0	27.2	78.9	184.2	398.1	1,035.9
1958	31.7	*	*	*	*	3.6	9.7	31.1	96.8	205.6	438.6	1,136.8
1957	34.6	*	*	*	*	3.4	11.5	37.2	102.1	241.0	489.7	1,016.2
1956	34.6	*	2.0	*	*	3.7	11.6	38.4	111.2	225.7	484.1	1,077.8
1955	35.3	*	*	*	*	4.6	13.3	49.3	110.8	225.4	474.2	1,047.1
1954	37.5	*	*	*	*	4.2	16.0	54.8	113.8	245.7	486.5	1,184.4
1953	40.0	*	*	*	1.7	5.0	18.2	49.3	125.0	267.5	543.0	1,309.7
1952	44.9	*	*	*	2.1	7.2	21.4	63.1	156.2	276.8	599.1	1,355.2
1951	47.4	*	*	*	1.6	6.6	21.7	65.4	175.4	289.6	651.4	1,457.1
1950	51.8	...	*	*	2.9	6.8	26.9	67.8	179.6	380.3	728.2	1,627.3

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES OF HEART (430-434)												
BOTH SEXES												
1959	13.8	5.5	1.0	0.3	0.9	2.1	4.6	12.2	26.0	57.3	139.5	346.7
1958	14.2	5.5	0.8	0.3	0.9	1.8	4.5	12.0	27.4	59.7	145.8	368.6
1957	12.8	4.8	0.8	0.4	0.9	1.8	4.2	11.2	25.7	55.8	127.2	302.4
1956	12.6	4.3	0.6	0.3	0.8	1.7	3.9	10.9	25.7	56.9	128.4	293.2
1955	12.6	3.8	0.8	0.3	0.8	1.6	4.3	11.4	25.5	57.0	130.1	284.2
1954	13.0	4.4	0.6	0.3	1.1	1.8	4.6	12.3	27.2	58.5	131.5	295.1
1953	14.3	5.4	0.6	0.4	1.0	1.9	5.5	13.6	30.7	63.7	146.1	329.1
1952	14.6	5.0	0.7	0.3	1.2	2.1	5.7	14.6	31.0	66.3	149.4	324.5
1951	15.1	5.1	0.6	0.3	1.2	2.1	6.0	15.1	32.1	69.8	158.7	354.9
1950	15.9	2.2	0.7	0.5	1.2	2.4	6.2	16.0	35.1	74.4	169.3	386.4
MALE												
1959	16.4	5.9	1.1	0.4	1.0	2.1	6.0	16.9	35.9	76.6	167.5	382.3
1958	16.8	5.9	0.8	0.4	1.0	1.8	5.9	16.6	37.9	78.4	178.4	412.2
1957	15.4	5.3	0.9	0.4	1.1	1.9	5.5	15.1	35.8	74.8	155.6	329.1
1956	15.2	4.5	0.6	0.3	0.9	1.8	4.6	15.0	35.6	75.1	160.2	324.2
1955	15.5	4.1	0.8	0.4	0.9	1.9	5.5	15.7	35.2	78.7	160.2	307.2
1954	15.9	4.9	0.6	0.3	1.3	2.1	5.8	17.0	38.0	76.8	157.3	337.2
1953	17.7	6.0	0.6	0.4	1.2	2.2	7.0	19.1	42.8	86.2	176.8	352.9
1952	18.0	5.6	0.8	0.4	1.3	2.2	7.1	20.2	43.7	88.6	179.9	353.5
1951	18.5	5.7	0.5	0.3	1.4	2.3	7.3	20.5	43.5	92.0	194.4	388.7
1950	19.3	2.4	0.7	0.5	1.3	2.6	7.9	21.2	47.1	97.4	201.8	451.8
FEMALE												
1959	11.3	5.1	0.9	0.3	0.8	2.0	3.3	7.6	16.7	40.4	117.7	323.4
1958	11.6	5.1	0.8	0.3	0.9	1.8	3.3	7.5	17.6	43.2	120.1	339.5
1957	10.2	4.2	0.8	0.3	0.8	1.6	2.9	7.4	16.2	38.9	104.7	284.2
1956	10.0	4.1	0.6	0.2	0.7	1.6	3.2	6.9	16.2	40.6	103.0	271.7
1955	9.7	3.4	0.8	0.3	0.7	1.3	3.1	7.3	16.2	37.4	105.8	267.9
1954	10.1	3.9	0.6	0.3	0.9	1.5	3.4	7.7	16.6	41.8	110.6	264.9
1953	11.0	4.7	0.6	0.4	0.9	1.7	4.0	8.2	18.9	43.2	120.7	311.9
1952	11.2	4.2	0.6	0.3	1.1	1.9	4.4	9.1	18.5	45.8	124.1	303.7
1951	11.8	4.5	0.7	0.3	1.1	1.9	4.7	9.7	20.8	49.3	128.7	330.8
1950	12.5	2.1	0.7	0.5	1.1	2.3	4.6	10.8	23.1	53.0	141.7	340.8

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES OF HEART (430-434)												
WHITE, BOTH SEXES												
1959	13.1	4.5	0.7	0.3	0.7	1.5	3.6	10.0	22.4	52.7	136.9	349.8
1958	13.3	4.5	0.7	0.3	0.7	1.3	3.4	9.5	23.0	54.7	143.2	374.4
1957	11.8	4.3	0.6	0.3	0.7	1.3	3.1	8.8	21.5	50.3	124.6	302.0
1956	11.7	3.6	0.5	0.3	0.6	1.2	2.9	8.3	22.0	52.2	126.0	292.8
1955	11.7	3.2	0.6	0.3	0.7	1.2	3.2	8.9	21.8	52.3	126.2	281.8
1954	12.1	3.7	0.4	0.3	0.8	1.3	3.4	9.6	23.7	54.3	128.5	294.5
1953	13.2	4.6	0.5	0.3	0.7	1.4	4.1	10.7	26.3	59.0	142.3	323.0
1952	13.4	4.2	0.6	0.3	0.9	1.4	4.2	11.1	26.4	61.4	145.8	317.7
1951	13.9	4.8	0.5	0.2	0.9	1.5	4.2	11.4	27.4	64.7	154.3	351.0
1950	14.6	2.2	0.7	0.5	0.9	1.7	4.6	12.3	30.2	68.2	165.1	385.4
WHITE, MALE												
1959	15.5	5.4	0.7	0.3	0.8	1.6	5.0	14.4	32.2	71.2	164.7	383.4
1958	15.8	5.1	0.6	0.4	0.8	1.4	4.6	13.7	32.8	72.2	173.8	414.9
1957	14.4	4.7	0.6	0.3	0.9	1.5	4.3	12.5	31.1	68.7	152.7	325.4
1956	14.3	3.6	0.5	0.3	0.7	1.5	3.6	12.1	31.5	70.2	156.8	317.2
1955	14.5	3.7	0.6	0.3	0.8	1.4	4.4	12.9	31.3	72.5	156.7	299.3
1954	15.0	3.8	0.5	0.3	1.1	1.6	4.6	13.8	34.3	72.4	154.2	331.9
1953	16.5	5.5	0.5	0.3	0.9	1.7	5.5	15.8	37.5	80.6	172.8	342.9
1952	16.7	5.0	0.7	0.3	1.0	1.6	5.7	16.4	38.1	83.2	174.4	340.3
1951	17.2	5.4	0.4	0.3	1.0	1.7	5.6	16.5	38.2	86.4	189.7	382.3
1950	18.1	2.5	0.6	0.5	1.1	2.1	6.2	17.5	41.9	90.3	197.0	447.4
WHITE, FEMALE												
1959	10.6	3.5	0.7	0.3	0.6	1.4	2.3	5.7	13.3	36.7	115.5	328.0
1958	10.9	3.8	0.7	0.3	0.6	1.2	2.3	5.3	13.7	39.3	119.3	347.7
1957	9.3	3.8	0.6	0.3	0.6	1.1	1.9	5.2	12.4	34.0	102.4	286.0
1956	9.2	3.5	0.4	0.2	0.6	1.0	2.2	4.6	12.9	36.2	101.6	275.8
1955	8.9	2.6	0.6	0.3	0.6	0.9	2.0	4.9	12.7	34.1	101.9	269.4
1954	9.3	3.5	0.4	0.3	0.5	0.9	2.2	5.4	13.5	37.8	107.7	267.7
1953	10.0	3.7	0.5	0.3	0.6	1.1	2.7	5.6	15.3	39.2	117.6	308.6
1952	10.1	3.5	0.5	0.2	0.7	1.2	2.7	5.9	14.9	41.4	122.3	301.4
1951	10.6	4.2	0.6	*	0.7	1.3	2.8	6.3	16.6	44.7	124.9	328.6
1950	11.2	1.8	0.7	0.4	0.7	1.4	3.0	7.0	18.4	47.7	138.1	342.4

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES OF HEART (430-434)												
ALL OTHER, BOTH SEXES												
1959	19.7	11.3	2.8	0.5	2.1	6.6	13.3	33.0	63.3	111.8	175.4	308.8
1958	21.2	11.7	1.7	0.5	2.7	6.1	14.2	36.0	74.8	120.1	181.5	298.5
1957	20.5	7.8	2.3	0.8	2.5	5.7	13.7	34.3	72.1	122.0	165.0	303.1
1956	19.5	9.0	1.5	*	2.2	5.6	12.6	35.6	67.7	112.7	162.7	298.4
1955	20.1	7.4	2.1	0.7	1.9	5.2	13.5	36.1	68.6	112.8	187.1	313.8
1954	20.3	9.2	1.5	*	3.1	6.1	14.8	39.0	68.4	108.5	177.0	296.4
1953	23.3	10.0	*	0.9	3.0	6.7	17.6	41.8	85.3	120.5	200.9	403.8
1952	24.6	9.8	1.6	*	3.7	7.7	19.2	48.0	88.1	124.7	201.8	408.2
1951	25.8	7.2	1.2	0.8	3.9	7.4	21.9	50.3	92.3	130.8	224.0	402.1
1950	26.6	*	*	0.9	3.7	8.5	20.8	51.4	94.4	158.5	233.6	397.5
ALL OTHER, MALE												
1959	23.0	8.7	3.5	*	2.3	6.5	15.4	41.0	73.9	140.2	203.5	369.0
1958	25.6	11.1	1.8	*	2.5	6.0	16.7	44.4	91.2	151.9	236.7	396.3
1957	24.1	9.4	2.6	*	2.7	5.8	16.3	40.7	86.5	147.0	192.6	373.1
1956	22.8	9.9	*	*	2.5	4.8	13.2	42.3	82.0	133.2	204.6	408.0
1955	23.9	*	2.2	*	1.9	5.9	14.8	42.6	79.4	152.2	206.4	404.2
1954	23.9	12.1	*	*	3.1	6.7	15.9	48.1	82.3	129.3	198.3	404.5
1953	28.0	9.0	*	*	3.2	6.8	20.3	50.9	106.1	151.2	231.6	481.0
1952	29.4	10.4	*	*	3.6	8.0	19.5	56.5	111.5	152.2	253.6	520.0
1951	29.5	*	*	*	3.8	7.3	22.5	57.9	108.5	157.8	260.4	468.4
1950	30.3	*	*	*	3.3	7.0	23.3	56.1	108.7	189.9	267.6	503.5
ALL OTHER, FEMALE												
1959	16.6	13.9	2.0	*	1.8	6.7	11.5	25.5	53.1	86.0	150.3	264.1
1958	17.1	12.4	*	*	2.9	6.3	11.9	28.1	59.1	90.8	131.4	236.8
1957	17.2	*	1.9	*	2.2	5.6	11.3	28.2	58.1	99.3	139.3	262.2
1956	16.3	8.0	*	*	1.9	6.4	12.0	29.1	53.7	93.3	123.2	222.2
1955	16.5	8.0	*	*	2.0	4.5	12.2	29.9	57.9	76.6	167.4	250.0
1954	16.9	*	*	*	3.1	5.5	13.6	30.1	54.5	89.2	155.6	231.3
1953	18.9	11.1	*	*	2.8	6.7	15.1	32.9	64.0	91.6	170.2	351.6
1952	20.1	9.3	*	*	3.8	7.5	18.8	39.6	64.0	98.9	152.6	331.0
1951	22.3	*	*	*	4.0	7.6	21.3	42.7	75.4	105.2	189.2	357.1
1950	23.2	*	*	*	4.0	9.8	18.5	46.8	79.8	126.9	200.1	321.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HYPERTENSIVE HEART DISEASE (440-443)												
BOTH SEXES												
1959	38.6	*	*	*	0.2	1.3	6.4	24.2	69.3	183.6	474.9	1,046.9
1958	42.7	*	*	*	0.2	1.3	6.9	27.3	78.9	205.7	530.6	1,130.0
1957	42.5	*	*	*	0.2	1.5	8.0	29.3	82.6	206.8	512.5	1,086.4
1956	43.3	*	*	*	0.2	1.5	7.4	29.6	83.3	210.5	545.9	1,114.6
1955	45.0	*	*	*	0.2	1.4	8.4	30.5	88.1	222.8	571.0	1,128.4
1954	46.6	*	*	*	0.2	1.9	8.9	34.7	93.5	237.1	578.4	1,115.9
1953	51.3	*	*	*	0.3	1.9	10.9	40.8	108.2	258.4	640.3	1,181.5
1952	53.9	*	*	*	0.4	2.1	11.7	44.4	114.4	277.3	672.1	1,242.7
1951	55.6	*	*	*	0.4	2.4	12.2	46.6	119.4	284.8	707.8	1,341.9
1950	56.5	*	*	*	0.4	2.3	12.6	48.3	123.6	302.9	725.6	1,356.6
MALE												
1959	34.2	*	*	*	*	1.3	6.3	25.7	74.2	180.4	425.2	901.4
1958	37.5	*	*	*	0.2	1.3	7.0	27.7	80.8	203.6	471.3	957.8
1957	38.3	*	*	*	0.2	1.4	7.9	29.7	86.4	206.9	459.5	959.1
1956	39.2	*	*	*	*	1.5	7.2	30.5	87.1	210.2	498.6	966.7
1955	41.1	*	*	*	0.2	1.5	8.5	31.6	93.1	218.9	526.2	987.4
1954	42.7	*	*	*	0.2	1.7	8.6	35.1	99.0	231.4	529.4	1,001.3
1953	47.9	*	*	*	0.3	1.9	10.6	41.3	116.2	256.8	590.1	1,055.0
1952	50.8	*	*	*	0.4	2.1	11.1	45.2	120.4	279.1	627.5	1,133.3
1951	52.3	*	*	*	0.3	2.2	11.5	45.5	125.6	283.1	670.4	1,247.3
1950	53.1	*	*	*	0.4	2.0	11.3	48.0	128.2	302.6	685.5	1,269.7
FEMALE												
1959	42.9	*	*	*	*	1.3	6.5	22.8	64.8	186.4	513.9	1,142.0
1958	47.6	*	*	*	0.2	1.3	6.8	26.9	77.2	207.5	577.3	1,244.8
1957	46.6	*	*	*	*	1.5	8.1	28.9	79.0	206.7	554.7	1,173.3
1956	47.2	*	*	*	*	1.4	7.7	28.7	79.7	210.8	583.7	1,217.4
1955	48.7	*	*	*	*	1.3	8.3	29.3	83.3	226.3	607.2	1,228.3
1954	50.3	*	*	*	0.3	2.0	9.1	34.2	88.2	242.2	618.0	1,197.9
1953	54.5	*	*	*	0.3	1.8	11.3	40.4	100.3	259.8	681.2	1,272.4
1952	56.9	*	*	*	0.3	2.1	12.3	43.7	108.4	275.6	709.0	1,321.0
1951	58.9	*	*	*	0.4	2.6	12.9	47.6	113.3	286.2	738.9	1,409.2
1950	59.9	*	*	*	0.4	2.5	13.8	48.7	119.0	303.1	759.7	1,417.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HYPERTENSIVE HEART DISEASE (440-443)												
WHITE, BOTH SEXES												
1959	34.4	*	*	*	*	0.5	2.5	13.9	50.3	159.1	457.1	1,029.9
1958	38.1	*	*	*	0.1	0.5	3.1	15.5	58.0	179.6	512.0	1,119.3
1957	37.6	*	*	*	*	0.6	3.4	17.4	59.9	181.1	493.2	1,073.3
1956	38.6	*	*	*	0.1	0.6	3.2	17.7	60.9	185.5	527.1	1,105.7
1955	40.4	*	*	*	*	0.6	3.9	18.3	67.1	197.5	552.8	1,120.2
1954	42.0	*	*	*	0.1	0.9	4.3	21.5	72.9	212.4	562.3	1,106.1
1953	46.3	*	*	*	0.2	0.9	5.5	26.6	85.4	233.8	621.4	1,161.5
1952	48.6	*	*	*	0.2	1.0	6.3	29.1	90.6	249.1	653.4	1,230.9
1951	50.5	*	*	*	0.2	1.2	6.5	31.4	96.3	258.0	689.2	1,326.4
1950	51.4	*	*	*	0.2	1.1	6.8	32.7	100.2	274.0	708.7	1,342.0
WHITE, MALE												
1959	30.0	*	*	*	*	0.6	2.9	16.0	56.4	156.6	404.0	878.5
1958	33.2	*	*	*	*	0.5	3.6	17.5	62.6	177.5	449.0	932.0
1957	33.5	*	*	*	*	0.6	4.0	19.7	65.4	180.8	435.8	922.8
1956	34.5	*	*	*	*	0.6	3.6	19.7	66.4	184.4	476.5	935.8
1955	36.4	*	*	*	*	0.7	4.4	20.1	72.7	193.7	502.3	964.3
1954	38.2	*	*	*	*	1.0	4.6	23.2	78.7	207.2	509.4	977.0
1953	42.8	*	*	*	*	0.9	6.0	28.4	93.9	230.5	567.3	1,014.2
1952	45.3	*	*	*	0.2	1.1	6.5	31.5	96.9	249.0	600.9	1,102.8
1951	47.1	*	*	*	0.2	1.2	6.9	32.7	102.9	255.0	645.5	1,215.6
1950	47.9	*	*	*	0.3	1.1	6.6	33.8	105.8	272.4	661.2	1,240.4
WHITE, FEMALE												
1959	38.7	*	*	*	*	0.4	2.2	11.8	44.7	161.2	498.3	1,127.8
1958	43.0	*	*	*	*	0.4	2.5	13.5	53.6	181.3	561.1	1,243.1
1957	41.7	*	*	*	*	0.5	2.9	15.1	54.7	181.3	538.0	1,175.7
1956	42.6	*	*	*	*	0.5	2.9	15.8	55.6	186.5	567.2	1,224.0
1955	44.4	*	*	*	*	0.4	3.4	16.5	61.8	200.8	593.4	1,230.6
1954	45.8	*	*	*	*	0.8	3.9	19.9	67.3	217.2	604.5	1,198.7
1953	49.8	*	*	*	*	0.9	4.9	24.9	77.1	236.8	665.3	1,267.9
1952	51.9	*	*	*	*	1.0	6.1	26.7	84.3	249.4	696.5	1,323.0
1951	53.8	*	*	*	0.3	1.2	6.2	30.2	89.8	260.8	725.5	1,405.4
1950	54.8	*	*	*	0.2	1.1	7.0	31.6	94.5	275.5	748.7	1,412.6

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HYPERTENSIVE HEART DISEASE (440-443)												
ALL OTHER, BOTH SEXES												
1959	72.2	*	*	*	0.7	7.8	40.1	122.5	268.6	477.2	721.0	1,254.4
1958	79.0	*	*	*	*	8.1	40.2	140.0	303.6	519.7	789.0	1,259.1
1957	81.9	*	*	*	0.8	8.7	47.7	143.0	332.1	514.9	786.4	1,226.6
1956	81.4	*	*	*	*	8.8	43.8	142.7	337.4	509.9	811.9	1,221.3
1955	82.5	*	*	*	*	8.5	47.1	147.4	331.0	525.3	834.0	1,229.3
1954	84.4	*	*	*	1.2	10.0	48.7	160.8	338.1	532.2	816.0	1,216.4
1953	92.8	*	*	*	1.2	9.9	58.5	176.5	385.4	552.0	917.1	1,426.9
1952	98.4	*	*	*	1.4	11.2	58.9	190.7	411.7	614.2	946.5	1,387.8
1951	99.0	*	*	*	1.1	12.7	61.6	190.8	414.2	604.3	983.9	1,529.8
1950	100.5	*	*	*	1.6	12.1	63.2	198.4	406.8	694.8	981.6	1,529.5
ALL OTHER, MALE												
1959	68.0	*	*	*	*	7.3	37.0	118.6	259.0	459.1	695.8	1,158.6
1958	73.2	*	*	*	*	8.2	37.0	126.1	273.5	511.1	756.1	1,296.3
1957	77.7	*	*	*	*	8.7	43.2	126.1	313.5	515.0	763.7	1,392.3
1956	78.2	*	*	*	*	9.3	38.5	134.3	318.1	514.0	784.6	1,340.0
1955	80.9	*	*	*	*	8.6	44.7	143.1	325.6	516.3	837.6	1,270.8
1954	81.6	*	*	*	*	8.4	43.7	149.9	336.0	518.8	795.8	1,313.6
1953	91.7	*	*	*	*	10.6	51.8	164.5	381.7	565.7	896.5	1,576.2
1952	98.1	*	*	*	*	10.7	52.2	175.5	407.5	636.7	989.1	1,520.0
1951	96.5	*	*	*	*	10.9	53.1	167.6	407.7	615.0	1,018.9	1,642.1
1950	98.7	*	*	*	*	9.8	53.8	183.5	395.6	695.4	1,024.8	1,612.4
ALL OTHER, FEMALE												
1959	76.1	*	*	*	*	8.3	42.9	126.2	277.9	493.6	743.5	1,325.6
1958	84.5	*	*	*	*	8.1	43.1	153.3	333.5	526.4	819.0	1,265.8
1957	85.8	*	*	*	*	8.8	51.8	159.1	350.3	514.7	807.6	1,143.2
1956	84.4	*	*	*	*	8.4	48.7	150.7	356.3	503.7	837.7	1,138.9
1955	83.9	*	*	*	*	8.5	49.3	151.6	336.3	533.6	824.2	1,200.0
1954	87.1	*	*	*	1.5	11.5	53.3	171.5	340.1	544.6	828.6	1,187.5
1953	93.8	*	*	*	*	9.3	64.8	188.2	388.4	537.7	928.9	1,325.8
1952	98.7	*	*	*	*	11.7	65.0	205.7	416.1	593.0	906.0	1,296.6
1951	101.4	*	*	*	1.6	14.3	69.5	213.6	420.9	594.2	950.5	1,453.6
1950	102.2	*	*	*	1.8	14.2	71.9	213.4	418.3	694.3	939.1	1,470.3

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER HYPERTENSIVE DISEASE (444-447)												
BOTH SEXES												
1959	7.4	*	*	*	0.2	1.2	3.1	5.9	11.1	27.2	89.4	241.3
1958	8.0	*	*	*	0.2	1.1	3.3	7.0	12.2	30.7	93.5	247.0
1957	6.5	*	*	*	0.1	0.9	2.8	5.3	9.3	25.6	79.4	216.4
1956	6.5	*	*	*	0.1	0.9	2.8	5.1	10.0	25.6	79.9	224.2
1955	6.8	*	*	*	0.2	1.0	3.1	5.3	10.3	26.4	85.8	226.9
1954	7.1	*	*	*	0.2	1.3	3.2	6.3	11.3	27.8	86.3	218.2
1953	7.8	*	*	*	0.2	1.3	4.0	7.7	13.0	30.7	93.6	240.4
1952	8.0	*	*	*	0.3	1.4	4.1	7.9	13.3	32.1	96.6	251.8
1951	8.6	*	*	*	0.3	1.4	4.2	8.1	14.7	34.6	110.2	274.8
1950	8.3	*	*	*	0.2	1.3	4.1	7.7	13.8	34.9	109.6	280.5
MALE												
1959	7.7	*	*	*	0.3	1.3	3.6	6.9	12.6	31.9	98.2	254.4
1958	8.1	*	*	*	0.2	1.2	3.5	8.0	14.0	33.7	104.1	265.7
1957	6.6	*	*	*	*	1.0	3.0	6.2	10.4	28.9	84.1	226.4
1956	6.7	*	*	*	*	0.9	3.1	6.3	11.3	28.3	84.6	243.1
1955	7.0	*	*	*	0.2	1.0	3.3	6.3	11.3	29.1	90.4	253.1
1954	7.2	*	*	*	0.2	1.2	3.4	7.2	12.6	30.3	91.4	233.2
1953	8.2	*	*	*	*	1.3	4.4	9.2	14.9	32.9	102.2	245.7
1952	8.2	*	*	*	0.3	1.2	4.2	9.0	15.0	34.0	101.6	258.6
1951	8.7	*	*	*	0.3	1.4	4.2	8.6	16.4	37.3	114.5	278.9
1950	8.5	*	*	*	0.2	1.3	4.2	8.8	15.4	38.4	112.8	298.1
FEMALE												
1959	7.2	*	*	*	*	1.2	2.7	4.9	9.7	23.2	82.5	232.8
1958	7.8	*	*	*	0.2	1.1	3.2	6.1	10.5	28.1	85.1	234.5
1957	6.4	*	*	*	*	0.9	2.6	4.3	8.4	22.7	75.6	209.5
1956	6.4	*	*	*	*	0.8	2.4	4.1	8.7	23.1	76.1	211.1
1955	6.6	*	*	*	*	0.9	2.8	4.3	9.3	23.9	82.1	208.2
1954	6.9	*	*	*	*	1.3	3.1	5.4	10.1	25.5	82.2	207.5
1953	7.5	*	*	*	0.2	1.3	3.5	6.1	11.2	28.7	86.4	236.6
1952	7.9	*	*	*	0.2	1.6	3.9	6.7	11.5	30.2	92.4	247.0
1951	8.6	*	*	*	0.3	1.5	4.1	7.6	13.0	32.1	106.5	271.9
1950	8.1	*	*	*	0.2	1.3	3.9	6.6	12.2	31.7	106.9	268.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER HYPERTENSIVE DISEASE (444-447)												
WHITE, BOTH SEXES												
1959	6.7	*	*	*	0.1	0.6	1.4	3.7	8.5	24.5	88.9	245.5
1958	7.2	*	*	*	0.2	0.5	1.6	4.5	9.9	28.0	92.1	249.7
1957	5.8	*	*	*	*	0.3	1.2	3.1	7.0	22.8	77.8	219.0
1956	5.8	*	*	*	*	0.4	1.2	3.2	7.6	22.9	77.7	226.6
1955	6.0	*	*	*	*	0.4	1.3	3.2	8.1	23.2	84.4	228.8
1954	6.2	*	*	*	0.1	0.6	1.5	3.9	9.3	24.9	84.5	216.0
1953	6.9	*	*	*	0.1	0.6	2.1	5.1	10.5	27.5	91.5	240.7
1952	7.2	*	*	*	0.2	0.6	2.4	5.6	10.7	29.0	94.3	252.1
1951	7.8	*	*	*	0.2	0.7	2.4	5.9	12.1	31.4	107.8	274.3
1950	7.6	*	*	*	0.2	0.7	2.4	5.5	11.3	32.0	108.1	285.3
WHITE, MALE												
1959	6.8	*	*	*	*	0.6	1.6	4.6	10.0	28.3	97.4	259.8
1958	7.3	*	*	*	*	0.5	1.8	5.6	11.5	30.8	102.4	265.2
1957	5.8	*	*	*	*	0.4	1.5	3.9	8.0	25.7	82.9	227.7
1956	6.0	*	*	*	*	0.4	1.5	4.2	8.7	25.3	82.6	248.7
1955	6.2	*	*	*	*	0.5	1.6	4.0	9.4	25.4	88.9	252.4
1954	6.4	*	*	*	*	0.6	1.8	4.9	10.5	27.5	89.6	229.8
1953	7.2	*	*	*	*	0.7	2.6	6.5	12.3	29.3	98.6	244.4
1952	7.3	*	*	*	*	0.6	2.6	6.8	12.4	31.2	98.5	254.5
1951	7.8	*	*	*	*	0.8	2.7	6.5	13.6	33.9	111.6	275.5
1950	7.8	*	*	*	*	0.8	2.7	6.7	13.3	35.2	110.6	305.7
WHITE, FEMALE												
1959	6.6	*	*	*	*	0.5	1.1	2.9	7.1	21.1	82.4	236.3
1958	7.1	*	*	*	*	0.4	1.4	3.5	8.5	25.5	84.2	239.5
1957	5.7	*	*	*	*	0.3	0.8	2.5	6.0	20.2	73.7	213.1
1956	5.7	*	*	*	*	0.3	0.9	2.3	6.6	20.7	73.9	211.3
1955	5.9	*	*	*	*	0.3	1.0	2.3	6.9	21.1	80.8	212.0
1954	6.1	*	*	*	*	0.5	1.3	2.9	8.2	22.5	80.3	206.1
1953	6.7	*	*	*	*	0.5	1.7	3.6	8.8	25.9	85.6	238.0
1952	7.0	*	*	*	*	0.7	2.1	4.4	9.1	27.1	90.9	250.3
1951	7.7	*	*	*	*	0.7	2.2	5.3	10.5	29.0	104.6	273.5
1950	7.4	*	*	*	*	0.7	2.1	4.3	9.3	29.2	106.0	271.1

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER HYPERTENSIVE DISEASE (444-447)												
ALL OTHER, BOTH SEXES												
1959	13.1	*	*	*	0.8	6.6	18.3	26.2	38.0	59.8	95.7	189.7
1958	14.0	*	*	*	*	6.6	18.7	30.7	36.9	63.4	112.0	213.6
1957	12.7	*	*	*	*	5.8	17.0	25.4	35.3	59.9	102.1	181.3
1956	12.5	*	*	*	*	5.2	16.4	23.5	36.6	58.0	110.1	195.1
1955	13.2	*	*	*	0.9	5.7	18.2	25.7	34.9	64.5	106.3	203.4
1954	13.9	*	*	*	*	7.3	17.9	29.2	35.1	62.5	113.9	241.8
1953	15.3	*	*	*	*	7.0	19.9	32.4	43.1	68.8	124.4	236.5
1952	15.3	*	*	*	1.0	7.9	19.1	29.8	45.3	68.2	130.1	249.0
1951	15.9	*	*	*	0.9	7.2	19.5	29.6	48.6	73.3	146.1	280.9
1950	14.7	*	*	*	*	6.3	18.7	28.9	44.5	73.8	132.3	223.3
ALL OTHER, MALE												
1959	14.4	*	*	*	*	7.5	20.8	28.6	39.0	73.2	109.0	193.1
1958	14.7	*	*	*	*	6.5	18.7	31.4	41.1	68.5	126.6	213.5
1957	13.2	*	*	*	*	6.1	16.5	28.6	36.4	66.8	100.0	211.5
1956	13.3	*	*	*	*	5.7	17.5	26.4	40.6	63.1	110.8	176.0
1955	13.9	*	*	*	*	5.5	18.8	29.1	32.9	72.1	109.6	262.5
1954	13.9	*	*	*	*	6.3	17.6	28.9	36.9	64.2	115.1	277.3
1953	16.5	*	*	*	*	6.6	20.9	34.8	45.4	75.0	150.9	261.9
1952	15.6	*	*	*	*	6.9	18.8	30.5	47.4	67.8	144.5	310.0
1951	16.0	*	*	*	*	6.4	18.1	29.1	50.7	77.7	155.7	321.1
1950	14.8	*	*	*	*	6.2	17.9	29.2	40.8	80.0	142.7	208.9
ALL OTHER, FEMALE												
1959	11.9	*	*	*	*	5.8	16.1	24.0	37.1	47.6	83.9	187.2
1958	13.3	*	*	*	*	6.6	18.6	30.0	32.9	58.6	98.7	171.1
1957	12.2	*	*	*	*	5.6	17.5	22.2	34.2	53.7	104.1	164.9
1956	11.7	*	*	*	*	4.8	15.4	20.8	32.6	53.1	109.4	208.3
1955	12.5	*	*	*	*	5.9	17.5	22.4	36.9	57.5	102.3	161.8
1954	13.9	*	*	*	*	8.1	18.2	29.4	33.3	60.9	111.9	225.0
1953	14.2	*	*	*	*	7.4	18.9	30.1	40.7	62.9	98.3	219.4
1952	15.0	*	*	*	*	8.8	19.4	29.1	43.0	68.6	116.4	206.9
1951	15.7	*	*	*	*	7.9	20.7	30.2	46.4	69.1	136.9	253.6
1950	14.6	*	*	*	*	6.4	19.5	28.6	48.2	67.6	122.0	233.6

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET GENERAL ARTERIOSCLEROSIS (450)												
BOTH SEXES												
1959	19.6	*	*	*	*	*	0.3	2.1	10.8	54.8	306.6	1,410.5
1958	19.9	*	*	*	*	*	0.3	1.7	10.4	58.7	313.9	1,465.7
1957	19.5	*	*	*	*	*	0.2	1.9	10.4	58.8	312.0	1,448.9
1956	19.1	*	*	*	*	*	0.2	1.8	9.8	56.7	319.1	1,436.4
1955	19.8	*	*	*	*	*	0.3	1.8	10.1	58.7	340.2	1,491.4
1954	18.8	*	*	*	*	*	0.3	1.9	10.3	57.6	324.0	1,444.0
1953	20.4	*	*	*	*	0.1	0.3	1.8	11.1	66.0	361.0	1,594.4
1952	20.4	*	*	*	*	*	0.3	2.1	11.8	66.4	370.8	1,598.2
1951	20.8	*	*	*	*	*	0.3	2.1	12.0	70.3	391.0	1,664.0
1950	20.4	*	*	*	*	*	0.3	2.0	12.3	70.0	392.0	1,727.7
MALE												
1959	18.9	*	*	*	*	*	0.4	2.8	13.9	66.3	338.5	1,414.4
1958	19.4	*	*	*	*	*	0.3	2.2	13.0	70.8	351.2	1,488.1
1957	19.1	*	*	*	*	*	0.2	2.4	13.1	72.1	341.9	1,462.3
1956	18.9	*	*	*	*	*	0.3	2.4	12.7	68.1	354.8	1,442.5
1955	19.9	*	*	*	*	*	0.3	2.6	12.8	71.7	383.8	1,506.6
1954	18.6	*	*	*	*	*	0.3	2.4	12.8	69.0	362.9	1,420.1
1953	20.6	*	*	*	*	*	0.4	2.3	14.2	80.4	405.2	1,589.3
1952	20.4	*	*	*	*	*	0.4	2.8	15.0	79.0	405.2	1,608.8
1951	21.1	*	*	*	*	*	0.2	2.7	14.9	84.7	429.6	1,727.7
1950	20.9	*	*	*	*	*	0.3	2.5	15.7	84.1	436.6	1,856.2
FEMALE												
1959	20.3	*	*	*	*	*	*	1.4	7.9	44.8	281.8	1,407.9
1958	20.4	*	*	*	*	*	0.2	1.3	7.9	47.9	284.6	1,450.8
1957	19.9	*	*	*	*	*	*	1.4	7.8	47.0	288.3	1,439.7
1956	19.4	*	*	*	*	*	0.2	1.3	7.0	46.5	290.5	1,432.1
1955	19.7	*	*	*	*	*	0.2	1.0	7.4	47.0	305.0	1,480.6
1954	18.9	*	*	*	*	*	0.3	1.3	7.8	47.2	292.2	1,461.2
1953	20.3	*	*	*	*	*	0.2	1.4	8.1	52.8	324.5	1,598.0
1952	20.3	*	*	*	*	*	0.2	1.4	8.7	54.9	342.3	1,590.6
1951	20.5	*	*	*	*	*	0.3	1.4	9.1	57.0	358.3	1,618.6
1950	19.8	*	*	*	*	*	0.2	1.4	8.9	56.9	354.0	1,638.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET GENERAL ARTERIOSCLEROSIS (450)												
WHITE, BOTH SEXES												
1959	20.5	*	*	*	*	*	0.2	1.7	9.4	52.0	310.2	1,465.2
1958	20.7	*	*	*	*	*	0.2	1.4	8.9	55.4	316.5	1,518.5
1957	20.3	*	*	*	*	*	0.1	1.6	8.9	55.9	316.3	1,499.3
1956	20.0	*	*	*	*	*	0.2	1.5	8.2	53.7	324.8	1,494.2
1955	20.6	*	*	*	*	*	0.2	1.4	8.4	56.3	345.2	1,547.7
1954	19.5	*	*	*	*	*	0.2	1.4	8.7	55.3	328.9	1,494.8
1953	21.3	*	*	*	*	0.1	0.2	1.3	9.2	63.5	366.3	1,658.2
1952	21.2	*	*	*	*	*	0.2	1.6	10.2	64.3	378.0	1,657.5
1951	21.7	*	*	*	*	*	0.2	1.5	10.2	68.4	399.6	1,735.5
1950	21.3	*	*	*	*	*	0.2	1.5	10.6	67.2	400.2	1,800.9
WHITE, MALE												
1959	19.6	*	*	*	*	*	0.3	2.4	12.7	62.9	342.6	1,474.8
1958	20.1	*	*	*	*	*	0.3	1.8	11.8	67.7	354.9	1,539.6
1957	19.8	*	*	*	*	*	0.2	2.2	11.7	69.2	347.7	1,509.3
1956	19.6	*	*	*	*	*	*	2.2	10.9	65.3	362.9	1,494.7
1955	20.5	*	*	*	*	*	0.2	2.1	11.2	68.4	392.0	1,559.9
1954	19.3	*	*	*	*	*	0.2	2.0	11.3	66.7	368.8	1,461.7
1953	21.3	*	*	*	*	*	0.3	1.7	12.1	77.5	412.7	1,641.4
1952	21.1	*	*	*	*	*	0.3	2.4	13.1	76.7	412.6	1,662.8
1951	21.9	*	*	*	*	*	*	2.0	13.1	82.9	438.7	1,797.9
1950	21.7	*	*	*	*	*	0.3	2.0	13.9	80.9	445.3	1,937.6
WHITE, FEMALE												
1959	21.3	*	*	*	*	*	*	1.0	6.2	42.5	285.3	1,458.9
1958	21.3	*	*	*	*	*	*	1.0	6.1	44.6	286.5	1,504.6
1957	20.9	*	*	*	*	*	*	1.0	6.2	44.1	291.5	1,492.6
1956	20.4	*	*	*	*	*	*	0.9	5.6	43.4	294.7	1,493.8
1955	20.7	*	*	*	*	*	*	0.8	5.8	45.4	308.0	1,539.0
1954	19.8	*	*	*	*	*	*	0.9	6.2	44.9	296.7	1,518.6
1953	21.2	*	*	*	*	*	*	0.9	6.4	50.7	328.6	1,670.4
1952	21.4	*	*	*	*	*	*	0.9	7.2	53.0	349.6	1,653.7
1951	21.6	*	*	*	*	*	*	1.0	7.4	55.2	367.1	1,691.0
1950	20.9	*	*	*	*	*	*	0.9	7.4	54.4	362.2	1,706.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
GENERAL ARTERIOSCLEROSIS (450)												
ALL OTHER, BOTH SEXES												
1959	12.8	*	*	*	*	*	0.9	5.9	25.6	88.6	256.7	742.6
1958	13.7	*	*	*	*	*	0.9	4.9	26.4	97.3	278.8	830.3
1957	13.0	*	*	*	*	*	*	5.0	26.7	94.0	252.9	820.3
1956	12.4	*	*	*	*	*	0.8	5.0	27.8	91.9	237.7	739.3
1955	12.9	*	*	*	*	*	1.2	5.6	29.1	87.0	269.5	803.4
1954	12.4	*	*	*	*	*	1.2	6.1	29.1	84.9	253.3	794.5
1953	13.6	*	*	*	*	*	*	6.8	34.6	95.4	282.9	809.6
1952	13.0	*	*	*	*	*	1.4	6.5	32.9	91.3	265.0	865.3
1951	12.8	*	*	*	*	*	1.0	7.4	34.6	92.3	263.1	797.9
1950	12.8	*	*	*	*	*	1.0	6.8	32.0	107.9	267.5	857.4
ALL OTHER, MALE												
1959	13.6	*	*	*	*	*	*	6.9	26.2	105.8	286.8	734.5
1958	14.2	*	*	*	*	*	*	5.7	25.8	107.2	303.6	940.7
1957	13.5	*	*	*	*	*	*	4.4	27.9	107.1	267.4	900.0
1956	13.0	*	*	*	*	*	*	5.1	32.0	100.3	249.2	812.0
1955	14.2	*	*	*	*	*	*	7.8	32.1	110.2	276.8	854.2
1954	13.4	*	*	*	*	*	*	6.7	30.6	96.3	284.9	886.4
1953	15.1	*	*	*	*	*	*	7.9	39.4	114.0	304.4	923.8
1952	14.4	*	*	*	*	*	*	6.5	37.9	106.3	304.5	925.0
1951	14.2	*	*	*	*	*	*	9.4	37.3	106.7	308.5	852.6
1950	14.4	*	*	*	*	*	*	7.1	37.3	124.5	315.2	905.3
ALL OTHER, FEMALE												
1959	12.0	*	*	*	*	*	*	4.8	24.9	73.0	229.8	748.7
1958	13.2	*	*	*	*	*	*	4.1	27.0	88.0	256.2	773.7
1957	12.5	*	*	*	*	*	*	5.6	25.5	81.9	239.3	786.5
1956	11.8	*	*	*	*	*	*	5.0	23.6	83.7	226.8	688.9
1955	11.7	*	*	*	*	*	*	3.6	26.1	65.6	260.6	767.6
1954	11.6	*	*	*	*	*	*	5.5	27.6	74.3	221.4	756.3
1953	12.1	*	*	*	*	*	*	5.7	29.6	77.8	260.3	732.3
1952	11.7	*	*	*	*	*	*	6.4	27.9	77.3	227.6	824.1
1951	11.4	*	*	*	*	*	*	5.4	31.8	78.6	219.8	760.7
1950	11.3	*	*	*	*	*	*	6.6	26.6	91.1	220.6	823.2

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)												
BOTH SEXES												
1959	10.3	1.1	0.2	0.2	0.9	1.5	3.4	8.8	24.7	52.8	92.9	137.3
1958	9.9	0.9	0.2	0.1	0.7	1.7	3.3	8.9	22.8	52.0	89.3	136.5
1957	9.0	1.4	0.1	0.1	0.8	1.5	3.2	8.0	23.2	45.8	74.0	117.4
1956	8.4	1.4	0.2	0.2	0.8	1.6	3.2	8.2	20.6	43.1	70.1	98.4
1955	7.3	1.4	*	0.2	0.9	1.5	3.0	7.3	18.1	35.8	60.8	95.4
1954	6.4	1.6	0.2	0.2	0.8	1.6	2.9	7.2	16.2	30.2	47.7	78.9
1953	6.0	1.8	0.2	0.2	0.9	1.5	2.7	6.9	15.6	27.6	46.6	76.1
1952	5.7	1.9	0.2	0.1	0.8	1.5	2.9	6.4	15.4	25.0	44.9	67.7
1951	5.4	2.1	0.2	0.1	0.9	1.4	2.9	6.1	13.9	24.5	42.5	67.7
1950	4.9	1.5	0.3	0.2	0.9	1.4	2.7	5.9	12.8	22.5	36.5	63.8
MALE												
1959	12.8	1.2	0.3	0.1	0.5	1.1	3.6	11.2	34.9	77.0	117.2	151.5
1958	12.5	1.4	*	*	0.5	1.3	3.5	11.2	33.1	74.6	115.8	155.8
1957	11.4	1.6	*	0.2	0.6	1.2	3.6	10.4	33.7	65.5	95.0	135.0
1956	10.7	1.8	0.3	0.1	0.4	1.2	3.3	10.0	30.7	62.5	90.6	116.2
1955	9.2	1.7	*	0.1	0.7	1.1	3.0	9.7	25.8	50.7	76.8	117.3
1954	8.0	1.9	*	*	0.6	1.4	3.0	9.3	22.7	42.7	62.0	87.2
1953	7.5	2.2	*	*	0.6	1.2	2.8	9.0	22.4	37.9	58.1	84.4
1952	7.0	2.2	*	*	0.5	1.2	3.0	8.3	22.1	33.5	57.2	80.6
1951	6.5	1.9	*	0.2	0.5	1.1	3.1	7.9	19.1	33.2	51.4	73.4
1950	5.8	1.6	*	*	0.5	1.0	3.0	7.3	17.1	29.5	45.7	70.5
FEMALE												
1959	7.9	1.0	*	0.2	1.3	1.8	3.1	6.4	15.1	31.6	73.9	128.0
1958	7.5	*	*	*	1.0	2.0	3.1	6.5	13.1	32.1	68.5	123.6
1957	6.6	1.2	*	0.1	1.1	1.9	2.9	5.7	13.3	28.3	57.4	105.5
1956	6.2	1.1	*	0.2	1.1	2.0	3.0	6.4	11.0	25.7	53.7	86.0
1955	5.5	*	*	0.2	1.1	1.8	2.9	5.0	10.7	22.3	47.9	80.0
1954	4.8	1.2	*	0.3	1.0	1.8	2.7	5.2	9.8	18.8	36.1	72.9
1953	4.6	1.4	*	0.3	1.2	1.8	2.7	4.8	9.0	18.2	37.2	70.1
1952	4.4	1.6	*	0.2	1.2	1.8	2.8	4.5	8.8	17.3	34.7	58.5
1951	4.3	2.2	*	*	1.3	1.6	2.7	4.2	8.7	16.4	35.0	63.6
1950	4.0	1.4	*	0.2	1.2	1.7	2.5	4.5	8.5	15.9	28.6	59.1

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)												
WHITE, BOTH SEXES												
1959	10.4	1.1	0.2	0.1	0.7	1.2	2.7	8.0	23.6	52.8	93.8	141.3
1958	9.9	0.9	*	0.1	0.7	1.3	2.8	7.6	21.9	51.8	90.8	140.7
1957	9.0	1.1	*	0.1	0.7	1.2	2.6	7.0	22.2	46.0	75.7	121.7
1956	8.4	1.1	0.2	0.2	0.6	1.3	2.6	7.3	20.0	43.6	71.1	101.6
1955	7.3	1.2	*	0.2	0.8	1.2	2.4	6.6	17.4	35.9	61.7	97.6
1954	6.3	1.4	*	0.2	0.7	1.5	2.4	6.5	15.6	30.5	48.6	80.9
1953	6.0	1.4	0.2	0.2	0.8	1.3	2.4	6.2	15.1	27.5	47.6	78.2
1952	5.7	1.7	0.2	0.1	0.7	1.2	2.5	5.9	14.9	25.5	45.9	69.8
1951	5.4	1.9	0.2	0.1	0.8	1.2	2.5	5.5	13.4	24.7	43.0	69.4
1950	4.9	1.5	0.2	0.2	0.7	1.2	2.3	5.4	12.4	22.6	37.0	65.2
WHITE, MALE												
1959	13.0	1.1	*	*	0.5	1.0	3.0	10.4	34.2	77.6	119.1	155.5
1958	12.7	1.3	*	*	0.5	1.1	3.1	10.3	32.6	75.3	118.9	161.1
1957	11.5	1.3	*	*	0.5	0.9	3.0	9.4	32.7	66.3	98.3	138.6
1956	10.9	1.4	*	0.1	0.3	1.0	2.9	9.4	30.3	63.5	92.4	120.9
1955	9.3	1.5	*	0.1	0.6	1.0	2.6	9.1	25.2	51.5	78.5	119.4
1954	8.1	1.7	*	*	0.5	1.3	2.5	8.4	22.3	43.6	64.1	89.7
1953	7.5	1.7	*	*	0.5	1.1	2.6	8.3	22.1	38.0	59.5	85.4
1952	7.2	1.8	*	*	0.5	1.0	2.7	7.9	21.6	34.6	58.6	82.6
1951	6.6	1.6	*	*	0.5	0.9	2.7	7.3	18.8	33.7	52.1	74.3
1950	5.9	1.6	*	*	0.4	0.9	2.6	6.8	16.7	29.9	46.5	72.9
WHITE, FEMALE												
1959	7.8	*	*	0.2	1.0	1.4	2.5	5.6	13.7	31.2	74.3	132.1
1958	7.2	*	*	*	0.8	1.5	2.5	5.0	11.8	31.1	68.9	127.2
1957	6.5	*	*	*	0.9	1.5	2.3	4.7	12.2	27.9	58.0	110.3
1956	6.0	*	*	0.2	0.9	1.6	2.3	5.3	10.2	25.8	54.3	88.2
1955	5.3	*	*	0.2	1.0	1.4	2.3	4.2	9.9	21.9	48.3	82.2
1954	4.7	*	*	0.2	0.9	1.6	2.3	4.5	9.1	18.6	36.1	74.6
1953	4.5	*	*	0.3	1.1	1.5	2.2	4.0	8.4	18.0	37.9	73.0
1952	4.2	1.6	*	0.2	0.9	1.4	2.3	3.9	8.2	17.1	35.4	60.5
1951	4.2	2.1	*	*	1.2	1.4	2.2	3.7	8.0	16.5	35.5	66.0
1950	3.9	*	*	0.3	1.0	1.5	2.1	4.0	8.2	15.8	29.1	59.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)												
ALL OTHER, BOTH SEXES												
1959	9.8	*	*	*	1.8	3.7	8.8	16.3	36.5	52.2	80.0	88.2
1958	9.9	*	*	*	1.4	4.5	8.0	20.7	32.3	55.0	68.5	86.4
1957	9.0	3.5	*	*	1.9	4.1	8.2	17.1	34.9	43.6	50.4	64.1
1956	8.1	*	*	*	1.9	4.3	8.0	16.1	27.3	36.4	55.2	59.0
1955	7.4	*	*	*	1.5	3.5	7.5	14.1	26.5	33.6	47.7	69.0
1954	6.6	*	*	*	1.5	3.0	6.9	14.4	23.0	26.7	34.4	52.7
1953	6.3	4.3	*	*	1.8	3.3	6.0	13.7	21.4	28.1	32.1	50.0
1952	5.8	*	*	*	1.8	3.7	6.2	11.6	22.0	19.9	31.0	42.9
1951	5.7	*	*	*	1.7	2.9	6.6	11.7	20.5	21.1	34.6	46.8
1950	5.2	*	*	*	1.7	2.9	6.0	10.4	17.5	21.1	27.5	46.9
ALL OTHER, MALE												
1959	11.0	*	*	*	*	2.3	9.1	18.6	42.5	69.9	93.1	106.9
1958	10.4	*	*	*	*	3.3	7.7	20.3	38.3	66.4	76.3	100.0
1957	10.5	*	*	*	*	3.6	8.8	19.5	44.6	56.2	52.6	92.3
1956	8.9	*	*	*	*	2.8	7.4	15.9	34.6	50.1	66.9	*
1955	8.0	*	*	*	*	2.0	6.9	15.4	32.9	41.2	54.4	91.7
1954	7.3	*	*	*	*	2.4	7.7	17.2	28.0	32.1	33.6	*
1953	7.1	*	*	*	*	2.2	5.3	15.7	26.5	36.0	38.6	*
1952	6.0	*	*	*	*	2.5	5.4	12.5	28.1	21.2	38.2	*
1951	6.3	*	*	*	*	2.2	6.6	13.8	22.9	27.2	41.5	*
1950	5.7	*	*	*	*	1.9	6.1	11.7	22.4	25.0	33.7	*
ALL OTHER, FEMALE												
1959	8.7	*	*	*	3.0	4.8	8.6	14.1	30.8	36.2	68.3	74.4
1958	9.3	*	*	*	2.0	5.5	8.3	21.1	26.7	44.5	61.4	78.9
1957	7.6	*	*	*	2.4	4.6	7.6	14.7	25.3	32.0	48.3	*
1956	7.4	*	*	*	2.7	5.7	8.6	16.3	20.1	23.7	44.2	58.3
1955	6.7	*	*	*	2.0	4.8	8.0	12.8	20.1	26.7	40.9	*
1954	5.9	*	*	*	2.1	3.6	6.1	11.8	18.1	21.8	34.9	*
1953	5.6	*	*	*	2.2	4.3	6.5	11.8	16.1	20.6	25.6	*
1952	5.6	*	*	*	2.8	4.8	7.0	10.7	15.7	18.8	24.1	*
1951	5.2	*	*	*	2.7	3.5	6.5	9.5	18.1	15.3	27.9	*
1950	4.7	*	*	*	2.2	3.7	5.9	9.2	12.5	17.2	21.5	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)												
BOTH SEXES												
1959	7.0	0.8	0.2	0.5	1.5	2.9	4.3	7.5	13.0	24.4	58.4	140.1
1958	8.0	0.7	0.2	0.5	1.9	3.0	4.7	8.4	14.4	29.5	68.5	168.4
1957	8.8	*	0.3	0.7	2.1	3.5	5.4	8.9	16.6	32.6	73.5	181.0
1956	9.1	0.8	0.4	0.7	1.9	3.3	5.2	9.2	16.8	34.2	82.6	196.7
1955	9.6	0.6	0.4	0.7	2.2	3.2	5.3	9.6	17.7	36.8	90.5	200.9
1954	10.6	0.9	0.6	0.8	2.2	3.7	5.4	10.8	20.1	40.7	101.6	214.7
1953	11.9	0.6	0.7	0.9	2.5	3.9	6.3	12.3	22.0	46.4	113.1	260.5
1952	13.3	0.8	0.7	0.9	2.3	3.9	7.0	14.2	25.6	52.2	134.0	286.5
1951	14.7	0.8	0.8	1.0	2.5	4.1	7.3	14.7	28.5	59.0	151.0	356.3
1950	16.4	0.8	1.0	1.0	2.6	4.5	8.2	16.3	30.9	69.8	174.3	411.0
MALE												
1959	7.7	1.0	*	0.4	1.7	3.5	5.1	9.1	14.9	28.2	69.2	166.8
1958	8.8	*	0.3	0.5	2.1	3.3	5.7	9.5	16.7	34.3	80.6	199.4
1957	9.5	*	0.3	0.7	2.6	4.3	6.1	10.2	18.4	37.9	82.1	207.4
1956	9.9	*	0.4	0.7	2.2	4.0	5.8	10.3	18.8	38.2	94.6	236.1
1955	10.4	*	0.5	0.7	2.7	3.7	6.2	10.7	19.5	41.8	105.5	227.7
1954	11.5	*	0.6	0.8	2.6	4.1	6.1	11.8	22.2	47.1	119.3	245.7
1953	12.8	*	0.7	0.9	2.7	4.4	6.8	13.6	23.9	52.5	132.9	303.5
1952	14.6	*	0.9	1.0	2.7	4.5	7.4	16.2	28.5	59.6	157.9	329.7
1951	15.8	*	1.0	1.1	2.7	4.6	7.8	16.4	31.5	66.6	172.2	390.6
1950	17.5	*	1.1	1.0	3.0	4.8	8.7	17.7	34.2	77.7	199.4	467.4
FEMALE												
1959	6.3	*	*	0.6	1.3	2.2	3.6	5.9	11.1	21.0	50.0	122.7
1958	7.2	*	*	0.4	1.6	2.7	3.7	7.3	12.2	25.2	58.9	147.7
1957	8.1	*	0.3	0.8	1.7	2.6	4.8	7.6	14.8	27.8	66.8	163.0
1956	8.4	*	0.3	0.7	1.6	2.6	4.6	8.0	14.8	30.6	73.0	169.4
1955	8.8	*	0.3	0.7	1.7	2.6	4.5	8.5	16.1	32.3	78.4	182.0
1954	9.7	*	0.6	0.7	1.8	3.2	4.8	9.9	18.1	34.8	87.1	192.5
1953	10.9	*	0.7	0.9	2.2	3.5	5.8	10.9	20.1	40.8	96.7	229.6
1952	12.1	*	0.5	0.9	1.9	3.3	6.6	12.3	22.8	45.4	114.2	255.6
1951	13.6	*	0.7	1.0	2.3	3.6	6.9	13.0	25.5	52.1	133.2	331.9
1950	15.3	*	0.9	0.9	2.2	4.2	7.8	15.0	27.6	62.4	152.9	371.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)												
WHITE, BOTH SEXES												
1959	6.1	0.6	0.2	0.5	1.4	2.5	3.3	5.7	10.5	20.3	54.2	133.1
1958	7.1	0.7	0.2	0.5	1.8	2.6	3.6	6.4	11.1	25.5	63.7	160.3
1957	7.8	*	0.3	0.7	2.0	2.9	4.1	6.9	13.5	28.3	68.1	171.9
1956	8.1	0.7	0.3	0.6	1.8	2.8	4.0	6.8	13.6	29.5	77.4	189.1
1955	8.7	*	0.4	0.7	2.1	2.7	4.1	7.4	14.4	32.6	84.8	193.2
1954	9.5	0.7	0.6	0.7	2.0	3.1	4.3	8.4	16.6	35.5	95.3	205.2
1953	10.6	*	0.6	0.9	2.3	3.3	4.9	9.5	18.1	40.6	104.9	251.0
1952	11.7	0.9	0.7	0.9	2.1	3.3	5.3	10.8	20.5	45.5	123.6	267.4
1951	13.1	*	0.8	1.0	2.4	3.4	5.5	11.3	23.5	51.3	141.2	337.3
1950	14.6	*	1.0	1.0	2.5	3.8	6.3	12.7	25.4	60.3	162.5	397.3
WHITE, MALE												
1959	6.9	*	*	0.4	1.7	3.2	4.1	7.3	12.6	23.4	64.4	155.5
1958	7.9	*	*	0.5	2.1	3.1	4.8	7.6	13.4	29.7	75.5	185.8
1957	8.6	*	*	0.6	2.5	3.8	5.0	8.6	15.3	33.6	75.8	191.6
1956	8.9	*	0.3	0.6	2.1	3.7	4.9	8.2	15.5	33.0	88.3	222.8
1955	9.6	*	0.5	0.7	2.7	3.4	5.1	8.8	16.4	37.2	98.8	221.1
1954	10.3	*	0.5	0.8	2.4	3.7	5.0	9.6	18.8	40.5	109.4	230.1
1953	11.6	*	0.6	0.9	2.8	3.9	5.7	11.2	20.1	45.7	122.7	291.8
1952	13.0	*	0.9	1.0	2.6	4.1	6.0	12.9	23.4	52.0	143.7	308.3
1951	14.2	*	0.9	1.1	2.7	4.1	6.4	13.4	26.8	57.9	159.3	362.4
1950	15.8	*	1.0	1.0	2.9	4.4	7.4	14.7	28.6	66.8	184.7	444.6
WHITE, FEMALE												
1959	5.4	*	*	0.6	1.2	1.8	2.5	4.1	8.6	17.6	46.3	118.7
1958	6.3	*	*	0.4	1.5	2.0	2.5	5.2	9.0	21.9	54.6	143.5
1957	7.1	*	*	0.7	1.6	2.1	3.3	5.2	11.9	23.7	62.1	158.4
1956	7.4	*	0.3	0.7	1.4	1.9	3.1	5.5	11.9	26.4	68.8	165.7
1955	7.8	*	*	0.7	1.6	2.0	3.1	6.1	12.5	28.6	73.7	173.5
1954	8.7	*	0.6	0.7	1.7	2.6	3.5	7.3	14.5	30.9	83.8	187.3
1953	9.6	*	0.6	0.9	1.9	2.8	4.0	7.8	16.2	36.0	90.4	221.6
1952	10.5	*	0.5	0.9	1.7	2.6	4.6	8.6	17.7	39.6	107.1	238.1
1951	12.0	*	0.7	1.0	2.1	2.7	4.7	9.1	20.4	45.3	126.2	319.3
1950	13.5	*	0.9	0.9	2.0	3.2	5.3	10.7	22.1	54.3	143.8	364.4

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)												
ALL OTHER, BOTH SEXES												
1959	13.6	*	*	0.7	2.0	6.0	13.0	24.2	38.4	72.9	117.4	225.0
1958	15.4	*	*	0.6	2.2	6.3	13.8	27.0	49.3	76.7	134.2	265.2
1957	16.9	*	*	0.9	3.0	7.8	17.0	28.2	49.8	83.0	150.0	287.5
1956	17.6	*	*	0.9	2.9	7.5	15.8	31.7	52.4	89.8	156.7	288.5
1955	17.6	*	*	0.7	2.8	6.8	16.2	30.3	55.9	86.2	171.9	294.8
1954	19.7	*	*	0.9	3.1	8.2	15.6	34.0	61.1	102.9	193.4	327.3
1953	22.3	*	*	0.8	3.6	8.9	18.6	38.5	68.9	115.7	232.9	376.9
1952	26.7	*	*	1.0	3.3	8.9	22.0	47.4	89.2	132.2	287.2	522.4
1951	28.1	*	*	1.0	3.6	9.7	22.8	47.3	91.4	151.3	296.3	587.2
1950	31.2	*	*	0.8	3.1	10.4	25.1	51.1	97.4	199.0	353.1	573.9
ALL OTHER, MALE												
1959	14.6	*	*	*	2.0	6.7	13.5	26.0	39.3	84.3	130.6	293.1
1958	16.3	*	*	*	2.6	5.1	14.1	27.6	51.7	88.9	146.0	366.7
1957	17.4	*	*	*	3.5	8.8	16.3	25.7	52.4	88.4	163.0	396.2
1956	18.4	*	*	1.1	3.0	6.9	13.9	30.9	56.4	99.5	176.9	396.0
1955	17.9	*	*	*	2.8	6.4	16.1	28.6	54.1	96.4	192.0	308.3
1954	21.5	*	*	1.1	3.5	7.9	15.3	32.6	60.8	125.6	251.3	445.5
1953	23.0	*	*	*	2.6	8.6	16.2	36.7	69.6	132.6	271.1	452.4
1952	28.3	*	*	*	3.7	8.3	19.9	47.0	90.8	149.6	350.9	600.0
1951	29.1	*	*	*	3.0	8.5	20.0	45.0	90.3	169.4	351.9	742.1
1950	32.1	*	*	*	3.3	8.7	20.8	45.9	100.3	219.9	405.3	733.9
ALL OTHER, FEMALE												
1959	12.7	*	*	*	2.0	5.4	12.5	22.6	37.5	62.5	105.6	174.4
1958	14.4	*	*	*	1.8	7.4	13.4	26.5	47.3	65.4	123.5	200.0
1957	16.4	*	*	1.0	2.6	6.9	17.7	30.5	47.3	78.1	137.9	218.9
1956	16.8	*	*	*	2.7	8.1	17.6	32.5	48.4	80.5	137.7	213.9
1955	17.3	*	*	*	2.8	7.2	16.2	31.9	57.7	76.8	151.5	285.3
1954	17.9	*	*	*	2.7	8.4	15.9	35.3	61.5	81.9	137.3	256.3
1953	21.6	*	*	*	4.4	9.1	20.9	40.4	68.1	99.7	195.0	325.8
1952	25.2	*	*	*	3.0	9.4	23.9	47.7	87.6	116.0	226.7	469.0
1951	27.2	*	*	*	4.2	10.8	25.4	49.6	92.6	134.1	243.2	482.1
1950	30.3	*	*	*	3.0	12.0	29.1	56.3	94.4	178.0	301.6	459.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)												
BOTH SEXES												
1959	31.2	239.2	15.3	2.5	2.7	3.8	7.7	15.5	33.5	86.5	281.6	924.3
1958	33.1	245.3	16.0	2.3	2.9	4.3	7.8	17.3	38.9	96.0	292.8	988.1
1957	35.8	239.9	18.3	3.8	5.5	5.6	10.2	20.6	44.3	106.9	298.6	979.8
1956	28.2	225.3	15.3	2.3	2.4	3.0	7.0	14.0	30.4	80.6	258.0	853.5
1955	27.1	221.6	14.9	2.5	2.7	3.1	6.8	14.2	29.5	76.4	249.8	791.1
1954	25.4	227.2	16.1	2.4	2.4	3.3	6.6	14.2	27.6	70.8	220.6	713.2
1953	33.0	257.5	18.8	3.1	3.2	4.2	8.8	18.6	38.2	96.6	310.7	1,000.7
1952	29.7	266.9	19.9	3.2	3.0	4.2	8.6	17.3	33.7	82.0	266.6	825.2
1951	31.4	252.0	19.0	3.2	3.4	4.0	8.7	18.9	38.3	93.8	299.5	935.1
1950	31.3	268.1	18.9	3.2	3.2	4.2	9.0	19.3	38.2	97.1	296.5	954.4
MALE												
1959	35.9	265.9	16.4	2.4	3.0	4.3	9.3	20.8	46.8	118.3	345.7	1,035.8
1958	38.6	271.4	16.2	2.3	3.3	4.8	10.0	23.6	54.7	131.7	361.0	1,107.0
1957	41.5	265.9	19.0	4.0	5.2	5.4	12.4	27.9	61.8	144.9	360.4	1,097.3
1956	32.3	248.0	15.9	2.3	2.7	3.4	8.4	18.9	42.6	106.8	311.8	930.0
1955	30.9	242.4	15.7	2.6	3.0	3.6	8.6	19.3	40.7	98.9	296.4	835.8
1954	29.4	248.1	16.8	2.6	2.8	3.9	8.3	19.6	38.4	92.3	258.9	774.0
1953	37.9	279.3	20.1	3.2	3.7	4.6	11.5	25.5	52.9	122.6	362.8	1,084.4
1952	33.9	287.5	20.4	3.3	3.3	4.9	11.2	24.2	46.3	102.8	306.4	860.8
1951	35.4	274.9	19.5	3.5	3.7	4.4	10.7	25.3	51.0	115.7	340.9	980.9
1950	35.2	293.0	20.1	3.3	3.5	4.2	10.9	25.3	50.6	120.2	331.2	1,028.6
FEMALE												
1959	26.6	211.6	14.1	2.5	2.4	3.4	6.1	10.3	21.0	58.8	231.6	851.4
1958	27.9	218.3	15.8	2.3	2.5	3.8	5.7	11.2	23.9	64.4	239.0	908.9
1957	30.3	212.9	17.7	3.7	5.8	5.7	8.1	13.4	27.7	73.1	249.4	899.6
1956	24.1	201.8	14.8	2.4	2.0	2.7	5.6	9.2	18.7	57.1	215.0	800.2
1955	23.4	200.1	14.1	2.5	2.4	2.7	5.0	9.2	18.8	56.1	212.1	759.5
1954	21.6	205.6	15.3	2.3	2.1	2.7	5.0	8.9	17.2	51.4	189.2	669.6
1953	28.2	235.0	17.5	3.0	2.8	3.8	6.2	11.8	23.9	72.8	267.7	940.5
1952	25.6	245.5	19.3	3.1	2.7	3.6	6.2	10.6	21.3	62.8	233.6	799.7
1951	27.5	228.3	18.4	3.0	3.2	3.7	6.8	12.5	25.6	73.6	264.7	902.5
1950	27.4	242.2	17.8	3.0	3.0	4.1	7.1	13.3	25.7	75.6	266.9	902.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)												
WHITE, BOTH SEXES												
1959	28.5	171.5	11.3	2.2	2.3	2.8	5.5	12.8	28.9	81.7	282.8	952.8
1958	30.1	177.7	12.3	2.0	2.6	3.1	5.8	14.1	33.2	89.7	290.6	1,008.2
1957	32.5	174.0	14.0	3.4	4.9	4.1	7.7	16.5	38.0	100.3	297.6	997.8
1956	25.6	166.4	11.4	2.0	2.1	2.1	5.1	11.2	26.3	75.8	258.7	874.0
1955	24.5	163.0	11.5	2.2	2.3	2.2	5.1	11.0	25.3	71.3	249.0	810.6
1954	22.8	168.5	12.3	2.1	2.0	2.3	5.0	11.2	23.6	65.8	220.4	724.6
1953	29.4	193.9	15.2	2.7	2.5	2.8	6.4	13.9	31.2	88.0	305.6	1,007.8
1952	26.3	197.2	15.7	2.8	2.4	2.9	6.2	13.6	28.5	75.6	264.8	839.2
1951	28.2	189.3	15.2	2.8	2.6	2.8	6.6	14.0	32.4	87.7	298.1	950.1
1950	28.0	211.3	14.6	2.7	2.7	2.9	6.7	14.6	31.9	89.3	293.8	966.0
WHITE, MALE												
1959	32.7	194.7	12.0	2.2	2.6	3.1	6.7	17.2	40.9	112.5	347.6	1,067.8
1958	34.9	198.2	12.5	2.1	2.9	3.4	7.4	19.2	47.5	123.7	358.6	1,125.3
1957	37.8	200.1	14.4	3.6	4.7	3.9	9.3	23.0	54.3	138.0	359.1	1,117.4
1956	29.4	186.3	11.8	1.9	2.5	2.5	6.2	15.6	37.6	101.1	313.6	949.7
1955	27.7	179.6	11.9	2.3	2.5	2.5	6.4	15.3	35.6	92.4	295.8	846.6
1954	26.2	185.9	12.9	2.2	2.3	2.8	6.4	15.6	33.4	86.7	258.7	782.3
1953	33.8	213.8	15.7	2.6	2.9	3.1	8.4	19.9	44.5	113.1	358.8	1,085.8
1952	30.0	213.3	15.8	2.9	2.7	3.5	8.1	19.8	40.5	95.5	303.4	871.5
1951	31.8	206.7	15.5	2.9	2.8	3.2	8.4	19.6	44.2	109.3	340.1	992.0
1950	31.6	232.1	15.3	2.8	3.0	3.0	8.6	20.3	43.6	111.8	329.2	1,034.1
WHITE, FEMALE												
1959	24.5	147.3	10.6	2.2	2.0	2.5	4.3	8.4	17.6	54.9	232.8	878.4
1958	25.5	156.3	12.1	1.9	2.2	2.8	4.3	9.1	19.7	59.7	237.7	930.8
1957	27.3	146.8	13.6	3.3	5.2	4.4	6.0	10.2	22.5	66.8	249.2	916.4
1956	21.9	145.5	10.9	2.1	1.7	1.7	4.0	7.0	15.5	53.1	215.2	821.4
1955	21.3	145.7	11.1	2.1	2.0	1.8	3.9	6.8	15.5	52.2	211.7	785.1
1954	19.4	150.3	11.8	2.0	1.7	1.9	3.6	6.8	14.2	46.9	189.4	683.2
1953	25.1	173.1	14.7	2.7	2.1	2.6	4.5	7.9	18.2	65.0	262.4	951.5
1952	22.8	180.3	15.5	2.8	2.0	2.3	4.3	7.4	16.8	57.3	233.1	815.9
1951	24.6	171.1	14.9	2.6	2.3	2.5	4.9	8.4	20.6	67.7	263.3	920.2
1950	24.5	189.5	13.9	2.6	2.4	2.7	4.8	8.9	20.2	68.4	264.0	918.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)												
ALL OTHER, BOTH SEXES												
1959	52.2	630.8	39.1	4.3	5.2	12.0	26.3	41.2	82.1	144.4	264.6	576.5
1958	57.3	649.9	39.0	4.2	5.5	14.3	25.2	48.3	99.6	171.6	322.6	747.0
1957	62.6	633.0	45.5	6.4	9.5	17.0	32.2	59.2	113.8	186.4	312.9	748.4
1956	48.7	579.9	40.7	4.8	4.5	10.6	23.1	40.0	76.7	137.7	248.9	604.9
1955	48.6	582.1	37.0	4.7	5.9	11.1	21.0	44.4	78.2	137.6	261.7	553.4
1954	47.7	598.8	40.6	5.0	5.8	11.1	20.9	42.9	74.9	131.0	223.8	560.0
1953	63.4	671.6	42.8	6.3	8.5	15.5	29.1	64.1	124.1	199.7	385.0	913.5
1952	58.2	729.3	48.2	6.1	7.6	15.4	30.3	53.3	97.9	158.4	292.9	653.1
1951	59.0	657.5	45.9	6.4	10.0	14.3	27.3	65.6	113.3	167.3	319.8	753.2
1950	59.0	651.4	50.4	6.4	7.2	15.2	29.6	64.6	114.0	203.6	336.4	817.2
ALL OTHER, MALE												
1959	61.7	685.7	43.6	4.2	5.6	14.4	32.3	55.6	108.1	186.4	321.5	675.9
1958	68.5	719.0	39.7	4.0	6.2	17.1	33.2	65.9	130.7	226.1	392.1	933.3
1957	71.9	667.9	48.3	6.7	8.8	18.7	39.7	75.4	143.0	227.2	377.8	857.7
1956	56.3	628.3	42.5	5.0	4.7	11.2	28.1	50.5	99.2	173.9	288.5	692.0
1955	57.6	640.2	40.6	4.7	6.6	12.9	27.9	57.4	98.8	176.0	303.2	704.2
1954	56.0	651.2	43.1	5.5	6.4	13.2	25.5	57.3	96.6	158.8	261.3	668.2
1953	73.7	714.5	49.6	6.9	9.6	18.2	38.5	79.1	153.5	234.0	417.5	1,066.7
1952	67.3	794.1	52.1	6.5	8.1	17.3	38.5	66.3	116.4	189.3	346.4	725.0
1951	67.2	726.4	48.5	7.3	10.3	15.6	31.4	78.9	135.9	191.7	354.7	842.1
1950	66.1	714.1	55.1	6.8	7.2	14.7	32.3	73.5	135.0	229.6	358.8	964.2
ALL OTHER, FEMALE												
1959	43.2	574.2	34.7	4.2	4.8	10.0	20.9	27.6	57.0	106.4	213.7	502.6
1958	46.6	581.0	38.3	4.5	5.0	12.0	17.9	31.6	69.8	121.6	259.5	634.2
1957	53.9	598.3	42.7	6.2	10.0	15.6	25.3	43.8	85.3	149.2	252.4	691.9
1956	41.6	531.8	38.9	4.6	4.3	10.2	18.4	29.9	54.7	104.0	211.6	544.4
1955	40.0	526.6	33.3	4.7	5.3	9.4	14.7	31.7	57.7	102.3	220.5	447.1
1954	40.0	546.8	38.2	4.4	5.2	9.2	16.6	29.0	53.2	105.2	186.5	503.1
1953	53.7	628.9	36.0	5.6	7.5	13.2	20.3	49.5	94.0	167.2	351.2	809.7
1952	49.6	665.3	44.3	5.6	7.2	13.8	22.6	40.6	78.8	129.4	242.2	603.4
1951	51.1	589.2	43.1	5.6	9.7	13.1	23.5	52.3	89.7	144.2	286.5	692.9
1950	52.3	589.0	45.6	6.1	7.2	15.7	27.0	55.7	92.5	177.3	314.3	712.2

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFLUENZA (480-483)												
BOTH SEXES												
1959	1.6	8.3	0.8	0.3	0.2	0.2	0.4	0.8	1.8	4.3	15.1	55.7
1958	2.6	11.0	1.1	0.2	0.4	0.4	0.6	1.4	3.2	7.8	24.0	94.3
1957	4.4	16.1	1.9	0.9	1.5	1.0	1.6	2.8	6.2	14.2	32.3	116.7
1956	1.4	7.5	0.8	0.1	0.2	0.1	0.3	0.6	1.3	3.9	14.6	57.0
1955	1.7	9.8	0.8	0.2	0.2	0.2	0.3	0.9	1.8	4.9	16.8	58.9
1954	1.7	11.6	1.0	0.2	0.2	0.2	0.4	0.8	1.6	5.0	16.3	52.0
1953	6.0	29.0	2.6	0.5	0.6	0.6	1.1	2.9	6.5	19.5	68.5	236.5
1952	3.6	24.0	2.0	0.5	0.5	0.5	1.0	1.9	3.7	10.5	37.0	128.4
1951	4.5	23.5	2.1	0.5	0.6	0.5	1.1	2.4	5.2	14.8	50.4	180.8
1950	4.4	23.3	2.3	0.5	0.5	0.6	1.1	2.4	5.3	14.3	50.2	170.4
MALE												
1959	1.7	9.5	0.9	0.3	0.2	0.2	0.5	0.9	2.1	5.0	15.5	61.1
1958	2.7	12.3	1.1	0.2	0.4	0.4	0.7	1.7	3.9	9.7	27.3	98.3
1957	4.7	17.5	2.0	0.9	1.3	0.8	1.7	3.3	7.5	18.3	35.1	130.0
1956	1.4	7.9	0.8	*	*	*	0.3	0.7	1.6	4.3	15.9	59.6
1955	1.6	10.5	0.7	0.2	0.2	0.2	0.4	1.0	2.0	5.5	16.9	55.3
1954	1.7	12.1	1.1	0.2	0.3	0.3	0.4	1.1	1.8	5.3	17.3	54.3
1953	6.2	31.0	2.6	0.5	0.6	0.4	1.2	3.2	7.5	22.2	75.1	256.7
1952	3.7	25.8	2.2	0.5	0.5	0.5	1.1	2.2	4.3	10.9	38.6	127.8
1951	4.6	24.0	2.3	0.6	0.6	0.6	1.2	2.7	5.9	16.3	50.9	182.8
1950	4.4	24.7	2.3	0.5	0.5	0.5	1.1	2.7	5.9	15.7	52.0	175.7
FEMALE												
1959	1.6	7.1	0.6	0.3	0.2	0.2	0.4	0.6	1.5	3.6	14.8	52.1
1958	2.4	9.8	1.1	0.2	0.4	0.4	0.5	1.1	2.5	6.0	21.4	91.7
1957	4.1	14.7	1.9	0.9	1.6	1.3	1.5	2.3	4.9	10.6	30.0	107.7
1956	1.4	7.1	0.7	*	*	*	0.3	0.6	1.0	3.6	13.5	55.1
1955	1.7	9.1	1.0	0.2	0.3	0.2	0.3	0.8	1.6	4.4	16.7	61.5
1954	1.6	11.2	1.0	0.2	*	0.2	0.4	0.6	1.5	4.8	15.4	50.4
1953	5.8	27.0	2.7	0.6	0.6	0.7	1.0	2.6	5.5	17.1	63.0	221.9
1952	3.6	22.2	1.8	0.6	0.5	0.5	0.9	1.7	3.2	10.0	35.6	128.9
1951	4.5	23.0	1.9	0.5	0.6	0.5	1.0	2.0	4.4	13.5	49.9	179.4
1950	4.4	21.9	2.3	0.5	0.5	0.8	1.1	2.1	4.7	13.0	48.7	166.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFLUENZA (480-483)												
WHITE, BOTH SEXES												
1959	1.4	5.4	0.6	0.3	0.2	0.2	0.3	0.6	1.4	3.6	14.8	57.1
1958	2.2	5.9	0.8	0.2	0.4	0.3	0.4	1.0	2.4	6.6	22.3	89.8
1957	3.9	9.6	1.5	0.9	1.4	0.9	1.2	2.2	5.0	12.8	30.7	116.3
1956	1.2	4.7	0.5	*	0.1	*	0.2	0.4	1.0	3.3	13.6	55.6
1955	1.4	5.7	0.5	0.1	0.1	0.2	0.3	0.6	1.3	3.9	16.0	57.5
1954	1.4	6.9	0.7	0.1	0.2	0.2	0.3	0.6	1.2	4.1	15.5	50.4
1953	5.2	21.1	2.0	0.4	0.4	0.4	0.7	1.9	4.5	16.1	65.2	230.4
1952	3.0	15.4	1.4	0.5	0.4	0.4	0.6	1.2	2.7	8.8	35.3	126.8
1951	4.0	15.5	1.6	0.4	0.4	0.4	0.7	1.5	4.0	13.3	48.5	180.3
1950	3.8	17.4	1.8	0.4	0.4	0.4	0.7	1.4	3.7	12.2	49.0	168.0
WHITE, MALE												
1959	1.5	6.2	0.7	0.3	*	*	0.4	0.7	1.8	4.4	15.3	62.9
1958	2.3	6.9	0.8	0.1	0.4	0.3	0.5	1.2	3.1	8.4	24.4	90.5
1957	4.2	10.8	1.5	0.9	1.2	0.7	1.3	2.8	6.6	16.9	33.5	129.9
1956	1.2	4.8	0.5	*	*	*	0.2	0.5	1.3	3.8	15.0	57.6
1955	1.3	6.1	0.4	*	*	*	0.3	0.7	1.4	4.4	16.4	53.4
1954	1.4	7.2	0.8	*	0.2	0.2	0.3	0.8	1.4	4.5	16.2	52.8
1953	5.4	22.9	1.8	0.4	0.4	0.3	0.9	2.2	5.5	18.8	72.3	249.6
1952	3.1	16.3	1.5	0.5	0.4	0.4	0.7	1.6	3.4	9.7	35.9	126.1
1951	4.1	15.5	1.9	0.4	0.4	0.4	0.9	2.0	4.8	15.1	49.4	181.0
1950	3.8	18.1	1.8	0.3	0.4	0.3	0.8	1.8	4.2	13.8	51.1	172.4
WHITE, FEMALE												
1959	1.4	4.6	0.5	0.3	*	0.2	0.3	0.5	1.2	2.9	14.4	53.4
1958	2.1	5.0	0.7	0.2	0.3	0.2	0.4	0.8	1.8	5.1	20.6	89.3
1957	3.6	8.3	1.6	0.9	1.6	1.1	1.2	1.5	3.6	9.1	28.6	107.0
1956	1.1	4.5	0.5	*	*	*	*	0.4	0.7	2.9	12.4	54.1
1955	1.4	5.4	0.7	*	*	*	0.2	0.5	1.1	3.5	15.8	60.5
1954	1.3	6.5	0.6	*	*	*	0.2	0.3	1.0	3.7	14.8	48.6
1953	5.0	19.3	2.2	0.5	0.5	0.5	0.6	1.6	3.6	13.7	59.5	216.4
1952	3.0	14.5	1.3	0.5	0.4	0.4	0.5	0.9	2.0	8.0	34.8	127.3
1951	3.9	15.4	1.4	0.4	0.4	0.4	0.5	1.0	3.3	11.7	47.8	179.8
1950	3.7	16.8	1.8	0.4	0.4	0.5	0.5	1.0	3.1	10.7	47.2	165.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFLUENZA (480-483)												
ALL OTHER, BOTH SEXES												
1959	2.9	25.1	1.6	0.5	*	*	1.2	2.4	5.8	11.9	19.3	38.2
1958	5.8	41.5	3.2	0.5	0.8	1.4	2.0	5.5	11.5	21.7	47.3	148.5
1957	8.4	55.3	4.4	1.2	2.1	2.5	4.7	9.2	18.6	32.0	53.6	120.3
1956	3.1	24.5	2.4	*	*	*	1.3	2.4	4.9	11.1	28.7	73.8
1955	4.1	35.0	2.4	*	0.9	0.9	1.2	3.9	8.2	16.7	27.7	75.9
1954	4.2	41.8	2.9	0.6	*	0.9	1.2	3.2	7.1	16.5	27.9	70.9
1953	12.8	80.2	6.8	1.3	2.0	2.2	4.6	12.5	30.2	60.3	115.8	311.5
1952	8.5	81.2	5.9	1.1	1.3	1.7	4.1	8.3	15.8	30.1	61.5	149.0
1951	9.2	75.6	5.5	1.2	1.8	1.7	3.9	10.6	19.7	32.5	77.4	187.2
1950	9.7	63.0	5.7	1.4	1.4	2.7	5.0	11.8	25.2	43.1	69.3	198.7
ALL OTHER, MALE												
1959	3.2	29.0	2.1	*	*	*	*	3.0	6.1	11.6	18.1	*
1958	6.6	45.0	3.0	*	*	*	2.4	7.0	12.8	25.8	63.3	192.6
1957	8.7	58.5	5.3	1.5	2.0	1.9	5.3	8.6	17.0	35.1	55.6	130.8
1956	3.1	26.8	2.7	*	*	*	*	2.6	4.7	11.1	27.7	84.0
1955	4.2	38.7	2.0	*	*	*	*	3.7	9.2	18.8	24.0	*
1954	4.2	43.5	2.8	*	*	*	*	3.3	6.6	15.3	31.9	*
1953	13.3	84.6	7.8	1.3	2.6	1.7	4.6	12.9	30.6	62.5	113.2	347.6
1952	8.8	90.5	7.2	1.2	*	*	4.1	8.1	14.1	25.4	75.5	150.0
1951	9.2	80.5	5.7	1.5	*	2.1	3.2	9.4	20.3	30.6	71.7	205.3
1950	9.6	70.6	5.8	1.6	*	1.8	4.2	10.7	26.1	41.0	65.4	214.3
ALL OTHER, FEMALE												
1959	2.7	21.2	*	*	*	*	*	*	5.6	12.1	20.5	*
1958	5.0	37.9	3.5	*	*	1.5	1.7	4.0	10.4	17.9	32.7	121.1
1957	8.2	52.1	3.6	*	2.1	3.0	4.2	9.7	20.1	29.2	51.7	116.2
1956	3.1	22.3	2.1	*	*	*	*	2.3	5.1	11.1	29.7	66.7
1955	3.9	31.6	2.8	*	*	*	*	4.1	7.2	14.8	31.1	73.5
1954	4.1	40.0	3.1	*	*	*	*	3.2	7.5	17.6	23.8	71.9
1953	12.4	75.7	5.7	1.2	1.6	2.7	4.6	12.0	29.8	58.0	117.4	287.1
1952	8.2	72.0	4.7	*	*	1.8	4.1	8.4	17.5	34.5	48.3	148.3
1951	9.3	70.7	5.3	*	2.0	*	4.6	11.8	19.1	34.4	82.9	175.0
1950	9.8	55.5	5.6	*	1.5	3.5	5.7	12.8	24.2	45.2	73.2	187.6

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)												
BOTH SEXES												
1959	29.6	230.8	14.5	2.2	2.5	3.6	7.2	14.7	31.7	82.3	266.4	868.6
1958	30.6	234.3	14.9	2.1	2.5	3.9	7.2	15.9	35.7	88.2	268.8	893.8
1957	31.4	223.7	16.4	2.9	4.0	4.5	8.6	17.7	38.1	92.7	266.3	863.1
1956	26.8	217.8	14.6	2.2	2.2	2.9	6.7	13.4	29.1	76.6	243.5	796.5
1955	25.4	211.8	14.1	2.4	2.5	2.9	6.4	13.3	27.7	71.5	233.0	732.2
1954	23.8	215.6	15.1	2.2	2.2	3.0	6.3	13.3	26.0	65.8	204.3	661.2
1953	27.0	228.5	16.2	2.6	2.6	3.6	7.6	15.7	31.7	77.1	242.2	764.3
1952	26.1	242.9	17.9	2.7	2.5	3.7	7.7	15.4	30.0	71.5	229.6	696.8
1951	26.9	228.5	16.9	2.7	2.9	3.5	7.7	16.5	33.1	79.0	249.1	754.2
1950	26.9	244.7	16.7	2.7	2.7	3.5	7.9	16.9	32.9	82.8	246.2	784.0
MALE												
1959	34.2	256.4	15.5	2.2	2.8	4.1	8.8	19.9	44.7	113.3	330.2	974.6
1958	35.8	259.1	15.1	2.2	2.9	4.4	9.4	21.8	50.8	122.0	333.8	1,008.7
1957	36.8	248.4	17.0	3.0	3.9	4.6	10.7	24.6	54.3	126.6	325.3	967.4
1956	30.9	240.2	15.1	2.1	2.6	3.3	8.1	18.2	41.1	102.5	295.9	870.3
1955	29.3	231.9	15.0	2.4	2.8	3.3	8.2	18.2	38.7	93.4	279.5	780.5
1954	27.6	236.0	15.8	2.4	2.4	3.6	7.9	18.5	36.6	86.9	241.6	719.7
1953	31.7	248.3	17.5	2.7	3.0	4.2	10.3	22.3	45.5	100.4	287.8	827.7
1952	30.2	261.7	18.2	2.8	2.8	4.4	10.1	22.0	42.0	91.9	267.7	733.0
1951	30.8	250.9	17.2	2.9	3.1	3.8	9.5	22.6	45.1	99.4	290.0	798.0
1950	30.8	268.3	17.8	2.9	3.0	3.7	9.8	22.7	44.7	104.5	279.1	852.9
FEMALE												
1959	25.0	204.5	13.5	2.2	2.2	3.2	5.7	9.7	19.4	55.2	216.8	799.3
1958	25.5	208.6	14.8	2.0	2.2	3.5	5.2	10.2	21.4	58.4	217.7	817.2
1957	26.2	198.2	15.8	2.7	4.1	4.4	6.5	11.1	22.7	62.5	219.4	791.9
1956	22.7	194.7	14.1	2.3	1.9	2.5	5.3	8.6	17.6	53.5	201.5	745.1
1955	21.7	191.0	13.2	2.3	2.1	2.5	4.7	8.4	17.2	51.7	195.4	698.0
1954	20.0	194.5	14.4	2.1	2.0	2.5	4.6	8.3	15.7	46.6	173.9	619.3
1953	22.5	208.0	14.9	2.5	2.2	3.1	5.1	9.3	18.3	55.8	204.8	718.7
1952	22.1	223.3	17.5	2.6	2.2	3.1	5.3	8.9	18.1	52.8	198.0	670.9
1951	23.0	205.3	16.5	2.5	2.7	3.2	5.9	10.5	21.2	60.1	214.7	723.1
1950	23.1	220.3	15.5	2.5	2.5	3.3	6.1	11.2	20.9	62.6	218.2	736.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)												
WHITE, BOTH SEXES												
1959	27.1	166.1	10.7	1.9	2.1	2.6	5.2	12.2	27.4	78.1	268.0	895.7
1958	28.0	171.7	11.6	1.9	2.2	2.8	5.4	13.1	30.8	83.1	268.3	918.4
1957	28.6	164.4	12.4	2.6	3.6	3.3	6.4	14.4	32.9	87.5	266.9	881.5
1956	24.5	161.7	10.9	1.9	2.0	2.0	4.9	10.8	25.3	72.5	245.1	818.5
1955	23.1	157.3	11.0	2.1	2.1	2.0	4.9	10.4	24.1	67.3	233.0	753.0
1954	21.4	161.6	11.6	1.9	1.8	2.2	4.7	10.6	22.5	61.7	204.9	674.2
1953	24.2	172.8	13.2	2.2	2.1	2.4	5.7	12.0	26.6	71.8	240.4	777.5
1952	23.3	181.7	14.2	2.4	2.0	2.5	5.6	12.3	25.8	66.8	229.5	712.4
1951	24.2	173.8	13.6	2.3	2.2	2.4	5.9	12.5	28.3	74.3	249.6	769.8
1950	24.3	193.8	12.8	2.3	2.3	2.5	6.0	13.2	28.2	77.1	244.8	797.9
WHITE, MALE												
1959	31.2	188.5	11.3	1.9	2.4	2.9	6.4	16.5	39.2	108.0	332.2	1,004.9
1958	32.6	191.3	11.7	1.9	2.5	3.0	7.0	18.0	44.4	115.4	334.2	1,034.8
1957	33.6	189.3	12.9	2.7	3.5	3.2	8.0	20.2	47.7	121.1	325.6	987.5
1956	28.2	181.6	11.3	1.8	2.3	2.5	6.0	15.1	36.3	97.3	298.6	892.1
1955	26.4	173.5	11.5	2.1	2.4	2.3	6.1	14.6	34.2	88.0	279.5	793.2
1954	24.8	178.7	12.1	2.1	2.0	2.7	6.1	14.8	32.0	82.2	242.5	729.4
1953	28.4	190.9	13.8	2.2	2.5	2.8	7.6	17.7	39.0	94.3	286.5	836.2
1952	26.9	197.0	14.3	2.4	2.3	3.1	7.4	18.2	37.1	85.8	267.5	745.5
1951	27.7	191.2	13.6	2.5	2.4	2.7	7.4	17.6	39.4	94.2	290.7	811.0
1950	27.8	214.1	13.5	2.5	2.6	2.7	7.8	18.5	39.4	98.0	278.1	861.8
WHITE, FEMALE												
1959	23.1	142.7	10.1	2.0	1.9	2.3	4.0	7.9	16.4	52.0	218.4	825.0
1958	23.5	151.4	11.4	1.8	1.9	2.5	3.9	8.3	18.0	54.7	217.1	841.4
1957	23.8	138.5	12.0	2.4	3.6	3.3	4.8	8.7	18.9	57.7	220.7	809.4
1956	20.8	141.0	10.5	2.1	1.6	1.6	3.9	6.6	14.9	50.2	202.8	767.3
1955	19.9	140.3	10.4	2.0	1.8	1.7	3.7	6.3	14.3	48.7	195.9	724.6
1954	18.1	143.8	11.1	1.8	1.6	1.7	3.4	6.4	13.3	43.2	174.6	634.6
1953	20.2	153.7	12.5	2.2	1.6	2.1	3.9	6.3	14.6	51.3	202.9	735.0
1952	19.8	165.8	14.2	2.3	1.7	2.0	3.8	6.5	14.7	49.3	198.3	688.6
1951	20.8	155.6	13.5	2.2	2.0	2.2	4.4	7.4	17.4	56.0	215.5	740.4
1950	20.7	172.7	12.1	2.1	2.1	2.2	4.3	7.9	17.1	57.7	216.8	753.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)												
ALL OTHER, BOTH SEXES												
1959	49.2	605.7	37.5	3.7	4.7	11.5	25.1	38.7	76.2	132.5	245.2	538.2
1958	51.5	608.5	35.8	3.8	4.8	12.9	23.2	42.8	88.1	149.9	275.3	598.5
1957	54.2	577.7	41.1	5.3	7.4	14.5	27.5	50.0	95.2	154.4	259.3	628.1
1956	45.6	555.3	38.3	4.4	4.2	10.0	21.7	37.6	71.8	126.6	220.1	531.1
1955	44.5	547.0	34.5	4.4	5.0	10.1	19.9	40.5	70.0	120.9	234.0	477.6
1954	43.6	557.0	37.7	4.3	5.1	10.2	19.7	39.7	67.8	114.5	195.9	489.1
1953	50.5	591.5	36.0	5.0	6.4	13.3	24.5	51.7	93.8	139.5	269.2	601.9
1952	49.7	648.1	42.3	4.9	6.3	13.8	26.2	45.1	82.1	128.3	231.4	504.1
1951	49.7	581.9	40.4	5.2	8.2	12.6	23.4	55.0	93.6	134.8	242.4	566.0
1950	49.3	588.4	44.6	5.0	5.8	12.5	24.6	52.8	88.8	160.4	267.0	618.5
ALL OTHER, MALE												
1959	58.5	656.7	41.5	3.7	5.1	13.7	31.0	52.6	102.0	174.7	303.5	634.5
1958	62.0	674.0	36.7	3.6	5.4	15.7	30.8	59.0	118.0	200.3	328.8	740.7
1957	63.2	609.4	43.0	5.2	6.8	16.8	34.4	66.7	126.0	192.1	322.2	726.9
1956	53.2	601.5	39.8	4.7	4.5	10.6	27.0	47.9	94.5	162.8	260.8	608.0
1955	53.3	601.5	38.6	4.4	5.8	12.1	26.5	53.8	89.6	157.2	279.2	625.0
1954	51.7	607.7	40.2	4.7	5.6	12.4	24.5	54.0	90.0	143.5	229.4	595.5
1953	60.3	629.9	41.8	5.6	7.0	16.5	33.9	66.2	123.0	171.5	304.4	719.0
1952	58.5	703.6	44.9	5.3	6.7	15.8	34.4	58.2	102.3	163.9	270.9	575.0
1951	58.0	645.9	42.9	5.8	8.8	13.5	28.3	69.5	115.6	161.2	283.0	636.8
1950	56.5	643.5	49.3	5.2	5.9	12.9	28.2	62.8	108.9	188.5	293.4	749.9
ALL OTHER, FEMALE												
1959	40.5	553.0	33.5	3.7	4.2	9.7	19.8	25.7	51.5	94.3	193.2	466.7
1958	41.6	543.1	34.8	3.9	4.2	10.4	16.3	27.6	59.5	103.8	226.8	513.2
1957	45.7	546.2	39.1	5.3	8.0	12.6	21.1	34.0	65.2	120.0	200.7	575.7
1956	38.4	509.5	36.8	4.0	3.9	9.5	16.8	27.7	49.6	92.8	181.9	477.8
1955	36.1	495.1	30.5	4.4	4.3	8.4	13.7	27.6	50.5	87.5	189.4	373.5
1954	35.9	506.8	35.2	4.0	4.6	8.2	15.2	25.8	45.7	87.7	162.7	431.3
1953	41.3	553.2	30.3	4.4	5.9	10.5	15.7	37.5	64.2	109.2	233.9	522.6
1952	41.4	593.3	39.6	4.5	6.0	12.0	18.5	32.1	61.3	95.0	194.0	455.2
1951	41.9	518.5	37.9	4.7	7.7	11.8	18.8	40.5	70.6	109.8	203.6	517.9
1950	42.4	533.6	40.0	4.8	5.7	12.2	21.3	42.8	68.3	132.1	241.1	524.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BRONCHITIS (500-502)												
BOTH SEXES												
1959	2.2	18.0	1.9	0.3	0.2	0.2	0.4	1.2	3.6	8.1	13.1	31.6
1958	2.3	19.3	2.5	0.2	0.1	0.2	0.4	1.1	3.9	8.0	14.6	33.1
1957	2.1	19.1	2.0	0.3	0.2	0.2	0.3	1.0	3.3	7.2	13.5	31.5
1956	1.9	20.4	2.1	0.3	*	0.1	0.2	0.7	3.0	5.4	12.9	33.6
1955	1.9	17.9	2.3	0.3	0.1	0.1	0.3	0.7	2.7	5.7	12.6	34.9
1954	1.8	18.8	2.1	0.2	*	0.1	0.2	0.8	2.5	5.3	13.5	32.4
1953	1.9	19.0	2.5	0.3	0.1	0.1	0.3	0.9	2.7	5.9	14.3	36.5
1952	1.9	19.4	2.6	0.3	0.1	0.2	0.4	0.9	2.4	5.9	13.5	34.9
1951	1.9	17.6	2.4	0.3	0.1	0.1	0.3	0.8	2.7	5.9	15.0	44.2
1950	2.0	18.6	2.5	0.3	0.1	0.2	0.3	0.8	2.7	6.1	15.8	58.6
MALE												
1959	3.1	21.8	2.1	0.3	0.2	*	0.4	1.6	6.2	14.4	19.8	42.0
1958	3.1	21.9	2.8	0.3	*	0.2	0.4	1.5	6.5	13.7	21.0	42.7
1957	2.8	22.1	2.2	0.3	*	*	0.4	1.6	5.5	11.8	18.3	38.0
1956	2.5	22.7	2.4	0.3	*	*	0.2	1.0	5.1	8.9	18.4	37.3
1955	2.4	21.1	2.7	0.3	*	*	0.3	0.9	4.5	9.1	15.4	38.1
1954	2.3	22.3	2.4	0.2	*	*	0.2	1.1	4.1	8.2	17.9	31.6
1953	2.5	21.4	3.1	0.3	*	*	0.4	1.3	4.6	9.1	17.9	39.8
1952	2.4	21.8	2.9	0.3	*	0.2	0.5	1.4	4.0	8.4	16.8	35.9
1951	2.4	20.0	2.8	0.3	*	*	0.4	1.3	4.3	8.7	17.7	54.7
1950	2.4	22.0	2.8	0.4	*	*	0.4	1.1	4.1	8.2	17.0	70.9
FEMALE												
1959	1.3	14.1	1.6	0.3	*	0.2	0.3	0.8	1.1	2.7	7.9	24.9
1958	1.5	16.6	2.1	0.2	*	0.2	0.4	0.6	1.3	2.9	9.6	26.7
1957	1.4	16.1	1.8	0.2	*	0.2	0.3	0.5	1.2	3.1	9.6	27.1
1956	1.3	18.1	1.7	0.2	*	*	0.2	0.3	1.0	2.3	8.5	31.1
1955	1.4	14.6	1.9	0.2	*	*	0.3	0.5	1.0	2.7	10.3	32.7
1954	1.3	15.2	1.8	0.2	*	*	*	0.5	0.9	2.6	9.9	32.9
1953	1.4	16.5	1.9	0.3	*	*	0.2	0.5	0.8	2.9	11.4	34.1
1952	1.4	17.0	2.2	0.3	*	*	0.3	0.3	1.0	3.5	10.8	34.1
1951	1.4	15.2	1.9	0.3	*	*	0.3	0.4	1.2	3.3	12.7	36.7
1950	1.6	15.0	2.2	0.2	*	0.2	0.2	0.6	1.2	4.1	14.9	50.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BRONCHITIS (500-502)												
WHITE, BOTH SEXES												
1959	2.2	15.8	1.8	0.3	0.2	0.2	0.3	1.2	3.6	8.3	13.5	33.4
1958	2.3	17.1	2.3	0.2	0.1	0.2	0.3	1.0	3.7	8.1	15.1	34.8
1957	2.1	16.7	2.0	0.3	0.1	0.1	0.3	0.9	3.2	7.4	13.7	32.3
1956	1.9	17.8	2.0	0.3	*	0.1	0.2	0.7	3.0	5.5	13.3	34.8
1955	1.8	15.3	2.3	0.2	*	0.1	0.2	0.6	2.7	6.0	13.0	36.1
1954	1.8	16.0	2.0	0.2	*	0.1	0.1	0.7	2.5	5.4	14.0	33.5
1953	1.9	16.8	2.4	0.3	0.1	*	0.3	0.8	2.6	5.9	14.7	37.6
1952	1.8	16.3	2.5	0.3	0.1	0.1	0.3	0.8	2.4	5.9	13.9	36.4
1951	1.9	15.5	2.3	0.3	0.1	0.1	0.3	0.8	2.7	6.0	15.1	45.3
1950	1.9	16.7	2.4	0.3	*	0.1	0.2	0.7	2.6	6.0	16.3	60.7
WHITE, MALE												
1959	3.1	19.7	2.0	0.3	*	*	0.4	1.6	6.4	15.0	20.6	44.5
1958	3.1	19.7	2.6	0.2	*	*	0.3	1.6	6.4	14.1	22.0	44.9
1957	2.8	19.9	2.2	0.3	*	*	0.3	1.4	5.4	12.2	18.7	39.5
1956	2.5	19.7	2.4	0.3	*	*	0.2	1.1	5.0	9.1	19.5	38.4
1955	2.4	18.2	2.7	0.3	*	*	0.2	0.8	4.6	9.5	16.0	39.1
1954	2.3	19.6	2.2	0.2	*	*	*	1.0	4.3	8.3	19.0	33.0
1953	2.5	18.8	3.0	0.3	*	*	0.3	1.2	4.5	9.2	18.4	41.0
1952	2.3	18.0	2.8	0.3	*	*	0.4	1.3	4.0	8.6	17.3	37.5
1951	2.4	17.5	2.7	0.3	*	*	0.3	1.2	4.4	8.9	18.0	57.0
1950	2.4	20.0	2.7	0.4	*	*	0.3	1.0	4.2	8.2	17.6	74.3
WHITE, FEMALE												
1959	1.3	11.9	1.5	0.3	*	*	0.3	0.8	1.1	2.6	8.0	26.2
1958	1.4	14.5	2.1	0.2	*	*	0.3	0.5	1.3	2.8	9.8	28.0
1957	1.3	13.5	1.7	0.2	*	*	0.2	0.4	1.1	3.0	9.8	27.4
1956	1.3	15.9	1.6	0.2	*	*	0.2	0.3	1.0	2.2	8.4	32.3
1955	1.3	12.3	2.0	0.2	*	*	0.2	0.5	0.8	2.8	10.6	34.0
1954	1.2	12.3	1.8	0.2	*	*	*	0.5	0.8	2.7	10.0	33.8
1953	1.3	14.7	1.8	0.3	*	*	*	0.4	0.8	2.9	11.6	35.0
1952	1.4	14.6	2.1	0.3	*	*	0.2	0.3	0.9	3.4	11.0	35.5
1951	1.4	13.4	1.9	0.3	*	*	0.2	0.4	1.0	3.2	12.8	37.0
1950	1.5	13.2	2.1	0.2	*	*	*	0.4	1.0	4.0	15.2	51.3

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BRONCHITIS (500-502)												
ALL OTHER, BOTH SEXES												
1959	2.2	30.8	2.7	*	*	*	0.9	1.2	2.8	5.5	7.9	*
1958	2.6	32.1	3.2	*	*	*	0.9	1.4	4.9	7.0	7.2	*
1957	2.5	33.6	2.2	*	*	*	0.9	2.2	4.0	5.0	9.6	*
1956	2.2	36.1	2.2	*	*	*	*	*	3.3	4.7	*	*
1955	2.1	33.9	2.2	*	*	*	0.9	1.1	3.8	3.0	*	*
1954	2.0	36.1	2.6	*	*	*	*	*	1.8	4.0	*	*
1953	2.3	32.8	3.2	*	*	*	*	1.4	3.7	5.3	9.0	*
1952	2.4	40.0	3.1	*	*	*	*	1.9	2.9	5.2	*	*
1951	2.2	31.4	2.8	*	*	*	*	1.5	3.3	5.5	12.4	*
1950	2.4	31.3	3.4	*	*	*	1.0	1.8	3.8	7.2	*	*
ALL OTHER, MALE												
1959	2.7	34.7	2.8	*	*	*	*	*	4.7	7.3	*	*
1958	3.0	35.3	3.7	*	*	*	*	*	8.0	9.6	*	*
1957	2.9	35.9	2.3	*	*	*	*	3.0	6.0	6.9	*	*
1956	2.6	41.5	2.5	*	*	*	*	*	5.4	6.7	*	*
1955	2.5	39.5	2.6	*	*	*	*	*	4.2	*	*	*
1954	2.4	39.5	3.6	*	*	*	*	*	*	6.3	*	*
1953	2.8	38.0	4.0	*	*	*	*	*	5.9	7.3	*	*
1952	2.9	47.7	3.0	*	*	*	*	3.0	3.8	*	*	*
1951	2.5	36.4	3.4	*	*	*	*	*	*	6.7	*	*
1950	2.6	35.6	4.2	*	*	*	*	*	*	8.0	*	*
ALL OTHER, FEMALE												
1959	1.7	26.8	2.6	*	*	*	*	*	*	*	*	*
1958	2.1	29.0	2.6	*	*	*	*	*	*	4.7	*	*
1957	2.1	31.3	2.1	*	*	*	*	*	*	*	*	*
1956	1.7	30.7	2.0	*	*	*	*	*	*	*	*	*
1955	1.8	28.5	*	*	*	*	*	*	3.4	*	*	*
1954	1.6	32.8	*	*	*	*	*	*	*	*	*	*
1953	1.8	27.7	2.4	*	*	*	*	*	*	*	*	*
1952	2.0	32.4	3.1	*	*	*	*	*	*	*	*	*
1951	1.9	26.6	*	*	*	*	*	*	*	*	*	*
1950	2.2	27.0	2.6	*	*	*	*	*	4.2	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ULCER OF STOMACH AND DUODEMA (540, 541)												
BOTH SEXES												
1959	6.0	1.6	*	*	0.2	0.7	2.7	7.1	14.9	28.2	52.1	66.8
1958	6.2	1.9	*	*	0.2	0.9	2.8	7.5	16.1	28.7	51.7	70.7
1957	6.1	1.7	*	*	0.1	0.7	2.8	7.6	16.0	28.6	47.4	72.0
1956	6.0	1.3	*	*	0.2	0.8	2.6	7.5	15.6	28.4	48.5	68.5
1955	5.9	1.5	*	*	0.2	0.8	2.9	7.8	16.0	28.2	44.9	55.8
1954	5.8	1.9	*	*	0.2	0.8	3.2	8.1	15.9	27.3	41.4	53.9
1953	5.8	2.0	*	*	0.2	0.9	3.1	8.4	16.6	26.6	41.9	50.1
1952	5.6	1.4	*	*	0.2	0.8	3.3	8.4	16.5	25.9	37.3	44.6
1951	5.5	1.2	*	*	0.2	1.0	3.3	8.4	16.5	25.6	35.3	43.2
1950	5.5	1.6	*	*	0.3	0.9	3.4	8.6	16.7	24.8	34.6	38.3
MALE												
1959	9.2	2.4	*	*	0.2	1.1	4.1	11.4	25.4	46.0	80.5	96.9
1958	9.5	1.9	*	*	0.2	1.4	4.1	11.9	27.1	47.4	81.7	100.0
1957	9.4	1.7	*	*	*	1.0	4.5	12.5	26.7	47.4	74.0	100.3
1956	9.4	1.7	*	*	0.3	1.3	4.3	12.4	26.5	47.4	75.3	100.0
1955	9.3	2.1	*	*	0.3	1.2	4.7	13.0	27.2	46.4	71.4	76.1
1954	9.4	2.2	*	*	0.3	1.3	5.2	13.7	27.3	45.5	66.6	79.3
1953	9.5	2.2	*	*	0.2	1.4	5.1	14.3	28.8	45.2	67.6	77.5
1952	9.2	2.2	*	*	0.3	1.4	5.4	14.3	28.3	43.7	58.8	66.3
1951	9.0	1.6	*	*	0.3	1.6	5.7	14.3	27.8	42.8	55.5	66.0
1950	8.9	2.2	*	*	0.4	1.5	5.7	14.3	28.5	41.0	55.8	57.8
FEMALE												
1959	3.0	*	*	*	*	0.3	1.3	3.0	5.1	12.6	30.0	47.1
1958	3.0	1.8	*	*	*	0.4	1.4	3.2	5.7	12.2	28.1	51.2
1957	2.9	1.7	*	*	*	0.4	1.1	2.9	5.8	12.0	26.3	52.6
1956	2.7	*	*	*	*	0.4	1.0	2.8	5.2	11.3	27.0	46.6
1955	2.6	*	*	*	*	0.4	1.1	2.6	5.3	11.8	23.5	41.4
1954	2.4	1.6	*	*	*	0.4	1.3	2.6	4.9	10.8	20.9	35.8
1953	2.3	1.8	*	*	*	0.3	1.1	2.7	4.8	9.7	20.8	30.3
1952	2.2	*	*	*	*	0.4	1.2	2.5	4.9	9.6	19.5	29.1
1951	2.1	*	*	*	*	0.4	1.0	2.5	5.3	9.7	18.3	26.9
1950	2.1	*	*	*	*	0.2	1.3	2.9	4.8	9.6	16.5	24.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ULCER OF STOMACH AND DUODEMA (540, 541)												
WHITE, BOTH SEXES												
1959	6.3	1.2	*	*	0.1	0.7	2.4	6.9	15.0	28.8	53.9	70.2
1958	6.4	1.3	*	*	0.1	0.8	2.5	7.4	16.1	29.2	53.0	74.8
1957	6.3	1.3	*	*	0.1	0.6	2.6	7.5	15.9	29.4	48.8	75.3
1956	6.2	0.9	*	*	0.2	0.7	2.5	7.4	15.6	29.2	50.1	72.8
1955	6.1	1.1	*	*	0.2	0.7	2.7	7.6	16.1	28.9	46.1	59.0
1954	6.0	1.5	*	*	0.2	0.7	3.0	7.9	15.8	28.1	42.9	56.9
1953	6.0	1.6	*	*	0.1	0.8	3.0	8.1	16.5	27.3	43.4	52.3
1952	5.8	1.0	*	*	0.1	0.8	3.1	8.2	16.7	26.9	38.4	46.3
1951	5.7	1.1	*	*	0.2	0.9	3.2	8.1	16.4	26.2	37.2	44.6
1950	5.6	1.4	*	*	0.2	0.8	3.2	8.3	16.8	25.2	35.4	41.0
WHITE, MALE												
1959	9.5	1.6	*	*	*	1.0	3.8	11.1	25.6	47.1	83.7	102.1
1958	9.9	1.4	*	*	*	1.3	3.9	11.8	27.3	48.5	84.4	107.0
1957	9.7	1.2	*	*	*	0.9	4.3	12.3	26.7	48.8	76.5	104.8
1956	9.8	1.2	*	*	0.2	1.2	4.1	12.4	26.7	49.2	78.3	106.3
1955	9.7	1.7	*	*	0.3	1.1	4.5	12.9	27.6	47.9	73.7	80.3
1954	9.7	1.7	*	*	0.3	1.2	4.9	13.6	27.4	47.1	69.4	83.0
1953	9.8	1.5	*	*	*	1.3	5.0	14.0	28.8	46.6	70.2	81.0
1952	9.5	1.5	*	*	0.2	1.3	5.2	14.2	28.7	45.2	60.7	69.6
1951	9.3	1.4	*	*	0.3	1.5	5.5	14.1	28.0	44.0	59.1	67.5
1950	9.1	1.9	*	*	0.3	1.4	5.4	14.1	28.6	42.1	57.3	61.4
WHITE, FEMALE												
1959	3.1	*	*	*	*	0.3	1.1	2.8	5.2	12.9	30.9	49.6
1958	3.1	1.2	*	*	*	0.3	1.2	3.0	5.5	12.3	28.5	53.6
1957	3.0	1.5	*	*	*	0.4	1.0	2.8	5.7	12.3	27.0	55.1
1956	2.8	*	*	*	*	0.3	0.9	2.6	5.0	11.4	27.7	49.5
1955	2.7	*	*	*	*	0.3	1.1	2.4	5.1	11.8	24.0	43.9
1954	2.4	*	*	*	*	0.3	1.1	2.4	4.7	10.9	21.6	38.2
1953	2.3	1.7	*	*	*	0.3	1.0	2.4	4.6	9.8	21.6	31.5
1952	2.2	*	*	*	*	0.3	1.1	2.3	4.8	10.0	20.0	29.5
1951	2.1	*	*	*	*	0.3	0.9	2.2	5.0	9.8	19.0	28.3
1950	2.1	*	*	*	*	0.2	1.1	2.6	4.9	9.6	16.9	26.8

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ULCER OF STOMACH AND DUODEMA (540, 541)												
ALL OTHER, BOTH SEXES												
1959	4.2	4.3	*	*	*	1.1	4.8	9.1	13.9	20.8	27.2	*
1958	4.5	5.2	*	*	*	1.7	4.8	8.7	15.6	21.8	33.6	*
1957	4.2	4.0	*	*	*	1.5	3.9	9.0	16.7	19.0	27.9	31.3
1956	4.0	3.8	*	*	*	1.7	3.6	8.7	15.9	18.6	26.5	*
1955	4.2	4.0	*	*	*	1.6	4.2	9.3	15.4	19.3	28.1	*
1954	4.3	4.4	*	*	*	1.5	5.0	9.8	16.7	18.1	20.1	*
1953	4.4	4.5	*	*	*	1.6	3.8	11.4	18.0	18.4	20.1	*
1952	4.0	*	*	*	*	1.5	4.7	9.7	14.3	14.9	22.1	*
1951	4.2	*	*	*	*	1.7	4.6	11.0	17.2	18.7	*	*
1950	4.4	*	*	*	*	1.7	5.5	10.7	15.9	18.5	23.1	*
ALL OTHER, MALE												
1959	6.5	7.0	*	*	*	1.7	7.1	14.0	23.6	33.6	39.6	*
1958	6.5	*	*	*	*	1.9	6.6	13.0	24.1	34.6	46.8	*
1957	6.6	*	*	*	*	2.3	6.2	14.6	26.3	30.9	41.5	*
1956	5.9	*	*	*	*	2.3	5.5	12.6	24.7	27.0	36.9	*
1955	6.3	*	*	*	*	2.3	6.9	14.2	22.9	28.7	41.6	*
1954	6.5	*	*	*	*	2.3	7.7	14.6	26.0	27.6	29.4	*
1953	6.9	*	*	*	*	2.9	6.0	17.2	28.3	29.1	31.6	*
1952	6.4	*	*	*	*	2.2	7.2	15.2	22.9	26.0	33.6	*
1951	6.4	*	*	*	*	2.7	7.3	16.6	24.9	29.1	*	*
1950	7.0	*	*	*	*	3.1	8.4	16.0	27.8	26.8	35.7	*
ALL OTHER, FEMALE												
1959	2.0	*	*	*	*	*	2.7	4.5	4.5	9.2	16.1	*
1958	2.7	*	*	*	*	1.5	3.2	4.7	7.5	10.1	21.6	*
1957	2.0	*	*	*	*	*	1.8	3.7	7.3	8.2	15.2	*
1956	2.2	*	*	*	*	*	1.9	5.0	7.2	10.9	16.7	*
1955	2.2	*	*	*	*	*	1.7	4.6	8.1	10.7	15.2	*
1954	2.2	*	*	*	*	*	2.5	5.1	7.3	9.4	*	*
1953	2.0	*	*	*	*	*	1.8	5.7	7.5	8.4	*	*
1952	1.7	*	*	*	*	*	2.4	4.2	5.5	*	*	*
1951	2.1	*	*	*	*	*	2.0	5.4	9.2	9.0	*	*
1950	1.9	*	*	*	*	*	2.8	5.4	*	10.2	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
APPENDICITIS (550-553)												
BOTH SEXES												
1959	1.0	*	0.3	0.4	0.4	0.3	0.6	1.1	2.0	3.8	6.7	11.4
1958	1.1	0.6	0.3	0.4	0.3	0.4	0.5	1.2	2.0	4.1	7.5	11.2
1957	1.2	*	0.4	0.5	0.3	0.3	0.6	1.3	2.4	4.4	7.7	10.6
1956	1.3	0.6	0.5	0.4	0.4	0.4	0.6	1.5	2.3	4.7	8.3	13.4
1955	1.4	*	0.6	0.4	0.4	0.5	0.7	1.7	2.7	5.3	8.4	12.6
1954	1.4	*	0.5	0.6	0.5	0.5	0.8	1.5	3.0	5.0	8.6	14.0
1953	1.5	*	0.6	0.7	0.6	0.7	1.0	2.0	3.0	5.1	7.6	10.1
1952	1.7	*	0.8	0.8	0.8	0.7	1.2	2.2	3.4	5.2	9.4	13.3
1951	1.9	*	1.1	0.9	0.9	0.8	1.4	2.2	3.4	5.8	9.9	11.7
1950	2.0	*	1.3	1.0	1.0	0.9	1.5	2.5	4.0	6.9	8.7	13.0
MALE												
1959	1.4	*	0.3	0.5	0.5	0.3	0.8	1.5	3.0	5.3	8.9	16.3
1958	1.3	*	0.3	0.4	0.4	0.4	0.6	1.6	2.7	5.7	10.5	12.5
1957	1.5	*	0.4	0.6	0.4	0.3	0.8	2.0	3.4	6.5	9.6	11.0
1956	1.6	*	0.4	0.5	0.6	0.6	0.8	2.1	3.3	6.4	10.5	15.0
1955	1.8	*	0.7	0.5	0.5	0.6	0.9	2.5	3.8	7.0	10.2	16.0
1954	1.8	*	0.6	0.7	0.6	0.7	1.0	2.2	4.2	7.0	11.2	16.4
1953	2.0	*	0.8	0.7	0.8	0.8	1.4	2.9	4.3	7.0	10.4	13.8
1952	2.2	*	1.0	1.0	0.9	0.8	1.5	3.1	4.5	7.1	12.0	16.5
1951	2.3	*	1.2	1.1	1.1	0.9	2.0	2.7	4.7	7.3	12.5	14.1
1950	2.6	*	1.3	1.1	1.2	1.2	2.1	3.3	5.3	9.1	12.1	17.3
FEMALE												
1959	0.7	*	*	0.3	0.3	0.2	0.4	0.7	1.1	2.4	4.9	8.1
1958	0.8	*	*	0.3	0.3	0.3	0.3	0.8	1.3	2.7	5.1	10.3
1957	0.8	*	0.4	0.3	0.3	0.2	0.4	0.7	1.5	2.6	6.2	10.3
1956	0.9	*	0.5	0.4	0.3	0.3	0.5	0.8	1.3	3.1	6.4	12.3
1955	1.0	*	0.5	0.4	0.4	0.4	0.5	1.0	1.7	3.7	6.9	10.2
1954	1.0	*	0.4	0.5	0.4	0.4	0.6	0.8	1.8	3.2	6.5	12.2
1953	1.0	*	0.5	0.6	0.4	0.6	0.6	1.1	1.6	3.3	5.4	7.5
1952	1.3	*	0.7	0.7	0.8	0.6	0.8	1.4	2.3	3.5	7.2	11.0
1951	1.4	*	0.9	0.8	0.7	0.7	0.8	1.6	2.1	4.3	7.7	10.0
1950	1.5	*	1.2	0.9	0.8	0.7	0.8	1.6	2.7	4.8	5.9	10.0

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
APPENDICITIS (550-553)												
WHITE, BOTH SEXES												
1959	1.0	*	0.3	0.4	0.3	0.2	0.5	0.9	1.8	3.5	6.5	11.3
1958	1.0	*	0.3	0.3	0.3	0.3	0.4	1.1	1.8	4.0	7.6	11.3
1957	1.1	*	0.4	0.4	0.3	0.2	0.5	1.2	2.1	4.3	7.8	11.2
1956	1.2	*	0.4	0.4	0.4	0.3	0.5	1.2	2.0	4.5	8.4	14.3
1955	1.3	*	0.6	0.4	0.4	0.4	0.6	1.5	2.6	5.2	8.4	12.7
1954	1.3	*	0.5	0.5	0.4	0.4	0.7	1.3	2.8	4.7	8.5	14.7
1953	1.5	*	0.6	0.6	0.5	0.6	0.8	1.8	2.8	5.0	7.9	10.3
1952	1.6	*	0.9	0.8	0.7	0.6	0.9	1.9	3.1	5.1	9.2	13.6
1951	1.7	*	1.1	0.9	0.8	0.6	1.2	1.8	3.1	5.6	10.0	12.1
1950	1.9	*	1.3	0.9	0.9	0.8	1.2	2.1	3.8	6.8	8.8	13.5
WHITE, MALE												
1959	1.3	*	0.3	0.5	0.4	0.3	0.7	1.4	2.8	5.0	8.8	16.9
1958	1.3	*	0.3	0.4	0.3	0.3	0.5	1.5	2.5	5.4	10.7	12.3
1957	1.5	*	0.4	0.5	0.3	0.3	0.7	1.8	3.1	6.4	9.9	11.9
1956	1.6	*	0.4	0.5	0.5	0.5	0.8	1.9	3.0	6.3	10.7	15.9
1955	1.7	*	0.7	0.4	0.5	0.5	0.9	2.2	3.6	6.9	10.2	16.3
1954	1.7	*	0.6	0.6	0.4	0.5	0.9	2.0	4.1	6.6	11.1	17.0
1953	2.0	*	0.8	0.7	0.7	0.8	1.2	2.7	4.1	6.9	10.8	13.8
1952	2.1	*	1.0	0.9	0.8	0.7	1.2	2.8	4.1	7.0	11.6	16.6
1951	2.2	*	1.2	0.9	0.9	0.8	1.8	2.5	4.3	7.2	12.4	14.3
1950	2.5	*	1.4	1.0	1.1	1.0	1.9	3.0	5.1	8.9	12.2	17.9
WHITE, FEMALE												
1959	0.7	*	*	0.3	0.2	0.2	0.3	0.5	0.9	2.2	4.8	7.7
1958	0.7	*	*	0.3	0.2	0.2	0.2	0.7	1.1	2.6	5.2	10.7
1957	0.7	*	0.4	0.3	0.2	*	0.2	0.5	1.2	2.5	6.1	10.7
1956	0.8	*	0.4	0.3	0.3	0.2	0.3	0.6	1.1	2.9	6.6	13.1
1955	1.0	*	0.4	0.4	0.3	0.3	0.4	0.8	1.5	3.6	7.0	10.1
1954	0.9	*	0.4	0.5	0.4	0.3	0.5	0.6	1.5	3.0	6.5	13.0
1953	0.9	*	0.5	0.5	0.3	0.4	0.4	1.0	1.5	3.2	5.6	7.8
1952	1.2	*	0.7	0.6	0.7	0.5	0.6	1.1	2.1	3.3	7.3	11.4
1951	1.3	*	0.9	0.8	0.6	0.5	0.6	1.1	2.0	4.2	8.0	10.5
1950	1.4	*	1.2	0.9	0.7	0.5	0.6	1.3	2.6	4.9	6.0	10.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
APPENDICITIS (550-553)												
ALL OTHER, BOTH SEXES												
1959	1.6	*	*	0.8	0.8	*	1.5	2.7	3.8	6.7	9.2	*
1958	1.5	*	*	0.7	*	1.1	1.4	2.1	4.0	5.8	*	*
1957	1.7	*	*	0.9	*	0.9	1.7	2.7	5.8	5.4	*	*
1956	1.8	*	*	0.7	*	1.3	1.4	3.5	5.0	6.5	*	*
1955	1.8	*	*	0.7	*	1.3	1.6	3.5	4.4	6.8	*	*
1954	2.1	*	*	1.0	1.3	1.4	1.9	3.3	5.6	8.2	9.8	*
1953	2.0	*	*	0.9	1.3	1.4	2.5	4.0	4.5	5.9	*	*
1952	2.7	*	*	1.3	1.5	1.6	3.4	5.0	7.0	6.1	11.5	*
1951	2.9	*	*	1.5	1.8	2.2	3.1	5.9	6.4	7.6	*	*
1950	3.0	*	*	1.6	2.0	2.1	3.6	5.6	6.1	8.0	*	*
ALL OTHER, MALE												
1959	1.9	*	*	*	*	*	2.1	3.0	5.0	8.8	*	*
1958	1.8	*	*	1.0	*	*	*	*	4.9	8.8	*	*
1957	2.0	*	*	1.1	*	*	1.7	3.1	6.8	7.9	*	*
1956	2.0	*	*	*	*	*	*	4.4	6.7	7.3	*	*
1955	2.2	*	*	*	*	1.7	*	4.7	5.7	8.8	*	*
1954	2.5	*	*	*	*	1.8	2.2	4.1	6.1	10.8	*	*
1953	2.5	*	*	*	1.9	*	2.7	5.3	6.1	7.3	*	*
1952	3.3	*	*	1.6	*	*	4.3	5.7	9.9	7.2	*	*
1951	3.4	*	*	2.1	2.3	1.9	3.4	5.5	9.1	8.9	*	*
1950	3.6	*	*	1.8	2.2	2.5	4.2	6.1	8.3	12.5	*	*
ALL OTHER, FEMALE												
1959	1.3	*	*	*	*	*	*	2.4	*	4.8	*	*
1958	1.3	*	*	*	*	*	*	2.2	3.2	*	*	*
1957	1.5	*	*	*	*	*	1.7	2.3	4.7	*	*	*
1956	1.5	*	*	*	*	*	1.8	2.6	3.3	5.7	*	*
1955	1.5	*	*	*	*	*	2.0	2.4	*	*	*	*
1954	1.7	*	*	*	*	*	*	2.5	5.0	5.8	*	*
1953	1.6	*	*	*	*	1.6	2.3	2.7	*	*	*	*
1952	2.2	*	*	*	1.5	1.9	2.6	4.3	3.9	*	*	*
1951	2.5	*	*	*	*	2.5	2.8	6.2	*	6.4	*	*
1950	2.4	*	*	1.3	1.9	1.8	3.1	5.1	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)												
BOTH SEXES												
1959	5.2	23.5	0.6	0.2	0.2	0.5	1.2	3.2	8.9	20.6	50.7	111.9
1958	5.1	23.5	0.6	0.2	0.3	0.5	1.4	3.6	8.5	20.2	50.8	101.9
1957	5.0	26.0	0.6	0.2	0.3	0.4	1.5	3.4	7.8	20.2	48.8	107.6
1956	5.1	28.1	0.6	0.2	0.3	0.6	1.3	3.4	8.5	19.6	51.3	97.5
1955	5.3	25.5	0.7	0.1	0.3	0.5	1.6	3.9	9.1	21.1	52.7	104.4
1954	5.3	28.1	0.7	0.2	0.3	0.7	1.6	4.1	8.9	21.3	51.7	96.2
1953	5.5	28.2	0.8	0.2	0.4	0.8	1.8	4.3	9.7	22.4	54.2	97.3
1952	5.4	27.9	1.0	0.2	0.4	0.8	1.9	4.3	9.7	22.7	51.4	102.3
1951	5.6	29.3	0.9	0.3	0.4	0.9	2.1	4.8	10.3	23.6	52.1	97.4
1950	5.9	30.5	1.0	0.3	0.4	1.0	2.1	5.4	10.8	24.4	58.1	111.3
MALE												
1959	5.2	28.3	0.7	0.2	0.3	0.5	1.1	3.1	10.0	22.7	51.6	120.8
1958	5.2	28.6	0.8	0.2	0.3	0.5	1.2	3.7	9.5	22.2	52.9	106.7
1957	5.2	30.6	0.9	0.2	0.3	0.5	1.3	3.3	9.0	22.2	51.1	119.6
1956	5.3	34.0	0.7	0.3	0.3	0.6	1.0	3.4	9.2	21.5	56.3	103.1
1955	5.5	28.8	1.0	0.2	0.4	0.4	1.4	3.8	10.5	23.0	60.4	109.7
1954	5.6	33.6	0.9	0.3	0.4	0.5	1.4	4.2	9.8	24.8	56.3	108.2
1953	5.9	34.3	1.1	0.3	0.3	0.8	1.4	4.2	11.0	25.1	61.2	99.0
1952	5.9	32.7	1.3	0.2	0.4	0.7	1.6	4.5	10.6	26.4	58.0	111.7
1951	6.0	35.1	1.3	0.3	0.5	0.8	1.8	4.6	11.5	27.1	58.3	105.1
1950	6.2	36.4	1.3	0.4	0.5	0.8	2.0	5.1	12.1	26.8	62.8	127.1
FEMALE												
1959	5.1	18.5	0.5	0.1	0.2	0.5	1.4	3.4	7.8	18.8	50.0	106.1
1958	5.0	18.2	0.5	*	0.2	0.6	1.5	3.6	7.6	18.4	49.2	98.6
1957	4.9	21.3	0.3	*	*	0.4	1.7	3.4	6.7	18.4	47.0	99.4
1956	4.9	22.0	0.4	*	0.3	0.7	1.5	3.4	7.8	17.8	47.3	93.6
1955	5.0	22.0	0.5	*	0.2	0.7	1.8	4.0	7.7	19.5	46.5	100.7
1954	4.9	22.4	0.5	*	0.3	0.7	1.7	4.1	8.0	18.2	47.8	87.5
1953	5.2	22.0	0.6	*	0.4	0.8	2.1	4.4	8.4	20.0	48.4	96.0
1952	5.1	23.0	0.7	*	0.4	0.9	2.1	4.0	8.8	19.3	45.9	95.5
1951	5.2	23.2	0.6	0.2	0.3	1.1	2.3	5.0	9.1	20.3	46.9	91.9
1950	5.6	24.4	0.6	0.2	0.4	1.1	2.3	5.7	9.5	22.2	54.0	100.3

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)												
WHITE, BOTH SEXES												
1959	5.1	22.9	0.6	0.1	0.2	0.4	0.9	2.8	8.2	20.3	51.8	115.2
1958	5.0	22.4	0.5	0.1	0.2	0.4	1.1	3.0	7.9	19.3	51.7	106.2
1957	5.0	25.3	0.6	0.1	0.2	0.4	1.2	2.8	7.2	19.8	49.6	111.2
1956	5.0	27.5	0.6	0.2	0.2	0.5	1.0	2.9	7.7	19.1	51.8	101.8
1955	5.2	24.7	0.7	0.1	0.2	0.4	1.2	3.2	8.4	20.7	53.5	110.0
1954	5.2	27.5	0.6	0.2	0.2	0.5	1.2	3.5	8.3	21.1	52.2	99.4
1953	5.4	26.7	0.7	0.2	0.3	0.6	1.4	3.7	9.0	22.0	54.9	99.7
1952	5.3	27.0	0.8	0.2	0.3	0.6	1.5	3.6	9.0	22.6	52.0	106.3
1951	5.5	28.9	0.7	0.2	0.3	0.7	1.7	4.1	9.6	23.2	53.0	100.4
1950	5.7	29.0	0.8	0.3	0.3	0.7	1.7	4.5	9.9	24.1	58.6	117.6
WHITE, MALE												
1959	5.1	27.2	0.6	0.2	0.2	0.4	0.9	2.7	9.3	22.1	52.6	124.8
1958	5.0	26.6	0.6	0.1	0.3	0.4	1.0	3.2	8.8	21.0	53.3	110.1
1957	5.1	28.8	0.9	0.2	0.3	0.4	1.0	2.9	8.3	21.5	51.2	121.9
1956	5.1	32.9	0.7	0.3	*	0.4	0.9	2.8	8.3	20.7	56.8	105.0
1955	5.4	28.0	1.0	0.2	0.3	0.4	1.1	3.1	9.5	22.1	60.6	115.0
1954	5.5	32.5	0.8	0.2	0.3	0.4	1.1	3.4	9.1	24.2	57.0	109.6
1953	5.7	31.9	0.9	0.3	0.3	0.6	1.2	3.7	10.1	24.3	62.4	100.0
1952	5.7	31.2	1.0	0.3	0.4	0.5	1.3	3.8	9.8	26.3	57.9	114.2
1951	5.8	34.6	1.1	0.3	0.4	0.5	1.5	4.0	10.8	26.6	58.4	108.0
1950	5.9	34.4	1.1	0.3	0.3	0.5	1.6	4.2	10.9	26.2	63.0	133.8
WHITE, FEMALE												
1959	5.1	18.5	0.5	*	*	0.4	0.9	2.9	7.2	18.8	51.1	108.9
1958	5.0	18.0	0.4	*	*	0.4	1.2	2.8	6.9	17.7	50.5	103.6
1957	4.9	21.8	*	*	*	0.3	1.4	2.7	6.1	18.3	48.3	103.9
1956	4.9	21.9	0.4	*	0.2	0.6	1.2	2.9	7.2	17.7	47.8	99.5
1955	5.0	21.3	0.4	*	*	0.5	1.3	3.2	7.3	19.4	47.9	106.5
1954	4.9	22.3	0.4	*	*	0.5	1.3	3.6	7.5	18.3	48.3	92.1
1953	5.1	21.3	0.5	*	0.3	0.6	1.7	3.7	7.9	19.9	48.8	99.5
1952	5.0	22.6	0.7	*	0.3	0.6	1.6	3.4	8.2	19.3	47.2	100.6
1951	5.1	22.9	0.4	0.2	0.2	0.9	1.9	4.1	8.5	20.1	48.5	94.9
1950	5.5	23.3	0.5	*	0.3	0.8	1.8	4.7	8.9	22.1	54.9	106.4

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)												
ALL OTHER, BOTH SEXES												
1959	5.4	26.6	*	*	*	1.2	4.0	7.6	16.0	24.6	36.1	72.1
1958	6.0	30.1	1.4	*	0.7	1.5	3.7	9.4	15.7	31.7	38.4	50.0
1957	5.6	29.9	*	*	*	1.2	4.1	8.7	15.0	24.9	38.2	62.5
1956	5.7	31.9	*	*	0.8	1.5	3.6	8.4	16.7	24.7	45.1	45.9
1955	6.0	30.1	1.1	*	*	1.5	4.6	10.8	17.1	26.5	40.6	36.2
1954	6.1	31.7	1.1	*	1.0	2.3	4.7	10.4	16.6	23.6	43.9	54.5
1953	6.7	38.2	1.8	*	1.0	2.4	4.5	10.0	18.3	27.9	44.4	67.3
1952	6.5	34.2	2.3	*	1.0	2.9	5.5	10.3	18.2	23.1	42.5	53.1
1951	6.9	31.9	2.4	*	0.9	2.8	5.5	12.1	19.3	27.9	38.2	61.7
1950	7.6	40.6	2.0	*	1.4	3.3	5.8	14.3	22.0	29.0	49.7	*
ALL OTHER, MALE												
1959	5.8	34.7	*	*	*	*	3.0	6.9	17.5	30.1	38.9	75.9
1958	6.7	40.8	1.8	*	*	*	3.4	8.1	17.0	36.7	47.5	*
1957	6.4	41.5	*	*	*	*	3.3	7.4	16.4	29.8	49.6	92.3
1956	6.5	41.2	*	*	*	1.7	2.5	8.9	19.0	30.7	50.0	80.0
1955	6.8	34.1	*	*	*	*	3.8	10.0	22.0	32.9	57.6	*
1954	7.1	40.7	*	*	*	2.2	4.0	11.4	18.5	31.3	47.9	90.9
1953	7.5	50.4	2.6	*	*	2.2	3.3	9.2	22.2	34.9	45.6	*
1952	7.3	43.2	3.5	*	*	2.4	4.5	10.9	20.1	27.2	59.1	*
1951	7.6	38.6	3.0	*	*	3.1	4.9	10.2	21.3	32.4	56.6	*
1950	8.7	49.9	2.7	*	2.0	3.1	5.4	13.4	26.7	34.8	59.5	*
ALL OTHER, FEMALE												
1959	5.0	18.5	*	*	*	*	4.8	8.2	14.5	19.7	33.5	69.2
1958	5.4	19.3	*	*	*	1.8	4.1	10.8	14.5	27.1	30.1	*
1957	4.9	18.4	*	*	*	*	4.8	10.0	13.6	20.5	27.6	*
1956	4.9	22.6	*	*	*	*	4.5	7.9	14.5	19.0	40.6	*
1955	5.2	26.2	*	*	*	1.9	5.4	11.7	12.2	20.6	24.2	*
1954	5.2	22.8	*	*	*	2.3	5.4	9.5	14.7	16.5	39.7	*
1953	5.9	26.0	*	*	*	2.5	5.5	10.8	14.3	21.4	43.0	*
1952	5.7	25.3	*	*	*	3.3	6.4	9.7	16.3	19.3	26.7	*
1951	6.2	25.2	*	*	*	2.6	6.0	14.1	17.2	23.7	20.7	*
1950	6.6	31.4	*	*	*	3.6	6.1	15.1	17.3	23.1	40.0	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)												
BOTH SEXES												
1959	4.4	62.6	3.4	0.3	0.4	0.6	1.1	2.1	4.9	10.9	26.4	57.7
1958	4.5	66.0	4.1	0.3	0.4	0.7	1.1	2.3	4.8	11.1	24.6	55.0
1957	4.7	77.3	3.4	0.3	0.4	0.6	1.2	2.2	4.9	10.6	24.9	51.7
1956	4.5	73.2	3.8	0.3	0.5	0.7	1.0	2.1	4.3	10.3	24.2	57.5
1955	4.7	86.8	4.4	0.3	0.4	0.6	1.0	2.1	4.6	9.2	24.0	49.4
1954	4.9	98.4	3.9	0.3	0.4	0.8	1.2	2.3	4.3	8.8	22.1	50.3
1953	5.4	119.6	4.9	0.4	0.5	0.7	1.3	2.3	4.1	8.8	21.1	50.2
1952	5.6	128.0	6.1	0.6	0.5	0.6	1.1	2.4	4.2	8.7	21.9	46.8
1951	5.2	119.6	5.5	0.4	0.5	0.8	1.2	2.1	3.8	8.4	20.7	49.4
1950	5.1	119.1	5.3	0.4	0.6	0.7	1.0	2.1	3.9	8.8	20.5	49.4
MALE												
1959	4.5	67.3	3.8	0.4	0.4	0.6	1.2	2.5	5.5	10.9	23.4	50.1
1958	4.7	71.4	4.3	0.3	0.5	0.8	1.2	2.7	5.3	12.0	22.0	53.2
1957	4.9	82.4	3.6	0.3	0.5	0.6	1.3	2.6	5.6	11.4	23.7	44.5
1956	4.8	80.0	4.1	0.3	0.4	0.7	1.1	2.5	5.2	11.6	22.6	56.9
1955	5.0	93.7	4.9	0.3	0.4	0.6	1.1	2.5	5.5	9.9	21.8	41.2
1954	5.2	107.1	4.2	0.4	0.3	0.7	1.2	2.9	5.4	9.9	20.8	40.1
1953	5.7	127.8	5.2	0.4	0.6	0.7	1.3	2.6	4.6	9.3	21.3	44.6
1952	6.1	145.6	6.3	0.6	0.6	0.7	1.1	2.6	5.0	8.5	20.2	46.2
1951	5.6	131.5	5.6	0.5	0.5	0.8	1.3	2.4	4.4	8.4	18.1	43.4
1950	5.4	132.0	5.4	0.4	0.6	0.7	1.0	2.5	4.5	8.7	18.6	47.3
FEMALE												
1959	4.3	57.7	2.9	0.2	0.5	0.6	1.0	1.7	4.3	10.9	28.7	62.6
1958	4.4	60.4	3.9	0.3	0.4	0.7	1.1	1.8	4.3	10.2	26.7	56.2
1957	4.5	71.9	3.2	0.3	0.4	0.6	1.0	1.9	4.1	9.8	25.8	56.7
1956	4.2	66.2	3.4	0.3	0.5	0.6	1.0	1.7	3.6	9.2	25.5	57.9
1955	4.5	79.6	3.9	0.3	0.5	0.5	1.0	1.7	3.9	8.5	25.7	55.2
1954	4.6	89.5	3.7	0.3	0.5	0.9	1.1	1.8	3.2	7.8	23.2	57.6
1953	5.1	111.1	4.5	0.4	0.5	0.8	1.3	1.9	3.7	8.4	20.9	54.2
1952	5.1	109.8	5.9	0.5	0.5	0.6	1.1	2.1	3.5	8.8	23.3	47.2
1951	4.9	107.2	5.3	0.4	0.5	0.8	1.2	1.9	3.1	8.4	22.9	53.6
1950	4.7	105.7	5.2	0.4	0.6	0.7	1.0	1.7	3.4	8.9	22.0	50.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)												
WHITE, BOTH SEXES												
1959	3.8	37.4	2.6	0.3	0.4	0.6	1.1	2.0	4.7	11.0	27.2	59.0
1958	3.9	43.1	3.3	0.2	0.5	0.7	1.1	2.2	4.7	11.0	25.3	56.0
1957	4.0	48.2	2.7	0.3	0.4	0.6	1.1	2.1	4.8	10.5	25.6	53.6
1956	3.9	48.2	3.2	0.3	0.5	0.7	1.0	2.1	4.3	10.3	24.9	59.6
1955	4.0	57.6	3.6	0.2	0.4	0.6	1.0	2.1	4.5	9.2	24.7	51.1
1954	4.1	69.3	2.9	0.3	0.4	0.7	1.2	2.2	4.2	8.9	22.6	51.1
1953	4.5	84.7	3.7	0.3	0.6	0.7	1.2	2.1	3.9	8.9	21.4	52.0
1952	4.6	89.3	4.7	0.5	0.5	0.6	1.0	2.1	4.0	8.4	22.2	49.3
1951	4.5	92.2	4.1	0.4	0.5	0.8	1.2	1.9	3.6	8.4	20.7	50.8
1950	4.4	96.7	4.4	0.3	0.6	0.7	0.9	1.9	3.6	8.6	20.1	50.0
WHITE, MALE												
1959	3.7	40.5	2.8	0.3	0.4	0.6	1.2	2.3	5.4	11.0	24.2	50.6
1958	4.0	46.4	3.5	0.2	0.5	0.7	1.2	2.7	5.4	12.1	22.5	53.2
1957	4.0	50.7	2.9	0.3	0.5	0.5	1.3	2.5	5.7	11.3	24.3	45.0
1956	4.1	52.6	3.5	0.3	0.4	0.7	1.0	2.5	5.1	11.6	23.3	59.3
1955	4.2	62.8	4.1	0.2	0.3	0.7	1.1	2.5	5.3	9.9	22.4	43.2
1954	4.4	76.1	3.1	0.3	0.4	0.6	1.2	2.8	5.2	9.9	21.4	39.4
1953	4.7	92.0	4.1	0.4	0.6	0.6	1.3	2.5	4.5	9.5	21.5	45.9
1952	4.9	101.7	4.9	0.5	0.6	0.7	1.0	2.4	4.8	8.3	20.5	48.2
1951	4.7	101.1	4.1	0.5	0.6	0.7	1.3	2.2	4.2	8.4	17.9	44.3
1950	4.6	107.0	4.4	0.3	0.6	0.6	0.9	2.3	4.1	8.5	18.1	47.2
WHITE, FEMALE												
1959	3.9	34.2	2.5	0.2	0.5	0.6	1.1	1.6	4.0	11.0	29.6	64.5
1958	3.9	39.7	3.2	0.2	0.4	0.7	1.1	1.8	4.0	10.0	27.5	57.9
1957	3.9	45.5	2.6	0.3	0.4	0.6	1.0	1.8	4.0	9.8	26.7	59.5
1956	3.7	43.7	2.9	0.3	0.5	0.6	0.9	1.6	3.5	9.1	26.1	59.9
1955	3.8	52.1	3.0	0.2	0.5	0.5	0.9	1.6	3.6	8.5	26.5	56.6
1954	3.9	62.3	2.7	0.3	0.5	0.8	1.1	1.6	3.3	7.9	23.6	59.5
1953	4.2	77.1	3.4	0.3	0.5	0.7	1.2	1.7	3.4	8.4	21.3	56.3
1952	4.2	76.3	4.4	0.5	0.4	0.6	0.9	1.9	3.3	8.6	23.6	50.0
1951	4.3	82.9	4.0	0.4	0.5	0.8	1.1	1.6	3.0	8.4	23.0	55.4
1950	4.2	85.9	4.4	0.4	0.6	0.7	0.8	1.4	3.1	8.7	21.7	51.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)												
ALL OTHER, BOTH SEXES												
1959	9.3	208.3	7.9	0.6	*	*	1.2	3.1	7.2	10.0	14.4	41.2
1958	9.1	202.9	8.6	0.5	*	1.0	1.4	2.4	6.0	12.0	15.4	42.4
1957	10.6	251.0	7.5	0.6	*	0.9	1.6	3.1	5.4	11.6	14.6	*
1956	9.5	223.6	7.7	0.6	*	*	1.5	2.7	5.1	11.1	14.6	*
1955	10.8	266.5	9.9	*	*	*	1.7	2.6	6.8	9.0	14.5	*
1954	11.3	282.5	10.6	0.6	*	1.3	1.4	3.9	5.3	8.2	15.2	40.0
1953	13.1	346.9	12.2	0.8	*	1.1	1.9	3.5	6.4	8.0	16.7	*
1952	14.6	385.0	15.7	0.9	*	1.0	2.5	4.5	6.7	11.7	17.3	*
1951	11.8	296.6	15.3	*	*	1.0	1.6	4.1	6.2	8.2	21.7	*
1950	10.7	270.3	12.0	0.7	*	1.1	2.1	4.3	7.4	11.7	26.6	*
ALL OTHER, MALE												
1959	10.4	225.7	10.1	*	*	*	*	3.8	6.9	10.6	*	*
1958	10.0	224.6	9.2	*	*	*	*	2.5	4.6	11.6	16.5	*
1957	11.8	276.3	7.9	*	*	*	1.8	2.8	5.2	12.9	17.0	*
1956	10.6	249.3	8.4	*	*	*	*	2.6	5.7	12.1	*	*
1955	11.7	288.9	10.2	*	*	*	*	3.0	6.8	9.1	*	*
1954	12.6	308.1	10.9	*	*	*	*	4.2	7.5	9.9	*	*
1953	14.0	365.8	13.1	*	*	*	2.0	3.2	5.7	7.0	18.4	*
1952	16.5	445.5	16.0	*	*	*	2.0	5.1	6.9	11.6	*	*
1951	13.2	333.2	16.3	*	*	*	*	3.8	7.3	8.3	20.8	*
1950	12.1	304.7	13.0	*	*	*	2.1	4.6	8.7	11.5	25.8	*
ALL OTHER, FEMALE												
1959	8.2	190.4	5.7	*	*	*	*	2.3	7.6	9.4	15.5	*
1958	8.3	181.4	8.0	*	*	*	*	2.4	7.5	12.2	14.4	*
1957	9.5	225.7	7.2	*	*	*	*	3.4	5.5	10.4	*	*
1956	8.4	198.2	7.1	*	*	*	1.9	2.8	4.4	10.1	15.2	*
1955	10.0	245.2	9.7	*	*	*	1.7	2.3	6.9	8.9	*	*
1954	10.1	257.2	10.3	*	*	*	*	3.6	*	6.6	17.5	*
1953	12.2	328.1	11.3	*	*	*	1.8	3.8	7.1	8.9	*	*
1952	12.7	325.3	15.4	*	*	*	3.0	4.0	6.5	11.8	18.1	*
1951	10.5	260.4	14.3	*	*	*	1.9	4.3	5.1	8.1	22.5	*
1950	9.3	236.1	10.9	*	*	*	2.1	4.0	6.2	11.9	27.3	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CIRRHOSIS OF LIVER (581)												
BOTH SEXES												
1959	10.9	0.9	0.2	0.1	0.3	2.8	11.5	26.1	31.8	34.9	33.0	28.6
1958	10.8	1.0	0.1	0.1	0.3	2.5	11.1	24.9	31.3	36.6	34.0	31.2
1957	11.3	1.0	0.2	0.1	0.3	2.9	12.1	25.6	33.0	38.2	35.3	32.3
1956	10.7	1.0	0.2	0.1	0.3	2.4	11.2	23.8	30.9	36.8	37.3	32.7
1955	10.2	1.0	0.2	0.1	0.3	2.1	10.5	22.8	28.8	36.1	35.9	37.0
1954	10.1	1.2	*	0.1	0.3	2.2	10.4	21.5	29.1	35.5	35.7	37.4
1953	10.4	1.2	0.1	0.1	0.3	2.3	10.7	23.0	29.9	34.7	38.8	37.8
1952	10.2	1.3	0.2	0.2	0.4	2.4	10.6	22.4	29.9	34.4	37.6	37.9
1951	9.8	1.3	*	0.1	0.3	2.3	10.1	21.1	28.7	34.0	38.7	40.6
1950	9.2	1.2	*	0.1	0.3	2.1	9.3	19.1	27.2	33.8	36.9	39.0
MALE												
1959	14.6	1.0	*	*	0.2	2.9	14.1	34.7	46.1	53.3	46.2	37.5
1958	14.5	1.0	*	*	0.2	2.6	13.4	33.9	45.1	56.2	46.1	41.0
1957	15.3	1.2	*	0.1	*	3.1	14.7	35.3	48.2	57.4	48.9	42.1
1956	14.2	*	*	*	0.3	2.4	13.6	31.9	44.2	54.6	48.6	42.8
1955	13.6	1.3	*	*	0.2	2.1	12.9	30.9	41.4	52.5	50.1	43.7
1954	13.5	1.1	*	*	0.4	2.2	12.7	29.6	42.1	51.5	47.5	47.4
1953	13.8	1.2	*	0.2	0.2	2.5	12.9	30.9	43.1	49.9	50.9	45.0
1952	13.8	*	*	0.2	0.2	2.4	13.4	30.1	43.6	50.4	50.2	49.5
1951	13.0	1.4	*	*	0.2	2.2	12.6	28.2	39.6	48.6	52.8	42.2
1950	12.1	1.4	*	*	0.2	2.1	11.7	25.6	38.1	48.0	45.9	43.9
FEMALE												
1959	7.3	*	*	*	0.3	2.7	9.1	17.7	18.5	18.8	22.6	22.8
1958	7.1	1.1	*	*	0.4	2.3	8.9	16.0	18.2	19.4	24.5	24.6
1957	7.5	*	*	0.1	0.4	2.8	9.7	16.2	18.6	21.2	24.4	25.5
1956	7.3	1.2	*	0.1	0.4	2.4	8.8	15.9	18.2	20.8	28.2	25.7
1955	6.9	*	*	0.1	0.3	2.1	8.2	14.8	16.6	21.4	24.3	32.3
1954	6.7	1.3	*	0.2	0.3	2.1	8.2	13.6	16.5	21.0	26.1	30.4
1953	7.1	*	*	*	0.4	2.1	8.7	15.3	17.1	20.8	28.9	32.6
1952	6.8	1.6	*	*	0.5	2.3	8.0	14.7	16.4	19.7	27.2	29.7
1951	6.8	1.2	*	*	0.3	2.3	7.7	14.1	17.8	20.4	26.9	39.4
1950	6.3	*	*	*	0.5	2.0	6.9	12.6	16.1	20.6	29.2	35.6

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CIRRHOSIS OF LIVER (581)												
WHITE, BOTH SEXES												
1959	11.1	0.9	0.2	0.1	0.2	2.0	10.7	25.9	32.2	36.0	34.3	30.1
1958	11.0	1.0	*	0.1	0.2	1.9	10.5	24.9	31.9	37.8	35.3	31.9
1957	11.6	0.9	*	0.1	0.2	2.4	11.5	25.6	33.1	39.6	36.1	33.9
1956	11.0	0.8	0.2	0.1	0.3	1.9	10.6	23.9	31.1	38.2	38.6	34.5
1955	10.5	0.9	*	0.1	0.2	1.8	10.2	22.9	29.4	37.4	36.8	38.5
1954	10.4	1.2	*	0.1	0.3	1.9	10.1	21.8	29.6	36.8	37.0	39.1
1953	10.7	0.9	*	0.1	0.3	2.0	10.7	23.1	30.6	36.0	40.4	39.3
1952	10.6	1.3	0.2	0.1	0.3	2.1	10.5	22.6	30.5	35.8	38.9	38.7
1951	10.2	1.2	*	0.2	0.3	2.1	10.0	21.2	29.1	35.1	39.9	42.2
1950	9.5	1.1	*	0.1	0.3	1.9	9.3	19.3	27.6	34.4	37.8	41.2
WHITE, MALE												
1959	14.9	*	*	*	0.2	2.2	13.3	34.7	47.0	54.9	48.3	38.7
1958	14.9	*	*	*	*	2.1	13.0	34.0	46.2	57.9	48.0	40.8
1957	15.8	*	*	*	*	2.6	14.1	35.7	48.7	59.6	50.0	44.1
1956	14.7	*	*	*	0.3	2.0	13.2	32.4	44.7	56.7	50.5	44.0
1955	14.2	*	*	*	0.2	1.8	12.7	31.4	42.4	54.3	51.7	44.6
1954	14.1	*	*	*	0.3	1.9	12.6	30.4	43.0	53.5	49.0	48.9
1953	14.3	*	*	*	*	2.2	13.0	31.2	44.2	51.7	53.5	47.0
1952	14.3	*	*	*	*	2.2	13.3	30.6	44.5	52.3	51.9	50.2
1951	13.5	1.4	*	*	0.2	2.1	12.6	28.5	40.2	50.2	54.9	43.5
1950	12.6	*	*	*	*	2.0	11.9	26.3	38.8	48.9	47.2	47.2
WHITE, FEMALE												
1959	7.3	*	*	*	0.2	1.8	8.2	17.2	18.3	19.4	23.4	24.6
1958	7.2	*	*	*	0.3	1.6	8.1	16.0	18.5	20.2	25.4	25.9
1957	7.5	*	*	*	0.3	2.2	9.0	15.9	18.3	22.0	25.3	26.9
1956	7.4	*	*	*	0.3	1.8	8.1	15.7	18.2	21.7	29.2	27.9
1955	7.0	*	*	*	0.3	1.7	7.7	14.5	16.9	22.2	25.0	34.2
1954	6.8	1.4	*	0.2	0.3	1.8	7.8	13.3	16.6	21.7	27.2	32.1
1953	7.2	*	*	*	0.3	1.8	8.4	15.1	17.4	21.7	29.8	33.7
1952	7.0	1.6	*	*	0.4	2.0	7.8	14.7	16.8	20.6	28.3	30.4
1951	6.9	*	*	*	0.3	2.1	7.4	13.9	18.0	21.1	27.5	41.3
1950	6.5	*	*	*	0.5	1.8	6.7	12.4	16.4	21.0	29.9	36.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CIRRHOSIS OF LIVER (581)												
ALL OTHER, BOTH SEXES												
1959	9.7	*	*	*	0.8	9.2	18.9	28.1	28.6	22.4	15.4	*
1958	8.5	*	*	*	0.9	7.2	16.0	24.7	24.2	21.9	16.1	*
1957	9.4	*	*	*	0.8	7.2	17.5	25.1	31.9	21.4	22.9	*
1956	8.5	*	*	*	*	6.4	16.3	22.6	28.3	19.6	18.3	*
1955	7.5	*	*	*	*	5.0	13.0	22.0	21.4	20.4	22.3	*
1954	7.1	*	*	*	*	4.6	12.6	19.0	23.9	19.2	17.6	*
1953	7.2	*	*	*	*	4.4	11.3	22.2	21.8	19.0	15.4	*
1952	7.2	*	*	*	0.8	4.6	11.6	20.4	23.0	17.9	18.6	*
1951	7.1	*	*	*	*	3.8	11.3	20.3	24.0	20.5	20.7	*
1950	6.5	*	*	*	*	3.2	9.6	17.0	21.6	25.8	23.1	*
ALL OTHER, MALE												
1959	11.7	*	*	*	*	8.9	21.4	34.5	37.2	34.3	19.4	*
1958	10.7	*	*	*	*	6.6	17.1	33.1	33.7	35.4	21.6	*
1957	11.6	*	*	*	*	7.4	19.6	31.5	42.5	31.4	35.6	*
1956	10.2	*	*	*	*	5.7	17.8	27.0	38.1	29.9	23.8	*
1955	9.0	*	*	*	*	4.4	14.1	26.0	29.3	30.4	29.6	*
1954	8.6	*	*	*	*	5.0	13.5	21.6	32.1	27.6	26.9	*
1953	8.9	*	*	*	*	4.7	11.6	27.7	30.2	28.8	*	*
1952	9.4	*	*	*	*	5.0	13.8	25.8	33.8	27.2	27.3	*
1951	8.7	*	*	*	*	3.4	12.7	24.8	32.3	29.7	24.5	*
1950	7.8	*	*	*	*	2.7	10.0	19.4	30.2	36.9	28.7	*
ALL OTHER, FEMALE												
1959	7.9	*	*	*	*	9.4	16.5	22.1	20.2	11.7	*	*
1958	6.5	*	*	*	*	7.8	15.0	16.8	15.1	9.6	*	*
1957	7.4	*	*	*	*	7.0	15.5	19.0	21.5	12.3	*	*
1956	6.9	*	*	*	*	7.0	14.9	18.3	18.6	10.1	*	*
1955	6.0	*	*	*	*	5.5	12.0	18.1	13.6	11.2	15.2	*
1954	5.7	*	*	*	*	4.3	11.7	16.5	15.8	11.5	*	*
1953	5.7	*	*	*	*	4.2	11.1	16.7	13.3	9.8	*	*
1952	5.1	*	*	*	*	4.3	9.5	15.0	11.8	9.2	*	*
1951	5.6	*	*	*	*	4.1	10.1	15.9	15.4	11.8	*	*
1950	5.2	*	*	*	*	3.6	9.2	14.6	12.9	14.7	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)												
BOTH SEXES												
1959	2.5	*	*	*	*	0.2	0.6	1.7	4.9	12.7	29.3	55.8
1958	2.7	*	*	*	*	0.3	0.7	1.8	5.2	13.4	33.0	56.6
1957	2.9	*	*	*	0.1	0.3	0.7	2.1	5.6	14.9	33.3	58.5
1956	3.1	*	*	*	*	0.3	0.8	2.2	6.5	16.3	35.8	65.2
1955	3.6	*	*	*	0.1	0.4	1.0	2.7	8.4	19.0	38.3	62.2
1954	3.5	*	*	*	0.1	0.4	1.0	3.0	8.1	18.7	37.6	59.7
1953	3.5	*	*	*	*	0.4	0.9	3.2	8.9	19.0	35.5	54.7
1952	3.8	*	*	*	*	0.4	1.2	3.7	9.4	20.6	37.7	57.5
1951	3.7	*	*	*	*	0.4	1.2	3.8	9.0	20.0	38.6	58.3
1950	3.9	*	*	*	*	0.4	1.2	4.1	10.2	21.8	38.9	57.9
MALE												
1959	2.1	*	*	*	*	*	0.4	1.5	4.4	11.6	26.4	48.5
1958	2.2	*	*	*	*	*	0.5	1.5	4.6	11.7	32.7	43.0
1957	2.4	*	*	*	*	*	0.5	1.8	4.9	14.2	28.8	46.6
1956	2.5	*	*	*	*	*	0.5	1.8	5.7	14.2	31.4	48.0
1955	2.8	*	*	*	*	0.3	0.7	2.3	6.8	16.4	32.4	46.2
1954	2.8	*	*	*	*	0.2	0.7	2.3	6.3	16.5	32.3	43.4
1953	2.7	*	*	*	*	*	0.7	2.4	7.3	15.4	30.9	41.9
1952	2.8	*	*	*	*	*	0.8	2.8	7.1	15.8	30.0	42.9
1951	2.8	*	*	*	*	0.3	0.9	3.0	6.8	15.4	31.3	42.6
1950	2.8	*	*	*	*	0.2	0.8	3.0	7.5	16.2	32.3	46.9
FEMALE												
1959	3.0	*	*	*	*	0.3	0.7	1.9	5.3	13.7	31.6	60.6
1958	3.2	*	*	*	*	0.4	0.8	2.0	5.8	14.9	33.2	65.7
1957	3.4	*	*	*	*	0.5	0.9	2.4	6.2	15.6	36.8	66.6
1956	3.8	*	*	*	*	0.4	1.1	2.5	7.3	18.1	39.4	77.2
1955	4.4	*	*	*	0.2	0.6	1.2	3.2	9.9	21.4	43.1	73.5
1954	4.3	*	*	*	*	0.5	1.2	3.6	9.9	20.8	41.8	71.3
1953	4.3	*	*	*	*	0.6	1.2	4.1	10.5	22.3	39.3	63.9
1952	4.8	*	*	*	*	0.6	1.5	4.5	11.8	24.9	44.1	68.0
1951	4.7	*	*	*	*	0.5	1.5	4.7	11.3	24.2	44.7	69.4
1950	4.9	*	*	*	*	0.6	1.6	5.2	12.9	26.9	44.5	65.6

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)												
WHITE, BOTH SEXES												
1959	2.7	*	*	*	*	0.2	0.5	1.6	5.0	13.1	30.3	59.3
1958	2.9	*	*	*	*	0.3	0.6	1.7	5.3	13.8	34.5	60.2
1957	3.1	*	*	*	0.1	0.3	0.7	2.1	5.7	15.6	34.8	61.5
1956	3.4	*	*	*	*	0.3	0.8	2.2	6.7	16.9	37.5	68.9
1955	3.9	*	*	*	0.1	0.4	0.9	2.8	8.6	20.0	40.4	65.6
1954	3.8	*	*	*	*	0.4	0.9	2.9	8.4	19.6	39.5	62.8
1953	3.8	*	*	*	*	0.4	0.9	3.4	9.3	19.9	36.9	56.8
1952	4.1	*	*	*	*	0.4	1.2	3.8	9.8	21.5	39.6	61.0
1951	4.0	*	*	*	*	0.4	1.2	4.0	9.3	21.1	40.4	62.0
1950	4.2	*	*	*	*	0.4	1.2	4.2	10.6	22.8	40.8	61.8
WHITE, MALE												
1959	2.3	*	*	*	*	*	0.4	1.5	4.6	12.0	27.8	51.2
1958	2.4	*	*	*	*	*	0.5	1.5	4.8	12.1	34.4	45.3
1957	2.6	*	*	*	*	*	0.5	1.9	5.1	15.0	30.2	48.6
1956	2.7	*	*	*	*	*	0.5	1.9	5.9	14.8	32.9	50.3
1955	3.1	*	*	*	*	0.2	0.7	2.4	7.1	17.2	34.5	48.6
1954	3.0	*	*	*	*	0.2	0.7	2.3	6.6	17.2	34.4	46.1
1953	3.0	*	*	*	*	*	0.7	2.5	7.6	16.3	32.5	42.5
1952	3.0	*	*	*	*	*	0.9	2.9	7.4	16.5	31.6	45.8
1951	3.0	*	*	*	*	0.3	0.9	3.2	7.0	16.3	32.9	45.1
1950	3.1	*	*	*	*	*	0.8	3.1	7.8	17.3	33.9	50.4
WHITE, FEMALE												
1959	3.1	*	*	*	*	0.3	0.6	1.8	5.3	14.0	32.3	64.5
1958	3.4	*	*	*	*	0.4	0.8	2.0	5.8	15.3	34.5	70.1
1957	3.6	*	*	*	*	0.5	0.9	2.3	6.2	16.1	38.4	70.2
1956	4.0	*	*	*	*	0.4	1.1	2.5	7.4	18.7	41.2	81.8
1955	4.7	*	*	*	0.2	0.6	1.1	3.1	10.1	22.4	45.2	77.6
1954	4.6	*	*	*	*	0.5	1.2	3.5	10.2	21.7	43.5	74.8
1953	4.6	*	*	*	*	0.6	1.2	4.2	10.9	23.3	40.6	67.1
1952	5.1	*	*	*	*	0.6	1.5	4.6	12.0	26.1	46.1	71.9
1951	5.0	*	*	*	*	0.5	1.4	4.9	11.5	25.6	46.7	74.1
1950	5.3	*	*	*	*	0.6	1.6	5.3	13.4	27.9	46.7	69.8

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)												
ALL OTHER, BOTH SEXES												
1959	1.3	*	*	*	*	*	0.9	2.5	3.9	8.2	15.4	*
1958	1.3	*	*	*	*	*	0.9	2.3	4.3	8.6	12.0	*
1957	1.3	*	*	*	*	*	0.8	2.2	4.6	7.3	12.5	*
1956	1.3	*	*	*	*	*	0.9	1.8	4.7	8.8	11.9	*
1955	1.4	*	*	*	*	*	1.1	2.4	5.8	7.4	8.2	*
1954	1.4	*	*	*	*	*	1.0	3.2	4.4	8.9	9.8	*
1953	1.3	*	*	*	*	*	*	2.1	4.5	8.3	15.0	*
1952	1.4	*	*	*	*	*	0.9	2.9	5.6	8.8	9.7	*
1951	1.3	*	*	*	*	*	1.3	2.0	5.6	6.8	11.5	*
1950	1.3	*	*	*	*	*	1.1	2.9	5.2	7.5	9.8	*
ALL OTHER, MALE												
1959	0.9	*	*	*	*	*	*	*	*	6.3	*	*
1958	1.0	*	*	*	*	*	*	2.2	*	7.0	*	*
1957	0.9	*	*	*	*	*	*	*	*	*	*	*
1956	0.9	*	*	*	*	*	*	*	*	6.7	*	*
1955	0.9	*	*	*	*	*	*	*	3.6	6.4	*	*
1954	0.8	*	*	*	*	*	*	*	*	7.1	*	*
1953	0.8	*	*	*	*	*	*	*	*	*	*	*
1952	0.9	*	*	*	*	*	*	*	*	6.9	*	*
1951	1.0	*	*	*	*	*	*	*	*	*	*	*
1950	0.8	*	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	1.7	*	*	*	*	*	*	3.5	5.3	9.8	21.1	*
1958	1.6	*	*	*	*	*	*	2.4	5.9	10.1	13.7	*
1957	1.7	*	*	*	*	*	*	2.8	6.0	9.4	13.8	*
1956	1.7	*	*	*	*	*	*	2.2	6.4	10.6	*	*
1955	1.8	*	*	*	*	*	*	3.4	7.9	8.4	*	*
1954	1.9	*	*	*	*	*	*	4.9	5.7	10.5	*	*
1953	1.7	*	*	*	*	*	*	3.1	6.0	11.1	19.8	*
1952	1.9	*	*	*	*	*	*	3.9	8.6	10.6	*	*
1951	1.6	*	*	*	*	*	*	3.0	8.0	7.8	*	*
1950	1.9	*	*	*	*	*	*	4.2	7.3	13.0	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
NEPHRITIS AND NEPHROSIS (590-594)												
BOTH SEXES												
1959	8.0	2.0	0.6	0.9	1.9	3.3	5.1	8.6	14.6	27.2	64.6	151.9
1958	9.2	2.0	0.9	0.9	2.3	3.5	5.5	9.8	16.3	33.2	76.8	186.2
1957	10.2	1.8	1.1	1.1	2.7	4.1	6.3	10.2	18.8	36.6	82.0	198.7
1956	10.6	2.1	1.0	1.1	2.4	3.9	6.2	10.7	19.1	38.8	91.7	215.1
1955	11.1	1.9	1.3	1.1	2.7	3.9	6.3	11.3	20.2	41.1	99.5	217.2
1954	12.2	2.5	1.5	1.2	2.8	4.4	6.5	12.5	22.6	45.3	111.7	235.9
1953	13.6	2.2	1.7	1.4	3.2	4.8	7.5	14.2	25.0	51.4	124.2	279.9
1952	15.3	2.7	1.9	1.7	3.2	4.9	8.5	16.5	29.0	57.9	145.7	309.0
1951	16.8	2.7	2.2	1.8	3.4	5.0	8.8	16.8	32.0	65.5	164.4	383.4
1950	18.7	2.7	2.5	1.7	3.5	5.5	9.9	18.8	34.9	77.0	189.3	442.2
MALE												
1959	8.9	2.5	0.8	0.8	2.2	4.1	6.0	10.5	17.0	31.6	76.6	180.6
1958	10.2	2.2	1.0	1.0	2.7	3.9	6.5	11.1	18.9	39.1	90.5	220.6
1957	11.1	1.8	1.2	1.1	3.2	5.0	7.0	11.8	21.2	43.0	92.6	226.7
1956	11.4	2.0	1.2	1.1	2.8	4.7	6.9	12.0	21.6	43.6	104.7	256.9
1955	12.1	2.5	1.7	1.1	3.3	4.5	7.3	12.8	22.2	47.0	117.2	243.1
1954	13.3	3.1	1.8	1.3	3.3	4.9	7.3	13.8	25.0	52.3	130.7	272.7
1953	14.8	2.3	2.0	1.5	3.7	5.2	8.2	15.8	27.6	58.3	146.4	323.9
1952	16.8	3.3	2.3	1.8	3.7	5.5	9.1	18.8	32.3	66.3	171.0	352.0
1951	18.0	2.9	2.4	1.8	3.7	5.5	9.6	18.9	35.2	74.2	187.3	421.9
1950	20.1	3.3	2.8	1.8	4.1	6.0	10.6	20.8	38.7	85.3	215.9	508.0
FEMALE												
1959	7.1	1.5	0.5	1.0	1.6	2.6	4.2	6.7	12.4	23.3	55.2	133.1
1958	8.3	1.9	0.8	0.8	1.9	3.2	4.5	8.5	13.9	28.1	66.0	163.2
1957	9.3	1.8	1.0	1.1	2.2	3.2	5.6	8.7	16.5	31.0	73.6	179.6
1956	9.7	2.1	0.9	1.1	2.0	3.1	5.5	9.4	16.8	34.4	81.3	186.0
1955	10.1	1.3	0.9	1.1	2.2	3.2	5.3	9.8	18.4	35.8	85.2	198.9
1954	11.1	1.8	1.3	1.1	2.3	3.9	5.7	11.3	20.2	38.9	96.2	209.6
1953	12.5	2.1	1.4	1.3	2.9	4.4	6.9	12.6	22.4	45.2	105.9	248.3
1952	13.9	2.0	1.5	1.6	2.8	4.3	8.0	14.1	25.8	50.2	124.6	278.2
1951	15.6	2.6	2.1	1.8	3.2	4.6	8.2	14.8	28.9	57.4	145.0	356.1
1950	17.3	2.1	2.1	1.5	2.8	5.1	9.3	16.7	31.0	69.3	166.7	396.4

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
NEPHRITIS AND NEPHROSIS (590-594)												
WHITE, BOTH SEXES												
1959	7.0	1.8	0.5	0.8	1.8	2.8	3.9	6.6	12.0	22.9	59.9	144.0
1958	8.2	1.9	0.7	0.8	2.1	3.0	4.2	7.5	12.8	29.0	71.8	177.1
1957	9.0	1.5	0.9	1.1	2.4	3.3	4.7	8.0	15.4	31.8	76.2	188.9
1956	9.3	1.8	0.9	1.0	2.2	3.2	4.7	8.0	15.6	33.6	85.8	206.4
1955	9.9	1.7	1.1	1.1	2.6	3.2	4.8	8.6	16.5	36.3	93.4	209.2
1954	10.9	2.2	1.4	1.2	2.6	3.7	5.1	9.8	18.7	39.6	105.1	224.6
1953	12.1	1.9	1.5	1.4	3.0	4.0	5.7	11.0	20.6	45.0	115.5	268.2
1952	13.5	2.6	1.8	1.6	2.9	4.0	6.4	12.5	23.2	50.6	134.7	289.1
1951	14.9	2.3	2.0	1.7	3.1	4.1	6.7	12.9	26.5	56.6	153.6	362.7
1950	16.6	1.9	2.3	1.6	3.2	4.6	7.4	14.5	28.7	66.6	176.7	428.1
WHITE, MALE												
1959	7.9	2.3	0.6	0.7	2.1	3.6	4.8	8.5	14.4	26.5	71.4	167.8
1958	9.1	1.9	0.8	0.9	2.5	3.5	5.4	8.9	15.3	33.9	85.2	206.0
1957	9.9	1.6	1.0	1.1	3.0	4.3	5.7	9.9	17.6	38.1	85.5	210.9
1956	10.2	1.7	1.0	1.0	2.6	4.2	5.8	9.6	17.9	37.6	97.7	241.7
1955	11.0	2.1	1.4	1.1	3.2	4.1	5.9	10.5	18.7	41.6	109.9	236.7
1954	11.9	2.9	1.6	1.2	3.1	4.3	6.0	11.2	21.3	45.2	120.3	254.6
1953	13.4	2.1	1.8	1.5	3.7	4.5	6.8	13.0	23.2	50.7	135.5	309.0
1952	15.0	3.0	2.1	1.8	3.5	4.8	7.3	15.1	26.6	57.8	156.1	330.0
1951	16.1	2.3	2.1	1.7	3.6	4.8	7.8	15.3	29.9	64.3	173.3	390.3
1950	18.1	2.4	2.5	1.8	4.0	5.3	8.8	17.1	32.5	73.3	200.2	484.5
WHITE, FEMALE												
1959	6.2	1.2	0.4	0.9	1.4	2.1	2.9	4.7	9.7	19.7	51.1	128.6
1958	7.3	1.9	0.6	0.8	1.7	2.4	3.0	6.1	10.4	24.6	61.4	157.9
1957	8.1	1.5	0.8	1.0	1.9	2.4	3.8	6.1	13.3	26.2	68.8	174.0
1956	8.5	1.9	0.8	1.0	1.8	2.2	3.7	6.5	13.5	30.0	76.4	181.8
1955	8.8	1.3	0.8	1.1	2.0	2.4	3.7	6.9	14.3	31.6	80.4	189.6
1954	9.9	1.5	1.3	1.1	2.1	3.1	4.2	8.3	16.3	34.4	92.8	203.1
1953	10.9	1.7	1.2	1.3	2.3	3.4	4.7	9.0	18.1	39.8	99.3	238.8
1952	12.0	2.1	1.4	1.5	2.3	3.3	5.6	9.9	20.0	44.0	117.1	259.7
1951	13.6	2.3	1.8	1.7	2.7	3.3	5.6	10.4	23.0	49.6	137.2	343.1
1950	15.2	*	2.1	1.5	2.5	3.8	6.1	12.0	24.8	60.3	156.9	388.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
NEPHRITIS AND NEPHROSIS (590-594)												
ALL OTHER, BOTH SEXES												
1959	15.8	3.3	1.5	1.5	3.0	7.1	15.8	27.5	42.8	78.6	128.5	248.5
1958	17.9	*	2.3	1.3	3.2	8.0	16.9	31.4	54.0	84.7	145.2	295.5
1957	19.7	*	2.3	1.5	4.1	10.0	19.7	32.0	55.8	94.5	164.6	312.5
1956	20.6	3.7	1.9	1.7	3.8	9.5	19.2	36.2	58.8	101.3	175.7	319.7
1955	20.8	*	2.4	1.4	3.8	9.0	19.3	36.5	63.9	98.1	186.3	315.5
1954	22.7	4.2	2.2	1.5	4.5	10.2	18.8	38.9	67.9	113.4	208.6	370.9
1953	26.3	*	3.4	1.6	5.3	11.6	23.2	44.8	78.3	128.8	250.9	423.1
1952	31.1	*	3.0	2.0	5.7	12.0	26.7	54.2	101.0	145.5	306.6	555.1
1951	33.3	5.7	4.3	2.5	5.8	13.2	27.9	54.6	103.2	171.0	324.0	634.0
1950	36.3	8.1	3.6	2.0	5.1	13.8	31.7	59.0	109.6	219.1	380.1	609.6
ALL OTHER, MALE												
1959	17.0	*	2.1	1.2	2.8	7.7	16.7	29.9	44.4	91.4	142.4	324.1
1958	19.1	*	2.3	1.7	3.8	6.9	16.5	32.8	56.5	100.0	158.3	400.0
1957	20.3	*	2.6	1.3	4.3	11.0	19.0	29.8	59.7	100.8	183.7	415.4
1956	21.7	*	2.7	2.2	4.3	9.0	17.2	34.9	62.2	115.1	195.4	440.0
1955	21.4	*	3.2	1.3	3.7	8.5	19.3	35.2	61.4	111.0	212.8	320.8
1954	24.8	*	2.7	1.6	5.1	10.3	18.5	38.2	68.5	135.8	268.9	504.5
1953	27.2	*	3.9	1.5	3.7	11.1	20.9	42.7	80.2	148.3	293.9	514.3
1952	32.8	*	3.7	2.1	5.1	11.0	24.8	54.3	101.7	166.9	374.5	630.0
1951	34.2	*	4.6	2.5	4.5	11.4	25.6	52.3	101.2	190.8	382.1	815.8
1950	37.8	*	4.8	1.8	5.2	12.1	27.3	56.7	112.3	241.1	435.1	782.1
ALL OTHER, FEMALE												
1959	14.7	*	*	1.9	3.2	6.5	15.0	25.3	41.3	67.0	116.1	192.3
1958	16.8	*	2.3	0.9	2.6	9.0	17.2	30.2	51.7	70.6	133.3	228.9
1957	19.1	*	2.0	1.8	4.0	9.1	20.4	34.1	52.1	88.7	146.9	248.6
1956	19.6	*	*	1.1	3.3	10.0	21.0	37.3	55.4	88.1	157.2	236.1
1955	20.3	*	*	1.4	3.9	9.4	19.2	37.7	66.5	86.3	159.8	311.8
1954	20.6	*	*	1.3	3.9	10.1	19.1	39.5	67.4	92.7	150.0	290.6
1953	25.4	*	2.9	1.8	6.7	12.0	25.2	46.8	76.2	110.3	208.3	361.3
1952	29.4	*	2.2	1.9	6.2	12.9	28.5	54.2	100.2	125.5	242.2	503.4
1951	32.4	*	3.9	2.5	6.9	14.8	30.1	56.9	105.3	152.3	268.5	510.7
1950	34.9	*	*	2.1	4.9	15.2	35.9	61.4	106.8	197.0	326.0	486.3

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)												
BOTH SEXES												
1959	1.0	1.2	0.4	0.4	0.4	0.4	0.8	1.1	1.7	2.8	6.1	11.8
1958	1.3	1.3	0.7	0.4	0.4	0.6	0.8	1.4	1.9	3.8	8.3	17.8
1957	1.3	1.3	0.8	0.4	0.5	0.6	0.8	1.3	2.2	4.1	8.5	17.7
1956	1.4	1.2	0.6	0.4	0.5	0.6	1.0	1.5	2.4	4.6	9.1	18.3
1955	1.5	1.3	0.9	0.4	0.5	0.7	1.0	1.7	2.5	4.3	9.0	16.3
1954	1.6	1.6	0.9	0.5	0.6	0.7	1.1	1.7	2.5	4.6	10.1	21.3
1953	1.8	1.6	1.1	0.6	0.8	0.9	1.3	1.9	3.0	5.0	11.1	19.4
1952	2.0	1.8	1.2	0.8	0.9	1.0	1.5	2.2	3.4	5.7	11.6	22.5
1951	2.1	2.0	1.4	0.8	0.9	1.0	1.5	2.1	3.6	6.4	13.3	27.1
1950	2.3	2.0	1.5	0.7	0.9	1.0	1.7	2.4	4.0	7.2	15.0	31.2
MALE												
1959	1.2	1.5	0.6	0.3	0.5	0.5	1.0	1.4	2.1	3.4	7.4	13.8
1958	1.4	1.4	0.7	0.5	0.5	0.6	0.8	1.6	2.2	4.7	9.8	21.2
1957	1.5	1.3	0.9	0.4	0.6	0.7	0.9	1.5	2.8	5.1	10.5	19.3
1956	1.6	1.0	0.7	0.5	0.6	0.6	1.2	1.7	2.7	5.4	10.0	20.8
1955	1.7	1.5	1.1	0.5	0.6	0.8	1.1	2.1	2.7	5.2	11.7	15.4
1954	1.8	2.2	1.2	0.5	0.8	0.8	1.2	2.0	2.8	5.2	11.3	27.0
1953	2.0	1.5	1.4	0.6	0.9	0.8	1.4	2.2	3.7	5.8	13.5	20.4
1952	2.2	2.4	1.4	0.8	1.0	1.0	1.7	2.7	3.8	6.7	13.1	22.3
1951	2.3	1.8	1.4	0.7	1.0	0.9	1.8	2.4	3.7	7.6	15.1	31.3
1950	2.6	2.2	1.7	0.8	1.2	1.2	1.9	3.2	4.5	7.6	16.5	40.5
FEMALE												
1959	0.9	*	0.3	0.4	0.4	0.4	0.7	0.8	1.3	2.3	5.2	10.5
1958	1.1	1.2	0.6	0.4	0.3	0.5	0.9	1.2	1.7	2.9	7.1	15.5
1957	1.2	1.3	0.7	0.4	0.4	0.5	0.8	1.1	1.7	3.2	6.8	16.6
1956	1.3	1.4	0.5	0.4	0.4	0.5	0.9	1.4	2.0	3.9	8.4	16.6
1955	1.3	1.2	0.6	0.4	0.5	0.6	0.9	1.3	2.3	3.6	6.8	16.9
1954	1.4	*	0.7	0.5	0.5	0.7	0.9	1.3	2.2	4.0	9.2	17.2
1953	1.5	1.7	0.8	0.5	0.7	0.9	1.1	1.7	2.3	4.3	9.1	18.7
1952	1.8	1.3	1.0	0.7	0.9	1.0	1.3	1.8	3.0	4.8	10.4	22.6
1951	2.0	2.2	1.4	0.9	0.8	1.0	1.3	1.9	3.4	5.3	11.8	24.2
1950	2.0	1.7	1.2	0.6	0.6	0.9	1.5	1.7	3.4	6.9	13.8	24.7

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)												
WHITE, BOTH SEXES												
1959	0.9	1.1	0.3	0.3	0.4	0.4	0.6	0.9	1.4	2.6	5.8	10.8
1958	1.1	1.2	0.5	0.4	0.3	0.4	0.6	1.1	1.7	3.4	8.1	16.8
1957	1.2	1.1	0.7	0.4	0.4	0.4	0.6	1.1	1.9	3.5	8.0	17.1
1956	1.2	1.1	0.5	0.4	0.4	0.4	0.8	1.2	2.0	4.0	8.4	17.3
1955	1.2	1.1	0.7	0.4	0.5	0.5	0.7	1.2	2.0	3.7	8.6	15.9
1954	1.4	1.5	0.9	0.4	0.5	0.6	0.8	1.3	2.1	4.1	9.8	19.4
1953	1.5	1.3	0.9	0.5	0.7	0.6	0.9	1.5	2.5	4.4	10.7	17.2
1952	1.7	1.7	1.1	0.7	0.7	0.7	1.1	1.8	2.7	5.1	11.1	21.7
1951	1.8	1.6	1.1	0.7	0.7	0.7	1.1	1.6	2.9	5.3	12.4	25.5
1950	2.0	1.6	1.4	0.6	0.8	0.8	1.1	1.8	3.3	6.3	14.2	30.8
WHITE, MALE												
1959	1.0	1.4	0.4	0.3	0.5	0.5	0.7	1.1	1.8	3.1	7.0	12.3
1958	1.2	1.2	0.6	0.4	0.5	0.4	0.6	1.2	1.9	4.2	9.7	20.3
1957	1.3	1.1	0.8	0.4	0.6	0.5	0.7	1.3	2.4	4.5	9.7	19.3
1956	1.4	*	0.6	0.4	0.5	0.5	0.9	1.4	2.4	4.6	9.4	18.9
1955	1.4	*	1.0	0.5	0.5	0.6	0.9	1.7	2.3	4.4	11.0	15.6
1954	1.6	2.2	1.1	0.5	0.7	0.6	1.0	1.6	2.4	4.8	10.9	24.5
1953	1.7	*	1.1	0.6	0.9	0.6	1.0	1.8	3.1	5.0	12.8	17.2
1952	1.9	2.1	1.2	0.8	0.9	0.8	1.3	2.2	3.2	5.8	12.4	21.7
1951	1.9	1.4	1.2	0.6	0.9	0.7	1.4	1.9	3.2	6.4	14.0	27.8
1950	2.3	1.8	1.5	0.8	1.1	1.0	1.4	2.4	3.9	6.6	15.5	39.9
WHITE, FEMALE												
1959	0.7	*	*	0.3	0.2	0.3	0.4	0.6	1.1	2.2	4.8	9.9
1958	1.0	1.2	0.4	0.3	0.2	0.4	0.5	0.9	1.4	2.7	6.9	14.4
1957	1.0	*	0.6	0.3	0.3	0.3	0.5	0.9	1.4	2.6	6.7	15.5
1956	1.1	1.2	0.5	0.4	0.4	0.3	0.6	1.0	1.6	3.5	7.6	16.1
1955	1.0	*	0.5	0.4	0.4	0.4	0.6	0.8	1.7	3.1	6.7	16.1
1954	1.2	*	0.7	0.4	0.4	0.5	0.6	1.0	1.8	3.5	8.9	15.8
1953	1.3	1.3	0.6	0.4	0.4	0.6	0.7	1.2	1.9	3.8	8.9	17.3
1952	1.5	*	0.9	0.6	0.6	0.7	1.0	1.3	2.2	4.4	10.0	21.6
1951	1.6	1.9	1.1	0.8	0.6	0.6	0.9	1.3	2.7	4.3	11.0	23.8
1950	1.7	*	1.2	0.5	0.5	0.6	0.9	1.3	2.7	6.0	13.1	24.5

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)												
ALL OTHER, BOTH SEXES												
1959	2.2	*	1.3	0.8	1.0	1.1	2.9	3.3	4.4	5.8	11.1	*
1958	2.6	*	1.8	0.8	1.0	1.7	3.1	4.4	4.6	8.0	11.0	30.3
1957	2.8	*	1.8	0.7	1.1	2.2	2.7	3.8	6.0	11.5	14.6	*
1956	3.0	*	1.3	0.8	1.0	2.0	3.4	4.4	6.4	11.5	19.0	*
1955	3.2	*	1.7	0.6	1.0	2.2	3.1	6.3	8.0	11.9	14.5	*
1954	3.0	*	1.4	0.6	1.4	2.1	3.2	4.9	6.8	10.5	15.2	43.6
1953	4.0	*	2.4	0.9	1.8	2.7	4.6	6.2	9.3	13.1	17.9	46.2
1952	4.4	*	2.2	1.0	2.3	3.1	4.8	6.8	11.7	13.3	19.5	*
1951	5.2	*	3.2	1.5	2.2	3.5	5.1	7.3	11.8	19.8	27.6	46.8
1950	5.2	*	2.4	1.2	1.9	3.3	6.6	7.9	12.1	20.1	27.0	*
ALL OTHER, MALE												
1959	2.4	*	1.9	*	*	*	3.3	4.0	5.0	7.1	*	*
1958	2.7	*	*	*	*	1.8	2.4	5.1	4.9	11.1	*	*
1957	3.0	*	*	*	*	2.2	2.8	4.1	7.3	12.4	20.7	*
1956	3.3	*	*	1.2	*	2.1	3.4	4.1	5.9	15.6	18.5	*
1955	3.5	*	2.3	*	*	2.1	3.2	6.6	7.3	14.6	20.8	*
1954	3.3	*	*	*	*	2.4	3.2	5.6	7.7	10.2	17.6	*
1953	4.2	*	2.8	*	*	2.5	4.8	6.0	10.6	15.7	22.8	*
1952	4.5	*	2.7	*	*	2.7	4.9	7.2	10.9	17.3	23.6	*
1951	5.1	*	3.1	1.4	*	3.0	5.6	7.3	10.8	21.4	30.2	*
1950	5.7	*	3.4	*	2.0	3.4	6.5	10.7	12.0	21.2	29.7	*
ALL OTHER, FEMALE												
1959	2.0	*	*	1.0	*	*	2.5	2.7	3.8	4.6	*	*
1958	2.4	*	2.0	*	*	1.6	3.8	3.7	4.4	5.2	*	*
1957	2.7	*	*	*	1.4	2.2	2.7	3.6	4.7	10.6	*	*
1956	2.7	*	*	*	*	1.9	3.4	4.8	6.9	7.7	19.6	*
1955	3.0	*	*	*	*	2.2	3.0	5.9	8.8	9.4	*	*
1954	2.7	*	*	*	*	1.8	3.2	4.2	5.9	10.8	*	*
1953	3.8	*	*	*	2.3	2.9	4.4	6.5	8.1	10.6	*	*
1952	4.2	*	*	*	3.1	3.5	4.6	6.4	12.6	9.5	*	*
1951	5.2	*	3.3	1.6	2.7	4.0	4.7	7.3	12.7	18.2	25.2	*
1950	4.6	*	*	*	1.9	3.3	6.7	5.1	12.3	18.9	24.4	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFECTIONS OF KIDNEY (600)												
BOTH SEXES												
1959	3.9	1.5	0.1	0.1	0.4	0.8	1.6	3.5	7.6	16.7	37.9	71.9
1958	4.0	1.1	0.1	0.1	0.4	0.9	1.7	3.6	8.2	16.9	38.9	78.4
1957	3.7	1.3	*	0.1	0.4	0.8	1.6	3.5	7.7	15.6	35.0	79.4
1956	3.4	1.4	*	0.1	0.3	0.8	1.4	3.2	6.9	14.7	33.8	73.3
1955	3.0	1.1	0.1	0.1	0.3	0.7	1.4	2.9	5.6	12.8	29.9	60.2
1954	2.7	1.1	0.2	0.1	0.3	0.7	1.3	2.6	5.6	11.1	26.1	48.7
1953	2.6	1.3	*	*	0.3	0.6	1.3	2.8	4.9	10.7	25.1	51.5
1952	2.4	1.2	*	0.1	0.4	0.7	1.2	2.6	4.9	9.7	22.9	42.5
1951	2.1	1.1	*	0.1	0.3	0.6	1.1	2.3	4.5	9.0	20.0	39.9
1950	2.1	1.2	*	0.1	0.3	0.5	1.0	2.1	4.5	8.7	22.1	39.7
MALE												
1959	3.9	2.0	*	*	0.3	0.8	1.4	3.0	7.6	19.0	45.5	92.1
1958	4.1	1.7	*	*	0.3	0.7	1.4	3.3	8.6	20.2	47.2	101.7
1957	3.8	1.8	*	*	0.5	0.6	1.2	3.3	8.4	18.2	41.9	92.3
1956	3.6	1.6	*	*	0.3	0.6	1.1	2.9	7.2	17.7	41.9	89.0
1955	3.1	1.3	*	*	*	0.7	1.3	2.6	5.8	15.0	35.3	73.6
1954	2.7	1.2	*	*	0.2	0.6	1.2	2.3	6.1	13.1	30.3	50.7
1953	2.7	1.5	*	*	0.3	0.5	1.2	2.6	5.2	12.5	32.0	61.9
1952	2.5	1.4	*	*	0.4	0.7	1.0	2.6	5.3	11.0	27.5	52.0
1951	2.3	*	*	*	0.2	0.5	0.9	2.2	4.9	10.9	23.6	54.3
1950	2.1	1.6	*	*	0.4	0.3	0.8	2.0	4.7	9.9	26.5	45.2
FEMALE												
1959	3.8	*	*	0.2	0.5	0.8	1.8	4.0	7.5	14.6	32.0	58.7
1958	3.8	*	*	*	0.5	1.0	1.9	3.9	7.8	14.0	32.4	62.8
1957	3.6	*	*	0.1	0.4	1.0	1.9	3.8	7.0	13.3	29.5	70.6
1956	3.3	1.1	*	0.2	0.4	0.9	1.7	3.5	6.6	12.0	27.3	62.3
1955	2.9	*	*	*	0.4	0.7	1.5	3.2	5.5	10.8	25.5	50.8
1954	2.6	*	*	*	0.4	0.7	1.5	2.9	5.1	9.3	22.7	47.3
1953	2.4	*	*	*	0.4	0.7	1.4	3.1	4.7	9.0	19.5	44.0
1952	2.3	*	*	0.2	0.4	0.7	1.4	2.5	4.5	8.4	19.0	35.7
1951	2.0	*	*	*	0.3	0.7	1.2	2.5	4.0	7.3	16.9	29.7
1950	2.0	*	*	*	0.3	0.6	1.2	2.2	4.2	7.6	18.4	35.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFECTIONS OF KIDNEY (600)												
WHITE, BOTH SEXES												
1959	3.5	1.2	*	0.1	0.3	0.7	1.1	2.5	5.9	15.0	37.4	72.4
1958	3.6	0.8	*	0.1	0.3	0.7	1.2	2.7	6.7	15.5	38.7	78.7
1957	3.4	0.8	*	0.1	0.4	0.7	1.2	2.5	6.2	14.2	34.4	79.6
1956	3.2	1.2	*	0.1	0.3	0.6	1.0	2.4	5.6	13.5	33.6	75.7
1955	2.8	1.0	*	0.1	0.3	0.5	1.0	2.1	4.8	12.0	29.9	61.2
1954	2.5	1.0	0.2	0.1	0.2	0.5	1.0	2.0	4.8	10.6	26.0	49.8
1953	2.3	0.9	*	*	0.3	0.5	0.9	2.2	4.2	9.8	24.6	52.0
1952	2.2	1.1	*	0.1	0.4	0.6	0.8	2.0	4.2	9.1	22.7	44.0
1951	2.0	1.1	*	0.1	0.2	0.5	0.8	1.9	4.0	8.5	19.8	40.9
1950	2.0	1.1	*	0.1	0.3	0.4	0.8	1.7	3.9	8.3	21.6	41.5
WHITE, MALE												
1959	3.5	1.5	*	*	0.2	0.7	1.0	2.2	6.0	16.7	44.2	91.4
1958	3.8	*	*	*	0.2	0.6	1.0	2.4	7.0	18.3	46.7	101.3
1957	3.5	*	*	*	0.4	0.5	1.0	2.3	6.7	16.6	41.2	92.6
1956	3.3	1.4	*	*	*	0.5	0.9	2.2	5.9	16.1	41.1	90.4
1955	2.9	*	*	*	*	0.6	1.0	2.1	5.0	14.0	35.4	73.5
1954	2.5	*	*	*	*	0.5	0.9	1.7	5.0	12.3	30.2	50.7
1953	2.5	*	*	*	0.3	0.3	0.8	2.1	4.4	11.4	30.9	59.3
1952	2.3	*	*	*	0.4	0.6	0.7	2.0	4.4	10.5	26.9	53.0
1951	2.1	*	*	*	*	0.4	0.6	1.8	4.3	10.4	23.2	54.0
1950	2.0	1.6	*	*	0.3	0.3	0.6	1.7	4.1	9.5	25.3	47.7
WHITE, FEMALE												
1959	3.5	*	*	0.2	0.4	0.6	1.2	2.7	5.8	13.6	32.1	60.1
1958	3.5	*	*	*	0.4	0.8	1.3	2.9	6.4	13.0	32.5	63.8
1957	3.3	*	*	0.2	0.4	0.8	1.3	2.7	5.7	12.1	29.1	70.7
1956	3.1	*	*	0.2	0.4	0.8	1.2	2.5	5.4	11.2	27.7	65.4
1955	2.7	*	*	*	0.4	0.5	1.0	2.2	4.7	10.3	25.5	52.5
1954	2.4	*	*	*	0.3	0.5	1.0	2.2	4.5	9.1	22.7	49.1
1953	2.2	*	*	*	0.3	0.6	1.0	2.2	4.0	8.3	19.5	46.6
1952	2.1	*	*	0.2	0.3	0.6	0.9	2.0	4.0	7.9	19.3	37.5
1951	1.9	*	*	*	0.3	0.6	0.9	1.9	3.6	6.8	16.9	31.6
1950	1.9	*	*	*	0.3	0.6	1.0	1.7	3.7	7.1	18.5	37.3

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFECTIONS OF KIDNEY (600)												
ALL OTHER, BOTH SEXES												
1959	6.8	3.3	*	*	0.7	2.1	6.0	13.6	25.1	36.7	45.6	66.2
1958	6.6	*	*	*	*	2.5	6.3	12.6	24.4	34.4	42.5	74.2
1957	6.4	4.2	*	*	*	2.1	5.0	13.7	24.1	32.1	42.9	76.6
1956	5.5	*	*	*	*	1.9	4.8	10.8	21.3	29.2	36.6	44.3
1955	4.7	*	*	*	*	2.4	5.2	10.2	15.1	21.3	29.7	48.3
1954	4.2	*	*	*	*	1.8	4.5	9.2	15.0	17.2	27.5	*
1953	4.4	*	*	*	*	1.8	4.4	9.1	13.9	21.2	32.9	46.2
1952	3.7	*	*	*	*	1.6	4.1	7.8	13.1	16.2	25.2	*
1951	3.2	*	*	*	*	1.2	3.7	6.8	11.1	15.0	23.5	*
1950	3.1	*	*	*	*	1.0	3.0	6.2	11.2	14.7	30.0	*
ALL OTHER, MALE												
1959	7.0	*	*	*	*	1.7	4.6	11.0	24.5	46.5	62.5	100.0
1958	7.0	*	*	*	*	2.2	5.2	11.9	25.4	42.9	54.0	111.1
1957	6.7	7.0	*	*	*	1.7	3.0	13.3	26.5	37.2	51.1	88.5
1956	5.8	*	*	*	*	*	3.5	9.0	22.3	36.7	52.3	*
1955	4.6	*	*	*	*	1.8	4.2	7.8	15.8	26.8	33.6	*
1954	4.4	*	*	*	*	*	3.4	8.1	18.1	22.7	31.9	*
1953	4.8	*	*	*	*	*	4.3	7.4	15.2	25.3	46.5	95.2
1952	4.0	*	*	*	*	*	3.0	7.7	15.9	17.9	36.4	*
1951	3.4	*	*	*	*	*	3.5	5.6	12.6	17.4	29.2	*
1950	3.1	*	*	*	*	*	2.6	5.1	12.2	15.0	43.6	*
ALL OTHER, FEMALE												
1959	6.7	*	*	*	*	2.4	7.2	16.1	25.7	27.9	30.4	*
1958	6.2	*	*	*	*	2.8	7.3	13.2	23.5	26.6	32.0	*
1957	6.2	*	*	*	*	2.5	6.8	14.1	21.8	27.5	35.2	70.3
1956	5.2	*	*	*	*	2.2	6.0	12.5	20.3	22.2	21.7	*
1955	4.8	*	*	*	*	2.8	6.2	12.5	14.4	16.3	25.8	*
1954	4.0	*	*	*	*	2.4	5.5	10.3	11.8	12.1	23.0	*
1953	4.0	*	*	*	*	2.1	4.5	10.8	12.6	17.3	19.8	*
1952	3.5	*	*	*	*	1.9	5.0	8.0	10.2	14.6	*	*
1951	3.1	*	*	*	*	1.8	3.8	8.0	9.4	12.7	18.0	*
1950	3.0	*	*	*	*	*	3.2	7.3	10.1	14.4	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HYPERPLASIA OF PROSTATE (610)												
BOTH SEXES												
1959	2.6	*	*	*	*	*	*	0.2	2.4	12.1	42.2	106.5
1958	2.7	*	*	*	*	*	*	0.2	2.5	12.2	42.9	119.5
1957	3.0	*	*	*	*	*	*	0.3	3.0	14.7	48.2	124.3
1956	3.2	*	*	*	*	*	*	0.4	3.3	15.9	52.9	130.5
1955	3.7	*	*	*	*	*	*	0.4	3.6	18.8	64.1	141.9
1954	3.8	*	*	*	*	*	*	0.5	3.9	19.9	65.2	146.4
1953	3.9	*	*	*	*	*	*	0.5	4.0	20.1	70.3	157.2
1952	4.0	*	*	*	*	*	*	0.6	4.0	21.9	71.7	170.2
1951	4.2	*	*	*	*	*	*	0.5	4.4	22.2	77.5	182.8
1950	4.2	*	*	*	*	*	*	0.4	4.6	23.1	79.3	194.3
MALE												
1959	5.3	*	*	*	*	*	*	0.4	4.9	25.9	96.1	269.3
1958	5.4	*	*	*	*	*	*	0.5	5.2	26.1	97.4	298.8
1957	6.1	*	*	*	*	*	*	0.7	6.1	31.2	108.8	306.5
1956	6.5	*	*	*	*	*	*	0.7	6.8	33.8	118.9	318.0
1955	7.5	*	*	*	*	*	*	0.8	7.3	39.7	143.6	342.1
1954	7.7	*	*	*	*	*	*	0.9	7.9	41.9	145.4	351.0
1953	7.9	*	*	*	*	*	*	0.9	8.0	42.1	156.1	375.8
1952	8.2	*	*	*	*	*	*	1.1	8.0	45.8	158.2	407.7
1951	8.5	*	*	*	*	*	*	0.9	8.8	46.3	169.9	439.8
1950	8.5	*	*	*	*	*	*	0.8	9.2	48.0	172.4	473.3
FEMALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HYPERPLASIA OF PROSTATE (610)												
WHITE, BOTH SEXES												
1959	2.6	*	*	*	*	*	*	0.1	1.9	10.9	41.2	108.0
1958	2.6	*	*	*	*	*	*	0.1	2.0	11.1	41.6	122.0
1957	2.9	*	*	*	*	*	*	0.2	2.3	13.1	47.1	126.2
1956	3.2	*	*	*	*	*	*	0.2	2.6	14.8	52.3	131.5
1955	3.7	*	*	*	*	*	*	0.3	3.0	17.5	64.2	144.6
1954	3.8	*	*	*	*	*	*	0.3	3.3	19.0	65.1	149.6
1953	3.9	*	*	*	*	*	*	0.3	3.2	18.6	69.8	162.0
1952	4.0	*	*	*	*	*	*	0.3	3.2	20.7	71.2	174.7
1951	4.2	*	*	*	*	*	*	0.3	3.7	20.8	76.1	188.4
1950	4.2	*	*	*	*	*	*	0.2	3.8	21.4	78.1	200.5
WHITE, MALE												
1959	5.2	*	*	*	*	*	*	0.3	3.8	23.4	94.4	274.8
1958	5.3	*	*	*	*	*	*	0.3	4.2	23.7	94.9	306.6
1957	5.9	*	*	*	*	*	*	0.4	4.7	27.8	107.1	311.6
1956	6.4	*	*	*	*	*	*	0.4	5.4	31.5	118.4	320.5
1955	7.5	*	*	*	*	*	*	0.5	6.1	36.9	144.6	348.6
1954	7.7	*	*	*	*	*	*	0.6	6.6	39.9	146.0	358.2
1953	7.8	*	*	*	*	*	*	0.6	6.4	38.9	155.7	386.2
1952	8.2	*	*	*	*	*	*	0.6	6.5	43.4	157.9	417.8
1951	8.4	*	*	*	*	*	*	0.6	7.3	43.5	167.9	452.3
1950	8.4	*	*	*	*	*	*	0.5	7.6	44.4	170.8	489.1
WHITE, FEMALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HYPERPLASIA OF PROSTATE (610)												
ALL OTHER, BOTH SEXES												
1959	2.9	*	*	*	*	*	*	*	7.8	26.5	55.7	88.2
1958	3.0	*	*	*	*	*	*	1.3	7.6	25.9	61.6	89.4
1957	3.6	*	*	*	*	*	*	1.8	10.3	34.1	63.2	100.0
1956	3.4	*	*	*	*	*	*	1.7	10.7	29.2	60.8	118.0
1955	3.6	*	*	*	*	*	*	1.5	10.4	34.6	63.7	108.6
1954	3.6	*	*	*	*	*	*	2.1	11.4	31.4	66.8	103.6
1953	4.1	*	*	*	*	*	*	2.2	13.6	38.6	78.2	98.1
1952	4.1	*	*	*	*	*	*	3.0	13.8	35.8	78.8	114.3
1951	4.3	*	*	*	*	*	*	2.0	13.7	38.5	97.2	114.9
1950	4.5	*	*	*	*	*	*	2.1	14.7	47.1	96.9	120.6
ALL OTHER, MALE												
1959	6.0	*	*	*	*	*	*	*	16.0	55.8	118.1	206.9
1958	6.2	*	*	*	*	*	*	2.6	15.4	54.3	129.5	218.5
1957	7.5	*	*	*	*	*	*	3.8	20.9	71.5	131.1	246.2
1956	7.1	*	*	*	*	*	*	3.4	21.6	60.9	125.4	288.0
1955	7.4	*	*	*	*	*	*	3.0	21.0	72.1	130.4	262.5
1954	7.4	*	*	*	*	*	*	4.3	22.8	65.3	137.0	259.1
1953	8.5	*	*	*	*	*	*	4.3	27.0	79.9	160.5	242.9
1952	8.5	*	*	*	*	*	*	6.1	27.2	74.0	161.8	280.0
1951	8.9	*	*	*	*	*	*	4.1	26.8	79.2	199.1	284.2
1950	9.3	*	*	*	*	*	*	4.3	29.0	93.9	195.2	289.3
ALL OTHER, FEMALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)												
BOTH SEXES												
1959	0.9	*	1.9	2.9	1.9	*	*	*	*	*
1958	0.9	*	1.7	3.1	1.9	*	*	*	*	*
1957	1.0	*	1.9	3.3	2.1	0.1	*	*	*	*
1956	1.0	*	2.0	3.1	2.1	*	*	*	*	*
1955	1.2	*	2.3	3.5	2.4	0.2	*	*	*	*
1954	1.3	*	2.4	3.9	2.8	0.2	*	*	*	*
1953	1.5	*	3.1	4.2	3.0	0.2	*	*	*	*
1952	1.7	*	3.5	4.9	3.0	0.2	*	*	*	*
1951	1.8	*	3.7	5.2	3.3	0.2	*	*	*	*
1950	2.0	*	3.7	5.4	3.7	0.2	*
MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
FEMALE												
1959	1.8	*	3.7	5.8	3.7	*	*	*	*	*
1958	1.8	*	3.4	6.1	3.7	*	*	*	*	*
1957	2.0	*	3.8	6.5	4.2	0.3	*	*	*	*
1956	2.0	*	4.0	6.1	4.2	*	*	*	*	*
1955	2.3	*	4.5	6.8	4.8	0.3	*	*	*	*
1954	2.6	*	4.7	7.6	5.5	0.4	*	*	*	*
1953	3.0	*	6.0	8.2	6.0	0.4	*	*	*	*
1952	3.3	*	6.7	9.4	5.9	0.3	*	*	*	*
1951	3.6	*	7.3	10.2	6.5	0.3	*	*	*	*
1950	3.9	*	7.4	10.6	7.4	0.3	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)												
WHITE, BOTH SEXES												
1959	0.6	*	1.2	1.9	1.3	*	*	*	*	*
1958	0.6	*	1.1	2.0	1.3	*	*	*	*	*
1957	0.7	*	1.2	2.1	1.4	*	*	*	*	*
1956	0.7	*	1.3	2.0	1.5	*	*	*	*	*
1955	0.8	*	1.5	2.3	1.7	0.1	*	*	*	*
1954	0.9	*	1.6	2.5	2.1	0.1	*	*	*	*
1953	1.0	*	2.0	2.9	2.3	*	*	*	*	*
1952	1.2	*	2.2	3.5	2.2	0.2	*	*	*	*
1951	1.3	*	2.5	3.7	2.4	0.1	*	*	*	*
1950	1.4	*	2.5	3.8	2.8	*	*
WHITE, MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
WHITE, FEMALE												
1959	1.2	*	2.3	3.8	2.6	*	*	*	*	*
1958	1.2	*	2.3	4.0	2.6	*	*	*	*	*
1957	1.3	*	2.4	4.1	2.8	*	*	*	*	*
1956	1.3	*	2.6	4.0	2.9	*	*	*	*	*
1955	1.5	*	2.9	4.5	3.3	0.2	*	*	*	*
1954	1.8	*	3.2	4.8	4.1	0.3	*	*	*	*
1953	2.1	*	3.8	5.7	4.5	*	*	*	*	*
1952	2.3	*	4.3	6.8	4.3	0.3	*	*	*	*
1951	2.6	*	5.0	7.3	4.8	0.3	*	*	*	*
1950	2.8	*	5.0	7.5	5.6	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)												
ALL OTHER, BOTH SEXES												
1959	3.4	*	7.1	10.8	6.9	*	*	*	*	*
1958	3.4	*	6.0	11.5	7.1	*	*	*	*	*
1957	4.0	*	7.3	13.2	8.0	*	*	*	*	*
1956	3.8	*	7.0	11.8	7.6	*	*	*	*	*
1955	4.3	*	7.9	13.3	8.9	*	*	*	*	*
1954	4.8	*	8.0	15.7	8.7	*	*	*	*	*
1953	5.4	*	11.3	15.2	9.4	*	*	*	*	*
1952	6.0	*	12.7	16.5	10.3	*	*	*	*	*
1951	6.4	*	12.5	18.1	11.1	*	*	*	*	*
1950	6.9	*	12.9	19.3	11.5	*	*
ALL OTHER, MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
ALL OTHER, FEMALE												
1959	6.5	*	13.6	20.1	13.1	*	*	*	*	*
1958	6.5	*	11.6	21.5	13.5	*	*	*	*	*
1957	7.8	*	13.9	24.6	15.4	*	*	*	*	*
1956	7.3	*	13.3	22.1	14.7	*	*	*	*	*
1955	8.4	*	14.9	24.8	17.1	*	*	*	*	*
1954	9.3	*	14.9	29.4	16.8	*	*	*	*	*
1953	10.4	*	20.9	28.3	18.2	*	*	*	*	*
1952	11.6	*	23.7	31.1	19.9	*	*	*	*	*
1951	12.5	*	23.6	34.1	21.4	*	*	*	*	*
1950	13.5	*	24.6	36.5	22.1	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ABORTION (650-652)												
BOTH SEXES												
1959	0.2	*	0.4	0.6	0.2	*	*	*	*	*
1958	0.1	*	0.3	0.6	0.2	*	*	*	*	*
1957	0.2	*	0.3	0.5	0.3	*	*	*	*	*
1956	0.1	*	0.3	0.5	0.2	*	*	*	*	*
1955	0.2	*	0.4	0.5	0.3	*	*	*	*	*
1954	0.2	*	0.4	0.6	0.3	*	*	*	*	*
1953	0.2	*	0.5	0.6	0.2	*	*	*	*	*
1952	0.2	*	0.4	0.6	0.3	*	*	*	*	*
1951	0.2	*	0.5	0.5	0.3	*	*	*	*	*
1950	0.2	*	0.5	0.6	0.3	*	*
MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
FEMALE												
1959	0.3	*	0.7	1.1	0.5	*	*	*	*	*
1958	0.3	*	0.6	1.2	0.4	*	*	*	*	*
1957	0.3	*	0.7	1.1	0.5	*	*	*	*	*
1956	0.3	*	0.5	0.9	0.4	*	*	*	*	*
1955	0.3	*	0.7	1.0	0.6	*	*	*	*	*
1954	0.4	*	0.8	1.1	0.6	*	*	*	*	*
1953	0.4	*	0.9	1.2	0.4	*	*	*	*	*
1952	0.4	*	0.8	1.2	0.7	*	*	*	*	*
1951	0.4	*	0.9	1.1	0.6	*	*	*	*	*
1950	0.4	*	1.0	1.1	0.7	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ABORTION (650-652)												
WHITE, BOTH SEXES												
1959	0.1	*	0.2	0.3	0.1	*	*	*	*	*
1958	0.1	*	0.2	0.3	0.2	*	*	*	*	*
1957	0.1	*	0.2	0.3	0.1	*	*	*	*	*
1956	0.1	*	0.2	0.4	0.1	*	*	*	*	*
1955	0.1	*	0.2	0.3	0.2	*	*	*	*	*
1954	0.1	*	0.2	0.3	0.2	*	*	*	*	*
1953	0.1	*	0.3	0.4	0.1	*	*	*	*	*
1952	0.1	*	0.2	0.5	0.3	*	*	*	*	*
1951	0.1	*	0.3	0.3	0.2	*	*	*	*	*
1950	0.1	*	0.3	0.4	0.3	*	*
WHITE, MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
WHITE, FEMALE												
1959	0.2	*	0.4	0.7	0.3	*	*	*	*	*
1958	0.2	*	0.3	0.7	0.3	*	*	*	*	*
1957	0.2	*	0.3	0.6	0.3	*	*	*	*	*
1956	0.2	*	0.3	0.7	0.3	*	*	*	*	*
1955	0.2	*	0.4	0.6	0.5	*	*	*	*	*
1954	0.2	*	0.5	0.6	0.4	*	*	*	*	*
1953	0.2	*	0.6	0.7	0.2	*	*	*	*	*
1952	0.3	*	0.5	0.9	0.5	*	*	*	*	*
1951	0.2	*	0.5	0.7	0.4	*	*	*	*	*
1950	0.3	*	0.7	0.7	0.5	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ABORTION (650-652)												
ALL OTHER, BOTH SEXES												
1959	0.7	*	1.8	2.6	1.0	*	*	*	*	*
1958	0.6	*	1.3	2.7	*	*	*	*	*	*
1957	0.7	*	1.6	2.3	1.2	*	*	*	*	*
1956	0.5	*	1.0	1.5	*	*	*	*	*	*
1955	0.7	*	1.4	2.2	1.0	*	*	*	*	*
1954	0.8	*	1.5	2.5	1.1	*	*	*	*	*
1953	0.8	*	1.6	2.5	1.0	*	*	*	*	*
1952	0.8	*	1.8	2.1	1.1	*	*	*	*	*
1951	0.8	*	1.9	2.3	1.2	*	*	*	*	*
1950	0.8	*	1.7	2.3	*	*	*
ALL OTHER, MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
ALL OTHER, FEMALE												
1959	1.4	*	3.5	4.8	1.9	*	*	*	*	*
1958	1.2	*	2.6	5.0	*	*	*	*	*	*
1957	1.4	*	3.1	4.3	2.4	*	*	*	*	*
1956	0.9	*	1.8	2.8	*	*	*	*	*	*
1955	1.3	*	2.7	4.1	1.9	*	*	*	*	*
1954	1.5	*	2.9	4.6	2.2	*	*	*	*	*
1953	1.5	*	3.0	4.7	2.0	*	*	*	*	*
1952	1.5	*	3.3	3.9	2.2	*	*	*	*	*
1951	1.6	*	3.6	4.3	2.3	*	*	*	*	*
1950	1.5	*	3.2	4.3	*	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)												
BOTH SEXES												
1959	0.7	*	1.5	2.4	1.7	*	*	*	*	*
1958	0.8	*	1.4	2.5	1.7	*	*	*	*	*
1957	0.9	*	1.6	2.8	1.9	0.1	*	*	*	*
1956	0.9	*	1.8	2.6	1.9	*	*	*	*	*
1955	1.0	*	1.9	3.0	2.1	0.1	*	*	*	*
1954	1.1	*	2.0	3.3	2.5	0.2	*	*	*	*
1953	1.3	*	2.6	3.6	2.8	0.2	*	*	*	*
1952	1.5	*	3.1	4.2	2.7	0.2	*	*	*	*
1951	1.6	*	3.3	4.7	3.0	0.1	*	*	*	*
1950	1.8	*	3.2	4.9	3.4	0.2	*
MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
FEMALE												
1959	1.5	*	3.0	4.6	3.3	*	*	*	*	*
1958	1.5	*	2.8	4.9	3.4	*	*	*	*	*
1957	1.7	*	3.2	5.4	3.7	0.2	*	*	*	*
1956	1.7	*	3.5	5.1	3.8	*	*	*	*	*
1955	2.0	*	3.8	5.8	4.1	0.3	*	*	*	*
1954	2.2	*	3.9	6.5	4.9	0.3	*	*	*	*
1953	2.6	*	5.1	7.1	5.5	0.4	*	*	*	*
1952	2.9	*	5.9	8.2	5.2	0.3	*	*	*	*
1951	3.2	*	6.4	9.1	5.9	0.3	*	*	*	*
1950	3.5	*	6.4	9.5	6.7	0.3	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)												
WHITE, BOTH SEXES												
1959	0.5	*	1.0	1.6	1.2	*	*	*	*	*
1958	0.5	*	1.0	1.7	1.2	*	*	*	*	*
1957	0.6	*	1.0	1.8	1.3	*	*	*	*	*
1956	0.6	*	1.2	1.7	1.4	*	*	*	*	*
1955	0.7	*	1.3	2.0	1.4	*	*	*	*	*
1954	0.8	*	1.4	2.1	1.9	0.1	*	*	*	*
1953	0.9	*	1.7	2.6	2.2	*	*	*	*	*
1952	1.0	*	2.0	3.0	1.9	0.2	*	*	*	*
1951	1.2	*	2.3	3.4	2.2	0.1	*	*	*	*
1950	1.2	*	2.2	3.5	2.6	*	*
WHITE, MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
WHITE, FEMALE												
1959	1.0	*	1.9	3.2	2.3	*	*	*	*	*
1958	1.0	*	1.9	3.4	2.3	*	*	*	*	*
1957	1.1	*	2.1	3.5	2.6	*	*	*	*	*
1956	1.2	*	2.3	3.3	2.7	*	*	*	*	*
1955	1.3	*	2.5	3.9	2.8	*	*	*	*	*
1954	1.5	*	2.7	4.2	3.7	0.3	*	*	*	*
1953	1.8	*	3.2	5.0	4.3	*	*	*	*	*
1952	2.0	*	3.9	5.9	3.8	0.3	*	*	*	*
1951	2.3	*	4.5	6.6	4.4	0.3	*	*	*	*
1950	2.5	*	4.3	6.8	5.1	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)												
ALL OTHER, BOTH SEXES												
1959	2.6	*	5.3	8.2	5.9	*	*	*	*	*
1958	2.7	*	4.7	8.8	6.4	*	*	*	*	*
1957	3.3	*	5.7	10.9	6.8	*	*	*	*	*
1956	3.3	*	6.0	10.3	6.9	*	*	*	*	*
1955	3.7	*	6.4	11.1	7.9	*	*	*	*	*
1954	4.0	*	6.5	13.3	7.6	*	*	*	*	*
1953	4.6	*	9.6	12.6	8.4	*	*	*	*	*
1952	5.2	*	11.0	14.4	9.2	*	*	*	*	*
1951	5.6	*	10.6	15.9	9.9	*	*	*	*	*
1950	6.1	*	11.2	17.0	10.6	*	*
ALL OTHER, MALE												
1959
1958
1957
1956
1955
1954
1953
1952
1951
1950
ALL OTHER, FEMALE												
1959	5.1	*	10.2	15.3	11.3	*	*	*	*	*
1958	5.3	*	9.0	16.5	12.3	*	*	*	*	*
1957	6.4	*	10.8	20.3	13.0	*	*	*	*	*
1956	6.4	*	11.5	19.2	13.4	*	*	*	*	*
1955	7.1	*	12.2	20.8	15.3	*	*	*	*	*
1954	7.8	*	12.1	24.8	14.6	*	*	*	*	*
1953	8.9	*	17.8	23.6	16.2	*	*	*	*	*
1952	10.2	*	20.5	27.1	17.7	*	*	*	*	*
1951	10.9	*	20.0	29.8	19.1	*	*	*	*	*
1950	12.0	*	21.3	32.2	20.5	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CONGENITAL MALFORMATIONS (750-759)												
BOTH SEXES												
1959	12.3	382.3	12.2	3.8	2.8	1.9	2.2	2.6	2.8	2.2	2.4	2.4
1958	12.4	384.7	12.6	3.3	2.5	1.8	2.1	2.6	2.6	2.4	2.7	3.4
1957	12.8	394.3	13.6	3.3	2.3	1.8	2.3	2.8	2.9	2.6	2.8	3.1
1956	12.6	409.1	13.1	2.8	2.0	1.6	2.0	2.7	2.6	2.4	2.2	3.4
1955	12.5	409.2	12.1	2.7	2.1	1.7	2.0	2.7	2.5	2.6	2.8	2.6
1954	12.5	414.5	12.3	2.7	1.9	1.5	1.9	2.7	2.4	2.4	3.6	*
1953	12.6	430.4	12.8	2.4	2.0	1.4	1.9	2.5	2.7	2.4	3.0	3.0
1952	12.7	445.2	12.6	2.4	1.9	1.4	1.8	2.4	2.3	2.0	2.8	*
1951	12.3	446.2	11.3	2.4	1.8	1.4	1.7	2.2	2.0	1.9	2.1	*
1950	12.2	447.2	11.1	2.4	1.8	1.9	2.2	2.6	2.3	2.1	2.0	*
MALE												
1959	13.7	414.0	12.2	4.2	3.0	2.1	2.6	2.7	3.3	2.5	3.0	*
1958	13.6	411.2	12.6	3.5	2.9	1.9	2.4	2.9	2.9	2.7	3.1	*
1957	14.1	424.3	13.6	3.3	2.6	2.0	2.5	3.1	3.2	3.0	3.6	*
1956	13.9	440.0	13.5	3.0	2.3	1.7	2.1	3.0	2.7	3.0	2.9	*
1955	13.7	439.1	12.4	2.8	2.5	1.8	2.0	3.1	2.6	3.0	3.4	*
1954	13.8	448.9	12.2	2.9	2.2	1.7	2.0	3.2	2.7	2.8	3.8	*
1953	14.0	465.9	12.9	2.6	2.4	1.5	2.2	2.8	3.1	2.7	3.5	*
1952	14.0	478.0	12.7	2.5	2.2	1.7	2.1	2.8	2.5	2.1	3.9	*
1951	13.4	475.6	11.0	2.5	2.0	1.5	1.8	2.4	2.4	2.5	2.6	*
1950	13.4	477.4	11.6	2.6	2.0	2.1	2.5	2.7	2.9	2.4	2.1	*
FEMALE												
1959	11.0	349.6	12.1	3.4	2.6	1.7	1.9	2.5	2.3	1.9	1.9	*
1958	11.2	357.4	12.6	3.2	2.0	1.6	1.8	2.3	2.2	2.1	2.4	*
1957	11.6	363.3	13.6	3.3	1.9	1.7	2.1	2.5	2.6	2.2	2.1	*
1956	11.3	377.1	12.8	2.6	1.7	1.5	1.8	2.4	2.5	1.9	1.6	*
1955	11.3	378.1	11.8	2.6	1.6	1.6	1.9	2.2	2.4	2.1	2.3	*
1954	11.1	378.9	12.5	2.5	1.6	1.3	1.7	2.2	2.1	2.0	3.4	*
1953	11.3	393.5	12.8	2.3	1.6	1.3	1.6	2.2	2.4	2.1	2.5	*
1952	11.4	411.1	12.5	2.4	1.6	1.2	1.5	2.1	2.1	1.9	2.0	*
1951	11.2	415.9	11.7	2.3	1.5	1.2	1.5	1.9	1.6	1.4	1.8	*
1950	11.1	416.0	10.5	2.1	1.6	1.7	1.9	2.5	1.8	1.8	2.0	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CONGENITAL MALFORMATIONS (750-759)												
WHITE, BOTH SEXES												
1959	12.2	391.4	11.9	3.8	2.8	1.9	2.3	2.6	2.9	2.2	2.4	2.4
1958	12.1	390.1	12.3	3.4	2.5	1.7	2.0	2.7	2.5	2.3	2.7	3.3
1957	12.6	403.7	13.4	3.3	2.3	1.8	2.3	2.7	2.9	2.6	2.9	3.4
1956	12.5	420.8	13.0	2.9	2.1	1.6	1.9	2.8	2.6	2.5	2.2	3.3
1955	12.4	419.5	11.9	2.8	2.1	1.7	2.0	2.6	2.6	2.6	2.9	*
1954	12.5	428.9	12.4	2.7	1.9	1.5	1.9	2.6	2.5	2.4	3.7	*
1953	12.7	446.4	12.8	2.4	2.0	1.4	1.9	2.6	2.7	2.4	3.0	*
1952	12.8	462.5	12.7	2.5	1.9	1.4	1.8	2.5	2.3	2.1	3.0	*
1951	12.4	465.3	11.4	2.4	1.8	1.4	1.7	2.2	2.0	1.9	2.2	*
1950	12.3	463.9	11.2	2.4	1.8	1.9	2.2	2.6	2.4	2.0	2.1	*
WHITE, MALE												
1959	13.5	423.6	11.9	4.2	3.0	2.1	2.7	2.7	3.4	2.5	3.0	*
1958	13.2	414.4	12.2	3.6	3.0	1.9	2.3	3.0	2.7	2.5	3.2	*
1957	13.8	433.1	13.1	3.3	2.7	1.9	2.5	3.1	3.1	2.9	3.8	*
1956	13.7	450.7	13.1	3.0	2.4	1.7	2.1	3.1	2.7	3.0	3.0	*
1955	13.6	449.3	12.1	2.9	2.5	1.7	2.0	3.1	2.6	3.0	3.5	*
1954	13.7	462.5	12.2	2.9	2.2	1.7	1.9	3.0	2.7	2.8	3.9	*
1953	14.0	479.7	12.7	2.5	2.4	1.5	2.2	2.8	3.1	2.7	3.5	*
1952	14.1	491.7	12.7	2.5	2.2	1.8	2.2	2.8	2.5	2.2	4.1	*
1951	13.4	492.1	10.8	2.4	2.1	1.6	1.9	2.4	2.4	2.5	2.7	*
1950	13.5	493.5	11.6	2.6	2.0	2.1	2.5	2.8	2.9	2.3	2.1	*
WHITE, FEMALE												
1959	10.9	358.0	11.9	3.4	2.7	1.7	1.9	2.5	2.4	1.9	2.0	*
1958	11.0	364.7	12.4	3.3	2.1	1.6	1.7	2.3	2.2	2.2	2.3	*
1957	11.5	373.0	13.6	3.4	2.0	1.7	2.1	2.4	2.7	2.3	2.1	*
1956	11.3	389.5	12.9	2.7	1.8	1.4	1.8	2.4	2.5	2.0	1.6	*
1955	11.2	388.3	11.6	2.7	1.7	1.6	2.0	2.2	2.5	2.2	2.3	*
1954	11.2	393.9	12.7	2.6	1.6	1.4	1.8	2.3	2.2	2.1	3.6	*
1953	11.4	411.4	13.0	2.3	1.6	1.4	1.6	2.3	2.4	2.2	2.6	*
1952	11.6	432.0	12.7	2.4	1.6	1.1	1.5	2.2	2.2	2.0	2.0	*
1951	11.4	437.5	12.0	2.4	1.6	1.3	1.5	2.0	1.6	1.4	1.8	*
1950	11.2	433.0	10.7	2.1	1.6	1.7	1.9	2.5	1.9	1.8	2.1	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CONGENITAL MALFORMATIONS (750-759)												
ALL OTHER, BOTH SEXES												
1959	13.8	330.3	13.5	3.7	2.4	1.8	1.9	2.3	2.4	2.4	*	*
1958	14.4	352.8	14.4	2.7	2.0	1.9	2.2	2.3	3.6	2.8	*	*
1957	14.3	338.4	15.0	3.1	2.1	2.2	2.2	2.9	3.4	2.5	*	*
1956	13.5	338.3	14.0	2.3	1.7	1.6	2.3	2.0	2.4	*	*	*
1955	13.4	345.9	13.5	1.8	1.7	1.9	1.8	2.9	1.8	*	*	*
1954	12.4	323.1	11.9	2.6	1.6	1.4	1.9	3.6	*	*	*	*
1953	12.2	325.8	12.7	2.3	2.3	1.3	1.7	1.9	2.9	*	*	*
1952	11.8	330.2	12.1	2.3	1.8	1.5	1.7	1.9	2.2	*	*	*
1951	11.3	322.6	10.9	2.5	1.2	1.0	1.2	1.9	*	*	*	*
1950	11.5	334.8	10.3	2.2	1.7	1.5	2.5	2.5	*	*	*	*
ALL OTHER, MALE												
1959	15.3	357.3	14.0	4.0	2.6	2.2	2.3	2.7	*	*	*	*
1958	16.3	391.3	15.0	3.1	2.0	2.1	2.4	2.2	4.9	*	*	*
1957	16.1	370.4	16.8	3.2	2.3	2.6	2.1	2.8	4.7	*	*	*
1956	15.3	374.3	16.1	2.6	2.2	1.7	2.3	2.3	*	*	*	*
1955	14.9	376.6	13.9	2.1	2.0	2.1	2.0	3.3	*	*	*	*
1954	14.4	360.5	12.5	3.0	1.7	1.7	2.6	5.2	*	*	*	*
1953	14.3	371.8	14.1	2.8	2.7	*	2.1	2.5	*	*	*	*
1952	13.8	384.7	13.0	2.3	2.2	*	*	*	*	*	*	*
1951	13.2	365.9	12.3	3.3	*	*	*	*	*	*	*	*
1950	12.8	366.0	11.8	2.6	*	*	3.0	*	*	*	*	*
ALL OTHER, FEMALE												
1959	12.4	302.3	12.9	3.5	2.2	*	*	2.0	*	*	*	*
1958	12.6	314.5	13.7	2.4	2.0	1.8	2.0	2.5	*	*	*	*
1957	12.6	306.6	13.2	3.0	1.8	1.9	2.2	3.1	*	*	*	*
1956	11.7	302.6	12.0	2.1	*	1.6	2.4	*	*	*	*	*
1955	11.9	316.7	13.1	1.5	1.5	1.7	1.6	2.4	*	*	*	*
1954	10.6	286.0	11.3	2.3	1.5	*	*	*	*	*	*	*
1953	10.2	280.0	11.4	1.9	1.9	*	*	*	*	*	*	*
1952	9.9	276.4	11.1	2.3	*	1.6	1.7	*	*	*	*	*
1951	9.5	279.7	9.4	1.7	*	*	*	*	*	*	*	*
1950	10.2	303.8	8.8	1.9	2.0	1.7	1.9	2.7	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CERTAIN DISEASES OF EARLY INFANCY (760-776)												
BOTH SEXES												
1959	38.5	1,666.7	*	*	*	*	*	*	*	*	*	*
1958	39.8	1,705.1	*	*	*	*	*	*	*	*	*	*
1957	39.1	1,660.9	*	*	*	*	*	*	*	*	*	*
1956	38.6	1,683.6	*	*	*	*	*	*	*	*	*	*
1955	39.0	1,705.7	0.2	*	*	*	*	*	*	*	*	*
1954	39.4	1,739.8	0.2	*	*	*	*	*	*	*	*	*
1953	40.1	1,799.7	*	*	*	*	*	*	*	*	*	*
1952	40.9	1,865.5	0.2	*	*	*	*	*	*	*	*	*
1951	41.2	1,911.0	0.3	*	*	*	*	*	*	*	*	*
1950	40.5	1,937.3	*	*	*	*	*	*	*	*	*	*
MALE												
1959	45.6	1,917.3	*	*	*	*	*	*	*	*	*	*
1958	46.9	1,946.0	*	*	*	*	*	*	*	*	*	*
1957	46.2	1,901.5	*	*	*	*	*	*	*	*	*	*
1956	45.9	1,942.7	*	*	*	*	*	*	*	*	*	*
1955	46.1	1,956.5	*	*	*	*	*	*	*	*	*	*
1954	46.8	2,000.8	*	*	*	*	*	*	*	*	*	*
1953	47.7	2,074.8	*	*	*	*	*	*	*	*	*	*
1952	48.5	2,143.5	*	*	*	*	*	*	*	*	*	*
1951	49.0	2,215.7	*	*	*	*	*	*	*	*	*	*
1950	47.8	2,234.0	*	*	*	*	*	*	*	*	*	*
FEMALE												
1959	31.5	1,407.9	*	*	*	*	*	*	*	*	*	*
1958	32.9	1,455.7	*	*	*	*	*	*	*	*	*	*
1957	32.2	1,411.6	*	*	*	*	*	*	*	*	*	*
1956	31.5	1,415.4	*	*	*	*	*	*	*	*	*	*
1955	32.0	1,445.5	*	*	*	*	*	*	*	*	*	*
1954	32.2	1,469.6	*	*	*	*	*	*	*	*	*	*
1953	32.7	1,514.8	*	*	*	*	*	*	*	*	*	*
1952	33.5	1,577.0	*	*	*	*	*	*	*	*	*	*
1951	33.4	1,596.0	0.3	*	*	*	*	*	*	*	*	*
1950	33.2	1,629.7	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CERTAIN DISEASES OF EARLY INFANCY (760-776)												
WHITE, BOTH SEXES												
1959	33.0	1,490.9	*	*	*	*	*	*	*	*	*	*
1958	34.1	1,518.3	*	*	*	*	*	*	*	*	*	*
1957	33.7	1,489.7	*	*	*	*	*	*	*	*	*	*
1956	33.5	1,520.0	*	*	*	*	*	*	*	*	*	*
1955	34.2	1,550.5	*	*	*	*	*	*	*	*	*	*
1954	34.7	1,586.5	0.2	*	*	*	*	*	*	*	*	*
1953	35.7	1,650.6	*	*	*	*	*	*	*	*	*	*
1952	36.6	1,718.4	0.2	*	*	*	*	*	*	*	*	*
1951	37.2	1,782.8	0.2	*	*	*	*	*	*	*	*	*
1950	36.6	1,802.1	*	*	*	*	*	*	*	*	*	*
WHITE, MALE												
1959	39.6	1,730.3	*	*	*	*	*	*	*	*	*	*
1958	40.6	1,748.6	*	*	*	*	*	*	*	*	*	*
1957	40.2	1,720.4	*	*	*	*	*	*	*	*	*	*
1956	40.1	1,760.9	*	*	*	*	*	*	*	*	*	*
1955	40.8	1,791.8	*	*	*	*	*	*	*	*	*	*
1954	41.6	1,838.0	*	*	*	*	*	*	*	*	*	*
1953	43.0	1,923.9	*	*	*	*	*	*	*	*	*	*
1952	43.8	1,986.0	*	*	*	*	*	*	*	*	*	*
1951	44.5	2,075.8	*	*	*	*	*	*	*	*	*	*
1950	43.7	2,094.2	*	*	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	26.6	1,242.1	*	*	*	*	*	*	*	*	*	*
1958	27.8	1,278.2	*	*	*	*	*	*	*	*	*	*
1957	27.3	1,249.0	*	*	*	*	*	*	*	*	*	*
1956	27.1	1,268.1	*	*	*	*	*	*	*	*	*	*
1955	27.7	1,298.6	*	*	*	*	*	*	*	*	*	*
1954	28.1	1,324.4	*	*	*	*	*	*	*	*	*	*
1953	28.5	1,364.6	*	*	*	*	*	*	*	*	*	*
1952	29.6	1,438.5	*	*	*	*	*	*	*	*	*	*
1951	29.9	1,477.8	*	*	*	*	*	*	*	*	*	*
1950	29.6	1,497.3	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
CERTAIN DISEASES OF EARLY INFANCY (760-776)												
ALL OTHER, BOTH SEXES												
1959	81.9	2,684.9	*	*	*	*	*	*	*	*	*	*
1958	85.3	2,823.0	*	*	*	*	*	*	*	*	*	*
1957	82.6	2,683.0	*	*	*	*	*	*	*	*	*	*
1956	80.0	2,668.1	*	*	*	*	*	*	*	*	*	*
1955	78.6	2,660.0	*	*	*	*	*	*	*	*	*	*
1954	78.0	2,709.0	*	*	*	*	*	*	*	*	*	*
1953	77.1	2,771.0	*	*	*	*	*	*	*	*	*	*
1952	77.0	2,841.2	*	*	*	*	*	*	*	*	*	*
1951	75.0	2,739.8	*	*	*	*	*	*	*	*	*	*
1950	73.5	2,849.8	*	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	94.7	3,020.3	*	*	*	*	*	*	*	*	*	*
1958	97.9	3,153.6	*	*	*	*	*	*	*	*	*	*
1957	95.1	3,007.0	*	*	*	*	*	*	*	*	*	*
1956	94.1	3,063.6	*	*	*	*	*	*	*	*	*	*
1955	90.8	3,005.0	*	*	*	*	*	*	*	*	*	*
1954	90.3	3,055.6	*	*	*	*	*	*	*	*	*	*
1953	87.9	3,071.8	*	*	*	*	*	*	*	*	*	*
1952	89.1	3,218.5	*	*	*	*	*	*	*	*	*	*
1951	87.9	3,141.8	*	*	*	*	*	*	*	*	*	*
1950	84.1	3,200.3	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	69.9	2,342.7	*	*	*	*	*	*	*	*	*	*
1958	73.4	2,493.4	*	*	*	*	*	*	*	*	*	*
1957	70.7	2,360.1	*	*	*	*	*	*	*	*	*	*
1956	66.6	2,275.5	*	*	*	*	*	*	*	*	*	*
1955	67.1	2,327.8	*	*	*	*	*	*	*	*	*	*
1954	66.4	2,365.2	*	*	*	*	*	*	*	*	*	*
1953	66.9	2,471.5	*	*	*	*	*	*	*	*	*	*
1952	65.5	2,468.9	*	*	*	*	*	*	*	*	*	*
1951	62.8	2,341.4	*	*	*	*	*	*	*	*	*	*
1950	63.3	2,501.4	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)												
BOTH SEXES												
1959	17.1	742.2	*	*	*	*	*	*	*	*	*	*
1958	18.2	779.7	*	*	*	*	*	*	*	*	*	*
1957	17.8	755.5	*	*	*	*	*	*	*	*	*	*
1956	17.7	772.2	*	*	*	*	*	*	*	*	*	*
1955	17.8	779.2	*	*	*	*	*	*	*	*	*	*
1954	17.8	785.1	*	*	*	*	*	*	*	*	*	*
1953	17.9	801.9	*	*	*	*	*	*	*	*	*	*
1952	17.6	801.9	*	*	*	*	*	*	*	*	*	*
1951	17.6	817.1	*	*	*	*	*	*	*	*	*	*
1950	16.6	796.8	*	*	*	*	*	*	*	*	*	*
MALE												
1959	20.8	873.5	*	*	*	*	*	*	*	*	*	*
1958	21.9	910.0	*	*	*	*	*	*	*	*	*	*
1957	21.6	889.5	*	*	*	*	*	*	*	*	*	*
1956	21.6	916.5	*	*	*	*	*	*	*	*	*	*
1955	21.6	918.3	*	*	*	*	*	*	*	*	*	*
1954	21.8	931.5	*	*	*	*	*	*	*	*	*	*
1953	22.0	955.4	*	*	*	*	*	*	*	*	*	*
1952	21.6	955.0	*	*	*	*	*	*	*	*	*	*
1951	21.7	981.6	*	*	*	*	*	*	*	*	*	*
1950	20.4	951.0	*	*	*	*	*	*	*	*	*	*
FEMALE												
1959	13.6	606.7	*	*	*	*	*	*	*	*	*	*
1958	14.6	644.8	*	*	*	*	*	*	*	*	*	*
1957	14.1	616.7	*	*	*	*	*	*	*	*	*	*
1956	13.8	622.8	*	*	*	*	*	*	*	*	*	*
1955	14.1	635.0	*	*	*	*	*	*	*	*	*	*
1954	13.9	633.5	*	*	*	*	*	*	*	*	*	*
1953	13.9	642.9	*	*	*	*	*	*	*	*	*	*
1952	13.6	643.1	*	*	*	*	*	*	*	*	*	*
1951	13.5	647.0	*	*	*	*	*	*	*	*	*	*
1950	13.0	636.9	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)												
WHITE, BOTH SEXES												
1959	15.5	697.6	*	*	*	*	*	*	*	*	*	*
1958	16.4	728.8	*	*	*	*	*	*	*	*	*	*
1957	16.0	705.7	*	*	*	*	*	*	*	*	*	*
1956	16.1	728.5	*	*	*	*	*	*	*	*	*	*
1955	16.3	740.6	*	*	*	*	*	*	*	*	*	*
1954	16.4	750.6	*	*	*	*	*	*	*	*	*	*
1953	16.7	770.6	*	*	*	*	*	*	*	*	*	*
1952	16.5	773.9	*	*	*	*	*	*	*	*	*	*
1951	16.7	800.2	*	*	*	*	*	*	*	*	*	*
1950	15.8	778.8	*	*	*	*	*	*	*	*	*	*
WHITE, MALE												
1959	18.9	827.9	*	*	*	*	*	*	*	*	*	*
1958	19.9	857.6	*	*	*	*	*	*	*	*	*	*
1957	19.5	835.3	*	*	*	*	*	*	*	*	*	*
1956	19.8	868.7	*	*	*	*	*	*	*	*	*	*
1955	20.1	880.9	*	*	*	*	*	*	*	*	*	*
1954	20.3	897.8	*	*	*	*	*	*	*	*	*	*
1953	20.7	927.2	*	*	*	*	*	*	*	*	*	*
1952	20.4	925.3	*	*	*	*	*	*	*	*	*	*
1951	20.7	965.4	*	*	*	*	*	*	*	*	*	*
1950	19.5	934.7	*	*	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	12.1	562.2	*	*	*	*	*	*	*	*	*	*
1958	12.9	594.6	*	*	*	*	*	*	*	*	*	*
1957	12.5	570.5	*	*	*	*	*	*	*	*	*	*
1956	12.4	581.9	*	*	*	*	*	*	*	*	*	*
1955	12.7	594.0	*	*	*	*	*	*	*	*	*	*
1954	12.6	597.1	*	*	*	*	*	*	*	*	*	*
1953	12.7	606.8	*	*	*	*	*	*	*	*	*	*
1952	12.7	615.5	*	*	*	*	*	*	*	*	*	*
1951	12.7	628.3	*	*	*	*	*	*	*	*	*	*
1950	12.2	616.1	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)												
ALL OTHER, BOTH SEXES												
1959	30.6	1,001.5	*	*	*	*	*	*	*	*	*	*
1958	32.8	1,084.1	*	*	*	*	*	*	*	*	*	*
1957	32.4	1,052.9	*	*	*	*	*	*	*	*	*	*
1956	31.0	1,035.5	*	*	*	*	*	*	*	*	*	*
1955	30.1	1,017.0	*	*	*	*	*	*	*	*	*	*
1954	28.9	1,003.2	*	*	*	*	*	*	*	*	*	*
1953	28.0	1,005.8	*	*	*	*	*	*	*	*	*	*
1952	26.8	987.9	*	*	*	*	*	*	*	*	*	*
1951	25.4	926.2	*	*	*	*	*	*	*	*	*	*
1950	23.7	918.2	*	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	35.8	1,142.7	*	*	*	*	*	*	*	*	*	*
1958	38.2	1,230.1	*	*	*	*	*	*	*	*	*	*
1957	38.6	1,220.6	*	*	*	*	*	*	*	*	*	*
1956	37.2	1,211.4	*	*	*	*	*	*	*	*	*	*
1955	35.0	1,157.9	*	*	*	*	*	*	*	*	*	*
1954	34.0	1,149.6	*	*	*	*	*	*	*	*	*	*
1953	32.6	1,139.3	*	*	*	*	*	*	*	*	*	*
1952	32.0	1,157.2	*	*	*	*	*	*	*	*	*	*
1951	30.5	1,089.1	*	*	*	*	*	*	*	*	*	*
1950	28.0	1,063.8	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	25.6	857.9	*	*	*	*	*	*	*	*	*	*
1958	27.6	938.6	*	*	*	*	*	*	*	*	*	*
1957	26.5	885.8	*	*	*	*	*	*	*	*	*	*
1956	25.2	860.9	*	*	*	*	*	*	*	*	*	*
1955	25.4	881.0	*	*	*	*	*	*	*	*	*	*
1954	24.1	858.0	*	*	*	*	*	*	*	*	*	*
1953	23.6	872.8	*	*	*	*	*	*	*	*	*	*
1952	21.8	820.9	*	*	*	*	*	*	*	*	*	*
1951	20.5	764.9	*	*	*	*	*	*	*	*	*	*
1950	19.6	773.6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFECTIONS OF NEWBORN (763-768)												
BOTH SEXES												
1959	2.9	124.3	*	*	*	*	*	*	*	*	*	*
1958	3.2	136.8	*	*	*	*	*	*	*	*	*	*
1957	3.0	127.0	*	*	*	*	*	*	*	*	*	*
1956	2.8	120.4	*	*	*	*	*	*	*	*	*	*
1955	2.6	111.9	*	*	*	*	*	*	*	*	*	*
1954	2.7	119.5	*	*	*	*	*	*	*	*	*	*
1953	2.6	118.8	*	*	*	*	*	*	*	*	*	*
1952	2.6	120.0	*	*	*	*	*	*	*	*	*	*
1951	2.6	119.0	*	*	*	*	*	*	*	*	*	*
1950	2.6	122.2	*	*	*	*	*	*	*	*	*	*
MALE												
1959	3.4	143.9	*	*	*	*	*	*	*	*	*	*
1958	3.8	156.3	*	*	*	*	*	*	*	*	*	*
1957	3.5	146.1	*	*	*	*	*	*	*	*	*	*
1956	3.3	137.8	*	*	*	*	*	*	*	*	*	*
1955	3.1	130.6	*	*	*	*	*	*	*	*	*	*
1954	3.2	137.1	*	*	*	*	*	*	*	*	*	*
1953	3.1	135.4	*	*	*	*	*	*	*	*	*	*
1952	3.1	137.7	*	*	*	*	*	*	*	*	*	*
1951	3.0	136.4	*	*	*	*	*	*	*	*	*	*
1950	2.9	137.8	*	*	*	*	*	*	*	*	*	*
FEMALE												
1959	2.3	104.0	*	*	*	*	*	*	*	*	*	*
1958	2.6	116.5	*	*	*	*	*	*	*	*	*	*
1957	2.4	107.1	*	*	*	*	*	*	*	*	*	*
1956	2.3	102.3	*	*	*	*	*	*	*	*	*	*
1955	2.0	92.6	*	*	*	*	*	*	*	*	*	*
1954	2.2	101.3	*	*	*	*	*	*	*	*	*	*
1953	2.2	101.7	*	*	*	*	*	*	*	*	*	*
1952	2.2	101.7	*	*	*	*	*	*	*	*	*	*
1951	2.1	101.0	*	*	*	*	*	*	*	*	*	*
1950	2.2	106.1	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFECTIONS OF NEWBORN (763-768)												
WHITE, BOTH SEXES												
1959	2.0	91.8	*	*	*	*	*	*	*	*	*	*
1958	2.3	102.7	*	*	*	*	*	*	*	*	*	*
1957	2.2	96.4	*	*	*	*	*	*	*	*	*	*
1956	2.0	91.7	*	*	*	*	*	*	*	*	*	*
1955	1.9	85.1	*	*	*	*	*	*	*	*	*	*
1954	2.0	93.2	*	*	*	*	*	*	*	*	*	*
1953	2.0	92.2	*	*	*	*	*	*	*	*	*	*
1952	2.0	94.5	*	*	*	*	*	*	*	*	*	*
1951	2.0	97.5	*	*	*	*	*	*	*	*	*	*
1950	2.1	101.3	*	*	*	*	*	*	*	*	*	*
WHITE, MALE												
1959	2.4	106.6	*	*	*	*	*	*	*	*	*	*
1958	2.7	117.6	*	*	*	*	*	*	*	*	*	*
1957	2.7	114.0	*	*	*	*	*	*	*	*	*	*
1956	2.4	105.1	*	*	*	*	*	*	*	*	*	*
1955	2.3	100.4	*	*	*	*	*	*	*	*	*	*
1954	2.5	108.5	*	*	*	*	*	*	*	*	*	*
1953	2.4	106.5	*	*	*	*	*	*	*	*	*	*
1952	2.4	108.3	*	*	*	*	*	*	*	*	*	*
1951	2.4	111.8	*	*	*	*	*	*	*	*	*	*
1950	2.4	116.0	*	*	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	1.6	76.5	*	*	*	*	*	*	*	*	*	*
1958	1.9	87.2	*	*	*	*	*	*	*	*	*	*
1957	1.7	78.0	*	*	*	*	*	*	*	*	*	*
1956	1.7	77.6	*	*	*	*	*	*	*	*	*	*
1955	1.5	69.2	*	*	*	*	*	*	*	*	*	*
1954	1.6	77.3	*	*	*	*	*	*	*	*	*	*
1953	1.6	77.3	*	*	*	*	*	*	*	*	*	*
1952	1.6	80.0	*	*	*	*	*	*	*	*	*	*
1951	1.7	82.7	*	*	*	*	*	*	*	*	*	*
1950	1.7	86.0	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
INFECTIONS OF NEWBORN (763-768)												
ALL OTHER, BOTH SEXES												
1959	9.5	312.1	*	*	*	*	*	*	*	*	*	*
1958	10.3	340.4	*	*	*	*	*	*	*	*	*	*
1957	9.5	309.6	*	*	*	*	*	*	*	*	*	*
1956	8.8	293.2	*	*	*	*	*	*	*	*	*	*
1955	8.2	276.8	*	*	*	*	*	*	*	*	*	*
1954	8.2	285.9	*	*	*	*	*	*	*	*	*	*
1953	8.1	291.9	*	*	*	*	*	*	*	*	*	*
1952	7.8	289.5	*	*	*	*	*	*	*	*	*	*
1951	7.1	257.7	*	*	*	*	*	*	*	*	*	*
1950	6.8	263.2	*	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	11.4	364.3	*	*	*	*	*	*	*	*	*	*
1958	12.2	393.1	*	*	*	*	*	*	*	*	*	*
1957	10.8	342.2	*	*	*	*	*	*	*	*	*	*
1956	10.4	339.3	*	*	*	*	*	*	*	*	*	*
1955	9.7	321.8	*	*	*	*	*	*	*	*	*	*
1954	9.5	322.6	*	*	*	*	*	*	*	*	*	*
1953	9.4	326.9	*	*	*	*	*	*	*	*	*	*
1952	9.4	337.8	*	*	*	*	*	*	*	*	*	*
1951	8.4	299.1	*	*	*	*	*	*	*	*	*	*
1950	7.6	287.9	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	7.7	259.3	*	*	*	*	*	*	*	*	*	*
1958	8.5	287.9	*	*	*	*	*	*	*	*	*	*
1957	8.3	277.1	*	*	*	*	*	*	*	*	*	*
1956	7.2	247.4	*	*	*	*	*	*	*	*	*	*
1955	6.7	233.1	*	*	*	*	*	*	*	*	*	*
1954	7.0	249.6	*	*	*	*	*	*	*	*	*	*
1953	7.0	257.0	*	*	*	*	*	*	*	*	*	*
1952	6.4	241.8	*	*	*	*	*	*	*	*	*	*
1951	5.8	216.7	*	*	*	*	*	*	*	*	*	*
1950	6.0	238.6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURITY UNQUALIFIED (769-776)												
BOTH SEXES												
1959	18.5	800.1	*	*	*	*	*	*	*	*	*	*
1958	18.4	788.6	*	*	*	*	*	*	*	*	*	*
1957	18.3	778.4	*	*	*	*	*	*	*	*	*	*
1956	18.1	791.0	*	*	*	*	*	*	*	*	*	*
1955	18.6	814.6	*	*	*	*	*	*	*	*	*	*
1954	18.9	835.2	*	*	*	*	*	*	*	*	*	*
1953	19.6	879.0	*	*	*	*	*	*	*	*	*	*
1952	20.7	943.6	0.1	*	*	*	*	*	*	*	*	*
1951	21.0	974.9	0.3	*	*	*	*	*	*	*	*	*
1950	21.3	1,018.3	*	*	*	*	*	*	*	*	*	*
MALE												
1959	21.4	899.9	*	*	*	*	*	*	*	*	*	*
1958	21.2	879.7	*	*	*	*	*	*	*	*	*	*
1957	21.0	865.9	*	*	*	*	*	*	*	*	*	*
1956	21.0	888.4	*	*	*	*	*	*	*	*	*	*
1955	21.4	907.7	*	*	*	*	*	*	*	*	*	*
1954	21.8	932.2	*	*	*	*	*	*	*	*	*	*
1953	22.6	984.0	*	*	*	*	*	*	*	*	*	*
1952	23.8	1,050.9	*	*	*	*	*	*	*	*	*	*
1951	24.3	1,097.7	*	*	*	*	*	*	*	*	*	*
1950	24.5	1,145.2	*	*	*	*	*	*	*	*	*	*
FEMALE												
1959	15.6	697.2	*	*	*	*	*	*	*	*	*	*
1958	15.7	694.4	*	*	*	*	*	*	*	*	*	*
1957	15.7	687.9	*	*	*	*	*	*	*	*	*	*
1956	15.3	690.2	*	*	*	*	*	*	*	*	*	*
1955	15.9	718.0	*	*	*	*	*	*	*	*	*	*
1954	16.1	734.7	*	*	*	*	*	*	*	*	*	*
1953	16.6	770.2	*	*	*	*	*	*	*	*	*	*
1952	17.7	832.2	*	*	*	*	*	*	*	*	*	*
1951	17.8	848.0	0.3	*	*	*	*	*	*	*	*	*
1950	18.1	886.7	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURITY UNQUALIFIED (769-776)												
WHITE, BOTH SEXES												
1959	15.5	701.6	*	*	*	*	*	*	*	*	*	*
1958	15.4	686.7	*	*	*	*	*	*	*	*	*	*
1957	15.6	687.6	*	*	*	*	*	*	*	*	*	*
1956	15.4	699.9	*	*	*	*	*	*	*	*	*	*
1955	16.0	724.8	*	*	*	*	*	*	*	*	*	*
1954	16.3	742.7	*	*	*	*	*	*	*	*	*	*
1953	17.0	787.8	*	*	*	*	*	*	*	*	*	*
1952	18.1	850.0	*	*	*	*	*	*	*	*	*	*
1951	18.5	885.0	0.2	*	*	*	*	*	*	*	*	*
1950	18.7	922.0	*	*	*	*	*	*	*	*	*	*
WHITE, MALE												
1959	18.2	795.9	*	*	*	*	*	*	*	*	*	*
1958	18.0	773.3	*	*	*	*	*	*	*	*	*	*
1957	18.0	771.1	*	*	*	*	*	*	*	*	*	*
1956	17.9	787.1	*	*	*	*	*	*	*	*	*	*
1955	18.5	810.5	*	*	*	*	*	*	*	*	*	*
1954	18.8	831.7	*	*	*	*	*	*	*	*	*	*
1953	19.9	890.1	*	*	*	*	*	*	*	*	*	*
1952	21.0	952.3	*	*	*	*	*	*	*	*	*	*
1951	21.4	998.6	*	*	*	*	*	*	*	*	*	*
1950	21.8	1,043.5	*	*	*	*	*	*	*	*	*	*
WHITE, FEMALE												
1959	12.9	603.5	*	*	*	*	*	*	*	*	*	*
1958	13.0	596.5	*	*	*	*	*	*	*	*	*	*
1957	13.2	600.5	*	*	*	*	*	*	*	*	*	*
1956	13.0	608.6	*	*	*	*	*	*	*	*	*	*
1955	13.5	635.4	*	*	*	*	*	*	*	*	*	*
1954	13.8	649.9	*	*	*	*	*	*	*	*	*	*
1953	14.2	680.5	*	*	*	*	*	*	*	*	*	*
1952	15.3	743.1	*	*	*	*	*	*	*	*	*	*
1951	15.5	766.8	*	*	*	*	*	*	*	*	*	*
1950	15.7	795.2	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURITY UNQUALIFIED (769-776)												
ALL OTHER, BOTH SEXES												
1959	41.8	1,371.2	*	*	*	*	*	*	*	*	*	*
1958	42.3	1,398.4	*	*	*	*	*	*	*	*	*	*
1957	40.6	1,320.5	*	*	*	*	*	*	*	*	*	*
1956	40.1	1,339.4	*	*	*	*	*	*	*	*	*	*
1955	40.4	1,366.3	*	*	*	*	*	*	*	*	*	*
1954	40.9	1,419.9	*	*	*	*	*	*	*	*	*	*
1953	41.0	1,473.3	*	*	*	*	*	*	*	*	*	*
1952	42.4	1,563.8	*	*	*	*	*	*	*	*	*	*
1951	42.6	1,555.9	*	*	*	*	*	*	*	*	*	*
1950	43.0	1,668.3	*	*	*	*	*	*	*	*	*	*
ALL OTHER, MALE												
1959	47.4	1,513.3	*	*	*	*	*	*	*	*	*	*
1958	47.5	1,530.4	*	*	*	*	*	*	*	*	*	*
1957	45.7	1,444.3	*	*	*	*	*	*	*	*	*	*
1956	46.5	1,512.9	*	*	*	*	*	*	*	*	*	*
1955	46.1	1,525.3	*	*	*	*	*	*	*	*	*	*
1954	46.8	1,583.5	*	*	*	*	*	*	*	*	*	*
1953	46.0	1,605.6	*	*	*	*	*	*	*	*	*	*
1952	47.7	1,723.4	*	*	*	*	*	*	*	*	*	*
1951	49.1	1,753.6	*	*	*	*	*	*	*	*	*	*
1950	48.6	1,848.6	*	*	*	*	*	*	*	*	*	*
ALL OTHER, FEMALE												
1959	36.5	1,225.5	*	*	*	*	*	*	*	*	*	*
1958	37.3	1,266.9	*	*	*	*	*	*	*	*	*	*
1957	35.9	1,197.2	*	*	*	*	*	*	*	*	*	*
1956	34.2	1,167.2	*	*	*	*	*	*	*	*	*	*
1955	35.0	1,213.7	*	*	*	*	*	*	*	*	*	*
1954	35.3	1,257.6	*	*	*	*	*	*	*	*	*	*
1953	36.3	1,341.7	*	*	*	*	*	*	*	*	*	*
1952	37.3	1,406.2	*	*	*	*	*	*	*	*	*	*
1951	36.5	1,359.9	*	*	*	*	*	*	*	*	*	*
1950	37.7	1,489.2	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)												
BOTH SEXES												
1959	10.8	60.5	2.4	0.5	1.4	2.5	4.5	9.0	16.1	32.4	75.6	308.1
1958	11.4	68.1	2.7	0.5	1.4	2.7	4.8	10.0	16.0	31.7	80.9	345.0
1957	11.2	63.8	2.4	0.5	1.7	2.5	4.3	8.6	16.6	31.9	81.0	379.2
1956	11.3	66.7	2.4	0.5	1.5	2.5	4.8	9.3	16.6	31.4	82.5	371.1
1955	12.1	74.3	3.0	0.6	1.9	2.6	4.8	10.1	17.1	34.2	87.7	401.2
1954	12.5	76.2	3.1	0.6	2.1	2.7	4.9	9.9	16.6	35.9	97.4	444.7
1953	13.7	84.1	3.5	0.7	1.7	2.5	4.9	10.6	18.1	39.4	110.6	543.7
1952	14.1	98.3	3.5	0.7	1.6	2.4	4.9	10.6	17.7	39.7	116.4	582.3
1951	14.8	102.4	3.9	0.8	1.7	2.5	5.1	10.3	18.3	41.2	128.0	661.9
1950	14.9	115.5	3.9	0.8	1.6	2.5	4.9	10.4	17.4	42.2	130.7	721.1
MALE												
1959	12.9	66.5	2.5	0.5	1.8	2.8	5.5	12.4	22.5	45.6	95.4	331.8
1958	13.4	74.6	2.9	0.5	1.8	2.9	5.8	13.4	22.4	44.2	101.6	372.4
1957	13.3	71.5	2.5	0.6	2.1	2.8	5.5	11.4	23.4	43.4	103.0	413.1
1956	13.5	72.0	2.6	0.6	1.9	2.9	6.1	12.8	23.4	43.8	103.5	403.4
1955	14.6	81.7	3.3	0.7	2.4	3.1	6.2	13.8	24.3	47.3	112.2	419.8
1954	15.1	83.9	3.5	0.7	2.7	3.2	6.3	13.6	24.0	49.1	125.0	457.9
1953	16.1	93.0	3.5	0.9	2.0	2.7	6.2	14.4	25.5	51.9	136.0	569.6
1952	16.5	109.9	3.6	0.8	2.0	2.7	6.2	14.0	24.7	53.0	144.8	580.6
1951	17.3	117.4	4.2	0.9	2.0	2.7	6.4	13.9	24.5	56.4	154.1	700.4
1950	17.1	128.2	4.0	0.9	1.8	2.6	5.8	13.6	24.0	55.8	154.5	782.4
FEMALE												
1959	8.8	54.4	2.2	0.4	1.0	2.3	3.6	5.7	10.1	20.8	60.2	292.6
1958	9.4	61.3	2.5	0.4	1.1	2.4	3.8	6.7	9.9	20.7	64.5	326.7
1957	9.2	55.9	2.3	0.4	1.4	2.3	3.2	5.8	10.1	21.6	63.5	356.1
1956	9.2	61.2	2.3	0.5	1.1	2.2	3.6	5.9	10.0	20.3	65.8	348.7
1955	9.7	66.7	2.7	0.5	1.5	2.1	3.5	6.5	10.3	22.3	68.0	388.0
1954	10.1	68.3	2.7	0.5	1.5	2.2	3.6	6.3	9.5	23.9	74.9	435.3
1953	11.4	74.9	3.5	0.6	1.5	2.2	3.7	6.8	10.9	27.9	89.8	525.1
1952	11.8	86.3	3.4	0.6	1.3	2.2	3.5	7.2	10.8	27.6	92.9	583.5
1951	12.3	87.0	3.6	0.7	1.4	2.3	3.9	6.7	12.2	27.1	106.0	634.4
1950	12.8	102.4	3.7	0.7	1.4	2.4	4.0	7.1	10.7	29.6	110.5	678.4

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)												
WHITE, BOTH SEXES												
1959	8.0	29.4	1.3	0.3	1.0	1.5	3.1	6.3	11.5	23.8	61.2	267.3
1958	8.3	33.7	1.4	0.3	0.9	1.5	3.2	7.1	11.4	23.2	66.2	297.5
1957	8.3	31.5	1.4	0.4	1.2	1.4	3.0	5.9	11.7	23.0	66.1	329.7
1956	8.3	32.3	1.3	0.3	1.0	1.5	3.3	6.5	11.6	23.3	67.4	325.7
1955	9.0	37.2	1.8	0.3	1.4	1.6	3.3	7.0	12.5	25.1	71.6	354.4
1954	9.3	36.7	1.9	0.4	1.5	1.7	3.3	6.8	12.0	26.6	79.0	393.6
1953	9.8	41.6	2.0	0.5	1.0	1.4	3.1	6.9	12.3	27.7	90.0	463.7
1952	10.1	47.7	1.9	0.5	1.0	1.4	3.2	6.7	12.2	28.1	95.5	496.4
1951	10.6	51.8	2.5	0.5	1.0	1.5	3.3	6.6	12.2	29.2	102.9	570.8
1950	10.7	60.9	2.3	0.5	0.9	1.4	3.2	6.5	11.9	29.0	104.5	618.7
WHITE, MALE												
1959	9.8	33.5	1.5	0.4	1.3	1.8	4.0	9.3	17.3	35.5	77.2	285.0
1958	10.1	38.1	1.3	0.4	1.2	1.7	4.1	10.3	16.9	34.3	84.0	307.0
1957	10.1	37.2	1.4	0.5	1.6	1.7	4.0	8.5	17.8	32.8	84.3	352.7
1956	10.3	35.7	1.4	0.3	1.3	1.7	4.5	9.8	17.8	34.1	85.7	355.3
1955	11.2	44.0	2.0	0.5	1.9	2.1	4.5	10.5	19.0	36.6	93.5	370.1
1954	11.5	41.2	2.1	0.5	2.2	2.1	4.5	10.4	18.4	38.3	102.1	398.2
1953	11.9	49.6	1.9	0.6	1.2	1.6	4.2	10.4	18.8	38.8	111.8	477.2
1952	12.2	54.5	1.9	0.5	1.3	1.8	4.4	9.9	18.6	40.4	119.6	492.1
1951	12.7	61.0	2.6	0.7	1.2	1.6	4.4	9.8	17.8	42.2	124.9	599.2
1950	12.5	67.5	2.6	0.6	1.1	1.7	4.1	9.3	17.6	40.7	122.0	663.3
WHITE, FEMALE												
1959	6.2	25.2	1.1	0.3	0.7	1.3	2.2	3.3	6.1	13.7	48.9	256.0
1958	6.6	29.1	1.4	0.3	0.7	1.4	2.3	3.9	6.2	13.3	52.3	291.2
1957	6.5	25.6	1.4	0.3	0.8	1.2	1.9	3.4	6.0	14.4	51.8	314.0
1956	6.4	28.8	1.3	0.3	0.7	1.2	2.2	3.3	5.8	13.6	52.9	305.1
1955	6.8	30.1	1.5	0.2	0.8	1.1	2.2	3.6	6.1	14.7	54.2	343.4
1954	7.1	32.0	1.5	0.3	0.9	1.3	2.1	3.3	5.8	16.0	60.4	390.3
1953	7.8	33.2	2.1	0.3	0.8	1.2	2.0	3.4	5.9	17.6	72.2	453.9
1952	8.1	40.5	1.9	0.4	0.7	1.0	2.0	3.6	5.9	16.9	75.7	499.4
1951	8.6	42.1	2.4	0.4	0.7	1.3	2.1	3.3	6.7	17.2	84.7	550.6
1950	8.9	53.9	2.1	0.4	0.7	1.2	2.3	3.6	6.2	18.2	89.8	587.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)												
ALL OTHER, BOTH SEXES												
1959	33.5	240.4	9.1	1.3	4.2	10.5	17.0	34.8	64.6	134.9	274.4	805.9
1958	36.0	273.6	10.9	1.4	5.0	11.7	18.6	37.8	65.1	134.5	284.9	916.7
1957	35.4	256.7	8.6	1.5	5.6	11.6	16.4	33.8	69.9	137.7	290.4	967.2
1956	35.6	273.4	9.3	2.0	4.6	11.1	17.7	36.2	72.2	129.1	297.0	919.7
1955	38.1	302.5	11.1	2.2	5.8	10.8	17.8	39.5	71.4	142.9	319.5	972.4
1954	39.6	326.1	11.3	2.0	6.2	11.2	18.7	39.8	71.2	146.8	364.3	1,063.6
1953	46.2	361.0	13.4	2.7	7.4	11.2	20.9	46.0	88.9	179.1	413.2	1,526.9
1952	47.8	434.0	14.3	2.6	6.7	11.2	19.5	47.4	85.7	178.6	423.5	1,642.9
1951	49.9	429.9	14.1	2.7	6.6	11.2	21.6	45.5	96.1	185.0	499.1	1,763.8
1950	51.4	484.5	15.1	3.2	6.5	11.4	20.3	47.9	83.4	221.2	527.2	1,938.1
ALL OTHER, MALE												
1959	37.9	261.3	9.0	1.3	5.2	11.5	19.1	41.6	77.0	164.4	328.5	858.6
1958	41.1	297.9	12.8	1.4	6.3	13.1	20.8	42.9	80.8	160.2	327.3	1,151.9
1957	40.3	280.8	9.6	1.7	6.2	12.6	18.3	39.0	83.6	168.1	343.0	1,134.6
1956	40.3	296.0	10.1	2.3	5.7	13.0	20.0	42.2	85.2	158.0	333.1	984.0
1955	42.6	319.9	11.5	2.4	5.8	12.4	21.1	45.5	83.9	173.5	356.0	1,029.2
1954	45.6	360.5	12.2	2.1	7.0	13.1	21.5	44.6	88.4	177.0	428.6	1,222.7
1953	51.6	382.1	14.2	3.0	8.5	12.1	23.7	52.6	104.8	206.7	462.3	1,747.6
1952	53.1	487.8	15.0	3.2	7.2	10.4	22.7	53.4	98.1	202.4	487.3	1,700.0
1951	56.5	490.5	16.1	2.6	7.4	12.1	24.2	52.2	108.5	224.5	557.5	1,963.2
1950	57.9	547.7	14.8	3.4	6.9	10.6	21.4	54.8	100.5	251.9	607.5	2,174.8
ALL OTHER, FEMALE												
1959	29.3	218.9	9.2	1.3	3.3	9.7	15.1	28.5	52.6	108.2	226.1	766.7
1958	31.2	249.3	9.0	1.4	3.8	10.4	16.6	33.1	50.2	110.8	246.4	773.7
1957	30.7	232.6	7.5	1.4	5.0	10.7	14.6	28.8	56.5	109.9	241.4	875.7
1956	31.2	251.1	8.5	1.6	3.5	9.5	15.5	30.3	59.3	102.0	263.0	875.0
1955	33.8	286.3	10.7	2.0	5.8	9.5	14.8	33.6	58.9	114.8	282.6	932.4
1954	34.0	292.0	10.4	1.8	5.4	9.5	16.1	35.1	54.1	118.9	300.8	987.5
1953	41.2	340.0	12.7	2.4	6.4	10.4	18.2	39.6	72.6	152.8	363.6	1,377.4
1952	42.9	380.9	13.7	2.1	6.2	12.0	16.6	41.5	72.9	156.3	362.9	1,603.4
1951	43.6	369.8	12.0	2.8	5.9	10.4	19.3	38.7	83.2	147.7	443.2	1,628.6
1950	45.2	421.6	15.3	3.1	6.2	12.0	19.2	41.0	65.9	190.3	448.1	1,769.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ALL OTHER DISEASES (RESIDUAL)												
BOTH SEXES												
1959	37.3	105.1	12.0	5.1	6.9	10.1	18.7	36.5	74.1	144.4	241.1	462.7
1958	36.6	105.6	12.3	5.0	6.6	10.0	18.6	35.7	72.2	139.3	245.7	444.0
1957	45.5	128.4	11.8	4.7	7.8	11.7	22.9	44.3	92.9	182.1	311.7	570.4
1956	43.0	123.7	11.4	4.6	7.1	11.3	21.5	41.4	88.0	169.2	307.6	555.2
1955	42.1	114.4	11.8	5.0	7.6	11.4	21.6	40.9	85.8	167.1	297.6	530.2
1954	41.2	113.1	12.8	4.9	7.5	11.5	21.2	42.6	84.0	160.1	289.1	509.2
1953	42.6	111.5	13.9	5.3	8.0	12.0	22.6	44.5	89.6	163.7	298.4	531.8
1952	43.2	111.5	14.9	5.4	8.5	12.9	23.8	46.1	90.5	166.1	295.5	545.4
1951	42.8	100.0	13.9	5.6	8.1	12.9	23.8	46.4	90.2	167.2	302.6	574.2
1950	41.0	103.2	14.7	5.4	8.4	12.9	23.6	44.1	86.0	158.6	290.5	581.2
MALE												
1959	45.9	115.7	12.8	5.5	7.8	11.1	21.7	46.1	103.5	205.8	314.9	540.8
1958	45.2	120.5	12.9	5.7	7.5	10.3	21.6	45.4	101.7	198.4	319.6	518.3
1957	54.7	152.1	12.9	5.0	8.8	11.4	25.2	54.0	125.5	248.6	391.6	643.3
1956	51.1	148.3	12.1	4.8	7.9	11.1	23.1	51.0	116.0	226.8	380.0	610.4
1955	49.4	134.6	12.8	5.4	8.5	11.6	23.8	49.6	111.0	217.0	360.0	587.1
1954	48.0	130.5	13.9	5.2	8.1	11.6	22.8	49.6	109.3	206.4	346.3	559.5
1953	49.8	130.6	15.2	5.8	9.2	12.1	24.9	52.5	117.0	206.1	354.9	592.4
1952	49.8	127.5	15.9	5.8	9.5	12.8	25.7	55.2	114.3	205.7	352.1	571.8
1951	49.2	114.1	14.8	6.1	8.9	13.2	25.7	54.8	112.9	206.7	351.8	648.0
1950	45.8	119.0	16.1	5.8	8.8	12.5	25.6	51.5	103.4	188.7	327.6	655.3
FEMALE												
1959	28.9	94.1	11.3	4.6	5.9	9.2	15.9	27.2	46.5	90.8	183.4	411.6
1958	28.1	90.1	11.6	4.3	5.6	9.8	15.8	26.3	44.4	87.2	187.5	394.4
1957	36.5	103.9	10.6	4.4	6.8	12.0	20.8	34.9	62.0	123.1	248.2	520.6
1956	35.1	98.4	10.7	4.3	6.4	11.5	20.0	32.1	61.2	117.6	249.7	516.8
1955	34.9	93.3	10.7	4.6	6.8	11.2	19.5	32.5	61.5	122.1	247.2	490.0
1954	34.5	95.0	11.6	4.6	7.0	11.5	19.6	35.7	59.4	118.1	242.4	473.2
1953	35.6	91.7	12.6	4.8	6.9	11.9	20.4	36.7	62.9	125.0	251.8	488.3
1952	36.8	94.9	13.8	5.0	7.5	13.1	21.8	37.1	67.1	129.8	248.5	526.5
1951	36.6	85.4	12.9	5.1	7.3	12.6	22.0	38.1	67.6	130.7	261.1	521.7
1950	36.2	86.9	13.3	5.1	8.0	13.2	21.7	36.7	68.4	130.7	258.9	529.6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ALL OTHER DISEASES (RESIDUAL)												
WHITE, BOTH SEXES												
1959	36.7	91.0	11.1	5.0	6.1	7.9	15.8	33.6	72.3	145.2	246.2	472.2
1958	35.9	91.2	11.4	4.9	5.8	8.0	15.8	33.1	70.0	140.0	250.3	456.2
1957	44.6	117.4	10.7	4.5	6.9	9.3	19.5	40.0	89.2	182.2	316.2	584.0
1956	42.4	113.9	10.8	4.5	6.3	9.1	18.3	37.5	84.9	169.9	312.5	573.5
1955	41.5	105.8	11.1	4.8	6.7	9.1	18.7	37.8	83.1	167.6	303.1	544.6
1954	40.5	103.4	12.0	4.7	6.5	9.6	18.2	38.8	81.6	161.2	294.3	520.6
1953	41.9	102.3	13.1	5.2	6.9	9.8	19.4	41.0	87.0	164.6	302.6	541.5
1952	42.5	100.4	14.3	5.3	7.5	10.8	20.9	42.5	87.9	166.9	299.0	558.2
1951	42.2	91.6	13.1	5.3	7.0	11.0	20.8	42.8	88.0	168.2	307.3	585.9
1950	40.3	94.7	13.8	5.3	7.3	11.1	20.7	41.1	83.8	158.3	293.4	595.7
WHITE, MALE												
1959	45.4	101.6	11.5	5.5	7.2	8.6	18.5	43.2	102.3	206.7	321.8	552.8
1958	44.7	103.3	12.0	5.6	6.8	8.2	18.6	42.2	99.6	200.2	327.0	534.8
1957	53.9	140.4	11.5	4.9	8.1	9.1	21.7	49.6	121.3	249.3	397.5	654.3
1956	50.6	137.3	11.5	4.8	7.1	9.0	20.3	46.7	113.0	229.1	385.8	624.2
1955	49.0	125.3	11.8	5.2	7.7	9.4	21.1	46.6	108.2	217.8	366.4	599.0
1954	47.4	119.4	12.9	5.1	7.2	9.8	20.0	45.6	106.6	208.0	353.4	571.3
1953	49.0	121.5	14.3	5.6	8.2	10.0	21.6	48.7	114.1	206.9	360.2	592.9
1952	49.0	116.5	15.0	5.7	8.5	10.9	22.9	51.1	111.6	206.7	354.6	581.4
1951	48.7	104.9	14.3	5.7	7.8	11.5	22.8	51.4	111.1	207.9	358.2	658.6
1950	45.1	109.2	15.1	5.6	7.8	10.9	22.8	48.3	101.0	188.4	329.8	670.6
WHITE, FEMALE												
1959	28.2	79.9	10.7	4.5	4.9	7.3	13.3	24.3	44.2	91.5	187.8	420.0
1958	27.4	78.7	10.7	4.2	4.8	7.7	13.2	24.3	42.2	87.0	190.5	404.2
1957	35.6	93.5	9.8	4.2	5.7	9.6	17.4	30.7	58.8	122.7	252.2	536.1
1956	34.3	89.4	10.0	4.1	5.6	9.2	16.5	28.5	58.1	117.0	254.5	538.2
1955	34.3	85.4	10.3	4.4	5.6	8.8	16.4	29.1	59.1	122.4	252.7	506.0
1954	33.9	86.8	11.1	4.4	5.9	9.4	16.5	32.1	57.5	118.8	246.6	484.2
1953	34.9	82.2	11.9	4.7	5.7	9.7	17.3	33.4	60.6	125.9	255.8	504.3
1952	36.1	83.5	13.5	4.9	6.5	10.7	18.9	34.0	64.6	130.5	253.3	541.5
1951	35.8	77.9	11.9	4.8	6.3	10.5	18.8	34.3	65.1	131.6	265.1	534.0
1950	35.6	79.5	12.5	5.1	6.7	11.3	18.6	33.9	66.5	130.4	262.7	543.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
ALL OTHER DISEASES (RESIDUAL)												
ALL OTHER, BOTH SEXES												
1959	42.1	186.5	17.8	5.7	12.7	27.3	44.1	63.7	92.8	135.5	169.8	347.1
1958	41.5	191.5	17.6	5.4	11.8	26.3	42.7	60.4	95.6	132.1	182.2	297.0
1957	52.7	193.9	18.7	5.8	14.4	30.9	52.7	85.5	133.2	181.4	248.9	398.4
1956	48.3	183.2	15.6	5.3	12.9	29.0	49.1	79.3	122.9	160.9	238.1	334.4
1955	46.6	167.0	16.0	5.9	14.2	30.4	46.5	71.0	116.7	161.6	219.1	355.2
1954	46.4	174.3	17.7	6.4	14.7	27.7	46.9	78.8	111.5	146.7	214.3	360.0
1953	48.7	171.2	19.3	6.4	15.8	30.0	50.3	78.0	121.5	153.7	236.3	413.5
1952	49.2	185.0	19.1	6.1	15.8	30.6	48.7	79.9	123.4	156.8	243.8	387.8
1951	48.5	154.1	19.4	8.1	16.1	29.0	50.5	80.8	118.6	155.4	232.7	431.9
1950	46.8	161.0	21.2	6.3	16.6	28.2	49.6	73.2	112.3	163.2	246.9	408.6
ALL OTHER, MALE												
1959	50.0	198.3	20.8	5.8	12.6	32.2	50.6	74.3	115.7	194.4	227.8	406.9
1958	49.7	226.0	18.6	5.9	12.7	27.2	48.2	76.4	123.8	177.8	224.5	344.4
1957	61.8	223.3	21.9	6.0	14.5	31.3	56.5	96.8	170.1	240.6	315.6	511.5
1956	55.5	215.8	16.5	5.4	14.0	28.9	48.7	92.9	149.5	200.0	303.8	444.0
1955	53.5	194.3	19.2	6.4	13.8	30.8	47.5	78.3	143.8	208.0	276.8	441.7
1954	53.1	202.8	20.6	6.6	14.8	27.5	47.8	88.3	141.0	187.2	251.3	409.1
1953	56.4	190.2	21.2	7.1	16.2	30.5	54.2	88.8	151.5	195.9	284.2	585.7
1952	56.3	202.3	22.5	6.3	17.0	29.3	51.2	93.5	147.6	194.0	319.1	450.0
1951	53.9	175.5	18.8	9.0	17.8	28.3	51.8	87.5	136.3	193.0	267.9	515.8
1950	52.2	186.7	22.9	7.1	16.4	26.9	50.6	82.1	131.9	191.3	297.3	476.8
ALL OTHER, FEMALE												
1959	34.6	174.2	14.9	5.7	12.8	23.2	38.1	53.7	70.7	82.2	118.0	302.6
1958	33.8	157.2	16.7	4.9	10.9	25.5	37.7	45.2	68.6	90.1	143.8	271.1
1957	44.1	164.6	15.5	5.6	14.3	30.5	49.2	74.8	97.2	127.2	186.9	329.7
1956	41.5	150.7	14.7	5.2	11.9	29.1	49.5	66.2	96.7	124.2	176.1	258.3
1955	40.0	140.7	12.8	5.4	14.5	30.0	45.6	64.0	89.7	118.8	162.9	294.1
1954	40.1	146.0	14.8	6.1	14.5	27.8	46.1	69.5	81.9	109.2	177.8	337.5
1953	41.4	152.3	17.5	5.6	15.4	29.6	46.7	67.4	91.0	113.8	189.3	296.8
1952	42.4	168.0	15.6	5.8	14.8	31.8	46.3	66.6	98.4	121.8	172.4	344.8
1951	43.4	132.9	19.9	7.2	14.6	29.6	49.3	74.0	100.2	119.9	199.1	375.0
1950	41.7	135.5	19.4	5.5	16.8	29.4	48.6	64.3	92.3	134.9	197.2	359.9

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)												
BOTH SEXES												
1959	52.2	94.7	30.5	18.9	56.5	43.5	40.7	49.4	58.3	87.7	211.2	648.1
1958	52.3	96.9	31.0	19.5	56.2	42.5	41.5	47.2	57.6	89.4	221.3	655.1
1957	55.9	91.7	30.9	19.8	59.6	46.1	43.3	51.3	63.4	97.8	249.1	746.6
1956	56.7	91.3	31.0	20.1	63.6	45.9	43.7	51.2	63.5	101.3	250.7	758.7
1955	56.9	88.1	32.6	20.2	60.6	45.6	44.0	52.1	64.4	101.4	262.1	782.9
1954	55.9	98.2	33.5	20.4	56.8	44.3	41.9	50.4	62.5	101.4	267.0	780.0
1953	60.1	103.0	36.0	22.0	61.8	47.7	45.0	54.6	67.7	107.6	288.6	842.8
1952	61.8	109.4	37.2	22.4	61.2	48.3	47.2	55.9	71.1	111.7	300.3	897.7
1951	62.5	112.5	36.3	23.5	57.8	48.9	49.3	55.7	71.8	117.1	310.7	944.5
1950	60.6	114.2	36.8	22.7	54.8	45.7	45.7	53.0	70.8	116.9	315.7	972.6
MALE												
1959	73.2	102.9	35.0	26.5	94.8	72.7	66.2	76.7	88.7	119.6	234.9	602.8
1958	73.2	109.0	35.2	26.7	94.8	72.3	67.8	73.8	86.7	118.9	240.6	609.6
1957	78.2	101.3	34.3	27.5	101.1	78.0	70.0	80.2	95.1	133.1	269.8	672.7
1956	79.8	102.9	35.3	27.9	108.1	78.0	70.7	80.8	95.6	138.0	270.2	684.1
1955	79.8	99.4	36.6	28.0	102.8	77.1	71.4	82.6	96.9	138.9	279.7	667.9
1954	78.5	110.3	38.0	28.4	98.3	76.1	67.9	79.0	94.0	136.6	283.7	687.5
1953	85.0	116.0	41.3	30.5	106.0	82.0	73.7	86.6	103.5	144.7	307.6	734.6
1952	86.8	120.4	42.2	30.7	104.2	82.8	77.1	88.3	108.4	151.0	311.4	767.0
1951	88.2	123.9	41.7	33.0	98.1	83.8	81.5	88.8	108.4	158.5	314.3	821.9
1950	84.7	128.4	42.6	31.5	92.4	77.6	75.0	83.9	106.9	155.6	319.7	822.5
FEMALE												
1959	31.8	86.1	25.9	11.0	19.0	15.5	16.4	22.8	29.9	59.9	192.8	677.7
1958	32.0	84.3	26.7	12.1	18.7	13.9	16.3	21.3	30.2	63.3	206.1	685.5
1957	34.3	81.8	27.4	11.9	19.9	15.4	17.6	23.1	33.3	66.5	232.6	797.0
1956	34.2	79.2	26.6	12.0	21.2	15.2	17.7	22.2	33.0	68.4	235.2	810.6
1955	34.5	76.5	28.4	12.0	20.4	15.4	17.6	22.3	33.1	67.6	247.9	864.4
1954	33.9	85.6	28.9	12.1	18.2	14.0	16.7	22.3	32.0	69.5	253.3	846.1
1953	35.9	89.5	30.5	13.2	21.0	15.3	17.3	23.0	32.7	73.6	273.0	920.6
1952	37.4	97.8	32.1	13.9	21.5	15.7	18.2	23.9	34.4	75.7	291.1	991.3
1951	37.4	100.7	30.5	13.7	19.6	15.8	18.0	22.9	35.4	78.9	307.5	1,031.7
1950	36.7	99.6	30.9	13.5	18.2	15.3	17.1	22.3	34.6	80.9	312.3	1,077.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)												
WHITE, BOTH SEXES												
1959	50.5	76.6	26.6	17.4	56.4	41.0	37.6	46.4	55.6	86.5	215.6	676.1
1958	50.5	77.1	26.7	18.4	55.9	40.0	38.7	44.5	54.8	87.6	224.7	681.5
1957	54.2	73.7	27.0	18.5	59.6	42.9	40.3	48.0	60.7	96.5	253.2	778.6
1956	54.9	72.7	27.8	18.8	63.5	43.0	40.0	48.1	60.5	100.2	255.6	793.1
1955	55.3	72.9	28.3	19.1	60.5	42.7	41.0	48.7	61.7	100.2	267.3	817.8
1954	54.3	80.0	29.8	19.4	56.3	41.4	39.2	47.1	60.3	100.3	271.5	811.9
1953	58.2	86.2	32.0	21.0	61.1	44.4	41.5	50.8	65.1	106.9	293.1	878.1
1952	59.9	90.3	33.1	21.5	60.3	45.2	43.9	51.9	68.2	111.0	305.5	937.5
1951	61.0	95.0	32.7	22.3	57.2	46.5	46.1	52.2	69.0	116.7	316.9	980.7
1950	59.3	97.6	33.4	21.5	54.4	43.4	42.6	50.2	68.7	115.9	323.7	1,018.4
WHITE, MALE												
1959	70.5	84.9	31.0	24.8	94.5	68.6	61.1	71.8	84.2	118.0	239.5	631.0
1958	70.5	89.5	31.5	25.4	94.0	67.9	63.4	69.4	82.4	116.9	243.7	633.2
1957	75.4	82.9	30.8	25.9	101.0	72.7	65.3	75.0	91.1	131.5	273.8	697.4
1956	76.9	83.2	32.1	26.5	107.6	72.9	64.6	75.8	91.2	137.0	275.2	706.3
1955	77.3	84.2	32.8	26.8	102.5	72.6	66.5	77.0	93.1	137.5	286.0	693.9
1954	75.9	93.1	34.7	27.1	97.5	71.4	63.9	73.7	91.0	135.2	288.2	703.9
1953	81.9	99.0	37.3	29.2	104.7	76.6	68.2	80.8	99.6	143.8	310.5	759.7
1952	83.7	100.5	38.2	29.6	102.6	77.7	71.5	82.0	104.3	150.6	315.1	790.9
1951	85.7	108.0	38.3	31.3	97.4	80.0	76.5	83.4	104.5	157.6	319.2	846.0
1950	82.5	111.5	39.5	30.1	91.8	74.0	70.1	79.1	103.8	154.8	326.6	855.3
WHITE, FEMALE												
1959	31.0	68.0	22.0	9.8	19.0	14.2	15.0	21.6	28.9	59.0	197.1	705.4
1958	31.1	64.2	21.7	11.1	18.6	12.8	15.0	20.1	28.9	61.9	209.8	713.4
1957	33.5	64.2	23.0	10.8	19.5	13.9	16.2	21.5	31.9	65.4	236.9	833.9
1956	33.5	61.7	23.2	10.8	21.1	14.0	16.1	21.0	31.3	67.3	240.1	853.5
1955	33.8	61.1	23.6	11.0	20.2	13.9	16.3	20.9	31.5	66.6	252.5	905.5
1954	33.2	66.4	24.8	11.4	17.8	12.7	15.2	20.9	30.6	68.6	258.0	889.3
1953	35.1	72.7	26.4	12.5	20.4	13.7	15.6	21.4	31.5	73.3	279.0	963.6
1952	36.6	79.6	27.7	12.9	21.0	14.1	17.0	22.1	32.8	74.8	297.5	1,042.9
1951	36.8	81.4	26.9	12.9	18.8	14.6	16.4	21.2	33.8	79.0	315.0	1,076.8
1950	36.3	83.1	27.0	12.5	17.6	14.0	15.7	21.5	33.5	79.9	321.2	1,131.7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
STATISTICAL WORKSHEET												
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)												
ALL OTHER, BOTH SEXES												
1959	65.3	199.2	54.6	28.8	56.8	62.7	68.4	77.8	86.7	102.8	151.5	305.9
1958	66.2	215.0	57.4	27.4	58.6	63.0	65.4	73.4	87.7	110.2	174.3	337.9
1957	70.3	198.8	55.5	29.4	60.1	71.9	69.1	83.3	93.1	114.5	192.5	350.0
1956	70.7	203.1	52.1	28.9	64.6	69.6	75.7	80.9	97.8	114.3	181.3	344.3
1955	70.0	181.9	59.9	28.0	61.3	69.3	70.4	85.4	95.7	116.0	187.5	356.9
1954	69.0	212.9	57.6	27.5	60.1	68.1	65.1	82.7	89.2	115.1	202.5	374.5
1953	75.3	212.4	62.7	29.2	66.8	75.7	75.5	90.5	99.4	115.3	223.5	409.6
1952	77.4	235.8	65.2	29.4	67.4	74.9	76.2	94.6	107.3	120.1	224.3	406.1
1951	75.6	225.8	61.3	32.0	62.4	69.0	77.0	89.2	107.3	122.1	218.9	506.4
1950	71.2	226.5	61.7	31.1	57.9	65.5	72.4	80.3	96.3	130.2	194.3	428.7
ALL OTHER, MALE												
1959	94.5	209.3	59.8	38.5	97.3	106.6	112.2	124.3	134.8	137.4	177.1	286.2
1958	94.7	228.4	58.1	35.9	100.9	110.0	107.1	116.8	132.3	141.9	200.7	355.6
1957	101.0	213.6	56.8	39.5	101.7	123.6	111.9	130.5	138.5	152.0	219.3	376.9
1956	103.6	224.3	56.5	37.6	112.3	121.8	124.1	129.2	143.8	149.9	205.4	416.0
1955	100.9	195.8	61.2	36.7	105.5	116.9	115.0	136.7	139.7	155.0	196.8	350.0
1954	99.7	221.8	59.9	37.5	105.0	117.7	103.5	130.4	129.4	152.6	223.5	477.3
1953	110.9	228.2	68.3	40.4	116.1	130.0	122.3	143.0	149.6	155.8	268.4	414.3
1952	113.4	256.8	69.3	38.6	116.5	127.5	127.5	148.8	158.1	156.4	260.9	465.0
1951	110.1	229.1	66.6	45.0	104.0	118.2	125.7	140.2	156.2	168.8	250.9	521.1
1950	103.9	245.5	65.0	41.8	97.4	110.1	119.1	129.8	143.5	165.6	223.0	439.3
ALL OTHER, FEMALE												
1959	37.6	188.4	49.4	19.1	19.0	24.8	28.7	34.0	40.5	71.4	128.6	320.5
1958	39.2	201.7	56.7	19.0	19.6	22.3	27.2	32.4	44.8	81.2	150.3	334.2
1957	41.3	184.0	54.3	19.4	22.3	27.1	29.8	38.2	48.9	80.2	167.6	340.5
1956	39.5	182.1	47.6	20.2	21.7	24.3	31.0	34.4	52.6	81.2	158.7	294.4
1955	40.7	168.8	58.7	19.2	21.8	27.9	29.0	35.5	52.1	80.2	177.3	361.8
1954	40.1	204.0	55.4	17.5	21.1	25.1	29.5	36.1	49.1	80.6	181.0	315.6
1953	41.8	196.6	57.2	17.9	24.7	28.4	32.1	38.9	48.4	77.2	179.3	406.5
1952	43.5	215.1	61.2	20.2	24.9	28.8	28.5	41.0	55.0	86.0	189.7	365.5
1951	42.9	222.5	55.9	19.1	25.6	25.7	32.0	38.5	56.5	78.0	188.3	496.4
1950	39.9	207.6	58.3	20.4	22.1	25.8	29.0	30.9	47.8	94.6	166.0	421.2

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MOTOR VEHICLE ACCIDENTS (E810-E835)												
BOTH SEXES												
1959	21.5	7.8	9.5	7.9	38.2	24.9	19.5	22.0	24.8	32.3	42.2	36.7
1958	21.3	7.2	9.5	8.1	37.0	24.4	20.2	21.2	24.6	33.5	43.0	37.2
1957	22.7	8.4	9.4	8.0	39.8	26.7	21.2	22.8	27.4	34.6	46.4	38.1
1956	23.7	8.7	9.5	8.4	43.1	27.3	22.0	23.6	27.4	36.5	46.5	42.5
1955	23.4	8.7	10.5	8.0	41.1	27.0	21.9	22.8	28.0	35.3	47.1	41.1
1954	22.1	8.1	11.0	8.0	36.4	25.1	20.0	22.1	26.2	34.1	48.6	43.3
1953	24.0	9.4	12.0	8.5	39.3	26.9	21.8	23.7	28.4	38.0	51.6	49.9
1952	24.3	9.0	11.9	8.6	38.8	26.9	22.4	24.2	29.7	38.7	50.3	44.2
1951	24.1	8.7	11.4	9.2	36.1	26.6	22.6	24.2	29.4	39.5	52.2	47.6
1950	23.1	8.4	11.5	8.8	34.4	24.6	20.3	22.2	29.0	39.1	52.7	45.1
MALE												
1959	32.3	7.4	10.7	10.6	62.5	40.5	30.5	32.8	35.9	46.7	65.2	64.8
1958	32.2	7.9	10.9	10.8	60.8	40.5	31.7	31.3	35.5	48.3	66.3	68.0
1957	34.4	8.9	10.7	10.5	65.9	44.3	32.9	33.8	39.6	50.8	73.9	70.0
1956	36.1	9.6	10.4	11.2	71.5	45.3	34.4	35.6	40.2	53.7	74.7	74.3
1955	35.5	9.3	11.6	10.6	68.2	44.7	34.7	34.0	40.6	52.4	73.0	74.5
1954	33.6	9.2	12.5	10.4	61.6	42.2	31.3	32.9	38.2	50.3	77.8	71.1
1953	36.6	10.7	14.1	11.1	65.4	45.2	34.5	35.4	41.9	56.6	82.8	83.0
1952	37.2	9.4	13.9	11.3	64.4	45.3	35.4	36.0	43.8	59.8	80.4	72.9
1951	37.0	9.3	13.8	12.1	59.8	44.4	36.0	36.4	43.0	59.5	83.5	85.2
1950	35.4	9.1	13.0	11.9	56.7	40.8	32.5	33.6	43.1	59.1	85.0	78.1
FEMALE												
1959	11.0	8.3	8.3	5.1	14.4	9.9	9.1	11.6	14.4	19.7	24.3	18.4
1958	10.8	6.5	8.2	5.4	13.9	8.9	9.2	11.3	14.3	20.5	24.6	16.7
1957	11.3	7.9	8.1	5.4	14.9	9.8	9.9	12.1	15.7	20.1	24.5	16.4
1956	11.6	7.7	8.6	5.5	15.9	10.0	10.0	11.8	15.3	21.0	23.9	20.4
1955	11.6	8.2	9.4	5.2	15.5	10.2	9.6	11.8	15.9	19.8	26.2	17.4
1954	10.8	7.0	9.4	5.5	13.1	8.8	9.2	11.5	14.7	19.4	24.8	23.5
1953	11.7	8.1	9.9	5.8	15.3	9.7	9.7	12.2	15.1	21.0	26.0	26.1
1952	11.7	8.7	9.8	5.9	15.2	9.6	9.9	12.6	15.8	19.3	25.5	23.6
1951	11.5	8.1	8.9	6.2	13.6	9.8	9.6	12.1	15.9	21.0	26.0	20.8
1950	10.9	7.6	10.0	5.7	12.6	9.3	8.5	10.9	14.9	20.6	25.2	22.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MOTOR VEHICLE ACCIDENTS (E810-E835)												
WHITE, BOTH SEXES												
1959	21.4	8.2	9.4	7.7	39.3	24.1	18.5	20.9	24.3	32.5	43.4	37.2
1958	21.2	7.6	9.3	8.2	38.0	23.4	19.2	20.3	24.1	33.5	44.0	38.9
1957	22.5	8.5	9.1	7.9	41.1	25.5	20.2	21.7	26.5	34.7	46.5	38.4
1956	23.3	8.2	9.2	8.3	44.2	25.9	20.5	22.4	26.4	36.6	47.1	43.8
1955	23.1	8.7	10.2	8.0	42.1	25.9	20.8	21.7	27.3	35.3	47.9	41.3
1954	21.8	7.7	10.8	8.0	37.4	24.0	19.0	20.8	25.4	34.4	49.2	42.8
1953	23.6	9.2	11.8	8.4	40.2	25.6	20.5	22.5	27.8	38.2	51.8	50.4
1952	24.0	9.2	11.7	8.7	39.5	25.7	21.4	22.8	29.0	39.1	50.8	43.3
1951	23.9	8.8	11.4	9.1	36.7	25.6	21.4	23.1	28.7	39.6	52.7	48.0
1950	22.9	8.5	11.7	8.9	35.3	23.8	19.4	21.2	28.5	39.3	53.2	45.7
WHITE, MALE												
1959	31.9	7.6	10.5	10.3	64.2	39.0	28.7	30.7	34.6	46.9	67.0	64.7
1958	31.8	8.3	10.7	10.7	62.3	38.7	29.9	29.7	34.2	47.8	68.1	71.2
1957	33.9	9.0	10.3	10.4	68.0	42.2	31.1	31.9	38.0	50.8	73.9	70.1
1956	35.3	8.9	9.9	11.2	73.1	42.8	31.8	33.5	38.6	53.4	75.7	75.2
1955	35.0	9.2	11.3	10.6	69.6	42.6	32.8	32.0	39.3	52.3	74.6	74.8
1954	33.1	8.6	12.4	10.4	63.1	40.3	29.6	30.6	36.7	50.4	79.1	69.1
1953	35.8	10.7	13.6	10.9	66.6	42.8	32.2	33.1	40.7	56.3	83.1	83.2
1952	36.5	9.5	13.6	11.2	65.6	43.2	33.4	33.6	42.4	60.3	81.2	70.4
1951	36.5	9.7	13.8	12.0	61.0	42.4	34.0	34.7	41.8	59.4	84.3	85.7
1950	35.1	9.1	13.2	12.0	58.3	39.1	30.9	31.6	41.9	59.1	86.4	79.3
WHITE, FEMALE												
1959	11.1	8.8	8.2	5.0	14.8	9.6	8.8	11.4	14.6	20.0	25.1	19.4
1958	10.9	6.8	7.9	5.6	14.3	8.6	8.9	11.2	14.5	20.9	25.2	17.6
1957	11.3	8.1	7.9	5.3	15.1	9.2	9.6	11.8	15.5	20.4	24.9	16.8
1956	11.6	7.4	8.5	5.4	16.4	9.6	9.6	11.5	14.8	21.6	24.5	21.9
1955	11.5	8.2	9.2	5.2	15.8	9.8	9.1	11.5	15.8	20.0	26.7	17.6
1954	10.8	6.7	9.1	5.5	13.3	8.4	8.7	11.2	14.5	19.9	25.2	23.9
1953	11.7	7.6	9.9	5.7	15.6	9.2	9.1	12.0	15.3	21.7	26.4	26.7
1952	11.7	8.8	9.8	6.0	15.3	9.0	9.8	12.2	15.9	19.7	25.8	23.9
1951	11.5	7.9	8.9	6.2	13.5	9.5	9.1	11.7	15.8	21.3	26.5	21.1
1950	10.9	7.8	10.1	5.6	12.6	9.0	8.1	10.8	15.0	20.9	25.4	22.3

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
MOTOR VEHICLE ACCIDENTS (E810-E835)												
ALL OTHER, BOTH SEXES												
1959	22.5	5.7	10.3	9.1	29.8	31.2	28.4	32.6	30.6	29.7	26.6	30.9
1958	22.2	5.4	10.8	7.6	29.6	31.7	28.7	29.6	29.9	33.7	28.8	*
1957	24.7	7.7	11.2	8.5	30.8	36.5	29.7	33.1	37.1	32.9	45.0	34.4
1956	26.8	11.5	11.5	8.9	34.9	38.2	34.8	34.7	39.3	34.5	37.3	*
1955	25.7	9.1	12.3	7.9	34.2	36.6	31.6	33.5	36.5	35.6	35.5	37.9
1954	24.5	11.0	12.2	7.9	29.8	33.9	28.9	34.3	35.8	31.2	39.8	49.1
1953	27.2	10.9	13.6	9.5	32.8	38.3	33.8	35.9	34.9	35.4	49.1	44.2
1952	26.8	8.1	12.9	8.1	33.5	37.3	31.2	37.7	38.7	33.7	44.2	55.1
1951	26.4	7.9	11.3	9.4	31.0	35.5	32.8	34.2	38.0	38.5	45.2	42.6
1950	24.2	7.9	10.6	8.6	27.4	32.1	28.5	31.6	35.7	37.2	44.3	*
ALL OTHER, MALE												
1959	35.4	*	12.0	12.5	49.5	53.2	47.2	52.9	49.5	44.4	42.4	*
1958	35.4	*	11.9	11.1	50.1	55.6	47.4	47.3	48.8	54.0	43.2	*
1957	38.4	8.7	13.3	11.2	50.3	62.1	48.4	51.9	57.0	51.2	73.3	*
1956	42.6	13.6	13.9	11.2	59.7	67.3	57.3	55.3	58.4	56.9	61.5	*
1955	40.1	10.0	13.7	10.3	57.4	63.1	51.3	53.1	55.8	54.4	52.8	*
1954	38.3	13.3	13.2	10.4	50.3	59.1	45.9	55.3	54.8	50.0	60.5	95.5
1953	43.4	10.7	17.4	12.7	55.6	66.7	54.5	57.8	56.1	59.9	79.8	*
1952	42.7	*	15.8	11.4	55.5	63.6	53.1	58.9	61.4	54.0	69.1	105.0
1951	41.7	*	13.5	13.1	50.5	62.1	53.7	52.9	57.6	60.9	73.6	*
1950	38.2	*	11.7	10.8	44.3	55.4	46.9	52.0	57.0	58.8	66.4	*
ALL OTHER, FEMALE												
1959	10.2	*	8.6	5.7	11.5	12.2	11.3	13.5	12.3	16.2	12.4	*
1958	9.7	*	9.6	4.1	10.7	11.1	11.6	12.8	11.7	15.1	15.7	*
1957	11.8	*	9.1	5.9	13.2	14.4	12.5	15.1	17.7	16.1	18.6	*
1956	11.9	9.5	9.2	6.6	12.5	12.9	13.9	14.8	20.6	13.8	14.5	*
1955	12.1	8.4	10.9	5.5	13.5	13.6	13.3	14.4	17.4	18.3	18.9	*
1954	11.4	8.8	11.2	5.4	11.9	12.0	13.2	13.9	16.8	13.9	19.8	*
1953	11.9	11.1	9.8	6.2	13.3	13.6	14.5	14.3	13.3	12.5	19.8	*
1952	11.8	*	10.0	4.8	14.5	14.3	10.9	16.8	15.3	14.6	20.7	*
1951	12.0	9.5	9.1	5.6	13.8	12.0	13.5	15.6	17.7	17.3	18.0	*
1950	10.8	*	9.4	6.4	12.1	11.4	11.5	11.1	13.9	15.4	22.5	*

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER ACCIDENTS (E800-E802, E840-E962)												
BOTH SEXES												
1959	30.7	86.8	21.0	11.0	18.3	18.6	21.2	27.4	33.5	55.4	169.0	611.4
1958	30.9	89.6	21.5	11.4	19.2	18.2	21.3	26.0	33.1	55.9	178.3	617.9
1957	33.2	83.3	21.5	11.8	19.8	19.3	22.1	28.5	36.0	63.3	202.7	708.4
1956	33.0	82.6	21.5	11.7	20.6	18.6	21.7	27.6	36.1	64.8	204.2	716.2
1955	33.5	79.4	22.0	12.2	19.4	18.5	22.1	29.4	36.4	66.1	215.0	741.9
1954	33.8	90.0	22.6	12.4	20.4	19.2	21.8	28.4	36.3	67.3	218.4	736.6
1953	36.1	93.6	24.0	13.5	22.5	20.8	23.1	30.9	39.3	69.5	237.0	792.9
1952	37.5	100.3	25.3	13.8	22.3	21.4	24.8	31.7	41.4	73.0	249.9	853.5
1951	38.4	103.8	24.9	14.3	21.8	22.3	26.7	31.5	42.4	77.6	258.5	896.9
1950	37.5	105.8	25.3	13.8	20.4	21.1	25.4	30.9	41.8	77.8	263.0	927.5
MALE												
1959	40.9	95.5	24.3	16.0	32.3	32.1	35.7	44.0	52.7	72.8	169.7	538.0
1958	41.0	101.1	24.3	15.9	34.0	31.8	36.2	42.5	51.2	70.6	174.3	541.6
1957	43.7	92.3	23.6	17.0	35.2	33.7	37.1	46.4	55.5	82.3	196.0	602.7
1956	43.6	93.3	24.9	16.7	36.6	32.7	36.3	45.2	55.4	84.4	195.5	609.8
1955	44.3	90.1	25.0	17.5	34.7	32.5	36.7	48.6	56.3	86.5	206.7	593.4
1954	44.8	101.1	25.5	17.9	36.8	33.8	36.7	46.1	55.9	86.2	205.9	616.4
1953	48.3	105.2	27.2	19.4	40.7	36.8	39.2	51.2	61.6	88.1	224.7	651.6
1952	49.7	111.1	28.3	19.4	39.8	37.5	41.7	52.4	64.6	91.2	231.0	694.1
1951	51.2	114.6	28.0	20.9	38.3	39.5	45.4	52.4	65.4	99.0	230.8	736.7
1950	49.3	119.3	29.5	19.7	35.7	36.9	42.5	50.3	63.8	96.5	234.7	744.4
FEMALE												
1959	20.8	77.9	17.7	5.9	4.6	5.6	7.4	11.3	15.4	40.2	168.5	659.3
1958	21.2	77.7	18.5	6.7	4.8	5.1	7.1	10.0	16.0	42.9	181.5	668.8
1957	23.0	73.9	19.3	6.5	5.0	5.6	7.7	11.0	17.6	46.4	208.1	780.6
1956	22.6	71.5	18.0	6.5	5.2	5.2	7.6	10.4	17.7	47.3	211.2	790.2
1955	23.0	68.3	19.0	6.8	4.9	5.2	8.1	10.5	17.2	47.8	221.7	847.0
1954	23.1	78.6	19.5	6.6	5.1	5.2	7.5	10.9	17.3	50.1	228.5	822.6
1953	24.2	81.5	20.6	7.4	5.7	5.6	7.7	10.8	17.6	52.6	247.0	894.5
1952	25.6	89.2	22.2	8.0	6.2	6.2	8.3	11.3	18.5	56.3	265.6	967.7
1951	25.9	92.6	21.6	7.5	6.1	6.1	8.4	10.8	19.5	57.9	281.5	1,010.8
1950	25.8	91.9	20.9	7.8	5.6	6.0	8.6	11.5	19.7	60.3	287.1	1,055.1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER ACCIDENTS (E800-E802, E840-E962)												
WHITE, BOTH SEXES												
1959	29.2	68.4	17.2	9.7	17.1	16.9	19.0	25.5	31.3	54.0	172.2	638.9
1958	29.3	69.5	17.4	10.2	17.9	16.5	19.5	24.2	30.8	54.1	180.7	642.6
1957	31.7	65.2	17.8	10.5	18.5	17.4	20.1	26.2	34.2	61.7	206.7	740.2
1956	31.6	64.5	18.5	10.5	19.3	17.1	19.5	25.7	34.1	63.6	208.5	749.3
1955	32.2	64.2	18.1	11.1	18.3	16.8	20.2	27.0	34.4	65.0	219.4	776.4
1954	32.5	72.4	19.1	11.4	19.0	17.4	20.2	26.3	34.9	65.9	222.3	769.0
1953	34.6	77.0	20.2	12.7	20.9	18.8	21.0	28.4	37.3	68.7	241.3	827.7
1952	35.9	81.1	21.3	12.8	20.8	19.5	22.4	29.1	39.2	71.9	254.7	894.2
1951	37.1	86.1	21.4	13.2	20.5	21.0	24.6	29.0	40.3	77.1	264.2	932.7
1950	36.4	89.1	21.7	12.6	19.1	19.7	23.2	29.0	40.2	76.6	270.5	972.7
WHITE, MALE												
1959	38.7	77.2	20.5	14.5	30.3	29.6	32.4	41.1	49.6	71.1	172.5	566.3
1958	38.7	81.2	20.8	14.7	31.8	29.1	33.5	39.7	48.2	69.1	175.6	562.0
1957	41.5	73.9	20.4	15.4	33.0	30.5	34.2	43.1	53.1	80.7	199.9	627.3
1956	41.6	74.3	22.2	15.4	34.5	30.2	32.9	42.3	52.7	83.6	199.5	631.1
1955	42.3	75.0	21.6	16.2	32.9	30.0	33.7	45.0	53.8	85.3	211.5	619.0
1954	42.8	84.5	22.4	16.7	34.4	31.0	34.3	43.1	54.3	84.9	209.1	634.8
1953	46.1	88.3	23.7	18.3	38.1	33.8	35.9	47.6	58.9	87.5	227.4	676.5
1952	47.2	91.0	24.6	18.4	37.0	34.5	38.1	48.4	62.0	90.3	233.9	720.6
1951	49.2	98.3	24.5	19.4	36.4	37.6	42.5	48.7	62.7	98.2	234.9	760.3
1950	47.4	102.3	26.3	18.1	33.4	34.9	39.2	47.4	61.9	95.7	240.3	776.0
WHITE, FEMALE												
1959	19.9	59.2	13.8	4.8	4.1	4.6	6.2	10.3	14.2	39.0	172.0	685.9
1958	20.1	57.4	13.8	5.5	4.3	4.2	6.1	9.0	14.4	41.0	184.6	695.8
1957	22.2	56.1	15.1	5.4	4.4	4.6	6.6	9.7	16.3	44.9	212.0	817.1
1956	21.9	54.3	14.7	5.4	4.6	4.4	6.5	9.4	16.5	45.6	215.6	831.6
1955	22.3	52.9	14.5	5.8	4.5	4.1	7.2	9.4	15.8	46.6	225.9	888.0
1954	22.4	59.7	15.6	5.9	4.6	4.2	6.5	9.7	16.1	48.7	232.8	865.4
1953	23.5	65.1	16.5	6.8	4.8	4.5	6.5	9.4	16.2	51.5	252.6	936.9
1952	24.9	70.8	17.9	6.9	5.6	5.1	7.2	9.9	16.9	55.1	271.8	1,019.0
1951	25.3	73.5	18.0	6.7	5.3	5.1	7.2	9.5	18.0	57.7	288.5	1,055.7
1950	25.4	75.4	17.0	6.9	5.0	5.0	7.6	10.6	18.5	59.0	295.9	1,109.4

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
OTHER ACCIDENTS (E800-E802, E840-E962)												
ALL OTHER, BOTH SEXES												
1959	42.8	193.5	44.3	19.7	27.0	31.5	40.0	45.2	56.1	73.1	124.9	275.0
1958	44.0	209.7	46.6	19.8	29.0	31.3	36.6	43.8	57.8	76.6	145.5	321.2
1957	45.6	191.1	44.3	20.9	29.3	35.4	39.4	50.2	56.0	81.6	147.5	315.6
1956	43.9	191.6	40.5	20.0	29.7	31.4	41.0	46.3	58.4	79.8	144.0	316.4
1955	44.2	172.8	47.6	20.1	27.0	32.6	38.8	51.9	59.2	80.4	152.0	319.0
1954	44.6	201.8	45.4	19.7	30.4	34.2	36.2	48.3	53.4	83.9	162.7	325.5
1953	48.1	201.5	49.1	19.7	34.0	37.4	41.8	54.6	64.6	79.9	174.4	365.4
1952	50.6	227.7	52.4	21.3	33.9	37.6	45.0	56.9	68.6	86.4	180.1	351.0
1951	49.2	217.9	49.9	22.6	31.4	33.5	44.2	55.0	69.3	83.7	173.7	463.8
1950	47.0	218.6	51.1	22.5	30.5	33.4	43.8	48.8	60.5	93.1	150.0	390.8
ALL OTHER, MALE												
1959	59.1	203.3	47.8	26.0	47.8	53.4	65.0	71.4	85.2	92.9	134.7	220.7
1958	59.3	222.5	46.2	24.8	50.8	54.4	59.6	69.5	83.5	87.9	157.6	322.2
1957	62.6	204.9	43.4	28.3	51.4	61.6	63.5	78.6	81.5	100.8	145.9	307.7
1956	61.1	210.7	42.7	26.4	52.5	54.5	66.8	73.9	85.4	93.0	143.8	352.0
1955	60.8	185.8	47.5	26.5	48.0	53.8	63.7	83.6	83.9	100.6	144.0	279.2
1954	61.4	208.5	46.7	27.1	54.7	58.5	57.6	75.1	74.6	102.6	163.0	381.8
1953	67.5	217.5	50.8	27.7	60.5	63.4	67.8	85.2	93.5	95.9	188.6	333.3
1952	70.7	248.2	53.5	27.2	61.0	64.0	74.5	89.9	96.7	102.4	191.8	360.0
1951	68.4	222.7	53.1	31.8	53.5	56.1	72.0	87.3	98.6	108.0	177.4	442.1
1950	65.7	236.6	53.3	31.0	53.1	54.7	72.3	77.8	86.4	106.8	156.6	375.0
ALL OTHER, FEMALE												
1959	27.4	183.1	40.8	13.3	7.5	12.6	17.4	20.5	28.2	55.1	116.1	315.4
1958	29.5	196.9	47.0	14.9	8.9	11.2	15.6	19.5	33.1	66.1	134.6	328.9
1957	29.5	177.4	45.2	13.5	9.1	12.7	17.3	23.1	31.2	64.1	149.0	329.7
1956	27.6	172.6	38.4	13.6	9.2	11.4	17.1	19.6	32.0	67.4	144.2	291.7
1955	28.6	160.5	47.7	13.7	8.3	14.2	15.7	21.1	34.7	61.8	158.3	347.1
1954	28.7	195.2	44.2	12.2	9.2	13.1	16.3	22.2	32.3	66.7	161.1	296.9
1953	29.9	185.5	47.4	11.7	11.4	14.8	17.6	24.6	35.1	64.8	159.5	387.1
1952	31.6	207.6	51.2	15.5	10.5	14.5	17.7	24.3	39.7	71.4	169.0	344.8
1951	30.9	213.1	46.8	13.5	11.9	13.7	18.5	22.9	38.8	60.7	170.3	478.6
1950	29.1	200.8	48.9	14.0	10.0	14.4	17.5	19.8	33.9	79.2	143.5	402.0

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SUICIDE (E963, E970-E979)												
BOTH SEXES												
1959	10.6	0.2	4.9	9.9	13.6	19.8	24.3	24.8	27.8	25.7
1958	10.7	0.2	4.8	9.8	13.7	20.7	24.1	25.1	27.7	25.8
1957	9.8	0.2	4.0	8.7	12.7	18.2	22.4	23.6	26.5	26.4
1956	10.0	0.2	4.0	8.5	12.1	18.5	24.2	25.2	28.0	23.5
1955	10.2	0.1	4.1	8.4	12.3	19.6	24.8	25.0	27.5	27.9
1954	10.1	0.1	4.2	8.7	12.5	19.3	23.9	25.1	25.6	25.1
1953	10.1	0.2	4.4	8.5	12.6	18.7	22.4	25.4	28.0	27.9
1952	10.0	0.2	4.2	8.5	12.4	18.3	22.5	25.6	27.9	31.2
1951	10.4	0.2	4.4	8.6	13.3	18.8	23.2	27.1	27.7	30.5
1950	11.4	0.2	4.5	9.1	14.3	20.9	26.8	29.6	31.1	28.8
MALE												
1959	16.6	0.4	7.7	14.4	20.5	31.0	39.4	42.6	54.2	59.4
1958	16.8	0.4	7.4	14.2	21.1	32.1	39.6	43.3	54.9	54.9
1957	15.4	0.3	6.4	12.7	19.1	28.6	35.9	41.4	52.5	58.8
1956	15.7	0.3	6.3	12.7	18.2	28.3	39.1	43.6	54.8	51.1
1955	16.0	0.2	6.3	12.4	18.8	29.7	40.3	42.8	51.7	57.9
1954	16.3	0.2	6.7	13.4	18.9	31.0	39.2	43.7	49.9	50.7
1953	16.1	0.3	6.6	12.7	19.5	29.3	36.4	44.7	53.5	59.9
1952	15.8	0.2	6.6	12.4	18.5	27.9	36.5	44.4	53.6	66.3
1951	16.2	0.3	6.6	12.6	19.5	28.6	37.6	47.2	51.2	66.0
1950	17.8	0.3	6.5	13.4	21.3	32.0	43.6	50.5	58.3	58.3
FEMALE												
1959	4.7	*	2.1	5.4	7.0	8.8	10.1	9.3	7.2	3.7
1958	4.7	*	2.3	5.6	6.7	9.7	9.4	9.0	6.4	6.4
1957	4.3	*	1.8	4.8	6.6	8.0	9.6	7.8	5.9	4.3
1956	4.4	*	1.9	4.5	6.3	9.0	9.9	8.9	6.5	4.3
1955	4.6	*	2.0	4.6	6.1	9.6	9.8	8.9	7.9	6.7
1954	4.1	*	1.8	4.3	6.3	7.7	9.0	8.3	5.9	6.8
1953	4.3	*	2.3	4.5	6.0	8.3	8.7	7.8	7.1	5.0
1952	4.4	*	2.0	4.7	6.4	8.7	8.7	8.4	6.7	6.0
1951	4.7	*	2.3	4.8	7.2	9.1	8.8	8.6	7.9	*
1950	5.1	*	2.6	4.9	7.5	9.9	9.9	10.1	8.1	8.2

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SUICIDE (E963, E970-E979)												
WHITE, BOTH SEXES												
1959	11.3	0.3	5.0	10.0	14.4	21.0	25.7	26.2	29.0	27.3
1958	11.5	0.2	5.0	10.0	14.5	22.1	25.5	26.5	29.2	27.3
1957	10.5	0.2	4.1	8.8	13.4	19.4	23.7	24.8	27.8	27.7
1956	10.8	0.2	4.1	8.8	12.8	19.8	25.6	26.8	29.5	25.0
1955	11.0	0.1	4.1	8.7	13.1	20.9	26.3	26.5	28.9	29.6
1954	10.9	0.1	4.3	8.9	13.3	20.5	25.3	26.5	26.9	26.8
1953	10.8	0.2	4.6	8.8	13.4	19.9	23.7	27.0	29.2	29.7
1952	10.8	0.1	4.4	8.7	13.2	19.6	23.6	27.3	29.4	32.9
1951	11.1	0.2	4.5	8.8	14.0	20.0	24.4	28.9	29.0	32.9
1950	12.2	0.2	4.7	9.4	15.2	22.3	28.3	31.1	32.9	30.6
WHITE, MALE												
1959	17.7	0.5	7.9	14.4	21.5	32.8	41.8	45.0	56.6	63.8
1958	18.0	0.4	7.6	14.3	22.1	34.0	41.9	45.8	58.3	58.5
1957	16.5	0.4	6.5	12.7	20.1	30.3	38.1	43.5	55.3	62.7
1956	16.9	0.3	6.4	12.9	19.1	30.2	41.5	46.4	58.1	54.3
1955	17.2	0.2	6.2	12.7	19.8	31.7	42.8	45.6	54.7	61.2
1954	17.5	0.2	6.9	13.5	19.9	33.0	41.6	46.2	52.6	54.3
1953	17.2	0.4	7.0	13.0	20.5	31.0	38.4	47.5	55.9	63.4
1952	16.9	0.2	6.9	12.6	19.6	29.8	38.4	47.2	56.7	70.0
1951	17.3	0.3	6.8	12.8	20.5	30.3	39.7	50.3	53.8	70.9
1950	19.0	0.3	6.6	13.8	22.4	34.1	45.9	53.2	61.9	61.9
WHITE, FEMALE												
1959	5.0	*	2.1	5.7	7.5	9.5	10.6	9.8	7.6	*
1958	5.1	*	2.4	5.9	7.1	10.4	10.0	9.6	6.6	6.7
1957	4.6	*	1.8	5.0	7.1	8.8	10.1	8.3	6.2	*
1956	4.8	*	2.0	4.7	6.7	9.6	10.5	9.4	6.8	4.6
1955	4.9	*	2.0	4.9	6.6	10.3	10.4	9.4	8.3	7.2
1954	4.5	*	1.8	4.4	6.8	8.2	9.5	8.7	6.3	7.1
1953	4.6	*	2.3	4.8	6.5	9.0	9.3	8.3	7.5	5.4
1952	4.7	*	2.1	5.0	6.9	9.4	9.2	9.0	7.0	6.3
1951	5.0	*	2.3	5.0	7.7	9.8	9.2	9.1	8.3	*
1950	5.5	*	2.7	5.2	8.2	10.5	10.7	10.6	8.4	8.9

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
SUICIDE (E963, E970-E979)												
ALL OTHER, BOTH SEXES												
1959	4.6	*	4.3	8.7	6.6	8.2	9.4	8.6	11.8	*
1958	4.4	*	3.5	7.8	7.3	7.8	9.0	8.0	7.2	*
1957	4.0	*	3.4	7.4	6.3	6.4	7.9	9.3	8.6	*
1956	3.8	*	3.4	6.4	6.6	6.4	7.7	6.2	*	*
1955	3.8	*	4.0	6.0	5.5	6.6	7.6	6.5	*	*
1954	4.1	*	3.4	7.4	6.1	7.1	7.5	7.9	*	*
1953	3.8	*	3.0	5.8	5.8	7.2	7.0	7.0	11.1	*
1952	3.7	*	2.7	6.5	5.5	5.7	8.3	5.5	*	*
1951	4.1	*	3.7	6.4	6.6	6.8	7.5	6.1	*	*
1950	4.3	*	3.4	6.3	6.6	7.9	9.1	8.7	*	*
ALL OTHER, MALE												
1959	7.5	*	6.6	14.4	11.0	13.9	14.8	14.9	22.9	*
1958	7.1	*	5.4	12.6	12.3	12.8	15.3	14.2	*	*
1957	6.8	*	5.6	13.0	10.6	11.8	11.9	17.4	15.6	*
1956	6.1	*	5.8	10.8	10.5	9.8	12.2	10.0	*	*
1955	6.1	*	6.5	9.6	9.6	10.5	12.7	10.8	*	*
1954	6.8	*	5.5	12.8	10.6	11.4	12.2	13.6	*	*
1953	6.4	*	4.2	10.1	10.3	12.6	12.0	12.2	21.9	*
1952	6.1	*	4.5	10.7	8.9	9.9	14.3	11.0	*	*
1951	6.6	*	5.2	10.7	10.8	11.4	12.0	10.1	*	*
1950	7.0	*	5.3	10.1	11.3	11.7	16.8	15.0	*	*
ALL OTHER, FEMALE												
1959	1.9	*	2.2	3.7	2.7	2.9	4.3	*	*	*
1958	1.8	*	1.7	3.6	2.8	3.1	*	*	*	*
1957	1.4	*	*	2.5	2.3	*	4.0	*	*	*
1956	1.6	*	*	2.6	3.0	3.0	3.3	*	*	*
1955	1.5	*	1.8	2.8	1.7	2.8	*	*	*	*
1954	1.5	*	1.5	2.8	2.0	2.8	*	*	*	*
1953	1.3	*	2.0	2.1	*	*	*	*	*	*
1952	1.3	*	*	2.7	2.5	*	*	*	*	*
1951	1.7	*	2.4	2.6	2.7	*	*	*	*	*
1950	1.7	*	1.7	2.8	2.2	4.0	*	*	*	*

* Figure does not meet standard of reliability or precision

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HOMICIDE (E964, E980-E985)												
BOTH SEXES												
1959	4.6	3.7	0.9	0.5	5.9	9.3	8.0	5.9	4.0	2.9	2.5	2.6
1958	4.5	3.3	0.7	0.5	5.8	8.7	7.4	6.3	4.0	2.9	2.6	3.3
1957	4.5	3.1	0.6	0.4	5.8	9.0	7.5	5.8	4.0	3.0	2.2	*
1956	4.6	3.3	0.7	0.4	6.1	9.0	7.6	5.6	4.1	2.8	2.2	*
1955	4.5	3.4	0.5	0.4	5.5	8.8	7.7	5.9	4.0	2.4	2.5	3.0
1954	4.8	3.7	0.6	0.5	6.0	9.1	8.2	6.5	3.5	2.8	2.7	*
1953	4.8	3.1	0.7	0.5	5.9	9.0	8.7	6.1	3.4	2.8	2.3	3.6
1952	5.2	3.7	0.6	0.5	6.2	10.0	9.1	6.1	3.8	3.0	2.7	*
1951	4.9	4.3	0.6	0.4	5.9	9.1	8.2	5.9	3.9	2.7	2.4	*
1950	5.3	4.4	0.6	0.5	6.3	9.9	8.8	6.1	4.0	3.2	2.6	*
MALE												
1959	7.0	4.2	0.9	0.6	9.0	13.9	12.2	9.4	6.6	4.6	3.9	*
1958	6.8	3.4	0.7	0.6	8.8	13.7	11.2	9.7	6.5	4.7	3.2	*
1957	6.8	3.3	0.5	0.5	8.9	14.1	11.4	9.2	6.3	5.0	3.4	*
1956	6.9	3.1	0.7	0.6	9.4	13.9	11.8	8.5	6.5	4.4	3.3	*
1955	6.9	3.7	0.5	0.5	8.5	13.8	12.0	9.2	6.6	4.0	3.8	*
1954	7.4	4.0	0.6	0.6	9.3	14.5	12.9	10.1	5.9	4.4	4.1	*
1953	7.5	3.1	0.7	0.5	9.6	14.0	13.8	9.8	5.5	4.4	3.9	*
1952	8.1	3.9	0.8	0.5	9.9	15.8	14.3	9.8	6.2	5.1	4.6	*
1951	7.5	3.9	0.6	0.4	9.0	14.2	13.0	9.4	6.5	4.4	3.5	*
1950	8.1	4.5	0.5	0.6	9.6	15.5	13.8	9.9	6.5	5.3	4.0	*
FEMALE												
1959	2.3	3.2	0.8	0.5	2.8	4.8	4.0	2.5	1.6	1.3	1.3	*
1958	2.3	3.3	0.8	0.4	3.0	3.9	3.8	2.9	1.6	1.4	2.1	*
1957	2.2	2.8	0.6	0.3	2.8	4.1	3.8	2.6	1.8	1.1	1.2	*
1956	2.3	3.5	0.7	0.3	2.9	4.3	3.6	2.7	1.7	1.3	1.3	*
1955	2.1	3.1	0.6	0.3	2.7	4.0	3.5	2.7	1.5	0.9	1.5	*
1954	2.3	3.3	0.6	0.4	2.9	4.0	3.6	3.0	1.1	1.3	1.5	*
1953	2.3	3.2	0.7	0.4	2.6	4.2	3.7	2.5	1.5	1.4	*	*
1952	2.3	3.5	0.5	0.4	2.8	4.4	4.1	2.5	1.4	1.1	1.0	*
1951	2.4	4.6	0.6	0.4	3.0	4.3	3.6	2.5	1.4	1.1	1.5	*
1950	2.4	4.2	0.7	0.5	3.1	4.5	3.8	2.3	1.4	1.3	1.4	*

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HOMICIDE (E964, E980-E985)												
WHITE, BOTH SEXES												
1959	2.5	3.1	0.8	0.5	2.8	3.9	3.8	3.4	2.7	2.2	2.0	*
1958	2.4	2.7	0.6	0.4	2.8	3.5	3.5	3.5	2.6	2.3	2.3	3.0
1957	2.3	2.7	0.4	0.3	2.6	3.7	3.5	3.0	2.7	2.1	1.7	*
1956	2.3	2.6	0.6	0.4	2.7	3.6	3.3	3.0	2.7	2.0	1.9	*
1955	2.3	2.9	0.4	0.4	2.4	3.5	3.8	3.2	2.6	1.8	2.3	*
1954	2.4	3.0	0.5	0.4	2.7	3.6	3.9	3.6	2.3	2.0	2.1	*
1953	2.4	2.9	0.6	0.4	2.5	3.7	3.9	3.5	2.4	2.1	2.0	3.1
1952	2.5	3.3	0.6	0.4	2.6	3.7	4.0	3.5	2.5	2.4	2.1	*
1951	2.5	3.9	0.5	0.4	2.4	3.7	3.8	3.3	2.7	2.2	2.1	*
1950	2.6	4.1	0.5	0.4	2.5	3.6	4.3	3.5	2.8	2.6	2.2	*
WHITE, MALE												
1959	3.5	3.2	0.8	0.5	4.2	5.5	5.4	5.1	4.3	3.4	3.0	*
1958	3.4	2.5	0.5	0.5	4.0	5.5	4.8	5.1	4.1	3.5	2.6	*
1957	3.2	3.1	0.3	0.4	3.9	5.6	4.8	4.3	4.0	3.4	2.5	*
1956	3.3	2.5	0.6	0.5	4.0	5.4	4.8	4.3	4.3	3.3	2.7	*
1955	3.4	3.3	0.4	0.4	3.5	5.2	5.7	4.6	4.1	3.0	3.4	*
1954	3.5	3.0	0.5	0.4	3.9	5.4	5.7	5.1	3.6	3.1	3.3	*
1953	3.5	2.7	0.6	0.4	3.8	5.4	5.7	5.2	3.7	3.1	3.3	*
1952	3.7	3.7	0.7	0.5	3.9	5.6	5.8	5.3	4.0	3.9	3.6	*
1951	3.6	3.6	0.4	0.4	3.6	5.5	5.6	4.9	4.3	3.5	3.0	*
1950	3.9	4.3	0.4	0.4	3.7	5.4	6.4	5.5	4.4	4.1	3.5	*
WHITE, FEMALE												
1959	1.4	2.9	0.7	0.4	1.4	2.4	2.2	1.8	1.1	1.1	1.3	*
1958	1.4	2.8	0.7	0.3	1.6	1.7	2.2	1.9	1.2	1.2	2.1	*
1957	1.3	2.2	0.5	0.3	1.4	1.9	2.2	1.7	1.5	0.9	1.1	*
1956	1.3	2.6	0.6	0.2	1.4	1.9	1.8	1.8	1.2	1.0	1.3	*
1955	1.2	2.5	0.4	0.3	1.3	1.7	1.9	1.9	1.1	0.7	1.4	*
1954	1.4	3.0	0.6	0.4	1.5	1.9	2.1	2.0	1.0	1.0	1.1	*
1953	1.4	3.1	0.6	0.3	1.2	2.0	2.1	1.7	1.1	1.2	*	*
1952	1.3	3.0	0.5	0.3	1.4	1.8	2.3	1.7	1.1	0.9	*	*
1951	1.4	4.2	0.5	0.4	1.3	1.9	2.0	1.7	1.1	1.0	1.3	*
1950	1.4	3.9	0.6	0.4	1.3	1.9	2.2	1.6	1.3	1.1	1.2	*

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

HIST290F. DEATH RATES FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years
HOMICIDE (E964, E980-E985)												
ALL OTHER, BOTH SEXES												
1959	21.8	7.2	1.4	1.0	28.3	51.3	44.9	29.6	18.2	10.8	8.5	*
1958	21.7	7.4	1.3	0.8	27.7	49.7	42.1	33.0	18.3	10.6	6.8	*
1957	22.5	5.6	1.6	1.1	28.1	51.6	42.4	33.2	18.0	13.5	8.2	*
1956	23.3	7.5	1.2	1.1	30.0	53.3	45.3	29.7	19.0	11.4	*	*
1955	22.8	6.5	1.2	1.0	27.9	52.9	41.6	31.3	20.5	9.0	*	*
1954	24.6	7.8	1.2	1.2	29.8	55.2	45.2	34.7	17.8	12.0	11.5	*
1953	25.0	4.7	*	1.1	31.3	53.7	50.2	31.7	16.4	11.1	*	*
1952	27.6	6.3	*	0.9	33.0	63.0	53.5	31.6	19.0	11.0	11.1	*
1951	25.6	6.6	1.3	0.8	31.9	55.8	47.0	30.9	19.5	8.6	*	*
1950	28.0	6.2	*	1.5	35.1	63.4	47.8	30.7	17.9	12.4	*	*
ALL OTHER, MALE												
1959	35.0	9.7	*	1.3	45.3	84.2	73.4	51.3	30.8	18.4	15.3	*
1958	34.9	8.3	*	*	44.2	83.4	68.8	54.7	31.8	18.1	*	*
1957	36.5	*	*	1.6	45.8	86.7	69.6	56.3	31.5	23.7	14.8	*
1956	37.1	*	*	1.7	48.8	88.0	73.7	49.0	30.5	18.3	*	*
1955	36.9	*	*	1.4	45.3	89.1	67.8	53.0	35.2	16.0	*	*
1954	40.6	10.1	*	1.6	49.6	94.8	76.2	57.6	33.2	19.9	*	*
1953	41.3	*	*	1.5	53.7	90.6	85.6	54.2	27.0	19.2	*	*
1952	45.4	*	*	*	55.8	106.1	90.1	53.1	32.7	19.1	18.2	*
1951	41.3	*	*	*	50.8	91.5	79.8	51.6	33.9	15.0	*	*
1950	45.5	*	*	1.8	56.5	105.6	80.4	52.8	32.3	20.9	*	*
ALL OTHER, FEMALE												
1959	9.4	*	*	*	12.4	22.9	19.2	9.2	6.0	*	*	*
1958	9.3	*	*	*	12.5	20.5	17.7	12.4	5.3	*	*	*
1957	9.2	*	*	*	12.0	21.3	17.4	11.0	4.9	*	*	*
1956	10.3	8.4	*	*	13.0	23.1	19.0	11.1	7.6	4.9	*	*
1955	9.5	*	*	*	12.3	21.5	17.2	10.2	6.0	*	*	*
1954	9.5	*	*	*	12.6	20.8	16.4	12.5	*	*	*	*
1953	9.6	*	*	*	12.2	21.6	17.2	9.7	5.6	*	*	*
1952	10.8	*	*	*	13.2	25.2	19.4	10.4	4.9	*	*	*
1951	10.7	*	*	*	15.3	24.3	16.6	10.4	4.5	*	*	*
1950	11.2	*	*	*	15.8	25.8	17.7	8.7	*	*	*	*

* Figure does not meet standard of reliability or precision

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)