

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

PAGE

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
ALL CAUSES													
BOTH SEXES													
1959	1,656,814	112,008	17,116	16,430	25,268	33,818	70,046	150,065	262,997	405,792	388,125	174,369	780
1958	1,647,886	113,789	17,561	15,537	24,381	34,217	70,495	149,495	263,378	404,850	383,172	170,240	771
1957	1,633,128	112,094	17,188	15,882	25,137	35,929	71,841	149,133	266,594	401,901	371,940	164,453	1,036
1956	1,564,476	108,183	16,603	14,899	24,587	35,468	69,424	142,839	258,343	382,144	357,751	153,261	974
1955	1,528,717	106,903	16,684	14,760	24,315	35,820	69,742	142,266	252,292	373,029	346,066	145,602	1,238
1954	1,481,091	106,791	16,931	14,655	23,652	36,716	69,647	143,500	249,275	360,225	326,417	132,368	914
1953	1,517,541	108,405	18,235	15,269	25,720	38,910	73,860	148,358	259,821	363,568	331,816	132,584	995
1952	1,496,838	109,413	19,499	15,803	27,196	41,284	75,782	148,992	258,840	353,596	320,715	124,630	1,088
1951	1,482,099	106,702	19,106	14,860	27,383	42,066	76,956	149,202	256,840	349,444	316,513	121,714	1,313
1950	1,452,454	103,825	18,148	14,607	28,298	42,467	76,937	148,087	254,150	342,296	305,851	116,516	1,272
MALE													
1959	943,088	64,291	9,464	10,130	17,887	21,237	42,821	96,719	169,379	241,200	196,556	72,929	475
1958	936,794	65,082	9,543	9,442	17,261	21,380	43,027	95,597	168,955	239,904	194,497	71,667	439
1957	930,728	64,366	9,363	9,684	17,592	22,266	43,645	95,093	171,120	238,274	188,840	69,847	638
1956	893,357	62,394	9,111	9,085	17,386	22,289	41,928	91,191	165,635	225,680	182,512	65,548	598
1955	872,638	61,416	9,218	9,082	16,989	22,412	42,056	90,693	161,391	219,781	176,576	62,291	733
1954	846,124	61,369	9,358	8,920	16,371	22,903	41,635	90,454	159,147	211,629	166,806	56,973	559
1953	867,100	62,377	10,116	9,351	17,590	23,973	44,389	93,520	165,992	212,066	169,553	57,557	616
1952	853,927	62,762	10,701	9,444	18,121	25,045	45,214	93,663	164,850	205,586	164,171	53,709	661
1951	845,233	61,465	10,494	9,058	18,016	25,258	45,853	93,113	162,474	203,619	161,779	53,294	810
1950	827,749	59,727	10,062	8,769	18,332	25,111	45,402	92,356	160,422	198,428	157,095	51,240	805
FEMALE													
1959	713,726	47,717	7,652	6,300	7,381	12,581	27,225	53,346	93,618	164,592	191,569	101,440	305
1958	711,092	48,707	8,018	6,095	7,120	12,837	27,468	53,898	94,423	164,946	188,675	98,573	332
1957	702,400	47,728	7,825	6,198	7,545	13,663	28,196	54,040	95,474	163,627	183,100	94,606	398
1956	671,119	45,789	7,492	5,814	7,201	13,179	27,496	51,648	92,708	156,464	175,239	87,713	376
1955	656,079	45,487	7,466	5,678	7,326	13,408	27,686	51,573	90,901	153,248	169,490	83,311	505
1954	634,967	45,422	7,573	5,735	7,281	13,813	28,012	53,046	90,128	148,596	159,611	75,395	355
1953	650,441	46,028	8,119	5,918	8,130	14,937	29,471	54,838	93,829	151,502	162,263	75,027	379
1952	642,911	46,651	8,798	6,359	9,075	16,239	30,568	55,329	93,990	148,010	156,544	70,921	427
1951	636,866	45,237	8,612	5,802	9,367	16,808	31,103	56,089	94,366	145,825	154,734	68,420	503
1950	624,705	44,098	8,086	5,838	9,966	17,356	31,535	55,731	93,728	143,868	148,756	65,276	467

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
ALL CAUSES													
WHITE, BOTH SEXES													
1959	1,460,840	83,493	12,824	13,485	20,596	25,352	54,657	124,393	226,705	367,389	366,020	165,410	516
1958	1,451,077	84,935	13,245	12,896	19,876	25,521	55,300	123,174	226,860	366,082	361,328	161,399	461
1957	1,437,652	84,412	13,013	13,052	20,356	26,848	56,041	122,393	229,608	364,100	351,235	155,971	623
1956	1,380,915	82,203	12,689	12,321	19,968	26,708	54,051	117,505	223,502	347,407	338,327	145,572	662
1955	1,350,869	81,682	12,761	12,296	19,671	27,112	54,515	116,715	219,693	339,484	327,541	138,485	914
1954	1,306,989	82,200	13,052	12,205	18,871	27,644	54,247	117,174	217,728	328,505	309,066	125,667	630
1953	1,335,835	84,041	14,123	12,703	20,369	29,199	57,188	120,471	226,836	330,863	313,945	125,419	678
1952	1,315,022	84,752	15,076	13,180	21,338	30,867	58,498	120,952	226,232	322,064	303,418	117,843	802
1951	1,302,969	83,678	14,854	12,174	21,305	31,645	59,512	121,041	224,675	318,501	299,555	115,048	981
1950	1,276,085	82,018	14,203	12,002	21,793	31,558	59,208	120,144	222,222	312,482	289,503	110,034	918
WHITE, MALE													
1959	834,651	48,525	7,098	8,411	14,915	16,356	34,432	82,407	149,053	219,616	184,776	68,740	322
1958	828,550	49,027	7,299	7,910	14,358	16,505	34,902	81,227	148,634	218,199	182,767	67,452	270
1957	823,067	49,047	7,134	8,021	14,622	17,112	35,201	80,523	150,285	217,214	177,695	65,816	397
1956	792,364	47,789	7,026	7,576	14,406	17,324	33,843	77,453	146,171	206,462	171,960	61,936	418
1955	774,758	47,457	7,128	7,652	14,088	17,549	33,914	76,805	142,978	201,104	166,553	58,987	543
1954	750,099	47,727	7,265	7,475	13,437	17,811	33,547	76,194	141,140	194,008	157,292	53,809	394
1953	766,703	49,019	7,857	7,829	14,398	18,532	35,485	78,216	147,030	193,820	159,861	54,227	429
1952	753,571	48,986	8,312	7,940	14,724	19,351	36,043	78,386	146,153	188,099	154,511	50,577	489
1951	747,049	48,509	8,199	7,478	14,604	19,631	36,795	77,829	144,253	186,377	152,534	50,227	613
1950	731,366	47,595	7,921	7,298	14,769	19,323	36,293	77,150	142,419	181,770	147,984	48,249	595
WHITE, FEMALE													
1959	626,189	34,968	5,726	5,074	5,681	8,996	20,225	41,986	77,652	147,773	181,244	96,670	194
1958	622,527	35,908	5,946	4,986	5,518	9,016	20,398	41,947	78,226	147,883	178,561	93,947	191
1957	614,585	35,365	5,879	5,031	5,734	9,736	20,840	41,870	79,323	146,886	173,540	90,155	226
1956	588,551	34,414	5,663	4,745	5,562	9,384	20,208	40,052	77,331	140,945	166,367	83,636	244
1955	576,111	34,225	5,633	4,644	5,583	9,563	20,601	39,910	76,715	138,380	160,988	79,498	371
1954	556,890	34,473	5,787	4,730	5,434	9,833	20,700	40,980	76,588	134,497	151,774	71,858	236
1953	569,132	35,022	6,266	4,874	5,971	10,667	21,703	42,255	79,806	137,043	154,084	71,192	249
1952	561,451	35,766	6,764	5,240	6,614	11,516	22,455	42,566	80,079	133,965	148,907	67,266	313
1951	555,920	35,169	6,655	4,696	6,701	12,014	22,717	43,212	80,422	132,124	147,021	64,821	368
1950	544,719	34,423	6,282	4,704	7,024	12,235	22,915	42,994	79,803	130,712	141,519	61,785	323

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
ALL CAUSES													
ALL OTHER, BOTH SEXES													
1959	195,974	28,515	4,292	2,945	4,672	8,466	15,389	25,672	36,292	38,403	22,105	8,959	264
1958	196,809	28,854	4,316	2,641	4,505	8,696	15,195	26,321	36,518	38,768	21,844	8,841	310
1957	195,476	27,682	4,175	2,830	4,781	9,081	15,800	26,740	36,986	37,801	20,705	8,482	413
1956	183,561	25,980	3,914	2,578	4,619	8,760	15,373	25,334	34,841	34,737	19,424	7,689	312
1955	177,848	25,221	3,923	2,464	4,644	8,708	15,227	25,551	32,599	33,545	18,525	7,117	324
1954	174,102	24,591	3,879	2,450	4,781	9,072	15,400	26,326	31,547	31,720	17,351	6,701	284
1953	181,706	24,364	4,112	2,566	5,351	9,711	16,672	27,887	32,985	32,705	17,871	7,165	317
1952	181,816	24,661	4,423	2,623	5,858	10,417	17,284	28,040	32,608	31,532	17,297	6,787	286
1951	179,130	23,024	4,252	2,686	6,078	10,421	17,444	28,161	32,165	30,943	16,958	6,666	332
1950	176,369	21,807	3,945	2,605	6,505	10,909	17,729	27,943	31,928	29,814	16,348	6,482	354
ALL OTHER, MALE													
1959	108,437	15,766	2,366	1,719	2,972	4,881	8,389	14,312	20,326	21,584	11,780	4,189	153
1958	108,244	16,055	2,244	1,532	2,903	4,875	8,125	14,370	20,321	21,705	11,730	4,215	169
1957	107,661	15,319	2,229	1,663	2,970	5,154	8,444	14,570	20,835	21,060	11,145	4,031	241
1956	100,993	14,605	2,085	1,509	2,980	4,965	8,085	13,738	19,464	19,218	10,552	3,612	180
1955	97,880	13,959	2,090	1,430	2,901	4,863	8,142	13,888	18,413	18,677	10,023	3,304	190
1954	96,025	13,642	2,093	1,445	2,934	5,092	8,088	14,260	18,007	17,621	9,514	3,164	165
1953	100,397	13,358	2,259	1,522	3,192	5,441	8,904	15,304	18,962	18,246	9,692	3,330	187
1952	100,356	13,776	2,389	1,504	3,397	5,694	9,171	15,277	18,697	17,487	9,660	3,132	172
1951	98,184	12,956	2,295	1,580	3,412	5,627	9,058	15,284	18,221	17,242	9,245	3,067	197
1950	96,383	12,132	2,141	1,471	3,563	5,788	9,109	15,206	18,003	16,658	9,111	2,991	210
ALL OTHER, FEMALE													
1959	87,537	12,749	1,926	1,226	1,700	3,585	7,000	11,360	15,966	16,819	10,325	4,770	111
1958	88,565	12,799	2,072	1,109	1,602	3,821	7,070	11,951	16,197	17,063	10,114	4,626	141
1957	87,815	12,363	1,946	1,167	1,811	3,927	7,356	12,170	16,151	16,741	9,560	4,451	172
1956	82,568	11,375	1,829	1,069	1,639	3,795	7,288	11,596	15,377	15,519	8,872	4,077	132
1955	79,968	11,262	1,833	1,034	1,743	3,845	7,085	11,663	14,186	14,868	8,502	3,813	134
1954	78,077	10,949	1,786	1,005	1,847	3,980	7,312	12,066	13,540	14,099	7,837	3,537	119
1953	81,309	11,006	1,853	1,044	2,159	4,270	7,768	12,583	14,023	14,459	8,179	3,835	130
1952	81,460	10,885	2,034	1,119	2,461	4,723	8,113	12,763	13,911	14,045	7,637	3,655	114
1951	80,946	10,068	1,957	1,106	2,666	4,794	8,386	12,877	13,944	13,701	7,713	3,599	135
1950	79,986	9,675	1,804	1,134	2,942	5,121	8,620	12,737	13,925	13,156	7,237	3,491	144

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS, ALL FORMS (001-019)													
BOTH SEXES													
1959	11,456	50	131	59	153	641	1,272	2,070	2,568	2,669	1,494	341	8
1958	12,361	60	142	68	212	778	1,457	2,264	2,768	2,803	1,502	301	6
1957	13,324	77	144	62	252	912	1,663	2,468	2,893	2,973	1,545	325	10
1956	14,061	94	223	68	256	990	1,804	2,584	3,066	3,025	1,608	336	7
1955	14,940	96	256	98	345	1,160	2,014	2,820	3,149	3,128	1,564	302	8
1954	16,392	147	277	96	466	1,391	2,318	3,061	3,385	3,297	1,641	299	14
1953	19,544	144	436	164	711	1,845	2,778	3,637	4,136	3,616	1,747	316	14
1952	24,621	222	580	253	1,243	2,751	3,814	4,540	4,888	4,134	1,887	286	23
1951	30,863	234	732	343	1,874	4,026	4,936	5,752	5,987	4,605	2,020	334	20
1950	33,959	268	823	440	2,497	4,542	5,600	6,227	6,342	4,855	2,071	275	19
MALE													
1959	8,263	33	62	24	53	321	738	1,534	2,079	2,136	1,057	218	8
1958	8,943	30	77	34	94	357	891	1,747	2,253	2,226	1,041	188	5
1957	9,649	37	79	35	116	432	1,018	1,855	2,389	2,371	1,104	205	8
1956	9,995	49	111	36	91	443	1,105	1,953	2,515	2,356	1,127	203	6
1955	10,542	61	136	53	136	574	1,195	2,169	2,575	2,417	1,035	187	4
1954	11,494	80	137	47	174	657	1,416	2,314	2,803	2,542	1,123	190	11
1953	13,574	75	235	82	299	890	1,761	2,791	3,335	2,761	1,150	186	9
1952	16,815	110	312	124	514	1,339	2,384	3,515	3,998	3,106	1,221	172	20
1951	20,720	120	378	148	776	1,946	3,136	4,427	4,832	3,393	1,327	223	14
1950	22,539	137	424	198	982	2,166	3,550	4,820	5,111	3,643	1,336	162	10
FEMALE													
1959	3,193	17	69	35	100	320	534	536	489	533	437	123	-
1958	3,418	30	65	34	118	421	566	517	515	577	461	113	1
1957	3,675	40	65	27	136	480	645	613	504	602	441	120	2
1956	4,066	45	112	32	165	547	699	631	551	669	481	133	1
1955	4,398	35	120	45	209	586	819	651	574	711	529	115	4
1954	4,898	67	140	49	292	734	902	747	582	755	518	109	3
1953	5,970	69	201	82	412	955	1,017	846	801	855	597	130	5
1952	7,806	112	268	129	729	1,412	1,430	1,025	890	1,028	666	114	3
1951	10,143	114	354	195	1,098	2,080	1,800	1,325	1,155	1,212	693	111	6
1950	11,420	131	399	242	1,515	2,376	2,050	1,407	1,231	1,212	735	113	9

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS, ALL FORMS (001-019)													
WHITE, BOTH SEXES													
1959	8,601	26	71	29	73	276	774	1,478	1,996	2,261	1,318	296	3
1958	9,178	34	76	41	86	367	881	1,623	2,147	2,326	1,337	255	5
1957	10,030	34	84	37	108	462	1,081	1,778	2,268	2,522	1,363	288	5
1956	10,502	48	113	42	99	459	1,179	1,806	2,413	2,592	1,460	289	2
1955	11,116	51	139	57	120	585	1,288	2,018	2,519	2,676	1,395	262	6
1954	12,138	88	144	54	203	715	1,482	2,173	2,696	2,838	1,466	273	6
1953	14,254	71	251	91	296	947	1,776	2,572	3,295	3,077	1,584	287	7
1952	17,678	107	335	133	485	1,450	2,405	3,231	3,970	3,585	1,719	251	7
1951	22,280	143	437	188	851	2,292	3,281	4,137	4,837	3,975	1,829	297	13
1950	24,136	149	529	241	1,065	2,485	3,682	4,462	5,164	4,213	1,889	247	10
WHITE, MALE													
1959	6,397	18	34	14	30	135	460	1,100	1,654	1,833	929	187	3
1958	6,810	17	47	26	43	172	547	1,263	1,770	1,850	918	153	4
1957	7,465	17	49	20	56	211	674	1,355	1,900	2,018	984	177	4
1956	7,699	25	59	24	40	201	729	1,387	2,009	2,039	1,016	168	2
1955	8,086	30	69	33	51	294	761	1,591	2,085	2,093	916	161	2
1954	8,811	50	65	31	92	354	914	1,676	2,255	2,199	998	172	5
1953	10,214	40	140	51	133	440	1,123	2,015	2,692	2,365	1,042	168	5
1952	12,482	52	181	77	202	692	1,494	2,560	3,263	2,705	1,104	146	6
1951	15,480	71	225	87	346	1,095	2,111	3,245	3,959	2,948	1,186	198	9
1950	16,787	72	257	116	413	1,184	2,416	3,561	4,240	3,170	1,205	147	6
WHITE, FEMALE													
1959	2,204	8	37	15	43	141	314	378	342	428	389	109	-
1958	2,368	17	29	15	43	195	334	360	377	476	419	102	1
1957	2,565	17	35	17	52	251	407	423	368	504	379	111	1
1956	2,803	23	54	18	59	258	450	419	404	553	444	121	-
1955	3,030	21	70	24	69	291	527	427	434	583	479	101	4
1954	3,327	38	79	23	111	361	568	497	441	639	468	101	1
1953	4,040	31	111	40	163	507	653	557	603	712	542	119	2
1952	5,196	55	154	56	283	758	911	671	707	880	615	105	1
1951	6,800	72	212	101	505	1,197	1,170	892	878	1,027	643	99	4
1950	7,349	77	272	125	652	1,301	1,266	901	924	1,043	684	100	4

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS, ALL FORMS (001-019)													
ALL OTHER, BOTH SEXES													
1959	2,855	24	60	30	80	365	498	592	572	408	176	45	5
1958	3,183	26	66	27	126	411	576	641	621	477	165	46	1
1957	3,294	43	60	25	144	450	582	690	625	451	182	37	5
1956	3,559	46	110	26	157	531	625	778	653	433	148	47	5
1955	3,824	45	117	41	225	575	726	802	630	452	169	40	2
1954	4,254	59	133	42	263	676	836	888	689	459	175	26	8
1953	5,290	73	185	73	415	898	1,002	1,065	841	539	163	29	7
1952	6,943	115	245	120	758	1,301	1,409	1,309	918	549	168	35	16
1951	8,583	91	295	155	1,023	1,734	1,655	1,615	1,150	630	191	37	7
1950	9,823	119	294	199	1,432	2,057	1,918	1,765	1,178	642	182	28	9
ALL OTHER, MALE													
1959	1,866	15	28	10	23	186	278	434	425	303	128	31	5
1958	2,133	13	30	8	51	185	344	484	483	376	123	35	1
1957	2,184	20	30	15	60	221	344	500	489	353	120	28	4
1956	2,296	24	52	12	51	242	376	566	506	317	111	35	4
1955	2,456	31	67	20	85	280	434	578	490	324	119	26	2
1954	2,683	30	72	16	82	303	502	638	548	343	125	18	6
1953	3,360	35	95	31	166	450	638	776	643	396	108	18	4
1952	4,333	58	131	47	312	647	890	955	735	401	117	26	14
1951	5,240	49	153	61	430	851	1,025	1,182	873	445	141	25	5
1950	5,752	65	167	82	569	982	1,134	1,259	871	473	131	15	4
ALL OTHER, FEMALE													
1959	989	9	32	20	57	179	220	158	147	105	48	14	-
1958	1,050	13	36	19	75	226	232	157	138	101	42	11	-
1957	1,110	23	30	10	84	229	238	190	136	98	62	9	1
1956	1,263	22	58	14	106	289	249	212	147	116	37	12	1
1955	1,368	14	50	21	140	295	292	224	140	128	50	14	-
1954	1,571	29	61	26	181	373	334	250	141	116	50	8	2
1953	1,930	38	90	42	249	448	364	289	198	143	55	11	3
1952	2,610	57	114	73	446	654	519	354	183	148	51	9	2
1951	3,343	42	142	94	593	883	630	433	277	185	50	12	2
1950	4,071	54	127	117	863	1,075	784	506	307	169	51	13	5

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)													
BOTH SEXES													
1959	10,627	19	41	15	115	575	1,198	1,963	2,423	2,528	1,420	323	7
1958	11,477	25	44	22	175	704	1,356	2,158	2,612	2,656	1,433	286	6
1957	12,384	36	47	23	209	828	1,556	2,329	2,741	2,822	1,468	315	10
1956	12,978	44	74	28	214	896	1,687	2,433	2,884	2,852	1,531	328	7
1955	13,676	39	83	29	288	1,065	1,851	2,630	2,942	2,965	1,483	293	8
1954	15,024	69	106	40	401	1,267	2,153	2,876	3,160	3,098	1,549	291	14
1953	17,814	51	145	65	609	1,664	2,556	3,414	3,909	3,411	1,677	299	14
1952	22,486	97	184	113	1,074	2,536	3,558	4,272	4,647	3,920	1,794	272	19
1951	28,296	107	260	169	1,665	3,737	4,622	5,457	5,678	4,350	1,916	316	19
1950	31,093	108	268	217	2,216	4,215	5,298	5,916	6,004	4,603	1,975	257	16
MALE													
1959	7,779	11	16	9	39	281	694	1,468	1,991	2,037	1,017	209	7
1958	8,409	14	25	10	76	323	825	1,690	2,136	2,130	999	176	5
1957	9,068	16	25	13	96	384	962	1,768	2,283	2,263	1,050	200	8
1956	9,359	22	38	17	73	398	1,040	1,857	2,391	2,236	1,084	197	6
1955	9,780	24	35	15	108	521	1,097	2,050	2,445	2,313	983	185	4
1954	10,670	39	49	19	149	591	1,324	2,194	2,644	2,399	1,065	186	11
1953	12,570	29	87	28	249	795	1,640	2,645	3,187	2,620	1,104	177	9
1952	15,586	54	91	46	429	1,232	2,232	3,349	3,835	2,972	1,166	164	16
1951	19,266	56	136	61	672	1,800	2,944	4,239	4,649	3,227	1,260	209	13
1950	20,885	57	143	76	840	1,992	3,371	4,604	4,873	3,480	1,287	154	8
FEMALE													
1959	2,848	8	25	6	76	294	504	495	432	491	403	114	-
1958	3,068	11	19	12	99	381	531	468	476	526	434	110	1
1957	3,316	20	22	10	113	444	594	561	458	559	418	115	2
1956	3,619	22	36	11	141	498	647	576	493	616	447	131	1
1955	3,896	15	48	14	180	544	754	580	497	652	500	108	4
1954	4,354	30	57	21	252	676	829	682	516	699	484	105	3
1953	5,244	22	58	37	360	869	916	769	722	791	573	122	5
1952	6,900	43	93	67	645	1,304	1,326	923	812	948	628	108	3
1951	9,030	51	124	108	993	1,937	1,678	1,218	1,029	1,123	656	107	6
1950	10,208	51	125	141	1,376	2,223	1,927	1,312	1,131	1,123	688	103	8

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)													
WHITE, BOTH SEXES													
1959	8,107	10	19	5	56	251	737	1,422	1,910	2,154	1,260	280	3
1958	8,668	16	23	11	72	341	830	1,564	2,058	2,224	1,280	244	5
1957	9,466	17	26	16	81	434	1,026	1,688	2,179	2,409	1,305	280	5
1956	9,859	26	38	18	81	426	1,122	1,722	2,293	2,455	1,393	283	2
1955	10,377	24	44	15	97	552	1,207	1,920	2,378	2,554	1,325	255	6
1954	11,289	42	53	20	173	667	1,406	2,058	2,534	2,675	1,390	265	6
1953	13,222	26	84	27	249	880	1,651	2,439	3,148	2,914	1,523	274	7
1952	16,436	48	97	48	417	1,365	2,272	3,089	3,800	3,413	1,639	241	7
1951	20,674	68	143	69	761	2,154	3,108	3,954	4,620	3,767	1,735	282	13
1950	22,357	58	156	98	943	2,343	3,517	4,278	4,923	3,995	1,806	232	8
WHITE, MALE													
1959	6,102	7	8	4	21	121	437	1,065	1,602	1,756	899	179	3
1958	6,486	8	15	6	34	161	509	1,231	1,707	1,783	884	144	4
1957	7,112	10	15	9	43	195	643	1,298	1,838	1,940	943	174	4
1956	7,293	13	20	11	30	183	694	1,329	1,924	1,944	980	163	2
1955	7,628	12	16	9	40	276	710	1,524	1,992	2,015	873	159	2
1954	8,268	22	20	13	79	326	869	1,598	2,142	2,080	946	168	5
1953	9,605	16	54	14	112	397	1,058	1,933	2,597	2,254	1,003	162	5
1952	11,765	24	52	24	174	651	1,414	2,469	3,155	2,599	1,056	141	6
1951	14,576	35	76	28	305	1,024	2,001	3,126	3,840	2,817	1,128	187	9
1950	15,754	30	71	42	351	1,107	2,315	3,430	4,073	3,028	1,162	140	5
WHITE, FEMALE													
1959	2,005	3	11	1	35	130	300	357	308	398	361	101	-
1958	2,182	8	8	5	38	180	321	333	351	441	396	100	1
1957	2,354	7	11	7	38	239	383	390	341	469	362	106	1
1956	2,566	13	18	7	51	243	428	393	369	511	413	120	-
1955	2,749	12	28	6	57	276	497	396	386	539	452	96	4
1954	3,021	20	33	7	94	341	537	460	392	595	444	97	1
1953	3,617	10	30	13	137	483	593	506	551	660	520	112	2
1952	4,671	24	45	24	243	714	858	620	645	814	583	100	1
1951	6,098	33	67	41	456	1,130	1,107	828	780	950	607	95	4
1950	6,603	28	85	56	592	1,236	1,202	848	850	967	644	92	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS OF RESPIRATORY SYSTEM (001-008)													
ALL OTHER, BOTH SEXES													
1959	2,520	9	22	10	59	324	461	541	513	374	160	43	4
1958	2,809	9	21	11	103	363	526	594	554	432	153	42	1
1957	2,918	19	21	7	128	394	530	641	562	413	163	35	5
1956	3,119	18	36	10	133	470	565	711	591	397	138	45	5
1955	3,299	15	39	14	191	513	644	710	564	411	158	38	2
1954	3,735	27	53	20	228	600	747	818	626	423	159	26	8
1953	4,592	25	61	38	360	784	905	975	761	497	154	25	7
1952	6,050	49	87	65	657	1,171	1,286	1,183	847	507	155	31	12
1951	7,622	39	117	100	904	1,583	1,514	1,503	1,058	583	181	34	6
1950	8,736	50	112	119	1,273	1,872	1,781	1,638	1,081	608	169	25	8
ALL OTHER, MALE													
1959	1,677	4	8	5	18	160	257	403	389	281	118	30	4
1958	1,923	6	10	4	42	162	316	459	429	347	115	32	1
1957	1,956	6	10	4	53	189	319	470	445	323	107	26	4
1956	2,066	9	18	6	43	215	346	528	467	292	104	34	4
1955	2,152	12	19	6	68	245	387	526	453	298	110	26	2
1954	2,402	17	29	6	70	265	455	596	502	319	119	18	6
1953	2,965	13	33	14	137	398	582	712	590	366	101	15	4
1952	3,821	30	39	22	255	581	818	880	680	373	110	23	10
1951	4,690	21	60	33	367	776	943	1,113	809	410	132	22	4
1950	5,131	27	72	34	489	885	1,056	1,174	800	452	125	14	3
ALL OTHER, FEMALE													
1959	843	5	14	5	41	164	204	138	124	93	42	13	-
1958	886	3	11	7	61	201	210	135	125	85	38	10	-
1957	962	13	11	3	75	205	211	171	117	90	56	9	1
1956	1,053	9	18	4	90	255	219	183	124	105	34	11	1
1955	1,147	3	20	8	123	268	257	184	111	113	48	12	-
1954	1,333	10	24	14	158	335	292	222	124	104	40	8	2
1953	1,627	12	28	24	223	386	323	263	171	131	53	10	3
1952	2,229	19	48	43	402	590	468	303	167	134	45	8	2
1951	2,932	18	57	67	537	807	571	390	249	173	49	12	2
1950	3,605	23	40	85	784	987	725	464	281	156	44	11	5

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS, OTHER FORMS (010-019)													
BOTH SEXES													
1959	829	31	90	44	38	66	74	107	145	141	74	18	1
1958	884	35	98	46	37	74	101	106	156	147	69	15	-
1957	940	41	97	39	43	84	107	139	152	151	77	10	-
1956	1,083	50	149	40	42	94	117	151	182	173	77	8	-
1955	1,264	57	173	69	57	95	163	190	207	163	81	9	-
1954	1,368	78	171	56	65	124	165	185	225	199	92	8	-
1953	1,730	93	291	99	102	181	222	223	227	205	70	17	-
1952	2,135	125	396	140	169	215	256	268	241	214	93	14	4
1951	2,567	127	472	174	209	289	314	295	309	255	104	18	1
1950	2,866	160	555	223	281	327	302	311	338	252	96	18	3
MALE													
1959	484	22	46	15	14	40	44	66	88	99	40	9	1
1958	534	16	52	24	18	34	66	57	117	96	42	12	-
1957	581	21	54	22	20	48	56	87	106	108	54	5	-
1956	636	27	73	19	18	45	65	96	124	120	43	6	-
1955	762	37	101	38	28	53	98	119	130	104	52	2	-
1954	824	41	88	28	25	66	92	120	159	143	58	4	-
1953	1,004	46	148	54	50	95	121	146	148	141	46	9	-
1952	1,229	56	221	78	85	107	152	166	163	134	55	8	4
1951	1,454	64	242	87	104	146	192	188	183	166	67	14	1
1950	1,654	80	281	122	142	174	179	216	238	163	49	8	2
FEMALE													
1959	345	9	44	29	24	26	30	41	57	42	34	9	-
1958	350	19	46	22	19	40	35	49	39	51	27	3	-
1957	359	20	43	17	23	36	51	52	46	43	23	5	-
1956	447	23	76	21	24	49	52	55	58	53	34	2	-
1955	502	20	72	31	29	42	65	71	77	59	29	7	-
1954	544	37	83	28	40	58	73	65	66	56	34	4	-
1953	726	47	143	45	52	86	101	77	79	64	24	8	-
1952	906	69	175	62	84	108	104	102	78	80	38	6	-
1951	1,113	63	230	87	105	143	122	107	126	89	37	4	-
1950	1,212	80	274	101	139	153	123	95	100	89	47	10	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS, OTHER FORMS (010-019)													
WHITE, BOTH SEXES													
1959	494	16	52	24	17	25	37	56	86	107	58	16	-
1958	510	18	53	30	14	26	51	59	89	102	57	11	-
1957	564	17	58	21	27	28	55	90	89	113	58	8	-
1956	643	22	75	24	18	33	57	84	120	137	67	6	-
1955	739	27	95	42	23	33	81	98	141	122	70	7	-
1954	849	46	91	34	30	48	76	115	162	163	76	8	-
1953	1,032	45	167	64	47	67	125	133	147	163	61	13	-
1952	1,242	59	238	85	68	85	133	142	170	172	80	10	-
1951	1,606	75	294	119	90	138	173	183	217	208	94	15	-
1950	1,779	91	373	143	122	142	165	184	241	218	83	15	2
WHITE, MALE													
1959	295	11	26	10	9	14	23	35	52	77	30	8	-
1958	324	9	32	20	9	11	38	32	63	67	34	9	-
1957	353	7	34	11	13	16	31	57	62	78	41	3	-
1956	406	12	39	13	10	18	35	58	85	95	36	5	-
1955	458	18	53	24	11	18	51	67	93	78	43	2	-
1954	543	28	45	18	13	28	45	78	113	119	52	4	-
1953	609	24	86	37	21	43	65	82	95	111	39	6	-
1952	717	28	129	53	28	41	80	91	108	106	48	5	-
1951	904	36	149	59	41	71	110	119	119	131	58	11	-
1950	1,033	42	186	74	62	77	101	131	167	142	43	7	1
WHITE, FEMALE													
1959	199	5	26	14	8	11	14	21	34	30	28	8	-
1958	186	9	21	10	5	15	13	27	26	35	23	2	-
1957	211	10	24	10	14	12	24	33	27	35	17	5	-
1956	237	10	36	11	8	15	22	26	35	42	31	1	-
1955	281	9	42	18	12	15	30	31	48	44	27	5	-
1954	306	18	46	16	17	20	31	37	49	44	24	4	-
1953	423	21	81	27	26	24	60	51	52	52	22	7	-
1952	525	31	109	32	40	44	53	51	62	66	32	5	-
1951	702	39	145	60	49	67	63	64	98	77	36	4	-
1950	746	49	187	69	60	65	64	53	74	76	40	8	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
TUBERCULOSIS, OTHER FORMS (010-019)													
ALL OTHER, BOTH SEXES													
1959	335	15	38	20	21	41	37	51	59	34	16	2	1
1958	374	17	45	16	23	48	50	47	67	45	12	4	-
1957	376	24	39	18	16	56	52	49	63	38	19	2	-
1956	440	28	74	16	24	61	60	67	62	36	10	2	-
1955	525	30	78	27	34	62	82	92	66	41	11	2	-
1954	519	32	80	22	35	76	89	70	63	36	16	-	-
1953	698	48	124	35	55	114	97	90	80	42	9	4	-
1952	893	66	158	55	101	130	123	126	71	42	13	4	4
1951	961	52	178	55	119	151	141	112	92	47	10	3	1
1950	1,087	69	182	80	159	185	137	127	97	34	13	3	1
ALL OTHER, MALE													
1959	189	11	20	5	5	26	21	31	36	22	10	1	1
1958	210	7	20	4	9	23	28	25	54	29	8	3	-
1957	228	14	20	11	7	32	25	30	44	30	13	2	-
1956	230	15	34	6	8	27	30	38	39	25	7	1	-
1955	304	19	48	14	17	35	47	52	37	26	9	-	-
1954	281	13	43	10	12	38	47	42	46	24	6	-	-
1953	395	22	62	17	29	52	56	64	53	30	7	3	-
1952	512	28	92	25	57	66	72	75	55	28	7	3	4
1951	550	28	93	28	63	75	82	69	64	35	9	3	1
1950	621	38	95	48	80	97	78	85	71	21	6	1	1
ALL OTHER, FEMALE													
1959	146	4	18	15	16	15	16	20	23	12	6	1	-
1958	164	10	25	12	14	25	22	22	13	16	4	1	-
1957	148	10	19	7	9	24	27	19	19	8	6	-	-
1956	210	13	40	10	16	34	30	29	23	11	3	1	-
1955	221	11	30	13	17	27	35	40	29	15	2	2	-
1954	238	19	37	12	23	38	42	28	17	12	10	-	-
1953	303	26	62	18	26	62	41	26	27	12	2	1	-
1952	381	38	66	30	44	64	51	51	16	14	6	1	-
1951	411	24	85	27	56	76	59	43	28	12	1	-	-
1950	466	31	87	32	79	88	59	42	26	13	7	2	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SYPHILIS AND ITS SEQUELAE (020-029)													
BOTH SEXES													
1959	3,069	21	1	5	12	35	129	445	918	989	433	78	3
1958	3,469	26	3	1	13	41	147	539	1,047	1,079	475	94	4
1957	3,825	27	-	1	17	50	194	657	1,144	1,152	489	91	3
1956	3,870	26	3	6	17	51	195	703	1,202	1,143	437	83	4
1955	3,834	34	5	2	10	56	238	754	1,156	1,098	420	56	5
1954	4,835	43	-	2	23	90	290	885	1,445	1,405	564	88	-
1953	5,273	56	6	6	26	71	384	1,053	1,611	1,432	539	84	5
1952	5,719	92	9	8	42	87	432	1,207	1,719	1,488	538	92	5
1951	6,274	129	11	11	44	123	510	1,379	1,911	1,529	552	68	7
1950	7,568	201	9	10	57	168	715	1,609	2,116	1,837	744	96	6
MALE													
1959	2,177	12	-	2	6	21	88	296	696	742	262	49	3
1958	2,489	19	1	1	8	29	90	365	806	796	318	53	3
1957	2,759	14	-	1	12	25	121	471	866	871	327	50	1
1956	2,829	11	2	6	13	27	122	516	929	864	288	49	2
1955	2,840	20	4	1	7	35	172	543	904	829	292	29	4
1954	3,533	24	-	1	9	49	183	625	1,135	1,078	383	46	-
1953	3,890	33	4	2	16	31	247	790	1,263	1,087	367	45	5
1952	4,262	53	2	4	17	47	277	905	1,372	1,148	378	56	3
1951	4,669	67	4	6	25	75	337	1,040	1,530	1,170	379	31	5
1950	5,550	114	7	5	30	97	452	1,194	1,678	1,411	510	49	3
FEMALE													
1959	892	9	1	3	6	14	41	149	222	247	171	29	-
1958	980	7	2	-	5	12	57	174	241	283	157	41	1
1957	1,066	13	-	-	5	25	73	186	278	281	162	41	2
1956	1,041	15	1	-	4	24	73	187	273	279	149	34	2
1955	994	14	1	1	3	21	66	211	252	269	128	27	1
1954	1,302	19	-	1	14	41	107	260	310	327	181	42	-
1953	1,383	23	2	4	10	40	137	263	348	345	172	39	-
1952	1,457	39	7	4	25	40	155	302	347	340	160	36	2
1951	1,605	62	7	5	19	48	173	339	381	359	173	37	2
1950	2,018	87	2	5	27	71	263	415	438	426	234	47	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SYPHILIS AND ITS SEQUELAE (020-029)													
WHITE, BOTH SEXES													
1959	2,106	8	-	3	6	19	58	216	592	782	355	65	2
1958	2,240	6	-	-	7	21	48	257	658	800	370	72	1
1957	2,540	17	-	1	8	25	77	329	759	868	385	70	1
1956	2,574	7	1	5	9	26	68	388	771	877	356	65	1
1955	2,438	9	2	-	5	26	95	372	742	809	334	41	3
1954	3,238	12	-	1	7	31	113	435	962	1,113	494	70	-
1953	3,442	14	-	3	19	22	140	520	1,082	1,122	447	71	2
1952	3,787	32	5	4	23	32	168	637	1,185	1,174	446	79	2
1951	4,115	40	6	3	23	48	206	732	1,329	1,201	469	56	2
1950	5,024	74	-	2	37	55	294	837	1,500	1,503	636	83	3
WHITE, MALE													
1959	1,528	3	-	1	2	10	39	152	470	599	212	38	2
1958	1,637	3	-	-	5	16	31	181	519	600	243	38	1
1957	1,868	8	-	1	6	15	44	244	596	661	254	39	-
1956	1,930	4	-	5	7	17	42	291	623	673	226	42	-
1955	1,834	8	2	-	3	19	68	272	586	619	233	22	2
1954	2,414	3	-	1	3	19	79	307	769	865	336	32	-
1953	2,603	8	-	1	13	10	94	408	864	869	296	38	2
1952	2,893	17	1	2	12	22	104	487	958	925	317	46	2
1951	3,148	19	2	2	16	24	141	573	1,095	929	319	27	1
1950	3,764	43	-	2	22	28	183	628	1,211	1,168	435	41	3
WHITE, FEMALE													
1959	578	5	-	2	4	9	19	64	122	183	143	27	-
1958	603	3	-	-	2	5	17	76	139	200	127	34	-
1957	672	9	-	-	2	10	33	85	163	207	131	31	1
1956	644	3	1	-	2	9	26	97	148	204	130	23	1
1955	604	1	-	-	2	7	27	100	156	190	101	19	1
1954	824	9	-	-	4	12	34	128	193	248	158	38	-
1953	839	6	-	2	6	12	46	112	218	253	151	33	-
1952	894	15	4	2	11	10	64	150	227	249	129	33	-
1951	967	21	4	1	7	24	65	159	234	272	150	29	1
1950	1,260	31	-	-	15	27	111	209	289	335	201	42	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SYPHILIS AND ITS SEQUELAE (020-029)													
ALL OTHER, BOTH SEXES													
1959	963	13	1	2	6	16	71	229	326	207	78	13	1
1958	1,229	20	3	1	6	20	99	282	389	279	105	22	3
1957	1,285	10	-	-	9	25	117	328	385	284	104	21	2
1956	1,296	19	2	1	8	25	127	315	431	266	81	18	3
1955	1,396	25	3	2	5	30	143	382	414	289	86	15	2
1954	1,597	31	-	1	16	59	177	450	483	292	70	18	-
1953	1,831	42	6	3	7	49	244	533	529	310	92	13	3
1952	1,932	60	4	4	19	55	264	570	534	314	92	13	3
1951	2,159	89	5	8	21	75	304	647	582	328	83	12	5
1950	2,544	127	9	8	20	113	421	772	616	334	108	13	3
ALL OTHER, MALE													
1959	649	9	-	1	4	11	49	144	226	143	50	11	1
1958	852	16	1	1	3	13	59	184	287	196	75	15	2
1957	891	6	-	-	6	10	77	227	270	210	73	11	1
1956	899	7	2	1	6	10	80	225	306	191	62	7	2
1955	1,006	12	2	1	4	16	104	271	318	210	59	7	2
1954	1,119	21	-	-	6	30	104	318	366	213	47	14	-
1953	1,287	25	4	1	3	21	153	382	399	218	71	7	3
1952	1,369	36	1	2	5	25	173	418	414	223	61	10	1
1951	1,521	48	2	4	9	51	196	467	435	241	60	4	4
1950	1,786	71	7	3	8	69	269	566	467	243	75	8	-
ALL OTHER, FEMALE													
1959	314	4	1	1	2	5	22	85	100	64	28	2	-
1958	377	4	2	-	3	7	40	98	102	83	30	7	1
1957	394	4	-	-	3	15	40	101	115	74	31	10	1
1956	397	12	-	-	2	15	47	90	125	75	19	11	1
1955	390	13	1	1	1	14	39	111	96	79	27	8	-
1954	478	10	-	1	10	29	73	132	117	79	23	4	-
1953	544	17	2	2	4	28	91	151	130	92	21	6	-
1952	563	24	3	2	14	30	91	152	120	91	31	3	2
1951	638	41	3	4	12	24	108	180	147	87	23	8	1
1950	758	56	2	5	12	44	152	206	149	91	33	5	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DYSENTERY, ALL FORMS (045-048)													
BOTH SEXES													
1959	402	171	52	18	6	11	21	19	29	29	34	12	-
1958	407	187	51	15	7	9	14	21	22	41	30	10	-
1957	440	217	54	16	7	6	16	14	26	39	31	14	-
1956	476	216	39	26	5	6	13	24	38	46	50	11	2
1955	556	271	64	27	6	10	15	16	30	50	41	26	-
1954	655	301	75	16	7	9	26	37	44	60	56	23	1
1953	882	487	109	31	11	18	18	37	35	54	55	26	1
1952	937	517	115	29	16	14	25	32	51	53	54	31	-
1951	1,040	563	153	25	16	14	24	37	47	70	71	20	-
1950	923	521	116	25	10	24	30	32	52	51	42	20	-
MALE													
1959	214	88	25	12	3	5	12	14	18	14	20	3	-
1958	245	117	29	7	1	5	9	14	12	29	16	6	-
1957	227	109	27	10	6	4	8	8	12	22	16	5	-
1956	260	113	20	17	3	2	9	13	27	27	22	5	2
1955	291	138	38	13	4	4	13	7	18	28	16	12	-
1954	353	156	45	6	2	6	15	20	21	43	27	12	-
1953	480	271	58	17	5	8	8	26	21	30	26	9	1
1952	503	283	62	17	6	6	18	18	24	29	29	11	-
1951	555	317	72	13	9	9	12	22	27	35	33	6	-
1950	541	306	62	17	5	16	20	21	34	30	19	11	-
FEMALE													
1959	188	83	27	6	3	6	9	5	11	15	14	9	-
1958	162	70	22	8	6	4	5	7	10	12	14	4	-
1957	213	108	27	6	1	2	8	6	14	17	15	9	-
1956	216	103	19	9	2	4	4	11	11	19	28	6	-
1955	265	133	26	14	2	6	2	9	12	22	25	14	-
1954	302	145	30	10	5	3	11	17	23	17	29	11	1
1953	402	216	51	14	6	10	10	11	14	24	29	17	-
1952	434	234	53	12	10	8	7	14	27	24	25	20	-
1951	485	246	81	12	7	5	12	15	20	35	38	14	-
1950	382	215	54	8	5	8	10	11	18	21	23	9	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DYSENTERY, ALL FORMS (045-048)													
WHITE, BOTH SEXES													
1959	258	95	34	14	5	6	17	15	18	15	28	11	-
1958	263	111	30	12	3	7	10	9	15	30	26	10	-
1957	282	128	35	13	7	4	10	10	15	21	29	10	-
1956	307	124	24	20	5	3	12	18	18	33	40	10	-
1955	366	169	39	20	4	8	11	14	17	30	32	22	-
1954	400	167	39	11	5	6	15	19	31	37	51	19	-
1953	597	314	74	21	8	12	10	23	28	41	42	23	1
1952	630	328	65	23	12	11	14	23	42	43	45	24	-
1951	771	417	106	19	12	12	19	25	34	51	58	18	-
1950	693	401	85	20	7	17	20	16	40	36	35	16	-
WHITE, MALE													
1959	139	53	14	10	3	3	10	10	10	7	16	3	-
1958	153	67	14	5	1	4	7	8	7	19	15	6	-
1957	134	55	17	7	6	3	6	5	5	12	14	4	-
1956	162	63	12	14	3	-	8	10	13	19	15	5	-
1955	188	81	23	9	3	4	10	6	10	18	14	10	-
1954	212	88	26	2	2	4	9	12	13	26	22	8	-
1953	319	168	38	11	4	7	4	20	16	23	20	7	1
1952	334	181	37	13	5	4	9	13	19	22	23	8	-
1951	411	231	47	10	9	8	8	17	20	26	29	6	-
1950	403	238	44	13	5	11	13	10	27	20	15	7	-
WHITE, FEMALE													
1959	119	42	20	4	2	3	7	5	8	8	12	8	-
1958	110	44	16	7	2	3	3	1	8	11	11	4	-
1957	148	73	18	6	1	1	4	5	10	9	15	6	-
1956	145	61	12	6	2	3	4	8	5	14	25	5	-
1955	178	88	16	11	1	4	1	8	7	12	18	12	-
1954	188	79	13	9	3	2	6	7	18	11	29	11	-
1953	278	146	36	10	4	5	6	3	12	18	22	16	-
1952	296	147	28	10	7	7	5	10	23	21	22	16	-
1951	360	186	59	9	3	4	11	8	14	25	29	12	-
1950	290	163	41	7	2	6	7	6	13	16	20	9	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DYSENTERY, ALL FORMS (045-048)													
ALL OTHER, BOTH SEXES													
1959	144	76	18	4	1	5	4	4	11	14	6	1	-
1958	144	76	21	3	4	2	4	12	7	11	4	-	-
1957	158	89	19	3	-	2	6	4	11	18	2	4	-
1956	169	92	15	6	-	3	1	6	20	13	10	1	2
1955	190	102	25	7	2	2	4	2	13	20	9	4	-
1954	255	134	36	5	2	3	11	18	13	23	5	4	1
1953	285	173	35	10	3	6	8	14	7	13	13	3	-
1952	307	189	50	6	4	3	11	9	9	10	9	7	-
1951	269	146	47	6	4	2	5	12	13	19	13	2	-
1950	230	120	31	5	3	7	10	16	12	15	7	4	-
ALL OTHER, MALE													
1959	75	35	11	2	-	2	2	4	8	7	4	-	-
1958	92	50	15	2	-	1	2	6	5	10	1	-	-
1957	93	54	10	3	-	1	2	3	7	10	2	1	-
1956	98	50	8	3	-	2	1	3	14	8	7	-	2
1955	103	57	15	4	1	-	3	1	8	10	2	2	-
1954	141	68	19	4	-	2	6	8	8	17	5	4	-
1953	161	103	20	6	1	1	4	6	5	7	6	2	-
1952	169	102	25	4	1	2	9	5	5	7	6	3	-
1951	144	86	25	3	-	1	4	5	7	9	4	-	-
1950	138	68	18	4	-	5	7	11	7	10	4	4	-
ALL OTHER, FEMALE													
1959	69	41	7	2	1	3	2	-	3	7	2	1	-
1958	52	26	6	1	4	1	2	6	2	1	3	-	-
1957	65	35	9	-	-	1	4	1	4	8	-	3	-
1956	71	42	7	3	-	1	-	3	6	5	3	1	-
1955	87	45	10	3	1	2	1	1	5	10	7	2	-
1954	114	66	17	1	2	1	5	10	5	6	-	-	1
1953	124	70	15	4	2	5	4	8	2	6	7	1	-
1952	138	87	25	2	3	1	2	4	4	3	3	4	-
1951	125	60	22	3	4	1	1	7	6	10	9	2	-
1950	92	52	13	1	3	2	3	5	5	5	3	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)													
BOTH SEXES													
1959	130	15	18	23	13	7	10	9	11	11	9	4	-
1958	139	17	29	19	9	9	7	9	11	10	11	8	-
1957	213	24	27	28	12	15	22	24	24	15	15	7	-
1956	240	28	30	35	13	20	23	23	12	22	24	10	-
1955	235	28	40	32	15	19	21	20	19	19	16	6	-
1954	262	27	29	42	33	22	31	17	17	17	19	8	-
1953	295	35	43	43	29	30	21	30	14	27	16	7	-
1952	351	39	58	52	28	30	22	26	37	27	21	11	-
1951	347	38	57	37	29	36	32	31	26	21	24	15	1
1950	346	29	70	70	33	16	31	21	21	24	23	8	-
MALE													
1959	68	6	10	12	6	5	6	6	8	6	1	2	-
1958	69	8	20	7	5	5	3	5	4	5	5	2	-
1957	98	13	14	15	6	4	10	11	10	8	5	2	-
1956	121	12	14	22	6	10	16	11	5	12	8	5	-
1955	128	19	23	16	10	12	7	13	8	11	5	4	-
1954	130	14	15	20	17	12	20	9	7	5	5	6	-
1953	150	23	17	17	20	13	11	13	7	21	4	4	-
1952	186	20	34	22	19	17	14	13	23	12	9	3	-
1951	186	25	30	22	12	16	22	19	12	9	9	9	1
1950	182	20	40	39	16	8	20	13	7	9	7	3	-
FEMALE													
1959	62	9	8	11	7	2	4	3	3	5	8	2	-
1958	70	9	9	12	4	4	4	4	7	5	6	6	-
1957	115	11	13	13	6	11	12	13	14	7	10	5	-
1956	119	16	16	13	7	10	7	12	7	10	16	5	-
1955	107	9	17	16	5	7	14	7	11	8	11	2	-
1954	132	13	14	22	16	10	11	8	10	12	14	2	-
1953	145	12	26	26	9	17	10	17	7	6	12	3	-
1952	165	19	24	30	9	13	8	13	14	15	12	8	-
1951	161	13	27	15	17	20	10	12	14	12	15	6	-
1950	164	9	30	31	17	8	11	8	14	15	16	5	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)													
WHITE, BOTH SEXES													
1959	107	12	14	20	12	6	7	6	8	10	8	4	-
1958	111	10	22	16	8	7	4	6	10	9	11	8	-
1957	173	19	20	20	10	10	20	16	23	14	15	6	-
1956	196	19	22	23	11	18	18	19	12	20	24	10	-
1955	203	22	35	24	14	17	18	17	18	17	15	6	-
1954	217	15	23	39	25	18	26	15	14	15	19	8	-
1953	261	28	38	39	24	24	19	30	13	24	15	7	-
1952	303	30	47	46	26	25	18	19	34	26	21	11	-
1951	296	35	45	29	25	29	25	27	23	18	24	15	1
1950	294	23	59	65	25	11	22	17	18	23	23	8	-
WHITE, MALE													
1959	59	6	8	11	6	4	5	4	6	6	1	2	-
1958	57	6	15	6	5	4	2	4	3	5	5	2	-
1957	75	9	10	11	5	3	8	5	10	8	5	1	-
1956	95	8	11	12	5	9	12	9	5	11	8	5	-
1955	108	17	18	12	9	11	4	11	8	9	5	4	-
1954	112	10	12	19	13	11	18	9	5	4	5	6	-
1953	135	18	17	16	19	10	10	13	6	19	3	4	-
1952	158	14	26	19	19	12	13	10	22	11	9	3	-
1951	156	23	21	18	10	13	16	17	11	8	9	9	1
1950	161	16	34	38	13	8	15	11	7	9	7	3	-
WHITE, FEMALE													
1959	48	6	6	9	6	2	2	2	2	4	7	2	-
1958	54	4	7	10	3	3	2	2	7	4	6	6	-
1957	98	10	10	9	5	7	12	11	13	6	10	5	-
1956	101	11	11	11	6	9	6	10	7	9	16	5	-
1955	95	5	17	12	5	6	14	6	10	8	10	2	-
1954	105	5	11	20	12	7	8	6	9	11	14	2	-
1953	126	10	21	23	5	14	9	17	7	5	12	3	-
1952	145	16	21	27	7	13	5	9	12	15	12	8	-
1951	140	12	24	11	15	16	9	10	12	10	15	6	-
1950	133	7	25	27	12	3	7	6	11	14	16	5	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SCARLET FEVER AND STREPTOCOCCAL SORE THROAT (050,051)													
ALL OTHER, BOTH SEXES													
1959	23	3	4	3	1	1	3	3	3	1	1	-	-
1958	28	7	7	3	1	2	3	3	1	1	-	-	-
1957	40	5	7	8	2	5	2	8	1	1	-	1	-
1956	44	9	8	12	2	2	5	4	-	2	-	-	-
1955	32	6	5	8	1	2	3	3	1	2	1	-	-
1954	45	12	6	3	8	4	5	2	3	2	-	-	-
1953	34	7	5	4	5	6	2	-	1	3	1	-	-
1952	48	9	11	6	2	5	4	7	3	1	-	-	-
1951	51	3	12	8	4	7	7	4	3	3	-	-	-
1950	52	6	11	5	8	5	9	4	3	1	-	-	-
ALL OTHER, MALE													
1959	9	-	2	1	-	1	1	2	2	-	-	-	-
1958	12	2	5	1	-	1	1	1	1	-	-	-	-
1957	23	4	4	4	1	1	2	6	-	-	-	1	-
1956	26	4	3	10	1	1	4	2	-	1	-	-	-
1955	20	2	5	4	1	1	3	2	-	2	-	-	-
1954	18	4	3	1	4	1	2	-	2	1	-	-	-
1953	15	5	-	1	1	3	1	-	1	2	1	-	-
1952	28	6	8	3	-	5	1	3	1	1	-	-	-
1951	30	2	9	4	2	3	6	2	1	1	-	-	-
1950	21	4	6	1	3	-	5	2	-	-	-	-	-
ALL OTHER, FEMALE													
1959	14	3	2	2	1	-	2	1	1	1	1	-	-
1958	16	5	2	2	1	1	2	2	-	1	-	-	-
1957	17	1	3	4	1	4	-	2	1	1	-	-	-
1956	18	5	5	2	1	1	1	2	-	1	-	-	-
1955	12	4	-	4	-	1	-	1	1	-	1	-	-
1954	27	8	3	2	4	3	3	2	1	1	-	-	-
1953	19	2	5	3	4	3	1	-	-	1	-	-	-
1952	20	3	3	3	2	-	3	4	2	-	-	-	-
1951	21	1	3	4	2	4	1	2	2	2	-	-	-
1950	31	2	5	4	5	5	4	2	3	1	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SEPTICEMIA (053)													
BOTH SEXES													
1959	1,760	423	137	68	38	55	85	151	225	296	210	71	1
1958	1,663	433	139	47	46	58	84	120	210	249	205	71	1
1957	1,351	433	119	58	41	32	56	105	148	188	143	25	3
1956	1,118	329	144	37	23	45	53	64	118	165	105	34	1
1955	938	311	94	56	26	25	42	71	99	114	75	25	-
1954	897	293	101	45	28	38	46	71	78	108	65	24	-
1953	783	232	107	45	24	35	37	70	87	76	55	15	-
1952	726	246	100	41	25	34	36	57	67	57	46	17	-
1951	677	192	102	37	25	32	27	53	78	74	38	19	-
1950	616	179	78	30	24	26	48	50	54	57	49	19	2
MALE													
1959	982	237	78	42	19	24	38	81	146	171	111	34	1
1958	924	249	78	19	15	22	39	65	129	156	117	35	-
1957	805	251	64	36	22	11	34	64	92	127	87	14	3
1956	640	201	67	21	15	28	31	37	77	85	63	14	1
1955	538	186	51	27	16	9	19	39	62	72	45	12	-
1954	491	161	62	23	16	16	23	43	46	58	32	11	-
1953	467	140	62	26	10	17	19	40	65	54	27	7	-
1952	410	135	56	16	16	11	16	31	49	40	30	10	-
1951	367	108	59	20	11	9	14	30	36	51	20	9	-
1950	334	99	45	13	6	10	21	35	39	29	30	5	2
FEMALE													
1959	778	186	59	26	19	31	47	70	79	125	99	37	-
1958	739	184	61	28	31	36	45	55	81	93	88	36	1
1957	546	182	55	22	19	21	22	41	56	61	56	11	-
1956	478	128	77	16	8	17	22	27	41	80	42	20	-
1955	400	125	43	29	10	16	23	32	37	42	30	13	-
1954	406	132	39	22	12	22	23	28	32	50	33	13	-
1953	316	92	45	19	14	18	18	30	22	22	28	8	-
1952	316	111	44	25	9	23	20	26	18	17	16	7	-
1951	310	84	43	17	14	23	13	23	42	23	18	10	-
1950	282	80	33	17	18	16	27	15	15	28	19	14	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SEPTICEMIA (053)													
WHITE, BOTH SEXES													
1959	1,383	285	103	56	22	37	62	107	189	266	193	63	-
1958	1,343	307	111	40	33	47	60	96	176	221	186	66	-
1957	1,082	317	95	53	32	23	42	82	119	164	130	24	1
1956	902	243	118	25	20	36	35	46	96	154	97	31	1
1955	762	235	82	42	20	17	31	58	83	99	71	24	-
1954	728	220	86	40	23	27	36	55	60	96	61	24	-
1953	628	182	91	36	19	25	26	46	68	70	51	14	-
1952	589	186	83	33	18	24	33	45	56	52	43	16	-
1951	543	152	85	33	17	22	17	42	59	64	34	18	-
1950	504	157	68	25	15	14	30	41	47	45	43	18	1
WHITE, MALE													
1959	778	161	62	34	9	19	32	59	122	155	98	27	-
1958	749	181	64	13	11	19	27	51	110	135	106	32	-
1957	647	183	49	34	19	6	26	53	77	110	76	13	1
1956	531	155	57	16	14	23	21	28	66	81	57	12	1
1955	454	148	46	24	11	5	16	34	54	62	42	12	-
1954	405	125	54	21	14	14	16	35	34	52	29	11	-
1953	368	107	53	19	8	12	13	27	50	49	24	6	-
1952	344	105	47	13	12	8	16	28	42	37	27	9	-
1951	301	85	49	18	7	6	9	25	30	44	19	9	-
1950	286	87	41	10	4	7	16	30	35	24	26	5	1
WHITE, FEMALE													
1959	605	124	41	22	13	18	30	48	67	111	95	36	-
1958	594	126	47	27	22	28	33	45	66	86	80	34	-
1957	435	134	46	19	13	17	16	29	42	54	54	11	-
1956	371	88	61	9	6	13	14	18	30	73	40	19	-
1955	308	87	36	18	9	12	15	24	29	37	29	12	-
1954	323	95	32	19	9	13	20	20	26	44	32	13	-
1953	260	75	38	17	11	13	13	19	18	21	27	8	-
1952	245	81	36	20	6	16	17	17	14	15	16	7	-
1951	242	67	36	15	10	16	8	17	29	20	15	9	-
1950	218	70	27	15	11	7	14	11	12	21	17	13	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SEPTICEMIA (053)													
ALL OTHER, BOTH SEXES													
1959	377	138	34	12	16	18	23	44	36	30	17	8	1
1958	320	126	28	7	13	11	24	24	34	28	19	5	1
1957	269	116	24	5	9	9	14	23	29	24	13	1	2
1956	216	86	26	12	3	9	18	18	22	11	8	3	-
1955	176	76	12	14	6	8	11	13	16	15	4	1	-
1954	169	73	15	5	5	11	10	16	18	12	4	-	-
1953	155	50	16	9	5	10	11	24	19	6	4	1	-
1952	137	60	17	8	7	10	3	12	11	5	3	1	-
1951	134	40	17	4	8	10	10	11	19	10	4	1	-
1950	112	22	10	5	9	12	18	9	7	12	6	1	1
ALL OTHER, MALE													
1959	204	76	16	8	10	5	6	22	24	16	13	7	1
1958	175	68	14	6	4	3	12	14	19	21	11	3	-
1957	158	68	15	2	3	5	8	11	15	17	11	1	2
1956	109	46	10	5	1	5	10	9	11	4	6	2	-
1955	84	38	5	3	5	4	3	5	8	10	3	-	-
1954	86	36	8	2	2	2	7	8	12	6	3	-	-
1953	99	33	9	7	2	5	6	13	15	5	3	1	-
1952	66	30	9	3	4	3	-	3	7	3	3	1	-
1951	66	23	10	2	4	3	5	5	6	7	1	-	-
1950	48	12	4	3	2	3	5	5	4	5	4	-	1
ALL OTHER, FEMALE													
1959	173	62	18	4	6	13	17	22	12	14	4	1	-
1958	145	58	14	1	9	8	12	10	15	7	8	2	1
1957	111	48	9	3	6	4	6	12	14	7	2	-	-
1956	107	40	16	7	2	4	8	9	11	7	2	1	-
1955	92	38	7	11	1	4	8	8	8	5	1	1	-
1954	83	37	7	3	3	9	3	8	6	6	1	-	-
1953	56	17	7	2	3	5	5	11	4	1	1	-	-
1952	71	30	8	5	3	7	3	9	4	2	-	-	-
1951	68	17	7	2	4	7	5	6	13	3	3	1	-
1950	64	10	6	2	7	9	13	4	3	7	2	1	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DIPHTHERIA (055)													
BOTH SEXES													
1959	72	1	35	27	3	1	1	3	-	1	-	-	-
1958	74	3	27	33	1	3	1	3	2	1	-	-	-
1957	81	2	41	25	1	-	3	3	1	3	1	1	-
1956	103	7	40	36	-	2	4	11	2	-	1	-	-
1955	150	6	60	53	1	4	4	8	7	6	1	-	-
1954	145	6	69	46	-	2	11	6	2	3	-	-	-
1953	156	8	64	49	7	3	7	5	5	7	1	-	-
1952	217	12	96	55	3	8	11	11	12	6	3	-	-
1951	302	12	149	79	2	8	11	18	7	7	6	3	-
1950	410	21	210	115	8	3	17	17	5	10	3	1	-
MALE													
1959	39	1	19	13	2	1	-	2	-	1	-	-	-
1958	36	2	11	15	1	2	1	2	2	-	-	-	-
1957	36	1	21	8	1	-	1	1	-	1	1	1	-
1956	63	6	24	21	-	1	2	8	-	-	1	-	-
1955	85	4	32	25	1	3	3	7	5	4	1	-	-
1954	81	4	44	23	-	1	4	2	2	1	-	-	-
1953	90	7	35	22	4	3	6	4	3	5	1	-	-
1952	118	6	47	28	2	6	5	8	10	5	1	-	-
1951	179	7	90	44	-	4	8	15	4	4	2	1	-
1950	215	12	105	60	1	2	10	12	3	8	1	1	-
FEMALE													
1959	33	-	16	14	1	-	1	1	-	-	-	-	-
1958	38	1	16	18	-	1	-	1	-	1	-	-	-
1957	45	1	20	17	-	-	2	2	1	2	-	-	-
1956	40	1	16	15	-	1	2	3	2	-	-	-	-
1955	65	2	28	28	-	1	1	1	2	2	-	-	-
1954	64	2	25	23	-	1	7	4	-	2	-	-	-
1953	66	1	29	27	3	-	1	1	2	2	-	-	-
1952	99	6	49	27	1	2	6	3	2	1	2	-	-
1951	123	5	59	35	2	4	3	3	3	3	4	2	-
1950	195	9	105	55	7	1	7	5	2	2	2	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DIPHTHERIA (055)													
WHITE, BOTH SEXES													
1959	49	-	24	18	1	1	1	3	-	1	-	-	-
1958	47	2	14	23	1	2	1	3	1	-	-	-	-
1957	50	-	21	17	1	-	2	3	1	3	1	1	-
1956	63	5	18	22	-	1	4	10	2	-	1	-	-
1955	94	6	33	24	1	4	4	8	7	6	1	-	-
1954	91	4	31	34	-	1	11	6	2	2	-	-	-
1953	102	4	36	30	6	2	7	5	5	6	1	-	-
1952	161	7	63	40	2	7	10	11	12	6	3	-	-
1951	228	11	102	60	1	6	9	17	6	7	6	3	-
1950	323	17	156	89	6	2	17	17	5	10	3	1	-
WHITE, MALE													
1959	24	-	12	7	1	1	-	2	-	1	-	-	-
1958	24	1	6	11	1	1	1	2	1	-	-	-	-
1957	22	-	9	7	1	-	1	1	-	1	1	1	-
1956	36	4	12	10	-	-	2	7	-	-	1	-	-
1955	58	4	17	13	1	3	3	7	5	4	1	-	-
1954	50	3	19	19	-	1	4	2	2	-	-	-	-
1953	58	3	19	12	4	2	6	4	3	4	1	-	-
1952	85	2	29	18	2	6	4	8	10	5	1	-	-
1951	129	7	56	32	-	3	6	14	4	4	2	1	-
1950	172	10	79	46	1	1	10	12	3	8	1	1	-
WHITE, FEMALE													
1959	25	-	12	11	-	-	1	1	-	-	-	-	-
1958	23	1	8	12	-	1	-	1	-	-	-	-	-
1957	28	-	12	10	-	-	1	2	1	2	-	-	-
1956	27	1	6	12	-	1	2	3	2	-	-	-	-
1955	36	2	16	11	-	1	1	1	2	2	-	-	-
1954	41	1	12	15	-	-	7	4	-	2	-	-	-
1953	44	1	17	18	2	-	1	1	2	2	-	-	-
1952	76	5	34	22	-	1	6	3	2	1	2	-	-
1951	99	4	46	28	1	3	3	3	2	3	4	2	-
1950	151	7	77	43	5	1	7	5	2	2	2	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DIPHTHERIA (055)													
ALL OTHER, BOTH SEXES													
1959	23	1	11	9	2	-	-	-	-	-	-	-	-
1958	27	1	13	10	-	1	-	-	1	1	-	-	-
1957	31	2	20	8	-	-	1	-	-	-	-	-	-
1956	40	2	22	14	-	1	-	1	-	-	-	-	-
1955	56	-	27	29	-	-	-	-	-	-	-	-	-
1954	54	2	38	12	-	1	-	-	-	1	-	-	-
1953	54	4	28	19	1	1	-	-	-	1	-	-	-
1952	56	5	33	15	1	1	1	-	-	-	-	-	-
1951	74	1	47	19	1	2	2	1	1	-	-	-	-
1950	87	4	54	26	2	1	-	-	-	-	-	-	-
ALL OTHER, MALE													
1959	15	1	7	6	1	-	-	-	-	-	-	-	-
1958	12	1	5	4	-	1	-	-	1	-	-	-	-
1957	14	1	12	1	-	-	-	-	-	-	-	-	-
1956	27	2	12	11	-	1	-	1	-	-	-	-	-
1955	27	-	15	12	-	-	-	-	-	-	-	-	-
1954	31	1	25	4	-	-	-	-	-	1	-	-	-
1953	32	4	16	10	-	1	-	-	-	1	-	-	-
1952	33	4	18	10	-	-	1	-	-	-	-	-	-
1951	50	-	34	12	-	1	2	1	-	-	-	-	-
1950	43	2	26	14	-	1	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	8	-	4	3	1	-	-	-	-	-	-	-	-
1958	15	-	8	6	-	-	-	-	-	1	-	-	-
1957	17	1	8	7	-	-	1	-	-	-	-	-	-
1956	13	-	10	3	-	-	-	-	-	-	-	-	-
1955	29	-	12	17	-	-	-	-	-	-	-	-	-
1954	23	1	13	8	-	1	-	-	-	-	-	-	-
1953	22	-	12	9	1	-	-	-	-	-	-	-	-
1952	23	1	15	5	1	1	-	-	-	-	-	-	-
1951	24	1	13	7	1	1	-	-	1	-	-	-	-
1950	44	2	28	12	2	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
WHOOPING COUGH (056)													
BOTH SEXES													
1959	269	183	80	3	-	-	-	-	1	1	-	1	-
1958	177	130	43	3	-	-	-	-	1	-	-	-	-
1957	183	123	50	7	-	1	-	1	1	-	-	-	-
1956	266	172	80	9	1	3	-	-	-	1	-	-	-
1955	467	334	124	5	3	-	-	-	1	-	-	-	-
1954	373	263	103	6	-	-	1	-	-	-	-	-	-
1953	270	185	68	10	2	1	2	1	1	-	-	-	-
1952	402	262	121	12	3	1	2	1	-	-	-	-	-
1951	951	651	261	29	1	2	1	3	3	-	-	-	-
1950	1,118	745	328	39	3	-	-	-	-	2	1	-	-
MALE													
1959	103	71	30	1	-	-	-	-	1	-	-	-	-
1958	86	60	22	3	-	-	-	-	1	-	-	-	-
1957	83	53	25	2	-	1	-	1	1	-	-	-	-
1956	127	84	38	1	1	2	-	-	-	1	-	-	-
1955	205	154	47	2	2	-	-	-	-	-	-	-	-
1954	165	111	53	1	-	-	-	-	-	-	-	-	-
1953	117	85	27	3	-	-	1	-	1	-	-	-	-
1952	196	130	60	5	-	-	1	-	-	-	-	-	-
1951	475	352	102	15	1	1	-	2	2	-	-	-	-
1950	495	350	124	18	2	-	-	-	-	1	-	-	-
FEMALE													
1959	166	112	50	2	-	-	-	-	-	1	-	1	-
1958	91	70	21	-	-	-	-	-	-	-	-	-	-
1957	100	70	25	5	-	-	-	-	-	-	-	-	-
1956	139	88	42	8	-	1	-	-	-	-	-	-	-
1955	262	180	77	3	1	-	-	-	1	-	-	-	-
1954	208	152	50	5	-	-	1	-	-	-	-	-	-
1953	153	100	41	7	2	1	1	1	-	-	-	-	-
1952	206	132	61	7	3	1	1	1	-	-	-	-	-
1951	476	299	159	14	-	1	1	1	1	-	-	-	-
1950	623	395	204	21	1	-	-	-	-	1	1	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
WHOOPING COUGH (056)													
WHITE, BOTH SEXES													
1959	123	78	40	2	-	-	-	-	1	1	-	1	-
1958	91	66	22	2	-	-	-	-	1	-	-	-	-
1957	87	61	20	4	-	1	-	1	-	-	-	-	-
1956	101	65	26	5	1	3	-	-	-	1	-	-	-
1955	223	152	66	3	1	-	-	-	1	-	-	-	-
1954	222	162	56	3	-	-	1	-	-	-	-	-	-
1953	163	104	43	9	2	1	2	1	1	-	-	-	-
1952	206	124	69	6	3	1	2	1	-	-	-	-	-
1951	453	318	114	16	-	-	1	2	2	-	-	-	-
1950	705	476	200	24	2	-	-	-	-	2	1	-	-
WHITE, MALE													
1959	46	30	14	1	-	-	-	-	1	-	-	-	-
1958	44	29	12	2	-	-	-	-	1	-	-	-	-
1957	41	27	11	1	-	1	-	1	-	-	-	-	-
1956	50	32	13	1	1	2	-	-	-	1	-	-	-
1955	91	67	22	1	1	-	-	-	-	-	-	-	-
1954	95	65	30	-	-	-	-	-	-	-	-	-	-
1953	68	45	18	3	-	-	1	-	1	-	-	-	-
1952	99	59	36	3	-	-	1	-	-	-	-	-	-
1951	226	176	40	6	-	-	-	2	2	-	-	-	-
1950	304	222	68	12	1	-	-	-	-	1	-	-	-
WHITE, FEMALE													
1959	77	48	26	1	-	-	-	-	-	1	-	1	-
1958	47	37	10	-	-	-	-	-	-	-	-	-	-
1957	46	34	9	3	-	-	-	-	-	-	-	-	-
1956	51	33	13	4	-	1	-	-	-	-	-	-	-
1955	132	85	44	2	-	-	-	-	1	-	-	-	-
1954	127	97	26	3	-	-	1	-	-	-	-	-	-
1953	95	59	25	6	2	1	1	1	-	-	-	-	-
1952	107	65	33	3	3	1	1	1	-	-	-	-	-
1951	227	142	74	10	-	-	1	-	-	-	-	-	-
1950	401	254	132	12	1	-	-	-	-	1	1	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
WHOOPING COUGH (056)													
ALL OTHER, BOTH SEXES													
1959	146	105	40	1	-	-	-	-	-	-	-	-	-
1958	86	64	21	1	-	-	-	-	-	-	-	-	-
1957	96	62	30	3	-	-	-	-	1	-	-	-	-
1956	165	107	54	4	-	-	-	-	-	-	-	-	-
1955	244	182	58	2	2	-	-	-	-	-	-	-	-
1954	151	101	47	3	-	-	-	-	-	-	-	-	-
1953	107	81	25	1	-	-	-	-	-	-	-	-	-
1952	196	138	52	6	-	-	-	-	-	-	-	-	-
1951	498	333	147	13	1	2	-	1	1	-	-	-	-
1950	413	269	128	15	1	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1959	57	41	16	-	-	-	-	-	-	-	-	-	-
1958	42	31	10	1	-	-	-	-	-	-	-	-	-
1957	42	26	14	1	-	-	-	-	1	-	-	-	-
1956	77	52	25	-	-	-	-	-	-	-	-	-	-
1955	114	87	25	1	1	-	-	-	-	-	-	-	-
1954	70	46	23	1	-	-	-	-	-	-	-	-	-
1953	49	40	9	-	-	-	-	-	-	-	-	-	-
1952	97	71	24	2	-	-	-	-	-	-	-	-	-
1951	249	176	62	9	1	1	-	-	-	-	-	-	-
1950	191	128	56	6	1	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	89	64	24	1	-	-	-	-	-	-	-	-	-
1958	44	33	11	-	-	-	-	-	-	-	-	-	-
1957	54	36	16	2	-	-	-	-	-	-	-	-	-
1956	88	55	29	4	-	-	-	-	-	-	-	-	-
1955	130	95	33	1	1	-	-	-	-	-	-	-	-
1954	81	55	24	2	-	-	-	-	-	-	-	-	-
1953	58	41	16	1	-	-	-	-	-	-	-	-	-
1952	99	67	28	4	-	-	-	-	-	-	-	-	-
1951	249	157	85	4	-	1	-	1	1	-	-	-	-
1950	222	141	72	9	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MENINGOCOCCAL INFECTIONS (057)													
BOTH SEXES													
1959	699	203	238	87	43	25	19	21	28	20	9	4	2
1958	746	242	265	78	34	13	18	31	27	31	6	1	-
1957	785	245	278	98	45	17	18	21	25	29	5	4	-
1956	762	252	258	91	44	16	20	21	25	19	11	5	-
1955	907	257	331	104	60	23	24	37	30	23	13	5	-
1954	1,015	303	349	154	62	15	28	32	34	28	9	1	-
1953	1,325	393	452	192	92	28	31	51	48	25	13	-	-
1952	1,386	395	512	213	89	27	33	41	37	29	9	1	-
1951	1,124	303	397	141	96	24	35	52	32	32	12	-	-
1950	974	252	344	125	64	31	38	45	35	30	7	3	-
MALE													
1959	386	113	138	51	26	12	11	7	11	10	4	2	1
1958	378	130	134	38	16	4	9	15	16	15	1	-	-
1957	436	143	162	49	26	8	11	9	10	14	3	1	-
1956	435	146	155	43	24	12	11	14	13	8	7	2	-
1955	514	150	182	54	43	14	15	23	20	7	4	2	-
1954	571	171	195	87	45	8	15	16	18	13	2	1	-
1953	769	235	259	115	54	13	16	29	26	12	10	-	-
1952	755	211	293	108	49	16	15	22	18	16	6	1	-
1951	620	180	209	69	61	13	23	31	16	15	3	-	-
1950	559	139	205	67	37	19	22	28	22	16	3	1	-
FEMALE													
1959	313	90	100	36	17	13	8	14	17	10	5	2	1
1958	368	112	131	40	18	9	9	16	11	16	5	1	-
1957	349	102	116	49	19	9	7	12	15	15	2	3	-
1956	327	106	103	48	20	4	9	7	12	11	4	3	-
1955	393	107	149	50	17	9	9	14	10	16	9	3	-
1954	444	132	154	67	17	7	13	16	16	15	7	-	-
1953	556	158	193	77	38	15	15	22	22	13	3	-	-
1952	631	184	219	105	40	11	18	19	19	13	3	-	-
1951	504	123	188	72	35	11	12	21	16	17	9	-	-
1950	415	113	139	58	27	12	16	17	13	14	4	2	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MENINGOCOCCAL INFECTIONS (057)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	571	174	199	66	34	18	13	15	23	16	9	4	-
1958	619	202	224	61	29	11	13	20	24	29	5	1	-
1957	665	219	242	76	32	11	15	17	19	26	5	3	-
1956	641	212	229	72	33	14	14	12	21	18	11	5	-
1955	752	221	278	87	42	20	16	30	24	18	11	5	-
1954	860	262	308	119	45	10	21	30	30	25	9	1	-
1953	1,111	345	392	145	67	21	22	44	40	24	11	-	-
1952	1,185	351	448	174	70	22	22	30	33	27	8	-	-
1951	945	254	349	111	76	20	28	39	27	29	12	-	-
1950	843	218	313	108	51	26	31	31	28	27	7	3	-
<b>WHITE, MALE</b>													
1959	323	103	116	38	20	9	10	6	8	7	4	2	-
1958	310	104	114	31	14	4	7	9	13	13	1	-	-
1957	375	130	146	34	20	6	8	7	9	11	3	1	-
1956	366	121	137	39	17	10	7	7	11	8	7	2	-
1955	435	129	153	49	31	12	11	21	17	6	4	2	-
1954	480	150	172	67	32	5	11	14	15	11	2	1	-
1953	654	210	233	86	35	10	12	24	24	12	8	-	-
1952	653	192	257	87	40	13	9	16	18	15	6	-	-
1951	513	154	183	51	44	10	17	24	13	14	3	-	-
1950	475	119	186	53	31	14	18	18	17	15	3	1	-
<b>WHITE, FEMALE</b>													
1959	248	71	83	28	14	9	3	9	15	9	5	2	-
1958	309	98	110	30	15	7	6	11	11	16	4	1	-
1957	290	89	96	42	12	5	7	10	10	15	2	2	-
1956	275	91	92	33	16	4	7	5	10	10	4	3	-
1955	317	92	125	38	11	8	5	9	7	12	7	3	-
1954	380	112	136	52	13	5	10	16	15	14	7	-	-
1953	457	135	159	59	32	11	10	20	16	12	3	-	-
1952	532	159	191	87	30	9	13	14	15	12	2	-	-
1951	432	100	166	60	32	10	11	15	14	15	9	-	-
1950	368	99	127	55	20	12	13	13	11	12	4	2	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MENINGOCOCCAL INFECTIONS (057)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	128	29	39	21	9	7	6	6	5	4	-	-	2
1958	127	40	41	17	5	2	5	11	3	2	1	-	-
1957	120	26	36	22	13	6	3	4	6	3	-	1	-
1956	121	40	29	19	11	2	6	9	4	1	-	-	-
1955	155	36	53	17	18	3	8	7	6	5	2	-	-
1954	155	41	41	35	17	5	7	2	4	3	-	-	-
1953	214	48	60	47	25	7	9	7	8	1	2	-	-
1952	201	44	64	39	19	5	11	11	4	2	1	1	-
1951	179	49	48	30	20	4	7	13	5	3	-	-	-
1950	131	34	31	17	13	5	7	14	7	3	-	-	-
<b>ALL OTHER, MALE</b>													
1959	63	10	22	13	6	3	1	1	3	3	-	-	1
1958	68	26	20	7	2	-	2	6	3	2	-	-	-
1957	61	13	16	15	6	2	3	2	1	3	-	-	-
1956	69	25	18	4	7	2	4	7	2	-	-	-	-
1955	79	21	29	5	12	2	4	2	3	1	-	-	-
1954	91	21	23	20	13	3	4	2	3	2	-	-	-
1953	115	25	26	29	19	3	4	5	2	-	2	-	-
1952	102	19	36	21	9	3	6	6	-	1	-	1	-
1951	107	26	26	18	17	3	6	7	3	1	-	-	-
1950	84	20	19	14	6	5	4	10	5	1	-	-	-
<b>ALL OTHER, FEMALE</b>													
1959	65	19	17	8	3	4	5	5	2	1	-	-	1
1958	59	14	21	10	3	2	3	5	-	-	1	-	-
1957	59	13	20	7	7	4	-	2	5	-	-	1	-
1956	52	15	11	15	4	-	2	2	2	1	-	-	-
1955	76	15	24	12	6	1	4	5	3	4	2	-	-
1954	64	20	18	15	4	2	3	-	1	1	-	-	-
1953	99	23	34	18	6	4	5	2	6	1	-	-	-
1952	99	25	28	18	10	2	5	5	4	1	1	-	-
1951	72	23	22	12	3	1	1	6	2	2	-	-	-
1950	47	14	12	3	7	-	3	4	2	2	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ACUTE POLIOMYELITIS (080)													
BOTH SEXES													
1959	454	17	70	106	84	113	37	18	5	4	-	-	-
1958	255	16	41	46	55	64	20	8	4	1	-	-	-
1957	221	10	43	47	39	45	19	13	2	2	1	-	-
1956	566	29	98	110	124	140	43	16	4	-	2	-	-
1955	1,043	39	117	282	194	264	102	30	12	1	1	-	1
1954	1,368	69	185	394	228	335	99	37	16	5	-	-	-
1953	1,450	52	197	441	245	344	127	27	9	7	1	-	-
1952	3,145	116	464	980	595	712	205	43	22	8	-	-	-
1951	1,551	67	271	498	308	293	79	23	8	4	-	-	-
1950	1,904	81	260	602	387	394	126	35	14	4	1	-	-
MALE													
1959	312	13	45	75	53	84	24	12	3	3	-	-	-
1958	163	9	25	28	31	48	12	7	2	1	-	-	-
1957	128	7	26	25	22	26	10	8	2	2	-	-	-
1956	311	15	60	61	51	85	26	10	1	-	2	-	-
1955	580	24	60	154	97	151	66	18	8	-	1	-	1
1954	792	38	110	224	126	196	64	22	9	3	-	-	-
1953	851	31	106	273	148	186	73	20	8	5	1	-	-
1952	1,853	66	276	580	339	415	127	26	17	7	-	-	-
1951	929	35	158	319	173	167	55	13	7	2	-	-	-
1950	1,141	50	149	365	233	229	75	24	12	3	1	-	-
FEMALE													
1959	142	4	25	31	31	29	13	6	2	1	-	-	-
1958	92	7	16	18	24	16	8	1	2	-	-	-	-
1957	93	3	17	22	17	19	9	5	-	-	1	-	-
1956	255	14	38	49	73	55	17	6	3	-	-	-	-
1955	463	15	57	128	97	113	36	12	4	1	-	-	-
1954	576	31	75	170	102	139	35	15	7	2	-	-	-
1953	599	21	91	168	97	158	54	7	1	2	-	-	-
1952	1,292	50	188	400	256	297	78	17	5	1	-	-	-
1951	622	32	113	179	135	126	24	10	1	2	-	-	-
1950	763	31	111	237	154	165	51	11	2	1	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ACUTE POLIOMYELITIS (080)													
WHITE, BOTH SEXES													
1959	372	11	46	86	69	102	33	17	4	4	-	-	-
1958	212	10	30	39	48	54	18	8	4	1	-	-	-
1957	189	7	30	38	37	42	18	12	2	2	1	-	-
1956	484	21	71	88	111	133	39	16	3	-	2	-	-
1955	999	34	109	270	186	259	98	29	11	1	1	-	1
1954	1,299	59	170	372	221	325	96	36	15	5	-	-	-
1953	1,365	39	168	419	237	337	122	26	9	7	1	-	-
1952	3,008	106	420	944	573	696	200	41	20	8	-	-	-
1951	1,449	55	235	468	295	288	78	19	8	3	-	-	-
1950	1,811	73	235	579	371	380	120	34	14	4	1	-	-
WHITE, MALE													
1959	255	10	28	61	44	76	20	11	2	3	-	-	-
1958	136	6	18	25	27	40	10	7	2	1	-	-	-
1957	107	5	19	19	20	24	9	7	2	2	-	-	-
1956	263	8	45	49	45	80	23	10	1	-	2	-	-
1955	557	20	54	147	94	150	65	17	8	-	1	-	1
1954	753	31	103	211	123	190	62	21	9	3	-	-	-
1953	803	22	90	261	144	181	71	20	8	5	1	-	-
1952	1,769	56	250	556	329	408	123	25	15	7	-	-	-
1951	870	27	138	301	164	165	55	11	7	2	-	-	-
1950	1,094	45	137	356	222	222	73	23	12	3	1	-	-
WHITE, FEMALE													
1959	117	1	18	25	25	26	13	6	2	1	-	-	-
1958	76	4	12	14	21	14	8	1	2	-	-	-	-
1957	82	2	11	19	17	18	9	5	-	-	1	-	-
1956	221	13	26	39	66	53	16	6	2	-	-	-	-
1955	442	14	55	123	92	109	33	12	3	1	-	-	-
1954	546	28	67	161	98	135	34	15	6	2	-	-	-
1953	562	17	78	158	93	156	51	6	1	2	-	-	-
1952	1,239	50	170	388	244	288	77	16	5	1	-	-	-
1951	579	28	97	167	131	123	23	8	1	1	-	-	-
1950	717	28	98	223	149	158	47	11	2	1	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ACUTE POLIOMYELITIS (080)													
ALL OTHER, BOTH SEXES													
1959	82	6	24	20	15	11	4	1	1	-	-	-	-
1958	43	6	11	7	7	10	2	-	-	-	-	-	-
1957	32	3	13	9	2	3	1	1	-	-	-	-	-
1956	82	8	27	22	13	7	4	-	1	-	-	-	-
1955	44	5	8	12	8	5	4	1	1	-	-	-	-
1954	69	10	15	22	7	10	3	1	1	-	-	-	-
1953	85	13	29	22	8	7	5	1	-	-	-	-	-
1952	137	10	44	36	22	16	5	2	2	-	-	-	-
1951	102	12	36	30	13	5	1	4	-	1	-	-	-
1950	93	8	25	23	16	14	6	1	-	-	-	-	-
ALL OTHER, MALE													
1959	57	3	17	14	9	8	4	1	1	-	-	-	-
1958	27	3	7	3	4	8	2	-	-	-	-	-	-
1957	21	2	7	6	2	2	1	1	-	-	-	-	-
1956	48	7	15	12	6	5	3	-	-	-	-	-	-
1955	23	4	6	7	3	1	1	1	-	-	-	-	-
1954	39	7	7	13	3	6	2	1	-	-	-	-	-
1953	48	9	16	12	4	5	2	-	-	-	-	-	-
1952	84	10	26	24	10	7	4	1	2	-	-	-	-
1951	59	8	20	18	9	2	-	2	-	-	-	-	-
1950	47	5	12	9	11	7	2	1	-	-	-	-	-
ALL OTHER, FEMALE													
1959	25	3	7	6	6	3	-	-	-	-	-	-	-
1958	16	3	4	4	3	2	-	-	-	-	-	-	-
1957	11	1	6	3	-	1	-	-	-	-	-	-	-
1956	34	1	12	10	7	2	1	-	1	-	-	-	-
1955	21	1	2	5	5	4	3	-	1	-	-	-	-
1954	30	3	8	9	4	4	1	-	1	-	-	-	-
1953	37	4	13	10	4	2	3	1	-	-	-	-	-
1952	53	-	18	12	12	9	1	1	-	-	-	-	-
1951	43	4	16	12	4	3	1	2	-	1	-	-	-
1950	46	3	13	14	5	7	4	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MEASLES (085)</b>													
<b>BOTH SEXES</b>													
1959	385	72	180	101	8	8	4	3	2	3	3	1	-
1958	552	132	283	105	11	5	10	2	1	1	1	1	-
1957	389	77	175	110	11	5	1	1	-	3	5	1	-
1956	530	123	248	121	12	7	4	4	3	5	2	1	-
1955	345	81	153	82	9	6	6	2	-	4	1	1	-
1954	518	109	239	128	9	3	7	4	4	5	9	-	1
1953	462	123	203	96	16	6	6	3	4	2	1	2	-
1952	618	142	261	147	21	12	7	9	3	7	7	2	-
1951	683	162	304	147	20	18	11	6	2	4	4	4	1
1950	468	113	181	98	31	18	5	5	4	5	6	1	1
<b>MALE</b>													
1959	180	31	93	45	5	1	1	2	1	-	1	-	-
1958	289	71	147	52	6	4	5	1	1	1	-	1	-
1957	200	39	85	62	6	2	-	1	-	2	3	-	-
1956	257	56	138	45	5	3	3	2	3	1	-	1	-
1955	168	46	79	34	4	3	1	-	-	1	-	-	-
1954	243	50	118	53	7	2	4	1	2	4	2	-	-
1953	233	65	95	46	13	3	5	2	1	2	1	-	-
1952	295	78	117	74	6	4	4	5	2	2	3	-	-
1951	354	98	150	78	10	4	8	1	-	2	2	1	-
1950	236	53	97	46	21	7	3	3	3	1	1	1	-
<b>FEMALE</b>													
1959	205	41	87	56	3	7	3	1	1	3	2	1	-
1958	263	61	136	53	5	1	5	1	-	-	1	-	-
1957	189	38	90	48	5	3	1	-	-	1	2	1	-
1956	273	67	110	76	7	4	1	2	-	4	2	-	-
1955	177	35	74	48	5	3	5	2	-	3	1	1	-
1954	275	59	121	75	2	1	3	3	2	1	7	-	1
1953	229	58	108	50	3	3	1	1	3	-	-	2	-
1952	323	64	144	73	15	8	3	4	1	5	4	2	-
1951	329	64	154	69	10	14	3	5	2	2	2	3	1
1950	232	60	84	52	10	11	2	2	1	4	5	-	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MEASLES (085)													
WHITE, BOTH SEXES													
1959	293	52	129	81	8	8	4	3	2	2	3	1	-
1958	426	81	222	95	9	3	10	2	1	1	1	1	-
1957	301	48	132	95	11	5	-	1	-	3	5	1	-
1956	412	99	174	107	10	5	3	3	3	5	2	1	-
1955	255	51	109	69	9	5	5	2	-	3	1	1	-
1954	402	67	185	110	8	2	7	4	4	5	9	-	1
1953	384	96	167	86	14	6	6	3	2	2	1	1	-
1952	468	102	185	122	17	12	7	6	2	6	7	2	-
1951	545	117	245	123	14	16	11	5	2	4	4	4	-
1950	374	85	140	82	25	16	5	5	4	5	6	1	-
WHITE, MALE													
1959	137	21	68	37	5	1	1	2	1	-	1	-	-
1958	220	43	115	46	5	2	5	1	1	1	-	1	-
1957	154	26	61	53	6	2	-	1	-	2	3	-	-
1956	198	46	100	37	5	2	2	1	3	1	-	1	-
1955	126	30	58	30	4	3	1	-	-	-	-	-	-
1954	188	33	89	46	6	1	4	1	2	4	2	-	-
1953	202	53	81	42	12	3	5	2	1	2	1	-	-
1952	230	61	86	59	6	4	4	4	1	2	3	-	-
1951	288	73	125	64	8	4	8	1	-	2	2	1	-
1950	194	42	75	39	19	7	3	3	3	1	1	1	-
WHITE, FEMALE													
1959	156	31	61	44	3	7	3	1	1	2	2	1	-
1958	206	38	107	49	4	1	5	1	-	-	1	-	-
1957	147	22	71	42	5	3	-	-	-	1	2	1	-
1956	214	53	74	70	5	3	1	2	-	4	2	-	-
1955	129	21	51	39	5	2	4	2	-	3	1	1	-
1954	214	34	96	64	2	1	3	3	2	1	7	-	1
1953	182	43	86	44	2	3	1	1	1	-	-	1	-
1952	238	41	99	63	11	8	3	2	1	4	4	2	-
1951	257	44	120	59	6	12	3	4	2	2	2	3	-
1950	180	43	65	43	6	9	2	2	1	4	5	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MEASLES (085)													
ALL OTHER, BOTH SEXES													
1959	92	20	51	20	-	-	-	-	-	1	-	-	-
1958	126	51	61	10	2	2	-	-	-	-	-	-	-
1957	88	29	43	15	-	-	1	-	-	-	-	-	-
1956	118	24	74	14	2	2	1	1	-	-	-	-	-
1955	90	30	44	13	-	1	1	-	-	1	-	-	-
1954	116	42	54	18	1	1	-	-	-	-	-	-	-
1953	78	27	36	10	2	-	-	-	2	-	-	1	-
1952	150	40	76	25	4	-	-	3	1	1	-	-	-
1951	138	45	59	24	6	2	-	1	-	-	-	-	1
1950	94	28	41	16	6	2	-	-	-	-	-	-	1
ALL OTHER, MALE													
1959	43	10	25	8	-	-	-	-	-	-	-	-	-
1958	69	28	32	6	1	2	-	-	-	-	-	-	-
1957	46	13	24	9	-	-	-	-	-	-	-	-	-
1956	59	10	38	8	-	1	1	1	-	-	-	-	-
1955	42	16	21	4	-	-	-	-	-	1	-	-	-
1954	55	17	29	7	1	1	-	-	-	-	-	-	-
1953	31	12	14	4	1	-	-	-	-	-	-	-	-
1952	65	17	31	15	-	-	-	1	1	-	-	-	-
1951	66	25	25	14	2	-	-	-	-	-	-	-	-
1950	42	11	22	7	2	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	49	10	26	12	-	-	-	-	-	1	-	-	-
1958	57	23	29	4	1	-	-	-	-	-	-	-	-
1957	42	16	19	6	-	-	1	-	-	-	-	-	-
1956	59	14	36	6	2	1	-	-	-	-	-	-	-
1955	48	14	23	9	-	1	1	-	-	-	-	-	-
1954	61	25	25	11	-	-	-	-	-	-	-	-	-
1953	47	15	22	6	1	-	-	-	2	-	-	1	-
1952	85	23	45	10	4	-	-	2	-	1	-	-	-
1951	72	20	34	10	4	2	-	1	-	-	-	-	1
1950	52	17	19	9	4	2	-	-	-	-	-	-	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>OTHER INFECTIVE AND PARASITIC DISEASES</b>													
<b>BOTH SEXES</b>													
1959	5,485	884	477	401	237	334	405	585	707	755	513	185	2
1958	5,227	847	467	335	241	312	429	560	670	692	496	177	1
1957	4,721	819	407	328	227	298	355	564	586	589	416	128	4
1956	4,419	731	423	276	200	312	370	468	553	555	387	143	1
1955	4,151	650	356	316	200	268	371	456	511	529	341	151	2
1954	4,187	682	393	288	223	317	413	487	481	468	307	127	1
1953	4,190	642	421	349	218	324	371	443	505	488	312	116	1
1952	4,311	683	462	365	231	299	410	479	512	425	324	120	1
1951	4,069	607	434	319	219	300	384	469	489	457	276	112	3
1950	3,759	545	391	258	220	301	377	458	440	402	269	95	3
<b>MALE</b>													
1959	2,937	498	234	214	121	160	208	310	410	445	267	69	1
1958	2,797	490	248	170	107	126	207	309	404	402	262	72	-
1957	2,607	468	214	183	129	121	197	309	347	356	227	53	3
1956	2,406	421	220	154	89	161	198	251	344	307	203	57	1
1955	2,234	372	183	167	108	124	196	260	297	299	172	55	1
1954	2,280	382	225	168	119	149	211	286	283	259	140	57	1
1953	2,382	375	233	203	111	165	204	263	318	289	171	49	1
1952	2,343	373	255	198	111	133	197	275	321	244	177	58	1
1951	2,207	335	242	172	105	126	215	266	287	258	153	47	1
1950	2,123	301	226	147	117	141	197	279	288	239	146	40	2
<b>FEMALE</b>													
1959	2,548	386	243	187	116	174	197	275	297	310	246	116	1
1958	2,430	357	219	165	134	186	222	251	266	290	234	105	1
1957	2,114	351	193	145	98	177	158	255	239	233	189	75	1
1956	2,013	310	203	122	111	151	172	217	209	248	184	86	-
1955	1,917	278	173	149	92	144	175	196	214	230	169	96	1
1954	1,907	300	168	120	104	168	202	201	198	209	167	70	-
1953	1,808	267	188	146	107	159	167	180	187	199	141	67	-
1952	1,968	310	207	167	120	166	213	204	191	181	147	62	-
1951	1,862	272	192	147	114	174	169	203	202	199	123	65	2
1950	1,636	244	165	111	103	160	180	179	152	163	123	55	1

- Quantity zero

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and ICD Nos. 030-039, 041, 042, 044, 049, 052-.054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>OTHER INFECTIVE AND PARASITIC DISEASES</b>													
<b>WHITE, BOTH SEXES</b>													
1959	4,282	559	346	324	171	240	286	445	588	681	474	167	1
1958	4,214	567	365	274	187	239	326	437	573	619	458	169	-
1957	3,779	578	299	284	173	212	268	441	478	532	392	121	1
1956	3,489	498	322	207	141	221	272	373	451	500	366	137	1
1955	3,294	444	264	245	146	190	270	361	437	468	324	143	2
1954	3,296	483	314	228	147	220	303	389	390	413	287	121	1
1953	3,299	448	325	272	152	221	282	338	417	440	296	108	-
1952	3,429	490	345	287	171	228	318	366	422	382	303	116	1
1951	3,201	419	332	251	152	220	270	362	403	418	262	110	2
1950	2,947	400	307	197	147	206	280	351	373	349	244	92	1
<b>WHITE, MALE</b>													
1959	2,299	318	176	172	83	123	150	237	338	399	243	60	-
1958	2,267	335	199	133	82	99	158	250	345	357	241	68	-
1957	2,098	331	160	155	101	84	153	243	285	324	212	49	1
1956	1,919	299	165	118	68	122	145	208	273	277	189	54	1
1955	1,784	265	135	134	74	90	149	205	252	263	163	53	1
1954	1,780	273	184	130	76	102	156	229	220	226	129	54	1
1953	1,892	261	181	154	75	118	161	207	267	260	163	45	-
1952	1,882	271	192	157	84	98	165	213	264	221	162	54	1
1951	1,733	234	182	136	73	90	155	203	233	237	144	46	-
1950	1,696	220	181	117	77	98	158	219	245	210	130	40	1
<b>WHITE, FEMALE</b>													
1959	1,983	241	170	152	88	117	136	208	250	282	231	107	1
1958	1,947	232	166	141	105	140	168	187	228	262	217	101	-
1957	1,681	247	139	129	72	128	115	198	193	208	180	72	-
1956	1,570	199	157	89	73	99	127	165	178	223	177	83	-
1955	1,510	179	129	111	72	100	121	156	185	205	161	90	1
1954	1,516	210	130	98	71	118	147	160	170	187	158	67	-
1953	1,407	187	144	118	77	103	121	131	150	180	133	63	-
1952	1,547	219	153	130	87	130	153	153	158	161	141	62	-
1951	1,468	185	150	115	79	130	115	159	170	181	118	64	2
1950	1,251	180	126	80	70	108	122	132	128	139	114	52	-

- Quantity zero

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and ICD Nos. 030-039, 041, 042, 044, 049, 052-.054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>OTHER INFECTIVE AND PARASITIC DISEASES</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	1,203	325	131	77	66	94	119	140	119	74	39	18	1
1958	1,013	280	102	61	54	73	103	123	97	73	38	8	1
1957	942	241	108	44	54	86	87	123	108	57	24	7	3
1956	930	233	101	69	59	91	98	95	102	55	21	6	-
1955	857	206	92	71	54	78	101	95	74	61	17	8	-
1954	891	199	79	60	76	97	110	98	91	55	20	6	-
1953	891	194	96	77	66	103	89	105	88	48	16	8	1
1952	882	193	117	78	60	71	92	113	90	43	21	4	-
1951	868	188	102	68	67	80	114	107	86	39	14	2	1
1950	812	145	84	61	73	95	97	107	67	53	25	3	2
<b>ALL OTHER, MALE</b>													
1959	638	180	58	42	38	37	58	73	72	46	24	9	1
1958	530	155	49	37	25	27	49	59	59	45	21	4	-
1957	509	137	54	28	28	37	44	66	62	32	15	4	2
1956	487	122	55	36	21	39	53	43	71	30	14	3	-
1955	450	107	48	33	34	34	47	55	45	36	9	2	-
1954	500	109	41	38	43	47	55	57	63	33	11	3	-
1953	490	114	52	49	36	47	43	56	51	29	8	4	1
1952	461	102	63	41	27	35	32	62	57	23	15	4	-
1951	474	101	60	36	32	36	60	63	54	21	9	1	1
1950	427	81	45	30	40	43	39	60	43	29	16	-	1
<b>ALL OTHER, FEMALE</b>													
1959	565	145	73	35	28	57	61	67	47	28	15	9	-
1958	483	125	53	24	29	46	54	64	38	28	17	4	1
1957	433	104	54	16	26	49	43	57	46	25	9	3	1
1956	443	111	46	33	38	52	45	52	31	25	7	3	-
1955	407	99	44	38	20	44	54	40	29	25	8	6	-
1954	391	90	38	22	33	50	55	41	28	22	9	3	-
1953	401	80	44	28	30	56	46	49	37	19	8	4	-
1952	421	91	54	37	33	36	60	51	33	20	6	-	-
1951	394	87	42	32	35	44	54	44	32	18	5	1	-
1950	385	64	39	31	33	52	58	47	24	24	9	3	1

- Quantity zero

NOTE: Other infective and parasitic diseases include ICD Nos. 030-044, 049, 052-054, 058-074, 081-084, and 086-138 in 1959; and

ICD Nos. 030-039, 041, 042, 044, 049, 052-.054, 059-074, 081-083, 086-096, 120-138 in 1950-1958

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
MALIGNANT NEOPLASMS (ICD NOS. 140-205)													
BOTH SEXES													
1959	260,047	279	1,702	2,436	1,991	4,546	14,169	35,185	60,604	76,269	49,812	12,997	57
1958	254,426	261	1,719	2,277	1,998	4,516	14,197	34,415	59,397	74,685	48,486	12,411	64
1957	253,183	279	1,698	2,212	1,898	4,734	13,766	34,330	59,782	73,869	48,149	12,379	87
1956	247,357	301	1,669	2,247	1,839	4,771	14,027	33,272	59,178	70,855	47,298	11,825	75
1955	240,681	289	1,636	2,125	1,934	4,610	13,931	32,833	57,220	69,645	45,111	11,239	108
1954	234,669	306	1,613	1,963	1,881	4,714	13,820	32,603	56,003	67,410	43,699	10,554	103
1953	229,079	299	1,523	1,845	1,852	4,665	13,823	32,075	55,978	64,589	42,266	10,061	103
1952	223,277	291	1,597	1,726	1,957	4,844	13,537	31,415	54,875	62,576	40,836	9,518	105
1951	215,525	269	1,645	1,625	1,996	4,696	13,419	30,981	53,073	59,893	38,841	8,947	140
1950	210,733	274	1,521	1,635	1,907	4,763	13,440	30,372	52,232	58,277	37,802	8,371	139
MALE													
1959	139,078	150	932	1,432	1,205	2,058	5,641	16,770	33,733	43,972	26,898	6,258	29
1958	134,778	141	937	1,257	1,189	2,066	5,553	16,210	32,820	42,642	26,072	5,865	26
1957	134,400	142	959	1,279	1,116	2,084	5,334	15,960	33,076	42,611	25,891	5,909	39
1956	130,765	163	884	1,252	1,081	2,107	5,378	15,474	32,704	40,249	25,654	5,781	38
1955	126,268	161	915	1,272	1,146	2,036	5,122	15,256	31,262	39,504	24,153	5,387	54
1954	122,539	174	874	1,101	1,101	2,107	5,203	14,869	30,573	38,009	23,345	5,134	49
1953	118,827	154	837	1,075	1,073	2,024	5,248	14,677	30,522	35,901	22,422	4,849	45
1952	114,807	145	862	963	1,119	2,046	4,973	14,110	29,590	34,752	21,655	4,547	45
1951	110,034	146	933	906	1,182	2,012	4,817	13,879	28,312	33,022	20,433	4,324	68
1950	106,971	156	831	916	1,054	2,057	4,831	13,515	27,665	31,849	20,079	3,951	67
FEMALE													
1959	120,969	129	770	1,004	786	2,488	8,528	18,415	26,871	32,297	22,914	6,739	28
1958	119,648	120	782	1,020	809	2,450	8,644	18,205	26,577	32,043	22,414	6,546	38
1957	118,783	137	739	933	782	2,650	8,432	18,370	26,706	31,258	22,258	6,470	48
1956	116,592	138	785	995	758	2,664	8,649	17,798	26,474	30,606	21,644	6,044	37
1955	114,413	128	721	853	788	2,574	8,809	17,577	25,958	30,141	20,958	5,852	54
1954	112,130	132	739	862	780	2,607	8,617	17,734	25,430	29,401	20,354	5,420	54
1953	110,252	145	686	770	779	2,641	8,575	17,398	25,456	28,688	19,844	5,212	58
1952	108,470	146	735	763	838	2,798	8,564	17,305	25,285	27,824	19,181	4,971	60
1951	105,491	123	712	719	814	2,684	8,602	17,102	24,761	26,871	18,408	4,623	72
1950	103,762	118	690	719	853	2,706	8,609	16,857	24,567	26,428	17,723	4,420	72

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
MALIGNANT NEOPLASMS (ICD NOS. 140-205)													
WHITE, BOTH SEXES													
1959	236,641	239	1,553	2,220	1,738	3,901	12,095	30,696	54,086	70,448	47,268	12,358	39
1958	231,812	234	1,560	2,038	1,776	3,844	12,166	29,910	53,149	69,102	46,134	11,857	42
1957	230,953	256	1,560	2,003	1,681	4,127	11,822	29,792	53,543	68,404	45,866	11,836	63
1956	225,802	260	1,519	2,029	1,623	4,132	11,939	28,880	53,153	65,887	45,020	11,303	57
1955	220,428	257	1,508	1,937	1,719	3,999	11,862	28,491	51,750	64,907	43,146	10,770	82
1954	214,803	274	1,488	1,792	1,632	4,081	11,760	28,240	50,616	62,921	41,792	10,126	81
1953	210,293	261	1,401	1,696	1,639	4,049	11,846	27,832	50,996	60,296	40,545	9,659	73
1952	205,035	249	1,465	1,549	1,740	4,232	11,535	27,237	50,086	58,560	39,144	9,148	90
1951	198,160	246	1,509	1,501	1,778	4,072	11,478	26,934	48,441	56,126	37,362	8,595	118
1950	193,697	239	1,400	1,487	1,695	4,115	11,483	26,435	47,662	54,694	36,357	8,013	117
WHITE, MALE													
1959	126,478	128	852	1,312	1,071	1,818	4,861	14,627	30,114	40,420	25,361	5,896	18
1958	123,011	130	860	1,133	1,073	1,815	4,819	14,130	29,440	39,306	24,698	5,590	17
1957	122,675	128	870	1,158	1,005	1,875	4,640	13,841	29,591	39,350	24,570	5,619	28
1956	119,657	142	805	1,133	959	1,877	4,694	13,504	29,352	37,361	24,291	5,508	31
1955	115,910	143	849	1,162	1,024	1,832	4,434	13,317	28,204	36,756	23,011	5,136	42
1954	112,350	152	807	1,010	952	1,895	4,504	12,918	27,612	35,330	22,224	4,909	37
1953	109,410	133	770	988	957	1,841	4,606	12,791	27,763	33,404	21,473	4,650	34
1952	105,790	123	789	864	1,008	1,856	4,327	12,305	26,956	32,486	20,677	4,359	40
1951	101,605	134	864	850	1,062	1,800	4,263	12,132	25,801	30,896	19,604	4,145	54
1950	98,799	135	766	829	959	1,848	4,238	11,817	25,301	29,843	19,227	3,780	56
WHITE, FEMALE													
1959	110,163	111	701	908	667	2,083	7,234	16,069	23,972	30,028	21,907	6,462	21
1958	108,801	104	700	905	703	2,029	7,347	15,780	23,709	29,796	21,436	6,267	25
1957	108,278	128	690	845	676	2,252	7,182	15,951	23,952	29,054	21,296	6,217	35
1956	106,145	118	714	896	664	2,255	7,245	15,376	23,801	28,526	20,729	5,795	26
1955	104,518	114	659	775	695	2,167	7,428	15,174	23,546	28,151	20,135	5,634	40
1954	102,453	122	681	782	680	2,186	7,256	15,322	23,004	27,591	19,568	5,217	44
1953	100,883	128	631	708	682	2,208	7,240	15,041	23,233	26,892	19,072	5,009	39
1952	99,245	126	676	685	732	2,376	7,208	14,932	23,130	26,074	18,467	4,789	50
1951	96,555	112	645	651	716	2,272	7,215	14,802	22,640	25,230	17,758	4,450	64
1950	94,898	104	634	658	736	2,267	7,245	14,618	22,361	24,851	17,130	4,233	61

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
MALIGNANT NEOPLASMS (ICD NOS. 140-205)													
ALL OTHER, BOTH SEXES													
1959	23,406	40	149	216	253	645	2,074	4,489	6,518	5,821	2,544	639	18
1958	22,614	27	159	239	222	672	2,031	4,505	6,248	5,583	2,352	554	22
1957	22,230	23	138	209	217	607	1,944	4,538	6,239	5,465	2,283	543	24
1956	21,555	41	150	218	216	639	2,088	4,392	6,025	4,968	2,278	522	18
1955	20,253	32	128	188	215	611	2,069	4,342	5,470	4,738	1,965	469	26
1954	19,866	32	125	171	249	633	2,060	4,363	5,387	4,489	1,907	428	22
1953	18,786	38	122	149	213	616	1,977	4,243	4,982	4,293	1,721	402	30
1952	18,242	42	132	177	217	612	2,002	4,178	4,789	4,016	1,692	370	15
1951	17,365	23	136	124	218	624	1,941	4,047	4,632	3,767	1,479	352	22
1950	17,036	35	121	148	212	648	1,957	3,937	4,570	3,583	1,445	358	22
ALL OTHER, MALE													
1959	12,600	22	80	120	134	240	780	2,143	3,619	3,552	1,537	362	11
1958	11,767	11	77	124	116	251	734	2,080	3,380	3,336	1,374	275	9
1957	11,725	14	89	121	111	209	694	2,119	3,485	3,261	1,321	290	11
1956	11,108	21	79	119	122	230	684	1,970	3,352	2,888	1,363	273	7
1955	10,358	18	66	110	122	204	688	1,939	3,058	2,748	1,142	251	12
1954	10,189	22	67	91	149	212	699	1,951	2,961	2,679	1,121	225	12
1953	9,417	21	67	87	116	183	642	1,886	2,759	2,497	949	199	11
1952	9,017	22	73	99	111	190	646	1,805	2,634	2,266	978	188	5
1951	8,429	12	69	56	120	212	554	1,747	2,511	2,126	829	179	14
1950	8,172	21	65	87	95	209	593	1,698	2,364	2,006	852	171	11
ALL OTHER, FEMALE													
1959	10,806	18	69	96	119	405	1,294	2,346	2,899	2,269	1,007	277	7
1958	10,847	16	82	115	106	421	1,297	2,425	2,868	2,247	978	279	13
1957	10,505	9	49	88	106	398	1,250	2,419	2,754	2,204	962	253	13
1956	10,447	20	71	99	94	409	1,404	2,422	2,673	2,080	915	249	11
1955	9,895	14	62	78	93	407	1,381	2,403	2,412	1,990	823	218	14
1954	9,677	10	58	80	100	421	1,361	2,412	2,426	1,810	786	203	10
1953	9,369	17	55	62	97	433	1,335	2,357	2,223	1,796	772	203	19
1952	9,225	20	59	78	106	422	1,356	2,373	2,155	1,750	714	182	10
1951	8,936	11	67	68	98	412	1,387	2,300	2,121	1,641	650	173	8
1950	8,864	14	56	61	117	439	1,364	2,239	2,206	1,577	593	187	11

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)</b>													
<b>BOTH SEXES</b>													
1959	6,053	3	10	18	31	54	252	1,021	1,537	1,684	1,130	311	2
1958	5,766	-	2	17	23	55	252	898	1,525	1,577	1,092	325	-
1957	5,861	1	6	17	26	53	233	907	1,427	1,696	1,155	337	3
1956	5,589	2	8	20	23	58	224	831	1,455	1,550	1,109	307	2
1955	5,522	3	10	18	22	54	260	799	1,384	1,610	1,054	307	1
1954	5,382	5	8	11	21	45	241	779	1,349	1,605	1,016	300	2
1953	5,079	2	7	10	19	39	214	702	1,318	1,455	1,020	290	3
1952	5,029	-	3	10	25	59	216	703	1,299	1,459	978	272	5
1951	4,890	2	7	11	30	38	220	674	1,272	1,426	966	241	3
1950	5,138	1	2	13	23	55	227	702	1,300	1,492	1,066	251	6
<b>MALE</b>													
1959	4,710	2	7	11	23	43	170	806	1,254	1,360	814	219	1
1958	4,421	-	1	11	19	30	168	684	1,240	1,257	790	221	-
1957	4,609	1	4	14	17	34	172	713	1,170	1,399	850	233	2
1956	4,393	1	5	15	15	35	157	654	1,196	1,255	839	220	1
1955	4,340	1	5	10	14	34	179	614	1,151	1,304	809	218	1
1954	4,210	3	3	7	13	30	165	601	1,117	1,290	765	214	2
1953	3,995	2	4	5	13	23	153	540	1,094	1,187	767	205	2
1952	3,924	-	3	3	20	37	143	553	1,102	1,139	736	183	5
1951	3,842	1	5	7	17	21	157	517	1,062	1,140	747	165	3
1950	4,072	1	1	9	16	26	170	544	1,082	1,214	825	178	6
<b>FEMALE</b>													
1959	1,343	1	3	7	8	11	82	215	283	324	316	92	1
1958	1,345	-	1	6	4	25	84	214	285	320	302	104	-
1957	1,252	-	2	3	9	19	61	194	257	297	305	104	1
1956	1,196	1	3	5	8	23	67	177	259	295	270	87	1
1955	1,182	2	5	8	8	20	81	185	233	306	245	89	-
1954	1,172	2	5	4	8	15	76	178	232	315	251	86	-
1953	1,084	-	3	5	6	16	61	162	224	268	253	85	1
1952	1,105	-	-	7	5	22	73	150	197	320	242	89	-
1951	1,048	1	2	4	13	17	63	157	210	286	219	76	-
1950	1,066	-	1	4	7	29	57	158	218	278	241	73	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	5,529	3	9	13	26	43	196	885	1,389	1,582	1,085	297	1
1958	5,288	-	2	14	19	39	206	788	1,379	1,484	1,043	314	-
1957	5,341	1	6	15	22	41	191	779	1,272	1,585	1,107	322	-
1956	5,116	2	7	12	17	43	191	704	1,313	1,454	1,074	298	1
1955	5,103	2	8	15	17	48	217	697	1,273	1,509	1,022	294	1
1954	4,976	5	7	8	17	41	191	691	1,214	1,530	977	294	1
1953	4,715	1	7	10	17	33	180	606	1,207	1,381	987	285	1
1952	4,634	-	3	8	17	46	173	606	1,195	1,373	940	268	5
1951	4,503	2	7	11	21	31	173	565	1,154	1,355	946	235	3
1950	4,755	1	2	13	21	50	191	599	1,174	1,421	1,038	239	6
<b>WHITE, MALE</b>													
1959	4,305	2	6	7	19	36	133	695	1,135	1,283	779	210	-
1958	4,069	-	1	9	15	23	140	606	1,124	1,187	752	212	-
1957	4,199	1	4	13	15	25	140	610	1,044	1,306	814	227	-
1956	4,048	1	5	8	11	28	136	562	1,084	1,182	816	214	1
1955	4,036	1	5	8	12	30	149	543	1,056	1,235	786	210	1
1954	3,907	3	3	6	11	29	132	535	1,007	1,233	736	211	1
1953	3,725	1	4	5	12	20	132	474	1,008	1,126	741	201	1
1952	3,639	-	3	2	13	27	116	481	1,017	1,086	708	181	5
1951	3,560	1	5	7	13	16	124	441	969	1,090	731	160	3
1950	3,799	1	1	9	15	23	145	475	984	1,162	807	171	6
<b>WHITE, FEMALE</b>													
1959	1,224	1	3	6	7	7	63	190	254	299	306	87	1
1958	1,219	-	1	5	4	16	66	182	255	297	291	102	-
1957	1,142	-	2	2	7	16	51	169	228	279	293	95	-
1956	1,068	1	2	4	6	15	55	142	229	272	258	84	-
1955	1,067	1	3	7	5	18	68	154	217	274	236	84	-
1954	1,069	2	4	2	6	12	59	156	207	297	241	83	-
1953	990	-	3	5	5	13	48	132	199	255	246	84	-
1952	995	-	-	6	4	19	57	125	178	287	232	87	-
1951	943	1	2	4	8	15	49	124	185	265	215	75	-
1950	956	-	1	4	6	27	46	124	190	259	231	68	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF BUCCAL CAVITY AND PHARYNX (140-148)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	524	-	1	5	5	11	56	136	148	102	45	14	1
1958	478	-	-	3	4	16	46	110	146	93	49	11	-
1957	520	-	-	2	4	12	42	128	155	111	48	15	3
1956	473	-	1	8	6	15	33	127	142	96	35	9	1
1955	419	1	2	3	5	6	43	102	111	101	32	13	-
1954	406	-	1	3	4	4	50	88	135	75	39	6	1
1953	364	1	-	-	2	6	34	96	111	74	33	5	2
1952	395	-	-	2	8	13	43	97	104	86	38	4	-
1951	387	-	-	-	9	7	47	109	118	71	20	6	-
1950	383	-	-	-	2	5	36	103	126	71	28	12	-
<b>ALL OTHER, MALE</b>													
1959	405	-	1	4	4	7	37	111	119	77	35	9	1
1958	352	-	-	2	4	7	28	78	116	70	38	9	-
1957	410	-	-	1	2	9	32	103	126	93	36	6	2
1956	345	-	-	7	4	7	21	92	112	73	23	6	-
1955	304	-	-	2	2	4	30	71	95	69	23	8	-
1954	303	-	-	1	2	1	33	66	110	57	29	3	1
1953	270	1	-	-	1	3	21	66	86	61	26	4	1
1952	285	-	-	1	7	10	27	72	85	53	28	2	-
1951	282	-	-	-	4	5	33	76	93	50	16	5	-
1950	273	-	-	-	1	3	25	69	98	52	18	7	-
<b>ALL OTHER, FEMALE</b>													
1959	119	-	-	1	1	4	19	25	29	25	10	5	-
1958	126	-	-	1	-	9	18	32	30	23	11	2	-
1957	110	-	-	1	2	3	10	25	29	18	12	9	1
1956	128	-	1	1	2	8	12	35	30	23	12	3	1
1955	115	1	2	1	3	2	13	31	16	32	9	5	-
1954	103	-	1	2	2	3	17	22	25	18	10	3	-
1953	94	-	-	-	1	3	13	30	25	13	7	1	1
1952	110	-	-	1	1	3	16	25	19	33	10	2	-
1951	105	-	-	-	5	2	14	33	25	21	4	1	-
1950	110	-	-	-	1	2	11	34	28	19	10	5	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)													
BOTH SEXES													
1959	89,490	19	42	40	135	672	2,894	9,371	19,805	29,268	21,533	5,687	24
1958	88,144	24	53	35	135	642	2,857	9,163	19,544	29,240	20,975	5,447	29
1957	88,096	24	37	44	111	673	2,887	9,002	19,802	29,234	20,776	5,475	31
1956	87,740	30	38	41	141	687	3,043	9,124	20,036	28,514	20,884	5,178	24
1955	85,872	41	31	48	137	670	2,857	8,904	19,518	28,508	20,097	5,023	38
1954	85,493	41	33	51	140	712	3,063	9,250	19,630	28,174	19,588	4,780	31
1953	84,444	21	42	41	139	702	2,960	9,153	20,046	27,349	19,433	4,510	48
1952	83,238	30	49	34	155	745	3,007	9,137	19,961	26,770	19,023	4,282	45
1951	81,807	16	47	52	178	803	3,049	9,216	19,857	26,441	18,009	4,084	55
1950	81,726	17	47	43	168	808	3,074	9,262	20,057	26,149	18,185	3,869	47
MALE													
1959	49,052	8	26	27	71	350	1,500	5,313	11,542	16,728	11,022	2,453	12
1958	48,185	10	27	23	83	356	1,490	5,193	11,325	16,615	10,782	2,269	12
1957	48,394	13	20	31	61	331	1,512	5,045	11,517	16,811	10,662	2,377	14
1956	48,321	22	26	19	95	339	1,583	5,142	11,654	16,280	10,858	2,289	14
1955	47,304	33	20	33	87	344	1,417	5,046	11,362	16,404	10,354	2,181	23
1954	47,016	28	19	30	80	355	1,570	5,170	11,467	16,164	10,052	2,064	17
1953	46,511	12	25	28	82	355	1,538	5,147	11,800	15,574	9,967	1,961	22
1952	45,771	15	31	19	84	353	1,556	5,071	11,680	15,362	9,740	1,835	25
1951	44,905	12	31	34	102	374	1,524	5,067	11,605	15,092	9,213	1,818	33
1950	44,988	8	24	25	96	426	1,578	5,183	11,707	14,848	9,396	1,673	24
FEMALE													
1959	40,438	11	16	13	64	322	1,394	4,058	8,263	12,540	10,511	3,234	12
1958	39,959	14	26	12	52	286	1,367	3,970	8,219	12,625	10,193	3,178	17
1957	39,702	11	17	13	50	342	1,375	3,957	8,285	12,423	10,114	3,098	17
1956	39,419	8	12	22	46	348	1,460	3,982	8,382	12,234	10,026	2,889	10
1955	38,568	8	11	15	50	326	1,440	3,858	8,156	12,104	9,743	2,842	15
1954	38,477	13	14	21	60	357	1,493	4,080	8,163	12,010	9,536	2,716	14
1953	37,933	9	17	13	57	347	1,422	4,006	8,246	11,775	9,466	2,549	26
1952	37,467	15	18	15	71	392	1,451	4,066	8,281	11,408	9,283	2,447	20
1951	36,902	4	16	18	76	429	1,525	4,149	8,252	11,349	8,796	2,266	22
1950	36,738	9	23	18	72	382	1,496	4,079	8,350	11,301	8,789	2,196	23

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)													
WHITE, BOTH SEXES													
1959	81,475	17	39	35	104	535	2,408	8,005	17,425	26,974	20,485	5,431	17
1958	80,326	22	47	21	108	532	2,436	7,736	17,201	26,984	19,993	5,226	20
1957	80,451	23	35	34	87	558	2,436	7,617	17,508	27,075	19,823	5,230	25
1956	80,261	23	31	38	109	565	2,535	7,769	17,766	26,550	19,906	4,952	17
1955	78,732	33	27	36	116	554	2,397	7,526	17,399	26,496	19,287	4,832	29
1954	78,332	38	30	43	109	595	2,580	7,805	17,587	26,230	18,706	4,584	25
1953	77,714	15	34	35	114	601	2,502	7,771	18,061	25,525	18,674	4,344	38
1952	76,561	26	43	29	123	644	2,532	7,700	18,039	24,998	18,266	4,125	36
1951	75,443	12	42	48	146	672	2,566	7,877	17,989	24,772	17,335	3,938	46
1950	75,273	14	40	39	144	665	2,596	7,831	18,133	24,553	17,510	3,706	42
WHITE, MALE													
1959	44,187	7	23	22	52	276	1,228	4,478	10,069	15,298	10,414	2,313	7
1958	43,585	10	25	14	65	291	1,259	4,368	9,936	15,203	10,238	2,168	8
1957	43,791	12	19	25	49	266	1,272	4,208	10,093	15,488	10,103	2,246	10
1956	43,788	16	22	17	69	273	1,313	4,333	10,232	15,061	10,266	2,175	11
1955	42,968	27	17	24	73	286	1,163	4,232	10,020	15,143	9,880	2,085	18
1954	42,648	26	18	26	62	294	1,301	4,299	10,202	14,917	9,528	1,961	14
1953	42,472	9	21	24	67	312	1,306	4,330	10,564	14,427	9,524	1,870	18
1952	41,731	13	27	16	70	304	1,295	4,223	10,469	14,255	9,287	1,751	21
1951	41,162	10	26	33	84	308	1,300	4,290	10,433	14,086	8,821	1,744	27
1950	41,170	8	20	24	84	351	1,330	4,362	10,548	13,840	8,989	1,593	21
WHITE, FEMALE													
1959	37,288	10	16	13	52	259	1,180	3,527	7,356	11,676	10,071	3,118	10
1958	36,741	12	22	7	43	241	1,177	3,368	7,265	11,781	9,755	3,058	12
1957	36,660	11	16	9	38	292	1,164	3,409	7,415	11,587	9,720	2,984	15
1956	36,473	7	9	21	40	292	1,222	3,436	7,534	11,489	9,640	2,777	6
1955	35,764	6	10	12	43	268	1,234	3,294	7,379	11,353	9,407	2,747	11
1954	35,684	12	12	17	47	301	1,279	3,506	7,385	11,313	9,178	2,623	11
1953	35,242	6	13	11	47	289	1,196	3,441	7,497	11,098	9,150	2,474	20
1952	34,830	13	16	13	53	340	1,237	3,477	7,570	10,743	8,979	2,374	15
1951	34,281	2	16	15	62	364	1,266	3,587	7,556	10,686	8,514	2,194	19
1950	34,103	6	20	15	60	314	1,266	3,469	7,585	10,713	8,521	2,113	21

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF DIGESTIVE ORGANS AND PERITONEUM, NOT SPECIFIED AS SECONDARY (150-156a, 157-159)													
ALL OTHER, BOTH SEXES													
1959	8,015	2	3	5	31	137	486	1,366	2,380	2,294	1,048	256	7
1958	7,818	2	6	14	27	110	421	1,427	2,343	2,256	982	221	9
1957	7,645	1	2	10	24	115	451	1,385	2,294	2,159	953	245	6
1956	7,479	7	7	3	32	122	508	1,355	2,270	1,964	978	226	7
1955	7,140	8	4	12	21	116	460	1,378	2,119	2,012	810	191	9
1954	7,161	3	3	8	31	117	483	1,445	2,043	1,944	882	196	6
1953	6,730	6	8	6	25	101	458	1,382	1,985	1,824	759	166	10
1952	6,677	4	6	5	32	101	475	1,437	1,922	1,772	757	157	9
1951	6,364	4	5	4	32	131	483	1,339	1,868	1,669	674	146	9
1950	6,453	3	7	4	24	143	478	1,431	1,924	1,596	675	163	5
ALL OTHER, MALE													
1959	4,865	1	3	5	19	74	272	835	1,473	1,430	608	140	5
1958	4,600	-	2	9	18	65	231	825	1,389	1,412	544	101	4
1957	4,603	1	1	6	12	65	240	837	1,424	1,323	559	131	4
1956	4,533	6	4	2	26	66	270	809	1,422	1,219	592	114	3
1955	4,336	6	3	9	14	58	254	814	1,342	1,261	474	96	5
1954	4,368	2	1	4	18	61	269	871	1,265	1,247	524	103	3
1953	4,039	3	4	4	15	43	232	817	1,236	1,147	443	91	4
1952	4,040	2	4	3	14	49	261	848	1,211	1,107	453	84	4
1951	3,743	2	5	1	18	66	224	777	1,172	1,006	392	74	6
1950	3,818	-	4	1	12	75	248	821	1,159	1,008	407	80	3
ALL OTHER, FEMALE													
1959	3,150	1	-	-	12	63	214	531	907	864	440	116	2
1958	3,218	2	4	5	9	45	190	602	954	844	438	120	5
1957	3,042	-	1	4	12	50	211	548	870	836	394	114	2
1956	2,946	1	3	1	6	56	238	546	848	745	386	112	4
1955	2,804	2	1	3	7	58	206	564	777	751	336	95	4
1954	2,793	1	2	4	13	56	214	574	778	697	358	93	3
1953	2,691	3	4	2	10	58	226	565	749	677	316	75	6
1952	2,637	2	2	2	18	52	214	589	711	665	304	73	5
1951	2,621	2	-	3	14	65	259	562	696	663	282	72	3
1950	2,635	3	3	3	12	68	230	610	765	588	268	83	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)</b>													
<b>BOTH SEXES</b>													
1959	37,413	4	9	13	34	260	1,656	6,138	12,216	12,026	4,357	695	5
1958	35,423	8	15	24	41	261	1,618	5,945	11,506	11,259	4,128	615	3
1957	33,895	4	16	16	51	248	1,474	5,651	11,252	10,640	3,927	605	11
1956	32,197	8	9	24	45	241	1,459	5,406	10,824	9,961	3,686	530	4
1955	29,914	6	8	20	58	244	1,350	5,156	9,988	9,097	3,471	505	11
1954	27,627	4	11	14	58	229	1,305	4,813	9,296	8,343	3,078	461	15
1953	26,388	10	12	15	57	225	1,306	4,696	9,171	7,567	2,883	435	11
1952	24,330	7	10	16	62	218	1,223	4,289	8,445	7,017	2,638	398	7
1951	22,513	2	10	27	68	191	1,121	4,218	7,735	6,373	2,385	372	11
1950	21,219	3	13	24	54	224	1,102	3,977	7,339	5,832	2,270	369	12
<b>MALE</b>													
1959	31,914	3	4	10	21	184	1,266	5,166	10,874	10,500	3,394	488	4
1958	30,125	4	7	13	31	194	1,223	5,048	10,291	9,768	3,156	388	2
1957	28,884	2	9	12	36	174	1,131	4,825	10,085	9,196	3,007	397	10
1956	27,357	6	5	14	35	170	1,135	4,654	9,603	8,572	2,804	357	2
1955	25,292	4	3	11	36	169	1,040	4,471	8,862	7,728	2,638	320	10
1954	23,257	2	5	10	46	165	1,032	4,132	8,264	7,060	2,232	297	12
1953	22,128	6	7	11	38	157	1,015	4,058	8,123	6,317	2,113	274	9
1952	20,093	2	7	10	39	153	958	3,667	7,381	5,736	1,880	256	4
1951	18,480	1	4	15	39	127	854	3,583	6,728	5,250	1,653	219	7
1950	17,270	3	6	15	34	150	855	3,398	6,312	4,678	1,583	226	10
<b>FEMALE</b>													
1959	5,499	1	5	3	13	76	390	972	1,342	1,526	963	207	1
1958	5,298	4	8	11	10	67	395	897	1,215	1,491	972	227	1
1957	5,011	2	7	4	15	74	343	826	1,167	1,444	920	208	1
1956	4,840	2	4	10	10	71	324	752	1,221	1,389	882	173	2
1955	4,622	2	5	9	22	75	310	685	1,126	1,369	833	185	1
1954	4,370	2	6	4	12	64	273	681	1,032	1,283	846	164	3
1953	4,260	4	5	4	19	68	291	638	1,048	1,250	770	161	2
1952	4,237	5	3	6	23	65	265	622	1,064	1,281	758	142	3
1951	4,033	1	6	12	29	64	267	635	1,007	1,123	732	153	4
1950	3,949	-	7	9	20	74	247	579	1,027	1,154	687	143	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)													
WHITE, BOTH SEXES													
1959	34,328	2	9	12	29	223	1,389	5,414	11,154	11,302	4,133	656	5
1958	32,612	7	14	23	37	223	1,359	5,256	10,579	10,572	3,954	586	2
1957	31,249	2	14	13	41	216	1,296	4,986	10,310	10,015	3,769	578	9
1956	29,774	7	8	20	42	205	1,252	4,809	9,955	9,426	3,536	510	4
1955	27,732	6	7	18	49	209	1,163	4,566	9,251	8,634	3,334	486	9
1954	25,630	3	10	10	50	191	1,125	4,272	8,637	7,923	2,961	438	10
1953	24,498	9	9	14	52	190	1,133	4,174	8,520	7,196	2,777	413	11
1952	22,630	7	9	13	52	187	1,046	3,840	7,881	6,663	2,546	380	6
1951	20,953	2	10	21	63	168	1,002	3,784	7,189	6,042	2,303	359	10
1950	19,829	3	13	22	45	184	973	3,574	6,883	5,558	2,201	362	11
WHITE, MALE													
1959	29,332	1	4	9	19	158	1,052	4,576	9,949	9,890	3,214	456	4
1958	27,758	4	7	13	28	162	1,021	4,462	9,488	9,187	3,018	367	1
1957	26,648	1	8	11	30	158	997	4,256	9,247	8,679	2,879	373	9
1956	25,322	5	5	12	33	145	981	4,141	8,849	8,120	2,685	344	2
1955	23,517	4	3	11	30	143	893	3,979	8,239	7,364	2,534	309	8
1954	21,581	1	5	6	39	136	893	3,689	7,670	6,717	2,137	281	7
1953	20,560	5	6	10	34	137	884	3,619	7,556	6,012	2,027	261	9
1952	18,698	2	7	7	34	130	821	3,288	6,899	5,457	1,809	241	3
1951	17,223	1	4	13	36	114	772	3,231	6,266	4,975	1,593	212	6
1950	16,156	3	6	13	29	125	756	3,066	5,927	4,463	1,534	224	10
WHITE, FEMALE													
1959	4,996	1	5	3	10	65	337	838	1,205	1,412	919	200	1
1958	4,854	3	7	10	9	61	338	794	1,091	1,385	936	219	1
1957	4,601	1	6	2	11	58	299	730	1,063	1,336	890	205	-
1956	4,452	2	3	8	9	60	271	668	1,106	1,306	851	166	2
1955	4,215	2	4	7	19	66	270	587	1,012	1,270	800	177	1
1954	4,049	2	5	4	11	55	232	583	967	1,206	824	157	3
1953	3,938	4	3	4	18	53	249	555	964	1,184	750	152	2
1952	3,932	5	2	6	18	57	225	552	982	1,206	737	139	3
1951	3,730	1	6	8	27	54	230	553	923	1,067	710	147	4
1950	3,673	-	7	9	16	59	217	508	956	1,095	667	138	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF RESPIRATORY SYSTEM, NOT SPECIFIED AS SECONDARY (160-164)													
ALL OTHER, BOTH SEXES													
1959	3,085	2	-	1	5	37	267	724	1,062	724	224	39	-
1958	2,811	1	1	1	4	38	259	689	927	687	174	29	1
1957	2,646	2	2	3	10	32	178	665	942	625	158	27	2
1956	2,423	1	1	4	3	36	207	597	869	535	150	20	-
1955	2,182	-	1	2	9	35	187	590	737	463	137	19	2
1954	1,997	1	1	4	8	38	180	541	659	420	117	23	5
1953	1,890	1	3	1	5	35	173	522	651	371	106	22	-
1952	1,700	-	1	3	10	31	177	449	564	354	92	18	1
1951	1,560	-	-	6	5	23	119	434	546	331	82	13	1
1950	1,390	-	-	2	9	40	129	403	456	274	69	7	1
ALL OTHER, MALE													
1959	2,582	2	-	1	2	26	214	590	925	610	180	32	-
1958	2,367	-	-	-	3	32	202	586	803	581	138	21	1
1957	2,236	1	1	1	6	16	134	569	838	517	128	24	1
1956	2,035	1	-	2	2	25	154	513	754	452	119	13	-
1955	1,775	-	-	-	6	26	147	492	623	364	104	11	2
1954	1,676	1	-	4	7	29	139	443	594	343	95	16	5
1953	1,568	1	1	1	4	20	131	439	567	305	86	13	-
1952	1,395	-	-	3	5	23	137	379	482	279	71	15	1
1951	1,257	-	-	2	3	13	82	352	462	275	60	7	1
1950	1,114	-	-	2	5	25	99	332	385	215	49	2	-
ALL OTHER, FEMALE													
1959	503	-	-	-	3	11	53	134	137	114	44	7	-
1958	444	1	1	1	1	6	57	103	124	106	36	8	-
1957	410	1	1	2	4	16	44	96	104	108	30	3	1
1956	388	-	1	2	1	11	53	84	115	83	31	7	-
1955	407	-	1	2	3	9	40	98	114	99	33	8	-
1954	321	-	1	-	1	9	41	98	65	77	22	7	-
1953	322	-	2	-	1	15	42	83	84	66	20	9	-
1952	305	-	1	-	5	8	40	70	82	75	21	3	-
1951	303	-	-	4	2	10	37	82	84	56	22	6	-
1950	276	-	-	-	4	15	30	71	71	59	20	5	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF BREAST (170)</b>													
<b>BOTH SEXES</b>													
1959	23,085	1	1	2	10	506	2,395	5,000	5,679	5,202	3,232	1,053	4
1958	22,675	3	-	1	18	449	2,506	4,939	5,470	5,127	3,198	961	3
1957	22,720	1	1	1	10	489	2,348	4,967	5,486	5,162	3,251	998	6
1956	22,261	6	1	2	12	457	2,499	4,749	5,447	5,005	3,126	950	7
1955	21,945	2	2	1	19	468	2,483	4,745	5,342	4,846	3,116	905	16
1954	20,909	2	-	1	16	504	2,292	4,525	5,112	4,714	2,899	838	6
1953	20,566	1	1	3	21	479	2,349	4,452	4,994	4,580	2,864	810	12
1952	19,892	3	3	3	18	495	2,328	4,250	5,086	4,331	2,647	717	11
1951	19,179	-	-	1	16	447	2,184	4,083	4,812	4,202	2,707	716	11
1950	18,973	-	-	1	19	463	2,276	4,111	4,720	4,177	2,517	673	16
<b>MALE</b>													
1959	214	-	-	-	-	4	16	31	47	62	39	14	1
1958	215	-	-	-	-	2	15	21	60	58	41	18	-
1957	261	-	-	-	-	1	12	32	61	93	50	11	1
1956	207	1	-	-	1	-	13	28	52	52	45	14	1
1955	237	1	-	-	-	3	12	30	66	70	39	15	1
1954	230	-	-	-	-	4	19	30	51	67	45	14	-
1953	224	1	1	-	1	2	18	33	58	62	37	11	-
1952	228	1	-	-	-	4	13	26	52	66	53	13	-
1951	221	-	-	-	1	5	13	35	64	67	28	8	-
1950	239	-	-	-	-	3	16	37	56	77	41	8	1
<b>FEMALE</b>													
1959	22,871	1	1	2	10	502	2,379	4,969	5,632	5,140	3,193	1,039	3
1958	22,460	3	-	1	18	447	2,491	4,918	5,410	5,069	3,157	943	3
1957	22,459	1	1	1	10	488	2,336	4,935	5,425	5,069	3,201	987	5
1956	22,054	5	1	2	11	457	2,486	4,721	5,395	4,953	3,081	936	6
1955	21,708	1	2	1	19	465	2,471	4,715	5,276	4,776	3,077	890	15
1954	20,679	2	-	1	16	500	2,273	4,495	5,061	4,647	2,854	824	6
1953	20,342	-	-	3	20	477	2,331	4,419	4,936	4,518	2,827	799	12
1952	19,664	2	3	3	18	491	2,315	4,224	5,034	4,265	2,594	704	11
1951	18,958	-	-	1	15	442	2,171	4,048	4,748	4,135	2,679	708	11
1950	18,734	-	-	1	19	460	2,260	4,074	4,664	4,100	2,476	665	15

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF BREAST (170)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	21,337	1	1	1	8	429	2,113	4,542	5,216	4,911	3,106	1,008	1
1958	21,001	1	-	1	13	376	2,204	4,512	5,039	4,858	3,075	920	2
1957	21,035	1	1	-	8	425	2,065	4,496	5,077	4,875	3,118	964	5
1956	20,574	6	1	2	9	400	2,184	4,274	5,047	4,734	3,007	903	7
1955	20,372	2	1	1	17	392	2,182	4,329	4,986	4,573	2,997	880	12
1954	19,348	2	-	1	10	412	1,986	4,112	4,757	4,459	2,796	808	5
1953	19,130	1	-	3	15	394	2,085	4,044	4,698	4,350	2,761	773	6
1952	18,490	3	3	2	10	427	2,049	3,888	4,753	4,114	2,537	694	10
1951	17,825	-	-	1	13	386	1,929	3,678	4,506	3,994	2,623	685	10
1950	17,661	-	-	1	17	400	2,035	3,737	4,418	3,954	2,436	648	15
<b>WHITE, MALE</b>													
1959	189	-	-	-	-	4	13	28	40	58	35	11	-
1958	195	-	-	-	-	2	13	17	55	52	39	17	-
1957	230	-	-	-	-	1	9	28	51	83	47	10	1
1956	181	1	-	-	1	-	12	22	47	46	40	11	1
1955	221	1	-	-	-	3	11	27	63	64	38	13	1
1954	196	-	-	-	-	4	14	24	42	62	39	11	-
1953	201	1	-	-	1	2	12	30	55	54	36	10	-
1952	201	1	-	-	-	3	11	21	45	59	49	12	-
1951	201	-	-	-	1	4	11	25	60	65	27	8	-
1950	216	-	-	-	-	2	13	34	47	72	40	7	1
<b>WHITE, FEMALE</b>													
1959	21,148	1	1	1	8	425	2,100	4,514	5,176	4,853	3,071	997	1
1958	20,806	1	-	1	13	374	2,191	4,495	4,984	4,806	3,036	903	2
1957	20,805	1	1	-	8	424	2,056	4,468	5,026	4,792	3,071	954	4
1956	20,393	5	1	2	8	400	2,172	4,252	5,000	4,688	2,967	892	6
1955	20,151	1	1	1	17	389	2,171	4,302	4,923	4,509	2,959	867	11
1954	19,152	2	-	1	10	408	1,972	4,088	4,715	4,397	2,757	797	5
1953	18,929	-	-	3	14	392	2,073	4,014	4,643	4,296	2,725	763	6
1952	18,289	2	3	2	10	424	2,038	3,867	4,708	4,055	2,488	682	10
1951	17,624	-	-	1	12	382	1,918	3,653	4,446	3,929	2,596	677	10
1950	17,445	-	-	1	17	398	2,022	3,703	4,371	3,882	2,396	641	14

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF BREAST (170)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	1,748	-	-	1	2	77	282	458	463	291	126	45	3
1958	1,674	2	-	-	5	73	302	427	431	269	123	41	1
1957	1,685	-	-	1	2	64	283	471	409	287	133	34	1
1956	1,687	-	-	-	3	57	315	475	400	271	119	47	-
1955	1,573	-	1	-	2	76	301	416	356	273	119	25	4
1954	1,561	-	-	-	6	92	306	413	355	255	103	30	1
1953	1,436	-	1	-	6	85	264	408	296	230	103	37	6
1952	1,402	-	-	1	8	68	279	362	333	217	110	23	1
1951	1,354	-	-	-	3	61	255	405	306	208	84	31	1
1950	1,312	-	-	-	2	63	241	374	302	223	81	25	1
<b>ALL OTHER, MALE</b>													
1959	25	-	-	-	-	-	3	3	7	4	4	3	1
1958	20	-	-	-	-	-	2	4	5	6	2	1	-
1957	31	-	-	-	-	-	3	4	10	10	3	1	-
1956	26	-	-	-	-	-	1	6	5	6	5	3	-
1955	16	-	-	-	-	-	1	3	3	6	1	2	-
1954	34	-	-	-	-	-	5	6	9	5	6	3	-
1953	23	-	1	-	-	-	6	3	3	8	1	1	-
1952	27	-	-	-	-	1	2	5	7	7	4	1	-
1951	20	-	-	-	-	1	2	10	4	2	1	-	-
1950	23	-	-	-	-	1	3	3	9	5	1	1	-
<b>ALL OTHER, FEMALE</b>													
1959	1,723	-	-	1	2	77	279	455	456	287	122	42	2
1958	1,654	2	-	-	5	73	300	423	426	263	121	40	1
1957	1,654	-	-	1	2	64	280	467	399	277	130	33	1
1956	1,661	-	-	-	3	57	314	469	395	265	114	44	-
1955	1,557	-	1	-	2	76	300	413	353	267	118	23	4
1954	1,527	-	-	-	6	92	301	407	346	250	97	27	1
1953	1,413	-	-	-	6	85	258	405	293	222	102	36	6
1952	1,375	-	-	1	8	67	277	357	326	210	106	22	1
1951	1,334	-	-	-	3	60	253	395	302	206	83	31	1
1950	1,289	-	-	-	2	62	238	371	293	218	80	24	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)</b>													
<b>BOTH SEXES</b>													
1959	38,109	3	16	43	211	792	2,644	4,996	7,654	10,810	8,546	2,387	7
1958	38,390	1	22	32	210	883	2,666	5,146	7,799	10,783	8,504	2,332	12
1957	38,472	1	26	36	200	944	2,628	5,376	7,853	10,699	8,405	2,283	21
1956	37,741	-	19	33	189	987	2,667	5,173	7,769	10,242	8,402	2,243	17
1955	37,509	2	22	34	209	949	2,840	5,312	7,830	10,237	7,961	2,093	20
1954	36,947	4	15	29	223	911	2,818	5,310	7,773	9,920	7,961	1,959	24
1953	36,241	3	13	29	183	956	2,890	5,336	7,675	9,759	7,534	1,848	15
1952	35,379	2	26	25	206	983	2,816	5,306	7,570	9,405	7,226	1,795	19
1951	34,491	1	16	26	225	952	2,935	5,381	7,415	9,008	6,946	1,558	28
1950	34,742	-	21	37	217	1,012	3,036	5,540	7,583	8,933	6,815	1,514	34
<b>MALE</b>													
1959	15,017	1	10	12	100	183	226	378	1,781	5,141	5,523	1,661	1
1958	15,062	1	15	5	112	207	217	398	1,825	5,155	5,506	1,617	4
1957	15,245	-	18	10	88	198	195	404	1,849	5,255	5,607	1,615	6
1956	14,946	-	10	11	83	231	188	382	1,853	4,988	5,597	1,597	6
1955	14,489	1	16	10	94	222	195	417	1,901	4,939	5,207	1,481	6
1954	14,235	3	9	6	113	231	187	389	1,876	4,805	5,199	1,411	6
1953	13,478	2	7	12	89	210	196	368	1,727	4,656	4,899	1,306	6
1952	13,097	2	12	7	93	198	175	408	1,822	4,447	4,658	1,268	7
1951	12,483	-	10	8	102	243	194	420	1,727	4,212	4,478	1,080	9
1950	12,305	-	13	10	88	228	181	413	1,754	4,179	4,390	1,033	16
<b>FEMALE</b>													
1959	23,092	2	6	31	111	609	2,418	4,618	5,873	5,669	3,023	726	6
1958	23,328	-	7	27	98	676	2,449	4,748	5,974	5,628	2,998	715	8
1957	23,227	1	8	26	112	746	2,433	4,972	6,004	5,444	2,798	668	15
1956	22,795	-	9	22	106	756	2,479	4,791	5,916	5,254	2,805	646	11
1955	23,020	1	6	24	115	727	2,645	4,895	5,929	5,298	2,754	612	14
1954	22,712	1	6	23	110	680	2,631	4,921	5,897	5,115	2,762	548	18
1953	22,763	1	6	17	94	746	2,694	4,968	5,948	5,103	2,635	542	9
1952	22,282	-	14	18	113	785	2,641	4,898	5,748	4,958	2,568	527	12
1951	22,008	1	6	18	123	709	2,741	4,961	5,688	4,796	2,468	478	19
1950	22,437	-	8	27	129	784	2,855	5,127	5,829	4,754	2,425	481	18

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	33,148	3	11	31	186	642	2,121	4,110	6,446	9,498	7,884	2,212	4
1958	33,479	1	19	27	188	706	2,135	4,212	6,589	9,540	7,891	2,164	7
1957	33,551	1	24	30	170	776	2,098	4,391	6,671	9,405	7,828	2,144	13
1956	32,861	-	18	30	162	809	2,102	4,225	6,525	9,072	7,794	2,115	9
1955	32,857	2	20	29	180	772	2,229	4,317	6,720	9,189	7,422	1,965	12
1954	32,410	3	15	25	194	743	2,248	4,324	6,612	8,910	7,468	1,852	16
1953	31,802	3	12	24	160	770	2,304	4,338	6,647	8,709	7,083	1,745	7
1952	31,071	1	23	22	181	784	2,229	4,270	6,564	8,499	6,795	1,686	17
1951	30,351	1	11	23	196	767	2,315	4,402	6,457	8,146	6,546	1,466	21
1950	30,573	-	18	36	181	816	2,388	4,557	6,592	8,116	6,431	1,418	20
<b>WHITE, MALE</b>													
1959	13,147	1	6	11	96	179	202	264	1,392	4,379	5,074	1,543	-
1958	13,356	1	14	5	110	201	195	313	1,430	4,473	5,102	1,509	3
1957	13,498	-	17	9	86	195	169	291	1,468	4,513	5,224	1,522	4
1956	13,320	-	10	10	80	224	169	282	1,433	4,382	5,214	1,512	4
1955	12,958	1	15	10	91	214	167	306	1,523	4,377	4,852	1,398	4
1954	12,741	2	9	6	110	225	168	271	1,470	4,251	4,880	1,346	3
1953	12,156	2	7	11	85	203	175	271	1,392	4,124	4,640	1,245	1
1952	11,837	1	11	6	88	189	154	287	1,492	4,026	4,371	1,205	7
1951	11,282	-	7	8	93	231	174	298	1,421	3,785	4,239	1,021	5
1950	11,165	-	12	10	85	222	161	287	1,464	3,798	4,140	977	9
<b>WHITE, FEMALE</b>													
1959	20,001	2	5	20	90	463	1,919	3,846	5,054	5,119	2,810	669	4
1958	20,123	-	5	22	78	505	1,940	3,899	5,159	5,067	2,789	655	4
1957	20,053	1	7	21	84	581	1,929	4,100	5,203	4,892	2,604	622	9
1956	19,541	-	8	20	82	585	1,933	3,943	5,092	4,690	2,580	603	5
1955	19,899	1	5	19	89	558	2,062	4,011	5,197	4,812	2,570	567	8
1954	19,669	1	6	19	84	518	2,080	4,053	5,142	4,659	2,588	506	13
1953	19,646	1	5	13	75	567	2,129	4,067	5,255	4,585	2,443	500	6
1952	19,234	-	12	16	93	595	2,075	3,983	5,072	4,473	2,424	481	10
1951	19,069	1	4	15	103	536	2,141	4,104	5,036	4,361	2,307	445	16
1950	19,408	-	6	26	96	594	2,227	4,270	5,128	4,318	2,291	441	11

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MALIGNANT NEOPLASM OF GENITAL ORGANS (171-179)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	4,961	-	5	12	25	150	523	886	1,208	1,312	662	175	3
1958	4,911	-	3	5	22	177	531	934	1,210	1,243	613	168	5
1957	4,921	-	2	6	30	168	530	985	1,182	1,294	577	139	8
1956	4,880	-	1	3	27	178	565	948	1,244	1,170	608	128	8
1955	4,652	-	2	5	29	177	611	995	1,110	1,048	539	128	8
1954	4,537	1	-	4	29	168	570	986	1,161	1,010	493	107	8
1953	4,439	-	1	5	23	186	586	998	1,028	1,050	451	103	8
1952	4,308	1	3	3	25	199	587	1,036	1,006	906	431	109	2
1951	4,140	-	5	3	29	185	620	979	958	862	400	92	7
1950	4,169	-	3	1	36	196	648	983	991	817	384	96	14
<b>ALL OTHER, MALE</b>													
1959	1,870	-	4	1	4	4	24	114	389	762	449	118	1
1958	1,706	-	1	-	2	6	22	85	395	682	404	108	1
1957	1,747	-	1	1	2	3	26	113	381	742	383	93	2
1956	1,626	-	-	1	3	7	19	100	420	606	383	85	2
1955	1,531	-	1	-	3	8	28	111	378	562	355	83	2
1954	1,494	1	-	-	3	6	19	118	406	554	319	65	3
1953	1,322	-	-	1	4	7	21	97	335	532	259	61	5
1952	1,260	1	1	1	5	9	21	121	330	421	287	63	-
1951	1,201	-	3	-	9	12	20	122	306	427	239	59	4
1950	1,140	-	1	-	3	6	20	126	290	381	250	56	7
<b>ALL OTHER, FEMALE</b>													
1959	3,091	-	1	11	21	146	499	772	819	550	213	57	2
1958	3,205	-	2	5	20	171	509	849	815	561	209	60	4
1957	3,174	-	1	5	28	165	504	872	801	552	194	46	6
1956	3,254	-	1	2	24	171	546	848	824	564	225	43	6
1955	3,121	-	1	5	26	169	583	884	732	486	184	45	6
1954	3,043	-	-	4	26	162	551	868	755	456	174	42	5
1953	3,117	-	1	4	19	179	565	901	693	518	192	42	3
1952	3,048	-	2	2	20	190	566	915	676	485	144	46	2
1951	2,939	-	2	3	20	173	600	857	652	435	161	33	3
1950	3,029	-	2	1	33	190	628	857	701	436	134	40	7

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)													
BOTH SEXES													
1959	12,493	18	194	105	19	72	317	1,206	2,696	4,069	3,006	789	2
1958	12,122	15	171	111	22	64	341	1,214	2,531	4,016	2,872	763	2
1957	12,147	26	168	94	25	71	348	1,222	2,730	3,869	2,832	759	3
1956	11,718	24	166	102	12	62	328	1,150	2,692	3,727	2,733	719	3
1955	11,502	18	162	102	19	77	305	1,124	2,623	3,751	2,659	661	1
1954	11,193	17	163	81	19	85	343	1,139	2,583	3,578	2,568	608	9
1953	10,915	21	140	92	23	64	360	1,168	2,627	3,490	2,344	584	2
1952	10,679	16	146	77	26	77	343	1,131	2,615	3,392	2,316	536	4
1951	10,158	28	167	76	27	77	295	1,140	2,498	3,132	2,169	544	5
1950	10,044	29	156	81	31	68	334	1,082	2,552	3,166	2,104	433	8
MALE													
1959	8,353	11	106	46	12	36	195	851	1,922	2,798	1,924	450	2
1958	8,036	8	94	48	12	40	218	824	1,802	2,761	1,813	414	2
1957	8,023	8	88	46	18	37	212	843	1,935	2,673	1,745	415	3
1956	7,786	18	82	45	6	33	194	796	1,945	2,566	1,699	400	2
1955	7,553	9	91	50	7	41	191	783	1,850	2,567	1,610	353	1
1954	7,434	10	96	36	8	56	225	781	1,868	2,408	1,587	354	5
1953	7,210	15	81	44	12	30	223	777	1,924	2,341	1,426	336	1
1952	7,068	11	64	33	14	41	215	796	1,881	2,277	1,432	303	1
1951	6,682	17	94	34	16	42	183	799	1,774	2,063	1,349	307	4
1950	6,595	14	82	30	17	38	203	756	1,802	2,097	1,278	273	5
FEMALE													
1959	4,140	7	88	59	7	36	122	355	774	1,271	1,082	339	-
1958	4,086	7	77	63	10	24	123	390	729	1,255	1,059	349	-
1957	4,124	18	80	48	7	34	136	379	795	1,196	1,087	344	-
1956	3,932	6	84	57	6	29	134	354	747	1,161	1,034	319	1
1955	3,949	9	71	52	12	36	114	341	773	1,184	1,049	308	-
1954	3,759	7	67	45	11	29	118	358	715	1,170	981	254	4
1953	3,705	6	59	48	11	34	137	391	703	1,149	918	248	1
1952	3,611	5	82	44	12	36	128	335	734	1,115	884	233	3
1951	3,476	11	73	42	11	35	112	341	724	1,069	820	237	1
1950	3,449	15	74	51	14	30	131	326	750	1,069	826	160	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)													
WHITE, BOTH SEXES													
1959	11,614	16	168	89	15	58	259	1,056	2,449	3,830	2,905	767	2
1958	11,258	14	148	94	15	47	286	1,026	2,316	3,788	2,779	745	-
1957	11,297	22	149	81	19	59	293	1,060	2,482	3,652	2,740	737	3
1956	10,929	21	144	83	12	44	268	1,002	2,477	3,541	2,636	698	3
1955	10,835	18	145	92	15	63	260	1,006	2,428	3,583	2,585	639	1
1954	10,455	17	146	67	14	70	292	982	2,377	3,399	2,496	587	8
1953	10,200	20	121	82	22	56	305	1,006	2,438	3,300	2,277	571	2
1952	10,004	11	133	64	21	68	289	1,002	2,419	3,224	2,242	528	3
1951	9,520	25	149	67	23	60	242	994	2,335	2,974	2,118	528	5
1950	9,407	26	136	72	26	51	273	960	2,386	3,003	2,051	415	8
WHITE, MALE													
1959	7,819	10	94	40	9	28	167	764	1,769	2,637	1,861	438	2
1958	7,511	8	83	41	9	30	188	713	1,666	2,618	1,751	404	-
1957	7,507	6	76	39	14	31	181	740	1,784	2,535	1,693	405	3
1956	7,315	15	72	37	6	25	167	713	1,808	2,450	1,633	387	2
1955	7,150	9	79	45	6	35	164	713	1,716	2,472	1,566	344	1
1954	6,989	10	85	29	5	48	196	688	1,731	2,301	1,550	341	5
1953	6,792	15	68	40	12	25	196	690	1,805	2,218	1,391	331	1
1952	6,656	7	57	30	10	33	185	723	1,747	2,176	1,386	301	1
1951	6,293	16	86	31	12	35	154	715	1,668	1,959	1,317	296	4
1950	6,226	11	72	25	15	32	174	683	1,707	2,002	1,238	262	5
WHITE, FEMALE													
1959	3,795	6	74	49	6	30	92	292	680	1,193	1,044	329	-
1958	3,747	6	65	53	6	17	98	313	650	1,170	1,028	341	-
1957	3,790	16	73	42	5	28	112	320	698	1,117	1,047	332	-
1956	3,614	6	72	46	6	19	101	289	669	1,091	1,003	311	1
1955	3,685	9	66	47	9	28	96	293	712	1,111	1,019	295	-
1954	3,466	7	61	38	9	22	96	294	646	1,098	946	246	3
1953	3,408	5	53	42	10	31	109	316	633	1,082	886	240	1
1952	3,348	4	76	34	11	35	104	279	672	1,048	856	227	2
1951	3,227	9	63	36	11	25	88	279	667	1,015	801	232	1
1950	3,181	15	64	47	11	19	99	277	679	1,001	813	153	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF URINARY ORGANS (180, 181)													
ALL OTHER, BOTH SEXES													
1959	879	2	26	16	4	14	58	150	247	239	101	22	-
1958	864	1	23	17	7	17	55	188	215	228	93	18	2
1957	850	4	19	13	6	12	55	162	248	217	92	22	-
1956	789	3	22	19	-	18	60	148	215	186	97	21	-
1955	667	-	17	10	4	14	45	118	195	168	74	22	-
1954	738	-	17	14	5	15	51	157	206	179	72	21	1
1953	715	1	19	10	1	8	55	162	189	190	67	13	-
1952	675	5	13	13	5	9	54	129	196	168	74	8	1
1951	638	3	18	9	4	17	53	146	163	158	51	16	-
1950	637	3	20	9	5	17	61	122	166	163	53	18	-
ALL OTHER, MALE													
1959	534	1	12	6	3	8	28	87	153	161	63	12	-
1958	525	-	11	7	3	10	30	111	136	143	62	10	2
1957	516	2	12	7	4	6	31	103	151	138	52	10	-
1956	471	3	10	8	-	8	27	83	137	116	66	13	-
1955	403	-	12	5	1	6	27	70	134	95	44	9	-
1954	445	-	11	7	3	8	29	93	137	107	37	13	-
1953	418	-	13	4	-	5	27	87	119	123	35	5	-
1952	412	4	7	3	4	8	30	73	134	101	46	2	-
1951	389	1	8	3	4	7	29	84	106	104	32	11	-
1950	369	3	10	5	2	6	29	73	95	95	40	11	-
ALL OTHER, FEMALE													
1959	345	1	14	10	1	6	30	63	94	78	38	10	-
1958	339	1	12	10	4	7	25	77	79	85	31	8	-
1957	334	2	7	6	2	6	24	59	97	79	40	12	-
1956	318	-	12	11	-	10	33	65	78	70	31	8	-
1955	264	-	5	5	3	8	18	48	61	73	30	13	-
1954	293	-	6	7	2	7	22	64	69	72	35	8	1
1953	297	1	6	6	1	3	28	75	70	67	32	8	-
1952	263	1	6	10	1	1	24	56	62	67	28	6	1
1951	249	2	10	6	-	10	24	62	57	54	19	5	-
1950	268	-	10	4	3	11	32	49	71	68	13	7	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)													
BOTH SEXES													
1959	28,663	92	475	846	644	967	2,158	4,292	6,244	6,970	4,509	1,459	7
1958	27,978	90	493	759	625	975	2,078	4,087	6,195	6,820	4,470	1,378	8
1957	28,425	80	497	701	601	1,050	2,017	4,205	6,439	6,786	4,633	1,406	10
1956	27,271	101	490	763	584	995	1,995	3,985	6,123	6,449	4,365	1,413	8
1955	26,958	89	465	755	594	939	2,054	4,033	6,121	6,475	4,123	1,296	14
1954	26,561	86	478	677	612	977	2,074	4,011	6,054	6,251	4,116	1,212	13
1953	25,870	92	469	636	580	959	2,065	3,934	5,906	6,043	3,937	1,239	10
1952	25,783	85	439	586	620	1,023	1,988	3,949	5,842	6,025	3,999	1,222	5
1951	24,692	90	468	528	619	963	1,998	3,783	5,619	5,641	3,829	1,136	18
1950	22,135	80	421	552	622	899	1,848	3,350	5,073	5,033	3,202	1,046	9
MALE													
1959	15,418	45	254	466	399	543	1,173	2,369	3,499	3,745	2,276	645	4
1958	14,833	56	250	417	369	543	1,096	2,250	3,415	3,665	2,173	596	3
1957	15,155	43	279	411	349	579	1,032	2,308	3,565	3,756	2,224	606	3
1956	14,484	54	251	411	338	538	1,075	2,145	3,504	3,361	2,165	637	5
1955	14,405	51	248	412	363	508	1,024	2,230	3,447	3,481	2,057	576	8
1954	14,184	46	263	390	352	527	1,029	2,118	3,399	3,420	2,072	563	5
1953	13,713	51	252	331	327	500	1,095	2,171	3,257	3,198	1,959	569	3
1952	13,586	52	238	323	358	527	982	2,045	3,252	3,242	2,025	540	2
1951	12,863	49	265	291	367	494	965	1,964	3,029	2,969	1,899	564	7
1950	11,663	44	228	298	350	468	935	1,776	2,786	2,685	1,642	449	2
FEMALE													
1959	13,245	47	221	380	245	424	985	1,923	2,745	3,225	2,233	814	3
1958	13,145	34	243	342	256	432	982	1,837	2,780	3,155	2,297	782	5
1957	13,270	37	218	290	252	471	985	1,897	2,874	3,030	2,409	800	7
1956	12,787	47	239	352	246	457	920	1,840	2,619	3,088	2,200	776	3
1955	12,553	38	217	343	231	431	1,030	1,803	2,674	2,994	2,066	720	6
1954	12,377	40	215	287	260	450	1,045	1,893	2,655	2,831	2,044	649	8
1953	12,157	41	217	305	253	459	970	1,763	2,649	2,845	1,978	670	7
1952	12,197	33	201	263	262	496	1,006	1,904	2,590	2,783	1,974	682	3
1951	11,829	41	203	237	252	469	1,033	1,819	2,590	2,672	1,930	572	11
1950	10,472	36	193	254	272	431	913	1,574	2,287	2,348	1,560	597	7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)													
WHITE, BOTH SEXES													
1959	26,222	79	434	764	561	882	1,942	3,824	5,623	6,439	4,277	1,393	4
1958	25,616	80	435	670	559	868	1,857	3,652	5,606	6,311	4,241	1,331	6
1957	26,057	76	441	627	540	956	1,796	3,726	5,828	6,295	4,407	1,359	6
1956	25,053	91	440	690	514	891	1,782	3,537	5,597	5,981	4,163	1,359	8
1955	24,724	80	421	688	533	850	1,823	3,556	5,568	6,018	3,938	1,236	13
1954	24,453	78	432	612	539	887	1,831	3,556	5,498	5,852	3,975	1,180	13
1953	23,890	84	433	578	513	884	1,824	3,518	5,435	5,639	3,782	1,193	7
1952	23,899	76	394	520	566	932	1,773	3,513	5,408	5,676	3,854	1,183	4
1951	22,830	82	416	483	564	883	1,777	3,339	5,177	5,301	3,699	1,094	15
1950	20,466	71	376	489	553	821	1,639	3,024	4,674	4,721	3,079	1,011	8
WHITE, MALE													
1959	14,137	36	235	419	347	500	1,066	2,141	3,172	3,456	2,151	612	2
1958	13,622	50	223	368	329	490	992	2,028	3,107	3,403	2,048	581	3
1957	13,951	40	240	367	318	531	925	2,066	3,263	3,495	2,117	588	1
1956	13,372	50	230	372	292	494	979	1,938	3,225	3,129	2,048	610	5
1955	13,257	46	229	377	329	468	928	1,999	3,143	3,234	1,957	540	7
1954	13,162	41	241	356	306	491	916	1,921	3,130	3,201	2,002	552	5
1953	12,683	45	231	301	288	465	991	1,950	2,988	2,976	1,893	552	3
1952	12,690	48	213	284	324	491	901	1,865	3,022	3,053	1,960	527	2
1951	11,959	44	236	273	338	455	884	1,744	2,811	2,791	1,832	546	5
1950	10,825	39	204	258	309	435	850	1,617	2,581	2,519	1,577	435	1
WHITE, FEMALE													
1959	12,085	43	199	345	214	382	876	1,683	2,451	2,983	2,126	781	2
1958	11,994	30	212	302	230	378	865	1,624	2,499	2,908	2,193	750	3
1957	12,106	36	201	260	222	425	871	1,660	2,565	2,800	2,290	771	5
1956	11,681	41	210	318	222	397	803	1,599	2,372	2,852	2,115	749	3
1955	11,467	34	192	311	204	382	895	1,557	2,425	2,784	1,981	696	6
1954	11,291	37	191	256	233	396	915	1,635	2,368	2,651	1,973	628	8
1953	11,207	39	202	277	225	419	833	1,568	2,447	2,663	1,889	641	4
1952	11,209	28	181	236	242	441	872	1,648	2,386	2,623	1,894	656	2
1951	10,871	38	180	210	226	428	893	1,595	2,366	2,510	1,867	548	10
1950	9,641	32	172	231	244	386	789	1,407	2,093	2,202	1,502	576	7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MALIGNANT NEOPLASM OF OTHER AND UNSPECIFIED SITES (156b, 165, 190-199)													
ALL OTHER, BOTH SEXES													
1959	2,441	13	41	82	83	85	216	468	621	531	232	66	3
1958	2,362	10	58	89	66	107	221	435	589	509	229	47	2
1957	2,368	4	56	74	61	94	221	479	611	491	226	47	4
1956	2,218	10	50	73	70	104	213	448	526	468	202	54	-
1955	2,234	9	44	67	61	89	231	477	553	457	185	60	1
1954	2,108	8	46	65	73	90	243	455	556	399	141	32	-
1953	1,980	8	36	58	67	75	241	416	471	404	155	46	3
1952	1,884	9	45	66	54	91	215	436	434	349	145	39	1
1951	1,862	8	52	45	55	80	221	444	442	340	130	42	3
1950	1,669	9	45	63	69	78	209	326	399	312	123	35	1
ALL OTHER, MALE													
1959	1,281	9	19	47	52	43	107	228	327	289	125	33	2
1958	1,211	6	27	49	40	53	104	222	308	262	125	15	-
1957	1,204	3	39	44	31	48	107	242	302	261	107	18	2
1956	1,112	4	21	39	46	44	96	207	279	232	117	27	-
1955	1,148	5	19	35	34	40	96	231	304	247	100	36	1
1954	1,022	5	22	34	46	36	113	197	269	219	70	11	-
1953	1,030	6	21	30	39	35	104	221	269	222	66	17	-
1952	896	4	25	39	34	36	81	180	230	189	65	13	-
1951	904	5	29	18	29	39	81	220	218	178	67	18	2
1950	838	5	24	40	41	33	85	159	205	166	65	14	1
ALL OTHER, FEMALE													
1959	1,160	4	22	35	31	42	109	240	294	242	107	33	1
1958	1,151	4	31	40	26	54	117	213	281	247	104	32	2
1957	1,164	1	17	30	30	46	114	237	309	230	119	29	2
1956	1,106	6	29	34	24	60	117	241	247	236	85	27	-
1955	1,086	4	25	32	27	49	135	246	249	210	85	24	-
1954	1,086	3	24	31	27	54	130	258	287	180	71	21	-
1953	950	2	15	28	28	40	137	195	202	182	89	29	3
1952	988	5	20	27	20	55	134	256	204	160	80	26	1
1951	958	3	23	27	26	41	140	224	224	162	63	24	1
1950	831	4	21	23	28	45	124	167	194	146	58	21	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>LEUKEMIA AND ALEUKEMIA (204)</b>													
<b>BOTH SEXES</b>													
1959	12,270	117	834	1,116	470	500	777	1,226	1,960	2,958	1,910	398	4
1958	11,980	94	846	1,065	515	471	792	1,186	1,948	2,812	1,854	394	3
1957	11,718	108	855	1,009	477	542	804	1,219	1,926	2,697	1,757	323	1
1956	11,396	84	838	983	440	572	755	1,157	2,034	2,555	1,671	304	3
1955	10,816	102	794	897	468	533	780	1,124	1,884	2,452	1,478	300	4
1954	10,443	115	807	875	423	538	694	1,175	1,759	2,393	1,419	244	1
1953	9,918	128	744	783	421	509	734	1,106	1,784	2,170	1,303	235	1
1952	9,841	125	810	775	468	514	726	1,145	1,785	2,122	1,176	188	7
1951	9,357	105	831	716	462	551	737	1,080	1,746	1,880	1,056	188	5
1950	8,845	127	766	691	414	562	729	1,012	1,618	1,808	973	140	5
<b>MALE</b>													
1959	7,123	67	453	672	286	267	437	692	1,174	1,794	1,069	210	2
1958	6,956	47	470	581	309	264	449	668	1,138	1,701	1,081	247	1
1957	6,745	57	473	550	281	309	417	693	1,165	1,640	995	165	-
1956	6,523	34	443	557	258	330	404	641	1,176	1,562	947	169	2
1955	6,303	49	436	560	291	305	430	647	1,098	1,480	835	169	3
1954	5,989	63	418	472	242	303	373	653	1,048	1,436	833	147	1
1953	5,740	57	400	467	273	287	404	629	1,049	1,311	730	132	1
1952	5,612	54	447	439	282	272	392	616	1,044	1,293	675	97	1
1951	5,458	58	466	388	289	302	377	629	1,049	1,177	614	106	3
1950	5,058	75	420	389	233	302	375	565	946	1,124	548	79	2
<b>FEMALE</b>													
1959	5,147	50	381	444	184	233	340	534	786	1,164	841	188	2
1958	5,024	47	376	484	206	207	343	518	810	1,111	773	147	2
1957	4,973	51	382	459	196	233	387	526	761	1,057	762	158	1
1956	4,873	50	395	426	182	242	351	516	858	993	724	135	1
1955	4,513	53	358	337	177	228	350	477	786	972	643	131	1
1954	4,454	52	389	403	181	235	321	522	711	957	586	97	-
1953	4,178	71	344	316	148	222	330	477	735	859	573	103	-
1952	4,229	71	363	336	186	242	334	529	741	829	501	91	6
1951	3,899	47	365	328	173	249	360	451	697	703	442	82	2
1950	3,787	52	346	302	181	260	354	447	672	684	425	61	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>LEUKEMIA AND ALEUKEMIA (204)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	11,468	98	768	1,045	417	442	692	1,115	1,806	2,834	1,858	389	4
1958	11,193	84	791	982	459	396	710	1,075	1,811	2,684	1,818	381	2
1957	10,982	100	805	936	429	493	727	1,106	1,782	2,577	1,709	317	1
1956	10,646	69	780	908	398	510	676	1,041	1,892	2,445	1,629	295	3
1955	10,189	91	751	836	423	495	706	1,014	1,771	2,369	1,438	292	3
1954	9,794	98	757	820	373	483	622	1,069	1,649	2,295	1,391	236	1
1953	9,325	111	693	734	375	460	656	998	1,686	2,102	1,282	227	1
1952	9,252	104	757	718	434	463	651	1,048	1,691	2,044	1,153	182	7
1951	8,819	97	790	672	416	492	669	993	1,649	1,817	1,036	183	5
1950	8,357	110	729	642	379	522	654	935	1,526	1,756	960	139	5
<b>WHITE, MALE</b>													
1959	6,683	59	417	634	262	237	405	633	1,093	1,705	1,031	205	2
1958	6,524	43	442	542	277	217	412	613	1,059	1,624	1,054	240	1
1957	6,296	53	442	507	251	280	374	629	1,075	1,558	966	161	-
1956	6,093	29	406	515	235	293	370	580	1,091	1,492	918	162	2
1955	5,930	43	413	520	258	284	391	591	1,029	1,423	811	165	2
1954	5,589	51	390	442	209	266	337	593	978	1,367	815	140	1
1953	5,404	49	374	435	243	262	362	579	993	1,264	715	127	1
1952	5,291	43	415	406	266	248	362	574	981	1,241	660	94	1
1951	5,141	54	448	365	260	271	336	582	985	1,135	601	101	3
1950	4,783	64	398	364	217	280	337	524	889	1,089	540	79	2
<b>WHITE, FEMALE</b>													
1959	4,785	39	351	411	155	205	287	482	713	1,129	827	184	2
1958	4,669	41	349	440	182	179	298	462	752	1,060	764	141	1
1957	4,686	47	363	429	178	213	353	477	707	1,019	743	156	1
1956	4,553	40	374	393	163	217	306	461	801	953	711	133	1
1955	4,259	48	338	316	165	211	315	423	742	946	627	127	1
1954	4,205	47	367	378	164	217	285	476	671	928	576	96	-
1953	3,921	62	319	299	132	198	294	419	693	838	567	100	-
1952	3,961	61	342	312	168	215	289	474	710	803	493	88	6
1951	3,678	43	342	307	156	221	333	411	664	682	435	82	2
1950	3,574	46	331	278	162	242	317	411	637	667	420	60	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
LEUKEMIA AND ALEUKEMIA (204)													
ALL OTHER, BOTH SEXES													
1959	802	19	66	71	53	58	85	111	154	124	52	9	-
1958	787	10	55	83	56	75	82	111	137	128	36	13	1
1957	736	8	50	73	48	49	77	113	144	120	48	6	-
1956	750	15	58	75	42	62	79	116	142	110	42	9	-
1955	627	11	43	61	45	38	74	110	113	83	40	8	1
1954	649	17	50	55	50	55	72	106	110	98	28	8	-
1953	593	17	51	49	46	49	78	108	98	68	21	8	-
1952	589	21	53	57	34	51	75	97	94	78	23	6	-
1951	538	8	41	44	46	59	68	87	97	63	20	5	-
1950	488	17	37	49	35	40	75	77	92	52	13	1	-
ALL OTHER, MALE													
1959	440	8	36	38	24	30	32	59	81	89	38	5	-
1958	432	4	28	39	32	47	37	55	79	77	27	7	-
1957	449	4	31	43	30	29	43	64	90	82	29	4	-
1956	430	5	37	42	23	37	34	61	85	70	29	7	-
1955	373	6	23	40	33	21	39	56	69	57	24	4	1
1954	400	12	28	30	33	37	36	60	70	69	18	7	-
1953	336	8	26	32	30	25	42	50	56	47	15	5	-
1952	321	11	32	33	16	24	30	42	63	52	15	3	-
1951	317	4	18	23	29	31	41	47	64	42	13	5	-
1950	275	11	22	25	16	22	38	41	57	35	8	-	-
ALL OTHER, FEMALE													
1959	362	11	30	33	29	28	53	52	73	35	14	4	-
1958	355	6	27	44	24	28	45	56	58	51	9	6	1
1957	287	4	19	30	18	20	34	49	54	38	19	2	-
1956	320	10	21	33	19	25	45	55	57	40	13	2	-
1955	254	5	20	21	12	17	35	54	44	26	16	4	-
1954	249	5	22	25	17	18	36	46	40	29	10	1	-
1953	257	9	25	17	16	24	36	58	42	21	6	3	-
1952	268	10	21	24	18	27	45	55	31	26	8	3	-
1951	221	4	23	21	17	28	27	40	33	21	7	-	-
1950	213	6	15	24	19	18	37	36	35	17	5	1	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)													
BOTH SEXES													
1959	12,471	22	121	253	437	723	1,076	1,935	2,813	3,282	1,589	218	2
1958	11,948	26	117	233	409	716	1,087	1,837	2,879	3,051	1,393	196	4
1957	11,849	34	92	294	397	664	1,027	1,781	2,867	3,086	1,413	193	1
1956	11,444	46	100	279	393	712	1,057	1,697	2,798	2,852	1,322	181	7
1955	10,643	26	142	250	408	676	1,002	1,636	2,530	2,669	1,152	149	3
1954	10,114	32	98	224	369	713	990	1,601	2,447	2,432	1,054	152	2
1953	9,658	21	95	236	409	732	945	1,528	2,457	2,176	948	110	1
1952	9,106	23	111	200	377	730	890	1,505	2,272	2,055	833	108	2
1951	8,438	25	99	188	371	674	880	1,406	2,119	1,790	774	108	4
1950	7,911	17	95	193	359	672	814	1,336	1,990	1,687	670	76	2
MALE													
1959	7,277	13	72	188	293	448	658	1,164	1,640	1,844	837	118	2
1958	6,945	15	73	159	254	430	677	1,124	1,724	1,662	730	95	2
1957	7,084	18	68	205	266	421	651	1,097	1,729	1,788	751	90	-
1956	6,748	27	62	180	250	431	629	1,032	1,721	1,613	700	98	5
1955	6,345	12	96	186	254	410	634	1,018	1,525	1,531	604	74	1
1954	5,984	19	61	150	247	436	603	995	1,483	1,359	560	70	1
1953	5,828	8	60	177	238	460	606	954	1,490	1,255	524	55	1
1952	5,428	8	60	129	229	461	539	928	1,376	1,190	456	52	-
1951	5,100	8	58	129	249	404	550	865	1,274	1,052	452	57	2
1950	4,781	11	57	140	220	416	518	843	1,220	947	376	32	1
FEMALE													
1959	5,194	9	49	65	144	275	418	771	1,173	1,438	752	100	-
1958	5,003	11	44	74	155	286	410	713	1,155	1,389	663	101	2
1957	4,765	16	24	89	131	243	376	684	1,138	1,298	662	103	1
1956	4,696	19	38	99	143	281	428	665	1,077	1,239	622	83	2
1955	4,298	14	46	64	154	266	368	618	1,005	1,138	548	75	2
1954	4,130	13	37	74	122	277	387	606	964	1,073	494	82	1
1953	3,830	13	35	59	171	272	339	574	967	921	424	55	-
1952	3,678	15	51	71	148	269	351	577	896	865	377	56	2
1951	3,338	17	41	59	122	270	330	541	845	738	322	51	2
1950	3,130	6	38	53	139	256	296	493	770	740	294	44	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)													
WHITE, BOTH SEXES													
1959	11,520	20	114	230	392	647	975	1,745	2,578	3,078	1,535	205	1
1958	11,039	25	104	206	378	657	973	1,653	2,629	2,881	1,340	190	3
1957	10,990	30	85	267	365	603	920	1,631	2,613	2,925	1,365	185	1
1956	10,588	41	90	246	360	665	949	1,519	2,581	2,684	1,275	173	5
1955	9,884	23	128	222	369	616	885	1,480	2,354	2,536	1,123	146	2
1954	9,405	30	91	206	326	659	885	1,429	2,285	2,323	1,022	147	2
1953	9,019	17	92	216	371	661	857	1,377	2,304	2,094	922	108	-
1952	8,494	21	100	173	336	681	793	1,370	2,136	1,969	811	102	2
1951	7,916	25	84	175	336	613	805	1,302	1,985	1,725	756	107	3
1950	7,376	14	86	173	329	606	734	1,218	1,876	1,612	651	75	2
WHITE, MALE													
1959	6,679	12	67	170	267	400	595	1,048	1,495	1,714	802	108	1
1958	6,391	14	65	141	240	399	599	1,010	1,575	1,559	696	92	1
1957	6,555	15	64	187	242	388	573	1,013	1,566	1,693	727	87	-
1956	6,218	25	55	162	232	395	567	933	1,583	1,499	671	93	3
1955	5,873	11	88	167	225	369	568	927	1,415	1,444	587	72	-
1954	5,537	18	56	139	210	402	547	898	1,382	1,281	537	66	1
1953	5,417	6	59	162	215	415	548	848	1,402	1,203	506	53	-
1952	5,047	8	56	113	203	431	482	843	1,284	1,133	447	47	-
1951	4,784	8	52	120	225	366	508	806	1,188	1,010	443	57	1
1950	4,459	9	53	126	205	378	472	769	1,154	898	362	32	1
WHITE, FEMALE													
1959	4,841	8	47	60	125	247	380	697	1,083	1,364	733	97	-
1958	4,648	11	39	65	138	258	374	643	1,054	1,322	644	98	2
1957	4,435	15	21	80	123	215	347	618	1,047	1,232	638	98	1
1956	4,370	16	35	84	128	270	382	586	998	1,185	604	80	2
1955	4,011	12	40	55	144	247	317	553	939	1,092	536	74	2
1954	3,868	12	35	67	116	257	338	531	903	1,042	485	81	1
1953	3,602	11	33	54	156	246	309	529	902	891	416	55	-
1952	3,447	13	44	60	133	250	311	527	852	836	364	55	2
1951	3,132	17	32	55	111	247	297	496	797	715	313	50	2
1950	2,917	5	33	47	124	228	262	449	722	714	289	43	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
LYMPHOSARCOMA AND OTHER NEOPLASMS OF LYMPHATIC AND HEMATOPOIETIC TISSUES (200-203, 205)													
ALL OTHER, BOTH SEXES													
1959	951	2	7	23	45	76	101	190	235	204	54	13	1
1958	909	1	13	27	31	59	114	184	250	170	53	6	1
1957	859	4	7	27	32	61	107	150	254	161	48	8	-
1956	856	5	10	33	33	47	108	178	217	168	47	8	2
1955	759	3	14	28	39	60	117	156	176	133	29	3	1
1954	709	2	7	18	43	54	105	172	162	109	32	5	-
1953	639	4	3	20	38	71	88	151	153	82	26	2	1
1952	612	2	11	27	41	49	97	135	136	86	22	6	-
1951	522	-	15	13	35	61	75	104	134	65	18	1	1
1950	535	3	9	20	30	66	80	118	114	75	19	1	-
ALL OTHER, MALE													
1959	598	1	5	18	26	48	63	116	145	130	35	10	1
1958	554	1	8	18	14	31	78	114	149	103	34	3	1
1957	529	3	4	18	24	33	78	84	163	95	24	3	-
1956	530	2	7	18	18	36	62	99	138	114	29	5	2
1955	472	1	8	19	29	41	66	91	110	87	17	2	1
1954	447	1	5	11	37	34	56	97	101	78	23	4	-
1953	411	2	1	15	23	45	58	106	88	52	18	2	1
1952	381	-	4	16	26	30	57	85	92	57	9	5	-
1951	316	-	6	9	24	38	42	59	86	42	9	-	1
1950	322	2	4	14	15	38	46	74	66	49	14	-	-
ALL OTHER, FEMALE													
1959	353	1	2	5	19	28	38	74	90	74	19	3	-
1958	355	-	5	9	17	28	36	70	101	67	19	3	-
1957	330	1	3	9	8	28	29	66	91	66	24	5	-
1956	326	3	3	15	15	11	46	79	79	54	18	3	-
1955	287	2	6	9	10	19	51	65	66	46	12	1	-
1954	262	1	2	7	6	20	49	75	61	31	9	1	-
1953	228	2	2	5	15	26	30	45	65	30	8	-	-
1952	231	2	7	11	15	19	40	50	44	29	13	1	-
1951	206	-	9	4	11	23	33	45	48	23	9	1	-
1950	213	1	5	6	15	28	34	44	48	26	5	1	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)</b>													
<b>BOTH SEXES</b>													
1959	4,949	127	124	240	177	284	562	924	921	912	494	183	1
1958	4,961	135	132	207	175	322	632	889	920	895	484	168	2
1957	5,068	134	117	210	178	334	651	926	953	865	521	179	-
1956	4,887	157	132	190	190	325	633	890	951	788	477	151	3
1955	5,168	163	147	210	203	356	701	946	930	852	503	152	5
1954	5,094	168	146	207	175	388	708	984	944	755	487	131	1
1953	5,263	158	147	236	174	411	789	1,016	964	737	491	138	2
1952	5,133	149	151	189	201	401	870	973	964	687	406	138	4
1951	5,166	147	138	170	199	431	823	1,036	950	717	408	144	3
1950	5,374	137	147	185	198	413	876	1,019	1,054	769	441	126	9
<b>MALE</b>													
1959	2,196	63	75	138	87	113	191	351	453	458	211	56	-
1958	2,251	51	71	95	92	123	232	402	475	435	217	57	1
1957	2,296	77	60	112	93	128	234	407	466	437	220	62	-
1956	2,139	90	65	104	80	121	188	372	469	396	203	50	1
1955	2,228	86	66	98	92	127	212	385	464	413	229	54	2
1954	2,181	89	79	101	95	147	218	369	469	372	203	39	-
1953	2,247	90	76	116	82	155	217	362	494	381	217	57	-
1952	2,084	82	77	96	93	156	254	366	438	305	163	52	2
1951	2,090	69	74	85	97	139	241	362	455	351	180	36	1
1950	2,122	74	79	103	93	132	233	349	479	366	174	37	3
<b>FEMALE</b>													
1959	2,753	64	49	102	90	171	371	573	468	454	283	127	1
1958	2,710	84	61	112	83	199	400	487	445	460	267	111	1
1957	2,772	57	57	98	85	206	417	519	487	428	301	117	-
1956	2,748	67	67	86	110	204	445	518	482	392	274	101	2
1955	2,940	77	81	112	111	229	489	561	466	439	274	98	3
1954	2,913	79	67	106	80	241	490	615	475	383	284	92	1
1953	3,016	68	71	120	92	256	572	654	470	356	274	81	2
1952	3,049	67	74	93	108	245	616	607	526	382	243	86	2
1951	3,076	78	64	85	102	292	582	674	495	366	228	108	2
1950	3,252	63	68	82	105	281	643	670	575	403	267	89	6

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	4,241	108	106	202	147	228	419	761	809	832	460	169	-
1958	4,259	117	110	176	143	255	478	753	801	817	453	155	1
1957	4,329	115	101	172	141	262	495	762	820	798	494	169	-
1956	4,179	133	110	166	155	248	479	739	845	717	445	141	1
1955	4,360	127	125	177	171	267	506	780	827	768	465	144	3
1954	4,320	140	119	172	149	303	538	798	837	692	450	122	-
1953	4,492	133	112	205	147	324	585	839	870	680	465	132	-
1952	4,279	127	129	163	158	303	625	777	854	626	383	131	3
1951	4,392	133	122	139	159	328	611	866	843	670	382	138	1
1950	4,574	122	128	166	168	313	644	845	941	694	424	121	8
<b>WHITE, MALE</b>													
1959	1,944	56	63	120	72	96	164	301	404	420	199	49	-
1958	1,997	45	58	81	84	104	193	353	422	401	200	55	1
1957	2,036	65	53	95	71	110	201	360	404	408	210	59	-
1956	1,921	77	53	93	68	111	161	337	421	361	193	46	-
1955	1,990	72	56	84	80	109	186	341	425	372	211	54	-
1954	1,925	76	65	85	82	127	187	312	422	347	187	35	-
1953	2,007	76	59	101	69	137	181	309	462	351	205	57	-
1952	1,854	70	66	81	73	138	219	312	401	289	155	49	1
1951	1,873	63	63	71	75	117	207	320	417	334	171	34	1
1950	1,923	66	66	93	82	115	211	303	440	339	171	34	3
<b>WHITE, FEMALE</b>													
1959	2,297	52	43	82	75	132	255	460	405	412	261	120	-
1958	2,262	72	52	95	59	151	285	400	379	416	253	100	-
1957	2,293	50	48	77	70	152	294	402	416	390	284	110	-
1956	2,258	56	57	73	87	137	318	402	424	356	252	95	1
1955	2,370	55	69	93	91	158	320	439	402	396	254	90	3
1954	2,395	64	54	87	67	176	351	486	415	345	263	87	-
1953	2,485	57	53	104	78	187	404	530	408	329	260	75	-
1952	2,425	57	63	82	85	165	406	465	453	337	228	82	2
1951	2,519	70	59	68	84	211	404	546	426	336	211	104	-
1950	2,651	56	62	73	86	198	433	542	501	355	253	87	5

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>BENIGN NEOPLASMS AND NEOPLASMS OF UNSPECIFIED NATURE (210-239)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	708	19	18	38	30	56	143	163	112	80	34	14	1
1958	702	18	22	31	32	67	154	136	119	78	31	13	1
1957	739	19	16	38	37	72	156	164	133	67	27	10	-
1956	708	24	22	24	35	77	154	151	106	71	32	10	2
1955	808	36	22	33	32	89	195	166	103	84	38	8	2
1954	774	28	27	35	26	85	170	186	107	63	37	9	1
1953	771	25	35	31	27	87	204	177	94	57	26	6	2
1952	854	22	22	26	43	98	245	196	110	61	23	7	1
1951	774	14	16	31	40	103	212	170	107	47	26	6	2
1950	800	15	19	19	30	100	232	174	113	75	17	5	1
<b>ALL OTHER, MALE</b>													
1959	252	7	12	18	15	17	27	50	49	38	12	7	-
1958	254	6	13	14	8	19	39	49	53	34	17	2	-
1957	260	12	7	17	22	18	33	47	62	29	10	3	-
1956	218	13	12	11	12	10	27	35	48	35	10	4	1
1955	238	14	10	14	12	18	26	44	39	41	18	-	2
1954	256	13	14	16	13	20	31	57	47	25	16	4	-
1953	240	14	17	15	13	18	36	53	32	30	12	-	-
1952	230	12	11	15	20	18	35	54	37	16	8	3	1
1951	217	6	11	14	22	22	34	42	38	17	9	2	-
1950	199	8	13	10	11	17	22	46	39	27	3	3	-
<b>ALL OTHER, FEMALE</b>													
1959	456	12	6	20	15	39	116	113	63	42	22	7	1
1958	448	12	9	17	24	48	115	87	66	44	14	11	1
1957	479	7	9	21	15	54	123	117	71	38	17	7	-
1956	490	11	10	13	23	67	127	116	58	36	22	6	1
1955	570	22	12	19	20	71	169	122	64	43	20	8	-
1954	518	15	13	19	13	65	139	129	60	38	21	5	1
1953	531	11	18	16	14	69	168	124	62	27	14	6	2
1952	624	10	11	11	23	80	210	142	73	45	15	4	-
1951	557	8	5	17	18	81	178	128	69	30	17	4	2
1950	601	7	6	9	19	83	210	128	74	48	14	2	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ASTHMA (241)													
BOTH SEXES													
1959	4,984	31	63	65	69	172	372	717	1,174	1,396	769	155	1
1958	5,035	31	59	56	77	158	326	703	1,203	1,426	812	183	1
1957	6,677	52	87	62	83	180	392	821	1,577	2,000	1,189	231	3
1956	5,965	60	85	38	84	144	282	695	1,396	1,908	1,035	234	4
1955	5,960	49	87	48	86	168	332	688	1,433	1,815	1,008	243	3
1954	6,136	63	91	33	86	175	354	751	1,477	1,836	1,044	224	2
1953	6,737	80	80	53	70	168	384	803	1,693	2,060	1,106	235	5
1952	6,943	91	93	43	88	193	395	882	1,773	2,078	1,072	233	2
1951	6,871	64	104	32	77	182	364	908	1,733	2,001	1,127	269	10
1950	4,322	73	104	39	76	173	301	679	1,083	1,054	593	145	2
MALE													
1959	3,266	22	26	39	20	70	159	439	865	1,025	518	82	1
1958	3,384	20	31	31	23	66	144	408	891	1,098	574	97	1
1957	4,536	31	50	36	32	66	187	526	1,188	1,506	795	117	2
1956	4,038	39	48	18	35	50	124	445	1,048	1,462	663	102	4
1955	3,985	24	43	25	32	57	156	437	1,051	1,367	655	135	3
1954	4,066	38	51	18	34	73	168	471	1,096	1,349	647	120	1
1953	4,465	45	38	34	24	56	204	489	1,258	1,476	709	127	5
1952	4,601	56	57	19	29	62	182	580	1,303	1,516	690	107	-
1951	4,539	37	54	21	37	65	164	582	1,318	1,425	700	130	6
1950	2,695	41	60	23	24	51	144	422	751	731	373	74	1
FEMALE													
1959	1,718	9	37	26	49	102	213	278	309	371	251	73	-
1958	1,651	11	28	25	54	92	182	295	312	328	238	86	-
1957	2,141	21	37	26	51	114	205	295	389	494	394	114	1
1956	1,927	21	37	20	49	94	158	250	348	446	372	132	-
1955	1,975	25	44	23	54	111	176	251	382	448	353	108	-
1954	2,070	25	40	15	52	102	186	280	381	487	397	104	1
1953	2,272	35	42	19	46	112	180	314	435	584	397	108	-
1952	2,342	35	36	24	59	131	213	302	470	562	382	126	2
1951	2,332	27	50	11	40	117	200	326	415	576	427	139	4
1950	1,627	32	44	16	52	122	157	257	332	323	220	71	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ASTHMA (241)													
WHITE, BOTH SEXES													
1959	4,323	23	40	37	37	101	248	595	1,038	1,322	735	146	1
1958	4,405	19	44	36	52	101	211	566	1,091	1,321	786	178	-
1957	5,881	37	60	42	52	110	280	662	1,400	1,870	1,143	222	3
1956	5,355	44	62	26	62	99	210	565	1,266	1,797	994	229	1
1955	5,402	40	70	34	56	110	248	596	1,316	1,724	977	229	2
1954	5,577	41	70	24	64	130	278	647	1,357	1,749	999	217	1
1953	6,151	57	58	41	49	118	313	686	1,549	1,973	1,073	230	4
1952	6,300	77	72	36	66	137	309	751	1,641	1,963	1,031	215	2
1951	6,305	54	77	22	57	134	294	782	1,614	1,910	1,095	257	9
1950	3,900	53	81	28	59	129	228	608	1,005	1,003	568	136	2
WHITE, MALE													
1959	2,906	15	16	20	12	37	105	364	784	971	502	79	1
1958	3,046	12	23	20	14	44	99	343	816	1,022	559	94	-
1957	4,065	21	30	26	22	45	133	430	1,072	1,413	757	114	2
1956	3,670	29	36	14	24	29	91	365	954	1,389	638	100	1
1955	3,658	20	34	18	16	36	122	385	971	1,297	632	125	2
1954	3,738	24	41	13	27	53	128	410	1,016	1,289	618	118	1
1953	4,134	32	29	25	13	38	169	424	1,168	1,418	690	124	4
1952	4,243	45	44	16	19	38	147	500	1,222	1,440	668	104	-
1951	4,235	32	41	17	30	48	131	511	1,246	1,367	678	128	6
1950	2,462	30	47	14	19	37	108	379	707	694	356	70	1
WHITE, FEMALE													
1959	1,417	8	24	17	25	64	143	231	254	351	233	67	-
1958	1,359	7	21	16	38	57	112	223	275	299	227	84	-
1957	1,816	16	30	16	30	65	147	232	328	457	386	108	1
1956	1,685	15	26	12	38	70	119	200	312	408	356	129	-
1955	1,744	20	36	16	40	74	126	211	345	427	345	104	-
1954	1,839	17	29	11	37	77	150	237	341	460	381	99	-
1953	2,017	25	29	16	36	80	144	262	381	555	383	106	-
1952	2,057	32	28	20	47	99	162	251	419	523	363	111	2
1951	2,070	22	36	5	27	86	163	271	368	543	417	129	3
1950	1,438	23	34	14	40	92	120	229	298	309	212	66	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ASTHMA (241)													
ALL OTHER, BOTH SEXES													
1959	661	8	23	28	32	71	124	122	136	74	34	9	-
1958	630	12	15	20	25	57	115	137	112	105	26	5	1
1957	796	15	27	20	31	70	112	159	177	130	46	9	-
1956	610	16	23	12	22	45	72	130	130	111	41	5	3
1955	558	9	17	14	30	58	84	92	117	91	31	14	1
1954	559	22	21	9	22	45	76	104	120	87	45	7	1
1953	586	23	22	12	21	50	71	117	144	87	33	5	1
1952	643	14	21	7	22	56	86	131	132	115	41	18	-
1951	566	10	27	10	20	48	70	126	119	91	32	12	1
1950	422	20	23	11	17	44	73	71	78	51	25	9	-
ALL OTHER, MALE													
1959	360	7	10	19	8	33	54	75	81	54	16	3	-
1958	338	8	8	11	9	22	45	65	75	76	15	3	1
1957	471	10	20	10	10	21	54	96	116	93	38	3	-
1956	368	10	12	4	11	21	33	80	94	73	25	2	3
1955	327	4	9	7	16	21	34	52	80	70	23	10	1
1954	328	14	10	5	7	20	40	61	80	60	29	2	-
1953	331	13	9	9	11	18	35	65	90	58	19	3	1
1952	358	11	13	3	10	24	35	80	81	76	22	3	-
1951	304	5	13	4	7	17	33	71	72	58	22	2	-
1950	233	11	13	9	5	14	36	43	44	37	17	4	-
ALL OTHER, FEMALE													
1959	301	1	13	9	24	38	70	47	55	20	18	6	-
1958	292	4	7	9	16	35	70	72	37	29	11	2	-
1957	325	5	7	10	21	49	58	63	61	37	8	6	-
1956	242	6	11	8	11	24	39	50	36	38	16	3	-
1955	231	5	8	7	14	37	50	40	37	21	8	4	-
1954	231	8	11	4	15	25	36	43	40	27	16	5	1
1953	255	10	13	3	10	32	36	52	54	29	14	2	-
1952	285	3	8	4	12	32	51	51	51	39	19	15	-
1951	262	5	14	6	13	31	37	55	47	33	10	10	1
1950	189	9	10	2	12	30	37	28	34	14	8	5	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
DIABETES (260)													
BOTH SEXES													
1959	28,080	14	40	127	203	586	917	2,308	5,759	9,708	6,985	1,426	7
1958	27,501	14	54	109	203	575	977	2,230	5,598	9,666	6,610	1,461	4
1957	27,180	12	42	146	253	620	1,064	2,182	5,684	9,526	6,324	1,314	13
1956	26,184	19	29	93	215	561	884	2,110	5,549	9,241	6,177	1,296	10
1955	25,488	23	42	115	216	568	877	2,047	5,441	8,971	5,927	1,251	10
1954	25,151	18	44	124	213	526	885	2,090	5,490	8,780	5,840	1,132	9
1953	25,796	17	42	124	249	565	907	2,119	5,888	8,875	5,919	1,081	10
1952	25,474	26	63	150	270	540	875	2,190	5,678	8,947	5,720	999	16
1951	25,047	13	60	125	265	526	856	2,191	5,835	8,704	5,526	931	15
1950	24,419	23	38	136	248	511	898	2,146	5,635	8,439	5,465	867	13
MALE													
1959	11,232	5	17	49	89	334	494	1,131	2,305	3,612	2,648	545	3
1958	10,961	6	28	46	97	331	549	1,036	2,213	3,616	2,476	562	1
1957	10,605	3	22	58	109	344	570	997	2,135	3,483	2,389	489	6
1956	10,329	8	16	38	86	315	477	941	2,213	3,390	2,350	492	3
1955	10,082	15	24	56	95	299	492	918	2,130	3,284	2,307	456	6
1954	9,835	10	24	59	97	293	496	947	2,073	3,170	2,229	435	2
1953	9,889	8	24	49	114	323	489	895	2,133	3,166	2,245	438	5
1952	9,957	15	37	56	114	267	462	952	2,122	3,255	2,261	408	8
1951	9,511	8	28	41	106	260	429	946	2,186	3,126	2,022	354	5
1950	9,348	16	15	57	97	246	457	887	2,026	3,069	2,138	333	7
FEMALE													
1959	16,848	9	23	78	114	252	423	1,177	3,454	6,096	4,337	881	4
1958	16,540	8	26	63	106	244	428	1,194	3,385	6,050	4,134	899	3
1957	16,575	9	20	88	144	276	494	1,185	3,549	6,043	3,935	825	7
1956	15,855	11	13	55	129	246	407	1,169	3,336	5,851	3,827	804	7
1955	15,406	8	18	59	121	269	385	1,129	3,311	5,687	3,620	795	4
1954	15,316	8	20	65	116	233	389	1,143	3,417	5,610	3,611	697	7
1953	15,907	9	18	75	135	242	418	1,224	3,755	5,709	3,674	643	5
1952	15,517	11	26	94	156	273	413	1,238	3,556	5,692	3,459	591	8
1951	15,536	5	32	84	159	266	427	1,245	3,649	5,578	3,504	577	10
1950	15,071	7	23	79	151	265	441	1,259	3,609	5,370	3,327	534	6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
DIABETES (260)													
WHITE, BOTH SEXES													
1959	24,770	12	33	110	171	489	705	1,724	4,753	8,830	6,590	1,349	4
1958	24,378	13	46	84	158	475	761	1,683	4,719	8,802	6,256	1,378	3
1957	24,234	8	36	122	199	517	810	1,631	4,842	8,769	6,036	1,255	9
1956	23,500	18	27	82	166	468	697	1,622	4,718	8,538	5,915	1,244	5
1955	22,956	14	38	97	178	487	666	1,565	4,729	8,304	5,668	1,203	7
1954	22,714	13	36	102	173	450	688	1,575	4,791	8,175	5,629	1,077	5
1953	23,314	15	40	107	207	469	688	1,589	5,204	8,291	5,674	1,024	6
1952	23,151	25	59	122	218	458	680	1,711	5,049	8,364	5,497	954	14
1951	22,766	13	54	104	208	446	634	1,696	5,241	8,148	5,323	887	12
1950	22,147	21	31	115	192	421	644	1,646	5,059	7,922	5,265	824	7
WHITE, MALE													
1959	10,073	4	14	44	79	284	427	921	1,986	3,289	2,497	527	1
1958	9,920	5	25	37	77	285	458	867	1,969	3,322	2,352	522	1
1957	9,623	2	20	49	83	290	483	840	1,897	3,202	2,291	462	4
1956	9,438	7	16	35	68	274	407	808	1,948	3,145	2,251	477	2
1955	9,242	10	21	47	78	258	409	794	1,912	3,070	2,197	442	4
1954	9,009	8	18	45	78	258	424	801	1,859	2,959	2,143	415	1
1953	9,089	7	22	41	98	275	413	759	1,930	2,984	2,141	416	3
1952	9,167	15	34	49	91	227	403	818	1,928	3,040	2,164	391	7
1951	8,711	8	25	31	79	229	359	799	1,981	2,914	1,945	336	5
1950	8,580	14	14	52	76	211	369	739	1,854	2,876	2,049	322	4
WHITE, FEMALE													
1959	14,697	8	19	66	92	205	278	803	2,767	5,541	4,093	822	3
1958	14,458	8	21	47	81	190	303	816	2,750	5,480	3,904	856	2
1957	14,611	6	16	73	116	227	327	791	2,945	5,567	3,745	793	5
1956	14,062	11	11	47	98	194	290	814	2,770	5,393	3,664	767	3
1955	13,714	4	17	50	100	229	257	771	2,817	5,234	3,471	761	3
1954	13,705	5	18	57	95	192	264	774	2,932	5,216	3,486	662	4
1953	14,225	8	18	66	109	194	275	830	3,274	5,307	3,533	608	3
1952	13,984	10	25	73	127	231	277	893	3,121	5,324	3,333	563	7
1951	14,055	5	29	73	129	217	275	897	3,260	5,234	3,378	551	7
1950	13,567	7	17	63	116	210	275	907	3,205	5,046	3,216	502	3

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
DIABETES (260)													
ALL OTHER, BOTH SEXES													
1959	3,310	2	7	17	32	97	212	584	1,006	878	395	77	3
1958	3,123	1	8	25	45	100	216	547	879	864	354	83	1
1957	2,946	4	6	24	54	103	254	551	842	757	288	59	4
1956	2,684	1	2	11	49	93	187	488	831	703	262	52	5
1955	2,532	9	4	18	38	81	211	482	712	667	259	48	3
1954	2,437	5	8	22	40	76	197	515	699	605	211	55	4
1953	2,482	2	2	17	42	96	219	530	684	584	245	57	4
1952	2,323	1	4	28	52	82	195	479	629	583	223	45	2
1951	2,281	-	6	21	57	80	222	495	594	556	203	44	3
1950	2,272	2	7	21	56	90	254	500	576	517	200	43	6
ALL OTHER, MALE													
1959	1,159	1	3	5	10	50	67	210	319	323	151	18	2
1958	1,041	1	3	9	20	46	91	169	244	294	124	40	-
1957	982	1	2	9	26	54	87	157	238	281	98	27	2
1956	891	1	-	3	18	41	70	133	265	245	99	15	1
1955	840	5	3	9	17	41	83	124	218	214	110	14	2
1954	826	2	6	14	19	35	72	146	214	211	86	20	1
1953	800	1	2	8	16	48	76	136	203	182	104	22	2
1952	790	-	3	7	23	40	59	134	194	215	97	17	1
1951	800	-	3	10	27	31	70	147	205	212	77	18	-
1950	768	2	1	5	21	35	88	148	172	193	89	11	3
ALL OTHER, FEMALE													
1959	2,151	1	4	12	22	47	145	374	687	555	244	59	1
1958	2,082	-	5	16	25	54	125	378	635	570	230	43	1
1957	1,964	3	4	15	28	49	167	394	604	476	190	32	2
1956	1,793	-	2	8	31	52	117	355	566	458	163	37	4
1955	1,692	4	1	9	21	40	128	358	494	453	149	34	1
1954	1,611	3	2	8	21	41	125	369	485	394	125	35	3
1953	1,682	1	-	9	26	48	143	394	481	402	141	35	2
1952	1,533	1	1	21	29	42	136	345	435	368	126	28	1
1951	1,481	-	3	11	30	49	152	348	389	344	126	26	3
1950	1,504	-	6	16	35	55	166	352	404	324	111	32	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)</b>													
<b>BOTH SEXES</b>													
1959	2,208	837	397	149	71	73	121	119	179	175	74	12	1
1958	2,247	877	444	137	66	75	77	150	180	154	66	21	-
1957	2,025	769	368	156	68	53	92	140	163	137	69	9	1
1956	1,933	747	360	142	41	52	88	137	156	137	61	12	-
1955	1,873	676	361	130	57	62	103	127	147	129	65	13	3
1954	1,829	645	342	137	50	72	91	132	138	128	80	13	1
1953	2,059	687	437	148	58	81	123	159	164	129	59	14	-
1952	1,918	645	379	147	67	77	133	157	137	110	53	11	2
1951	1,881	565	397	143	77	90	133	145	145	125	48	12	1
1950	1,839	573	366	144	86	88	112	147	152	113	45	11	2
<b>MALE</b>													
1959	1,305	494	256	89	47	35	66	68	99	105	39	7	-
1958	1,307	504	259	80	38	47	45	88	97	102	37	10	-
1957	1,159	447	204	83	45	22	51	81	109	79	37	1	-
1956	1,147	440	213	68	29	28	49	86	101	92	35	6	-
1955	1,100	409	197	71	39	28	61	79	92	76	36	9	3
1954	1,059	365	207	81	24	37	56	81	85	71	44	8	-
1953	1,221	389	255	94	32	44	80	97	106	83	34	7	-
1952	1,086	358	218	81	34	47	75	103	79	56	28	6	1
1951	1,166	326	246	97	48	56	88	93	97	85	26	4	-
1950	1,095	323	211	84	53	48	72	95	103	73	25	8	-
<b>FEMALE</b>													
1959	903	343	141	60	24	38	55	51	80	70	35	5	1
1958	940	373	185	57	28	28	32	62	83	52	29	11	-
1957	866	322	164	73	23	31	41	59	54	58	32	8	1
1956	786	307	147	74	12	24	39	51	55	45	26	6	-
1955	773	267	164	59	18	34	42	48	55	53	29	4	-
1954	770	280	135	56	26	35	35	51	53	57	36	5	1
1953	838	298	182	54	26	37	43	62	58	46	25	7	-
1952	832	287	161	66	33	30	58	54	58	54	25	5	1
1951	715	239	151	46	29	34	45	52	48	40	22	8	1
1950	744	250	155	60	33	40	40	52	49	40	20	3	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	1,585	570	292	117	46	48	77	84	132	146	63	10	-
1958	1,622	601	328	113	50	46	48	102	129	126	61	18	-
1957	1,431	511	275	120	45	32	64	88	114	118	57	7	-
1956	1,366	499	266	118	29	34	49	91	106	111	52	11	-
1955	1,339	458	280	102	39	49	65	79	101	101	53	9	3
1954	1,282	432	245	96	31	46	63	91	102	100	64	11	1
1953	1,529	482	340	116	39	56	74	112	130	114	53	13	-
1952	1,387	452	273	120	44	40	91	111	112	89	44	9	2
1951	1,357	381	294	120	47	55	89	106	113	100	42	10	-
1950	1,322	384	269	110	64	59	70	103	117	98	37	10	1
<b>WHITE, MALE</b>													
1959	941	328	184	72	30	24	48	48	77	90	33	7	-
1958	958	352	192	66	30	30	28	61	74	80	35	10	-
1957	832	293	158	65	30	15	39	53	77	70	31	1	-
1956	829	308	157	54	20	20	27	66	64	77	31	5	-
1955	806	278	162	57	26	24	39	51	72	57	31	6	3
1954	758	249	147	59	13	24	40	60	67	58	35	6	-
1953	922	273	196	77	23	32	49	71	91	73	30	7	-
1952	796	262	157	64	23	27	53	71	64	45	24	5	1
1951	869	225	188	83	30	37	59	72	78	70	23	4	-
1950	794	227	151	67	39	31	44	67	77	64	20	7	-
<b>WHITE, FEMALE</b>													
1959	644	242	108	45	16	24	29	36	55	56	30	3	-
1958	664	249	136	47	20	16	20	41	55	46	26	8	-
1957	599	218	117	55	15	17	25	35	37	48	26	6	-
1956	537	191	109	64	9	14	22	25	42	34	21	6	-
1955	533	180	118	45	13	25	26	28	29	44	22	3	-
1954	524	183	98	37	18	22	23	31	35	42	29	5	1
1953	607	209	144	39	16	24	25	41	39	41	23	6	-
1952	591	190	116	56	21	13	38	40	48	44	20	4	1
1951	488	156	106	37	17	18	30	34	35	30	19	6	-
1950	528	157	118	43	25	28	26	36	40	34	17	3	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MENINGITIS, EXCEPT MENINGOCOCCAL AND TUBERCULOUS (340)													
ALL OTHER, BOTH SEXES													
1959	623	267	105	32	25	25	44	35	47	29	11	2	1
1958	625	276	116	24	16	29	29	48	51	28	5	3	-
1957	594	258	93	36	23	21	28	52	49	19	12	2	1
1956	567	248	94	24	12	18	39	46	50	26	9	1	-
1955	534	218	81	28	18	13	38	48	46	28	12	4	-
1954	547	213	97	41	19	26	28	41	36	28	16	2	-
1953	530	205	97	32	19	25	49	47	34	15	6	1	-
1952	531	193	106	27	23	37	42	46	25	21	9	2	-
1951	524	184	103	23	30	35	44	39	32	25	6	2	1
1950	517	189	97	34	22	29	42	44	35	15	8	1	1
ALL OTHER, MALE													
1959	364	166	72	17	17	11	18	20	22	15	6	-	-
1958	349	152	67	14	8	17	17	27	23	22	2	-	-
1957	327	154	46	18	15	7	12	28	32	9	6	-	-
1956	318	132	56	14	9	8	22	20	37	15	4	1	-
1955	294	131	35	14	13	4	22	28	20	19	5	3	-
1954	301	116	60	22	11	13	16	21	18	13	9	2	-
1953	299	116	59	17	9	12	31	26	15	10	4	-	-
1952	290	96	61	17	11	20	22	32	15	11	4	1	-
1951	297	101	58	14	18	19	29	21	19	15	3	-	-
1950	301	96	60	17	14	17	28	28	26	9	5	1	-
ALL OTHER, FEMALE													
1959	259	101	33	15	8	14	26	15	25	14	5	2	1
1958	276	124	49	10	8	12	12	21	28	6	3	3	-
1957	267	104	47	18	8	14	16	24	17	10	6	2	1
1956	249	116	38	10	3	10	17	26	13	11	5	-	-
1955	240	87	46	14	5	9	16	20	26	9	7	1	-
1954	246	97	37	19	8	13	12	20	18	15	7	-	-
1953	231	89	38	15	10	13	18	21	19	5	2	1	-
1952	241	97	45	10	12	17	20	14	10	10	5	1	-
1951	227	83	45	9	12	16	15	18	13	10	3	2	1
1950	216	93	37	17	8	12	14	16	9	6	3	-	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)													
BOTH SEXES													
1959	910,635	533	464	902	2,023	6,141	23,773	69,871	144,283	254,735	275,154	132,337	419
1958	907,316	554	449	872	2,041	6,395	24,150	70,265	145,779	255,459	271,996	128,952	404
1957	891,825	580	485	1,013	2,278	6,527	24,687	69,701	147,186	252,731	262,663	123,486	488
1956	854,152	576	472	1,006	2,159	6,537	23,758	67,278	142,902	241,531	252,473	114,966	494
1955	831,353	560	456	1,022	2,262	6,658	24,015	66,872	139,784	235,237	244,726	109,086	675
1954	798,091	594	457	1,125	2,367	6,981	24,132	68,495	139,173	226,980	229,014	98,311	462
1953	814,662	625	487	1,223	2,549	7,481	26,044	71,313	145,460	230,431	231,761	96,823	465
1952	796,871	589	528	1,275	2,750	7,627	26,447	71,807	145,575	223,818	224,859	91,077	519
1951	786,799	526	481	1,248	2,834	7,648	26,479	71,532	143,974	221,420	221,721	88,294	642
1950	769,751	360	491	1,344	2,991	7,705	26,249	71,366	142,823	217,591	213,804	84,375	652
MALE													
1959	507,896	301	260	460	1,044	3,555	16,183	49,798	96,765	150,453	135,299	53,554	224
1958	505,683	311	242	453	1,094	3,641	16,333	49,131	97,063	150,387	134,216	52,609	203
1957	498,395	336	256	515	1,218	3,739	16,557	48,540	98,090	148,327	129,644	50,894	279
1956	478,884	319	247	501	1,113	3,788	15,863	46,946	95,106	141,750	125,281	47,690	280
1955	467,131	310	256	521	1,183	3,833	16,007	46,356	92,997	137,994	121,708	45,585	381
1954	449,037	338	254	569	1,215	3,936	15,740	46,742	92,007	132,643	114,257	41,072	264
1953	458,410	367	250	599	1,308	4,168	16,749	48,205	95,969	133,936	115,694	40,893	272
1952	448,329	354	297	625	1,452	4,180	16,776	48,157	95,752	129,473	112,584	38,388	291
1951	442,864	303	263	629	1,400	4,184	16,640	47,325	93,948	128,663	111,314	37,816	379
1950	434,007	223	269	630	1,505	4,158	16,326	47,205	92,965	125,921	107,941	36,477	387
FEMALE													
1959	402,739	232	204	442	979	2,586	7,590	20,073	47,518	104,282	139,855	78,783	195
1958	401,633	243	207	419	947	2,754	7,817	21,134	48,716	105,072	137,780	76,343	201
1957	393,430	244	229	498	1,060	2,788	8,130	21,161	49,096	104,404	133,019	72,592	209
1956	375,268	257	225	505	1,046	2,749	7,895	20,332	47,796	99,781	127,192	67,276	214
1955	364,222	250	200	501	1,079	2,825	8,008	20,516	46,787	97,243	123,018	63,501	294
1954	349,054	256	203	556	1,152	3,045	8,392	21,753	47,166	94,337	114,757	57,239	198
1953	356,252	258	237	624	1,241	3,313	9,295	23,108	49,491	96,495	116,067	55,930	193
1952	348,542	235	231	650	1,298	3,447	9,671	23,650	49,823	94,345	112,275	52,689	228
1951	343,935	223	218	619	1,434	3,464	9,839	24,207	50,026	92,757	110,407	50,478	263
1950	335,744	137	222	714	1,486	3,547	9,923	24,161	49,858	91,670	105,863	47,898	265

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)													
WHITE, BOTH SEXES													
1959	821,293	384	321	687	1,526	4,362	18,044	56,978	123,182	229,873	259,809	125,849	278
1958	816,770	402	324	675	1,581	4,476	18,416	56,702	124,186	230,313	256,839	122,625	231
1957	802,174	431	334	754	1,709	4,566	18,434	56,136	125,320	228,276	248,440	117,467	307
1956	769,796	410	321	796	1,622	4,685	17,932	54,196	122,224	218,731	239,072	109,472	335
1955	749,944	403	328	800	1,708	4,849	18,073	53,709	120,258	213,427	231,859	104,039	491
1954	719,064	434	336	873	1,780	5,042	18,061	54,793	120,343	206,355	217,110	93,638	299
1953	732,481	462	369	954	1,899	5,375	19,505	56,753	125,753	209,360	219,674	92,057	320
1952	715,166	437	417	984	2,053	5,484	19,837	57,142	125,718	203,207	213,038	86,450	399
1951	706,668	418	394	946	2,140	5,544	19,755	57,014	124,694	201,288	210,116	83,877	482
1950	691,296	274	391	1,065	2,266	5,622	19,576	56,911	123,637	198,286	202,701	80,100	467
WHITE, MALE													
1959	461,196	228	178	360	827	2,690	13,193	42,836	85,403	137,184	127,518	50,629	150
1958	458,680	233	176	356	884	2,770	13,478	42,068	85,570	136,963	126,369	49,698	115
1957	451,870	248	176	391	952	2,813	13,426	41,503	86,332	135,407	122,326	48,109	187
1956	435,234	232	173	403	865	2,922	13,021	40,132	84,142	129,664	118,284	45,195	201
1955	424,689	235	185	415	927	3,023	12,978	39,501	82,480	126,358	114,998	43,310	279
1954	407,932	251	187	442	960	3,060	12,778	39,684	81,745	121,764	107,946	38,936	179
1953	415,306	281	182	481	1,026	3,179	13,576	40,571	85,131	122,650	109,310	38,731	188
1952	405,676	267	239	478	1,135	3,253	13,646	40,654	84,866	118,426	106,201	36,286	225
1951	401,354	238	206	487	1,113	3,249	13,470	39,901	83,550	117,873	105,152	35,824	291
1950	393,299	178	214	510	1,182	3,237	13,210	39,815	82,648	115,520	101,945	34,557	283
WHITE, FEMALE													
1959	360,097	156	143	327	699	1,672	4,851	14,142	37,779	92,689	132,291	75,220	128
1958	358,090	169	148	319	697	1,706	4,938	14,634	38,616	93,350	130,470	72,927	116
1957	350,304	183	158	363	757	1,753	5,008	14,633	38,988	92,869	126,114	69,358	120
1956	334,562	178	148	393	757	1,763	4,911	14,064	38,082	89,067	120,788	64,277	134
1955	325,255	168	143	385	781	1,826	5,095	14,208	37,778	87,069	116,861	60,729	212
1954	311,132	183	149	431	820	1,982	5,283	15,109	38,598	84,591	109,164	54,702	120
1953	317,175	181	187	473	873	2,196	5,929	16,182	40,622	86,710	110,364	53,326	132
1952	309,490	170	178	506	918	2,231	6,191	16,488	40,852	84,781	106,837	50,164	174
1951	305,314	180	188	459	1,027	2,295	6,285	17,113	41,144	83,415	104,964	48,053	191
1950	297,997	96	177	555	1,084	2,385	6,366	17,096	40,989	82,766	100,756	45,543	184

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
MAJOR CARDIOVASCULAR-RENAL DISEASES (330-334, 400-468, 592-594)													
ALL OTHER, BOTH SEXES													
1959	89,342	149	143	215	497	1,779	5,729	12,893	21,101	24,862	15,345	6,488	141
1958	90,546	152	125	197	460	1,919	5,734	13,563	21,593	25,146	15,157	6,327	173
1957	89,651	149	151	259	569	1,961	6,253	13,565	21,866	24,455	14,223	6,019	181
1956	84,356	166	151	210	537	1,852	5,826	13,082	20,678	22,800	13,401	5,494	159
1955	81,409	157	128	222	554	1,809	5,942	13,163	19,526	21,810	12,867	5,047	184
1954	79,027	160	121	252	587	1,939	6,071	13,702	18,830	20,625	11,904	4,673	163
1953	82,181	163	118	269	650	2,106	6,539	14,560	19,707	21,071	12,087	4,766	145
1952	81,705	152	111	291	697	2,143	6,610	14,665	19,857	20,611	11,821	4,627	120
1951	80,131	108	87	302	694	2,104	6,724	14,518	19,280	20,132	11,605	4,417	160
1950	78,455	86	100	279	725	2,083	6,673	14,455	19,186	19,305	11,103	4,275	185
ALL OTHER, MALE													
1959	46,700	73	82	100	217	865	2,990	6,962	11,362	13,269	7,781	2,925	74
1958	47,003	78	66	97	210	871	2,855	7,063	11,493	13,424	7,847	2,911	88
1957	46,525	88	80	124	266	926	3,131	7,037	11,758	12,920	7,318	2,785	92
1956	43,650	87	74	98	248	866	2,842	6,814	10,964	12,086	6,997	2,495	79
1955	42,442	75	71	106	256	810	3,029	6,855	10,517	11,636	6,710	2,275	102
1954	41,105	87	67	127	255	876	2,962	7,058	10,262	10,879	6,311	2,136	85
1953	43,104	86	68	118	282	989	3,173	7,634	10,838	11,286	6,384	2,162	84
1952	42,653	87	58	147	317	927	3,130	7,503	10,886	11,047	6,383	2,102	66
1951	41,510	65	57	142	287	935	3,170	7,424	10,398	10,790	6,162	1,992	88
1950	40,708	45	55	120	323	921	3,116	7,390	10,317	10,401	5,996	1,920	104
ALL OTHER, FEMALE													
1959	42,642	76	61	115	280	914	2,739	5,931	9,739	11,593	7,564	3,563	67
1958	43,543	74	59	100	250	1,048	2,879	6,500	10,100	11,722	7,310	3,416	85
1957	43,126	61	71	135	303	1,035	3,122	6,528	10,108	11,535	6,905	3,234	89
1956	40,706	79	77	112	289	986	2,984	6,268	9,714	10,714	6,404	2,999	80
1955	38,967	82	57	116	298	999	2,913	6,308	9,009	10,174	6,157	2,772	82
1954	37,922	73	54	125	332	1,063	3,109	6,644	8,568	9,746	5,593	2,537	78
1953	39,077	77	50	151	368	1,117	3,366	6,926	8,869	9,785	5,703	2,604	61
1952	39,052	65	53	144	380	1,216	3,480	7,162	8,971	9,564	5,438	2,525	54
1951	38,621	43	30	160	407	1,169	3,554	7,094	8,882	9,342	5,443	2,425	72
1950	37,747	41	45	159	402	1,162	3,557	7,065	8,869	8,904	5,107	2,355	81

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)													
BOTH SEXES													
1959	898,336	502	435	725	1,672	5,485	22,752	68,365	142,293	252,105	272,513	131,079	410
1958	893,489	524	412	711	1,621	5,701	23,050	68,603	143,605	252,349	269,013	127,504	396
1957	876,793	561	439	780	1,812	5,713	23,416	67,963	144,707	249,379	259,557	121,982	484
1956	838,848	544	414	795	1,755	5,751	22,563	65,524	140,415	238,101	249,109	113,398	479
1955	815,532	539	394	814	1,799	5,901	22,807	65,069	137,197	231,637	241,166	107,545	664
1954	781,018	561	371	906	1,917	6,104	22,909	66,492	136,293	223,110	225,161	96,746	448
1953	795,888	604	393	978	2,038	6,544	24,647	69,086	142,355	226,148	227,622	95,023	450
1952	776,114	561	431	1,029	2,273	6,694	24,899	69,264	142,010	219,133	220,112	89,203	505
1951	764,281	501	363	995	2,294	6,679	24,888	68,943	140,085	216,269	216,545	86,099	620
1950	745,074	336	364	1,112	2,425	6,639	24,481	68,534	138,696	211,769	208,091	82,004	623
MALE													
1959	501,196	281	245	386	846	3,158	15,595	48,893	95,652	149,031	133,927	52,962	220
1958	498,185	294	221	363	854	3,262	15,673	48,200	95,838	148,689	132,669	51,923	199
1957	490,381	326	231	405	942	3,243	15,859	47,554	96,747	146,491	128,109	50,195	279
1956	470,724	300	213	394	881	3,318	15,211	45,970	93,742	139,939	123,567	46,918	271
1955	458,660	291	217	419	907	3,397	15,316	45,363	91,605	136,054	119,854	44,861	376
1954	439,890	321	211	447	959	3,456	15,071	45,664	90,443	130,511	112,226	40,325	256
1953	448,407	353	201	472	1,034	3,660	16,006	46,976	94,299	131,624	113,501	40,016	265
1952	437,113	338	235	493	1,181	3,657	15,970	46,720	93,785	126,914	110,050	37,488	282
1951	430,920	285	193	494	1,118	3,655	15,808	45,883	91,804	125,875	108,622	36,816	367
1950	420,922	205	198	510	1,180	3,600	15,402	45,677	90,677	122,794	104,936	35,370	373
FEMALE													
1959	397,140	221	190	339	826	2,327	7,157	19,472	46,641	103,074	138,586	78,117	190
1958	395,304	230	191	348	767	2,439	7,377	20,403	47,767	103,660	136,344	75,581	197
1957	386,412	235	208	375	870	2,470	7,557	20,409	47,960	102,888	131,448	71,787	205
1956	368,124	244	201	401	874	2,433	7,352	19,554	46,673	98,162	125,542	66,480	208
1955	356,872	248	177	395	892	2,504	7,491	19,706	45,592	95,583	121,312	62,684	288
1954	341,128	240	160	459	958	2,648	7,838	20,828	45,850	92,599	112,935	56,421	192
1953	347,481	251	192	506	1,004	2,884	8,641	22,110	48,056	94,524	114,121	55,007	185
1952	339,001	223	196	536	1,092	3,037	8,929	22,544	48,225	92,219	110,062	51,715	223
1951	333,361	216	170	501	1,176	3,024	9,080	23,060	48,281	90,394	107,923	49,283	253
1950	324,152	131	166	602	1,245	3,039	9,079	22,857	48,019	88,975	103,155	46,634	250

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)													
WHITE, BOTH SEXES													
1959	811,675	362	298	541	1,233	3,862	17,341	55,939	121,706	227,850	257,526	124,744	273
1958	805,887	379	297	539	1,222	3,946	17,651	55,549	122,650	227,825	254,248	121,352	229
1957	790,295	416	300	557	1,324	3,953	17,573	54,918	123,463	225,583	245,754	116,147	307
1956	757,699	386	276	620	1,293	4,094	17,114	53,016	120,367	215,996	236,128	108,080	329
1955	737,253	384	281	620	1,318	4,268	17,247	52,443	118,319	210,478	228,739	102,669	487
1954	705,390	411	265	685	1,408	4,374	17,202	53,383	118,145	203,239	213,729	92,253	296
1953	717,464	443	293	735	1,477	4,663	18,536	55,188	123,387	205,901	216,080	90,453	308
1952	698,816	411	334	770	1,660	4,775	18,788	55,402	123,074	199,437	208,940	84,832	393
1951	688,695	400	294	724	1,692	4,818	18,674	55,216	121,714	197,155	205,583	81,958	467
1950	671,528	264	282	857	1,781	4,814	18,361	54,916	120,505	193,604	197,706	77,986	452
WHITE, MALE													
1959	455,892	213	166	298	657	2,374	12,762	42,174	84,548	136,096	126,334	50,122	148
1958	452,703	221	162	284	679	2,452	12,982	41,390	84,673	135,609	125,025	49,111	115
1957	445,435	240	160	297	721	2,423	12,916	40,749	85,312	133,906	121,011	47,513	187
1956	428,703	218	150	317	671	2,535	12,528	39,430	83,114	128,222	116,800	44,522	196
1955	417,767	218	155	327	685	2,664	12,470	38,758	81,400	124,767	113,384	42,660	279
1954	400,591	240	153	340	745	2,674	12,281	38,887	80,520	120,074	106,214	38,287	176
1953	407,185	268	143	365	782	2,773	13,012	39,654	83,837	120,794	107,426	37,949	182
1952	396,729	253	186	364	907	2,828	13,057	39,613	83,374	116,368	104,053	35,506	220
1951	391,701	226	149	369	867	2,819	12,853	38,834	81,864	115,639	102,833	34,965	283
1950	382,686	169	154	400	897	2,780	12,506	38,663	80,879	113,025	99,349	33,587	277
WHITE, FEMALE													
1959	355,783	149	132	243	576	1,488	4,579	13,765	37,158	91,754	131,192	74,622	125
1958	353,184	158	135	255	543	1,494	4,669	14,159	37,977	92,216	129,223	72,241	114
1957	344,860	176	140	260	603	1,530	4,657	14,169	38,151	91,677	124,743	68,634	120
1956	328,996	168	126	303	622	1,559	4,586	13,586	37,253	87,774	119,328	63,558	133
1955	319,486	166	126	293	633	1,604	4,777	13,685	36,919	85,711	115,355	60,009	208
1954	304,799	171	112	345	663	1,700	4,921	14,496	37,625	83,165	107,515	53,966	120
1953	310,279	175	150	370	695	1,890	5,524	15,534	39,550	85,107	108,654	52,504	126
1952	302,087	158	148	406	753	1,947	5,731	15,789	39,700	83,069	104,887	49,326	173
1951	296,994	174	145	355	825	1,999	5,821	16,382	39,850	81,516	102,750	46,993	184
1950	288,842	95	128	457	884	2,034	5,855	16,253	39,626	80,579	98,357	44,399	175

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>DISEASES OF CARDIOVASCULAR SYSTEM (330-334, 400-468)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	86,661	140	137	184	439	1,623	5,411	12,426	20,587	24,255	14,987	6,335	137
1958	87,602	145	115	172	399	1,755	5,399	13,054	20,955	24,524	14,765	6,152	167
1957	86,498	145	139	223	488	1,760	5,843	13,045	21,244	23,796	13,803	5,835	177
1956	81,149	158	138	175	462	1,657	5,449	12,508	20,048	22,105	12,981	5,318	150
1955	78,279	155	113	194	481	1,633	5,560	12,626	18,878	21,159	12,427	4,876	177
1954	75,628	150	106	221	509	1,730	5,707	13,109	18,148	19,871	11,432	4,493	152
1953	78,424	161	100	243	561	1,881	6,111	13,898	18,968	20,247	11,542	4,570	142
1952	77,298	150	97	259	613	1,919	6,111	13,862	18,936	19,696	11,172	4,371	112
1951	75,586	101	69	271	602	1,861	6,214	13,727	18,371	19,114	10,962	4,141	153
1950	73,546	72	82	255	644	1,825	6,120	13,618	18,191	18,165	10,385	4,018	171
<b>ALL OTHER, MALE</b>													
1959	45,304	68	79	88	189	784	2,833	6,719	11,104	12,935	7,593	2,840	72
1958	45,482	73	59	79	175	810	2,691	6,810	11,165	13,080	7,644	2,812	84
1957	44,946	86	71	108	221	820	2,943	6,805	11,435	12,585	7,098	2,682	92
1956	42,021	82	63	77	210	783	2,683	6,540	10,628	11,717	6,767	2,396	75
1955	40,893	73	62	92	222	733	2,846	6,605	10,205	11,287	6,470	2,201	97
1954	39,299	81	58	107	214	782	2,790	6,777	9,923	10,437	6,012	2,038	80
1953	41,222	85	58	107	252	887	2,994	7,322	10,462	10,830	6,075	2,067	83
1952	40,384	85	49	129	274	829	2,913	7,107	10,411	10,546	5,997	1,982	62
1951	39,219	59	44	125	251	836	2,955	7,049	9,940	10,236	5,789	1,851	84
1950	38,236	36	44	110	283	820	2,896	7,014	9,798	9,769	5,587	1,783	96
<b>ALL OTHER, FEMALE</b>													
1959	41,357	72	58	96	250	839	2,578	5,707	9,483	11,320	7,394	3,495	65
1958	42,120	72	56	93	224	945	2,708	6,244	9,790	11,444	7,121	3,340	83
1957	41,552	59	68	115	267	940	2,900	6,240	9,809	11,211	6,705	3,153	85
1956	39,128	76	75	98	252	874	2,766	5,968	9,420	10,388	6,214	2,922	75
1955	37,386	82	51	102	259	900	2,714	6,021	8,673	9,872	5,957	2,675	80
1954	36,329	69	48	114	295	948	2,917	6,332	8,225	9,434	5,420	2,455	72
1953	37,202	76	42	136	309	994	3,117	6,576	8,506	9,417	5,467	2,503	59
1952	36,914	65	48	130	339	1,090	3,198	6,755	8,525	9,150	5,175	2,389	50
1951	36,367	42	25	146	351	1,025	3,259	6,678	8,431	8,878	5,173	2,290	69
1950	35,310	36	38	145	361	1,005	3,224	6,604	8,393	8,396	4,798	2,235	75

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
CEREBROVASCULAR DISEASES (ICD 330-334)													
BOTH SEXES													
1959	191,376	168	165	226	408	1,091	3,567	9,949	23,174	51,665	68,180	32,695	88
1958	190,758	187	160	244	401	1,138	3,657	10,263	23,684	52,242	67,067	31,621	94
1957	187,709	197	169	252	433	1,152	3,889	10,762	24,220	52,337	64,419	29,766	113
1956	177,845	177	164	219	399	1,132	3,728	10,437	23,869	49,917	60,666	27,013	124
1955	174,142	212	136	209	390	1,188	3,774	10,574	23,810	49,528	58,932	25,239	150
1954	167,777	185	133	213	401	1,097	4,026	11,310	24,678	47,950	55,392	22,288	104
1953	169,800	206	134	174	360	1,210	4,301	12,175	25,981	49,041	55,028	21,086	104
1952	166,331	182	158	195	410	1,196	4,209	12,408	26,328	47,984	53,144	20,013	104
1951	163,550	163	121	160	379	1,140	4,247	12,435	26,258	48,059	51,967	18,462	159
1950	156,751	160	119	113	363	1,009	4,015	12,209	25,958	46,258	49,153	17,250	144
MALE													
1959	91,021	105	81	141	207	510	1,729	5,201	12,405	26,906	31,067	12,631	38
1958	91,076	105	99	130	223	530	1,708	5,254	12,812	27,311	30,724	12,140	40
1957	90,423	116	86	139	238	575	1,866	5,485	13,116	27,224	29,809	11,709	60
1956	85,618	102	76	111	199	561	1,778	5,283	12,792	25,974	28,026	10,653	63
1955	84,292	114	82	119	185	596	1,812	5,247	12,883	25,750	27,304	10,130	70
1954	81,174	100	75	119	191	526	1,927	5,556	13,120	24,945	25,784	8,777	54
1953	82,280	125	69	97	198	591	2,020	5,928	13,894	25,248	25,620	8,434	56
1952	80,964	120	84	106	215	568	1,943	6,051	14,104	24,600	25,024	8,097	52
1951	79,512	99	71	79	180	549	1,991	5,989	13,941	24,865	24,164	7,490	94
1950	76,672	103	72	60	192	491	1,855	5,879	13,742	23,723	23,258	7,220	77
FEMALE													
1959	100,355	63	84	85	201	581	1,838	4,748	10,769	24,759	37,113	20,064	50
1958	99,682	82	61	114	178	608	1,949	5,009	10,872	24,931	36,343	19,481	54
1957	97,286	81	83	113	195	577	2,023	5,277	11,104	25,113	34,610	18,057	53
1956	92,227	75	88	108	200	571	1,950	5,154	11,077	23,943	32,640	16,360	61
1955	89,850	98	54	90	205	592	1,962	5,327	10,927	23,778	31,628	15,109	80
1954	86,603	85	58	94	210	571	2,099	5,754	11,558	23,005	29,608	13,511	50
1953	87,520	81	65	77	162	619	2,281	6,247	12,087	23,793	29,408	12,652	48
1952	85,367	62	74	89	195	628	2,266	6,357	12,224	23,384	28,120	11,916	52
1951	84,038	64	50	81	199	591	2,256	6,446	12,317	23,194	27,803	10,972	65
1950	80,079	57	47	53	171	518	2,160	6,330	12,216	22,535	25,895	10,030	67

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
CEREBROVASCULAR DISEASES (ICD 330-334)													
WHITE, BOTH SEXES													
1959	167,661	127	129	188	318	761	2,240	6,823	17,532	44,656	63,870	30,966	51
1958	167,044	140	120	198	318	748	2,340	7,062	17,987	45,182	62,877	30,021	51
1957	164,477	147	131	207	335	730	2,435	7,449	18,584	45,552	60,603	28,242	62
1956	156,013	125	120	198	317	744	2,420	7,095	18,413	43,753	57,154	25,596	78
1955	152,834	146	109	178	291	817	2,366	7,297	18,497	43,455	55,591	23,996	91
1954	147,006	137	102	171	313	742	2,562	7,724	19,532	42,253	52,234	21,173	63
1953	148,964	149	106	146	276	790	2,801	8,420	20,779	43,435	51,972	20,022	68
1952	145,677	126	135	163	307	833	2,733	8,677	21,102	42,465	50,154	18,913	69
1951	143,450	133	103	136	273	777	2,790	8,758	21,214	42,668	48,996	17,497	105
1950	137,482	121	100	93	269	670	2,565	8,536	20,938	41,363	46,429	16,297	101
WHITE, MALE													
1959	79,795	77	64	116	172	375	1,143	3,732	9,668	23,483	29,049	11,896	20
1958	79,666	81	73	105	184	372	1,174	3,782	9,973	23,793	28,680	11,427	22
1957	79,360	83	69	118	192	410	1,254	3,961	10,345	23,891	27,930	11,074	33
1956	75,305	76	56	105	163	384	1,236	3,794	10,195	22,900	26,279	10,075	42
1955	74,103	82	64	102	140	436	1,167	3,819	10,281	22,742	25,637	9,592	41
1954	71,333	77	58	94	147	380	1,311	3,954	10,608	22,159	24,221	8,295	29
1953	72,306	95	54	85	156	422	1,394	4,180	11,316	22,451	24,122	7,993	38
1952	71,109	87	75	89	163	435	1,339	4,363	11,501	21,897	23,501	7,628	31
1951	69,984	81	55	63	127	396	1,402	4,301	11,497	22,228	22,697	7,074	63
1950	67,462	82	64	49	152	352	1,248	4,207	11,263	21,286	21,880	6,822	57
WHITE, FEMALE													
1959	87,866	50	65	72	146	386	1,097	3,091	7,864	21,173	34,821	19,070	31
1958	87,378	59	47	93	134	376	1,166	3,280	8,014	21,389	34,197	18,594	29
1957	85,117	64	62	89	143	320	1,181	3,488	8,239	21,661	32,673	17,168	29
1956	80,708	49	64	93	154	360	1,184	3,301	8,218	20,853	30,875	15,521	36
1955	78,731	64	45	76	151	381	1,199	3,478	8,216	20,713	29,954	14,404	50
1954	75,673	60	44	77	166	362	1,251	3,770	8,924	20,094	28,013	12,878	34
1953	76,658	54	52	61	120	368	1,407	4,240	9,463	20,984	27,850	12,029	30
1952	74,568	39	60	74	144	398	1,394	4,314	9,601	20,568	26,653	11,285	38
1951	73,466	52	48	73	146	381	1,388	4,457	9,717	20,440	26,299	10,423	42
1950	70,020	39	36	44	117	318	1,317	4,329	9,675	20,077	24,549	9,475	44

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
CEREBROVASCULAR DISEASES (ICD 330-334)													
ALL OTHER, BOTH SEXES													
1959	23,715	41	36	38	90	330	1,327	3,126	5,642	7,009	4,310	1,729	37
1958	23,714	47	40	46	83	390	1,317	3,201	5,697	7,060	4,190	1,600	43
1957	23,232	50	38	45	98	422	1,454	3,313	5,636	6,785	3,816	1,524	51
1956	21,832	52	44	21	82	388	1,308	3,342	5,456	6,164	3,512	1,417	46
1955	21,308	66	27	31	99	371	1,408	3,277	5,313	6,073	3,341	1,243	59
1954	20,771	48	31	42	88	355	1,464	3,586	5,146	5,697	3,158	1,115	41
1953	20,836	57	28	28	84	420	1,500	3,755	5,202	5,606	3,056	1,064	36
1952	20,654	56	23	32	103	363	1,476	3,731	5,226	5,519	2,990	1,100	35
1951	20,100	30	18	24	106	363	1,457	3,677	5,044	5,391	2,971	965	54
1950	19,269	39	19	20	94	339	1,450	3,673	5,020	4,895	2,724	953	43
ALL OTHER, MALE													
1959	11,226	28	17	25	35	135	586	1,469	2,737	3,423	2,018	735	18
1958	11,410	24	26	25	39	158	534	1,472	2,839	3,518	2,044	713	18
1957	11,063	33	17	21	46	165	612	1,524	2,771	3,333	1,879	635	27
1956	10,313	26	20	6	36	177	542	1,489	2,597	3,074	1,747	578	21
1955	10,189	32	18	17	45	160	645	1,428	2,602	3,008	1,667	538	29
1954	9,841	23	17	25	44	146	616	1,602	2,512	2,786	1,563	482	25
1953	9,974	30	15	12	42	169	626	1,748	2,578	2,797	1,498	441	18
1952	9,855	33	9	17	52	133	604	1,688	2,603	2,703	1,523	469	21
1951	9,528	18	16	16	53	153	589	1,688	2,444	2,637	1,467	416	31
1950	9,210	21	8	11	40	139	607	1,672	2,479	2,437	1,378	398	20
ALL OTHER, FEMALE													
1959	12,489	13	19	13	55	195	741	1,657	2,905	3,586	2,292	994	19
1958	12,304	23	14	21	44	232	783	1,729	2,858	3,542	2,146	887	25
1957	12,169	17	21	24	52	257	842	1,789	2,865	3,452	1,937	889	24
1956	11,519	26	24	15	46	211	766	1,853	2,859	3,090	1,765	839	25
1955	11,119	34	9	14	54	211	763	1,849	2,711	3,065	1,674	705	30
1954	10,930	25	14	17	44	209	848	1,984	2,634	2,911	1,595	633	16
1953	10,862	27	13	16	42	251	874	2,007	2,624	2,809	1,558	623	18
1952	10,799	23	14	15	51	230	872	2,043	2,623	2,816	1,467	631	14
1951	10,572	12	2	8	53	210	868	1,989	2,600	2,754	1,504	549	23
1950	10,059	18	11	9	54	200	843	2,001	2,541	2,458	1,346	555	23

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
DISEASES OF HEART (ICD 400-402, 410-413)													
BOTH SEXES													
1959	641,044	288	238	443	1,008	3,767	17,583	55,042	111,959	185,877	182,230	82,318	291
1958	637,246	299	220	430	1,000	3,901	17,756	54,845	113,050	185,184	180,305	79,980	276
1957	629,404	307	246	476	1,165	3,969	18,081	54,247	114,055	183,628	175,483	77,402	345
1956	604,028	312	220	513	1,161	4,010	17,411	52,188	110,568	175,612	169,343	72,366	324
1955	585,751	276	239	550	1,179	4,117	17,603	51,776	107,779	170,284	163,078	68,395	475
1954	561,374	318	211	631	1,305	4,294	17,446	52,345	106,192	164,166	152,391	61,765	310
1953	571,798	334	227	753	1,443	4,636	18,787	53,934	110,756	165,638	154,248	60,733	309
1952	556,724	314	239	789	1,627	4,786	19,092	53,930	110,045	160,067	148,822	56,648	365
1951	547,323	270	215	794	1,657	4,861	19,047	53,641	108,283	156,925	145,947	55,277	406
1950	537,629	128	211	953	1,822	4,977	18,949	53,628	107,534	154,890	141,303	52,801	433
MALE													
1959	375,906	151	143	222	550	2,367	12,985	41,615	78,672	113,299	91,875	33,869	158
1958	372,912	161	108	216	550	2,444	13,116	40,849	78,604	112,521	90,986	33,214	143
1957	368,795	177	127	239	624	2,412	13,209	40,249	79,459	111,196	88,557	32,340	206
1956	355,200	163	115	261	624	2,504	12,670	38,919	76,996	106,437	85,944	30,373	194
1955	345,172	144	124	277	626	2,544	12,752	38,374	75,150	103,275	82,863	28,762	281
1954	331,847	184	120	308	689	2,614	12,398	38,379	73,924	99,133	77,655	26,257	186
1953	337,834	189	116	361	756	2,765	13,151	39,196	76,811	99,717	78,550	26,035	187
1952	328,816	180	131	374	888	2,802	13,202	38,881	76,078	96,022	75,973	24,073	212
1951	323,882	154	110	392	857	2,816	13,010	38,206	74,434	94,508	75,150	24,001	244
1950	317,824	76	110	434	906	2,832	12,752	38,185	73,706	92,957	72,713	22,881	272
FEMALE													
1959	265,138	137	95	221	458	1,400	4,598	13,427	33,287	72,578	90,355	48,449	133
1958	264,334	138	112	214	450	1,457	4,640	13,996	34,446	72,663	89,319	46,766	133
1957	260,609	130	119	237	541	1,557	4,872	13,998	34,596	72,432	86,926	45,062	139
1956	248,828	149	105	252	537	1,506	4,741	13,269	33,572	69,175	83,399	41,993	130
1955	240,579	132	115	273	553	1,573	4,851	13,402	32,629	67,009	80,215	39,633	194
1954	229,527	134	91	323	616	1,680	5,048	13,966	32,268	65,033	74,736	35,508	124
1953	233,964	145	111	392	687	1,871	5,636	14,738	33,945	65,921	75,698	34,698	122
1952	227,908	134	108	415	739	1,984	5,890	15,049	33,967	64,045	72,849	32,575	153
1951	223,441	116	105	402	800	2,045	6,037	15,435	33,849	62,417	70,797	31,276	162
1950	219,805	52	101	519	916	2,145	6,197	15,443	33,828	61,933	68,590	29,920	161

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
DISEASES OF HEART (ICD 400-402, 410-413)													
WHITE, BOTH SEXES													
1959	585,128	196	144	308	732	2,745	14,186	46,675	98,354	170,302	172,872	78,406	208
1958	580,549	209	157	312	739	2,828	14,346	46,052	99,028	169,469	171,071	76,174	164
1957	572,606	232	153	314	840	2,891	14,314	45,391	99,655	168,185	166,631	73,773	227
1956	550,721	224	135	372	848	2,989	13,874	43,830	97,078	161,113	160,954	69,071	233
1955	534,718	200	155	394	863	3,095	14,084	43,233	95,263	156,596	155,076	65,386	373
1954	512,199	230	143	470	941	3,187	13,809	43,690	94,164	151,268	145,097	58,986	214
1953	520,143	250	160	545	1,029	3,447	14,790	44,700	98,052	152,366	146,790	57,797	217
1952	505,711	234	174	570	1,187	3,530	15,063	44,610	97,369	147,132	141,603	53,944	295
1951	497,391	214	165	555	1,228	3,619	14,896	44,404	95,987	144,458	138,919	52,630	316
1950	488,498	103	153	722	1,335	3,725	14,846	44,439	95,323	142,782	134,511	50,241	318
WHITE, MALE													
1959	345,557	115	84	162	417	1,839	11,098	36,872	71,012	104,773	87,004	32,064	117
1958	342,504	117	81	164	429	1,913	11,275	36,037	70,946	103,896	86,090	31,472	84
1957	338,279	134	80	162	472	1,875	11,177	35,442	71,466	102,816	83,905	30,606	144
1956	326,605	118	80	191	471	2,000	10,827	34,288	69,604	98,586	81,479	28,817	144
1955	317,586	111	80	204	479	2,062	10,858	33,654	68,112	95,805	78,611	27,389	221
1954	305,284	135	82	231	537	2,083	10,521	33,659	67,046	92,160	73,722	24,969	139
1953	309,748	147	76	267	573	2,156	11,086	34,119	69,528	92,458	74,536	24,673	129
1952	301,174	138	95	264	681	2,219	11,175	33,879	68,863	88,833	72,035	22,818	174
1951	297,062	121	82	288	676	2,237	10,917	33,280	67,500	87,601	71,364	22,801	195
1950	291,480	64	76	338	686	2,247	10,726	33,236	66,907	86,285	69,000	21,712	203
WHITE, FEMALE													
1959	239,571	81	60	146	315	906	3,088	9,803	27,342	65,529	85,868	46,342	91
1958	238,045	92	76	148	310	915	3,071	10,015	28,082	65,573	84,981	44,702	80
1957	234,327	98	73	152	368	1,016	3,137	9,949	28,189	65,369	82,726	43,167	83
1956	224,116	106	55	181	377	989	3,047	9,542	27,474	62,527	79,475	40,254	89
1955	217,132	89	75	190	384	1,033	3,226	9,579	27,151	60,791	76,465	37,997	152
1954	206,915	95	61	239	404	1,104	3,288	10,031	27,118	59,108	71,375	34,017	75
1953	210,395	103	84	278	456	1,291	3,704	10,581	28,524	59,908	72,254	33,124	88
1952	204,537	96	79	306	506	1,311	3,888	10,731	28,506	58,299	69,568	31,126	121
1951	200,329	93	83	267	552	1,382	3,979	11,124	28,487	56,857	67,555	29,829	121
1950	197,018	39	77	384	649	1,478	4,120	11,203	28,416	56,497	65,511	28,529	115

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
DISEASES OF HEART (ICD 400-402, 410-413)													
ALL OTHER, BOTH SEXES													
1959	55,916	92	94	135	276	1,022	3,397	8,367	13,605	15,575	9,358	3,912	83
1958	56,697	90	63	118	261	1,073	3,410	8,793	14,022	15,715	9,234	3,806	112
1957	56,798	75	93	162	325	1,078	3,767	8,856	14,400	15,443	8,852	3,629	118
1956	53,307	88	85	141	313	1,021	3,537	8,358	13,490	14,499	8,389	3,295	91
1955	51,033	76	84	156	316	1,022	3,519	8,543	12,516	13,688	8,002	3,009	102
1954	49,175	88	68	161	364	1,107	3,637	8,655	12,028	12,898	7,294	2,779	96
1953	51,655	84	67	208	414	1,189	3,997	9,234	12,704	13,272	7,458	2,936	92
1952	51,013	80	65	219	440	1,256	4,029	9,320	12,676	12,935	7,219	2,704	70
1951	49,932	56	50	239	429	1,242	4,151	9,237	12,296	12,467	7,028	2,647	90
1950	49,131	25	58	231	487	1,252	4,103	9,189	12,211	12,108	6,792	2,560	115
ALL OTHER, MALE													
1959	30,349	36	59	60	133	528	1,887	4,743	7,660	8,526	4,871	1,805	41
1958	30,408	44	27	52	121	531	1,841	4,812	7,658	8,625	4,896	1,742	59
1957	30,516	43	47	77	152	537	2,032	4,807	7,993	8,380	4,652	1,734	62
1956	28,595	45	35	70	153	504	1,843	4,631	7,392	7,851	4,465	1,556	50
1955	27,586	33	44	73	147	482	1,894	4,720	7,038	7,470	4,252	1,373	60
1954	26,563	49	38	77	152	531	1,877	4,720	6,878	6,973	3,933	1,288	47
1953	28,086	42	40	94	183	609	2,065	5,077	7,283	7,259	4,014	1,362	58
1952	27,642	42	36	110	207	583	2,027	5,002	7,215	7,189	3,938	1,255	38
1951	26,820	33	28	104	181	579	2,093	4,926	6,934	6,907	3,786	1,200	49
1950	26,344	12	34	96	220	585	2,026	4,949	6,799	6,672	3,713	1,169	69
ALL OTHER, FEMALE													
1959	25,567	56	35	75	143	494	1,510	3,624	5,945	7,049	4,487	2,107	42
1958	26,289	46	36	66	140	542	1,569	3,981	6,364	7,090	4,338	2,064	53
1957	26,282	32	46	85	173	541	1,735	4,049	6,407	7,063	4,200	1,895	56
1956	24,712	43	50	71	160	517	1,694	3,727	6,098	6,648	3,924	1,739	41
1955	23,447	43	40	83	169	540	1,625	3,823	5,478	6,218	3,750	1,636	42
1954	22,612	39	30	84	212	576	1,760	3,935	5,150	5,925	3,361	1,491	49
1953	23,569	42	27	114	231	580	1,932	4,157	5,421	6,013	3,444	1,574	34
1952	23,371	38	29	109	233	673	2,002	4,318	5,461	5,746	3,281	1,449	32
1951	23,112	23	22	135	248	663	2,058	4,311	5,362	5,560	3,242	1,447	41
1950	22,787	13	24	135	267	667	2,077	4,240	5,412	5,436	3,079	1,391	46

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)</b>													
<b>BOTH SEXES</b>													
1959	18,290	6	34	266	525	1,113	2,433	3,979	4,402	3,346	1,672	510	4
1958	18,796	12	38	250	503	1,202	2,477	4,130	4,418	3,384	1,838	541	3
1957	20,174	6	38	257	629	1,307	2,656	4,377	4,729	3,676	1,924	570	5
1956	20,029	18	44	332	659	1,348	2,613	4,240	4,589	3,545	1,995	640	6
1955	19,757	16	49	362	660	1,423	2,717	4,104	4,441	3,471	1,900	608	6
1954	19,553	15	56	452	740	1,504	2,670	4,039	4,270	3,312	1,912	576	7
1953	21,110	25	72	542	829	1,652	3,014	4,384	4,439	3,465	2,072	604	12
1952	21,337	16	66	582	925	1,703	3,222	4,123	4,433	3,402	2,213	647	5
1951	21,636	20	76	631	895	1,750	3,253	4,310	4,334	3,438	2,247	676	6
1950	22,316	19	64	716	1,019	1,861	3,322	4,171	4,384	3,531	2,424	787	18
<b>MALE</b>													
1959	8,769	1	22	119	250	534	1,224	1,997	2,153	1,589	688	190	2
1958	9,030	5	16	111	259	613	1,284	1,994	2,158	1,598	779	212	1
1957	9,812	4	20	117	299	618	1,361	2,171	2,414	1,738	856	212	2
1956	9,874	9	21	156	324	680	1,318	2,146	2,345	1,723	892	258	2
1955	9,775	8	29	171	320	683	1,412	2,047	2,297	1,723	859	225	1
1954	9,605	10	34	206	363	740	1,330	1,989	2,155	1,667	874	235	2
1953	10,484	10	37	246	414	807	1,464	2,192	2,343	1,779	952	233	7
1952	10,784	10	29	259	503	826	1,651	2,130	2,338	1,718	1,056	260	4
1951	10,805	11	44	299	446	864	1,636	2,158	2,263	1,765	1,031	286	2
1950	11,278	12	28	317	484	956	1,655	2,104	2,347	1,866	1,188	310	11
<b>FEMALE</b>													
1959	9,521	5	12	147	275	579	1,209	1,982	2,249	1,757	984	320	2
1958	9,766	7	22	139	244	589	1,193	2,136	2,260	1,786	1,059	329	2
1957	10,362	2	18	140	330	689	1,295	2,206	2,315	1,938	1,068	358	3
1956	10,155	9	23	176	335	668	1,295	2,094	2,244	1,822	1,103	382	4
1955	9,982	8	20	191	340	740	1,305	2,057	2,144	1,748	1,041	383	5
1954	9,948	5	22	246	377	764	1,340	2,050	2,115	1,645	1,038	341	5
1953	10,626	15	35	296	415	845	1,550	2,192	2,096	1,686	1,120	371	5
1952	10,553	6	37	323	422	877	1,571	1,993	2,095	1,684	1,157	387	1
1951	10,831	9	32	332	449	886	1,617	2,152	2,071	1,673	1,216	390	4
1950	11,038	7	36	399	535	905	1,667	2,067	2,037	1,665	1,236	477	7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	16,741	3	16	165	393	917	2,176	3,673	4,158	3,184	1,580	473	3
1958	17,210	7	30	167	388	986	2,183	3,807	4,158	3,227	1,750	505	2
1957	18,356	5	20	161	479	1,064	2,340	4,011	4,414	3,504	1,825	528	5
1956	18,233	12	29	220	493	1,139	2,293	3,916	4,289	3,340	1,891	607	4
1955	17,966	12	28	249	500	1,226	2,403	3,746	4,168	3,253	1,808	567	6
1954	17,689	12	39	322	563	1,243	2,355	3,698	4,018	3,114	1,782	539	4
1953	19,040	16	47	388	637	1,414	2,665	4,007	4,153	3,205	1,957	545	6
1952	19,132	8	44	414	724	1,450	2,900	3,726	4,095	3,110	2,059	597	5
1951	19,521	14	54	438	721	1,501	2,930	3,940	4,038	3,175	2,100	604	6
1950	19,959	10	38	532	793	1,611	2,967	3,774	4,026	3,243	2,234	717	14
<b>WHITE, MALE</b>													
1959	8,000	1	10	78	189	451	1,100	1,835	2,024	1,495	645	171	1
1958	8,274	2	13	77	211	521	1,145	1,842	2,020	1,513	732	197	1
1957	8,935	4	12	76	240	516	1,202	1,981	2,249	1,650	810	193	2
1956	8,987	7	18	102	251	588	1,181	1,978	2,188	1,601	834	238	1
1955	8,917	5	18	124	246	610	1,261	1,867	2,147	1,612	814	212	1
1954	8,694	8	23	150	292	629	1,166	1,807	2,029	1,565	811	213	1
1953	9,508	4	24	184	327	700	1,314	2,016	2,193	1,640	892	209	5
1952	9,708	6	21	182	406	732	1,497	1,920	2,164	1,569	974	233	4
1951	9,791	7	29	215	382	760	1,479	1,978	2,102	1,630	950	257	2
1950	10,096	7	15	242	384	835	1,486	1,904	2,138	1,712	1,077	287	9
<b>WHITE, FEMALE</b>													
1959	8,741	2	6	87	204	466	1,076	1,838	2,134	1,689	935	302	2
1958	8,936	5	17	90	177	465	1,038	1,965	2,138	1,714	1,018	308	1
1957	9,421	1	8	85	239	548	1,138	2,030	2,165	1,854	1,015	335	3
1956	9,246	5	11	118	242	551	1,112	1,938	2,101	1,739	1,057	369	3
1955	9,049	7	10	125	254	616	1,142	1,879	2,021	1,641	994	355	5
1954	8,995	4	16	172	271	614	1,189	1,891	1,989	1,549	971	326	3
1953	9,532	12	23	204	310	714	1,351	1,991	1,960	1,565	1,065	336	1
1952	9,424	2	23	232	318	718	1,403	1,806	1,931	1,541	1,085	364	1
1951	9,730	7	25	223	339	741	1,451	1,962	1,936	1,545	1,150	347	4
1950	9,863	3	23	290	409	776	1,481	1,870	1,888	1,531	1,157	430	5

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>RHEUMATIC FEVER AND CHRONIC RHEUMATIC HEART DISEASE (400-402, 410-416)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	1,549	3	18	101	132	196	257	306	244	162	92	37	1
1958	1,586	5	8	83	115	216	294	323	260	157	88	36	1
1957	1,818	1	18	96	150	243	316	366	315	172	99	42	-
1956	1,796	6	15	112	166	209	320	324	300	205	104	33	2
1955	1,791	4	21	113	160	197	314	358	273	218	92	41	-
1954	1,864	3	17	130	177	261	315	341	252	198	130	37	3
1953	2,070	9	25	154	192	238	349	377	286	260	115	59	6
1952	2,205	8	22	168	201	253	322	397	338	292	154	50	-
1951	2,115	6	22	193	174	249	323	370	296	263	147	72	-
1950	2,357	9	26	184	226	250	355	397	358	288	190	70	4
<b>ALL OTHER, MALE</b>													
1959	769	-	12	41	61	83	124	162	129	94	43	19	1
1958	756	3	3	34	48	92	139	152	138	85	47	15	-
1957	877	-	8	41	59	102	159	190	165	88	46	19	-
1956	887	2	3	54	73	92	137	168	157	122	58	20	1
1955	858	3	11	47	74	73	151	180	150	111	45	13	-
1954	911	2	11	56	71	111	164	182	126	102	63	22	1
1953	976	6	13	62	87	107	150	176	150	139	60	24	2
1952	1,076	4	8	77	97	94	154	210	174	149	82	27	-
1951	1,014	4	15	84	64	104	157	180	161	135	81	29	-
1950	1,182	5	13	75	100	121	169	200	209	154	111	23	2
<b>ALL OTHER, FEMALE</b>													
1959	780	3	6	60	71	113	133	144	115	68	49	18	-
1958	830	2	5	49	67	124	155	171	122	72	41	21	1
1957	941	1	10	55	91	141	157	176	150	84	53	23	-
1956	909	4	12	58	93	117	183	156	143	83	46	13	1
1955	933	1	10	66	86	124	163	178	123	107	47	28	-
1954	953	1	6	74	106	150	151	159	126	96	67	15	2
1953	1,094	3	12	92	105	131	199	201	136	121	55	35	4
1952	1,129	4	14	91	104	159	168	187	164	143	72	23	-
1951	1,101	2	7	109	110	145	166	190	135	128	66	43	-
1950	1,175	4	13	109	126	129	186	197	149	134	79	47	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)													
BOTH SEXES													
1959	474,143	13	12	25	178	1,656	11,903	41,774	87,867	143,886	131,674	54,954	201
1958	461,373	9	9	26	168	1,723	11,869	40,807	87,001	140,321	127,255	52,016	169
1957	452,507	47	14	38	189	1,644	11,811	39,528	86,856	138,002	123,835	50,312	231
1956	427,516	62	19	40	195	1,636	11,383	37,762	83,508	130,201	116,809	45,691	210
1955	405,830	47	20	35	208	1,678	11,060	37,044	80,172	123,897	109,407	41,937	325
1954	379,788	78	20	38	178	1,617	10,754	36,656	77,607	116,599	99,567	36,475	199
1953	373,597	60	26	45	224	1,696	10,986	36,389	78,596	114,413	96,842	34,139	181
1952	352,135	49	18	42	224	1,699	10,690	35,755	76,624	106,733	89,683	30,405	213
1951	336,619	35	12	26	269	1,591	10,443	34,628	73,770	102,507	84,987	28,107	244
1950	321,003	39	16	30	235	1,532	10,008	33,999	71,449	98,485	79,449	25,522	239
MALE													
1959	295,463	8	9	15	123	1,312	9,966	34,176	65,178	91,719	69,389	23,453	115
1958	288,607	5	5	15	116	1,322	9,909	33,216	64,406	89,490	67,428	22,603	92
1957	282,514	34	10	25	114	1,263	9,850	32,259	64,283	87,448	65,266	21,816	146
1956	268,559	31	13	19	133	1,276	9,524	30,936	61,947	82,704	61,832	20,007	137
1955	256,318	23	8	17	132	1,302	9,203	30,230	59,661	78,943	58,218	18,371	210
1954	241,826	45	13	24	107	1,250	8,897	29,836	57,768	74,385	53,282	16,090	129
1953	238,542	36	13	28	122	1,279	9,096	29,477	58,324	72,682	52,062	15,306	117
1952	225,524	30	10	22	130	1,280	8,809	28,830	57,001	67,602	48,173	13,505	132
1951	217,348	22	9	21	153	1,193	8,570	27,938	54,795	65,500	46,229	12,753	165
1950	207,864	26	12	15	141	1,120	8,175	27,398	53,104	62,798	43,296	11,623	156
FEMALE													
1959	178,680	5	3	10	55	344	1,937	7,598	22,689	52,167	62,285	31,501	86
1958	172,766	4	4	11	52	401	1,960	7,591	22,595	50,831	59,827	29,413	77
1957	169,993	13	4	13	75	381	1,961	7,269	22,573	50,554	58,569	28,496	85
1956	158,957	31	6	21	62	360	1,859	6,826	21,561	47,497	54,977	25,684	73
1955	149,512	24	12	18	76	376	1,857	6,814	20,511	44,954	51,189	23,566	115
1954	137,962	33	7	14	71	367	1,857	6,820	19,839	42,214	46,285	20,385	70
1953	135,055	24	13	17	102	417	1,890	6,912	20,272	41,731	44,780	18,833	64
1952	126,611	19	8	20	94	419	1,881	6,925	19,623	39,131	41,510	16,900	81
1951	119,271	13	3	5	116	398	1,873	6,690	18,975	37,007	38,758	15,354	79
1950	113,139	13	4	15	94	412	1,833	6,601	18,345	35,687	36,153	13,899	83

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	443,814	9	8	20	130	1,318	10,303	37,272	80,136	135,132	126,448	52,875	163
1958	431,994	5	8	20	127	1,335	10,341	36,304	79,503	131,855	122,327	50,057	112
1957	423,622	36	10	27	132	1,284	10,132	35,099	79,287	129,899	119,149	48,395	172
1956	401,212	42	15	37	150	1,307	9,817	33,737	76,622	122,692	112,574	44,058	161
1955	381,631	30	16	28	141	1,326	9,632	33,004	73,995	117,175	105,495	40,520	269
1954	357,660	57	12	34	127	1,303	9,288	32,794	71,857	110,591	96,202	35,238	157
1953	351,254	44	22	35	147	1,317	9,509	32,297	72,865	108,369	93,584	32,924	141
1952	331,790	38	15	32	169	1,333	9,243	31,977	71,245	101,437	86,776	29,334	191
1951	317,520	28	10	22	199	1,274	8,981	31,001	68,760	97,541	82,328	27,163	213
1950	303,133	34	13	24	175	1,221	8,672	30,561	66,735	93,900	76,942	24,661	195
<b>WHITE, MALE</b>													
1959	277,674	6	7	13	101	1,096	8,927	31,378	60,482	86,608	66,521	22,439	96
1958	271,556	3	5	12	93	1,103	8,971	30,468	59,939	84,545	64,672	21,684	61
1957	265,752	27	8	21	84	1,056	8,812	29,584	59,745	82,776	62,644	20,875	120
1956	253,385	21	10	18	107	1,092	8,572	28,471	57,852	78,428	59,481	19,222	111
1955	242,246	18	4	13	102	1,123	8,320	27,756	55,900	75,048	56,074	17,713	175
1954	228,766	36	7	21	84	1,072	8,037	27,486	54,205	70,855	51,358	15,496	109
1953	225,416	28	10	20	94	1,064	8,182	26,997	54,801	69,221	50,206	14,704	89
1952	213,533	24	8	15	106	1,073	7,956	26,552	53,680	64,451	46,552	12,995	121
1951	206,090	18	8	18	123	1,015	7,692	25,733	51,741	62,544	44,758	12,298	142
1950	197,338	22	10	11	103	943	7,389	25,317	50,238	60,087	41,876	11,213	129
<b>WHITE, FEMALE</b>													
1959	166,140	3	1	7	29	222	1,376	5,894	19,654	48,524	59,927	30,436	67
1958	160,438	2	3	8	34	232	1,370	5,836	19,564	47,310	57,655	28,373	51
1957	157,870	9	2	6	48	228	1,320	5,515	19,542	47,123	56,505	27,520	52
1956	147,827	21	5	19	43	215	1,245	5,266	18,770	44,264	53,093	24,836	50
1955	139,385	12	12	15	39	203	1,312	5,248	18,095	42,127	49,421	22,807	94
1954	128,894	21	5	13	43	231	1,251	5,308	17,652	39,736	44,844	19,742	48
1953	125,838	16	12	15	53	253	1,327	5,300	18,064	39,148	43,378	18,220	52
1952	118,257	14	7	17	63	260	1,287	5,425	17,565	36,986	40,224	16,339	70
1951	111,430	10	2	4	76	259	1,289	5,268	17,019	34,997	37,570	14,865	71
1950	105,795	12	3	13	72	278	1,283	5,244	16,497	33,813	35,066	13,448	66

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ARTERIOSCLEROTIC HEART DISEASE, INCLUDING CORONARY DISEASE (420)													
ALL OTHER, BOTH SEXES													
1959	30,329	4	4	5	48	338	1,600	4,502	7,731	8,754	5,226	2,079	38
1958	29,379	4	1	6	41	388	1,528	4,503	7,498	8,466	4,928	1,959	57
1957	28,885	11	4	11	57	360	1,679	4,429	7,569	8,103	4,686	1,917	59
1956	26,304	20	4	3	45	329	1,566	4,025	6,886	7,509	4,235	1,633	49
1955	24,199	17	4	7	67	352	1,428	4,040	6,177	6,722	3,912	1,417	56
1954	22,128	21	8	4	51	314	1,466	3,862	5,750	6,008	3,365	1,237	42
1953	22,343	16	4	10	77	379	1,477	4,092	5,731	6,044	3,258	1,215	40
1952	20,345	11	3	10	55	366	1,447	3,778	5,379	5,296	2,907	1,071	22
1951	19,099	7	2	4	70	317	1,462	3,627	5,010	4,966	2,659	944	31
1950	17,870	5	3	6	60	311	1,336	3,438	4,714	4,585	2,507	861	44
ALL OTHER, MALE													
1959	17,789	2	2	2	22	216	1,039	2,798	4,696	5,111	2,868	1,014	19
1958	17,051	2	-	3	23	219	938	2,748	4,467	4,945	2,756	919	31
1957	16,762	7	2	4	30	207	1,038	2,675	4,538	4,672	2,622	941	26
1956	15,174	10	3	1	26	184	952	2,465	4,095	4,276	2,351	785	26
1955	14,072	5	4	4	30	179	883	2,474	3,761	3,895	2,144	658	35
1954	13,060	9	6	3	23	178	860	2,350	3,563	3,530	1,924	594	20
1953	13,126	8	3	8	28	215	914	2,480	3,523	3,461	1,856	602	28
1952	11,991	6	2	7	24	207	853	2,278	3,321	3,151	1,621	510	11
1951	11,258	4	1	3	30	178	878	2,205	3,054	2,956	1,471	455	23
1950	10,526	4	2	4	38	177	786	2,081	2,866	2,711	1,420	410	27
ALL OTHER, FEMALE													
1959	12,540	2	2	3	26	122	561	1,704	3,035	3,643	2,358	1,065	19
1958	12,328	2	1	3	18	169	590	1,755	3,031	3,521	2,172	1,040	26
1957	12,123	4	2	7	27	153	641	1,754	3,031	3,431	2,064	976	33
1956	11,130	10	1	2	19	145	614	1,560	2,791	3,233	1,884	848	23
1955	10,127	12	-	3	37	173	545	1,566	2,416	2,827	1,768	759	21
1954	9,068	12	2	1	28	136	606	1,512	2,187	2,478	1,441	643	22
1953	9,217	8	1	2	49	164	563	1,612	2,208	2,583	1,402	613	12
1952	8,354	5	1	3	31	159	594	1,500	2,058	2,145	1,286	561	11
1951	7,841	3	1	1	40	139	584	1,422	1,956	2,010	1,188	489	8
1950	7,344	1	1	2	22	134	550	1,357	1,848	1,874	1,087	451	17

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)													
BOTH SEXES													
1959	56,099	43	31	41	61	227	620	1,933	5,040	12,626	21,108	14,340	29
1958	58,595	54	44	38	77	242	715	2,108	5,531	13,470	21,743	14,535	38
1957	62,618	63	69	50	101	258	770	2,442	6,248	14,902	22,709	14,979	27
1956	63,088	66	63	44	96	270	805	2,450	6,322	15,039	23,081	14,815	37
1955	65,622	71	54	39	97	293	947	2,745	6,596	15,537	24,182	15,015	46
1954	66,091	59	50	46	113	299	1,004	2,956	7,000	16,130	23,975	14,428	31
1953	73,330	60	49	51	119	377	1,128	3,280	8,095	18,031	26,546	15,552	42
1952	76,684	79	52	75	154	385	1,340	3,501	8,759	19,102	27,830	15,347	60
1951	80,635	46	42	52	157	442	1,392	3,849	9,491	20,049	29,016	16,042	57
1950	85,159	...	39	69	207	462	1,590	4,304	10,484	21,407	30,097	16,437	63
MALE													
1959	27,731	19	21	24	41	139	363	1,208	3,145	7,044	10,045	5,669	13
1958	28,834	28	23	24	44	149	441	1,306	3,314	7,488	10,312	5,686	19
1957	31,387	30	30	26	66	142	464	1,502	3,851	8,356	10,934	5,971	15
1956	31,857	35	32	28	55	161	501	1,540	3,818	8,488	11,288	5,887	24
1955	33,171	34	29	28	55	162	573	1,689	4,019	8,796	11,719	6,049	18
1954	33,825	34	29	24	67	176	585	1,773	4,323	9,122	11,811	5,863	18
1953	37,637	36	24	26	72	202	661	2,088	5,046	10,150	12,882	6,427	23
1952	39,657	41	32	41	84	194	768	2,115	5,404	10,908	13,786	6,249	35
1951	42,088	25	19	27	89	239	786	2,326	5,866	11,537	14,374	6,774	26
1950	44,447	...	23	31	92	226	892	2,696	6,517	12,196	14,860	6,871	43
FEMALE													
1959	28,368	24	10	17	20	88	257	725	1,895	5,582	11,063	8,671	16
1958	29,761	26	21	14	33	93	274	802	2,217	5,982	11,431	8,849	19
1957	31,231	33	39	24	35	116	306	940	2,397	6,546	11,775	9,008	12
1956	31,231	31	31	16	41	109	304	910	2,504	6,551	11,793	8,928	13
1955	32,451	37	25	11	42	131	374	1,056	2,577	6,741	12,463	8,966	28
1954	32,266	25	21	22	46	123	419	1,183	2,677	7,008	12,164	8,565	13
1953	35,693	24	25	25	47	175	467	1,192	3,049	7,881	13,664	9,125	19
1952	37,027	38	20	34	70	191	572	1,386	3,355	8,194	14,044	9,098	25
1951	38,547	21	23	25	68	203	606	1,523	3,625	8,512	14,642	9,268	31
1950	40,712	...	16	38	115	236	698	1,608	3,967	9,211	15,237	9,566	20

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)													
WHITE, BOTH SEXES													
1959	50,159	28	22	34	45	114	391	1,374	3,851	10,873	19,802	13,607	18
1958	52,071	41	29	29	62	143	451	1,457	4,159	11,567	20,359	13,752	22
1957	55,661	45	46	33	70	156	478	1,653	4,777	12,791	21,306	14,288	18
1956	56,260	53	28	33	68	159	501	1,667	4,892	13,073	21,643	14,113	30
1955	58,792	56	36	31	77	173	600	1,858	5,161	13,607	22,798	14,359	36
1954	59,013	44	36	35	86	180	628	1,989	5,510	14,134	22,599	13,755	17
1953	65,654	48	32	40	79	227	709	2,265	6,453	15,851	25,077	14,842	31
1952	68,522	62	42	56	99	226	852	2,403	6,958	16,868	26,267	14,644	45
1951	72,079	35	36	39	100	270	895	2,637	7,535	17,758	27,415	15,319	40
1950	76,288	...	26	56	143	282	1,027	3,042	8,463	19,060	28,473	15,671	45
WHITE, MALE													
1959	24,646	12	16	21	33	76	250	919	2,494	6,096	9,380	5,340	9
1958	25,431	21	19	21	34	100	301	956	2,577	6,459	9,599	5,335	9
1957	27,750	21	21	14	51	87	316	1,064	3,025	7,245	10,241	5,656	9
1956	28,268	29	17	21	40	101	341	1,111	3,063	7,436	10,518	5,573	18
1955	29,563	27	22	23	45	106	389	1,246	3,229	7,752	10,961	5,749	14
1954	30,085	28	22	15	56	114	402	1,290	3,468	8,062	11,048	5,569	11
1953	33,433	29	13	21	55	120	459	1,500	4,070	8,957	12,070	6,123	16
1952	35,303	32	22	30	57	132	532	1,555	4,398	9,662	12,918	5,939	26
1951	37,456	18	17	20	54	155	541	1,663	4,764	10,248	13,496	6,459	21
1950	39,747	...	14	26	68	135	637	1,989	5,401	10,934	13,982	6,530	31
WHITE, FEMALE													
1959	25,513	16	6	13	12	38	141	455	1,357	4,777	10,422	8,267	9
1958	26,640	20	10	8	28	43	150	501	1,582	5,108	10,760	8,417	13
1957	27,911	24	25	19	19	69	162	589	1,752	5,546	11,065	8,632	9
1956	27,992	24	11	12	28	58	160	556	1,829	5,637	11,125	8,540	12
1955	29,229	29	14	8	32	67	211	612	1,932	5,855	11,837	8,610	22
1954	28,928	16	14	20	30	66	226	699	2,042	6,072	11,551	8,186	6
1953	32,221	19	19	19	24	107	250	765	2,383	6,894	13,007	8,719	15
1952	33,219	30	20	26	42	94	320	848	2,560	7,206	13,349	8,705	19
1951	34,623	17	19	19	46	115	354	974	2,771	7,510	13,919	8,860	19
1950	36,541	...	12	30	75	147	390	1,053	3,062	8,126	14,491	9,141	14

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
NONRHEUMATIC CHRONIC ENDOCARDITIS AND OTHER MYOCARDIAL DEGENERATION (421, 422)													
ALL OTHER, BOTH SEXES													
1959	5,940	15	9	7	16	113	229	559	1,189	1,753	1,306	733	11
1958	6,524	13	15	9	15	99	264	651	1,372	1,903	1,384	783	16
1957	6,957	18	23	17	31	102	292	789	1,471	2,111	1,403	691	9
1956	6,828	13	35	11	28	111	304	783	1,430	1,966	1,438	702	7
1955	6,830	15	18	8	20	120	347	887	1,435	1,930	1,384	656	10
1954	7,078	15	14	11	27	119	376	967	1,490	1,996	1,376	673	14
1953	7,676	12	17	11	40	150	419	1,015	1,642	2,180	1,469	710	11
1952	8,162	17	10	19	55	159	488	1,098	1,801	2,234	1,563	703	15
1951	8,556	11	6	13	57	172	497	1,212	1,956	2,291	1,601	723	17
1950	8,871	...	13	13	64	180	563	1,262	2,021	2,347	1,624	766	18
ALL OTHER, MALE													
1959	3,085	7	5	3	8	63	113	289	651	948	665	329	4
1958	3,403	7	4	3	10	49	140	350	737	1,029	713	351	10
1957	3,637	9	9	12	15	55	148	438	826	1,111	693	315	6
1956	3,589	6	15	7	15	60	160	429	755	1,052	770	314	6
1955	3,608	7	7	5	10	56	184	443	790	1,044	758	300	4
1954	3,740	6	7	9	11	62	183	483	855	1,060	763	294	7
1953	4,204	7	11	5	17	82	202	588	976	1,193	812	304	7
1952	4,354	9	10	11	27	62	236	560	1,006	1,246	868	310	9
1951	4,632	7	2	7	35	84	245	663	1,102	1,289	878	315	5
1950	4,700	...	9	5	24	91	255	707	1,116	1,262	878	341	12
ALL OTHER, FEMALE													
1959	2,855	8	4	4	8	50	116	270	538	805	641	404	7
1958	3,121	6	11	6	5	50	124	301	635	874	671	432	6
1957	3,320	9	14	5	16	47	144	351	645	1,000	710	376	3
1956	3,239	7	20	4	13	51	144	354	675	914	668	388	1
1955	3,222	8	11	3	10	64	163	444	645	886	626	356	6
1954	3,338	9	7	2	16	57	193	484	635	936	613	379	7
1953	3,472	5	6	6	23	68	217	427	666	987	657	406	4
1952	3,808	8	-	8	28	97	252	538	795	988	695	393	6
1951	3,924	4	4	6	22	88	252	549	854	1,002	723	408	12
1950	4,171	...	4	8	40	89	308	555	905	1,085	746	425	6

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES OF HEART (430-434)													
BOTH SEXES													
1959	24,353	224	161	107	208	472	1,104	2,461	3,996	6,187	6,307	3,113	13
1958	24,548	223	128	113	210	427	1,070	2,379	4,150	6,302	6,351	3,170	25
1957	21,733	191	124	121	206	418	973	2,188	3,855	5,750	5,372	2,513	22
1956	21,002	166	93	91	173	409	892	2,083	3,807	5,706	5,230	2,337	15
1955	20,659	141	114	104	176	379	968	2,153	3,723	5,576	5,121	2,180	24
1954	20,903	161	82	88	223	424	1,025	2,281	3,897	5,563	4,991	2,151	17
1953	22,612	189	78	105	211	466	1,224	2,469	4,342	5,882	5,347	2,274	25
1952	22,657	169	99	82	250	495	1,259	2,611	4,312	5,949	5,291	2,122	18
1951	23,176	169	85	76	261	504	1,306	2,655	4,381	6,089	5,439	2,186	25
1950	23,958	70	90	128	271	581	1,335	2,769	4,692	6,207	5,550	2,229	36
MALE													
1959	14,226	122	91	62	118	238	700	1,680	2,670	3,858	3,321	1,357	9
1958	14,386	122	64	64	110	210	675	1,622	2,783	3,878	3,423	1,418	17
1957	12,942	109	67	66	121	224	627	1,458	2,611	3,627	2,909	1,109	14
1956	12,536	88	49	54	93	211	515	1,416	2,579	3,559	2,902	1,060	10
1955	12,566	79	58	56	97	217	613	1,461	2,518	3,653	2,817	977	20
1954	12,637	91	43	50	131	245	639	1,560	2,686	3,479	2,677	1,025	11
1953	13,803	107	40	55	118	255	769	1,722	2,988	3,796	2,918	1,020	15
1952	13,846	98	59	48	134	261	768	1,793	3,017	3,805	2,887	965	11
1951	14,019	96	38	41	141	266	786	1,795	2,959	3,851	3,038	995	13
1950	14,469	38	46	66	147	298	834	1,834	3,152	3,921	3,040	1,070	23
FEMALE													
1959	10,127	102	70	45	90	234	404	781	1,326	2,329	2,986	1,756	4
1958	10,162	101	64	49	100	217	395	757	1,367	2,424	2,928	1,752	8
1957	8,791	82	57	55	85	194	346	730	1,244	2,123	2,463	1,404	8
1956	8,466	78	44	37	80	198	377	667	1,228	2,147	2,328	1,277	5
1955	8,093	62	56	48	79	162	355	692	1,205	1,923	2,304	1,203	4
1954	8,266	70	39	38	92	179	386	721	1,211	2,084	2,314	1,126	6
1953	8,809	82	38	50	93	211	455	747	1,354	2,086	2,429	1,254	10
1952	8,811	71	40	34	116	234	491	818	1,295	2,144	2,404	1,157	7
1951	9,157	73	47	35	120	238	520	860	1,422	2,238	2,401	1,191	12
1950	9,489	32	44	62	124	283	501	935	1,540	2,286	2,510	1,159	13

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES OF HEART (430-434)													
WHITE, BOTH SEXES													
1959	20,477	156	98	85	149	301	777	1,824	3,149	5,256	5,772	2,903	7
1958	20,483	155	90	93	136	268	724	1,700	3,183	5,328	5,821	2,973	12
1957	17,895	146	76	87	140	271	643	1,555	2,955	4,781	4,910	2,319	12
1956	17,451	117	62	77	116	264	591	1,439	2,992	4,834	4,794	2,155	10
1955	17,096	102	73	77	126	246	650	1,512	2,928	4,724	4,642	1,998	18
1954	17,393	115	54	74	145	268	681	1,601	3,134	4,768	4,559	1,988	6
1953	18,686	142	59	76	136	295	819	1,751	3,428	5,024	4,877	2,064	15
1952	18,592	125	70	64	157	300	824	1,797	3,403	5,086	4,835	1,922	9
1951	19,007	137	65	52	162	318	816	1,815	3,464	5,209	4,953	1,997	19
1950	19,761	59	76	101	176	370	877	1,927	3,728	5,299	5,075	2,051	22
WHITE, MALE													
1959	12,029	96	51	48	86	160	521	1,296	2,185	3,303	3,028	1,250	5
1958	12,004	90	44	52	77	138	481	1,215	2,204	3,290	3,094	1,311	8
1957	10,756	82	39	49	86	155	439	1,091	2,078	3,070	2,649	1,012	6
1956	10,516	61	35	47	62	154	363	1,040	2,090	3,065	2,636	958	5
1955	10,506	61	36	40	74	146	445	1,089	2,060	3,102	2,559	880	14
1954	10,630	61	29	42	95	165	460	1,146	2,227	3,024	2,441	936	4
1953	11,516	86	29	38	81	175	544	1,289	2,415	3,276	2,654	919	10
1952	11,490	75	43	36	92	167	555	1,317	2,434	3,295	2,608	861	7
1951	11,696	78	28	33	96	181	544	1,313	2,409	3,335	2,762	906	11
1950	12,137	35	37	55	106	216	587	1,375	2,590	3,375	2,770	976	15
WHITE, FEMALE													
1959	8,448	60	47	37	63	141	256	528	964	1,953	2,744	1,653	2
1958	8,479	65	46	41	59	130	243	485	979	2,038	2,727	1,662	4
1957	7,139	64	37	38	54	116	204	464	877	1,711	2,261	1,307	6
1956	6,935	56	27	30	54	110	228	399	902	1,769	2,158	1,197	5
1955	6,590	41	37	37	52	100	205	423	868	1,622	2,083	1,118	4
1954	6,763	54	25	32	50	103	221	455	907	1,744	2,118	1,052	2
1953	7,170	56	30	38	55	120	275	462	1,013	1,748	2,223	1,145	5
1952	7,102	50	27	28	65	133	269	480	969	1,791	2,227	1,061	2
1951	7,311	59	37	19	66	137	272	502	1,055	1,874	2,191	1,091	8
1950	7,624	24	39	46	70	154	290	552	1,138	1,924	2,305	1,075	7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES OF HEART (430-434)													
ALL OTHER, BOTH SEXES													
1959	3,876	68	63	22	59	171	327	637	847	931	535	210	6
1958	4,065	68	38	20	74	159	346	679	967	974	530	197	13
1957	3,838	45	48	34	66	147	330	633	900	969	462	194	10
1956	3,551	49	31	14	57	145	301	644	815	872	436	182	5
1955	3,563	39	41	27	50	133	318	641	795	852	479	182	6
1954	3,510	46	28	14	78	156	344	680	763	795	432	163	11
1953	3,926	47	19	29	75	171	405	718	914	858	470	210	10
1952	4,065	44	29	18	93	195	435	814	909	863	456	200	9
1951	4,169	32	20	24	99	186	490	840	917	880	486	189	6
1950	4,197	11	14	27	95	211	458	842	964	908	475	178	14
ALL OTHER, MALE													
1959	2,197	26	40	14	32	78	179	384	485	555	293	107	4
1958	2,382	32	20	12	33	72	194	407	579	588	329	107	9
1957	2,186	27	28	17	35	69	188	367	533	557	260	97	8
1956	2,020	27	14	7	31	57	152	376	489	494	266	102	5
1955	2,060	18	22	16	23	71	168	372	458	551	258	97	6
1954	2,007	30	14	8	36	80	179	414	459	455	236	89	7
1953	2,287	21	11	17	37	80	225	433	573	520	264	101	5
1952	2,356	23	16	12	42	94	213	476	583	510	279	104	4
1951	2,323	18	10	8	45	85	242	482	550	516	276	89	2
1950	2,332	3	9	11	41	82	247	459	562	546	270	94	8
ALL OTHER, FEMALE													
1959	1,679	42	23	8	27	93	148	253	362	376	242	103	2
1958	1,683	36	18	8	41	87	152	272	388	386	201	90	4
1957	1,652	18	20	17	31	78	142	266	367	412	202	97	2
1956	1,531	22	17	7	26	88	149	268	326	378	170	80	-
1955	1,503	21	19	11	27	62	150	269	337	301	221	85	-
1954	1,503	16	14	6	42	76	165	266	304	340	196	74	4
1953	1,639	26	8	12	38	91	180	285	341	338	206	109	5
1952	1,709	21	13	6	51	101	222	338	326	353	177	96	5
1951	1,846	14	10	16	54	101	248	358	367	364	210	100	4
1950	1,865	8	5	16	54	129	211	383	402	362	205	84	6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HYPERTENSIVE HEART DISEASE (440-443)													
BOTH SEXES													
1959	68,159	2	-	4	36	299	1,523	4,895	10,654	19,832	21,469	9,401	44
1958	73,934	1	1	3	42	307	1,625	5,421	11,950	21,707	23,118	9,718	41
1957	72,372	-	1	10	40	342	1,871	5,712	12,367	21,298	21,643	9,028	60
1956	72,393	-	1	6	38	347	1,718	5,653	12,342	21,121	22,228	8,883	56
1955	73,883	1	2	10	38	344	1,911	5,730	12,847	21,803	22,468	8,655	74
1954	75,039	5	3	7	51	450	1,993	6,413	13,418	22,562	21,946	8,135	56
1953	81,149	-	2	10	60	445	2,435	7,412	15,284	23,847	23,441	8,164	49
1952	83,911	1	4	8	74	504	2,581	7,940	15,917	24,881	23,805	8,127	69
1951	85,257	-	-	9	75	574	2,653	8,199	16,307	24,842	24,258	8,266	74
1950	85,193	-	2	10	90	541	2,694	8,385	16,525	25,260	23,783	7,826	77
MALE													
1959	29,717	1	-	2	18	144	732	2,554	5,526	9,089	8,432	3,200	19
1958	32,055	1	-	2	21	150	807	2,711	5,943	10,067	9,044	3,295	14
1957	32,140	-	-	5	24	165	907	2,859	6,300	10,027	8,592	3,232	29
1956	32,374	-	-	4	19	176	812	2,881	6,307	9,963	9,030	3,161	21
1955	33,342	-	-	5	22	180	951	2,947	6,655	10,160	9,250	3,140	32
1954	33,954	4	1	4	21	203	947	3,221	6,992	10,480	9,011	3,044	26
1953	37,368	-	2	6	30	222	1,161	3,717	8,110	11,310	9,736	3,049	25
1952	39,005	1	1	4	37	241	1,206	4,013	8,318	11,989	10,071	3,094	30
1951	39,622	-	-	4	28	254	1,232	3,989	8,551	11,855	10,478	3,193	38
1950	39,766	-	1	5	42	232	1,196	4,153	8,586	12,176	10,329	3,007	39
FEMALE													
1959	38,442	1	-	2	18	155	791	2,341	5,128	10,743	13,037	6,201	25
1958	41,879	-	1	1	21	157	818	2,710	6,007	11,640	14,074	6,423	27
1957	40,232	-	1	5	16	177	964	2,853	6,067	11,271	13,051	5,796	31
1956	40,019	-	1	2	19	171	906	2,772	6,035	11,158	13,198	5,722	35
1955	40,541	1	2	5	16	164	960	2,783	6,192	11,643	13,218	5,515	42
1954	41,085	1	2	3	30	247	1,046	3,192	6,426	12,082	12,935	5,091	30
1953	43,781	-	-	4	30	223	1,274	3,695	7,174	12,537	13,705	5,115	24
1952	44,906	-	3	4	37	263	1,375	3,927	7,599	12,892	13,734	5,033	39
1951	45,635	-	-	5	47	320	1,421	4,210	7,756	12,987	13,780	5,073	36
1950	45,427	-	1	5	48	309	1,498	4,232	7,939	13,084	13,454	4,819	38

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HYPERTENSIVE HEART DISEASE (440-443)													
WHITE, BOTH SEXES													
1959	53,937	-	-	4	15	95	539	2,532	7,060	15,857	19,270	8,548	17
1958	58,791	1	-	3	26	96	647	2,784	8,025	17,492	20,814	8,887	16
1957	57,072	-	1	6	19	116	721	3,073	8,222	17,210	19,441	8,243	20
1956	57,565	-	1	5	21	120	672	3,071	8,283	17,174	20,052	8,138	28
1955	59,233	-	2	9	19	124	799	3,113	9,011	17,837	20,333	7,942	44
1954	60,444	2	2	5	20	193	857	3,608	9,645	18,661	19,955	7,466	30
1953	65,509	-	-	6	30	194	1,088	4,380	11,153	19,917	21,295	7,422	24
1952	67,675	1	3	4	38	221	1,244	4,707	11,668	20,631	21,666	7,447	45
1951	69,264	-	-	4	46	256	1,274	5,011	12,190	20,775	22,123	7,547	38
1950	69,357	-	-	9	48	241	1,303	5,135	12,371	21,280	21,787	7,141	42
WHITE, MALE													
1959	23,208	-	-	2	8	56	300	1,444	3,827	7,271	7,430	2,864	6
1958	25,239	1	-	2	14	51	377	1,556	4,206	8,089	7,993	2,945	5
1957	25,086	-	-	2	11	61	408	1,722	4,369	8,075	7,561	2,870	7
1956	25,449	-	-	3	11	65	370	1,688	4,411	8,056	8,010	2,826	9
1955	26,354	-	-	4	12	77	443	1,696	4,776	8,291	8,203	2,835	17
1954	27,109	2	1	3	10	103	456	1,930	5,117	8,654	8,064	2,755	14
1953	29,875	-	-	4	16	97	587	2,317	6,049	9,364	8,714	2,718	9
1952	31,140	1	1	1	20	115	635	2,535	6,187	9,856	8,983	2,790	16
1951	32,029	-	-	2	21	126	661	2,593	6,484	9,844	9,398	2,881	19
1950	32,162	-	-	4	25	118	627	2,651	6,540	10,177	9,295	2,706	19
WHITE, FEMALE													
1959	30,729	-	-	2	7	39	239	1,088	3,233	8,586	11,840	5,684	11
1958	33,552	-	-	1	12	45	270	1,228	3,819	9,403	12,821	5,942	11
1957	31,986	-	1	4	8	55	313	1,351	3,853	9,135	11,880	5,373	13
1956	32,116	-	1	2	10	55	302	1,383	3,872	9,118	12,042	5,312	19
1955	32,879	-	2	5	7	47	356	1,417	4,235	9,546	12,130	5,107	27
1954	33,335	-	1	2	10	90	401	1,678	4,528	10,007	11,891	4,711	16
1953	35,634	-	-	2	14	97	501	2,063	5,104	10,553	12,581	4,704	15
1952	36,535	-	2	3	18	106	609	2,172	5,481	10,775	12,683	4,657	29
1951	37,235	-	-	2	25	130	613	2,418	5,706	10,931	12,725	4,666	19
1950	37,195	-	-	5	23	123	676	2,484	5,831	11,103	12,492	4,435	23

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HYPERTENSIVE HEART DISEASE (440-443)													
ALL OTHER, BOTH SEXES													
1959	14,222	2	-	-	21	204	984	2,363	3,594	3,975	2,199	853	27
1958	15,143	-	1	-	16	211	978	2,637	3,925	4,215	2,304	831	25
1957	15,300	-	-	4	21	226	1,150	2,639	4,145	4,088	2,202	785	40
1956	14,828	-	-	1	17	227	1,046	2,582	4,059	3,947	2,176	745	28
1955	14,650	1	-	1	19	220	1,112	2,617	3,836	3,966	2,135	713	30
1954	14,595	3	1	2	31	257	1,136	2,805	3,773	3,901	1,991	669	26
1953	15,640	-	2	4	30	251	1,347	3,032	4,131	3,930	2,146	742	25
1952	16,236	-	1	4	36	283	1,337	3,233	4,249	4,250	2,139	680	24
1951	15,993	-	-	5	29	318	1,379	3,188	4,117	4,067	2,135	719	36
1950	15,836	-	2	1	42	300	1,391	3,250	4,154	3,980	1,996	685	35
ALL OTHER, MALE													
1959	6,509	1	-	-	10	88	432	1,110	1,699	1,818	1,002	336	13
1958	6,816	-	-	-	7	99	430	1,155	1,737	1,978	1,051	350	9
1957	7,054	-	-	3	13	104	499	1,137	1,931	1,952	1,031	362	22
1956	6,925	-	-	1	8	111	442	1,193	1,896	1,907	1,020	335	12
1955	6,988	-	-	1	10	103	508	1,251	1,879	1,869	1,047	305	15
1954	6,845	2	-	1	11	100	491	1,291	1,875	1,826	947	289	12
1953	7,493	-	2	2	14	125	574	1,400	2,061	1,946	1,022	331	16
1952	7,865	-	-	3	17	126	571	1,478	2,131	2,133	1,088	304	14
1951	7,593	-	-	2	7	128	571	1,396	2,067	2,011	1,080	312	19
1950	7,604	-	1	1	17	114	569	1,502	2,046	1,999	1,034	301	20
ALL OTHER, FEMALE													
1959	7,713	1	-	-	11	116	552	1,253	1,895	2,157	1,197	517	14
1958	8,327	-	1	-	9	112	548	1,482	2,188	2,237	1,253	481	16
1957	8,246	-	-	1	8	122	651	1,502	2,214	2,136	1,171	423	18
1956	7,903	-	-	-	9	116	604	1,389	2,163	2,040	1,156	410	16
1955	7,662	1	-	-	9	117	604	1,366	1,957	2,097	1,088	408	15
1954	7,750	1	1	1	20	157	645	1,514	1,898	2,075	1,044	380	14
1953	8,147	-	-	2	16	126	773	1,632	2,070	1,984	1,124	411	9
1952	8,371	-	1	1	19	157	766	1,755	2,118	2,117	1,051	376	10
1951	8,400	-	-	3	22	190	808	1,792	2,050	2,056	1,055	407	17
1950	8,232	-	1	-	25	186	822	1,748	2,108	1,981	962	384	15

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER HYPERTENSIVE DISEASE (444-447)													
BOTH SEXES													
1959	13,119	1	-	4	47	285	738	1,184	1,703	2,939	4,042	2,167	9
1958	13,798	-	2	4	48	263	789	1,395	1,851	3,242	4,072	2,124	8
1957	11,128	-	2	4	30	220	655	1,024	1,398	2,640	3,351	1,798	6
1956	10,948	-	-	3	28	211	637	984	1,475	2,566	3,252	1,787	5
1955	11,162	-	1	2	37	233	696	995	1,497	2,580	3,376	1,740	5
1954	11,383	-	1	4	39	308	724	1,161	1,624	2,644	3,276	1,591	11
1953	12,387	-	-	1	41	306	885	1,391	1,837	2,834	3,425	1,661	6
1952	12,506	-	2	7	54	339	899	1,407	1,847	2,876	3,421	1,647	7
1951	13,256	-	-	6	57	339	909	1,433	2,009	3,018	3,777	1,693	15
1950	12,563	-	1	4	50	314	875	1,337	1,848	2,912	3,592	1,618	12
MALE													
1959	6,678	-	-	2	31	149	414	682	938	1,605	1,948	903	6
1958	6,959	-	1	3	23	131	404	783	1,031	1,668	1,998	914	3
1957	5,568	-	1	1	14	111	348	595	756	1,401	1,573	763	5
1956	5,558	-	-	1	12	109	355	592	818	1,341	1,532	795	3
1955	5,664	-	-	1	25	120	373	591	808	1,349	1,589	805	3
1954	5,731	-	1	2	20	143	375	658	887	1,374	1,555	709	7
1953	6,374	-	-	-	19	151	486	830	1,038	1,449	1,687	710	4
1952	6,280	-	2	3	27	145	460	802	1,039	1,461	1,631	706	4
1951	6,592	-	-	3	28	159	451	757	1,117	1,563	1,790	714	10
1950	6,381	-	-	2	25	152	447	765	1,034	1,544	1,699	706	7
FEMALE													
1959	6,441	1	-	2	16	136	324	502	765	1,334	2,094	1,264	3
1958	6,839	-	1	1	25	132	385	612	820	1,574	2,074	1,210	5
1957	5,560	-	1	3	16	109	307	429	642	1,239	1,778	1,035	1
1956	5,390	-	-	2	16	102	282	392	657	1,225	1,720	992	2
1955	5,498	-	1	1	12	113	323	404	689	1,231	1,787	935	2
1954	5,652	-	-	2	19	165	349	503	737	1,270	1,721	882	4
1953	6,013	-	-	1	22	155	399	561	799	1,385	1,738	951	2
1952	6,226	-	-	4	27	194	439	605	808	1,415	1,790	941	3
1951	6,664	-	-	3	29	180	458	676	892	1,455	1,987	979	5
1950	6,182	-	1	2	25	162	428	572	814	1,368	1,893	912	5

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER HYPERTENSIVE DISEASE (444-447)													
WHITE, BOTH SEXES													
1959	10,536	1	-	4	25	114	289	678	1,194	2,441	3,750	2,038	2
1958	11,117	-	1	4	33	93	335	817	1,374	2,728	3,745	1,983	4
1957	8,764	-	1	3	18	70	244	556	958	2,164	3,065	1,682	3
1956	8,676	-	-	3	14	77	245	558	1,035	2,117	2,957	1,668	2
1955	8,823	-	1	2	13	86	267	539	1,092	2,093	3,104	1,622	4
1954	8,985	-	1	4	20	122	306	652	1,232	2,186	2,998	1,458	6
1953	9,808	-	-	1	25	128	427	834	1,375	2,344	3,134	1,538	2
1952	9,982	-	2	5	29	139	465	902	1,380	2,404	3,127	1,525	4
1951	10,690	-	-	6	33	159	473	938	1,526	2,525	3,460	1,561	9
1950	10,252	-	1	4	33	158	463	864	1,394	2,489	3,323	1,518	5
WHITE, MALE													
1959	5,301	-	-	2	17	59	172	414	682	1,315	1,791	847	2
1958	5,591	-	1	3	18	53	187	495	770	1,403	1,822	838	1
1957	4,372	-	-	1	10	38	157	337	532	1,148	1,438	708	3
1956	4,384	-	-	1	6	41	154	358	576	1,107	1,388	751	2
1955	4,463	-	-	1	9	54	159	337	618	1,088	1,452	742	3
1954	4,568	-	1	2	12	68	177	409	681	1,148	1,418	648	4
1953	5,026	-	-	-	9	73	254	534	793	1,191	1,515	655	2
1952	5,030	-	2	1	19	64	255	545	791	1,234	1,472	644	3
1951	5,330	-	-	3	18	84	256	515	860	1,309	1,625	653	7
1950	5,244	-	-	2	17	80	258	526	823	1,314	1,555	667	2
WHITE, FEMALE													
1959	5,235	1	-	2	8	55	117	264	512	1,126	1,959	1,191	-
1958	5,526	-	-	1	15	40	148	322	604	1,325	1,923	1,145	3
1957	4,392	-	1	2	8	32	87	219	426	1,016	1,627	974	-
1956	4,292	-	-	2	8	36	91	200	459	1,010	1,569	917	-
1955	4,360	-	1	1	4	32	108	202	474	1,005	1,652	880	1
1954	4,417	-	-	2	8	54	129	243	551	1,038	1,580	810	2
1953	4,782	-	-	1	16	55	173	300	582	1,153	1,619	883	-
1952	4,952	-	-	4	10	75	210	357	589	1,170	1,655	881	1
1951	5,360	-	-	3	15	75	217	423	666	1,216	1,835	908	2
1950	5,008	-	1	2	16	78	205	338	571	1,175	1,768	851	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER HYPERTENSIVE DISEASE (444-447)													
ALL OTHER, BOTH SEXES													
1959	2,583	-	-	-	22	171	449	506	509	498	292	129	7
1958	2,681	-	1	-	15	170	454	578	477	514	327	141	4
1957	2,364	-	1	1	12	150	411	468	440	476	286	116	3
1956	2,272	-	-	-	14	134	392	426	440	449	295	119	3
1955	2,339	-	-	-	24	147	429	456	405	487	272	118	1
1954	2,398	-	-	-	19	186	418	509	392	458	278	133	5
1953	2,579	-	-	-	16	178	458	557	462	490	291	123	4
1952	2,524	-	-	2	25	200	434	505	467	472	294	122	3
1951	2,566	-	-	-	24	180	436	495	483	493	317	132	6
1950	2,311	-	-	-	17	156	412	473	454	423	269	100	7
ALL OTHER, MALE													
1959	1,377	-	-	-	14	90	242	268	256	290	157	56	4
1958	1,368	-	-	-	5	78	217	288	261	265	176	76	2
1957	1,196	-	1	-	4	73	191	258	224	253	135	55	2
1956	1,174	-	-	-	6	68	201	234	242	234	144	44	1
1955	1,201	-	-	-	16	66	214	254	190	261	137	63	-
1954	1,163	-	-	-	8	75	198	249	206	226	137	61	3
1953	1,348	-	-	-	10	78	232	296	245	258	172	55	2
1952	1,250	-	-	2	8	81	205	257	248	227	159	62	1
1951	1,262	-	-	-	10	75	195	242	257	254	165	61	3
1950	1,137	-	-	-	8	72	189	239	211	230	144	39	5
ALL OTHER, FEMALE													
1959	1,206	-	-	-	8	81	207	238	253	208	135	73	3
1958	1,313	-	1	-	10	92	237	290	216	249	151	65	2
1957	1,168	-	-	1	8	77	220	210	216	223	151	61	1
1956	1,098	-	-	-	8	66	191	192	198	215	151	75	2
1955	1,138	-	-	-	8	81	215	202	215	226	135	55	1
1954	1,235	-	-	-	11	111	220	260	186	232	141	72	2
1953	1,231	-	-	-	6	100	226	261	217	232	119	68	2
1952	1,274	-	-	-	17	119	229	248	219	245	135	60	2
1951	1,304	-	-	-	14	105	241	253	226	239	152	71	3
1950	1,174	-	-	-	9	84	223	234	243	193	125	61	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
GENERAL ARTERIOSCLEROSIS (450)													
BOTH SEXES													
1959	34,622	1	-	-	4	9	60	422	1,659	5,922	13,862	12,666	17
1958	34,483	-	1	1	4	15	65	342	1,570	6,190	13,678	12,605	12
1957	33,272	-	-	-	4	9	42	371	1,556	6,059	13,177	12,040	14
1956	32,018	-	1	1	6	11	56	353	1,447	5,686	12,994	11,448	15
1955	32,486	-	1	-	4	12	63	342	1,469	5,744	13,388	11,439	24
1954	30,225	1	1	-	3	15	67	343	1,479	5,478	12,292	10,527	19
1953	32,351	-	-	-	4	29	63	334	1,574	6,089	13,215	11,017	26
1952	31,685	-	-	1	7	18	66	373	1,649	5,959	13,134	10,452	26
1951	31,884	-	1	-	-	14	58	365	1,638	6,133	13,398	10,250	27
1950	30,734	-	-	-	2	13	59	341	1,640	5,836	12,848	9,967	28
MALE													
1959	16,460	1	-	-	3	6	43	283	1,037	3,340	6,713	5,021	13
1958	16,588	-	1	1	2	9	36	214	958	3,501	6,739	5,119	8
1957	16,044	-	-	-	3	6	28	228	956	3,496	6,394	4,928	5
1956	15,557	-	1	-	3	5	31	231	916	3,226	6,425	4,717	2
1955	16,094	-	1	-	3	5	39	245	918	3,327	6,748	4,791	17
1954	14,801	1	1	-	1	8	36	222	907	3,125	6,177	4,317	6
1953	16,094	-	-	-	3	16	39	207	992	3,541	6,686	4,593	17
1952	15,640	-	-	1	3	7	44	248	1,037	3,392	6,504	4,392	12
1951	15,980	-	-	-	-	7	26	237	1,012	3,548	6,715	4,423	12
1950	15,676	-	-	-	-	8	33	216	1,049	3,382	6,578	4,396	14
FEMALE													
1959	18,162	-	-	-	1	3	17	139	622	2,582	7,149	7,645	4
1958	17,895	-	-	-	2	6	29	128	612	2,689	6,939	7,486	4
1957	17,228	-	-	-	1	3	14	143	600	2,563	6,783	7,112	9
1956	16,461	-	-	1	3	6	25	122	531	2,460	6,569	6,731	13
1955	16,392	-	-	-	1	7	24	97	551	2,417	6,640	6,648	7
1954	15,424	-	-	-	2	7	31	121	572	2,353	6,115	6,210	13
1953	16,257	-	-	-	1	13	24	127	582	2,548	6,529	6,424	9
1952	16,045	-	-	-	4	11	22	125	612	2,567	6,630	6,060	14
1951	15,904	-	1	-	-	7	32	128	626	2,585	6,683	5,827	15
1950	15,058	-	-	-	2	5	26	125	591	2,454	6,270	5,571	14

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
GENERAL ARTERIOSCLEROSIS (450)													
WHITE, BOTH SEXES													
1959	32,107	1	-	-	4	4	39	309	1,317	5,184	13,079	12,161	9
1958	31,863	-	-	1	2	10	42	250	1,229	5,401	12,864	12,057	7
1957	30,845	-	-	-	3	6	28	278	1,223	5,313	12,469	11,515	10
1956	29,763	-	1	1	4	9	36	262	1,113	4,975	12,357	10,997	8
1955	30,193	-	1	-	2	9	35	242	1,132	5,087	12,698	10,973	14
1954	28,076	1	1	-	3	11	39	236	1,154	4,856	11,674	10,090	11
1953	30,066	-	-	-	3	20	44	218	1,203	5,410	12,553	10,596	19
1952	29,538	-	-	1	6	12	35	263	1,309	5,327	12,535	10,028	22
1951	29,821	-	1	-	-	10	35	242	1,294	5,512	12,827	9,875	25
1950	28,713	-	-	-	1	6	36	229	1,313	5,218	12,304	9,583	23
WHITE, MALE													
1959	15,156	1	-	-	3	3	31	218	865	2,921	6,300	4,808	6
1958	15,264	-	-	1	1	6	26	162	794	3,086	6,317	4,865	6
1957	14,822	-	-	-	2	4	22	188	784	3,090	6,033	4,694	5
1956	14,408	-	1	-	2	4	19	186	725	2,854	6,101	4,514	2
1955	14,867	-	1	-	1	4	24	177	733	2,928	6,402	4,586	11
1954	13,681	1	1	-	1	6	23	164	736	2,786	5,838	4,122	3
1953	14,859	-	-	-	2	11	27	140	779	3,149	6,339	4,399	13
1952	14,487	-	-	1	2	4	26	193	839	3,036	6,169	4,207	10
1951	14,864	-	-	-	-	4	19	159	823	3,199	6,388	4,261	11
1950	14,567	-	-	-	-	6	24	158	856	3,024	6,260	4,227	12
WHITE, FEMALE													
1959	16,951	-	-	-	1	1	8	91	452	2,263	6,779	7,353	3
1958	16,599	-	-	-	1	4	16	88	435	2,315	6,547	7,192	1
1957	16,023	-	-	-	1	2	6	90	439	2,223	6,436	6,821	5
1956	15,355	-	-	1	2	5	17	76	388	2,121	6,256	6,483	6
1955	15,326	-	-	-	1	5	11	65	399	2,159	6,296	6,387	3
1954	14,395	-	-	-	2	5	16	72	418	2,070	5,836	5,968	8
1953	15,207	-	-	-	1	9	17	78	424	2,261	6,214	6,197	6
1952	15,051	-	-	-	4	8	9	70	470	2,291	6,366	5,821	12
1951	14,957	-	1	-	-	6	16	83	471	2,313	6,439	5,614	14
1950	14,146	-	-	-	1	-	12	71	457	2,194	6,044	5,356	11

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
GENERAL ARTERIOSCLEROSIS (450)													
ALL OTHER, BOTH SEXES													
1959	2,515	-	-	-	-	5	21	113	342	738	783	505	8
1958	2,620	-	1	-	2	5	23	92	341	789	814	548	5
1957	2,427	-	-	-	1	3	14	93	333	746	708	525	4
1956	2,255	-	-	-	2	2	20	91	334	711	637	451	7
1955	2,293	-	-	-	2	3	28	100	337	657	690	466	10
1954	2,149	-	-	-	-	4	28	107	325	622	618	437	8
1953	2,285	-	-	-	1	9	19	116	371	679	662	421	7
1952	2,147	-	-	-	1	6	31	110	340	632	599	424	4
1951	2,063	-	-	-	-	4	23	123	344	621	571	375	2
1950	2,021	-	-	-	1	7	23	112	327	618	544	384	5
ALL OTHER, MALE													
1959	1,304	-	-	-	-	3	12	65	172	419	413	213	7
1958	1,324	-	1	-	1	3	10	52	164	415	422	254	2
1957	1,222	-	-	-	1	2	6	40	172	406	361	234	-
1956	1,149	-	-	-	1	1	12	45	191	372	324	203	-
1955	1,227	-	-	-	2	1	15	68	185	399	346	205	6
1954	1,120	-	-	-	-	2	13	58	171	339	339	195	3
1953	1,235	-	-	-	1	5	12	67	213	392	347	194	4
1952	1,153	-	-	-	1	3	18	55	198	356	335	185	2
1951	1,116	-	-	-	-	3	7	78	189	349	327	162	1
1950	1,109	-	-	-	-	2	9	58	193	358	318	169	2
ALL OTHER, FEMALE													
1959	1,211	-	-	-	-	2	9	48	170	319	370	292	1
1958	1,296	-	-	-	1	2	13	40	177	374	392	294	3
1957	1,205	-	-	-	-	1	8	53	161	340	347	291	4
1956	1,106	-	-	-	1	1	8	46	143	339	313	248	7
1955	1,066	-	-	-	-	2	13	32	152	258	344	261	4
1954	1,029	-	-	-	-	2	15	49	154	283	279	242	5
1953	1,050	-	-	-	-	4	7	49	158	287	315	227	3
1952	994	-	-	-	-	3	13	55	142	276	264	239	2
1951	947	-	-	-	-	1	16	45	155	272	244	213	1
1950	912	-	-	-	1	5	14	54	134	260	226	215	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)													
BOTH SEXES													
1959	18,175	44	32	52	205	333	804	1,768	3,798	5,702	4,199	1,233	5
1958	17,204	38	29	32	168	384	783	1,758	3,450	5,491	3,891	1,174	6
1957	15,280	57	22	48	180	363	749	1,559	3,478	4,715	3,127	976	6
1956	14,009	55	29	59	161	387	731	1,562	3,056	4,320	2,854	784	11
1955	11,991	51	17	53	189	351	671	1,382	2,642	3,501	2,392	732	10
1954	10,259	57	25	58	169	390	646	1,333	2,320	2,872	1,810	575	4
1953	9,552	64	32	50	190	363	611	1,252	2,207	2,546	1,706	526	5
1952	8,868	65	32	37	175	355	633	1,146	2,141	2,247	1,591	443	3
1951	8,268	68	26	35	201	325	627	1,069	1,897	2,134	1,456	417	13
1950	7,397	48	33	42	188	326	583	1,019	1,716	1,873	1,195	368	6
MALE													
1959	11,131	24	21	21	55	126	424	1,112	2,600	3,881	2,324	538	5
1958	10,650	28	12	13	56	148	409	1,100	2,433	3,688	2,222	536	5
1957	9,551	33	17	26	63	139	408	997	2,460	3,174	1,776	455	3
1956	8,791	35	21	21	43	139	377	945	2,220	2,961	1,640	380	9
1955	7,438	33	10	22	68	132	340	906	1,846	2,353	1,350	373	5
1954	6,337	36	14	18	58	165	335	849	1,605	1,934	1,055	265	3
1953	5,825	39	16	14	58	137	310	815	1,564	1,669	958	244	1
1952	5,413	38	18	9	48	135	321	738	1,527	1,439	918	220	2
1951	4,954	32	12	20	53	124	330	694	1,300	1,391	803	188	7
1950	4,369	26	16	14	57	117	315	632	1,146	1,188	688	167	3
FEMALE													
1959	7,044	20	11	31	150	207	380	656	1,198	1,821	1,875	695	-
1958	6,554	10	17	19	112	236	374	658	1,017	1,803	1,669	638	1
1957	5,729	24	5	22	117	224	341	562	1,018	1,541	1,351	521	3
1956	5,218	20	8	38	118	248	354	617	836	1,359	1,214	404	2
1955	4,553	18	7	31	121	219	331	476	796	1,148	1,042	359	5
1954	3,922	21	11	40	111	225	311	484	715	938	755	310	1
1953	3,727	25	16	36	132	226	301	437	643	877	748	282	4
1952	3,455	27	14	28	127	220	312	408	614	808	673	223	1
1951	3,314	36	14	15	148	201	297	375	597	743	653	229	6
1950	3,028	22	17	28	131	209	268	387	570	685	507	201	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)													
WHITE, BOTH SEXES													
1959	16,243	37	25	41	154	238	587	1,454	3,309	5,267	3,955	1,173	3
1958	15,314	30	19	24	130	267	588	1,368	3,032	5,045	3,691	1,117	3
1957	13,603	37	15	33	128	256	552	1,244	3,043	4,369	2,986	935	5
1956	12,526	37	20	46	110	275	539	1,271	2,728	4,038	2,706	748	8
1955	10,685	38	15	46	149	261	495	1,132	2,335	3,247	2,270	692	5
1954	9,124	43	18	40	131	312	486	1,081	2,063	2,676	1,726	546	2
1953	8,483	44	27	43	144	278	474	1,016	1,978	2,346	1,631	500	2
1952	7,908	51	23	31	131	261	492	950	1,914	2,109	1,521	422	3
1951	7,343	53	25	27	158	253	480	874	1,693	1,992	1,381	395	12
1950	6,583	40	28	38	143	255	451	848	1,537	1,752	1,139	347	5
WHITE, MALE													
1959	10,083	20	18	18	48	98	318	938	2,321	3,604	2,190	507	3
1958	9,678	23	7	11	47	108	320	914	2,190	3,431	2,116	509	2
1957	8,602	23	11	16	45	96	306	821	2,185	2,961	1,705	431	2
1956	8,001	24	13	20	29	106	292	804	2,014	2,775	1,553	365	6
1955	6,748	25	10	20	56	108	262	771	1,656	2,204	1,282	351	3
1954	5,725	27	11	13	48	137	249	701	1,449	1,821	1,015	253	1
1953	5,246	26	13	13	42	111	251	681	1,421	1,545	914	229	-
1952	4,929	28	14	9	42	106	262	633	1,380	1,368	876	209	2
1951	4,461	24	12	15	46	98	259	579	1,184	1,302	759	176	7
1950	3,933	23	14	11	42	95	250	536	1,030	1,116	654	159	3
WHITE, FEMALE													
1959	6,160	17	7	23	106	140	269	516	988	1,663	1,765	666	-
1958	5,636	7	12	13	83	159	268	454	842	1,614	1,575	608	1
1957	5,001	14	4	17	83	160	246	423	858	1,408	1,281	504	3
1956	4,525	13	7	26	81	169	247	467	714	1,263	1,153	383	2
1955	3,937	13	5	26	93	153	233	361	679	1,043	988	341	2
1954	3,399	16	7	27	83	175	237	380	614	855	711	293	1
1953	3,237	18	14	30	102	167	223	335	557	801	717	271	2
1952	2,979	23	9	22	89	155	230	317	534	741	645	213	1
1951	2,882	29	13	12	112	155	221	295	509	690	622	219	5
1950	2,650	17	14	27	101	160	201	312	507	636	485	188	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES OF CIRCULATORY SYSTEM (451-468)													
ALL OTHER, BOTH SEXES													
1959	1,932	7	7	11	51	95	217	314	489	435	244	60	2
1958	1,890	8	10	8	38	117	195	390	418	446	200	57	3
1957	1,677	20	7	15	52	107	197	315	435	346	141	41	1
1956	1,483	18	9	13	51	112	192	291	328	282	148	36	3
1955	1,306	13	2	7	40	90	176	250	307	254	122	40	5
1954	1,135	14	7	18	38	78	160	252	257	196	84	29	2
1953	1,069	20	5	7	46	85	137	236	229	200	75	26	3
1952	960	14	9	6	44	94	141	196	227	138	70	21	-
1951	925	15	1	8	43	72	147	195	204	142	75	22	1
1950	814	8	5	4	45	71	132	171	179	121	56	21	1
ALL OTHER, MALE													
1959	1,048	4	3	3	7	28	106	174	279	277	134	31	2
1958	972	5	5	2	9	40	89	186	243	257	106	27	3
1957	949	10	6	10	18	43	102	176	275	213	71	24	1
1956	790	11	8	1	14	33	85	141	206	186	87	15	3
1955	690	8	-	2	12	24	78	135	190	149	68	22	2
1954	612	9	3	5	10	28	86	148	156	113	40	12	2
1953	579	13	3	1	16	26	59	134	143	124	44	15	1
1952	484	10	4	-	6	29	59	105	147	71	42	11	-
1951	493	8	-	5	7	26	71	115	116	89	44	12	-
1950	436	3	2	3	15	22	65	96	116	72	34	8	-
ALL OTHER, FEMALE													
1959	884	3	4	8	44	67	111	140	210	158	110	29	-
1958	918	3	5	6	29	77	106	204	175	189	94	30	-
1957	728	10	1	5	34	64	95	139	160	133	70	17	-
1956	693	7	1	12	37	79	107	150	122	96	61	21	-
1955	616	5	2	5	28	66	98	115	117	105	54	18	3
1954	523	5	4	13	28	50	74	104	101	83	44	17	-
1953	490	7	2	6	30	59	78	102	86	76	31	11	2
1952	476	4	5	6	38	65	82	91	80	67	28	10	-
1951	432	7	1	3	36	46	76	80	88	53	31	10	1
1950	378	5	3	1	30	49	67	75	63	49	22	13	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)													
BOTH SEXES													
1959	12,299	31	29	177	351	656	1,021	1,506	1,990	2,630	2,641	1,258	9
1958	13,827	30	37	161	420	694	1,100	1,662	2,174	3,110	2,983	1,448	8
1957	15,032	19	46	233	466	814	1,271	1,738	2,479	3,352	3,106	1,504	4
1956	15,304	32	58	211	404	786	1,195	1,754	2,487	3,430	3,364	1,568	15
1955	15,821	21	62	208	463	757	1,208	1,803	2,587	3,600	3,560	1,541	11
1954	17,073	33	86	219	450	877	1,223	2,003	2,880	3,870	3,853	1,565	14
1953	18,774	21	94	245	511	937	1,397	2,227	3,105	4,283	4,139	1,800	15
1952	20,757	28	97	246	477	933	1,548	2,543	3,565	4,685	4,747	1,874	14
1951	22,518	25	118	253	540	969	1,591	2,589	3,889	5,151	5,176	2,195	22
1950	24,677	24	127	232	566	1,066	1,768	2,832	4,127	5,822	5,713	2,371	29
MALE													
1959	6,700	20	15	74	198	397	588	905	1,113	1,422	1,372	592	4
1958	7,498	17	21	90	240	379	660	931	1,225	1,698	1,547	686	4
1957	8,014	10	25	110	276	496	698	986	1,343	1,836	1,535	699	-
1956	8,160	19	34	107	232	470	652	976	1,364	1,811	1,714	772	9
1955	8,471	19	39	102	276	436	691	993	1,392	1,940	1,854	724	5
1954	9,147	17	43	122	256	480	669	1,078	1,564	2,132	2,031	747	8
1953	10,003	14	49	127	274	508	743	1,229	1,670	2,312	2,193	877	7
1952	11,216	16	62	132	271	523	806	1,437	1,967	2,559	2,534	900	9
1951	11,944	18	70	135	282	529	832	1,442	2,144	2,788	2,692	1,000	12
1950	13,085	18	71	120	325	558	924	1,528	2,288	3,127	3,005	1,107	14
FEMALE													
1959	5,599	11	14	103	153	259	433	601	877	1,208	1,269	666	5
1958	6,329	13	16	71	180	315	440	731	949	1,412	1,436	762	4
1957	7,018	9	21	123	190	318	573	752	1,136	1,516	1,571	805	4
1956	7,144	13	24	104	172	316	543	778	1,123	1,619	1,650	796	6
1955	7,350	2	23	106	187	321	517	810	1,195	1,660	1,706	817	6
1954	7,926	16	43	97	194	397	554	925	1,316	1,738	1,822	818	6
1953	8,771	7	45	118	237	429	654	998	1,435	1,971	1,946	923	8
1952	9,541	12	35	114	206	410	742	1,106	1,598	2,126	2,213	974	5
1951	10,574	7	48	118	258	440	759	1,147	1,745	2,363	2,484	1,195	10
1950	11,592	6	56	112	241	508	844	1,304	1,839	2,695	2,708	1,264	15

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)													
WHITE, BOTH SEXES													
1959	9,618	22	23	146	293	500	703	1,039	1,476	2,023	2,283	1,105	5
1958	10,883	23	27	136	359	530	765	1,153	1,536	2,488	2,591	1,273	2
1957	11,879	15	34	197	385	613	861	1,218	1,857	2,693	2,686	1,320	-
1956	12,097	24	45	176	329	591	818	1,180	1,857	2,735	2,944	1,392	6
1955	12,691	19	47	180	390	581	826	1,266	1,939	2,949	3,120	1,370	4
1954	13,674	23	71	188	372	668	859	1,410	2,198	3,116	3,381	1,385	3
1953	15,017	19	76	219	422	712	969	1,565	2,366	3,459	3,594	1,604	12
1952	16,350	26	83	214	393	709	1,049	1,740	2,644	3,770	4,098	1,618	6
1951	17,973	18	100	222	448	726	1,081	1,798	2,980	4,133	4,533	1,919	15
1950	19,768	10	109	208	485	808	1,215	1,995	3,132	4,682	4,995	2,114	15
WHITE, MALE													
1959	5,304	15	12	62	170	316	431	662	855	1,088	1,184	507	2
1958	5,977	12	14	72	205	318	496	678	897	1,354	1,344	587	-
1957	6,435	8	16	94	231	390	510	754	1,020	1,501	1,315	596	-
1956	6,531	14	23	86	194	387	493	702	1,028	1,442	1,484	673	5
1955	6,922	17	30	88	242	359	508	743	1,080	1,591	1,614	650	-
1954	7,341	11	34	102	215	386	497	797	1,225	1,690	1,732	649	3
1953	8,121	13	39	116	244	406	564	917	1,294	1,856	1,884	782	6
1952	8,947	14	53	114	228	425	589	1,041	1,492	2,058	2,148	780	5
1951	9,653	12	57	118	246	430	617	1,067	1,686	2,234	2,319	859	8
1950	10,613	9	60	110	285	457	704	1,152	1,769	2,495	2,596	970	6
WHITE, FEMALE													
1959	4,314	7	11	84	123	184	272	377	621	935	1,099	598	3
1958	4,906	11	13	64	154	212	269	475	639	1,134	1,247	686	2
1957	5,444	7	18	103	154	223	351	464	837	1,192	1,371	724	-
1956	5,566	10	22	90	135	204	325	478	829	1,293	1,460	719	1
1955	5,769	2	17	92	148	222	318	523	859	1,358	1,506	720	4
1954	6,333	12	37	86	157	282	362	613	973	1,426	1,649	736	-
1953	6,896	6	37	103	178	306	405	648	1,072	1,603	1,710	822	6
1952	7,403	12	30	100	165	284	460	699	1,152	1,712	1,950	838	1
1951	8,320	6	43	104	202	296	464	731	1,294	1,899	2,214	1,060	7
1950	9,155	1	49	98	200	351	511	843	1,363	2,187	2,399	1,144	9

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CHRONIC AND UNSPECIFIED NEPHRITIS AND OTHER RENAL SCLEROSIS (592-594)													
ALL OTHER, BOTH SEXES													
1959	2,681	9	6	31	58	156	318	467	514	607	358	153	4
1958	2,944	7	10	25	61	164	335	509	638	622	392	175	6
1957	3,153	4	12	36	81	201	410	520	622	659	420	184	4
1956	3,207	8	13	35	75	195	377	574	630	695	420	176	9
1955	3,130	2	15	28	73	176	382	537	648	651	440	171	7
1954	3,399	10	15	31	78	209	364	593	682	754	472	180	11
1953	3,757	2	18	26	89	225	428	662	739	824	545	196	3
1952	4,407	2	14	32	84	224	499	803	921	915	649	256	8
1951	4,545	7	18	31	92	243	510	791	909	1,018	643	276	7
1950	4,909	14	18	24	81	258	553	837	995	1,140	718	257	14
ALL OTHER, MALE													
1959	1,396	5	3	12	28	81	157	243	258	334	188	85	2
1958	1,521	5	7	18	35	61	164	253	328	344	203	99	4
1957	1,579	2	9	16	45	106	188	232	323	335	220	103	-
1956	1,629	5	11	21	38	83	159	274	336	369	230	99	4
1955	1,549	2	9	14	34	77	183	250	312	349	240	74	5
1954	1,806	6	9	20	41	94	172	281	339	442	299	98	5
1953	1,882	1	10	11	30	102	179	312	376	456	309	95	1
1952	2,269	2	9	18	43	98	217	396	475	501	386	120	4
1951	2,291	6	13	17	36	99	215	375	458	554	373	141	4
1950	2,472	9	11	10	40	101	220	376	519	632	409	137	8
ALL OTHER, FEMALE													
1959	1,285	4	3	19	30	75	161	224	256	273	170	68	2
1958	1,423	2	3	7	26	103	171	256	310	278	189	76	2
1957	1,574	2	3	20	36	95	222	288	299	324	200	81	4
1956	1,578	3	2	14	37	112	218	300	294	326	190	77	5
1955	1,581	-	6	14	39	99	199	287	336	302	200	97	2
1954	1,593	4	6	11	37	115	192	312	343	312	173	82	6
1953	1,875	1	8	15	59	123	249	350	363	368	236	101	2
1952	2,138	-	5	14	41	126	282	407	446	414	263	136	4
1951	2,254	1	5	14	56	144	295	416	451	464	270	135	3
1950	2,437	5	7	14	41	157	333	461	476	508	309	120	6

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)													
BOTH SEXES													
1959	55,039	9,747	2,443	851	627	879	1,824	3,123	5,147	9,345	12,730	8,300	23
1958	57,439	9,918	2,523	763	662	1,002	1,847	3,437	5,885	10,128	12,756	8,498	20
1957	61,001	9,611	2,817	1,241	1,194	1,306	2,382	4,011	6,631	11,009	12,609	8,142	48
1956	47,103	8,634	2,312	736	507	723	1,608	2,671	4,499	8,082	10,507	6,802	22
1955	44,510	8,318	2,196	761	571	751	1,540	2,664	4,308	7,477	9,828	6,068	28
1954	40,991	8,287	2,304	705	506	785	1,490	2,618	3,965	6,742	8,368	5,199	22
1953	52,238	9,074	2,639	864	668	1,003	1,954	3,379	5,401	8,916	11,374	6,915	51
1952	46,265	9,104	2,744	861	627	1,009	1,904	3,099	4,686	7,355	9,442	5,397	37
1951	48,169	8,313	2,647	808	737	961	1,901	3,325	5,223	8,186	10,264	5,760	44
1950	47,120	8,436	2,466	771	715	990	1,935	3,347	5,102	8,100	9,717	5,506	35
MALE													
1959	31,237	5,504	1,334	429	346	481	1,082	2,071	3,487	5,958	6,855	3,677	13
1958	32,931	5,582	1,299	393	369	547	1,155	2,306	4,021	6,513	6,928	3,808	10
1957	34,840	5,422	1,484	658	553	622	1,419	2,686	4,505	7,024	6,740	3,698	29
1956	26,684	4,834	1,219	361	284	398	955	1,785	3,086	5,062	5,646	3,041	13
1955	25,067	4,632	1,175	394	311	420	958	1,793	2,911	4,592	5,210	2,658	13
1954	23,314	4,602	1,227	381	275	455	918	1,792	2,712	4,179	4,407	2,353	13
1953	29,593	5,007	1,432	448	367	534	1,256	2,296	3,695	5,401	5,987	3,134	36
1952	26,043	4,994	1,438	453	335	566	1,210	2,148	3,196	4,416	4,917	2,350	20
1951	26,876	4,610	1,386	440	382	511	1,145	2,216	3,474	4,847	5,328	2,511	26
1950	26,323	4,695	1,331	411	377	488	1,159	2,192	3,390	4,837	4,990	2,436	17
FEMALE													
1959	23,802	4,243	1,109	422	281	398	742	1,052	1,660	3,387	5,875	4,623	10
1958	24,508	4,336	1,224	370	293	455	692	1,131	1,864	3,615	5,828	4,690	10
1957	26,161	4,189	1,333	583	641	684	963	1,325	2,126	3,985	5,869	4,444	19
1956	20,419	3,800	1,093	375	223	325	653	886	1,413	3,020	4,861	3,761	9
1955	19,443	3,686	1,021	367	260	331	582	871	1,397	2,885	4,618	3,410	15
1954	17,677	3,685	1,077	324	231	330	572	826	1,253	2,563	3,961	2,846	9
1953	22,645	4,067	1,207	416	301	469	698	1,083	1,706	3,515	5,387	3,781	15
1952	20,222	4,110	1,306	408	292	443	694	951	1,490	2,939	4,525	3,047	17
1951	21,293	3,703	1,261	368	355	450	756	1,109	1,749	3,339	4,936	3,249	18
1950	20,797	3,741	1,135	360	338	502	776	1,155	1,712	3,263	4,727	3,070	18

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)													
WHITE, BOTH SEXES													
1959	44,764	5,956	1,557	661	480	566	1,178	2,329	4,049	8,142	11,923	7,908	15
1958	46,465	6,155	1,666	583	509	631	1,233	2,527	4,597	8,736	11,814	8,005	9
1957	49,298	5,971	1,855	979	940	866	1,605	2,918	5,211	9,529	11,733	7,663	28
1956	38,221	5,468	1,484	547	389	448	1,057	1,947	3,576	7,016	9,840	6,433	16
1955	35,883	5,262	1,467	585	418	466	1,043	1,876	3,402	6,438	9,158	5,747	21
1954	32,737	5,305	1,532	527	360	502	1,003	1,869	3,129	5,782	7,822	4,891	15
1953	41,553	5,924	1,849	650	456	608	1,285	2,277	4,071	7,494	10,473	6,440	26
1952	36,666	5,844	1,884	663	436	619	1,217	2,195	3,676	6,259	8,780	5,077	16
1951	38,645	5,407	1,861	607	483	603	1,290	2,229	4,097	7,060	9,570	5,406	32
1950	37,820	5,790	1,674	576	529	613	1,284	2,289	3,938	6,934	9,033	5,140	20
WHITE, MALE													
1959	25,334	3,447	840	334	269	308	705	1,551	2,778	5,220	6,392	3,481	9
1958	26,549	3,504	863	308	287	342	769	1,702	3,191	5,638	6,383	3,556	6
1957	28,313	3,505	973	522	440	398	960	2,006	3,624	6,163	6,230	3,475	17
1956	21,695	3,125	787	262	225	264	632	1,337	2,495	4,417	5,271	2,868	12
1955	20,095	2,961	775	306	230	265	641	1,291	2,341	3,955	4,831	2,489	10
1954	18,618	2,987	818	282	199	298	632	1,299	2,173	3,620	4,096	2,206	8
1953	23,570	3,335	974	330	256	319	829	1,623	2,866	4,596	5,511	2,910	21
1952	20,649	3,231	972	346	241	362	789	1,590	2,587	3,782	4,536	2,205	8
1951	21,587	3,012	968	326	259	328	807	1,559	2,785	4,220	4,952	2,351	20
1950	21,231	3,249	896	308	289	317	817	1,590	2,692	4,177	4,628	2,256	12
WHITE, FEMALE													
1959	19,430	2,509	717	327	211	258	473	778	1,271	2,922	5,531	4,427	6
1958	19,916	2,651	803	275	222	289	464	825	1,406	3,098	5,431	4,449	3
1957	20,985	2,466	882	457	500	468	645	912	1,587	3,366	5,503	4,188	11
1956	16,526	2,343	697	285	164	184	425	610	1,081	2,599	4,569	3,565	4
1955	15,788	2,301	692	279	188	201	402	585	1,061	2,483	4,327	3,258	11
1954	14,119	2,318	714	245	161	204	371	570	956	2,162	3,726	2,685	7
1953	17,983	2,589	875	320	200	289	456	654	1,205	2,898	4,962	3,530	5
1952	16,017	2,613	912	317	195	257	428	605	1,089	2,477	4,244	2,872	8
1951	17,058	2,395	893	281	224	275	483	670	1,312	2,840	4,618	3,055	12
1950	16,589	2,541	778	268	240	296	467	699	1,246	2,757	4,405	2,884	8

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFLUENZA AND PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (480-493)													
ALL OTHER, BOTH SEXES													
1959	10,275	3,791	886	190	147	313	646	794	1,098	1,203	807	392	8
1958	10,974	3,763	857	180	153	371	614	910	1,288	1,392	942	493	11
1957	11,703	3,640	962	262	254	440	777	1,093	1,420	1,480	876	479	20
1956	8,882	3,166	828	189	118	275	551	724	923	1,066	667	369	6
1955	8,627	3,056	729	176	153	285	497	788	906	1,039	670	321	7
1954	8,254	2,982	772	178	146	283	487	749	836	960	546	308	7
1953	10,685	3,150	790	214	212	395	669	1,102	1,330	1,422	901	475	25
1952	9,599	3,260	860	198	191	390	687	904	1,010	1,096	662	320	21
1951	9,524	2,906	786	201	254	358	611	1,096	1,126	1,126	694	354	12
1950	9,300	2,646	792	195	186	377	651	1,058	1,164	1,166	684	366	15
ALL OTHER, MALE													
1959	5,903	2,057	494	95	77	173	377	520	709	738	463	196	4
1958	6,382	2,078	436	85	82	205	386	604	830	875	545	252	4
1957	6,527	1,917	511	136	113	224	459	680	881	861	510	223	12
1956	4,989	1,709	432	99	59	134	323	448	591	645	375	173	1
1955	4,972	1,671	400	88	81	155	317	502	570	637	379	169	3
1954	4,696	1,615	409	99	76	157	286	493	539	559	311	147	5
1953	6,023	1,672	458	118	111	215	427	673	829	805	476	224	15
1952	5,394	1,763	466	107	94	204	421	558	609	634	381	145	12
1951	5,289	1,598	418	114	123	183	338	657	689	627	376	160	6
1950	5,092	1,446	435	103	88	171	342	602	698	660	362	180	5
ALL OTHER, FEMALE													
1959	4,372	1,734	392	95	70	140	269	274	389	465	344	196	4
1958	4,592	1,685	421	95	71	166	228	306	458	517	397	241	7
1957	5,176	1,723	451	126	141	216	318	413	539	619	366	256	8
1956	3,893	1,457	396	90	59	141	228	276	332	421	292	196	5
1955	3,655	1,385	329	88	72	130	180	286	336	402	291	152	4
1954	3,558	1,367	363	79	70	126	201	256	297	401	235	161	2
1953	4,662	1,478	332	96	101	180	242	429	501	617	425	251	10
1952	4,205	1,497	394	91	97	186	266	346	401	462	281	175	9
1951	4,235	1,308	368	87	131	175	273	439	437	499	318	194	6
1950	4,208	1,200	357	92	98	206	309	456	466	506	322	186	10

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFLUENZA (480-483)													
BOTH SEXES													
1959	2,845	340	122	101	51	48	103	155	281	460	684	500	-
1958	4,442	446	173	64	92	92	139	279	482	820	1,044	811	-
1957	7,463	646	299	299	318	245	376	551	923	1,466	1,362	970	8
1956	2,323	287	113	36	33	34	71	116	190	394	593	454	2
1955	2,755	369	118	50	50	58	79	171	265	481	661	452	1
1954	2,680	424	144	55	50	57	83	154	235	480	617	379	2
1953	9,460	1,022	369	149	131	142	252	523	917	1,803	2,506	1,634	12
1952	5,631	820	277	146	103	119	212	342	516	938	1,309	840	9
1951	6,946	776	295	133	122	128	231	416	705	1,294	1,726	1,114	6
1950	6,597	734	295	121	109	153	235	416	709	1,191	1,647	983	4
MALE													
1959	1,452	197	73	51	26	23	54	92	159	252	308	217	-
1958	2,342	252	90	28	48	48	76	170	288	481	523	338	-
1957	3,938	357	156	155	135	91	193	321	544	887	656	438	5
1956	1,166	153	62	18	17	15	37	62	113	206	288	195	-
1955	1,336	201	49	25	20	29	45	94	145	255	297	176	-
1954	1,362	224	77	28	32	30	41	98	129	242	295	165	1
1953	4,837	555	185	71	64	51	135	287	521	979	1,239	742	8
1952	2,829	448	158	74	51	55	115	193	294	469	620	349	3
1951	3,490	403	165	73	59	71	125	238	405	683	795	468	5
1950	3,287	396	151	57	54	55	117	231	394	632	784	416	-
FEMALE													
1959	1,393	143	49	50	25	25	49	63	122	208	376	283	-
1958	2,100	194	83	36	44	44	63	109	194	339	521	473	-
1957	3,525	289	143	144	183	154	183	230	379	579	706	532	3
1956	1,157	134	51	18	16	19	34	54	77	188	305	259	2
1955	1,419	168	69	25	30	29	34	77	120	226	364	276	1
1954	1,318	200	67	27	18	27	42	56	106	238	322	214	1
1953	4,623	467	184	78	67	91	117	236	396	824	1,267	892	4
1952	2,802	372	119	72	52	64	97	149	222	469	689	491	6
1951	3,456	373	130	60	63	57	106	178	300	611	931	646	1
1950	3,310	338	144	64	55	98	118	185	315	559	863	567	4

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFLUENZA (480-483)													
WHITE, BOTH SEXES													
1959	2,267	189	85	77	36	35	74	108	203	361	625	474	-
1958	3,340	206	102	44	71	55	90	176	333	644	906	713	-
1957	5,884	328	205	252	263	181	262	381	691	1,212	1,212	893	4
1956	1,753	153	65	18	25	17	39	72	131	308	516	409	-
1955	2,030	185	70	38	27	34	51	102	170	355	590	408	-
1954	1,961	216	88	32	33	34	55	98	156	359	549	340	1
1953	7,294	646	244	106	80	85	146	309	593	1,374	2,235	1,472	4
1952	4,227	457	171	109	71	77	119	202	353	730	1,170	767	1
1951	5,456	442	201	95	77	85	143	239	509	1,075	1,558	1,026	6
1950	5,065	478	205	78	73	86	126	223	452	944	1,506	894	-
WHITE, MALE													
1959	1,147	110	49	39	19	15	39	64	119	206	282	205	-
1958	1,732	122	57	20	38	32	48	106	207	381	435	286	-
1957	3,146	189	100	125	109	68	132	243	439	754	581	404	2
1956	888	80	35	12	14	8	24	39	85	165	252	174	-
1955	971	100	29	19	10	19	29	62	92	187	267	157	-
1954	1,007	116	50	13	22	20	30	70	92	188	257	149	-
1953	3,746	357	113	49	34	31	84	177	356	764	1,110	669	2
1952	2,122	247	94	54	35	37	70	125	220	384	537	319	-
1951	2,767	226	116	49	41	46	91	160	302	583	719	429	5
1950	2,546	253	105	33	38	34	73	143	259	514	718	376	-
WHITE, FEMALE													
1959	1,120	79	36	38	17	20	35	44	84	155	343	269	-
1958	1,608	84	45	24	33	23	42	70	126	263	471	427	-
1957	2,738	139	105	127	154	113	130	138	252	458	631	489	2
1956	865	73	30	6	11	9	15	33	46	143	264	235	-
1955	1,059	85	41	19	17	15	22	40	78	168	323	251	-
1954	954	100	38	19	11	14	25	28	64	171	292	191	1
1953	3,548	289	131	57	46	54	62	132	237	610	1,125	803	2
1952	2,105	210	77	55	36	40	49	77	133	346	633	448	1
1951	2,689	216	85	46	36	39	52	79	207	492	839	597	1
1950	2,519	225	100	45	35	52	53	80	193	430	788	518	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFLUENZA (480-483)													
ALL OTHER, BOTH SEXES													
1959	578	151	37	24	15	13	29	47	78	99	59	26	-
1958	1,102	240	71	20	21	37	49	103	149	176	138	98	-
1957	1,579	318	94	47	55	64	114	170	232	254	150	77	4
1956	570	134	48	18	8	17	32	44	59	86	77	45	2
1955	725	184	48	12	23	24	28	69	95	126	71	44	1
1954	719	208	56	23	17	23	28	56	79	121	68	39	1
1953	2,166	376	125	43	51	57	106	214	324	429	271	162	8
1952	1,404	363	106	37	32	42	93	140	163	208	139	73	8
1951	1,490	334	94	38	45	43	88	177	196	219	168	88	-
1950	1,532	256	90	43	36	67	109	193	257	247	141	89	4
ALL OTHER, MALE													
1959	305	87	24	12	7	8	15	28	40	46	26	12	-
1958	610	130	33	8	10	16	28	64	81	100	88	52	-
1957	792	168	56	30	26	23	61	78	105	133	75	34	3
1956	278	73	27	6	3	7	13	23	28	41	36	21	-
1955	365	101	20	6	10	10	16	32	53	68	30	19	-
1954	355	108	27	15	10	10	11	28	37	54	38	16	1
1953	1,091	198	72	22	30	20	51	110	165	215	129	73	6
1952	707	201	64	20	16	18	45	68	74	85	83	30	3
1951	723	177	49	24	18	25	34	78	103	100	76	39	-
1950	741	143	46	24	16	21	44	88	135	118	66	40	-
ALL OTHER, FEMALE													
1959	273	64	13	12	8	5	14	19	38	53	33	14	-
1958	492	110	38	12	11	21	21	39	68	76	50	46	-
1957	787	150	38	17	29	41	53	92	127	121	75	43	1
1956	292	61	21	12	5	10	19	21	31	45	41	24	2
1955	360	83	28	6	13	14	12	37	42	58	41	25	1
1954	364	100	29	8	7	13	17	28	42	67	30	23	-
1953	1,075	178	53	21	21	37	55	104	159	214	142	89	2
1952	697	162	42	17	16	24	48	72	89	123	56	43	5
1951	767	157	45	14	27	18	54	99	93	119	92	49	-
1950	791	113	44	19	20	46	65	105	122	129	75	49	4

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)													
BOTH SEXES													
1959	52,194	9,407	2,321	750	576	831	1,721	2,968	4,866	8,885	12,046	7,800	23
1958	52,997	9,472	2,350	699	570	910	1,708	3,158	5,403	9,308	11,712	7,687	20
1957	53,538	8,965	2,518	942	876	1,061	2,006	3,460	5,708	9,543	11,247	7,172	40
1956	44,780	8,347	2,199	700	474	689	1,537	2,555	4,309	7,688	9,914	6,348	20
1955	41,755	7,949	2,078	711	521	693	1,461	2,493	4,043	6,996	9,167	5,616	27
1954	38,311	7,863	2,160	650	456	728	1,407	2,464	3,730	6,262	7,751	4,820	20
1953	42,778	8,052	2,270	715	537	861	1,702	2,856	4,484	7,113	8,868	5,281	39
1952	40,634	8,284	2,467	715	524	890	1,692	2,757	4,170	6,417	8,133	4,557	28
1951	41,223	7,537	2,352	675	615	833	1,670	2,909	4,518	6,892	8,538	4,646	38
1950	40,523	7,702	2,171	650	606	837	1,700	2,931	4,393	6,909	8,070	4,523	31
MALE													
1959	29,785	5,307	1,261	378	320	458	1,028	1,979	3,328	5,706	6,547	3,460	13
1958	30,589	5,330	1,209	365	321	499	1,079	2,136	3,733	6,032	6,405	3,470	10
1957	30,902	5,065	1,328	503	418	531	1,226	2,365	3,961	6,137	6,084	3,260	24
1956	25,518	4,681	1,157	343	267	383	918	1,723	2,973	4,856	5,358	2,846	13
1955	23,731	4,431	1,126	369	291	391	913	1,699	2,766	4,337	4,913	2,482	13
1954	21,952	4,378	1,150	353	243	425	877	1,694	2,583	3,937	4,112	2,188	12
1953	24,756	4,452	1,247	377	303	483	1,121	2,009	3,174	4,422	4,748	2,392	28
1952	23,214	4,546	1,280	379	284	511	1,095	1,955	2,902	3,947	4,297	2,001	17
1951	23,386	4,207	1,221	367	323	440	1,020	1,978	3,069	4,164	4,533	2,043	21
1950	23,036	4,299	1,180	354	323	433	1,042	1,961	2,996	4,205	4,206	2,020	17
FEMALE													
1959	22,409	4,100	1,060	372	256	373	693	989	1,538	3,179	5,499	4,340	10
1958	22,408	4,142	1,141	334	249	411	629	1,022	1,670	3,276	5,307	4,217	10
1957	22,636	3,900	1,190	439	458	530	780	1,095	1,747	3,406	5,163	3,912	16
1956	19,262	3,666	1,042	357	207	306	619	832	1,336	2,832	4,556	3,502	7
1955	18,024	3,518	952	342	230	302	548	794	1,277	2,659	4,254	3,134	14
1954	16,359	3,485	1,010	297	213	303	530	770	1,147	2,325	3,639	2,632	8
1953	18,022	3,600	1,023	338	234	378	581	847	1,310	2,691	4,120	2,889	11
1952	17,420	3,738	1,187	336	240	379	597	802	1,268	2,470	3,836	2,556	11
1951	17,837	3,330	1,131	308	292	393	650	931	1,449	2,728	4,005	2,603	17
1950	17,487	3,403	991	296	283	404	658	970	1,397	2,704	3,864	2,503	14

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)													
WHITE, BOTH SEXES													
1959	42,497	5,767	1,472	584	444	531	1,104	2,221	3,846	7,781	11,298	7,434	15
1958	43,125	5,949	1,564	539	438	576	1,143	2,351	4,264	8,092	10,908	7,292	9
1957	43,414	5,643	1,650	727	677	685	1,343	2,537	4,520	8,317	10,521	6,770	24
1956	36,468	5,315	1,419	529	364	431	1,018	1,875	3,445	6,708	9,324	6,024	16
1955	33,853	5,077	1,397	547	391	432	992	1,774	3,232	6,083	8,568	5,339	21
1954	30,776	5,089	1,444	495	327	468	948	1,771	2,973	5,423	7,273	4,551	14
1953	34,259	5,278	1,605	544	376	523	1,139	1,968	3,478	6,120	8,238	4,968	22
1952	32,439	5,387	1,713	554	365	542	1,098	1,993	3,323	5,529	7,610	4,310	15
1951	33,189	4,965	1,660	512	406	518	1,147	1,990	3,588	5,985	8,012	4,380	26
1950	32,755	5,312	1,469	498	456	527	1,158	2,066	3,486	5,990	7,527	4,246	20
WHITE, MALE													
1959	24,187	3,337	791	295	250	293	666	1,487	2,659	5,014	6,110	3,276	9
1958	24,817	3,382	806	288	249	310	721	1,596	2,984	5,257	5,948	3,270	6
1957	25,167	3,316	873	397	331	330	828	1,763	3,185	5,409	5,649	3,071	15
1956	20,807	3,045	752	250	211	256	608	1,298	2,410	4,252	5,019	2,694	12
1955	19,124	2,861	746	287	220	246	612	1,229	2,249	3,768	4,564	2,332	10
1954	17,611	2,871	768	269	177	278	602	1,229	2,081	3,432	3,839	2,057	8
1953	19,824	2,978	861	281	222	288	745	1,446	2,510	3,832	4,401	2,241	19
1952	18,527	2,984	878	292	206	325	719	1,465	2,367	3,398	3,999	1,886	8
1951	18,820	2,786	852	277	218	282	716	1,399	2,483	3,637	4,233	1,922	15
1950	18,685	2,996	791	275	251	283	744	1,447	2,433	3,663	3,910	1,880	12
WHITE, FEMALE													
1959	18,310	2,430	681	289	194	238	438	734	1,187	2,767	5,188	4,158	6
1958	18,308	2,567	758	251	189	266	422	755	1,280	2,835	4,960	4,022	3
1957	18,247	2,327	777	330	346	355	515	774	1,335	2,908	4,872	3,699	9
1956	15,661	2,270	667	279	153	175	410	577	1,035	2,456	4,305	3,330	4
1955	14,729	2,216	651	260	171	186	380	545	983	2,315	4,004	3,007	11
1954	13,165	2,218	676	226	150	190	346	542	892	1,991	3,434	2,494	6
1953	14,435	2,300	744	263	154	235	394	522	968	2,288	3,837	2,727	3
1952	13,912	2,403	835	262	159	217	379	528	956	2,131	3,611	2,424	7
1951	14,369	2,179	808	235	188	236	431	591	1,105	2,348	3,779	2,458	11
1950	14,070	2,316	678	223	205	244	414	619	1,053	2,327	3,617	2,366	8

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
PNEUMONIA, EXCEPT PNEUMONIA OF NEWBORN (490-493)													
ALL OTHER, BOTH SEXES													
1959	9,697	3,640	849	166	132	300	617	747	1,020	1,104	748	366	8
1958	9,872	3,523	786	160	132	334	565	807	1,139	1,216	804	395	11
1957	10,124	3,322	868	215	199	376	663	923	1,188	1,226	726	402	16
1956	8,312	3,032	780	171	110	258	519	680	864	980	590	324	4
1955	7,902	2,872	681	164	130	261	469	719	811	913	599	277	6
1954	7,535	2,774	716	155	129	260	459	693	757	839	478	269	6
1953	8,519	2,774	665	171	161	338	563	888	1,006	993	630	313	17
1952	8,195	2,897	754	161	159	348	594	764	847	888	523	247	13
1951	8,034	2,572	692	163	209	315	523	919	930	907	526	266	12
1950	7,768	2,390	702	152	150	310	542	865	907	919	543	277	11
ALL OTHER, MALE													
1959	5,598	1,970	470	83	70	165	362	492	669	692	437	184	4
1958	5,772	1,948	403	77	72	189	358	540	749	775	457	200	4
1957	5,735	1,749	455	106	87	201	398	602	776	728	435	189	9
1956	4,711	1,636	405	93	56	127	310	425	563	604	339	152	1
1955	4,607	1,570	380	82	71	145	301	470	517	569	349	150	3
1954	4,341	1,507	382	84	66	147	275	465	502	505	273	131	4
1953	4,932	1,474	386	96	81	195	376	563	664	590	347	151	9
1952	4,687	1,562	402	87	78	186	376	490	535	549	298	115	9
1951	4,566	1,421	369	90	105	158	304	579	586	527	300	121	6
1950	4,351	1,303	389	79	72	150	298	514	563	542	296	140	5
ALL OTHER, FEMALE													
1959	4,099	1,670	379	83	62	135	255	255	351	412	311	182	4
1958	4,100	1,575	383	83	60	145	207	267	390	441	347	195	7
1957	4,389	1,573	413	109	112	175	265	321	412	498	291	213	7
1956	3,601	1,396	375	78	54	131	209	255	301	376	251	172	3
1955	3,295	1,302	301	82	59	116	168	249	294	344	250	127	3
1954	3,194	1,267	334	71	63	113	184	228	255	334	205	138	2
1953	3,587	1,300	279	75	80	143	187	325	342	403	283	162	8
1952	3,508	1,335	352	74	81	162	218	274	312	339	225	132	4
1951	3,468	1,151	323	73	104	157	219	340	344	380	226	145	6
1950	3,417	1,087	313	73	78	160	244	351	344	377	247	137	6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
BRONCHITIS (500-502)													
BOTH SEXES													
1959	3,840	735	303	89	39	38	90	240	549	878	593	284	2
1958	3,973	780	387	79	27	48	89	214	583	843	636	285	2
1957	3,581	767	308	92	35	40	80	197	492	740	568	262	-
1956	3,192	783	309	85	17	28	55	129	445	544	526	268	3
1955	3,057	672	341	76	20	30	65	129	400	561	495	268	-
1954	2,872	685	304	59	13	27	40	140	353	502	513	236	-
1953	3,073	668	354	78	25	29	64	160	379	540	524	252	-
1952	2,970	662	354	81	26	38	79	156	340	526	478	228	2
1951	2,926	582	330	67	21	29	74	149	371	517	513	272	1
1950	2,989	584	327	75	21	39	69	145	360	510	519	338	2
MALE													
1959	2,655	452	174	46	21	17	50	163	463	725	393	149	2
1958	2,653	450	221	43	11	22	46	149	480	679	403	147	2
1957	2,348	451	175	53	16	15	45	150	399	573	343	128	-
1956	2,084	443	184	55	11	17	28	99	366	423	334	122	2
1955	1,930	403	201	43	12	17	33	82	325	423	270	121	-
1954	1,825	413	176	27	8	16	27	97	290	370	305	96	-
1953	1,961	383	222	43	15	15	39	113	321	400	295	115	-
1952	1,843	378	202	47	17	20	50	127	273	361	269	98	1
1951	1,823	335	200	36	13	10	41	114	292	366	276	140	-
1950	1,783	352	189	48	8	16	44	94	277	331	256	168	-
FEMALE													
1959	1,185	283	129	43	18	21	40	77	86	153	200	135	-
1958	1,320	330	166	36	16	26	43	65	103	164	233	138	-
1957	1,233	316	133	39	19	25	35	47	93	167	225	134	-
1956	1,108	340	125	30	6	11	27	30	79	121	192	146	1
1955	1,127	269	140	33	8	13	32	47	75	138	225	147	-
1954	1,047	272	128	32	5	11	13	43	63	132	208	140	-
1953	1,112	285	132	35	10	14	25	47	58	140	229	137	-
1952	1,127	284	152	34	9	18	29	29	67	165	209	130	1
1951	1,103	247	130	31	8	19	33	35	79	151	237	132	1
1950	1,206	232	138	27	13	23	25	51	83	179	263	170	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
BRONCHITIS (500-502)													
WHITE, BOTH SEXES													
1959	3,409	550	242	78	33	33	67	216	511	832	569	277	1
1958	3,484	594	317	66	21	33	68	188	519	786	615	276	1
1957	3,121	574	262	84	26	28	59	157	442	700	541	248	-
1956	2,800	586	264	75	15	22	43	117	405	508	507	256	2
1955	2,676	494	298	65	16	23	43	109	356	538	478	256	-
1954	2,523	505	254	56	10	20	28	122	333	473	496	226	-
1953	2,689	514	295	71	20	18	51	136	339	502	503	240	-
1952	2,567	483	299	72	22	27	60	124	310	490	460	220	-
1951	2,569	443	282	60	20	23	55	124	338	480	486	258	-
1950	2,614	457	274	63	15	27	46	116	321	469	501	323	2
WHITE, MALE													
1959	2,398	348	142	40	16	16	38	146	432	696	378	145	1
1958	2,369	348	180	37	9	14	36	139	429	642	392	142	1
1957	2,089	348	151	50	15	12	33	123	362	547	325	123	-
1956	1,854	330	159	49	10	13	23	92	334	398	328	116	2
1955	1,710	300	175	37	11	11	21	70	301	407	262	115	-
1954	1,620	315	142	27	6	11	18	83	277	348	300	93	-
1953	1,732	294	185	41	13	9	34	99	289	375	283	110	-
1952	1,608	272	175	41	16	14	39	102	253	342	259	95	-
1951	1,623	255	171	32	12	9	32	95	276	344	262	135	-
1950	1,582	280	156	41	5	11	30	82	259	308	248	162	-
WHITE, FEMALE													
1959	1,011	202	100	38	17	17	29	70	79	136	191	132	-
1958	1,115	246	137	29	12	19	32	49	90	144	223	134	-
1957	1,032	226	111	34	11	16	26	34	80	153	216	125	-
1956	946	256	105	26	5	9	20	25	71	110	179	140	-
1955	966	194	123	28	5	12	22	39	55	131	216	141	-
1954	903	190	112	29	4	9	10	39	56	125	196	133	-
1953	957	220	110	30	7	9	17	37	50	127	220	130	-
1952	959	211	124	31	6	13	21	22	57	148	201	125	-
1951	946	188	111	28	8	14	23	29	62	136	224	123	-
1950	1,032	177	118	22	10	16	16	34	62	161	253	161	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
BRONCHITIS (500-502)													
ALL OTHER, BOTH SEXES													
1959	431	185	61	11	6	5	23	24	38	46	24	7	1
1958	489	186	70	13	6	15	21	26	64	57	21	9	1
1957	460	193	46	8	9	12	21	40	50	40	27	14	-
1956	392	197	45	10	2	6	12	12	40	36	19	12	1
1955	381	178	43	11	4	7	22	20	44	23	17	12	-
1954	349	180	50	3	3	7	12	18	20	29	17	10	-
1953	384	154	59	7	5	11	13	24	40	38	21	12	-
1952	403	179	55	9	4	11	19	32	30	36	18	8	2
1951	357	139	48	7	1	6	19	25	33	37	27	14	1
1950	375	127	53	12	6	12	23	29	39	41	18	15	-
ALL OTHER, MALE													
1959	257	104	32	6	5	1	12	17	31	29	15	4	1
1958	284	102	41	6	2	8	10	10	51	37	11	5	1
1957	259	103	24	3	1	3	12	27	37	26	18	5	-
1956	230	113	25	6	1	4	5	7	32	25	6	6	-
1955	220	103	26	6	1	6	12	12	24	16	8	6	-
1954	205	98	34	-	2	5	9	14	13	22	5	3	-
1953	229	89	37	2	2	6	5	14	32	25	12	5	-
1952	235	106	27	6	1	6	11	25	20	19	10	3	1
1951	200	80	29	4	1	1	9	19	16	22	14	5	-
1950	201	72	33	7	3	5	14	12	18	23	8	6	-
ALL OTHER, FEMALE													
1959	174	81	29	5	1	4	11	7	7	17	9	3	-
1958	205	84	29	7	4	7	11	16	13	20	10	4	-
1957	201	90	22	5	8	9	9	13	13	14	9	9	-
1956	162	84	20	4	1	2	7	5	8	11	13	6	1
1955	161	75	17	5	3	1	10	8	20	7	9	6	-
1954	144	82	16	3	1	2	3	4	7	7	12	7	-
1953	155	65	22	5	3	5	8	10	8	13	9	7	-
1952	168	73	28	3	3	5	8	7	10	17	8	5	1
1951	157	59	19	3	-	5	10	6	17	15	13	9	1
1950	174	55	20	5	3	7	9	17	21	18	10	9	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ULCER OF STOMACH AND DUODEMA (540, 541)													
BOTH SEXES													
1959	10,674	66	15	17	37	161	636	1,439	2,296	3,045	2,357	600	5
1958	10,801	75	15	12	37	206	649	1,485	2,433	3,025	2,251	608	5
1957	10,365	69	13	13	30	170	644	1,484	2,391	2,949	2,002	598	2
1956	10,028	51	10	12	33	196	600	1,443	2,313	2,847	1,975	546	2
1955	9,730	58	12	9	48	189	658	1,460	2,340	2,760	1,766	428	2
1954	9,391	69	9	8	40	193	720	1,501	2,281	2,599	1,572	393	6
1953	9,240	70	13	9	32	207	682	1,532	2,352	2,459	1,534	346	4
1952	8,770	48	12	9	40	203	721	1,494	2,294	2,327	1,322	292	8
1951	8,487	41	9	1	48	228	722	1,473	2,250	2,234	1,210	266	5
1950	8,214	50	5	14	58	206	740	1,483	2,233	2,066	1,134	221	4
MALE													
1959	7,975	49	9	9	23	122	478	1,135	1,890	2,318	1,596	344	2
1958	8,130	39	10	7	27	156	478	1,166	1,989	2,343	1,567	344	4
1957	7,866	35	6	7	19	120	511	1,202	1,946	2,297	1,383	338	2
1956	7,737	34	6	6	26	152	482	1,173	1,918	2,247	1,364	327	2
1955	7,563	41	6	4	34	141	526	1,212	1,945	2,155	1,255	242	2
1954	7,429	41	2	7	31	148	574	1,257	1,927	2,062	1,134	241	5
1953	7,427	39	8	5	22	166	558	1,286	2,009	1,992	1,115	224	3
1952	7,050	39	6	5	32	158	584	1,267	1,952	1,877	944	181	5
1951	6,840	27	5	1	34	183	612	1,255	1,890	1,792	868	169	4
1950	6,647	36	4	9	40	179	602	1,234	1,911	1,651	841	137	3
FEMALE													
1959	2,699	17	6	8	14	39	158	304	406	727	761	256	3
1958	2,671	36	5	5	10	50	171	319	444	682	684	264	1
1957	2,499	34	7	6	11	50	133	282	445	652	619	260	-
1956	2,291	17	4	6	7	44	118	270	395	600	611	219	-
1955	2,167	17	6	5	14	48	132	248	395	605	511	186	-
1954	1,962	28	7	1	9	45	146	244	354	537	438	152	1
1953	1,813	31	5	4	10	41	124	246	343	467	419	122	1
1952	1,720	9	6	4	8	45	137	227	342	450	378	111	3
1951	1,647	14	4	-	14	45	110	218	360	442	342	97	1
1950	1,567	14	1	5	18	27	138	249	322	415	293	84	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ULCER OF STOMACH AND DUODEMA (540, 541)													
WHITE, BOTH SEXES													
1959	9,855	40	12	15	30	133	518	1,263	2,110	2,872	2,274	583	5
1958	9,937	45	15	8	25	162	531	1,321	2,231	2,848	2,153	594	4
1957	9,577	46	13	12	23	132	549	1,317	2,183	2,798	1,924	578	2
1956	9,297	30	10	11	28	153	514	1,285	2,122	2,703	1,904	536	1
1955	8,982	37	10	9	36	148	558	1,295	2,161	2,614	1,694	418	2
1954	8,651	47	7	8	28	154	604	1,330	2,095	2,466	1,523	384	5
1953	8,501	49	12	7	23	167	595	1,337	2,159	2,328	1,487	334	3
1952	8,112	29	10	9	27	165	614	1,330	2,146	2,224	1,272	280	6
1951	7,809	32	7	1	35	185	620	1,290	2,079	2,108	1,193	254	5
1950	7,525	38	4	12	40	165	619	1,308	2,071	1,960	1,087	218	3
WHITE, MALE													
1959	7,355	28	6	7	19	102	395	1,004	1,735	2,185	1,539	333	2
1958	7,528	24	10	6	19	133	401	1,047	1,836	2,209	1,502	338	3
1957	7,269	21	6	6	16	92	439	1,070	1,784	2,180	1,327	326	2
1956	7,213	20	6	5	22	124	419	1,061	1,771	2,147	1,316	321	1
1955	7,017	28	5	4	27	113	447	1,088	1,813	2,051	1,203	236	2
1954	6,883	28	1	7	23	121	488	1,131	1,782	1,965	1,099	234	4
1953	6,861	24	8	3	16	132	492	1,140	1,856	1,892	1,079	217	2
1952	6,537	22	5	5	21	132	505	1,139	1,832	1,790	907	176	3
1951	6,336	21	3	1	25	151	533	1,117	1,764	1,697	860	160	4
1950	6,108	27	3	7	30	143	513	1,103	1,767	1,574	805	134	2
WHITE, FEMALE													
1959	2,500	12	6	8	11	31	123	259	375	687	735	250	3
1958	2,409	21	5	2	6	29	130	274	395	639	651	256	1
1957	2,308	25	7	6	7	40	110	247	399	618	597	252	-
1956	2,084	10	4	6	6	29	95	224	351	556	588	215	-
1955	1,965	9	5	5	9	35	111	207	348	563	491	182	-
1954	1,768	19	6	1	5	33	116	199	313	501	424	150	1
1953	1,640	25	4	4	7	35	103	197	303	436	408	117	1
1952	1,575	7	5	4	6	33	109	191	314	434	365	104	3
1951	1,473	11	4	-	10	34	87	173	315	411	333	94	1
1950	1,417	11	1	5	10	22	106	205	304	386	282	84	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ULCER OF STOMACH AND DUODEMA (540, 541)													
ALL OTHER, BOTH SEXES													
1959	819	26	3	2	7	28	118	176	186	173	83	17	-
1958	864	30	-	4	12	44	118	164	202	177	98	14	1
1957	788	23	-	1	7	38	95	167	208	151	78	20	-
1956	731	21	-	1	5	43	86	158	191	144	71	10	1
1955	748	21	2	-	12	41	100	165	179	146	72	10	-
1954	740	22	2	-	12	39	116	171	186	133	49	9	1
1953	739	21	1	2	9	40	87	195	193	131	47	12	1
1952	658	19	2	-	13	38	107	164	148	103	50	12	2
1951	678	9	2	-	13	43	102	183	171	126	17	12	-
1950	689	12	1	2	18	41	121	175	162	106	47	3	1
ALL OTHER, MALE													
1959	620	21	3	2	4	20	83	131	155	133	57	11	-
1958	602	15	-	1	8	23	77	119	153	134	65	6	1
1957	597	14	-	1	3	28	72	132	162	117	56	12	-
1956	524	14	-	1	4	28	63	112	147	100	48	6	1
1955	546	13	1	-	7	28	79	124	132	104	52	6	-
1954	546	13	1	-	8	27	86	126	145	97	35	7	1
1953	566	15	-	2	6	34	66	146	153	100	36	7	1
1952	513	17	1	-	11	26	79	128	120	87	37	5	2
1951	504	6	2	-	9	32	79	138	126	95	8	9	-
1950	539	9	1	2	10	36	89	131	144	77	36	3	1
ALL OTHER, FEMALE													
1959	199	5	-	-	3	8	35	45	31	40	26	6	-
1958	262	15	-	3	4	21	41	45	49	43	33	8	-
1957	191	9	-	-	4	10	23	35	46	34	22	8	-
1956	207	7	-	-	1	15	23	46	44	44	23	4	-
1955	202	8	1	-	5	13	21	41	47	42	20	4	-
1954	194	9	1	-	4	12	30	45	41	36	14	2	-
1953	173	6	1	-	3	6	21	49	40	31	11	5	-
1952	145	2	1	-	2	12	28	36	28	16	13	7	-
1951	174	3	-	-	4	11	23	45	45	31	9	3	-
1950	150	3	-	-	8	5	32	44	18	29	11	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
APPENDICITIS (550-553)													
BOTH SEXES													
1959	1,837	19	45	147	85	61	143	223	304	406	302	102	-
1958	1,845	24	44	124	71	84	110	234	299	432	327	96	-
1957	1,982	14	60	147	71	66	136	254	365	455	325	88	1
1956	2,103	23	69	141	92	104	146	279	336	468	336	107	2
1955	2,273	19	83	134	93	119	163	323	394	518	329	97	1
1954	2,278	19	69	173	103	121	183	276	429	473	327	102	3
1953	2,398	18	88	184	123	161	221	366	417	468	280	70	2
1952	2,710	17	115	220	175	169	257	399	473	465	332	87	1
1951	2,846	16	147	237	190	192	302	382	462	503	340	72	3
1950	3,080	19	163	242	224	219	314	427	534	576	286	75	1
MALE													
1959	1,185	14	27	88	53	32	95	153	220	268	177	58	-
1958	1,146	13	25	75	40	43	70	153	200	282	202	43	-
1957	1,267	10	30	92	39	39	89	188	248	314	180	37	1
1956	1,336	14	33	85	60	67	93	203	236	303	191	49	2
1955	1,427	13	50	78	55	71	102	230	270	327	179	51	1
1954	1,455	10	41	100	55	77	115	200	299	315	191	50	2
1953	1,583	14	56	104	83	93	150	263	300	307	171	40	2
1952	1,693	11	67	134	93	91	167	276	314	303	192	45	-
1951	1,761	9	88	135	112	107	212	241	317	306	196	36	2
1950	1,956	13	87	132	133	136	222	285	357	368	182	41	-
FEMALE													
1959	652	5	18	59	32	29	48	70	84	138	125	44	-
1958	699	11	19	49	31	41	40	81	99	150	125	53	-
1957	715	4	30	55	32	27	47	66	117	141	145	51	-
1956	767	9	36	56	32	37	53	76	100	165	145	58	-
1955	846	6	33	56	38	48	61	93	124	191	150	46	-
1954	823	9	28	73	48	44	68	76	130	158	136	52	1
1953	815	4	32	80	40	68	71	103	117	161	109	30	-
1952	1,017	6	48	86	82	78	90	123	159	162	140	42	1
1951	1,085	7	59	102	78	85	90	141	145	197	144	36	1
1950	1,124	6	76	110	91	83	92	142	177	208	104	34	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
APPENDICITIS (550-553)													
WHITE, BOTH SEXES													
1959	1,520	16	36	111	61	48	106	171	253	350	274	94	-
1958	1,553	19	34	94	52	55	75	194	247	385	308	90	-
1957	1,664	11	51	109	54	43	95	204	293	412	306	86	-
1956	1,782	19	54	114	75	71	112	216	276	418	320	105	2
1955	1,951	15	75	106	74	86	125	260	343	467	310	90	-
1954	1,913	16	62	138	70	85	139	219	367	413	303	99	2
1953	2,055	14	78	155	90	125	163	298	369	426	271	66	-
1952	2,265	12	104	178	137	128	179	314	401	423	306	82	1
1951	2,372	12	132	189	144	137	232	284	398	452	321	69	2
1950	2,608	15	148	194	171	166	234	335	472	530	271	72	-
WHITE, MALE													
1959	1,001	11	23	70	39	26	71	125	187	233	161	55	-
1958	978	9	23	54	32	29	51	134	169	248	190	39	-
1957	1,089	8	28	70	30	26	69	160	206	284	171	37	-
1956	1,155	11	29	69	50	49	81	164	196	276	180	48	2
1955	1,238	10	48	60	46	51	88	189	237	295	166	48	-
1954	1,243	8	39	81	37	56	90	165	265	277	175	48	2
1953	1,382	10	51	91	61	79	120	218	267	282	166	37	-
1952	1,432	8	61	108	75	76	120	228	262	279	173	42	-
1951	1,491	8	77	103	85	85	175	195	271	277	180	34	1
1950	1,677	10	81	104	106	107	178	235	314	332	171	39	-
WHITE, FEMALE													
1959	519	5	13	41	22	22	35	46	66	117	113	39	-
1958	575	10	11	40	20	26	24	60	78	137	118	51	-
1957	575	3	23	39	24	17	26	44	87	128	135	49	-
1956	627	8	25	45	25	22	31	52	80	142	140	57	-
1955	713	5	27	46	28	35	37	71	106	172	144	42	-
1954	670	8	23	57	33	29	49	54	102	136	128	51	-
1953	673	4	27	64	29	46	43	80	102	144	105	29	-
1952	833	4	43	70	62	52	59	86	139	144	133	40	1
1951	881	4	55	86	59	52	57	89	127	175	141	35	1
1950	931	5	67	90	65	59	56	100	158	198	100	33	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
APPENDICITIS (550-553)													
ALL OTHER, BOTH SEXES													
1959	317	3	9	36	24	13	37	52	51	56	28	8	-
1958	292	5	10	30	19	29	35	40	52	47	19	6	-
1957	318	3	9	38	17	23	41	50	72	43	19	2	1
1956	321	4	15	27	17	33	34	63	60	50	16	2	-
1955	322	4	8	28	19	33	38	63	51	51	19	7	1
1954	365	3	7	35	33	36	44	57	62	60	24	3	1
1953	343	4	10	29	33	36	58	68	48	42	9	4	2
1952	445	5	11	42	38	41	78	85	72	42	26	5	-
1951	474	4	15	48	46	55	70	98	64	51	19	3	1
1950	472	4	15	48	53	53	80	92	62	46	15	3	1
ALL OTHER, MALE													
1959	184	3	4	18	14	6	24	28	33	35	16	3	-
1958	168	4	2	21	8	14	19	19	31	34	12	4	-
1957	178	2	2	22	9	13	20	28	42	30	9	-	1
1956	181	3	4	16	10	18	12	39	40	27	11	1	-
1955	189	3	2	18	9	20	14	41	33	32	13	3	1
1954	212	2	2	19	18	21	25	35	34	38	16	2	-
1953	201	4	5	13	22	14	30	45	33	25	5	3	2
1952	261	3	6	26	18	15	47	48	52	24	19	3	-
1951	270	1	11	32	27	22	37	46	46	29	16	2	1
1950	279	3	6	28	27	29	44	50	43	36	11	2	-
ALL OTHER, FEMALE													
1959	133	-	5	18	10	7	13	24	18	21	12	5	-
1958	124	1	8	9	11	15	16	21	21	13	7	2	-
1957	140	1	7	16	8	10	21	22	30	13	10	2	-
1956	140	1	11	11	7	15	22	24	20	23	5	1	-
1955	133	1	6	10	10	13	24	22	18	19	6	4	-
1954	153	1	5	16	15	15	19	22	28	22	8	1	1
1953	142	-	5	16	11	22	28	23	15	17	4	1	-
1952	184	2	5	16	20	26	31	37	20	18	7	2	-
1951	204	3	4	16	19	33	33	52	18	22	3	1	-
1950	193	1	9	20	26	24	36	42	19	10	4	1	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)													
BOTH SEXES													
1959	9,106	956	95	54	56	110	292	651	1,366	2,228	2,292	1,005	1
1958	8,853	951	96	50	62	125	328	722	1,290	2,134	2,215	876	4
1957	8,571	1,042	92	53	55	105	351	658	1,170	2,080	2,062	894	9
1956	8,453	1,078	86	55	59	145	299	646	1,252	1,962	2,090	777	4
1955	8,654	956	109	39	56	129	361	736	1,322	2,068	2,073	801	4
1954	8,498	1,024	99	57	66	156	354	767	1,281	2,028	1,960	701	5
1953	8,727	995	119	63	78	196	392	783	1,370	2,071	1,984	672	4
1952	8,481	953	143	52	80	198	417	760	1,350	2,034	1,821	669	4
1951	8,627	966	131	66	83	220	453	851	1,410	2,057	1,785	600	5
1950	8,894	960	124	70	94	226	454	933	1,447	2,035	1,903	642	6
MALE													
1959	4,531	585	57	32	30	53	127	306	746	1,143	1,023	429	-
1958	4,462	589	60	33	38	54	142	361	701	1,100	1,015	367	2
1957	4,380	623	68	41	37	55	146	320	653	1,074	955	403	5
1956	4,339	663	56	44	27	64	118	322	664	1,020	1,020	337	4
1955	4,474	550	74	27	36	49	158	352	749	1,066	1,061	349	3
1954	4,476	623	66	39	38	64	156	383	695	1,122	959	329	2
1953	4,575	615	81	44	33	92	157	380	768	1,107	1,010	286	2
1952	4,497	568	94	33	41	83	177	398	731	1,133	931	305	3
1951	4,568	589	93	40	52	88	198	406	786	1,134	911	269	2
1950	4,649	583	85	47	51	90	209	442	813	1,078	946	301	4
FEMALE													
1959	4,575	371	38	22	26	57	165	345	620	1,085	1,269	576	1
1958	4,391	362	36	17	24	71	186	361	589	1,034	1,200	509	2
1957	4,191	419	24	12	18	50	205	338	517	1,006	1,107	491	4
1956	4,114	415	30	11	32	81	181	324	588	942	1,070	440	-
1955	4,180	406	35	12	20	80	203	384	573	1,002	1,012	452	1
1954	4,022	401	33	18	28	92	198	384	586	906	1,001	372	3
1953	4,152	380	38	19	45	104	235	403	602	964	974	386	2
1952	3,984	385	49	19	39	115	240	362	619	901	890	364	1
1951	4,059	377	38	26	31	132	255	445	624	923	874	331	3
1950	4,245	377	39	23	43	136	245	491	634	957	957	341	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)													
WHITE, BOTH SEXES													
1959	8,044	796	76	40	40	78	195	505	1,152	2,023	2,182	956	1
1958	7,702	777	66	37	42	85	237	544	1,087	1,877	2,103	843	4
1957	7,525	870	74	40	36	74	253	497	983	1,882	1,955	854	7
1956	7,421	904	74	48	39	106	214	494	1,051	1,771	1,969	749	2
1955	7,587	798	87	30	41	91	252	544	1,124	1,868	1,969	780	3
1954	7,436	866	79	43	41	98	244	585	1,096	1,855	1,853	671	5
1953	7,605	816	85	50	53	136	289	611	1,174	1,872	1,880	637	2
1952	7,411	800	102	42	56	126	293	585	1,162	1,874	1,725	643	3
1951	7,518	825	90	53	60	149	330	648	1,218	1,869	1,702	571	3
1950	7,693	795	92	56	57	143	327	699	1,222	1,869	1,802	626	5
WHITE, MALE													
1959	3,974	481	44	24	24	39	92	241	631	1,024	967	407	-
1958	3,838	471	40	22	26	39	103	287	593	958	949	348	2
1957	3,801	504	58	30	24	40	108	253	552	961	888	379	4
1956	3,762	551	49	39	16	44	89	243	551	906	955	317	2
1955	3,883	461	62	22	23	37	115	265	622	947	989	338	2
1954	3,878	522	52	30	23	38	111	285	592	1,012	902	309	2
1953	3,961	497	57	35	23	66	120	302	648	987	958	268	-
1952	3,912	472	63	31	32	55	128	306	626	1,042	866	289	2
1951	3,973	504	67	32	38	52	145	321	678	1,028	851	256	1
1950	3,982	482	64	37	27	54	152	332	675	978	886	292	3
WHITE, FEMALE													
1959	4,070	315	32	16	16	39	103	264	521	999	1,215	549	1
1958	3,864	306	26	15	16	46	134	257	494	919	1,154	495	2
1957	3,724	366	16	10	12	34	145	244	431	921	1,067	475	3
1956	3,659	353	25	9	23	62	125	251	500	865	1,014	432	-
1955	3,704	337	25	8	18	54	137	279	502	921	980	442	1
1954	3,558	344	27	13	18	60	133	300	504	843	951	362	3
1953	3,644	319	28	15	30	70	169	309	526	885	922	369	2
1952	3,499	328	39	11	24	71	165	279	536	832	859	354	1
1951	3,545	321	23	21	22	97	185	327	540	841	851	315	2
1950	3,711	313	28	19	30	89	175	367	547	891	916	334	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HERNIA AND INTESTINAL OBSTRUCTION (560, 561, 570)													
ALL OTHER, BOTH SEXES													
1959	1,062	160	19	14	16	32	97	146	214	205	110	49	-
1958	1,151	174	30	13	20	40	91	178	203	257	112	33	-
1957	1,046	172	18	13	19	31	98	161	187	198	107	40	2
1956	1,032	174	12	7	20	39	85	152	201	191	121	28	2
1955	1,067	158	22	9	15	38	109	192	198	200	104	21	1
1954	1,062	158	20	14	25	58	110	182	185	173	107	30	-
1953	1,122	179	34	13	25	60	103	172	196	199	104	35	2
1952	1,070	153	41	10	24	72	124	175	188	160	96	26	1
1951	1,109	141	41	13	23	71	123	203	192	188	83	29	2
1950	1,201	165	32	14	37	83	127	234	225	166	101	16	1
ALL OTHER, MALE													
1959	557	104	13	8	6	14	35	65	115	119	56	22	-
1958	624	118	20	11	12	15	39	74	108	142	66	19	-
1957	579	119	10	11	13	15	38	67	101	113	67	24	1
1956	577	112	7	5	11	20	29	79	113	114	65	20	2
1955	591	89	12	5	13	12	43	87	127	119	72	11	1
1954	598	101	14	9	15	26	45	98	103	110	57	20	-
1953	614	118	24	9	10	26	37	78	120	120	52	18	2
1952	585	96	31	2	9	28	49	92	105	91	65	16	1
1951	595	85	26	8	14	36	53	85	108	106	60	13	1
1950	667	101	21	10	24	36	57	110	138	100	60	9	1
ALL OTHER, FEMALE													
1959	505	56	6	6	10	18	62	81	99	86	54	27	-
1958	527	56	10	2	8	25	52	104	95	115	46	14	-
1957	467	53	8	2	6	16	60	94	86	85	40	16	1
1956	455	62	5	2	9	19	56	73	88	77	56	8	-
1955	476	69	10	4	2	26	66	105	71	81	32	10	-
1954	464	57	6	5	10	32	65	84	82	63	50	10	-
1953	508	61	10	4	15	34	66	94	76	79	52	17	-
1952	485	57	10	8	15	44	75	83	83	69	31	10	-
1951	514	56	15	5	9	35	70	118	84	82	23	16	1
1950	534	64	11	4	13	47	70	124	87	66	41	7	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)													
BOTH SEXES													
1959	7,758	2,550	543	104	100	134	268	417	752	1,179	1,192	518	1
1958	7,838	2,669	640	93	102	174	271	448	726	1,168	1,074	473	-
1957	7,971	3,096	523	102	93	143	275	436	727	1,088	1,052	430	6
1956	7,508	2,806	571	101	97	162	236	408	644	1,038	985	458	2
1955	7,754	3,257	650	82	90	139	236	397	677	897	944	379	6
1954	7,892	3,590	565	99	91	183	266	433	618	838	840	367	2
1953	8,510	4,215	681	106	113	177	289	409	584	814	773	347	2
1952	8,743	4,367	841	149	110	154	248	420	589	780	776	306	3
1951	8,036	3,945	763	111	113	193	264	377	517	735	711	304	3
1950	7,622	3,748	689	90	126	172	216	364	525	732	671	285	4
MALE													
1959	3,872	1,393	312	66	42	67	141	244	413	551	464	178	1
1958	3,997	1,469	342	48	56	88	139	264	390	595	423	183	-
1957	4,089	1,681	278	55	48	66	154	246	411	552	444	150	4
1956	3,941	1,560	317	55	43	87	120	239	373	551	409	186	1
1955	4,033	1,790	367	41	36	72	125	236	390	458	384	131	3
1954	4,160	1,987	303	52	33	76	135	268	382	447	354	122	1
1953	4,454	2,291	374	58	57	82	146	233	323	408	352	129	1
1952	4,708	2,529	444	78	61	78	124	233	343	367	325	126	-
1951	4,210	2,206	399	66	55	90	135	209	302	353	283	111	1
1950	4,036	2,115	359	46	61	82	111	217	301	349	281	112	2
FEMALE													
1959	3,886	1,157	231	38	58	67	127	173	339	628	728	340	-
1958	3,841	1,200	298	45	46	86	132	184	336	573	651	290	-
1957	3,882	1,415	245	47	45	77	121	190	316	536	608	280	2
1956	3,567	1,246	254	46	54	75	116	169	271	487	576	272	1
1955	3,721	1,467	283	41	54	67	111	161	287	439	560	248	3
1954	3,732	1,603	262	47	58	107	131	165	236	391	486	245	1
1953	4,056	1,924	307	48	56	95	143	176	261	406	421	218	1
1952	4,035	1,838	397	71	49	76	124	187	246	413	451	180	3
1951	3,826	1,739	364	45	58	103	129	168	215	382	428	193	2
1950	3,586	1,633	330	44	65	90	105	147	224	383	390	173	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)													
WHITE, BOTH SEXES													
1959	5,933	1,298	364	79	86	120	238	358	655	1,096	1,148	490	1
1958	6,088	1,494	451	70	92	148	238	402	648	1,071	1,029	445	-
1957	5,993	1,653	364	77	83	120	237	379	660	996	1,011	412	1
1956	5,785	1,585	414	76	88	143	199	359	583	952	946	439	1
1955	5,833	1,858	454	63	82	128	196	350	598	829	907	362	6
1954	5,939	2,183	363	78	81	149	234	365	559	778	803	345	1
1953	6,308	2,588	456	78	102	149	246	349	515	757	734	332	2
1952	6,339	2,646	561	118	95	129	191	343	520	699	737	298	2
1951	6,126	2,634	501	92	103	168	229	309	455	680	664	289	2
1950	5,943	2,650	501	70	114	145	170	293	449	665	617	266	3
WHITE, MALE													
1959	2,875	716	198	49	36	58	122	208	368	509	445	165	1
1958	3,066	820	241	36	50	76	123	241	361	550	400	168	-
1957	3,022	888	195	42	43	56	133	221	379	503	421	140	1
1956	3,003	882	232	38	39	74	106	216	339	506	391	179	1
1955	3,024	1,036	267	33	31	70	106	210	351	425	365	127	3
1954	3,103	1,223	199	39	31	58	118	232	340	412	339	111	1
1953	3,313	1,435	253	46	51	67	124	206	292	384	331	123	1
1952	3,382	1,540	301	62	55	68	102	190	307	328	307	122	-
1951	3,172	1,473	259	54	51	78	122	177	265	326	261	105	1
1950	3,107	1,498	256	33	56	65	89	179	256	316	255	103	1
WHITE, FEMALE													
1959	3,058	582	166	30	50	62	116	150	287	587	703	325	-
1958	3,022	674	210	34	42	72	115	161	287	521	629	277	-
1957	2,971	765	169	35	40	64	104	158	281	493	590	272	-
1956	2,782	703	182	38	49	69	93	143	244	446	555	260	-
1955	2,809	822	187	30	51	58	90	140	247	404	542	235	3
1954	2,836	960	164	39	50	91	116	133	219	366	464	234	-
1953	2,995	1,153	203	32	51	82	122	143	223	373	403	209	1
1952	2,957	1,106	260	56	40	61	89	153	213	371	430	176	2
1951	2,954	1,161	242	38	52	90	107	132	190	354	403	184	1
1950	2,836	1,152	245	37	58	80	81	114	193	349	362	163	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
GASTRITIS, DUODENITIS, ENTERITIS, AND COLITIS, EXCEPT DIARRHEA OF NEWBORN (543, 571, 572)													
ALL OTHER, BOTH SEXES													
1959	1,825	1,252	179	25	14	14	30	59	97	83	44	28	-
1958	1,750	1,175	189	23	10	26	33	46	78	97	45	28	-
1957	1,978	1,443	159	25	10	23	38	57	67	92	41	18	5
1956	1,723	1,221	157	25	9	19	37	49	61	86	39	19	1
1955	1,921	1,399	196	19	8	11	40	47	79	68	37	17	-
1954	1,953	1,407	202	21	10	34	32	68	59	60	37	22	1
1953	2,202	1,627	225	28	11	28	43	60	69	57	39	15	-
1952	2,404	1,721	280	31	15	25	57	77	69	81	39	8	1
1951	1,910	1,311	262	19	10	25	35	68	62	55	47	15	1
1950	1,679	1,098	188	20	12	27	46	71	76	67	54	19	1
ALL OTHER, MALE													
1959	997	677	114	17	6	9	19	36	45	42	19	13	-
1958	931	649	101	12	6	12	16	23	29	45	23	15	-
1957	1,067	793	83	13	5	10	21	25	32	49	23	10	3
1956	938	678	85	17	4	13	14	23	34	45	18	7	-
1955	1,009	754	100	8	5	2	19	26	39	33	19	4	-
1954	1,057	764	104	13	2	18	17	36	42	35	15	11	-
1953	1,141	856	121	12	6	15	22	27	31	24	21	6	-
1952	1,326	989	143	16	6	10	22	43	36	39	18	4	-
1951	1,038	733	140	12	4	12	13	32	37	27	22	6	-
1950	929	617	103	13	5	17	22	38	45	33	26	9	1
ALL OTHER, FEMALE													
1959	828	575	65	8	8	5	11	23	52	41	25	15	-
1958	819	526	88	11	4	14	17	23	49	52	22	13	-
1957	911	650	76	12	5	13	17	32	35	43	18	8	2
1956	785	543	72	8	5	6	23	26	27	41	21	12	1
1955	912	645	96	11	3	9	21	21	40	35	18	13	-
1954	896	643	98	8	8	16	15	32	17	25	22	11	1
1953	1,061	771	104	16	5	13	21	33	38	33	18	9	-
1952	1,078	732	137	15	9	15	35	34	33	42	21	4	1
1951	872	578	122	7	6	13	22	36	25	28	25	9	1
1950	750	481	85	7	7	10	24	33	31	34	28	10	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>CIRRHOSIS OF LIVER (581)</b>													
<b>BOTH SEXES</b>													
1959	19,242	36	30	28	69	649	2,749	5,265	4,893	3,771	1,491	257	4
1958	18,638	42	21	37	68	573	2,618	4,932	4,731	3,863	1,482	268	3
1957	19,317	42	24	41	64	687	2,832	4,990	4,940	3,937	1,489	268	3
1956	17,924	37	28	33	72	566	2,576	4,556	4,577	3,691	1,518	261	9
1955	16,763	38	24	33	55	509	2,383	4,287	4,195	3,535	1,411	284	9
1954	16,201	45	19	42	70	518	2,337	3,979	4,180	3,376	1,355	273	7
1953	16,399	41	21	40	69	543	2,386	4,178	4,232	3,202	1,421	261	5
1952	15,934	45	24	40	78	562	2,346	3,996	4,165	3,086	1,333	248	11
1951	15,075	43	13	36	63	537	2,199	3,715	3,917	2,962	1,327	250	13
1950	13,855	37	16	32	76	488	1,997	3,316	3,631	2,819	1,209	225	9
<b>MALE</b>													
1959	12,667	21	11	13	29	330	1,640	3,451	3,432	2,686	917	133	4
1958	12,360	21	13	18	22	297	1,549	3,320	3,316	2,777	885	141	1
1957	12,853	25	12	20	18	352	1,678	3,396	3,512	2,780	915	142	3
1956	11,720	15	17	11	30	276	1,541	3,017	3,197	2,589	880	140	7
1955	11,055	24	11	13	24	246	1,437	2,879	2,959	2,436	881	139	6
1954	10,720	21	13	17	35	260	1,402	2,708	2,975	2,331	808	144	6
1953	10,729	22	8	22	24	285	1,404	2,782	3,010	2,198	840	130	4
1952	10,609	18	10	21	24	284	1,449	2,676	3,015	2,163	806	135	8
1951	9,827	23	6	18	25	255	1,354	2,469	2,698	2,035	826	108	10
1950	9,059	23	5	15	21	243	1,244	2,219	2,555	1,932	692	104	6
<b>FEMALE</b>													
1959	6,575	15	19	15	40	319	1,109	1,814	1,461	1,085	574	124	-
1958	6,278	21	8	19	46	276	1,069	1,612	1,415	1,086	597	127	2
1957	6,464	17	12	21	46	335	1,154	1,594	1,428	1,157	574	126	-
1956	6,204	22	11	22	42	290	1,035	1,539	1,380	1,102	638	121	2
1955	5,708	14	13	20	31	263	946	1,408	1,236	1,099	530	145	3
1954	5,481	24	6	25	35	258	935	1,271	1,205	1,045	547	129	1
1953	5,670	19	13	18	45	258	982	1,396	1,222	1,004	581	131	1
1952	5,325	27	14	19	54	278	897	1,320	1,150	923	527	113	3
1951	5,248	20	7	18	38	282	845	1,246	1,219	927	501	142	3
1950	4,796	14	11	17	55	245	753	1,097	1,076	887	517	121	3

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>CIRRHOSIS OF LIVER (581)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	17,331	30	22	21	47	411	2,286	4,723	4,511	3,584	1,444	250	2
1958	17,006	36	18	34	44	386	2,228	4,467	4,418	3,685	1,435	253	2
1957	17,557	32	17	31	42	501	2,410	4,527	4,542	3,767	1,425	260	3
1956	16,371	26	22	28	53	401	2,187	4,147	4,237	3,539	1,469	254	8
1955	15,436	28	19	27	46	380	2,076	3,897	3,947	3,381	1,354	273	8
1954	14,968	38	18	37	54	400	2,043	3,647	3,913	3,235	1,312	264	7
1953	15,184	27	19	32	50	430	2,125	3,797	3,998	3,067	1,385	251	3
1952	14,753	38	22	33	58	445	2,083	3,651	3,928	2,962	1,291	234	8
1951	13,930	35	11	33	51	443	1,945	3,375	3,678	2,824	1,282	240	13
1950	12,834	30	13	24	66	409	1,786	3,037	3,410	2,671	1,162	219	7
<b>WHITE, MALE</b>													
1959	11,552	17	8	9	22	223	1,390	3,128	3,188	2,550	889	126	2
1958	11,367	18	10	16	11	218	1,350	3,017	3,102	2,640	855	129	1
1957	11,802	18	9	16	15	263	1,451	3,112	3,250	2,661	867	137	3
1956	10,813	9	13	9	23	208	1,337	2,777	2,970	2,478	849	133	7
1955	10,277	19	9	11	20	193	1,277	2,652	2,790	2,326	844	131	5
1954	9,997	17	12	16	29	201	1,250	2,522	2,796	2,234	776	138	6
1953	10,005	13	8	18	19	229	1,276	2,546	2,847	2,099	822	126	2
1952	9,858	15	9	17	17	225	1,298	2,459	2,838	2,072	776	127	5
1951	9,145	21	6	15	23	215	1,218	2,262	2,534	1,938	800	103	10
1950	8,456	18	4	11	18	211	1,138	2,060	2,399	1,826	663	103	5
<b>WHITE, FEMALE</b>													
1959	5,779	13	14	12	25	188	896	1,595	1,323	1,034	555	124	-
1958	5,639	18	8	18	33	168	878	1,450	1,316	1,045	580	124	1
1957	5,755	14	8	15	27	238	959	1,415	1,292	1,106	558	123	-
1956	5,558	17	9	19	30	193	850	1,370	1,267	1,061	620	121	1
1955	5,159	9	10	16	26	187	799	1,245	1,157	1,055	510	142	3
1954	4,971	21	6	21	25	199	793	1,125	1,117	1,001	536	126	1
1953	5,179	14	11	14	31	201	849	1,251	1,151	968	563	125	1
1952	4,895	23	13	16	41	220	785	1,192	1,090	890	515	107	3
1951	4,785	14	5	18	28	228	727	1,113	1,144	886	482	137	3
1950	4,378	12	9	13	48	198	648	977	1,011	845	499	116	2

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>CIRRHOSIS OF LIVER (581)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	1,911	6	8	7	22	238	463	542	382	187	47	7	2
1958	1,632	6	3	3	24	187	390	465	313	178	47	15	1
1957	1,760	10	7	10	22	186	422	463	398	170	64	8	-
1956	1,553	11	6	5	19	165	389	409	340	152	49	7	1
1955	1,327	10	5	6	9	129	307	390	248	154	57	11	1
1954	1,233	7	1	5	16	118	294	332	267	141	43	9	-
1953	1,215	14	2	8	19	113	261	381	234	135	36	10	2
1952	1,181	7	2	7	20	117	263	345	237	124	42	14	3
1951	1,145	8	2	3	12	94	254	340	239	138	45	10	-
1950	1,021	7	3	8	10	79	211	279	221	148	47	6	2
<b>ALL OTHER, MALE</b>													
1959	1,115	4	3	4	7	107	250	323	244	136	28	7	2
1958	993	3	3	2	11	79	199	303	214	137	30	12	-
1957	1,051	7	3	4	3	89	227	284	262	119	48	5	-
1956	907	6	4	2	7	68	204	240	227	111	31	7	-
1955	778	5	2	2	4	53	160	227	169	110	37	8	1
1954	723	4	1	1	6	59	152	186	179	97	32	6	-
1953	724	9	-	4	5	56	128	236	163	99	18	4	2
1952	751	3	1	4	7	59	151	217	177	91	30	8	3
1951	682	2	-	3	2	40	136	207	164	97	26	5	-
1950	603	5	1	4	3	32	106	159	156	106	29	1	1
<b>ALL OTHER, FEMALE</b>													
1959	796	2	5	3	15	131	213	219	138	51	19	-	-
1958	639	3	-	1	13	108	191	162	99	41	17	3	1
1957	709	3	4	6	19	97	195	179	136	51	16	3	-
1956	646	5	2	3	12	97	185	169	113	41	18	-	1
1955	549	5	3	4	5	76	147	163	79	44	20	3	-
1954	510	3	-	4	10	59	142	146	88	44	11	3	-
1953	491	5	2	4	14	57	133	145	71	36	18	6	-
1952	430	4	1	3	13	58	112	128	60	33	12	6	-
1951	463	6	2	-	10	54	118	133	75	41	19	5	-
1950	418	2	2	4	7	47	105	120	65	42	18	5	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)													
BOTH SEXES													
1959	4,497	2	-	2	13	50	134	347	748	1,373	1,326	501	1
1958	4,720	5	-	-	12	60	157	355	791	1,415	1,436	487	2
1957	4,944	-	2	2	24	78	166	410	834	1,536	1,406	486	-
1956	5,262	1	-	2	10	70	186	417	964	1,632	1,459	520	1
1955	5,939	4	-	2	25	107	217	513	1,223	1,861	1,508	477	2
1954	5,690	3	1	1	20	89	214	547	1,167	1,784	1,425	435	4
1953	5,602	5	-	-	13	94	204	588	1,259	1,757	1,301	378	3
1952	5,903	3	1	1	15	92	260	658	1,315	1,844	1,335	376	3
1951	5,724	4	1	1	18	94	262	677	1,234	1,746	1,323	359	5
1950	5,870	2	1	-	16	97	263	706	1,360	1,816	1,275	334	-
MALE													
1959	1,831	2	-	1	1	19	49	148	331	584	524	172	-
1958	1,918	2	-	-	1	15	60	151	336	578	627	148	-
1957	1,998	-	1	1	6	17	55	176	358	688	539	157	-
1956	2,066	1	-	2	3	19	56	174	412	673	568	157	1
1955	2,294	2	-	-	3	31	81	212	486	761	570	147	1
1954	2,190	1	1	1	4	22	79	207	446	745	550	132	2
1953	2,127	3	-	-	4	19	72	212	507	679	510	121	-
1952	2,129	1	-	1	3	17	91	252	488	678	481	117	-
1951	2,110	3	1	-	5	33	100	261	461	647	490	109	-
1950	2,120	1	1	-	3	21	87	257	499	653	487	111	-
FEMALE													
1959	2,666	-	-	1	12	31	85	199	417	789	802	329	1
1958	2,802	3	-	-	11	45	97	204	455	837	809	339	2
1957	2,946	-	1	1	18	61	111	234	476	848	867	329	-
1956	3,196	-	-	-	7	51	130	243	552	959	891	363	-
1955	3,645	2	-	2	22	76	136	301	737	1,100	938	330	1
1954	3,500	2	-	-	16	67	135	340	721	1,039	875	303	2
1953	3,475	2	-	-	9	75	132	376	752	1,078	791	257	3
1952	3,774	2	1	-	12	75	169	406	827	1,166	854	259	3
1951	3,614	1	-	1	13	61	162	416	773	1,099	833	250	5
1950	3,750	1	-	-	13	76	176	449	861	1,163	788	223	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)													
WHITE, BOTH SEXES													
1959	4,244	2	-	1	13	45	112	298	696	1,305	1,279	492	1
1958	4,474	3	-	-	10	52	135	312	736	1,345	1,401	478	2
1957	4,704	-	1	2	22	65	146	370	777	1,478	1,371	472	-
1956	5,022	1	-	2	6	58	165	384	907	1,564	1,427	507	1
1955	5,694	2	-	1	22	91	192	471	1,156	1,805	1,487	465	2
1954	5,450	2	1	1	16	81	191	492	1,118	1,719	1,401	424	4
1953	5,385	3	-	-	12	88	189	552	1,211	1,698	1,266	363	3
1952	5,675	3	1	1	14	84	239	608	1,257	1,783	1,313	369	3
1951	5,515	3	-	1	15	88	233	643	1,178	1,700	1,298	353	3
1950	5,659	-	1	-	12	84	239	659	1,307	1,773	1,255	329	-
WHITE, MALE													
1959	1,749	2	-	-	1	18	42	134	315	559	511	167	-
1958	1,827	1	-	-	1	14	53	131	320	551	613	143	-
1957	1,919	-	-	1	6	17	50	162	339	669	524	151	-
1956	1,985	1	-	2	1	18	54	161	394	648	553	152	1
1955	2,215	-	-	-	2	26	75	201	465	738	564	143	1
1954	2,120	1	1	1	3	21	72	195	429	720	545	130	2
1953	2,060	3	-	-	3	17	69	203	491	661	499	114	-
1952	2,060	1	-	1	3	16	86	235	474	655	473	116	-
1951	2,035	2	-	-	4	29	90	252	444	628	479	107	-
1950	2,059	-	1	-	2	17	79	244	483	647	476	110	-
WHITE, FEMALE													
1959	2,495	-	-	1	12	27	70	164	381	746	768	325	1
1958	2,647	2	-	-	9	38	82	181	416	794	788	335	2
1957	2,785	-	1	1	16	48	96	208	438	809	847	321	-
1956	3,037	-	-	-	5	40	111	223	513	916	874	355	-
1955	3,479	2	-	1	20	65	117	270	691	1,067	923	322	1
1954	3,330	1	-	-	13	60	119	297	689	999	856	294	2
1953	3,325	-	-	-	9	71	120	349	720	1,037	767	249	3
1952	3,615	2	1	-	11	68	153	373	783	1,128	840	253	3
1951	3,480	1	-	1	11	59	143	391	734	1,072	819	246	3
1950	3,600	-	-	-	10	67	160	415	824	1,126	779	219	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>CHOLELITHIASIS, CHOLECYSTITIS, AND CHOLANGITIS (584, 585)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	253	-	-	1	-	5	22	49	52	68	47	9	-
1958	246	2	-	-	2	8	22	43	55	70	35	9	-
1957	240	-	1	-	2	13	20	40	57	58	35	14	-
1956	240	-	-	-	4	12	21	33	57	68	32	13	-
1955	245	2	-	1	3	16	25	42	67	56	21	12	-
1954	240	1	-	-	4	8	23	55	49	65	24	11	-
1953	217	2	-	-	1	6	15	36	48	59	35	15	-
1952	228	-	-	-	1	8	21	50	58	61	22	7	-
1951	209	1	1	-	3	6	29	34	56	46	25	6	2
1950	211	2	-	-	4	13	24	47	53	43	20	5	-
<b>ALL OTHER, MALE</b>													
1959	82	-	-	1	-	1	7	14	16	25	13	5	-
1958	91	1	-	-	-	1	7	20	16	27	14	5	-
1957	79	-	1	-	-	-	5	14	19	19	15	6	-
1956	81	-	-	-	2	1	2	13	18	25	15	5	-
1955	79	2	-	-	1	5	6	11	21	23	6	4	-
1954	70	-	-	-	1	1	7	12	17	25	5	2	-
1953	67	-	-	-	1	2	3	9	16	18	11	7	-
1952	69	-	-	-	-	1	5	17	14	23	8	1	-
1951	75	1	1	-	1	4	10	9	17	19	11	2	-
1950	61	1	-	-	1	4	8	13	16	6	11	1	-
<b>ALL OTHER, FEMALE</b>													
1959	171	-	-	-	-	4	15	35	36	43	34	4	-
1958	155	1	-	-	2	7	15	23	39	43	21	4	-
1957	161	-	-	-	2	13	15	26	38	39	20	8	-
1956	159	-	-	-	2	11	19	20	39	43	17	8	-
1955	166	-	-	1	2	11	19	31	46	33	15	8	-
1954	170	1	-	-	3	7	16	43	32	40	19	9	-
1953	150	2	-	-	-	4	12	27	32	41	24	8	-
1952	159	-	-	-	1	7	16	33	44	38	14	6	-
1951	134	-	-	-	2	2	19	25	39	27	14	4	2
1950	150	1	-	-	3	9	16	34	37	37	9	4	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
NEPHRITIS AND NEPHROSIS (590-594)													
BOTH SEXES													
1959	14,120	81	100	309	453	758	1,212	1,729	2,250	2,935	2,919	1,364	10
1958	16,030	82	142	301	514	824	1,298	1,939	2,467	3,508	3,344	1,601	10
1957	17,307	72	173	363	577	956	1,467	1,997	2,810	3,772	3,463	1,651	6
1956	17,681	79	155	346	507	924	1,433	2,044	2,836	3,891	3,735	1,714	17
1955	18,214	71	188	341	573	925	1,429	2,121	2,950	4,024	3,914	1,666	12
1954	19,588	90	221	352	582	1,052	1,460	2,312	3,237	4,307	4,238	1,720	17
1953	21,563	77	243	400	674	1,143	1,679	2,573	3,529	4,748	4,546	1,934	17
1952	23,888	91	265	450	669	1,165	1,878	2,943	4,036	5,196	5,159	2,021	15
1951	25,782	90	312	453	732	1,198	1,926	2,965	4,375	5,711	5,633	2,362	25
1950	28,147	86	320	406	764	1,313	2,132	3,252	4,661	6,425	6,205	2,551	32
MALE													
1959	7,727	51	64	133	258	456	699	1,042	1,269	1,592	1,518	641	4
1958	8,698	45	77	170	300	444	755	1,088	1,386	1,932	1,736	759	6
1957	9,289	37	97	182	338	572	804	1,135	1,545	2,082	1,732	764	1
1956	9,444	39	91	183	290	545	785	1,135	1,561	2,068	1,896	840	11
1955	9,819	47	125	174	335	529	813	1,191	1,585	2,181	2,060	773	6
1954	10,540	57	128	190	333	571	803	1,263	1,764	2,367	2,224	829	11
1953	11,551	41	146	216	366	604	899	1,423	1,927	2,569	2,416	936	8
1952	12,915	58	162	245	367	636	985	1,673	2,228	2,846	2,745	961	9
1951	13,665	48	169	229	381	636	1,024	1,653	2,399	3,106	2,928	1,080	12
1950	15,035	53	186	221	451	698	1,126	1,803	2,592	3,433	3,253	1,203	16
FEMALE													
1959	6,393	30	36	176	195	302	513	687	981	1,343	1,401	723	6
1958	7,332	37	65	131	214	380	543	851	1,081	1,576	1,608	842	4
1957	8,018	35	76	181	239	384	663	862	1,265	1,690	1,731	887	5
1956	8,237	40	64	163	217	379	648	909	1,275	1,823	1,839	874	6
1955	8,395	24	63	167	238	396	616	930	1,365	1,843	1,854	893	6
1954	9,048	33	93	162	249	481	657	1,049	1,473	1,940	2,014	891	6
1953	10,012	36	97	184	308	539	780	1,150	1,602	2,179	2,130	998	9
1952	10,973	33	103	205	302	529	893	1,270	1,808	2,350	2,414	1,060	6
1951	12,117	42	143	224	351	562	902	1,312	1,976	2,605	2,705	1,282	13
1950	13,112	33	134	185	313	615	1,006	1,449	2,069	2,992	2,952	1,348	16

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
NEPHRITIS AND NEPHROSIS (590-594)													
WHITE, BOTH SEXES													
1959	11,014	61	65	241	367	574	824	1,198	1,677	2,280	2,527	1,195	5
1958	12,597	66	92	244	426	616	887	1,347	1,769	2,821	2,920	1,406	3
1957	13,626	53	124	300	466	698	991	1,406	2,113	3,022	3,002	1,451	-
1956	13,931	59	116	280	407	678	975	1,390	2,129	3,107	3,264	1,519	7
1955	14,514	54	140	289	474	693	974	1,473	2,209	3,283	3,437	1,483	5
1954	15,667	69	180	300	469	790	1,021	1,634	2,479	3,476	3,729	1,516	4
1953	17,129	58	180	344	541	849	1,146	1,804	2,690	3,831	3,959	1,714	13
1952	18,760	76	212	384	527	862	1,271	2,024	2,994	4,189	4,466	1,749	6
1951	20,403	65	239	375	585	867	1,301	2,052	3,349	4,560	4,930	2,064	16
1950	22,425	53	264	346	633	972	1,433	2,285	3,542	5,170	5,432	2,278	17
WHITE, MALE													
1959	6,105	40	40	107	219	363	504	762	978	1,230	1,313	547	2
1958	6,921	34	52	133	250	361	563	788	1,027	1,545	1,516	651	1
1957	7,442	28	69	155	283	440	584	866	1,177	1,700	1,484	656	-
1956	7,526	29	64	139	236	437	587	825	1,190	1,641	1,642	730	6
1955	7,971	34	93	149	290	427	594	883	1,231	1,779	1,794	696	1
1954	8,456	46	102	161	273	448	595	934	1,382	1,889	1,904	718	4
1953	9,324	32	110	190	323	473	667	1,060	1,494	2,059	2,081	828	7
1952	10,285	46	129	210	308	506	714	1,216	1,696	2,287	2,333	835	5
1951	10,971	33	129	190	327	502	749	1,217	1,886	2,482	2,523	925	8
1950	12,124	34	148	193	387	557	837	1,339	2,011	2,740	2,814	1,057	7
WHITE, FEMALE													
1959	4,909	21	25	134	148	211	320	436	699	1,050	1,214	648	3
1958	5,676	32	40	111	176	255	324	559	742	1,276	1,404	755	2
1957	6,184	25	55	145	183	258	407	540	936	1,322	1,518	795	-
1956	6,405	30	52	141	171	241	388	565	939	1,466	1,622	789	1
1955	6,543	20	47	140	184	266	380	590	978	1,504	1,643	787	4
1954	7,211	23	78	139	196	342	426	700	1,097	1,587	1,825	798	-
1953	7,805	26	70	154	218	376	479	744	1,196	1,772	1,878	886	6
1952	8,475	30	83	174	219	356	557	808	1,298	1,902	2,133	914	1
1951	9,432	32	110	185	258	365	552	835	1,463	2,078	2,407	1,139	8
1950	10,301	19	116	153	246	415	596	946	1,531	2,430	2,618	1,221	10

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
NEPHRITIS AND NEPHROSIS (590-594)													
ALL OTHER, BOTH SEXES													
1959	3,106	20	35	68	86	184	388	531	573	655	392	169	5
1958	3,433	16	50	57	88	208	411	592	698	687	424	195	7
1957	3,681	19	49	63	111	258	476	591	697	750	461	200	6
1956	3,750	20	39	66	100	246	458	654	707	784	471	195	10
1955	3,700	17	48	52	99	232	455	648	741	741	477	183	7
1954	3,921	21	41	52	113	262	439	678	758	831	509	204	13
1953	4,434	19	63	56	133	294	533	769	839	917	587	220	4
1952	5,128	15	53	66	142	303	607	919	1,042	1,007	693	272	9
1951	5,379	25	73	78	147	331	625	913	1,026	1,151	703	298	9
1950	5,722	33	56	60	131	341	699	967	1,119	1,255	773	273	15
ALL OTHER, MALE													
1959	1,622	11	24	26	39	93	195	280	291	362	205	94	2
1958	1,777	11	25	37	50	83	192	300	359	387	220	108	5
1957	1,847	9	28	27	55	132	220	269	368	382	248	108	1
1956	1,918	10	27	44	54	108	198	310	371	427	254	110	5
1955	1,848	13	32	25	45	102	219	308	354	402	266	77	5
1954	2,084	11	26	29	60	123	208	329	382	478	320	111	7
1953	2,227	9	36	26	43	131	232	363	433	510	335	108	1
1952	2,630	12	33	35	59	130	271	457	532	559	412	126	4
1951	2,694	15	40	39	54	134	275	436	513	624	405	155	4
1950	2,911	19	38	28	64	141	289	464	581	693	439	146	9
ALL OTHER, FEMALE													
1959	1,484	9	11	42	47	91	193	251	282	293	187	75	3
1958	1,656	5	25	20	38	125	219	292	339	300	204	87	2
1957	1,834	10	21	36	56	126	256	322	329	368	213	92	5
1956	1,832	10	12	22	46	138	260	344	336	357	217	85	5
1955	1,852	4	16	27	54	130	236	340	387	339	211	106	2
1954	1,837	10	15	23	53	139	231	349	376	353	189	93	6
1953	2,207	10	27	30	90	163	301	406	406	407	252	112	3
1952	2,498	3	20	31	83	173	336	462	510	448	281	146	5
1951	2,685	10	33	39	93	197	350	477	513	527	298	143	5
1950	2,811	14	18	32	67	200	410	503	538	562	334	127	6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)													
BOTH SEXES													
1959	1,821	50	71	132	102	102	191	223	260	305	278	106	1
1958	2,203	52	105	140	94	130	198	277	293	398	361	153	2
1957	2,275	53	127	130	111	142	196	259	331	420	357	147	2
1956	2,377	47	97	135	103	138	238	290	349	461	371	146	2
1955	2,393	50	126	133	110	168	221	318	363	424	354	125	1
1954	2,515	57	135	133	132	175	237	309	357	437	385	155	3
1953	2,789	56	149	155	163	206	282	346	424	465	407	134	2
1952	3,131	63	168	204	192	232	330	400	471	511	412	147	1
1951	3,264	65	194	200	192	229	335	376	486	560	457	167	3
1950	3,470	62	193	174	198	247	364	420	534	603	492	180	3
MALE													
1959	1,027	31	49	59	60	59	111	137	156	170	146	49	-
1958	1,200	28	56	80	60	65	95	157	161	234	189	73	2
1957	1,275	27	72	72	62	76	106	149	202	246	197	65	1
1956	1,284	20	57	76	58	75	133	159	197	257	182	68	2
1955	1,348	28	86	72	59	93	122	198	193	241	206	49	1
1954	1,393	40	85	68	77	91	134	185	200	235	193	82	3
1953	1,548	27	97	89	92	96	156	194	257	257	223	59	1
1952	1,699	42	100	113	96	113	179	236	261	287	211	61	-
1951	1,721	30	99	94	99	107	192	211	255	318	236	80	-
1950	1,950	35	115	101	126	140	202	275	304	306	248	96	2
FEMALE													
1959	794	19	22	73	42	43	80	86	104	135	132	57	1
1958	1,003	24	49	60	34	65	103	120	132	164	172	80	-
1957	1,000	26	55	58	49	66	90	110	129	174	160	82	1
1956	1,093	27	40	59	45	63	105	131	152	204	189	78	-
1955	1,045	22	40	61	51	75	99	120	170	183	148	76	-
1954	1,122	17	50	65	55	84	103	124	157	202	192	73	-
1953	1,241	29	52	66	71	110	126	152	167	208	184	75	1
1952	1,432	21	68	91	96	119	151	164	210	224	201	86	1
1951	1,543	35	95	106	93	122	143	165	231	242	221	87	3
1950	1,520	27	78	73	72	107	162	145	230	297	244	84	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)													
WHITE, BOTH SEXES													
1959	1,396	39	42	95	74	74	121	159	201	257	244	90	-
1958	1,714	43	65	108	67	86	122	194	233	333	329	133	1
1957	1,747	38	90	103	81	85	130	188	256	329	316	131	-
1956	1,834	35	71	104	78	87	157	210	272	372	320	127	1
1955	1,823	35	93	109	84	112	148	207	270	334	317	113	1
1954	1,993	46	109	112	97	122	162	224	281	360	348	131	1
1953	2,112	39	104	125	119	137	177	239	324	372	365	110	1
1952	2,410	50	129	170	134	153	222	284	350	419	368	131	-
1951	2,430	47	139	153	137	141	220	254	369	427	397	145	1
1950	2,657	43	155	138	148	164	218	290	410	488	437	164	2
WHITE, MALE													
1959	801	25	28	45	49	47	73	100	123	142	129	40	-
1958	944	22	38	61	45	43	67	110	130	191	172	64	1
1957	1,007	20	53	61	52	50	74	112	157	199	169	60	-
1956	995	15	41	53	42	50	94	123	162	199	158	57	1
1955	1,049	17	63	61	48	68	86	140	151	188	180	46	1
1954	1,115	35	68	59	58	62	98	137	157	199	172	69	1
1953	1,203	19	71	74	79	67	103	143	200	203	197	46	1
1952	1,338	32	76	96	80	81	125	175	204	229	185	55	-
1951	1,318	21	72	72	81	72	132	150	200	248	204	66	-
1950	1,511	25	88	83	102	100	133	187	242	245	218	87	1
WHITE, FEMALE													
1959	595	14	14	50	25	27	48	59	78	115	115	50	-
1958	770	21	27	47	22	43	55	84	103	142	157	69	-
1957	740	18	37	42	29	35	56	76	99	130	147	71	-
1956	839	20	30	51	36	37	63	87	110	173	162	70	-
1955	774	18	30	48	36	44	62	67	119	146	137	67	-
1954	878	11	41	53	39	60	64	87	124	161	176	62	-
1953	909	20	33	51	40	70	74	96	124	169	168	64	-
1952	1,072	18	53	74	54	72	97	109	146	190	183	76	-
1951	1,112	26	67	81	56	69	88	104	169	179	193	79	1
1950	1,146	18	67	55	46	64	85	103	168	243	219	77	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ACUTE NEPHRITIS, AND NEPHRITIS WITH EDEMA INCLUDING NEPHROSIS (590, 591)													
ALL OTHER, BOTH SEXES													
1959	425	11	29	37	28	28	70	64	59	48	34	16	1
1958	489	9	40	32	27	44	76	83	60	65	32	20	1
1957	528	15	37	27	30	57	66	71	75	91	41	16	2
1956	543	12	26	31	25	51	81	80	77	89	51	19	1
1955	570	15	33	24	26	56	73	111	93	90	37	12	-
1954	522	11	26	21	35	53	75	85	76	77	37	24	2
1953	677	17	45	30	44	69	105	107	100	93	42	24	1
1952	721	13	39	34	58	79	108	116	121	92	44	16	1
1951	834	18	55	47	55	88	115	122	117	133	60	22	2
1950	813	19	38	36	50	83	146	130	124	115	55	16	1
ALL OTHER, MALE													
1959	226	6	21	14	11	12	38	37	33	28	17	9	-
1958	256	6	18	19	15	22	28	47	31	43	17	9	1
1957	268	7	19	11	10	26	32	37	45	47	28	5	1
1956	289	5	16	23	16	25	39	36	35	58	24	11	1
1955	299	11	23	11	11	25	36	58	42	53	26	3	-
1954	278	5	17	9	19	29	36	48	43	36	21	13	2
1953	345	8	26	15	13	29	53	51	57	54	26	13	-
1952	361	10	24	17	16	32	54	61	57	58	26	6	-
1951	403	9	27	22	18	35	60	61	55	70	32	14	-
1950	439	10	27	18	24	40	69	88	62	61	30	9	1
ALL OTHER, FEMALE													
1959	199	5	8	23	17	16	32	27	26	20	17	7	1
1958	233	3	22	13	12	22	48	36	29	22	15	11	-
1957	260	8	18	16	20	31	34	34	30	44	13	11	1
1956	254	7	10	8	9	26	42	44	42	31	27	8	-
1955	271	4	10	13	15	31	37	53	51	37	11	9	-
1954	244	6	9	12	16	24	39	37	33	41	16	11	-
1953	332	9	19	15	31	40	52	56	43	39	16	11	1
1952	360	3	15	17	42	47	54	55	64	34	18	10	1
1951	431	9	28	25	37	53	55	61	62	63	28	8	2
1950	374	9	11	18	26	43	77	42	62	54	25	7	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFECTIONS OF KIDNEY (600)													
BOTH SEXES													
1959	6,822	60	21	40	89	189	386	712	1,161	1,802	1,714	646	2
1958	6,889	46	21	25	82	201	399	715	1,240	1,786	1,697	674	3
1957	6,346	52	15	36	95	195	363	689	1,151	1,609	1,478	660	3
1956	5,767	53	17	40	73	185	331	606	1,021	1,477	1,377	584	3
1955	4,911	42	22	35	65	172	320	546	823	1,249	1,175	462	-
1954	4,300	41	23	25	60	158	297	488	800	1,061	991	355	1
1953	4,052	45	18	17	70	145	284	515	699	983	920	356	-
1952	3,691	42	16	36	76	167	258	456	680	866	810	278	6
1951	3,261	36	15	32	54	144	234	410	611	789	685	246	5
1950	3,129	37	12	23	75	118	220	367	596	726	725	229	1
MALE													
1959	3,401	41	11	11	30	92	163	298	566	958	902	327	2
1958	3,544	34	9	11	29	82	166	324	631	1,000	906	350	2
1957	3,229	36	10	14	48	74	137	317	612	884	784	311	2
1956	2,963	32	6	14	26	73	127	271	522	840	759	291	2
1955	2,509	24	9	17	19	83	145	246	417	695	620	234	-
1954	2,172	23	10	10	22	68	131	213	429	595	516	154	1
1953	2,116	27	8	7	32	54	131	236	364	550	528	179	-
1952	1,916	24	5	12	37	78	105	229	367	474	442	142	1
1951	1,708	18	4	15	20	53	100	191	337	458	369	139	4
1950	1,604	26	8	8	39	39	89	174	317	398	399	107	-
FEMALE													
1959	3,421	19	10	29	59	97	223	414	595	844	812	319	-
1958	3,345	12	12	14	53	119	233	391	609	786	791	324	1
1957	3,117	16	5	22	47	121	226	372	539	725	694	349	1
1956	2,804	21	11	26	47	112	204	335	499	637	618	293	1
1955	2,402	18	13	18	46	89	175	300	406	554	555	228	-
1954	2,128	18	13	15	38	90	166	275	371	466	475	201	-
1953	1,936	18	10	10	38	91	153	279	335	433	392	177	-
1952	1,775	18	11	24	39	89	153	227	313	392	368	136	5
1951	1,553	18	11	17	34	91	134	219	274	331	316	107	1
1950	1,525	11	4	15	36	79	131	193	279	328	326	122	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFECTIONS OF KIDNEY (600)													
WHITE, BOTH SEXES													
1959	5,479	40	17	34	68	135	239	449	825	1,496	1,575	601	-
1958	5,622	29	12	21	67	136	246	478	925	1,507	1,573	625	3
1957	5,145	28	13	33	77	140	243	436	850	1,354	1,358	611	2
1956	4,765	40	14	36	57	136	216	411	765	1,251	1,279	557	3
1955	4,075	32	17	33	52	111	196	365	648	1,088	1,099	434	-
1954	3,579	33	20	21	45	113	192	327	633	935	924	336	-
1953	3,312	29	13	14	61	98	182	358	550	832	843	332	-
1952	3,078	33	12	32	65	126	166	323	545	754	753	266	3
1951	2,739	31	13	28	47	113	152	296	501	688	634	233	3
1950	2,648	31	12	20	63	92	155	265	482	642	664	221	1
WHITE, MALE													
1959	2,733	26	9	11	23	71	109	195	405	774	812	298	-
1958	2,890	19	4	9	24	56	106	215	470	834	831	320	2
1957	2,623	16	8	11	39	54	102	197	449	743	715	288	1
1956	2,450	24	4	11	19	55	87	191	389	704	691	273	2
1955	2,111	18	5	16	18	61	97	178	326	598	578	216	-
1954	1,806	17	7	8	18	56	93	143	328	515	478	143	-
1953	1,724	16	5	4	29	35	83	173	282	463	475	159	-
1952	1,597	17	4	10	33	62	72	164	284	414	402	134	1
1951	1,442	15	4	12	17	46	62	144	273	401	338	128	2
1950	1,363	22	8	7	33	32	61	132	254	355	355	104	-
WHITE, FEMALE													
1959	2,746	14	8	23	45	64	130	254	420	722	763	303	-
1958	2,732	10	8	12	43	80	140	263	455	673	742	305	1
1957	2,522	12	5	22	38	86	141	239	401	611	643	323	1
1956	2,315	16	10	25	38	81	129	220	376	547	588	284	1
1955	1,964	14	12	17	34	50	99	187	322	490	521	218	-
1954	1,773	16	13	13	27	57	99	184	305	420	446	193	-
1953	1,588	13	8	10	32	63	99	185	268	369	368	173	-
1952	1,481	16	8	22	32	64	94	159	261	340	351	132	2
1951	1,297	16	9	16	30	67	90	152	228	287	296	105	1
1950	1,285	9	4	13	30	60	94	133	228	287	309	117	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFECTIONS OF KIDNEY (600)													
ALL OTHER, BOTH SEXES													
1959	1,343	20	4	6	21	54	147	263	336	306	139	45	2
1958	1,267	17	9	4	15	65	153	237	315	279	124	49	-
1957	1,201	24	2	3	18	55	120	253	301	255	120	49	1
1956	1,002	13	3	4	16	49	115	195	256	226	98	27	-
1955	836	10	5	2	13	61	124	181	175	161	76	28	-
1954	721	8	3	4	15	45	105	161	167	126	67	19	1
1953	740	16	5	3	9	47	102	157	149	151	77	24	-
1952	613	9	4	4	11	41	92	133	135	112	57	12	3
1951	522	5	2	4	7	31	82	114	110	101	51	13	2
1950	481	6	-	3	12	26	65	102	114	84	61	8	-
ALL OTHER, MALE													
1959	668	15	2	-	7	21	54	103	161	184	90	29	2
1958	654	15	5	2	5	26	60	109	161	166	75	30	-
1957	606	20	2	3	9	20	35	120	163	141	69	23	1
1956	513	8	2	3	7	18	40	80	133	136	68	18	-
1955	398	6	4	1	1	22	48	68	91	97	42	18	-
1954	366	6	3	2	4	12	38	70	101	80	38	11	1
1953	392	11	3	3	3	19	48	63	82	87	53	20	-
1952	319	7	1	2	4	16	33	65	83	60	40	8	-
1951	266	3	-	3	3	7	38	47	64	57	31	11	2
1950	241	4	-	1	6	7	28	42	63	43	44	3	-
ALL OTHER, FEMALE													
1959	675	5	2	6	14	33	93	160	175	122	49	16	-
1958	613	2	4	2	10	39	93	128	154	113	49	19	-
1957	595	4	-	-	9	35	85	133	138	114	51	26	-
1956	489	5	1	1	9	31	75	115	123	90	30	9	-
1955	438	4	1	1	12	39	76	113	84	64	34	10	-
1954	355	2	-	2	11	33	67	91	66	46	29	8	-
1953	348	5	2	-	6	28	54	94	67	64	24	4	-
1952	294	2	3	2	7	25	59	68	52	52	17	4	3
1951	256	2	2	1	4	24	44	67	46	44	20	2	-
1950	240	2	-	2	6	19	37	60	51	41	17	5	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HYPERPLASIA OF PROSTATE (610)													
BOTH SEXES													
1959	4,583	-	1	-	-	1	5	42	365	1,306	1,906	956	1
1958	4,627	1	-	-	-	1	6	48	379	1,290	1,870	1,028	4
1957	5,097	-	-	-	-	1	6	67	442	1,512	2,035	1,033	1
1956	5,368	1	-	1	-	2	9	67	489	1,600	2,153	1,040	6
1955	6,062	-	-	-	1	1	6	70	525	1,842	2,524	1,088	5
1954	6,095	-	-	-	-	2	10	84	558	1,897	2,474	1,067	3
1953	6,173	-	-	-	-	2	4	83	560	1,856	2,575	1,086	7
1952	6,286	-	-	-	-	3	6	100	556	1,966	2,539	1,113	3
1951	6,417	-	-	-	-	1	9	82	599	1,938	2,655	1,126	7
1950	6,359	-	-	-	1	-	13	73	619	1,930	2,598	1,121	4
MALE													
1959	4,583	-	1	-	-	1	5	42	365	1,306	1,906	956	1
1958	4,627	1	-	-	-	1	6	48	379	1,290	1,870	1,028	4
1957	5,097	-	-	-	-	1	6	67	442	1,512	2,035	1,033	1
1956	5,368	1	-	1	-	2	9	67	489	1,600	2,153	1,040	6
1955	6,062	-	-	-	1	1	6	70	525	1,842	2,524	1,088	5
1954	6,095	-	-	-	-	2	10	84	558	1,897	2,474	1,067	3
1953	6,173	-	-	-	-	2	4	83	560	1,856	2,575	1,086	7
1952	6,286	-	-	-	-	3	6	100	556	1,966	2,539	1,113	3
1951	6,417	-	-	-	-	1	9	82	599	1,938	2,655	1,126	7
1950	6,359	-	-	-	1	-	13	73	619	1,930	2,598	1,121	4
FEMALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HYPERPLASIA OF PROSTATE (610)													
WHITE, BOTH SEXES													
1959	4,008	-	1	-	-	-	5	25	260	1,085	1,736	896	-
1958	4,047	-	-	-	-	-	2	24	281	1,080	1,690	969	1
1957	4,419	-	-	-	-	1	4	33	313	1,241	1,858	969	-
1956	4,741	1	-	1	-	2	6	37	360	1,374	1,990	968	2
1955	5,424	-	-	-	-	-	4	44	404	1,581	2,361	1,025	5
1954	5,472	-	-	-	-	2	3	47	431	1,667	2,311	1,010	1
1953	5,478	-	-	-	-	2	4	46	414	1,581	2,392	1,035	4
1952	5,607	-	-	-	-	3	3	49	414	1,718	2,361	1,057	2
1951	5,715	-	-	-	-	1	3	48	463	1,679	2,444	1,072	5
1950	5,645	-	-	-	1	-	6	38	469	1,660	2,401	1,067	3
WHITE, MALE													
1959	4,008	-	1	-	-	-	5	25	260	1,085	1,736	896	-
1958	4,047	-	-	-	-	-	2	24	281	1,080	1,690	969	1
1957	4,419	-	-	-	-	1	4	33	313	1,241	1,858	969	-
1956	4,741	1	-	1	-	2	6	37	360	1,374	1,990	968	2
1955	5,424	-	-	-	-	-	4	44	404	1,581	2,361	1,025	5
1954	5,472	-	-	-	-	2	3	47	431	1,667	2,311	1,010	1
1953	5,478	-	-	-	-	2	4	46	414	1,581	2,392	1,035	4
1952	5,607	-	-	-	-	3	3	49	414	1,718	2,361	1,057	2
1951	5,715	-	-	-	-	1	3	48	463	1,679	2,444	1,072	5
1950	5,645	-	-	-	1	-	6	38	469	1,660	2,401	1,067	3
WHITE, FEMALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HYPERPLASIA OF PROSTATE (610)													
ALL OTHER, BOTH SEXES													
1959	575	-	-	-	-	1	-	17	105	221	170	60	1
1958	580	1	-	-	-	1	4	24	98	210	180	59	3
1957	678	-	-	-	-	-	2	34	129	271	177	64	1
1956	627	-	-	-	-	-	3	30	129	226	163	72	4
1955	638	-	-	-	1	1	2	26	121	261	163	63	-
1954	623	-	-	-	-	-	7	37	127	230	163	57	2
1953	695	-	-	-	-	-	-	37	146	275	183	51	3
1952	679	-	-	-	-	-	3	51	142	248	178	56	1
1951	702	-	-	-	-	-	6	34	136	259	211	54	2
1950	714	-	-	-	-	-	7	35	150	270	197	54	1
ALL OTHER, MALE													
1959	575	-	-	-	-	1	-	17	105	221	170	60	1
1958	580	1	-	-	-	1	4	24	98	210	180	59	3
1957	678	-	-	-	-	-	2	34	129	271	177	64	1
1956	627	-	-	-	-	-	3	30	129	226	163	72	4
1955	638	-	-	-	1	1	2	26	121	261	163	63	-
1954	623	-	-	-	-	-	7	37	127	230	163	57	2
1953	695	-	-	-	-	-	-	37	146	275	183	51	3
1952	679	-	-	-	-	-	3	51	142	248	178	56	1
1951	702	-	-	-	-	-	6	34	136	259	211	54	2
1950	714	-	-	-	-	-	7	35	150	270	197	54	1
ALL OTHER, FEMALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)													
BOTH SEXES													
1959	1,588	...	...	7	440	673	453	13	-	-	-	-	2
1958	1,581	...	...	2	395	720	452	12	-	-	-	-	-
1957	1,746	...	...	10	424	780	497	25	8	1	1	-	-
1956	1,702	...	...	9	435	739	492	15	8	4	-	-	-
1955	1,901	...	...	4	481	834	550	29	2	-	-	1	-
1954	2,105	...	...	6	502	929	627	33	2	3	2	-	1
1953	2,385	...	...	4	645	1,012	674	34	9	3	2	-	2
1952	2,610	...	...	8	733	1,160	661	31	7	5	4	1	-
1951	2,812	...	...	11	798	1,248	722	27	3	1	1	1	-
1950	2,960	...	...	16	822	1,293	799	27	2	...	...	...	1
MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
FEMALE													
1959	1,588	...	...	7	440	673	453	13	-	-	-	-	2
1958	1,581	...	...	2	395	720	452	12	-	-	-	-	-
1957	1,746	...	...	10	424	780	497	25	8	1	1	-	-
1956	1,702	...	...	9	435	739	492	15	8	4	-	-	-
1955	1,901	...	...	4	481	834	550	29	2	-	-	1	-
1954	2,105	...	...	6	502	929	627	33	2	3	2	-	1
1953	2,385	...	...	4	645	1,012	674	34	9	3	2	-	2
1952	2,610	...	...	8	733	1,160	661	31	7	5	4	1	-
1951	2,812	...	...	11	798	1,248	722	27	3	1	1	1	-
1950	2,960	...	...	16	822	1,293	799	27	2	...	...	...	1

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)													
WHITE, BOTH SEXES													
1959	927	...	...	4	240	393	284	6	-	-	-	-	-
1958	938	...	...	-	229	421	280	8	-	-	-	-	-
1957	997	...	...	2	229	439	304	15	7	-	1	-	-
1956	1,018	...	...	3	251	434	310	11	6	3	-	-	-
1955	1,134	...	...	1	277	492	340	21	2	-	-	1	-
1954	1,280	...	...	1	299	526	424	25	2	2	1	-	-
1953	1,479	...	...	2	363	627	457	19	9	1	1	-	-
1952	1,624	...	...	-	413	742	426	27	7	5	3	1	-
1951	1,778	...	...	4	480	794	473	22	2	1	1	1	-
1950	1,873	...	...	3	488	814	546	19	2	...	...	...	1
WHITE, MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
WHITE, FEMALE													
1959	927	...	...	4	240	393	284	6	-	-	-	-	-
1958	938	...	...	-	229	421	280	8	-	-	-	-	-
1957	997	...	...	2	229	439	304	15	7	-	1	-	-
1956	1,018	...	...	3	251	434	310	11	6	3	-	-	-
1955	1,134	...	...	1	277	492	340	21	2	-	-	1	-
1954	1,280	...	...	1	299	526	424	25	2	2	1	-	-
1953	1,479	...	...	2	363	627	457	19	9	1	1	-	-
1952	1,624	...	...	-	413	742	426	27	7	5	3	1	-
1951	1,778	...	...	4	480	794	473	22	2	1	1	1	-
1950	1,873	...	...	3	488	814	546	19	2	...	...	...	1

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
DELIVERIES AND COMPLICATIONS OF PREGNANCY, CHILDBIRTH AND THE PUERPERIUM (640-689)													
ALL OTHER, BOTH SEXES													
1959	661	...	...	3	200	280	169	7	-	-	-	-	2
1958	643	...	...	2	166	299	172	4	-	-	-	-	-
1957	749	...	...	8	195	341	193	10	1	1	-	-	-
1956	684	...	...	6	184	305	182	4	2	1	-	-	-
1955	767	...	...	3	204	342	210	8	-	-	-	-	-
1954	825	...	...	5	203	403	203	8	-	1	1	-	1
1953	906	...	...	2	282	385	217	15	-	2	1	-	2
1952	986	...	...	8	320	418	235	4	-	-	1	-	-
1951	1,034	...	...	7	318	454	249	5	1	-	-	-	-
1950	1,087	...	...	13	334	479	253	8	-	...	...	...	-
ALL OTHER, MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
ALL OTHER, FEMALE													
1959	661	...	...	3	200	280	169	7	-	-	-	-	2
1958	643	...	...	2	166	299	172	4	-	-	-	-	-
1957	749	...	...	8	195	341	193	10	1	1	-	-	-
1956	684	...	...	6	184	305	182	4	2	1	-	-	-
1955	767	...	...	3	204	342	210	8	-	-	-	-	-
1954	825	...	...	5	203	403	203	8	-	1	1	-	1
1953	906	...	...	2	282	385	217	15	-	2	1	-	2
1952	986	...	...	8	320	418	235	4	-	-	1	-	-
1951	1,034	...	...	7	318	454	249	5	1	-	-	-	-
1950	1,087	...	...	13	334	479	253	8	-	...	...	...	-

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ABORTION (650-652)													
BOTH SEXES													
1959	284	...	...	1	89	134	56	4	-	-	-	-	-
1958	259	...	...	-	71	140	48	-	-	-	-	-	-
1957	260	...	...	-	73	127	59	1	-	-	-	-	-
1956	221	...	...	4	58	115	43	1	-	-	-	-	-
1955	266	...	...	-	74	120	70	2	-	-	-	-	-
1954	287	...	...	1	83	133	68	2	-	-	-	-	-
1953	294	...	...	1	99	144	48	1	-	-	1	-	-
1952	320	...	...	-	88	151	76	3	1	1	-	-	-
1951	303	...	...	1	99	131	70	2	-	-	-	-	-
1950	316	...	...	3	108	132	72	-	-	...	...	...	1
MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
FEMALE													
1959	284	...	...	1	89	134	56	4	-	-	-	-	-
1958	259	...	...	-	71	140	48	-	-	-	-	-	-
1957	260	...	...	-	73	127	59	1	-	-	-	-	-
1956	221	...	...	4	58	115	43	1	-	-	-	-	-
1955	266	...	...	-	74	120	70	2	-	-	-	-	-
1954	287	...	...	1	83	133	68	2	-	-	-	-	-
1953	294	...	...	1	99	144	48	1	-	-	1	-	-
1952	320	...	...	-	88	151	76	3	1	1	-	-	-
1951	303	...	...	1	99	131	70	2	-	-	-	-	-
1950	316	...	...	3	108	132	72	-	-	...	...	...	1

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ABORTION (650-652)													
WHITE, BOTH SEXES													
1959	138	...	...	1	38	67	32	-	-	-	-	-	-
1958	136	...	...	-	34	70	32	-	-	-	-	-	-
1957	126	...	...	-	30	67	29	-	-	-	-	-	-
1956	138	...	...	2	33	76	27	-	-	-	-	-	-
1955	150	...	...	-	37	64	47	2	-	-	-	-	-
1954	156	...	...	-	44	70	42	-	-	-	-	-	-
1953	162	...	...	-	58	80	24	-	-	-	-	-	-
1952	196	...	...	-	44	98	50	2	1	1	-	-	-
1951	170	...	...	1	51	74	43	1	-	-	-	-	-
1950	193	...	...	-	64	75	53	-	-	...	...	...	1
WHITE, MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
WHITE, FEMALE													
1959	138	...	...	1	38	67	32	-	-	-	-	-	-
1958	136	...	...	-	34	70	32	-	-	-	-	-	-
1957	126	...	...	-	30	67	29	-	-	-	-	-	-
1956	138	...	...	2	33	76	27	-	-	-	-	-	-
1955	150	...	...	-	37	64	47	2	-	-	-	-	-
1954	156	...	...	-	44	70	42	-	-	-	-	-	-
1953	162	...	...	-	58	80	24	-	-	-	-	-	-
1952	196	...	...	-	44	98	50	2	1	1	-	-	-
1951	170	...	...	1	51	74	43	1	-	-	-	-	-
1950	193	...	...	-	64	75	53	-	-	...	...	...	1

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ABORTION (650-652)													
ALL OTHER, BOTH SEXES													
1959	146	...	...	-	51	67	24	4	-	-	-	-	-
1958	123	...	...	-	37	70	16	-	-	-	-	-	-
1957	134	...	...	-	43	60	30	1	-	-	-	-	-
1956	83	...	...	2	25	39	16	1	-	-	-	-	-
1955	116	...	...	-	37	56	23	-	-	-	-	-	-
1954	131	...	...	1	39	63	26	2	-	-	-	-	-
1953	132	...	...	1	41	64	24	1	-	-	1	-	-
1952	124	...	...	-	44	53	26	1	-	-	-	-	-
1951	133	...	...	-	48	57	27	1	-	-	-	-	-
1950	123	...	...	3	44	57	19	-	-	...	...	...	-
ALL OTHER, MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
ALL OTHER, FEMALE													
1959	146	...	...	-	51	67	24	4	-	-	-	-	-
1958	123	...	...	-	37	70	16	-	-	-	-	-	-
1957	134	...	...	-	43	60	30	1	-	-	-	-	-
1956	83	...	...	2	25	39	16	1	-	-	-	-	-
1955	116	...	...	-	37	56	23	-	-	-	-	-	-
1954	131	...	...	1	39	63	26	2	-	-	-	-	-
1953	132	...	...	1	41	64	24	1	-	-	1	-	-
1952	124	...	...	-	44	53	26	1	-	-	-	-	-
1951	133	...	...	-	48	57	27	1	-	-	-	-	-
1950	123	...	...	3	44	57	19	-	-	...	...	...	-

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)													
BOTH SEXES													
1959	1,304	...	...	6	351	539	397	9	-	-	-	-	2
1958	1,322	...	...	2	324	580	404	12	-	-	-	-	-
1957	1,486	...	...	10	351	653	438	24	8	1	1	-	-
1956	1,481	...	...	5	377	624	449	14	8	4	-	-	-
1955	1,635	...	...	4	407	714	480	27	2	-	-	1	-
1954	1,818	...	...	5	419	796	559	31	2	3	2	-	1
1953	2,091	...	...	3	546	868	626	33	9	3	1	-	2
1952	2,290	...	...	8	645	1,009	585	28	6	4	4	1	-
1951	2,509	...	...	10	699	1,117	652	25	3	1	1	1	-
1950	2,644	...	...	13	714	1,161	727	27	2	...	...	...	-
MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
FEMALE													
1959	1,304	...	...	6	351	539	397	9	-	-	-	-	2
1958	1,322	...	...	2	324	580	404	12	-	-	-	-	-
1957	1,486	...	...	10	351	653	438	24	8	1	1	-	-
1956	1,481	...	...	5	377	624	449	14	8	4	-	-	-
1955	1,635	...	...	4	407	714	480	27	2	-	-	1	-
1954	1,818	...	...	5	419	796	559	31	2	3	2	-	1
1953	2,091	...	...	3	546	868	626	33	9	3	1	-	2
1952	2,290	...	...	8	645	1,009	585	28	6	4	4	1	-
1951	2,509	...	...	10	699	1,117	652	25	3	1	1	1	-
1950	2,644	...	...	13	714	1,161	727	27	2	...	...	...	-

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)													
WHITE, BOTH SEXES													
1959	789	...	...	3	202	326	252	6	-	-	-	-	-
1958	802	...	...	-	195	351	248	8	-	-	-	-	-
1957	871	...	...	2	199	372	275	15	7	-	1	-	-
1956	880	...	...	1	218	358	283	11	6	3	-	-	-
1955	984	...	...	1	240	428	293	19	2	-	-	1	-
1954	1,124	...	...	1	255	456	382	25	2	2	1	-	-
1953	1,317	...	...	2	305	547	433	19	9	1	1	-	-
1952	1,428	...	...	-	369	644	376	25	6	4	3	1	-
1951	1,608	...	...	3	429	720	430	21	2	1	1	1	-
1950	1,680	...	...	3	424	739	493	19	2	...	...	...	-
WHITE, MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
WHITE, FEMALE													
1959	789	...	...	3	202	326	252	6	-	-	-	-	-
1958	802	...	...	-	195	351	248	8	-	-	-	-	-
1957	871	...	...	2	199	372	275	15	7	-	1	-	-
1956	880	...	...	1	218	358	283	11	6	3	-	-	-
1955	984	...	...	1	240	428	293	19	2	-	-	1	-
1954	1,124	...	...	1	255	456	382	25	2	2	1	-	-
1953	1,317	...	...	2	305	547	433	19	9	1	1	-	-
1952	1,428	...	...	-	369	644	376	25	6	4	3	1	-
1951	1,608	...	...	3	429	720	430	21	2	1	1	1	-
1950	1,680	...	...	3	424	739	493	19	2	...	...	...	-

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND THE PUERPERIUM (640-649, 660-689)													
ALL OTHER, BOTH SEXES													
1959	515	...	...	3	149	213	145	3	-	-	-	-	2
1958	520	...	...	2	129	229	156	4	-	-	-	-	-
1957	615	...	...	8	152	281	163	9	1	1	-	-	-
1956	601	...	...	4	159	266	166	3	2	1	-	-	-
1955	651	...	...	3	167	286	187	8	-	-	-	-	-
1954	694	...	...	4	164	340	177	6	-	1	1	-	1
1953	774	...	...	1	241	321	193	14	-	2	-	-	2
1952	862	...	...	8	276	365	209	3	-	-	1	-	-
1951	901	...	...	7	270	397	222	4	1	-	-	-	-
1950	964	...	...	10	290	422	234	8	-	...	...	...	-
ALL OTHER, MALE													
1959	...	...	...	...	...	...	...	...	...	...	...	...	...
1958	...	...	...	...	...	...	...	...	...	...	...	...	...
1957	...	...	...	...	...	...	...	...	...	...	...	...	...
1956	...	...	...	...	...	...	...	...	...	...	...	...	...
1955	...	...	...	...	...	...	...	...	...	...	...	...	...
1954	...	...	...	...	...	...	...	...	...	...	...	...	...
1953	...	...	...	...	...	...	...	...	...	...	...	...	...
1952	...	...	...	...	...	...	...	...	...	...	...	...	...
1951	...	...	...	...	...	...	...	...	...	...	...	...	...
1950	...	...	...	...	...	...	...	...	...	...	...	...	...
ALL OTHER, FEMALE													
1959	515	...	...	3	149	213	145	3	-	-	-	-	2
1958	520	...	...	2	129	229	156	4	-	-	-	-	-
1957	615	...	...	8	152	281	163	9	1	1	-	-	-
1956	601	...	...	4	159	266	166	3	2	1	-	-	-
1955	651	...	...	3	167	286	187	8	-	-	-	-	-
1954	694	...	...	4	164	340	177	6	-	1	1	-	1
1953	774	...	...	1	241	321	193	14	-	2	-	-	2
1952	862	...	...	8	276	365	209	3	-	-	1	-	-
1951	901	...	...	7	270	397	222	4	1	-	-	-	-
1950	964	...	...	10	290	422	234	8	-	...	...	...	-

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CONGENITAL MALFORMATIONS (750-759)													
BOTH SEXES													
1959	21,780	15,579	1,947	1,310	655	436	531	520	433	237	108	22	2
1958	21,411	15,555	1,979	1,114	559	411	484	523	389	250	117	29	1
1957	21,818	15,801	2,088	1,079	497	427	536	538	438	263	117	26	8
1956	21,065	15,676	1,978	886	428	375	457	514	382	245	89	27	8
1955	20,502	15,356	1,782	819	433	407	445	502	367	250	109	20	12
1954	20,081	15,116	1,767	794	387	365	419	500	345	229	136	16	7
1953	20,012	15,166	1,799	672	420	340	423	453	388	218	108	21	4
1952	19,768	15,185	1,743	651	393	346	397	435	324	179	100	14	1
1951	18,865	14,721	1,579	598	375	331	360	381	270	166	73	7	4
1950	18,425	14,074	1,439	575	396	446	474	457	312	171	67	11	3
MALE													
1959	11,896	8,569	997	729	345	240	305	266	247	125	59	13	1
1958	11,584	8,458	1,007	598	324	217	271	287	214	133	59	16	-
1957	11,811	8,651	1,064	545	281	225	283	296	235	143	68	14	6
1956	11,463	8,576	1,033	479	244	197	240	284	195	143	53	16	3
1955	11,130	8,392	927	435	255	208	222	291	189	141	59	6	5
1954	10,969	8,327	890	433	218	200	219	292	189	129	64	3	5
1953	10,952	8,354	918	363	242	177	243	248	216	118	57	15	1
1952	10,775	8,303	896	341	222	200	231	245	175	91	62	9	-
1951	10,180	7,975	782	319	211	178	196	211	161	103	40	4	-
1950	10,026	7,648	769	326	214	238	267	238	192	95	31	6	2
FEMALE													
1959	9,884	7,010	950	581	310	196	226	254	186	112	49	9	1
1958	9,827	7,097	972	516	235	194	213	236	175	117	58	13	1
1957	10,007	7,150	1,024	534	216	202	253	242	203	120	49	12	2
1956	9,602	7,100	945	407	184	178	217	230	187	102	36	11	5
1955	9,372	6,964	855	384	178	199	223	211	178	109	50	14	7
1954	9,112	6,789	877	361	169	165	200	208	156	100	72	13	2
1953	9,060	6,812	881	309	178	163	180	205	172	100	51	6	3
1952	8,993	6,882	847	310	171	146	166	190	149	88	38	5	1
1951	8,685	6,746	797	279	164	153	164	170	109	63	33	3	4
1950	8,399	6,426	670	249	182	208	207	219	120	76	36	5	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CONGENITAL MALFORMATIONS (750-759)													
WHITE, BOTH SEXES													
1959	19,060	13,594	1,642	1,143	588	390	485	475	401	217	103	20	2
1958	18,653	13,512	1,663	998	504	361	431	479	342	227	109	26	1
1957	19,141	13,855	1,771	951	442	370	484	484	395	243	113	26	7
1956	18,610	13,829	1,692	794	384	333	401	477	353	230	85	24	8
1955	18,122	13,540	1,516	751	388	359	402	451	346	235	105	19	10
1954	17,929	13,507	1,541	700	347	329	375	437	327	214	131	16	5
1953	17,955	13,638	1,564	592	363	307	383	420	357	207	103	18	3
1952	17,819	13,709	1,528	577	348	308	359	402	301	173	99	14	1
1951	17,039	13,295	1,393	521	345	307	333	350	257	156	71	7	4
1950	16,621	12,714	1,277	507	351	408	420	416	294	158	64	10	2
WHITE, MALE													
1959	10,428	7,497	838	640	310	213	278	241	229	114	56	11	1
1958	10,070	7,327	842	533	297	192	243	267	183	115	57	14	-
1957	10,347	7,588	886	479	251	194	259	271	206	128	66	14	5
1956	10,105	7,558	869	428	216	177	214	264	179	132	50	15	3
1955	9,839	7,409	790	395	230	183	199	262	174	130	57	6	4
1954	9,762	7,433	771	380	198	180	190	247	178	117	61	3	4
1953	9,780	7,484	788	315	211	159	220	227	199	110	53	13	1
1952	9,667	7,449	780	304	196	183	213	226	159	86	62	9	-
1951	9,138	7,170	676	268	194	162	182	193	153	97	39	4	-
1950	9,042	6,907	676	286	196	222	235	219	179	86	29	5	2
WHITE, FEMALE													
1959	8,632	6,097	804	503	278	177	207	234	172	103	47	9	1
1958	8,583	6,185	821	465	207	169	188	212	159	112	52	12	1
1957	8,794	6,267	885	472	191	176	225	213	189	115	47	12	2
1956	8,505	6,271	823	366	168	156	187	213	174	98	35	9	5
1955	8,283	6,131	726	356	158	176	203	189	172	105	48	13	6
1954	8,167	6,074	770	320	149	149	185	190	149	97	70	13	1
1953	8,175	6,154	776	277	152	148	163	193	158	97	50	5	2
1952	8,152	6,260	748	273	152	125	146	176	142	87	37	5	1
1951	7,901	6,125	717	253	151	145	151	157	104	59	32	3	4
1950	7,579	5,807	601	221	155	186	185	197	115	72	35	5	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CONGENITAL MALFORMATIONS (750-759)													
ALL OTHER, BOTH SEXES													
1959	2,720	1,985	305	167	67	46	46	45	32	20	5	2	-
1958	2,758	2,043	316	116	55	50	53	44	47	23	8	3	-
1957	2,677	1,946	317	128	55	57	52	54	43	20	4	-	1
1956	2,455	1,847	286	92	44	42	56	37	29	15	4	3	-
1955	2,380	1,816	266	68	45	48	43	51	21	15	4	1	2
1954	2,152	1,609	226	94	40	36	44	63	18	15	5	-	2
1953	2,057	1,528	235	80	57	33	40	33	31	11	5	3	1
1952	1,949	1,476	215	74	45	38	38	33	23	6	1	-	-
1951	1,826	1,426	186	77	30	24	27	31	13	10	2	-	-
1950	1,804	1,360	162	68	45	38	54	41	18	13	3	1	1
ALL OTHER, MALE													
1959	1,468	1,072	159	89	35	27	27	25	18	11	3	2	-
1958	1,514	1,131	165	65	27	25	28	20	31	18	2	2	-
1957	1,464	1,063	178	66	30	31	24	25	29	15	2	-	1
1956	1,358	1,018	164	51	28	20	26	20	16	11	3	1	-
1955	1,291	983	137	40	25	25	23	29	15	11	2	-	1
1954	1,207	894	119	53	20	20	29	45	11	12	3	-	1
1953	1,172	870	130	48	31	18	23	21	17	8	4	2	-
1952	1,108	854	116	37	26	17	18	19	16	5	-	-	-
1951	1,042	805	106	51	17	16	14	18	8	6	1	-	-
1950	984	741	93	40	18	16	32	19	13	9	2	1	-
ALL OTHER, FEMALE													
1959	1,252	913	146	78	32	19	19	20	14	9	2	-	-
1958	1,244	912	151	51	28	25	25	24	16	5	6	1	-
1957	1,213	883	139	62	25	26	28	29	14	5	2	-	-
1956	1,097	829	122	41	16	22	30	17	13	4	1	2	-
1955	1,089	833	129	28	20	23	20	22	6	4	2	1	1
1954	945	715	107	41	20	16	15	18	7	3	2	-	1
1953	885	658	105	32	26	15	17	12	14	3	1	1	1
1952	841	622	99	37	19	21	20	14	7	1	1	-	-
1951	784	621	80	26	13	8	13	13	5	4	1	-	-
1950	820	619	69	28	27	22	22	22	5	4	1	-	1

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CERTAIN DISEASES OF EARLY INFANCY (760-776)													
BOTH SEXES													
1959	67,934	67,916	9	9	-	-	-	-	-	-	-	-	-
1958	68,960	68,939	9	8	2	1	-	-	-	-	-	-	1
1957	66,569	66,552	13	3	1	-	-	-	-	-	-	-	-
1956	64,546	64,515	19	4	-	-	-	-	-	-	-	-	8
1955	64,043	64,016	23	4	-	-	-	-	-	-	-	-	-
1954	63,486	63,450	30	6	-	-	-	-	-	-	-	-	-
1953	63,444	63,423	16	4	-	-	-	-	-	-	-	-	1
1952	63,659	63,632	24	3	-	-	-	-	-	-	-	-	-
1951	63,096	63,044	36	6	-	-	-	1	-	-	-	-	9
1950	60,989	60,967	19	3	-	-	-	-	-	-	-	-	-
MALE													
1959	39,701	39,688	7	6	-	-	-	-	-	-	-	-	-
1958	40,042	40,029	6	5	1	-	-	-	-	-	-	-	1
1957	38,777	38,771	5	-	1	-	-	-	-	-	-	-	-
1956	37,880	37,863	14	-	-	-	-	-	-	-	-	-	3
1955	37,403	37,389	12	2	-	-	-	-	-	-	-	-	-
1954	37,136	37,115	15	6	-	-	-	-	-	-	-	-	-
1953	37,212	37,201	9	1	-	-	-	-	-	-	-	-	1
1952	37,247	37,233	11	3	-	-	-	-	-	-	-	-	-
1951	37,183	37,157	16	3	-	-	-	-	-	-	-	-	7
1950	35,807	35,791	15	1	-	-	-	-	-	-	-	-	-
FEMALE													
1959	28,233	28,228	2	3	-	-	-	-	-	-	-	-	-
1958	28,918	28,910	3	3	1	1	-	-	-	-	-	-	-
1957	27,792	27,781	8	3	-	-	-	-	-	-	-	-	-
1956	26,666	26,652	5	4	-	-	-	-	-	-	-	-	5
1955	26,640	26,627	11	2	-	-	-	-	-	-	-	-	-
1954	26,350	26,335	15	-	-	-	-	-	-	-	-	-	-
1953	26,232	26,222	7	3	-	-	-	-	-	-	-	-	-
1952	26,412	26,399	13	-	-	-	-	-	-	-	-	-	-
1951	25,913	25,887	20	3	-	-	-	1	-	-	-	-	2
1950	25,182	25,176	4	2	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CERTAIN DISEASES OF EARLY INFANCY (760-776)													
WHITE, BOTH SEXES													
1959	51,797	51,780	8	9	-	-	-	-	-	-	-	-	-
1958	52,614	52,594	9	8	1	1	-	-	-	-	-	-	1
1957	51,140	51,125	11	3	1	-	-	-	-	-	-	-	-
1956	49,975	49,947	18	3	-	-	-	-	-	-	-	-	7
1955	50,074	50,051	19	4	-	-	-	-	-	-	-	-	-
1954	49,992	49,959	27	6	-	-	-	-	-	-	-	-	-
1953	50,448	50,427	16	4	-	-	-	-	-	-	-	-	1
1952	50,956	50,932	21	3	-	-	-	-	-	-	-	-	-
1951	50,977	50,934	29	6	-	-	-	1	-	-	-	-	7
1950	49,412	49,391	18	3	-	-	-	-	-	-	-	-	-
WHITE, MALE													
1959	30,640	30,627	7	6	-	-	-	-	-	-	-	-	-
1958	30,928	30,915	6	5	1	-	-	-	-	-	-	-	1
1957	30,145	30,141	3	-	1	-	-	-	-	-	-	-	-
1956	29,546	29,530	13	-	-	-	-	-	-	-	-	-	3
1955	29,558	29,546	10	2	-	-	-	-	-	-	-	-	-
1954	29,557	29,537	14	6	-	-	-	-	-	-	-	-	-
1953	30,024	30,013	9	1	-	-	-	-	-	-	-	-	1
1952	30,102	30,088	11	3	-	-	-	-	-	-	-	-	-
1951	30,266	30,245	13	3	-	-	-	-	-	-	-	-	5
1950	29,326	29,311	14	1	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1959	21,157	21,153	1	3	-	-	-	-	-	-	-	-	-
1958	21,686	21,679	3	3	-	1	-	-	-	-	-	-	-
1957	20,995	20,984	8	3	-	-	-	-	-	-	-	-	-
1956	20,429	20,417	5	3	-	-	-	-	-	-	-	-	4
1955	20,516	20,505	9	2	-	-	-	-	-	-	-	-	-
1954	20,435	20,422	13	-	-	-	-	-	-	-	-	-	-
1953	20,424	20,414	7	3	-	-	-	-	-	-	-	-	-
1952	20,854	20,844	10	-	-	-	-	-	-	-	-	-	-
1951	20,711	20,689	16	3	-	-	-	1	-	-	-	-	2
1950	20,086	20,080	4	2	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
CERTAIN DISEASES OF EARLY INFANCY (760-776)													
ALL OTHER, BOTH SEXES													
1959	16,137	16,136	1	-	-	-	-	-	-	-	-	-	-
1958	16,346	16,345	-	-	1	-	-	-	-	-	-	-	-
1957	15,429	15,427	2	-	-	-	-	-	-	-	-	-	-
1956	14,571	14,568	1	1	-	-	-	-	-	-	-	-	1
1955	13,969	13,965	4	-	-	-	-	-	-	-	-	-	-
1954	13,494	13,491	3	-	-	-	-	-	-	-	-	-	-
1953	12,996	12,996	-	-	-	-	-	-	-	-	-	-	-
1952	12,703	12,700	3	-	-	-	-	-	-	-	-	-	-
1951	12,119	12,110	7	-	-	-	-	-	-	-	-	-	2
1950	11,577	11,576	1	-	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1959	9,061	9,061	-	-	-	-	-	-	-	-	-	-	-
1958	9,114	9,114	-	-	-	-	-	-	-	-	-	-	-
1957	8,632	8,630	2	-	-	-	-	-	-	-	-	-	-
1956	8,334	8,333	1	-	-	-	-	-	-	-	-	-	-
1955	7,845	7,843	2	-	-	-	-	-	-	-	-	-	-
1954	7,579	7,578	1	-	-	-	-	-	-	-	-	-	-
1953	7,188	7,188	-	-	-	-	-	-	-	-	-	-	-
1952	7,145	7,145	-	-	-	-	-	-	-	-	-	-	-
1951	6,917	6,912	3	-	-	-	-	-	-	-	-	-	2
1950	6,481	6,480	1	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	7,076	7,075	1	-	-	-	-	-	-	-	-	-	-
1958	7,232	7,231	-	-	1	-	-	-	-	-	-	-	-
1957	6,797	6,797	-	-	-	-	-	-	-	-	-	-	-
1956	6,237	6,235	-	1	-	-	-	-	-	-	-	-	1
1955	6,124	6,122	2	-	-	-	-	-	-	-	-	-	-
1954	5,915	5,913	2	-	-	-	-	-	-	-	-	-	-
1953	5,808	5,808	-	-	-	-	-	-	-	-	-	-	-
1952	5,558	5,555	3	-	-	-	-	-	-	-	-	-	-
1951	5,202	5,198	4	-	-	-	-	-	-	-	-	-	-
1950	5,096	5,096	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)</b>													
<b>BOTH SEXES</b>													
1959	30,249	30,246	2	1	-	-	-	-	-	-	-	-	-
1958	31,525	31,524	-	-	1	-	-	-	-	-	-	-	-
1957	30,273	30,273	-	-	-	-	-	-	-	-	-	-	-
1956	29,599	29,591	2	2	-	-	-	-	-	-	-	-	4
1955	29,255	29,245	10	-	-	-	-	-	-	-	-	-	-
1954	28,642	28,632	10	-	-	-	-	-	-	-	-	-	-
1953	28,263	28,260	2	-	-	-	-	-	-	-	-	-	1
1952	27,357	27,354	3	-	-	-	-	-	-	-	-	-	-
1951	26,962	26,956	1	-	-	-	-	-	-	-	-	-	5
1950	25,076	25,075	1	-	-	-	-	-	-	-	-	-	-
<b>MALE</b>													
1959	18,084	18,081	2	1	-	-	-	-	-	-	-	-	-
1958	18,718	18,718	-	-	-	-	-	-	-	-	-	-	-
1957	18,137	18,137	-	-	-	-	-	-	-	-	-	-	-
1956	17,866	17,863	2	-	-	-	-	-	-	-	-	-	1
1955	17,550	17,548	2	-	-	-	-	-	-	-	-	-	-
1954	17,283	17,279	4	-	-	-	-	-	-	-	-	-	-
1953	17,133	17,131	1	-	-	-	-	-	-	-	-	-	1
1952	16,590	16,588	2	-	-	-	-	-	-	-	-	-	-
1951	16,468	16,462	1	-	-	-	-	-	-	-	-	-	5
1950	15,237	15,236	1	-	-	-	-	-	-	-	-	-	-
<b>FEMALE</b>													
1959	12,165	12,165	-	-	-	-	-	-	-	-	-	-	-
1958	12,807	12,806	-	-	1	-	-	-	-	-	-	-	-
1957	12,136	12,136	-	-	-	-	-	-	-	-	-	-	-
1956	11,733	11,728	-	2	-	-	-	-	-	-	-	-	3
1955	11,705	11,697	8	-	-	-	-	-	-	-	-	-	-
1954	11,359	11,353	6	-	-	-	-	-	-	-	-	-	-
1953	11,130	11,129	1	-	-	-	-	-	-	-	-	-	-
1952	10,767	10,766	1	-	-	-	-	-	-	-	-	-	-
1951	10,494	10,494	-	-	-	-	-	-	-	-	-	-	-
1950	9,839	9,839	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	24,230	24,227	2	1	-	-	-	-	-	-	-	-	-
1958	25,247	25,247	-	-	-	-	-	-	-	-	-	-	-
1957	24,219	24,219	-	-	-	-	-	-	-	-	-	-	-
1956	23,944	23,937	2	1	-	-	-	-	-	-	-	-	4
1955	23,913	23,906	7	-	-	-	-	-	-	-	-	-	-
1954	23,645	23,636	9	-	-	-	-	-	-	-	-	-	-
1953	23,546	23,543	2	-	-	-	-	-	-	-	-	-	1
1952	22,940	22,938	2	-	-	-	-	-	-	-	-	-	-
1951	22,867	22,862	1	-	-	-	-	-	-	-	-	-	4
1950	21,346	21,345	1	-	-	-	-	-	-	-	-	-	-
<b>WHITE, MALE</b>													
1959	14,656	14,653	2	1	-	-	-	-	-	-	-	-	-
1958	15,163	15,163	-	-	-	-	-	-	-	-	-	-	-
1957	14,634	14,634	-	-	-	-	-	-	-	-	-	-	-
1956	14,571	14,568	2	-	-	-	-	-	-	-	-	-	1
1955	14,527	14,526	1	-	-	-	-	-	-	-	-	-	-
1954	14,432	14,428	4	-	-	-	-	-	-	-	-	-	-
1953	14,467	14,465	1	-	-	-	-	-	-	-	-	-	1
1952	14,021	14,019	2	-	-	-	-	-	-	-	-	-	-
1951	14,071	14,066	1	-	-	-	-	-	-	-	-	-	4
1950	13,083	13,082	1	-	-	-	-	-	-	-	-	-	-
<b>WHITE, FEMALE</b>													
1959	9,574	9,574	-	-	-	-	-	-	-	-	-	-	-
1958	10,084	10,084	-	-	-	-	-	-	-	-	-	-	-
1957	9,585	9,585	-	-	-	-	-	-	-	-	-	-	-
1956	9,373	9,369	-	1	-	-	-	-	-	-	-	-	3
1955	9,386	9,380	6	-	-	-	-	-	-	-	-	-	-
1954	9,213	9,208	5	-	-	-	-	-	-	-	-	-	-
1953	9,079	9,078	1	-	-	-	-	-	-	-	-	-	-
1952	8,919	8,919	-	-	-	-	-	-	-	-	-	-	-
1951	8,796	8,796	-	-	-	-	-	-	-	-	-	-	-
1950	8,263	8,263	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
BIRTH INJURIES, POSTNATAL ASPHYXIA, AND ATELECTASIS (760-762)													
ALL OTHER, BOTH SEXES													
1959	6,019	6,019	-	-	-	-	-	-	-	-	-	-	-
1958	6,278	6,277	-	-	1	-	-	-	-	-	-	-	-
1957	6,054	6,054	-	-	-	-	-	-	-	-	-	-	-
1956	5,655	5,654	-	1	-	-	-	-	-	-	-	-	-
1955	5,342	5,339	3	-	-	-	-	-	-	-	-	-	-
1954	4,997	4,996	1	-	-	-	-	-	-	-	-	-	-
1953	4,717	4,717	-	-	-	-	-	-	-	-	-	-	-
1952	4,417	4,416	1	-	-	-	-	-	-	-	-	-	-
1951	4,095	4,094	-	-	-	-	-	-	-	-	-	-	1
1950	3,730	3,730	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1959	3,428	3,428	-	-	-	-	-	-	-	-	-	-	-
1958	3,555	3,555	-	-	-	-	-	-	-	-	-	-	-
1957	3,503	3,503	-	-	-	-	-	-	-	-	-	-	-
1956	3,295	3,295	-	-	-	-	-	-	-	-	-	-	-
1955	3,023	3,022	1	-	-	-	-	-	-	-	-	-	-
1954	2,851	2,851	-	-	-	-	-	-	-	-	-	-	-
1953	2,666	2,666	-	-	-	-	-	-	-	-	-	-	-
1952	2,569	2,569	-	-	-	-	-	-	-	-	-	-	-
1951	2,397	2,396	-	-	-	-	-	-	-	-	-	-	1
1950	2,154	2,154	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	2,591	2,591	-	-	-	-	-	-	-	-	-	-	-
1958	2,723	2,722	-	-	1	-	-	-	-	-	-	-	-
1957	2,551	2,551	-	-	-	-	-	-	-	-	-	-	-
1956	2,360	2,359	-	1	-	-	-	-	-	-	-	-	-
1955	2,319	2,317	2	-	-	-	-	-	-	-	-	-	-
1954	2,146	2,145	1	-	-	-	-	-	-	-	-	-	-
1953	2,051	2,051	-	-	-	-	-	-	-	-	-	-	-
1952	1,848	1,847	1	-	-	-	-	-	-	-	-	-	-
1951	1,698	1,698	-	-	-	-	-	-	-	-	-	-	-
1950	1,576	1,576	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFECTIONS OF NEWBORN (763-768)													
BOTH SEXES													
1959	5,064	5,064	-	-	-	-	-	-	-	-	-	-	-
1958	5,531	5,530	1	-	-	-	-	-	-	-	-	-	-
1957	5,087	5,087	-	-	-	-	-	-	-	-	-	-	-
1956	4,613	4,613	-	-	-	-	-	-	-	-	-	-	-
1955	4,201	4,200	1	-	-	-	-	-	-	-	-	-	-
1954	4,361	4,359	2	-	-	-	-	-	-	-	-	-	-
1953	4,187	4,187	-	-	-	-	-	-	-	-	-	-	-
1952	4,094	4,094	-	-	-	-	-	-	-	-	-	-	-
1951	3,927	3,926	-	-	-	-	-	-	-	-	-	-	1
1950	3,846	3,846	-	-	-	-	-	-	-	-	-	-	-
MALE													
1959	2,979	2,979	-	-	-	-	-	-	-	-	-	-	-
1958	3,217	3,216	1	-	-	-	-	-	-	-	-	-	-
1957	2,979	2,979	-	-	-	-	-	-	-	-	-	-	-
1956	2,686	2,686	-	-	-	-	-	-	-	-	-	-	-
1955	2,495	2,495	-	-	-	-	-	-	-	-	-	-	-
1954	2,544	2,543	1	-	-	-	-	-	-	-	-	-	-
1953	2,427	2,427	-	-	-	-	-	-	-	-	-	-	-
1952	2,391	2,391	-	-	-	-	-	-	-	-	-	-	-
1951	2,288	2,287	-	-	-	-	-	-	-	-	-	-	1
1950	2,207	2,207	-	-	-	-	-	-	-	-	-	-	-
FEMALE													
1959	2,085	2,085	-	-	-	-	-	-	-	-	-	-	-
1958	2,314	2,314	-	-	-	-	-	-	-	-	-	-	-
1957	2,108	2,108	-	-	-	-	-	-	-	-	-	-	-
1956	1,927	1,927	-	-	-	-	-	-	-	-	-	-	-
1955	1,706	1,705	1	-	-	-	-	-	-	-	-	-	-
1954	1,817	1,816	1	-	-	-	-	-	-	-	-	-	-
1953	1,760	1,760	-	-	-	-	-	-	-	-	-	-	-
1952	1,703	1,703	-	-	-	-	-	-	-	-	-	-	-
1951	1,639	1,639	-	-	-	-	-	-	-	-	-	-	-
1950	1,639	1,639	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFECTIONS OF NEWBORN (763-768)													
WHITE, BOTH SEXES													
1959	3,188	3,188	-	-	-	-	-	-	-	-	-	-	-
1958	3,560	3,559	1	-	-	-	-	-	-	-	-	-	-
1957	3,307	3,307	-	-	-	-	-	-	-	-	-	-	-
1956	3,012	3,012	-	-	-	-	-	-	-	-	-	-	-
1955	2,748	2,747	1	-	-	-	-	-	-	-	-	-	-
1954	2,937	2,935	2	-	-	-	-	-	-	-	-	-	-
1953	2,818	2,818	-	-	-	-	-	-	-	-	-	-	-
1952	2,800	2,800	-	-	-	-	-	-	-	-	-	-	-
1951	2,787	2,787	-	-	-	-	-	-	-	-	-	-	-
1950	2,777	2,777	-	-	-	-	-	-	-	-	-	-	-
WHITE, MALE													
1959	1,886	1,886	-	-	-	-	-	-	-	-	-	-	-
1958	2,081	2,080	1	-	-	-	-	-	-	-	-	-	-
1957	1,997	1,997	-	-	-	-	-	-	-	-	-	-	-
1956	1,763	1,763	-	-	-	-	-	-	-	-	-	-	-
1955	1,655	1,655	-	-	-	-	-	-	-	-	-	-	-
1954	1,744	1,743	1	-	-	-	-	-	-	-	-	-	-
1953	1,662	1,662	-	-	-	-	-	-	-	-	-	-	-
1952	1,641	1,641	-	-	-	-	-	-	-	-	-	-	-
1951	1,629	1,629	-	-	-	-	-	-	-	-	-	-	-
1950	1,624	1,624	-	-	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1959	1,302	1,302	-	-	-	-	-	-	-	-	-	-	-
1958	1,479	1,479	-	-	-	-	-	-	-	-	-	-	-
1957	1,310	1,310	-	-	-	-	-	-	-	-	-	-	-
1956	1,249	1,249	-	-	-	-	-	-	-	-	-	-	-
1955	1,093	1,092	1	-	-	-	-	-	-	-	-	-	-
1954	1,193	1,192	1	-	-	-	-	-	-	-	-	-	-
1953	1,156	1,156	-	-	-	-	-	-	-	-	-	-	-
1952	1,159	1,159	-	-	-	-	-	-	-	-	-	-	-
1951	1,158	1,158	-	-	-	-	-	-	-	-	-	-	-
1950	1,153	1,153	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
INFECTIONS OF NEWBORN (763-768)													
ALL OTHER, BOTH SEXES													
1959	1,876	1,876	-	-	-	-	-	-	-	-	-	-	-
1958	1,971	1,971	-	-	-	-	-	-	-	-	-	-	-
1957	1,780	1,780	-	-	-	-	-	-	-	-	-	-	-
1956	1,601	1,601	-	-	-	-	-	-	-	-	-	-	-
1955	1,453	1,453	-	-	-	-	-	-	-	-	-	-	-
1954	1,424	1,424	-	-	-	-	-	-	-	-	-	-	-
1953	1,369	1,369	-	-	-	-	-	-	-	-	-	-	-
1952	1,294	1,294	-	-	-	-	-	-	-	-	-	-	-
1951	1,140	1,139	-	-	-	-	-	-	-	-	-	-	1
1950	1,069	1,069	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1959	1,093	1,093	-	-	-	-	-	-	-	-	-	-	-
1958	1,136	1,136	-	-	-	-	-	-	-	-	-	-	-
1957	982	982	-	-	-	-	-	-	-	-	-	-	-
1956	923	923	-	-	-	-	-	-	-	-	-	-	-
1955	840	840	-	-	-	-	-	-	-	-	-	-	-
1954	800	800	-	-	-	-	-	-	-	-	-	-	-
1953	765	765	-	-	-	-	-	-	-	-	-	-	-
1952	750	750	-	-	-	-	-	-	-	-	-	-	-
1951	659	658	-	-	-	-	-	-	-	-	-	-	1
1950	583	583	-	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	783	783	-	-	-	-	-	-	-	-	-	-	-
1958	835	835	-	-	-	-	-	-	-	-	-	-	-
1957	798	798	-	-	-	-	-	-	-	-	-	-	-
1956	678	678	-	-	-	-	-	-	-	-	-	-	-
1955	613	613	-	-	-	-	-	-	-	-	-	-	-
1954	624	624	-	-	-	-	-	-	-	-	-	-	-
1953	604	604	-	-	-	-	-	-	-	-	-	-	-
1952	544	544	-	-	-	-	-	-	-	-	-	-	-
1951	481	481	-	-	-	-	-	-	-	-	-	-	-
1950	486	486	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURITY UNQUALIFIED (769-776)													
BOTH SEXES													
1959	32,621	32,606	7	8	-	-	-	-	-	-	-	-	-
1958	31,904	31,885	8	8	1	1	-	-	-	-	-	-	1
1957	31,209	31,192	13	3	1	-	-	-	-	-	-	-	-
1956	30,334	30,311	17	2	-	-	-	-	-	-	-	-	4
1955	30,587	30,571	12	4	-	-	-	-	-	-	-	-	-
1954	30,483	30,459	18	6	-	-	-	-	-	-	-	-	-
1953	30,994	30,976	14	4	-	-	-	-	-	-	-	-	-
1952	32,208	32,185	20	3	-	-	-	-	-	-	-	-	-
1951	32,207	32,162	35	6	-	-	-	1	-	-	-	-	3
1950	32,067	32,046	18	3	-	-	-	-	-	-	-	-	-
MALE													
1959	18,638	18,628	5	5	-	-	-	-	-	-	-	-	-
1958	18,107	18,095	5	5	1	-	-	-	-	-	-	-	1
1957	17,661	17,655	5	-	1	-	-	-	-	-	-	-	-
1956	17,328	17,314	12	-	-	-	-	-	-	-	-	-	2
1955	17,358	17,346	10	2	-	-	-	-	-	-	-	-	-
1954	17,309	17,293	10	6	-	-	-	-	-	-	-	-	-
1953	17,652	17,643	8	1	-	-	-	-	-	-	-	-	-
1952	18,266	18,254	9	3	-	-	-	-	-	-	-	-	-
1951	18,427	18,408	15	3	-	-	-	-	-	-	-	-	1
1950	18,363	18,348	14	1	-	-	-	-	-	-	-	-	-
FEMALE													
1959	13,983	13,978	2	3	-	-	-	-	-	-	-	-	-
1958	13,797	13,790	3	3	-	1	-	-	-	-	-	-	-
1957	13,548	13,537	8	3	-	-	-	-	-	-	-	-	-
1956	13,006	12,997	5	2	-	-	-	-	-	-	-	-	2
1955	13,229	13,225	2	2	-	-	-	-	-	-	-	-	-
1954	13,174	13,166	8	-	-	-	-	-	-	-	-	-	-
1953	13,342	13,333	6	3	-	-	-	-	-	-	-	-	-
1952	13,942	13,931	11	-	-	-	-	-	-	-	-	-	-
1951	13,780	13,754	20	3	-	-	-	1	-	-	-	-	2
1950	13,704	13,698	4	2	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURETY UNQUALIFIED (769-776)													
WHITE, BOTH SEXES													
1959	24,379	24,365	6	8	-	-	-	-	-	-	-	-	-
1958	23,807	23,788	8	8	1	1	-	-	-	-	-	-	1
1957	23,614	23,599	11	3	1	-	-	-	-	-	-	-	-
1956	23,019	22,998	16	2	-	-	-	-	-	-	-	-	3
1955	23,413	23,398	11	4	-	-	-	-	-	-	-	-	-
1954	23,410	23,388	16	6	-	-	-	-	-	-	-	-	-
1953	24,084	24,066	14	4	-	-	-	-	-	-	-	-	-
1952	25,216	25,195	18	3	-	-	-	-	-	-	-	-	-
1951	25,323	25,285	28	6	-	-	-	1	-	-	-	-	3
1950	25,289	25,269	17	3	-	-	-	-	-	-	-	-	-
WHITE, MALE													
1959	14,098	14,088	5	5	-	-	-	-	-	-	-	-	-
1958	13,684	13,672	5	5	1	-	-	-	-	-	-	-	1
1957	13,514	13,510	3	-	1	-	-	-	-	-	-	-	-
1956	13,212	13,199	11	-	-	-	-	-	-	-	-	-	2
1955	13,376	13,365	9	2	-	-	-	-	-	-	-	-	-
1954	13,381	13,366	9	6	-	-	-	-	-	-	-	-	-
1953	13,895	13,886	8	1	-	-	-	-	-	-	-	-	-
1952	14,440	14,428	9	3	-	-	-	-	-	-	-	-	-
1951	14,566	14,550	12	3	-	-	-	-	-	-	-	-	1
1950	14,619	14,605	13	1	-	-	-	-	-	-	-	-	-
WHITE, FEMALE													
1959	10,281	10,277	1	3	-	-	-	-	-	-	-	-	-
1958	10,123	10,116	3	3	-	1	-	-	-	-	-	-	-
1957	10,100	10,089	8	3	-	-	-	-	-	-	-	-	-
1956	9,807	9,799	5	2	-	-	-	-	-	-	-	-	1
1955	10,037	10,033	2	2	-	-	-	-	-	-	-	-	-
1954	10,029	10,022	7	-	-	-	-	-	-	-	-	-	-
1953	10,189	10,180	6	3	-	-	-	-	-	-	-	-	-
1952	10,776	10,767	9	-	-	-	-	-	-	-	-	-	-
1951	10,757	10,735	16	3	-	-	-	1	-	-	-	-	2
1950	10,670	10,664	4	2	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER DISEASES PECULIAR TO EARLY INFANCY, AND IMMATURITY UNQUALIFIED (769-776)													
ALL OTHER, BOTH SEXES													
1959	8,242	8,241	1	-	-	-	-	-	-	-	-	-	-
1958	8,097	8,097	-	-	-	-	-	-	-	-	-	-	-
1957	7,595	7,593	2	-	-	-	-	-	-	-	-	-	-
1956	7,315	7,313	1	-	-	-	-	-	-	-	-	-	1
1955	7,174	7,173	1	-	-	-	-	-	-	-	-	-	-
1954	7,073	7,071	2	-	-	-	-	-	-	-	-	-	-
1953	6,910	6,910	-	-	-	-	-	-	-	-	-	-	-
1952	6,992	6,990	2	-	-	-	-	-	-	-	-	-	-
1951	6,884	6,877	7	-	-	-	-	-	-	-	-	-	-
1950	6,778	6,777	1	-	-	-	-	-	-	-	-	-	-
ALL OTHER, MALE													
1959	4,540	4,540	-	-	-	-	-	-	-	-	-	-	-
1958	4,423	4,423	-	-	-	-	-	-	-	-	-	-	-
1957	4,147	4,145	2	-	-	-	-	-	-	-	-	-	-
1956	4,116	4,115	1	-	-	-	-	-	-	-	-	-	-
1955	3,982	3,981	1	-	-	-	-	-	-	-	-	-	-
1954	3,928	3,927	1	-	-	-	-	-	-	-	-	-	-
1953	3,757	3,757	-	-	-	-	-	-	-	-	-	-	-
1952	3,826	3,826	-	-	-	-	-	-	-	-	-	-	-
1951	3,861	3,858	3	-	-	-	-	-	-	-	-	-	-
1950	3,744	3,743	1	-	-	-	-	-	-	-	-	-	-
ALL OTHER, FEMALE													
1959	3,702	3,701	1	-	-	-	-	-	-	-	-	-	-
1958	3,674	3,674	-	-	-	-	-	-	-	-	-	-	-
1957	3,448	3,448	-	-	-	-	-	-	-	-	-	-	-
1956	3,199	3,198	-	-	-	-	-	-	-	-	-	-	1
1955	3,192	3,192	-	-	-	-	-	-	-	-	-	-	-
1954	3,145	3,144	1	-	-	-	-	-	-	-	-	-	-
1953	3,153	3,153	-	-	-	-	-	-	-	-	-	-	-
1952	3,166	3,164	2	-	-	-	-	-	-	-	-	-	-
1951	3,023	3,019	4	-	-	-	-	-	-	-	-	-	-
1950	3,034	3,034	-	-	-	-	-	-	-	-	-	-	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)													
BOTH SEXES													
1959	19,066	2,467	384	161	328	581	1,080	1,818	2,475	3,500	3,419	2,767	86
1958	19,729	2,752	428	154	327	620	1,129	1,983	2,418	3,348	3,523	2,967	80
1957	19,164	2,557	366	171	379	598	1,015	1,669	2,483	3,281	3,419	3,151	75
1956	18,912	2,555	363	167	311	600	1,111	1,783	2,452	3,153	3,361	2,958	98
1955	19,914	2,790	443	174	405	624	1,100	1,897	2,499	3,345	3,452	3,077	108
1954	20,209	2,780	445	174	436	645	1,107	1,832	2,382	3,414	3,694	3,242	58
1953	21,693	2,963	494	203	363	587	1,094	1,924	2,558	3,634	4,050	3,757	66
1952	21,950	3,353	483	192	343	583	1,069	1,892	2,459	3,566	4,124	3,808	78
1951	22,647	3,379	548	204	354	591	1,120	1,806	2,504	3,594	4,386	4,077	84
1950	22,517	3,636	503	201	348	589	1,055	1,798	2,324	3,523	4,285	4,160	95
MALE													
1959	11,182	1,377	207	96	208	315	645	1,230	1,678	2,299	1,892	1,178	57
1958	11,487	1,535	233	88	203	330	670	1,310	1,647	2,185	1,950	1,281	55
1957	11,211	1,457	195	101	228	327	629	1,096	1,704	2,102	1,926	1,392	54
1956	11,140	1,403	196	93	194	336	687	1,211	1,691	2,077	1,874	1,319	59
1955	11,802	1,561	245	107	245	368	692	1,283	1,735	2,195	1,972	1,335	64
1954	11,975	1,556	252	108	274	374	693	1,244	1,692	2,222	2,127	1,392	41
1953	12,521	1,667	251	122	200	314	677	1,298	1,780	2,287	2,244	1,646	35
1952	12,638	1,909	252	109	197	312	674	1,247	1,703	2,277	2,324	1,585	49
1951	13,088	1,968	300	115	203	309	689	1,216	1,671	2,362	2,409	1,793	53
1950	12,823	2,054	268	113	192	300	616	1,177	1,610	2,246	2,328	1,853	66
FEMALE													
1959	7,884	1,090	177	65	120	266	435	588	797	1,201	1,527	1,589	29
1958	8,242	1,217	195	66	124	290	459	673	771	1,163	1,573	1,686	25
1957	7,953	1,100	171	70	151	271	386	573	779	1,179	1,493	1,759	21
1956	7,772	1,152	167	74	117	264	424	572	761	1,076	1,487	1,639	39
1955	8,112	1,229	198	67	160	256	408	614	764	1,150	1,480	1,742	44
1954	8,234	1,224	193	66	162	271	414	588	690	1,192	1,567	1,850	17
1953	9,172	1,296	243	81	163	273	417	626	778	1,347	1,806	2,111	31
1952	9,312	1,444	231	83	146	271	395	645	756	1,289	1,800	2,223	29
1951	9,559	1,411	248	89	151	282	431	590	833	1,232	1,977	2,284	31
1950	9,694	1,582	235	88	156	289	439	621	714	1,277	1,957	2,307	29

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)													
WHITE, BOTH SEXES													
1959	12,471	1,022	178	103	209	308	663	1,146	1,611	2,376	2,582	2,219	54
1958	12,836	1,168	188	95	189	317	676	1,270	1,576	2,257	2,691	2,362	47
1957	12,552	1,081	185	109	229	299	619	1,045	1,611	2,188	2,606	2,532	48
1956	12,421	1,062	173	90	191	312	689	1,129	1,584	2,154	2,565	2,397	75
1955	13,147	1,202	224	92	254	345	679	1,196	1,672	2,266	2,634	2,513	70
1954	13,358	1,156	230	104	279	359	670	1,138	1,587	2,338	2,805	2,657	35
1953	13,903	1,270	246	111	178	302	614	1,133	1,605	2,359	3,083	2,963	39
1952	14,055	1,413	227	106	175	299	626	1,088	1,575	2,330	3,167	3,003	46
1951	14,589	1,479	307	120	187	310	636	1,046	1,549	2,349	3,303	3,248	55
1950	14,423	1,668	266	103	179	307	608	1,013	1,472	2,256	3,213	3,292	46
WHITE, MALE													
1959	7,552	593	105	66	137	177	422	841	1,173	1,648	1,419	929	42
1958	7,663	674	92	58	120	172	428	917	1,134	1,565	1,495	970	38
1957	7,549	651	93	67	149	176	417	744	1,189	1,465	1,463	1,097	38
1956	7,568	598	93	48	123	180	457	836	1,183	1,491	1,441	1,073	45
1955	8,125	726	132	63	174	220	452	885	1,251	1,567	1,527	1,088	40
1954	8,151	662	136	70	191	218	451	860	1,199	1,599	1,617	1,123	25
1953	8,306	773	120	71	102	171	414	850	1,214	1,576	1,717	1,279	19
1952	8,381	826	118	57	113	189	426	797	1,190	1,599	1,788	1,245	33
1951	8,642	889	161	74	115	167	429	781	1,121	1,628	1,818	1,420	39
1950	8,366	945	151	62	107	176	389	728	1,090	1,522	1,715	1,447	34
WHITE, FEMALE													
1959	4,919	429	73	37	72	131	241	305	438	728	1,163	1,290	12
1958	5,173	494	96	37	69	145	248	353	442	692	1,196	1,392	9
1957	5,003	430	92	42	80	123	202	301	422	723	1,143	1,435	10
1956	4,853	464	80	42	68	132	232	293	401	663	1,124	1,324	30
1955	5,022	476	92	29	80	125	227	311	421	699	1,107	1,425	30
1954	5,207	494	94	34	88	141	219	278	388	739	1,188	1,534	10
1953	5,597	497	126	40	76	131	200	283	391	783	1,366	1,684	20
1952	5,674	587	109	49	62	110	200	291	385	731	1,379	1,758	13
1951	5,947	590	146	46	72	143	207	265	428	721	1,485	1,828	16
1950	6,057	723	115	41	72	131	219	285	382	734	1,498	1,845	12

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SYMPTOMS, SENILITY, AND ILL-DEFINED CONDITIONS (780-795)													
ALL OTHER, BOTH SEXES													
1959	6,595	1,445	206	58	119	273	417	672	864	1,124	837	548	32
1958	6,893	1,584	240	59	138	303	453	713	842	1,091	832	605	33
1957	6,612	1,476	181	62	150	299	396	624	872	1,093	813	619	27
1956	6,491	1,493	190	77	120	288	422	654	868	999	796	561	23
1955	6,767	1,588	219	82	151	279	421	701	827	1,079	818	564	38
1954	6,851	1,624	215	70	157	286	437	694	795	1,076	889	585	23
1953	7,790	1,693	248	92	185	285	480	791	953	1,275	967	794	27
1952	7,895	1,940	256	86	168	284	443	804	884	1,236	957	805	32
1951	8,058	1,900	241	84	167	281	484	760	955	1,245	1,083	829	29
1950	8,094	1,968	237	98	169	282	447	785	852	1,267	1,072	868	49
ALL OTHER, MALE													
1959	3,630	784	102	30	71	138	223	389	505	651	473	249	15
1958	3,824	861	141	30	83	158	242	393	513	620	455	311	17
1957	3,662	806	102	34	79	151	212	352	515	637	463	295	16
1956	3,572	805	103	45	71	156	230	375	508	586	433	246	14
1955	3,677	835	113	44	71	148	240	398	484	628	445	247	24
1954	3,824	894	116	38	83	156	242	384	493	623	510	269	16
1953	4,215	894	131	51	98	143	263	448	566	711	527	367	16
1952	4,257	1,083	134	52	84	123	248	450	513	678	536	340	16
1951	4,446	1,079	139	41	88	142	260	435	550	734	591	373	14
1950	4,457	1,109	117	51	85	124	227	449	520	724	613	406	32
ALL OTHER, FEMALE													
1959	2,965	661	104	28	48	135	194	283	359	473	364	299	17
1958	3,069	723	99	29	55	145	211	320	329	471	377	294	16
1957	2,950	670	79	28	71	148	184	272	357	456	350	324	11
1956	2,919	688	87	32	49	132	192	279	360	413	363	315	9
1955	3,090	753	106	38	80	131	181	303	343	451	373	317	14
1954	3,027	730	99	32	74	130	195	310	302	453	379	316	7
1953	3,575	799	117	41	87	142	217	343	387	564	440	427	11
1952	3,638	857	122	34	84	161	195	354	371	558	421	465	16
1951	3,612	821	102	43	79	139	224	325	405	511	492	456	15
1950	3,637	859	120	47	84	158	220	336	332	543	459	462	17

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ALL OTHER DISEASES (RESIDUAL)													
BOTH SEXES													
1959	65,787	4,281	1,929	1,752	1,613	2,323	4,463	7,371	11,387	15,597	10,898	4,155	18
1958	63,353	4,269	1,929	1,667	1,486	2,328	4,387	7,089	10,933	14,705	10,705	3,818	37
1957	77,533	5,145	1,807	1,529	1,693	2,751	5,365	8,644	13,909	18,756	13,162	4,740	32
1956	71,982	4,742	1,719	1,433	1,518	2,684	4,964	7,926	13,031	16,970	12,527	4,425	43
1955	69,146	4,292	1,730	1,501	1,595	2,721	4,905	7,698	12,514	16,358	11,710	4,067	55
1954	66,354	4,124	1,830	1,437	1,561	2,763	4,764	7,873	12,046	15,238	10,968	3,712	38
1953	67,406	3,929	1,954	1,481	1,661	2,861	5,034	8,081	12,661	15,111	10,923	3,675	35
1952	67,225	3,802	2,055	1,440	1,773	3,090	5,233	8,235	12,601	14,908	10,465	3,567	56
1951	65,681	3,299	1,938	1,409	1,730	3,072	5,185	8,177	12,313	14,590	10,369	3,537	62
1950	61,746	3,249	1,917	1,325	1,847	3,056	5,071	7,647	11,493	13,231	9,521	3,353	36
MALE													
1959	39,923	2,394	1,039	968	910	1,250	2,524	4,586	7,708	10,366	6,245	1,920	13
1958	38,633	2,479	1,035	962	841	1,166	2,486	4,441	7,474	9,811	6,133	1,783	22
1957	45,982	3,101	1,006	831	942	1,311	2,886	5,200	9,148	12,046	7,322	2,168	21
1956	42,196	2,890	931	773	823	1,289	2,618	4,826	8,394	10,747	6,881	1,996	28
1955	40,083	2,573	959	825	866	1,351	2,653	4,616	7,939	10,075	6,329	1,867	30
1954	38,120	2,421	1,012	776	810	1,352	2,517	4,548	7,717	9,347	5,894	1,701	25
1953	38,844	2,341	1,084	819	914	1,401	2,721	4,728	8,166	9,079	5,856	1,712	23
1952	38,227	2,214	1,121	783	954	1,485	2,791	4,899	7,899	8,834	5,652	1,561	34
1951	37,310	1,914	1,053	783	928	1,528	2,752	4,808	7,689	8,657	5,499	1,659	40
1950	34,283	1,907	1,065	712	957	1,448	2,710	4,455	6,927	7,591	4,936	1,552	23
FEMALE													
1959	25,864	1,887	890	784	703	1,073	1,939	2,785	3,679	5,231	4,653	2,235	5
1958	24,720	1,790	894	705	645	1,162	1,901	2,648	3,459	4,894	4,572	2,035	15
1957	31,551	2,044	801	698	751	1,440	2,479	3,444	4,761	6,710	5,840	2,572	11
1956	29,786	1,852	788	660	695	1,395	2,346	3,100	4,637	6,223	5,646	2,429	15
1955	29,063	1,719	771	676	729	1,370	2,252	3,082	4,575	6,283	5,381	2,200	25
1954	28,234	1,703	818	661	751	1,411	2,247	3,325	4,329	5,891	5,074	2,011	13
1953	28,562	1,588	870	662	747	1,460	2,313	3,353	4,495	6,032	5,067	1,963	12
1952	28,998	1,588	934	657	819	1,605	2,442	3,336	4,702	6,074	4,813	2,006	22
1951	28,371	1,385	885	626	802	1,544	2,433	3,369	4,624	5,933	4,870	1,878	22
1950	27,463	1,342	852	613	890	1,608	2,361	3,192	4,566	5,640	4,585	1,801	13

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ALL OTHER DISEASES (RESIDUAL)													
WHITE, BOTH SEXES													
1959	57,494	3,160	1,525	1,496	1,252	1,612	3,382	6,143	10,146	14,468	10,380	3,919	11
1958	55,392	3,160	1,542	1,438	1,161	1,647	3,347	5,952	9,697	13,634	10,173	3,622	19
1957	67,684	4,030	1,412	1,293	1,307	1,953	4,093	7,065	12,247	17,316	12,465	4,485	18
1956	63,177	3,742	1,401	1,226	1,178	1,935	3,791	6,491	11,553	15,725	11,889	4,221	25
1955	60,869	3,415	1,415	1,281	1,228	1,939	3,806	6,437	11,162	15,138	11,149	3,861	38
1954	58,324	3,256	1,493	1,210	1,189	2,055	3,670	6,499	10,802	14,163	10,445	3,514	28
1953	59,201	3,126	1,597	1,263	1,266	2,098	3,876	6,741	11,358	14,017	10,370	3,460	29
1952	59,114	2,975	1,715	1,242	1,375	2,316	4,128	6,880	11,328	13,823	9,914	3,377	41
1951	57,847	2,618	1,606	1,156	1,321	2,347	4,055	6,827	11,134	13,544	9,864	3,334	41
1950	54,374	2,595	1,584	1,134	1,418	2,356	3,979	6,448	10,346	12,296	9,019	3,170	29
WHITE, MALE													
1959	35,138	1,799	804	839	737	862	1,934	3,891	6,949	9,596	5,917	1,802	8
1958	34,006	1,826	831	837	673	839	1,926	3,741	6,688	9,123	5,821	1,690	11
1957	40,373	2,460	775	709	757	936	2,233	4,327	8,100	11,134	6,896	2,035	11
1956	37,279	2,303	763	668	648	942	2,059	4,001	7,503	10,005	6,486	1,885	16
1955	35,462	2,066	770	706	697	982	2,113	3,932	7,109	9,322	5,983	1,761	21
1954	33,663	1,918	816	658	635	1,025	1,980	3,788	6,930	8,688	5,595	1,611	19
1953	34,233	1,896	888	697	727	1,040	2,120	3,972	7,348	8,405	5,532	1,589	19
1952	33,713	1,765	920	680	756	1,139	2,231	4,112	7,127	8,184	5,301	1,471	27
1951	33,074	1,528	891	642	716	1,197	2,195	4,079	6,998	8,026	5,215	1,561	26
1950	30,265	1,529	884	604	756	1,134	2,174	3,783	6,245	7,041	4,636	1,463	16
WHITE, FEMALE													
1959	22,356	1,361	721	657	515	750	1,448	2,252	3,197	4,872	4,463	2,117	3
1958	21,386	1,334	711	601	488	808	1,421	2,211	3,009	4,511	4,352	1,932	8
1957	27,311	1,570	637	584	550	1,017	1,860	2,738	4,147	6,182	5,569	2,450	7
1956	25,898	1,439	638	558	530	993	1,732	2,490	4,050	5,720	5,403	2,336	9
1955	25,407	1,349	645	575	531	957	1,693	2,505	4,053	5,816	5,166	2,100	17
1954	24,661	1,338	677	552	554	1,030	1,690	2,711	3,872	5,475	4,850	1,903	9
1953	24,968	1,230	709	566	539	1,058	1,756	2,769	4,010	5,612	4,838	1,871	10
1952	25,401	1,210	795	562	619	1,177	1,897	2,768	4,201	5,639	4,613	1,906	14
1951	24,773	1,090	715	514	605	1,150	1,860	2,748	4,136	5,518	4,649	1,773	15
1950	24,109	1,066	700	530	662	1,222	1,805	2,665	4,101	5,255	4,383	1,707	13

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
ALL OTHER DISEASES (RESIDUAL)													
ALL OTHER, BOTH SEXES													
1959	8,293	1,121	404	256	361	711	1,081	1,228	1,241	1,129	518	236	7
1958	7,961	1,109	387	229	325	681	1,040	1,137	1,236	1,071	532	196	18
1957	9,849	1,115	395	236	386	798	1,272	1,579	1,662	1,440	697	255	14
1956	8,805	1,000	318	207	340	749	1,173	1,435	1,478	1,245	638	204	18
1955	8,277	877	315	220	367	782	1,099	1,261	1,352	1,220	561	206	17
1954	8,030	868	337	227	372	708	1,094	1,374	1,244	1,075	523	198	10
1953	8,205	803	357	218	395	763	1,158	1,340	1,303	1,094	553	215	6
1952	8,111	827	340	198	398	774	1,105	1,355	1,273	1,085	551	190	15
1951	7,834	681	332	253	409	725	1,130	1,350	1,179	1,046	505	203	21
1950	7,372	654	333	191	429	700	1,092	1,199	1,147	935	502	183	7
ALL OTHER, MALE													
1959	4,785	595	235	129	173	388	590	695	759	770	328	118	5
1958	4,627	653	204	125	168	327	560	700	786	688	312	93	11
1957	5,609	641	231	122	185	375	653	873	1,048	912	426	133	10
1956	4,917	587	168	105	175	347	559	825	891	742	395	111	12
1955	4,621	507	189	119	169	369	540	684	830	753	346	106	9
1954	4,457	503	196	118	175	327	537	760	787	659	299	90	6
1953	4,611	445	196	122	187	361	601	756	818	674	324	123	4
1952	4,514	449	201	103	198	346	560	787	772	650	351	90	7
1951	4,236	386	162	141	212	331	557	729	691	631	284	98	14
1950	4,018	378	181	108	201	314	536	672	682	550	300	89	7
ALL OTHER, FEMALE													
1959	3,508	526	169	127	188	323	491	533	482	359	190	118	2
1958	3,334	456	183	104	157	354	480	437	450	383	220	103	7
1957	4,240	474	164	114	201	423	619	706	614	528	271	122	4
1956	3,888	413	150	102	165	402	614	610	587	503	243	93	6
1955	3,656	370	126	101	198	413	559	577	522	467	215	100	8
1954	3,573	365	141	109	197	381	557	614	457	416	224	108	4
1953	3,594	358	161	96	208	402	557	584	485	420	229	92	2
1952	3,597	378	139	95	200	428	545	568	501	435	200	100	8
1951	3,598	295	170	112	197	394	573	621	488	415	221	105	7
1950	3,354	276	152	83	228	386	556	527	465	385	202	94	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)													
BOTH SEXES													
1959	92,080	3,857	4,891	6,511	13,269	9,956	9,710	9,972	8,965	9,475	9,550	5,820	104
1958	90,604	3,916	4,873	6,514	12,744	9,874	9,784	9,368	8,727	9,431	9,642	5,634	97
1957	95,307	3,674	4,749	6,454	12,973	10,828	10,121	10,005	9,490	10,076	10,520	6,204	213
1956	94,780	3,498	4,675	6,316	13,545	10,914	10,072	9,794	9,413	10,160	10,210	6,047	136
1955	93,443	3,308	4,791	6,099	12,742	10,904	10,007	9,808	9,391	9,929	10,313	6,005	146
1954	90,032	3,580	4,800	5,939	11,801	10,607	9,416	9,326	8,973	9,652	10,130	5,686	122
1953	95,032	3,629	5,049	6,136	12,837	11,411	10,011	9,911	9,568	9,927	10,567	5,824	162
1952	96,172	3,730	5,141	5,980	12,786	11,553	10,397	9,996	9,896	10,026	10,636	5,871	160
1951	95,871	3,711	5,058	5,892	12,366	11,644	10,719	9,809	9,801	10,218	10,647	5,818	188
1950	91,249	3,595	4,794	5,519	12,119	10,864	9,799	9,199	9,466	9,750	10,347	5,611	186
MALE													
1959	63,639	2,130	2,850	4,644	11,015	8,149	7,706	7,632	6,601	6,023	4,659	2,140	90
1958	62,489	2,242	2,814	4,529	10,589	8,226	7,823	7,225	6,373	5,878	4,617	2,097	76
1957	65,647	2,065	2,680	4,558	10,768	8,980	8,020	7,726	6,935	6,450	5,046	2,267	152
1956	65,823	2,006	2,709	4,462	11,240	9,070	7,995	7,644	6,916	6,541	4,893	2,237	110
1955	64,690	1,899	2,737	4,319	10,537	9,012	7,966	7,689	6,927	6,448	4,917	2,124	115
1954	62,315	2,046	2,770	4,206	9,834	8,880	7,500	7,244	6,638	6,184	4,828	2,090	95
1953	66,246	2,079	2,951	4,331	10,574	9,528	8,047	7,805	7,227	6,376	5,075	2,123	130
1952	66,697	2,092	2,967	4,168	10,451	9,618	8,362	7,847	7,486	6,486	4,998	2,094	128
1951	66,873	2,078	2,969	4,210	10,207	9,705	8,731	7,787	7,378	6,637	4,913	2,104	154
1950	63,415	2,057	2,823	3,901	10,088	9,002	7,941	7,258	7,158	6,260	4,817	1,948	162
FEMALE													
1959	28,441	1,727	2,041	1,867	2,254	1,807	2,004	2,340	2,364	3,452	4,891	3,680	14
1958	28,115	1,674	2,059	1,985	2,155	1,648	1,961	2,143	2,354	3,553	5,025	3,537	21
1957	29,660	1,609	2,069	1,896	2,205	1,848	2,101	2,279	2,555	3,626	5,474	3,937	61
1956	28,957	1,492	1,966	1,854	2,305	1,844	2,077	2,150	2,497	3,619	5,317	3,810	26
1955	28,753	1,409	2,054	1,780	2,205	1,892	2,041	2,119	2,464	3,481	5,396	3,881	31
1954	27,717	1,534	2,030	1,733	1,967	1,727	1,916	2,082	2,335	3,468	5,302	3,596	27
1953	28,786	1,550	2,098	1,805	2,263	1,883	1,964	2,106	2,341	3,551	5,492	3,701	32
1952	29,475	1,638	2,174	1,812	2,335	1,935	2,035	2,149	2,410	3,540	5,638	3,777	32
1951	28,998	1,633	2,089	1,682	2,159	1,939	1,988	2,022	2,423	3,581	5,734	3,714	34
1950	27,834	1,538	1,971	1,618	2,031	1,862	1,858	1,941	2,308	3,490	5,530	3,663	24

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)													
WHITE, BOTH SEXES													
1959	79,224	2,660	3,655	5,224	11,658	8,325	8,032	8,471	7,805	8,619	9,088	5,612	75
1958	77,920	2,671	3,611	5,346	11,126	8,242	8,194	7,985	7,593	8,537	9,133	5,411	71
1957	82,165	2,531	3,575	5,251	11,361	8,970	8,453	8,467	8,328	9,167	9,981	5,980	101
1956	81,906	2,389	3,615	5,183	11,845	9,116	8,262	8,330	8,237	9,275	9,724	5,837	93
1955	81,010	2,353	3,609	5,054	11,153	9,119	8,345	8,293	8,282	9,053	9,833	5,798	118
1954	78,095	2,520	3,705	4,955	10,276	8,864	7,897	7,884	7,977	8,808	9,636	5,480	93
1953	82,333	2,633	3,892	5,142	11,164	9,488	8,273	8,357	8,502	9,106	10,044	5,611	121
1952	83,393	2,676	3,977	5,020	11,094	9,661	8,667	8,393	8,789	9,195	10,129	5,672	120
1951	83,656	2,713	4,008	4,893	10,783	9,918	8,996	8,318	8,734	9,396	10,172	5,580	145
1950	80,034	2,675	3,824	4,576	10,621	9,241	8,206	7,883	8,483	9,004	9,952	5,419	150
WHITE, MALE													
1959	54,593	1,502	2,173	3,783	9,682	6,864	6,398	6,469	5,717	5,479	4,404	2,057	65
1958	53,669	1,582	2,175	3,765	9,252	6,904	6,579	6,155	5,533	5,329	4,338	2,001	56
1957	56,476	1,452	2,080	3,751	9,469	7,498	6,726	6,549	6,082	5,874	4,750	2,169	76
1956	56,645	1,396	2,134	3,725	9,840	7,608	6,570	6,497	6,059	5,985	4,626	2,133	72
1955	55,976	1,388	2,134	3,632	9,246	7,611	6,659	6,494	6,121	5,887	4,671	2,040	93
1954	53,949	1,496	2,201	3,535	8,596	7,481	6,338	6,121	5,916	5,647	4,562	1,985	71
1953	57,176	1,545	2,321	3,642	9,235	7,991	6,692	6,588	6,419	5,840	4,769	2,036	98
1952	57,605	1,522	2,347	3,538	9,095	8,113	6,968	6,594	6,659	5,962	4,711	2,001	95
1951	58,214	1,574	2,396	3,509	8,970	8,321	7,380	6,619	6,586	6,085	4,647	2,005	122
1950	55,412	1,560	2,310	3,267	8,891	7,718	6,680	6,195	6,416	5,784	4,592	1,866	133
WHITE, FEMALE													
1959	24,631	1,158	1,482	1,441	1,976	1,461	1,634	2,002	2,088	3,140	4,684	3,555	10
1958	24,251	1,089	1,436	1,581	1,874	1,338	1,615	1,830	2,060	3,208	4,795	3,410	15
1957	25,689	1,079	1,495	1,500	1,892	1,472	1,727	1,918	2,246	3,293	5,231	3,811	25
1956	25,261	993	1,481	1,458	2,005	1,508	1,692	1,833	2,178	3,290	5,098	3,704	21
1955	25,034	965	1,475	1,422	1,907	1,508	1,686	1,799	2,161	3,166	5,162	3,758	25
1954	24,146	1,024	1,504	1,420	1,680	1,383	1,559	1,763	2,061	3,161	5,074	3,495	22
1953	25,157	1,088	1,571	1,500	1,929	1,497	1,581	1,769	2,083	3,266	5,275	3,575	23
1952	25,788	1,154	1,630	1,482	1,999	1,548	1,699	1,799	2,130	3,233	5,418	3,671	25
1951	25,442	1,139	1,612	1,384	1,813	1,597	1,616	1,699	2,148	3,311	5,525	3,575	23
1950	24,622	1,115	1,514	1,309	1,730	1,523	1,526	1,688	2,067	3,220	5,360	3,553	17

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
STATISTICAL WORKSHEET													
ACCIDENTS AND ADVERSE EFFECTS (E800-E962)													
ALL OTHER, BOTH SEXES													
1959	12,856	1,197	1,236	1,287	1,611	1,631	1,678	1,501	1,160	856	462	208	29
1958	12,684	1,245	1,262	1,168	1,618	1,632	1,590	1,383	1,134	894	509	223	26
1957	13,142	1,143	1,174	1,203	1,612	1,858	1,668	1,538	1,162	909	539	224	112
1956	12,874	1,109	1,060	1,133	1,700	1,798	1,810	1,464	1,176	885	486	210	43
1955	12,433	955	1,182	1,045	1,589	1,785	1,662	1,515	1,109	876	480	207	28
1954	11,937	1,060	1,095	984	1,525	1,743	1,519	1,442	996	844	494	206	29
1953	12,699	996	1,157	994	1,673	1,923	1,738	1,554	1,066	821	523	213	41
1952	12,779	1,054	1,164	960	1,692	1,892	1,730	1,603	1,107	831	507	199	40
1951	12,215	998	1,050	999	1,583	1,726	1,723	1,491	1,067	822	475	238	43
1950	11,215	920	970	943	1,498	1,623	1,593	1,316	983	746	395	192	36
ALL OTHER, MALE													
1959	9,046	628	677	861	1,333	1,285	1,308	1,163	884	544	255	83	25
1958	8,820	660	639	764	1,337	1,322	1,244	1,070	840	549	279	96	20
1957	9,171	613	600	807	1,299	1,482	1,294	1,177	853	576	296	98	76
1956	9,178	610	575	737	1,400	1,462	1,425	1,147	857	556	267	104	38
1955	8,714	511	603	687	1,291	1,401	1,307	1,195	806	561	246	84	22
1954	8,366	550	569	671	1,238	1,399	1,162	1,123	722	537	266	105	24
1953	9,070	534	630	689	1,339	1,537	1,355	1,217	808	536	306	87	32
1952	9,092	570	620	630	1,356	1,505	1,394	1,253	827	524	287	93	33
1951	8,659	504	573	701	1,237	1,384	1,351	1,168	792	552	266	99	32
1950	8,003	497	513	634	1,197	1,284	1,261	1,063	742	476	225	82	29
ALL OTHER, FEMALE													
1959	3,810	569	559	426	278	346	370	338	276	312	207	125	4
1958	3,864	585	623	404	281	310	346	313	294	345	230	127	6
1957	3,971	530	574	396	313	376	374	361	309	333	243	126	36
1956	3,696	499	485	396	300	336	385	317	319	329	219	106	5
1955	3,719	444	579	358	298	384	355	320	303	315	234	123	6
1954	3,571	510	526	313	287	344	357	319	274	307	228	101	5
1953	3,629	462	527	305	334	386	383	337	258	285	217	126	9
1952	3,687	484	544	330	336	387	336	350	280	307	220	106	7
1951	3,556	494	477	298	346	342	372	323	275	270	209	139	11
1950	3,212	423	457	309	301	339	332	253	241	270	170	110	7

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MOTOR VEHICLE ACCIDENTS (E810-E835)</b>													
<b>BOTH SEXES</b>													
1959	37,910	319	1,523	2,719	8,969	5,699	4,659	4,446	3,817	3,487	1,910	330	32
1958	36,981	293	1,498	2,710	8,388	5,656	4,758	4,204	3,718	3,535	1,872	320	29
1957	38,702	337	1,448	2,604	8,667	6,280	4,950	4,450	4,095	3,560	1,958	317	36
1956	39,628	332	1,438	2,640	9,169	6,483	5,068	4,508	4,065	3,657	1,893	339	36
1955	38,426	328	1,547	2,406	8,656	6,470	4,978	4,286	4,086	3,455	1,854	315	45
1954	35,586	296	1,568	2,332	7,571	6,016	4,505	4,083	3,765	3,247	1,844	316	43
1953	37,955	332	1,687	2,368	8,169	6,441	4,861	4,308	4,010	3,508	1,890	345	36
1952	37,794	308	1,643	2,295	8,115	6,437	4,943	4,328	4,135	3,472	1,783	289	46
1951	36,996	287	1,588	2,300	7,713	6,336	4,917	4,260	4,016	3,444	1,789	293	53
1950	34,763	264	1,503	2,152	7,600	5,855	4,359	3,848	3,880	3,264	1,727	260	51
<b>MALE</b>													
1959	28,064	153	871	1,850	7,257	4,544	3,552	3,261	2,675	2,353	1,293	230	25
1958	27,492	163	868	1,827	6,789	4,607	3,652	3,064	2,608	2,387	1,272	234	21
1957	28,900	182	838	1,742	7,017	5,099	3,767	3,255	2,889	2,464	1,381	236	30
1956	29,804	187	800	1,785	7,432	5,269	3,888	3,365	2,909	2,543	1,352	243	31
1955	28,803	177	867	1,630	6,984	5,219	3,870	3,164	2,904	2,434	1,284	237	33
1954	26,725	171	908	1,547	6,158	4,931	3,452	3,015	2,694	2,279	1,324	216	30
1953	28,562	192	1,009	1,576	6,517	5,253	3,767	3,193	2,928	2,493	1,367	240	27
1952	28,536	163	977	1,529	6,459	5,260	3,836	3,195	3,025	2,568	1,290	199	35
1951	28,075	156	978	1,543	6,222	5,138	3,860	3,194	2,925	2,492	1,305	218	44
1950	26,516	146	865	1,467	6,195	4,727	3,439	2,905	2,884	2,377	1,281	185	45
<b>FEMALE</b>													
1959	9,846	166	652	869	1,712	1,155	1,107	1,185	1,142	1,134	617	100	7
1958	9,489	130	630	883	1,599	1,049	1,106	1,140	1,110	1,148	600	86	8
1957	9,802	155	610	862	1,650	1,181	1,183	1,195	1,206	1,096	577	81	6
1956	9,824	145	638	855	1,737	1,214	1,180	1,143	1,156	1,114	541	96	5
1955	9,623	151	680	776	1,672	1,251	1,108	1,122	1,182	1,021	570	78	12
1954	8,861	125	660	785	1,413	1,085	1,053	1,068	1,071	968	520	100	13
1953	9,393	140	678	792	1,652	1,188	1,094	1,115	1,082	1,015	523	105	9
1952	9,258	145	666	766	1,656	1,177	1,107	1,133	1,110	904	493	90	11
1951	8,921	131	610	757	1,491	1,198	1,057	1,066	1,091	952	484	75	9
1950	8,247	118	638	685	1,405	1,128	920	943	996	887	446	75	6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MOTOR VEHICLE ACCIDENTS (E810-E835)</b>													
<b>WHITE, BOTH SEXES</b>													
1959	33,483	285	1,290	2,312	8,123	4,888	3,963	3,817	3,408	3,240	1,829	309	19
1958	32,730	262	1,261	2,385	7,570	4,834	4,059	3,647	3,331	3,262	1,788	309	22
1957	34,084	293	1,211	2,255	7,840	5,336	4,233	3,839	3,632	3,299	1,832	295	19
1956	34,745	269	1,203	2,291	8,251	5,496	4,237	3,881	3,592	3,390	1,793	322	20
1955	33,854	280	1,304	2,111	7,768	5,526	4,232	3,692	3,663	3,186	1,763	293	36
1954	31,357	241	1,337	2,050	6,816	5,148	3,830	3,484	3,365	3,018	1,747	289	32
1953	33,372	281	1,436	2,045	7,348	5,468	4,084	3,692	3,636	3,256	1,775	322	29
1952	33,365	272	1,413	2,031	7,274	5,495	4,235	3,689	3,736	3,239	1,683	262	36
1951	32,724	252	1,394	2,007	6,926	5,449	4,183	3,688	3,638	3,185	1,691	273	38
1950	30,955	232	1,337	1,891	6,891	5,059	3,731	3,331	3,515	3,051	1,637	243	37
<b>WHITE, MALE</b>													
1959	24,675	135	735	1,571	6,579	3,903	3,002	2,766	2,350	2,177	1,232	211	14
1958	24,198	146	737	1,590	6,125	3,939	3,101	2,631	2,298	2,178	1,212	225	16
1957	25,416	157	697	1,513	6,375	4,355	3,207	2,787	2,538	2,270	1,282	218	17
1956	26,034	150	659	1,565	6,687	4,461	3,230	2,874	2,561	2,332	1,272	227	16
1955	25,336	151	732	1,438	6,281	4,463	3,287	2,700	2,582	2,237	1,218	220	27
1954	23,511	138	783	1,361	5,565	4,228	2,937	2,539	2,388	2,103	1,252	195	22
1953	25,012	167	848	1,359	5,876	4,465	3,163	2,701	2,625	2,287	1,276	223	22
1952	25,110	144	836	1,343	5,813	4,510	3,256	2,699	2,704	2,387	1,214	178	26
1951	24,797	142	862	1,338	5,621	4,411	3,283	2,753	2,633	2,293	1,227	203	31
1950	23,574	128	773	1,303	5,651	4,081	2,943	2,479	2,589	2,208	1,214	173	32
<b>WHITE, FEMALE</b>													
1959	8,808	150	555	741	1,544	985	961	1,051	1,058	1,063	597	98	5
1958	8,532	116	524	795	1,445	895	958	1,016	1,033	1,084	576	84	6
1957	8,668	136	514	742	1,465	981	1,026	1,052	1,094	1,029	550	77	2
1956	8,711	119	544	726	1,564	1,035	1,007	1,007	1,031	1,058	521	95	4
1955	8,518	129	572	673	1,487	1,063	945	992	1,081	949	545	73	9
1954	7,846	103	554	689	1,251	920	893	945	977	915	495	94	10
1953	8,360	114	588	686	1,472	1,003	921	991	1,011	969	499	99	7
1952	8,255	128	577	688	1,461	985	979	990	1,032	852	469	84	10
1951	7,927	110	532	669	1,305	1,038	900	935	1,005	892	464	70	7
1950	7,381	104	564	588	1,240	978	788	852	926	843	423	70	5

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
<b>MOTOR VEHICLE ACCIDENTS (E810-E835)</b>													
<b>ALL OTHER, BOTH SEXES</b>													
1959	4,427	34	233	407	846	811	696	629	409	247	81	21	13
1958	4,251	31	237	325	818	822	699	557	387	273	84	11	7
1957	4,618	44	237	349	827	944	717	611	463	261	126	22	17
1956	4,883	63	235	349	918	987	831	627	473	267	100	17	16
1955	4,572	48	243	295	888	944	746	594	423	269	91	22	9
1954	4,229	55	231	282	755	868	675	599	400	229	97	27	11
1953	4,583	51	251	323	821	973	777	616	374	252	115	23	7
1952	4,429	36	230	264	841	942	708	639	399	233	100	27	10
1951	4,272	35	194	293	787	887	734	572	378	259	98	20	15
1950	3,808	32	166	261	709	796	628	517	365	213	90	17	14
<b>ALL OTHER, MALE</b>													
1959	3,389	18	136	279	678	641	550	495	325	176	61	19	11
1958	3,294	17	131	237	664	668	551	433	310	209	60	9	5
1957	3,484	25	141	229	642	744	560	468	351	194	99	18	13
1956	3,770	37	141	220	745	808	658	491	348	211	80	16	15
1955	3,467	26	135	192	703	756	583	464	322	197	66	17	6
1954	3,214	33	125	186	593	703	515	476	306	176	72	21	8
1953	3,550	25	161	217	641	788	604	492	303	206	91	17	5
1952	3,426	19	141	186	646	750	580	496	321	181	76	21	9
1951	3,278	14	116	205	601	727	577	441	292	199	78	15	13
1950	2,942	18	92	164	544	646	496	426	295	169	67	12	13
<b>ALL OTHER, FEMALE</b>													
1959	1,038	16	97	128	168	170	146	134	84	71	20	2	2
1958	957	14	106	88	154	154	148	124	77	64	24	2	2
1957	1,134	19	96	120	185	200	157	143	112	67	27	4	4
1956	1,113	26	94	129	173	179	173	136	125	56	20	1	1
1955	1,105	22	108	103	185	188	163	130	101	72	25	5	3
1954	1,015	22	106	96	162	165	160	123	94	53	25	6	3
1953	1,033	26	90	106	180	185	173	124	71	46	24	6	2
1952	1,003	17	89	78	195	192	128	143	78	52	24	6	1
1951	994	21	78	88	186	160	157	131	86	60	20	5	2
1950	866	14	74	97	165	150	132	91	70	44	23	5	1

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER ACCIDENTS (E800-E802, E840-E962)													
BOTH SEXES													
1959	54,170	3,538	3,368	3,792	4,300	4,257	5,051	5,526	5,148	5,988	7,640	5,490	72
1958	53,623	3,623	3,375	3,804	4,356	4,218	5,026	5,164	5,009	5,896	7,770	5,314	68
1957	56,605	3,337	3,301	3,850	4,306	4,548	5,171	5,555	5,395	6,516	8,562	5,887	177
1956	55,152	3,166	3,237	3,676	4,376	4,431	5,004	5,286	5,348	6,503	8,317	5,708	100
1955	55,017	2,980	3,244	3,693	4,086	4,434	5,029	5,522	5,305	6,474	8,459	5,690	101
1954	54,446	3,284	3,232	3,607	4,230	4,591	4,911	5,243	5,208	6,405	8,286	5,370	79
1953	57,077	3,297	3,362	3,768	4,668	4,970	5,150	5,603	5,558	6,419	8,677	5,479	126
1952	58,378	3,422	3,498	3,685	4,671	5,116	5,454	5,668	5,761	6,554	8,853	5,582	114
1951	58,875	3,424	3,470	3,592	4,653	5,308	5,802	5,549	5,785	6,774	8,858	5,525	135
1950	56,486	3,331	3,291	3,367	4,519	5,009	5,440	5,351	5,586	6,486	8,620	5,351	135
MALE													
1959	35,575	1,977	1,979	2,794	3,758	3,605	4,154	4,371	3,926	3,670	3,366	1,910	65
1958	34,997	2,079	1,946	2,702	3,800	3,619	4,171	4,161	3,765	3,491	3,345	1,863	55
1957	36,747	1,883	1,842	2,816	3,751	3,881	4,253	4,471	4,046	3,986	3,665	2,031	122
1956	36,019	1,819	1,909	2,677	3,808	3,801	4,107	4,279	4,007	3,998	3,541	1,994	79
1955	35,887	1,722	1,870	2,689	3,553	3,793	4,096	4,525	4,023	4,014	3,633	1,887	82
1954	35,590	1,875	1,862	2,659	3,676	3,949	4,048	4,229	3,944	3,905	3,504	1,874	65
1953	37,684	1,887	1,942	2,755	4,057	4,275	4,280	4,612	4,299	3,883	3,708	1,883	103
1952	38,161	1,929	1,990	2,639	3,992	4,358	4,526	4,652	4,461	3,918	3,708	1,895	93
1951	38,798	1,922	1,991	2,667	3,985	4,567	4,871	4,593	4,453	4,145	3,608	1,886	110
1950	36,899	1,911	1,958	2,434	3,893	4,275	4,502	4,353	4,274	3,883	3,536	1,763	117
FEMALE													
1959	18,595	1,561	1,389	998	542	652	897	1,155	1,222	2,318	4,274	3,580	7
1958	18,626	1,544	1,429	1,102	556	599	855	1,003	1,244	2,405	4,425	3,451	13
1957	19,858	1,454	1,459	1,034	555	667	918	1,084	1,349	2,530	4,897	3,856	55
1956	19,133	1,347	1,328	999	568	630	897	1,007	1,341	2,505	4,776	3,714	21
1955	19,130	1,258	1,374	1,004	533	641	933	997	1,282	2,460	4,826	3,803	19
1954	18,856	1,409	1,370	948	554	642	863	1,014	1,264	2,500	4,782	3,496	14
1953	19,393	1,410	1,420	1,013	611	695	870	991	1,259	2,536	4,969	3,596	23
1952	20,217	1,493	1,508	1,046	679	758	928	1,016	1,300	2,636	5,145	3,687	21
1951	20,077	1,502	1,479	925	668	741	931	956	1,332	2,629	5,250	3,639	25
1950	19,587	1,420	1,333	933	626	734	938	998	1,312	2,603	5,084	3,588	18

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER ACCIDENTS (E800-E802, E840-E962)													
WHITE, BOTH SEXES													
1959	45,741	2,375	2,365	2,912	3,535	3,437	4,069	4,654	4,397	5,379	7,259	5,303	56
1958	45,190	2,409	2,350	2,961	3,556	3,408	4,135	4,338	4,262	5,275	7,345	5,102	49
1957	48,081	2,238	2,364	2,996	3,521	3,634	4,220	4,628	4,696	5,868	8,149	5,685	82
1956	47,161	2,120	2,412	2,892	3,594	3,620	4,025	4,449	4,645	5,885	7,931	5,515	73
1955	47,156	2,073	2,305	2,943	3,385	3,593	4,113	4,601	4,619	5,867	8,070	5,505	82
1954	46,738	2,279	2,368	2,905	3,460	3,716	4,067	4,400	4,612	5,790	7,889	5,191	61
1953	48,961	2,352	2,456	3,097	3,816	4,020	4,189	4,665	4,866	5,850	8,269	5,289	92
1952	50,028	2,404	2,564	2,989	3,820	4,166	4,432	4,704	5,053	5,956	8,446	5,410	84
1951	50,932	2,461	2,614	2,886	3,857	4,469	4,813	4,630	5,096	6,211	8,481	5,307	107
1950	49,079	2,443	2,487	2,685	3,730	4,182	4,475	4,552	4,968	5,953	8,315	5,176	113
WHITE, MALE													
1959	29,918	1,367	1,438	2,212	3,103	2,961	3,396	3,703	3,367	3,302	3,172	1,846	51
1958	29,471	1,436	1,438	2,175	3,127	2,965	3,478	3,524	3,235	3,151	3,126	1,776	40
1957	31,060	1,295	1,383	2,238	3,094	3,143	3,519	3,762	3,544	3,604	3,468	1,951	59
1956	30,611	1,246	1,475	2,160	3,153	3,147	3,340	3,623	3,498	3,653	3,354	1,906	56
1955	30,640	1,237	1,402	2,194	2,965	3,148	3,372	3,794	3,539	3,650	3,453	1,820	66
1954	30,438	1,358	1,418	2,174	3,031	3,253	3,401	3,582	3,528	3,544	3,310	1,790	49
1953	32,164	1,378	1,473	2,283	3,359	3,526	3,529	3,887	3,794	3,553	3,493	1,813	76
1952	32,495	1,378	1,511	2,195	3,282	3,603	3,712	3,895	3,955	3,575	3,497	1,823	69
1951	33,417	1,432	1,534	2,171	3,349	3,910	4,097	3,866	3,953	3,792	3,420	1,802	91
1950	31,838	1,432	1,537	1,964	3,240	3,637	3,737	3,716	3,827	3,576	3,378	1,693	101
WHITE, FEMALE													
1959	15,823	1,008	927	700	432	476	673	951	1,030	2,077	4,087	3,457	5
1958	15,719	973	912	786	429	443	657	814	1,027	2,124	4,219	3,326	9
1957	17,021	943	981	758	427	491	701	866	1,152	2,264	4,681	3,734	23
1956	16,550	874	937	732	441	473	685	826	1,147	2,232	4,577	3,609	17
1955	16,516	836	903	749	420	445	741	807	1,080	2,217	4,617	3,685	16
1954	16,300	921	950	731	429	463	666	818	1,084	2,246	4,579	3,401	12
1953	16,797	974	983	814	457	494	660	778	1,072	2,297	4,776	3,476	16
1952	17,533	1,026	1,053	794	538	563	720	809	1,098	2,381	4,949	3,587	15
1951	17,515	1,029	1,080	715	508	559	716	764	1,143	2,419	5,061	3,505	16
1950	17,241	1,011	950	721	490	545	738	836	1,141	2,377	4,937	3,483	12

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
OTHER ACCIDENTS (E800-E802, E840-E962)													
ALL OTHER, BOTH SEXES													
1959	8,429	1,163	1,003	880	765	820	982	872	751	609	381	187	16
1958	8,433	1,214	1,025	843	800	810	891	826	747	621	425	212	19
1957	8,524	1,099	937	854	785	914	951	927	699	648	413	202	95
1956	7,991	1,046	825	784	782	811	979	837	703	618	386	193	27
1955	7,861	907	939	750	701	841	916	921	686	607	389	185	19
1954	7,708	1,005	864	702	770	875	844	843	596	615	397	179	18
1953	8,116	945	906	671	852	950	961	938	692	569	408	190	34
1952	8,350	1,018	934	696	851	950	1,022	964	708	598	407	172	30
1951	7,943	963	856	706	796	839	989	919	689	563	377	218	28
1950	7,407	888	804	682	789	827	965	799	618	533	305	175	22
ALL OTHER, MALE													
1959	5,657	610	541	582	655	644	758	668	559	368	194	64	14
1958	5,526	643	508	527	673	654	693	637	530	340	219	87	15
1957	5,687	588	459	578	657	738	734	709	502	382	197	80	63
1956	5,408	573	434	517	655	654	767	656	509	345	187	88	23
1955	5,247	485	468	495	588	645	724	731	484	364	180	67	16
1954	5,152	517	444	485	645	696	647	647	416	361	194	84	16
1953	5,520	509	469	472	698	749	751	725	505	330	215	70	27
1952	5,666	551	479	444	710	755	814	757	506	343	211	72	24
1951	5,381	490	457	496	636	657	774	727	500	353	188	84	19
1950	5,061	479	421	470	653	638	765	637	447	307	158	70	16
ALL OTHER, FEMALE													
1959	2,772	553	462	298	110	176	224	204	192	241	187	123	2
1958	2,907	571	517	316	127	156	198	189	217	281	206	125	4
1957	2,837	511	478	276	128	176	217	218	197	266	216	122	32
1956	2,583	473	391	267	127	157	212	181	194	273	199	105	4
1955	2,614	422	471	255	113	196	192	190	202	243	209	118	3
1954	2,556	488	420	217	125	179	197	196	180	254	203	95	2
1953	2,596	436	437	199	154	201	210	213	187	239	193	120	7
1952	2,684	467	455	252	141	195	208	207	202	255	196	100	6
1951	2,562	473	399	210	160	182	215	192	189	210	189	134	9
1950	2,346	409	383	212	136	189	200	162	171	226	147	105	6

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SUICIDE (E963, E970-E979)													
BOTH SEXES													
1959	18,633	...	...	86	1,152	2,255	3,236	3,992	3,733	2,681	1,257	231	10
1958	18,519	...	...	77	1,088	2,273	3,236	4,115	3,645	2,648	1,209	222	6
1957	16,632	...	...	68	880	2,034	2,970	3,545	3,353	2,435	1,121	219	7
1956	16,727	...	...	54	859	2,024	2,797	3,547	3,580	2,533	1,139	187	7
1955	16,760	...	...	39	854	2,013	2,800	3,679	3,615	2,446	1,081	214	19
1954	16,356	...	...	37	865	2,094	2,815	3,559	3,427	2,389	973	183	14
1953	15,947	...	...	58	909	2,027	2,806	3,401	3,164	2,346	1,026	193	17
1952	15,567	...	...	40	881	2,023	2,725	3,261	3,131	2,297	989	204	16
1951	15,909	...	...	52	940	2,041	2,886	3,306	3,162	2,366	948	188	20
1950	17,145	...	...	38	997	2,156	3,072	3,628	3,585	2,467	1,021	166	15
MALE													
1959	14,441	...	...	77	898	1,619	2,385	3,086	2,937	2,147	1,074	211	7
1958	14,366	...	...	65	821	1,610	2,433	3,136	2,912	2,141	1,054	189	5
1957	12,951	...	...	55	684	1,463	2,188	2,751	2,617	2,008	981	198	6
1956	12,968	...	...	45	654	1,477	2,060	2,675	2,827	2,064	992	167	7
1955	12,961	...	...	29	641	1,445	2,092	2,770	2,885	1,989	909	184	17
1954	12,964	...	...	30	672	1,570	2,091	2,840	2,770	1,977	849	154	11
1953	12,534	...	...	48	663	1,476	2,128	2,640	2,539	1,970	883	173	14
1952	12,115	...	...	31	664	1,443	2,010	2,482	2,524	1,905	860	181	15
1951	12,300	...	...	43	684	1,457	2,094	2,504	2,560	1,975	800	169	14
1950	13,297	...	...	31	706	1,554	2,253	2,771	2,922	2,031	878	138	13
FEMALE													
1959	4,192	...	...	9	254	636	851	906	796	534	183	20	3
1958	4,153	...	...	12	267	663	803	979	733	507	155	33	1
1957	3,681	...	...	13	196	571	782	794	736	427	140	21	1
1956	3,759	...	...	9	205	547	737	872	753	469	147	20	-
1955	3,799	...	...	10	213	568	708	909	730	457	172	30	2
1954	3,392	...	...	7	193	524	724	719	657	412	124	29	3
1953	3,413	...	...	10	246	551	678	761	625	376	143	20	3
1952	3,452	...	...	9	217	580	715	779	607	392	129	23	1
1951	3,609	...	...	9	256	584	792	802	602	391	148	19	6
1950	3,848	...	...	7	291	602	819	857	663	436	143	28	2

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SUICIDE (E963, E970-E979)													
WHITE, BOTH SEXES													
1959	17,719	...	...	80	1,029	2,030	3,073	3,833	3,607	2,609	1,221	227	10
1958	17,684	...	...	72	992	2,071	3,058	3,968	3,529	2,583	1,188	217	6
1957	15,878	...	...	67	789	1,843	2,819	3,427	3,255	2,361	1,097	213	7
1956	16,034	...	...	52	770	1,858	2,639	3,432	3,487	2,485	1,122	184	5
1955	16,092	...	...	37	749	1,859	2,670	3,562	3,527	2,397	1,063	210	18
1954	15,652	...	...	36	779	1,904	2,672	3,436	3,343	2,331	956	181	14
1953	15,307	...	...	52	833	1,879	2,673	3,278	3,089	2,296	1,000	190	17
1952	14,963	...	...	33	812	1,860	2,599	3,164	3,045	2,259	976	199	16
1951	15,252	...	...	47	845	1,882	2,739	3,192	3,087	2,325	930	187	18
1950	16,468	...	...	35	909	2,001	2,927	3,499	3,492	2,417	1,010	163	15
WHITE, MALE													
1959	13,724	...	...	73	808	1,446	2,257	2,956	2,840	2,088	1,041	208	7
1958	13,707	...	...	61	750	1,458	2,290	3,019	2,815	2,086	1,038	185	5
1957	12,331	...	...	54	612	1,307	2,066	2,645	2,544	1,942	960	195	6
1956	12,427	...	...	44	582	1,347	1,939	2,588	2,754	2,027	977	164	5
1955	12,430	...	...	27	561	1,330	1,983	2,678	2,812	1,950	893	180	16
1954	12,396	...	...	29	607	1,418	1,972	2,742	2,702	1,929	833	153	11
1953	12,008	...	...	46	614	1,357	2,014	2,533	2,474	1,928	858	170	14
1952	11,624	...	...	26	612	1,317	1,913	2,399	2,449	1,868	848	177	15
1951	11,784	...	...	38	622	1,332	1,978	2,409	2,499	1,942	784	168	12
1950	12,755	...	...	29	641	1,436	2,133	2,675	2,835	1,988	870	135	13
WHITE, FEMALE													
1959	3,995	...	...	7	221	584	816	877	767	521	180	19	3
1958	3,977	...	...	11	242	613	768	949	714	497	150	32	1
1957	3,547	...	...	13	177	536	753	782	711	419	137	18	1
1956	3,607	...	...	8	188	511	700	844	733	458	145	20	-
1955	3,662	...	...	10	188	529	687	884	715	447	170	30	2
1954	3,256	...	...	7	172	486	700	694	641	402	123	28	3
1953	3,299	...	...	6	219	522	659	745	615	368	142	20	3
1952	3,339	...	...	7	200	543	686	765	596	391	128	22	1
1951	3,468	...	...	9	223	550	761	783	588	383	146	19	6
1950	3,713	...	...	6	268	565	794	824	657	429	140	28	2

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
SUICIDE (E963, E970-E979)													
ALL OTHER, BOTH SEXES													
1959	914	...	...	6	123	225	163	159	126	72	36	4	-
1958	835	...	...	5	96	202	178	147	116	65	21	5	-
1957	754	...	...	1	91	191	151	118	98	74	24	6	-
1956	693	...	...	2	89	166	158	115	93	48	17	3	2
1955	668	...	...	2	105	154	130	117	88	49	18	4	1
1954	704	...	...	1	86	190	143	123	84	58	17	2	-
1953	640	...	...	6	76	148	133	123	75	50	26	3	-
1952	604	...	...	7	69	163	126	97	86	38	13	5	-
1951	657	...	...	5	95	159	147	114	75	41	18	1	2
1950	677	...	...	3	88	155	145	129	93	50	11	3	-
ALL OTHER, MALE													
1959	717	...	...	4	90	173	128	130	97	59	33	3	-
1958	659	...	...	4	71	152	143	117	97	55	16	4	-
1957	620	...	...	1	72	156	122	106	73	66	21	3	-
1956	541	...	...	1	72	130	121	87	73	37	15	3	2
1955	531	...	...	2	80	115	109	92	73	39	16	4	1
1954	568	...	...	1	65	152	119	98	68	48	16	1	-
1953	526	...	...	2	49	119	114	107	65	42	25	3	-
1952	491	...	...	5	52	126	97	83	75	37	12	4	-
1951	516	...	...	5	62	125	116	95	61	33	16	1	2
1950	542	...	...	2	65	118	120	96	87	43	8	3	-
ALL OTHER, FEMALE													
1959	197	...	...	2	33	52	35	29	29	13	3	1	-
1958	176	...	...	1	25	50	35	30	19	10	5	1	-
1957	134	...	...	-	19	35	29	12	25	8	3	3	-
1956	152	...	...	1	17	36	37	28	20	11	2	-	-
1955	137	...	...	-	25	39	21	25	15	10	2	-	-
1954	136	...	...	-	21	38	24	25	16	10	1	1	-
1953	114	...	...	4	27	29	19	16	10	8	1	-	-
1952	113	...	...	2	17	37	29	14	11	1	1	1	-
1951	141	...	...	-	33	34	31	19	14	8	2	-	-
1950	135	...	...	1	23	37	25	33	6	7	3	-	-

- Quantity zero

... Not applicable

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HOMICIDE (E964, E980-E985)													
BOTH SEXES													
1959	8,159	150	137	186	1,375	2,125	1,913	1,195	618	309	112	23	16
1958	7,815	135	110	161	1,319	2,018	1,757	1,247	602	310	113	28	15
1957	7,641	123	88	138	1,255	2,118	1,750	1,140	600	305	91	12	21
1956	7,629	125	104	141	1,290	2,144	1,756	1,062	600	277	89	18	23
1955	7,418	129	76	131	1,162	2,102	1,749	1,106	586	232	99	23	23
1954	7,735	134	90	150	1,241	2,178	1,837	1,201	498	264	101	14	27
1953	7,640	110	93	127	1,236	2,149	1,928	1,113	487	259	84	25	29
1952	8,054	127	88	120	1,301	2,380	2,012	1,095	524	271	94	13	29
1951	7,495	141	79	111	1,264	2,178	1,788	1,043	537	234	83	13	24
1950	7,942	137	77	133	1,393	2,344	1,879	1,059	531	271	85	13	20
MALE													
1959	6,068	86	75	106	1,048	1,564	1,424	937	494	232	78	12	12
1958	5,804	69	52	96	979	1,560	1,297	952	478	230	61	18	12
1957	5,739	68	41	87	947	1,620	1,301	884	463	243	63	6	16
1956	5,705	60	50	98	977	1,619	1,333	804	468	210	59	13	14
1955	5,630	71	36	83	875	1,617	1,341	854	474	187	66	14	12
1954	5,886	74	47	86	933	1,688	1,426	922	416	199	70	8	17
1953	5,828	55	48	75	956	1,631	1,510	887	382	192	65	9	18
1952	6,202	68	54	74	992	1,840	1,553	871	426	219	74	9	22
1951	5,669	66	41	57	936	1,648	1,394	822	444	183	55	8	15
1950	6,089	72	33	75	1,051	1,799	1,464	860	436	214	61	6	18
FEMALE													
1959	2,091	64	62	80	327	561	489	258	124	77	34	11	4
1958	2,011	66	58	65	340	458	460	295	124	80	52	10	3
1957	1,902	55	47	51	308	498	449	256	137	62	28	6	5
1956	1,924	65	54	43	313	525	423	258	132	67	30	5	9
1955	1,788	58	40	48	287	485	408	252	112	45	33	9	11
1954	1,849	60	43	64	308	490	411	279	82	65	31	6	10
1953	1,812	55	45	52	280	518	418	226	105	67	19	16	11
1952	1,852	59	34	46	309	540	459	224	98	52	20	4	7
1951	1,826	75	38	54	328	530	394	221	93	51	28	5	9
1950	1,853	65	44	58	342	545	415	199	95	57	24	7	2

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)



TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HOMICIDE (E964, E980-E985)													
WHITE, BOTH SEXES													
1959	3,856	107	106	140	573	791	810	624	375	219	86	15	10
1958	3,651	92	82	125	554	730	732	626	365	224	93	24	4
1957	3,444	91	54	95	501	783	727	528	375	198	68	10	14
1956	3,383	84	79	97	502	768	674	525	372	189	73	11	9
1955	3,361	95	53	93	438	739	767	551	348	164	83	15	15
1954	3,480	95	68	108	485	766	783	595	299	176	73	11	21
1953	3,425	88	74	89	452	784	774	568	311	180	67	20	18
1952	3,502	99	71	89	473	789	798	559	328	195	69	13	19
1951	3,363	112	56	87	454	783	737	526	343	176	66	10	13
1950	3,538	112	58	88	484	774	826	556	348	200	69	10	13
WHITE, MALE													
1959	2,716	57	58	76	428	549	568	457	292	159	56	7	9
1958	2,555	45	36	77	393	557	497	451	276	160	46	14	3
1957	2,426	54	23	55	362	580	497	376	269	153	43	4	10
1956	2,421	42	41	65	369	563	487	369	286	142	45	7	5
1955	2,439	55	25	56	320	550	570	391	271	129	55	10	7
1954	2,477	49	34	57	348	561	570	426	231	129	52	7	13
1953	2,449	42	38	50	337	560	561	426	236	126	51	9	13
1952	2,562	56	42	55	342	588	568	424	255	155	54	9	14
1951	2,421	53	28	44	332	577	536	392	272	134	43	5	5
1950	2,586	60	26	48	357	567	613	428	269	154	49	4	11
WHITE, FEMALE													
1959	1,140	50	48	64	145	242	242	167	83	60	30	8	1
1958	1,096	47	46	48	161	173	235	175	89	64	47	10	1
1957	1,018	37	31	40	139	203	230	152	106	45	25	6	4
1956	962	42	38	32	133	205	187	156	86	47	28	4	4
1955	922	40	28	37	118	189	197	160	77	35	28	5	8
1954	1,003	46	34	51	137	205	213	169	68	47	21	4	8
1953	976	46	36	39	115	224	213	142	75	54	16	11	5
1952	940	43	29	34	131	201	230	135	73	40	15	4	5
1951	942	59	28	43	122	206	201	134	71	42	23	5	8
1950	952	52	32	40	127	207	213	128	79	46	20	6	2

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)

TABLE 290F. DEATHS FOR APPROXIMATELY 64 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 1950-59

CAUSE OF DEATH CODE, RACE, SEX, AND YEAR	Total	Under 1 year	1-4 Years	5-14 Years	15-24 Years	25-34 Years	35-44 Years	45-54 Years	55-64 Years	65-74 Years	75-84 Years	85+ Years	N.S.
HOMICIDE (E964, E980-E985)													
ALL OTHER, BOTH SEXES													
1959	4,303	43	31	46	802	1,334	1,103	571	243	90	26	8	6
1958	4,164	43	28	36	765	1,288	1,025	621	237	86	20	4	11
1957	4,197	32	34	43	754	1,335	1,023	612	225	107	23	2	7
1956	4,246	41	25	44	788	1,376	1,082	537	228	88	16	7	14
1955	4,057	34	23	38	724	1,363	982	555	238	68	16	8	8
1954	4,255	39	22	42	756	1,412	1,054	606	199	88	28	3	6
1953	4,215	22	19	38	784	1,365	1,154	545	176	79	17	5	11
1952	4,552	28	17	31	828	1,591	1,214	536	196	76	25	-	10
1951	4,132	29	23	24	810	1,395	1,051	517	194	58	17	3	11
1950	4,404	25	19	45	909	1,570	1,053	503	183	71	16	3	7
ALL OTHER, MALE													
1959	3,352	29	17	30	620	1,015	856	480	202	73	22	5	3
1958	3,249	24	16	19	586	1,003	800	501	202	70	15	4	9
1957	3,313	14	18	32	585	1,040	804	508	194	90	20	2	6
1956	3,284	18	9	33	608	1,056	846	435	182	68	14	6	9
1955	3,191	16	11	27	555	1,067	771	463	203	58	11	4	5
1954	3,409	25	13	29	585	1,127	856	496	185	70	18	1	4
1953	3,379	13	10	25	619	1,071	949	461	146	66	14	-	5
1952	3,640	12	12	19	650	1,252	985	447	171	64	20	-	8
1951	3,248	13	13	13	604	1,071	858	430	172	49	12	3	10
1950	3,503	12	7	27	694	1,232	851	432	167	60	12	2	7
ALL OTHER, FEMALE													
1959	951	14	14	16	182	319	247	91	41	17	4	3	3
1958	915	19	12	17	179	285	225	120	35	16	5	-	2
1957	884	18	16	11	169	295	219	104	31	17	3	-	1
1956	962	23	16	11	180	320	236	102	46	20	2	1	5
1955	866	18	12	11	169	296	211	92	35	10	5	4	3
1954	846	14	9	13	171	285	198	110	14	18	10	2	2
1953	836	9	9	13	165	294	205	84	30	13	3	5	6
1952	912	16	5	12	178	339	229	89	25	12	5	-	2
1951	884	16	10	11	206	324	193	87	22	9	5	-	1
1950	901	13	12	18	215	338	202	71	16	11	4	1	-

- Quantity zero

Source: CDC/NCHS, National Vital Statistics System, Unpublished trend table, and Vital Statistics of the United States (1950-1959)