

National Bureau of Standards

Certificate

Standard Reference Material 4339 Radioactivity Standard

Radionuclide Radium-228

Source identification 4339-

Source description Liquid in flame-sealed NBS

borosilicate-glass ampoule (1)*

Solution composition Radium-228 and progeny plus

approximately 13 μg of non-radioactive barium per gram of

1-molar nitric acid (2)

Mass grams

Radioactivity concentration 287.0 Bq g⁻¹

Reference time 1200 EST July 1, 1988

Overall uncertainty 2.76 percent (3)

Photon-emitting impurities 226 Ra: 0.9 ± 0.1 Bq g^{-1 (4)}

Half life 5.75 ± 0.03 years ⁽⁵⁾

Measuring instrument Calibrated germanium spectrometer

systems

This Standard Reference Material was prepared in the Center for Radiation Research, Ionizing Radiation Division, Radioactivity Group, Dale D. Hoppes, Group Leader.

Gaithersburg, MD 20899 July, 1988 Stanley D. Rasberry, Chief Office of Standard Reference Materials

*Notes on back

NOTES

(1) Approximately five milliliters of solution. Ampoule specifications:

- (2) Solution density 1.030 \pm 0.003 g/mL at 24.4 °C.
- (3) The overall uncertainty was formed by taking three times the quadratic combination of standard deviations of the mean, or assumed approximations thereof, for the following:

a)	photon-emission-rate measurements	0.26	percent
b)	gravimetric measurements	0.05	percent
c)	source positioning	0.20	percent
d)	detector efficiency	0.50	percent
e)	decay-scheme data	0.61	percent
f)	half-life	0.33	percent

- (4) The limit of detection for photon-emitting impurities is:
 - $0.3~\gamma~s^{-1}~g^{-1}$ for energies between 50 and 1900 KeV, provided that the impurity photons are separated in energy by 5 keV or more from photons emitted in the decay of radium-228 and progeny.
- (5) NCRP Report No. 58, 2nd edition, February 1985, p. 365.

For further information please contact Dr. Larry Lucas, (301) 975-5546.