Department of Commerce
Malcolus Baldrige
Secretary
onal Burnau of Standards
rnest Ambler. Director

National Bureau of Standards

Certificate of Analysis

Standard Reference Material 948

Plutonium Isotopic Standard

This Standard Reference Material (SRM) is certified as an isotopic standard for use in isotopic measurements of plutonium. SRM 948 consists of approximately 250 mg of plutonium in the form of plutonium sulfate tetrahydrate packaged in a glass microbottle.

	Pu	²³⁹ Pu	²⁴⁰ Pu	²⁴¹ Pu	Pu
Atom Percent*	0.010	91.736	7.922	0.299	0.0330
	± 0.001	± 0.010	± 0.010	± 0.001	± 0.0003

^{*}As of January 1, 1982, refer to Table 1 for quarterly decay-adjusted values.

The plutonium isotopic distribution was determined by thermal ionization mass spectrometry at the National Bureau of Standards (NBS) on samples from which americium and uranium were removed. Because high-purity plutonium isotopes have not been used to prepare known synthetic mixtures, the accuracy is dependent on uranium and plutonium exhibiting similar behavior. The observed mass spectrometer data were corrected for mass discrimination effects using data from the analysis of uranium isotopic SRM's that had been analyzed under similar conditions. In addition, the value for ²³⁸Pu was checked by alpha-count of ²³⁸Pu, using the known value for ²³⁹Pu as a ratio check.

SRM 948 contains uranium and americium isotopes, including growing-in daughters of plutonium that are isobaric with the plutonium isotopes. In addition, there may be radiation damage to the glass bottle and teflon cap liner resulting in small glass slivers. Therefore, in its use, a chemical separation that provides a purified plutonium fraction is essential to the attainment of high accuracy.

Measurements leading to the certification of this SRM were made in the Inorganic Analytical Research Division by E.L. Garner, L.A. Machlan, and W.R. Shields.

The technical and support aspects involved in the revision of this Certificate were coordinated through the Office of Standard Reference Materials by T.E. Gills.

August 19, 1982 Washington, D.C. 20234 (Revision of Certificate dated 12-3-71) George A. Uriano, Chief
Office of Standard Reference Materials

(over)

The decay-adjusted values for the plutonium isotopic composition, in atom percent, are tabulated below in Table 1. The half-life values, in years, used for the decay-adjustment are: ²³⁸Pu, 87.74; ²³⁹Pu, 24,119; ²⁴⁰Pu, 6,560; ²⁴¹Pu, 14.34; and ²⁴²Pu, 387,000.

Table 1
Decay-Adjusted Plutonium Isotopic Composition
Atom Percent

Date	²³⁸ Pu	²³⁹ Pu	Pu	²⁴¹ Pu	²⁴² Pu
January 1, 1982	0.010	91.736	7.922	0.299	0.0330
April 1, 1982	.010	91.739	7.922	.295	.0330
July 1, 1982	.010	91.743	7.923	.292	.0330
October 1, 1982	.010	91.746	7.923	.288	.0330
January 1, 1983	.010	91.749	7.923	.285	.0330
April 1, 1983	.010	91.753	7.923	.281	.0330
July 1, 1983	.010	91.756	7.923	.278	.0330
October 1, 1983	.010	91.759	7.923	.274	.0330
January 1, 1984	.010	91.762	7.923	.271	.0330
April 1, 1984	.010	91.765	7.923	.268	.0330
July 1, 1984	.010	91.769	7.924	.265	.0330
October 1, 1984	.010	91.772	7.924	.262	.0330
January 1, 1985	.010	91.775	7.924	.258	.0330
April 1, 1985	.010	91.778	7.924	.255	.0330
July 1, 1985	.010	91.781	7.924	.252	.0330
October 1, 1985	.010	91.784	7.924	.249	.0330
January 1, 1986	.010	91.787	7.924	.246	.0330
April 1, 1986	.010	91.789	7.924	.243	.0330
July 1, 1986	.010	91.792	7.924	.240	.0330
October 1, 1986	.010	91.795	7.925	.238	.0330
95% Confidence Limit:	±0.001	±0.010	±0.010	±0.001	±0.0003