

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
AARHUS	2	2	.	.	1	1	6
ABAEETETUBA	1	.	1	.	3	2	7
ABERDEEN	1	1	.	.	.	2	4
ABONY	1	3	.	.	4
ADELAIDE	2	27	22	4	17	3	8	1	11		95
AEQUATORIA	1	1
AFLAO	1	1
AFRICANA	6	6
AGAMA	2	2
AGBENI	.	.	1	1
AGONA	50	99	97	36	49	12	50	12	123		528
AGOUEVE	.	2	2
ALABAMA	1	3	4
ALACHUA	3	4	2	1	2	1	5	.	4		22
ALBANY	.	1	4	1	1	2	.	3	5		17
ALTONA	1		1
AMAGER	.	1	.	1	2		4
AMSTERDAM	1	4	1		6
ANATUM	10	19	26	8	30	4	16	17	27		157
ANECHO	.	.	.	1	.	.	.	1	.		2

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
APAPA	1	.	1	2
APEYEME	1	1
ARECHAVALETA	2	1	.	.	.	3
ARKANSAS	4	.	.	.	4
ASSEN	1	.	.	.	1
AZTECA	.	.	.	1	1
BAILDON	.	1	3	.	20	6	.	13	34	77	
BARDO	.	1	1	2	2	.	4	1	2	13	
BAREILLY	6	19	12	22	43	36	23	1	9	171	
BARRANQUILLA	1	.	.	.	1	
BELEM	1	.	.	.	1	
BENFICA	.	.	1	1	
BERGEN	1	.	1	
BERN	.	2	2	
BERTA	8	34	19	5	35	4	16	1	21	143	
BINZA	1	1	
BLEDAM	.	.	.	1	1	
BLOCKLEY	1	9	17	4	6	.	4	7	6	54	
BOCHUM	1	1	
BONAIRE	1	1	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
BONARIENSIS	1	.	1	.	1	1	4
BOVISMORBIFICANS	.	4	12	3	10	1	.	2	3	35	
BRAENDERUP	23	77	65	47	106	34	63	26	88	529	
BRANDENBURG	8	23	17	13	36	9	7	4	.	117	
BRAZIL	2	.	.	.	2	
BRAZOS	1	.	1	
BREDENEY	1	12	4	2	7	2	7	2	7	44	
BROUGHTON	1	1	
BUTANTAN	1	1	
BUZU	1	1	
CALABAR	.	.	1	1	
CALIFORNIA	1	.	1	
CAMBRIDGE	1	.	1	
CARACAS	1	1	
CARMEL	1	1	
CARNO	.	1	1	
CARRAU	1	1	1	4	2	.	.	1	2	12	
CERRO	1	2	11	6	7	.	11	8	10	56	
CHAILLEY	.	.	1	2	3	
CHAMELEON	1	1	1	.	.	1	.	1	.	5	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
CHESTER	.	4	9	.	6	.	3	.	7	29	
CHINGOL	2	2	
CHOLERAESUIS	5	2	11	3	2	.	.	.	2	25	
CHOLERAESUIS VAR KUN	.	.	.	2	5	.	1	1	.	9	
CLACKAMAS	3	3	
CLAIBORNEI	1	.	.	1	
COELN	2	2	
COLINDALE	.	.	.	1	1	2	
COLORADO	2	2	
CONCORD	.	1	.	1	1	3	
CORVALLIS	1	1	
COTHAM	2	2	
CUBANA	4	7	1	3	9	.	7	2	9	42	
CURACAO	1	1	
DAHRA	2	.	2	
DAYTONA	3	.	.	1	.	4	
DEGANIA	1	1	
DENVER	1	.	1	
DERBY	5	16	31	7	38	2	7	24	44	174	
DJELFA	.	1	1	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION											TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific			
DJUGU	1	1
DRYPOOL	5	.	.	.	5
DUBLIN	2	8	5	.	5	.	.	15	31	.	.	66
DUESSELDORF	2	2	.	.	1	5
DURBAN	.	3	3
DURHAM	1	1
EALING	.	1	2	.	2	.	.	.	1	.	.	6
EASTBOURNE	.	1	2	.	2	.	1	.	1	.	.	7
EBRIE	.	1	1
EDINBURG	1	4	1	.	.	6
EIMSBUETTTEL	.	.	1	1
ELOMRANE	.	.	1	1	.	.	2
EMEK	1	1	2	2	.	.	.	1	1	.	.	8
ENTEBBE	.	.	1	1
ENTERITIDIS	454	1484	860	231	794	87	110	567	756			5343
EPPENDORF	1	1	2
ERLANGEN	.	.	1	1
ESSEN	.	2	.	.	1	3
FARMSEN	1	1	1	3
FAYED	6	6

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
FISCHERKIEZ	1	1
FLINT	.	1	3	.	59	.	.	1	.	.	64
FLORIDA	.	.	.	1	1
GABON	.	.	.	1	1
GAMINARA	1	3	2	3	8	7	23	4	1	1	52
GATUNI	1	1
GIVE	3	5	14	9	12	8	26	3	17	.	97
GLOSTRUP	1	.	4	.	.	.	1	.	1	.	7
GOETTINGEN	1	.	.	1
GOLDCOAST	.	.	1	1
GROUP 51	1	1
GROUP 53	2	.	2
GROUP 58	1	1
GROUP 60	1	.	.	1	.	2
GROUP 61	1	.	2	1	.	1	5
GROUP 62	1	.	.	.	1
GROUP 63	.	.	1	1
GROUP A	1	.	.	1	.	.	1	.	.	.	3
GROUP B	21	60	44	128	64	31	39	27	24	.	438
GROUP C1	4	20	8	57	10	12	15	7	5	.	138

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION											TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific			
GROUP C2	2	2	.	25	4	2	5	8	.			48
GROUP D1	.	5	4	22	8	6	10	22	3			80
GROUP D2	1	.	.			1
GROUP E1	1	.	1	7	2	2	3	1	.			17
GROUP E2	1	.	.			1
GROUP E4	1	.	.	1			2
GROUP G	2	.	2	1	3	2	5	.	.			15
GROUP H	.	.	1	.	.	1	.	.	.			2
GROUP I	1	.	.	3	.	1	.	.	41			46
GROUP J	.	.	1			1
GROUP K	2	.	2			4
GROUP O	.	1	.	.	3			4
GROUP Q	.	1	.	.	1			2
GROUP R	1			1
GROUP S	1			1
GROUP V	.	3	1	.	1	2	.	.	.			7
GROUP W	1	2			3
GROUP X	1	1	2			4
GROUP Y	3	.	6	.	5	1	.	.	.			15
GROUP Z	1	1	2	2	1	.	.	.	7			14

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
GRUMPENSIS	.	.	1	1	2
GUILDFORD	1	1
HAARDT	.	.	3	3
HADAR	41	134	43	22	104	14	45	39	74	516	
HADEJIA	1	1	
HAGENBECK	.	.	1	1	
HAIFA	.	.	1	.	3	.	.	.	2	6	
HALLE	1	1	
HALMSTAD	2	.	.	2	
HAMBURG	.	.	.	1	1	
HARTFORD	11	15	48	16	33	7	4	2	4	140	
HATFIELD	1	1	
HATO	1	.	.	1	
HAVANA	1	6	2	2	8	1	10	6	10	46	
HEIDELBERG	127	387	310	173	286	86	92	82	273	1816	
HEVES	1	1	
HIDUDDIFY	1	1	
HOLCOMB	.	1	1	
HORSHAM	2	1	3	
HOUTEN	.	.	6	1	.	2	.	.	1	10	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
HVITTINGFOSS	3	2	7	3	6	4	12	.	1		38
I 4,5,12:I:-	.	1	36	2	5	44
IBADAN	1	2	2	.	3	.	18	1	.	.	27
IIIA 48:G,Z51:-	3	3
IIIA 53:Z4,Z23,Z32:-	2	.	.	2	.	.	4
IIIB 48:I:Z	.	.	1	1
IIIB 61:1,V:1,5,7	.	.	1	1	.	2
IIIB 61:K:1,5,7	2	.	1	3
IIIB 65:K:Z	1	.	1
INCHPARK	1	1
INDIANA	.	1	1	1	4	.	.	2	5	14	
INFANTIS	36	67	133	44	80	29	74	42	91	596	
INVERNESS	.	3	1	1	14	2	1	1	1	24	
IPSWICH	.	.	1	1	
IRENEA	.	1	1	
IRUMU	.	2	1	1	2	6	
ISANGI	.	.	1	1	
ISTANBUL	.	10	.	1	1	.	.	.	13	25	
ITAMI	4	.	.	3	7	
ITURI	2	1	.	.	.	3	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
IV 44:Z4,Z23:-	1	3	3	3	1	1	1	1	1	1	14
IV 45:G,Z51:-	.	1	1
JAJA	.	.	1	1
JAMAICA	2	2
JANGWANI	2	1	1	.	1	.	.	.	1	1	6
JAVA	9	36	108	46	52	23	20	13	7	314	
JAVIANA	50	68	65	56	555	93	200	56	54	1197	
JOHANNESBURG	1	5	6	2	18	6	3	1	2	44	
JOS	1	.	.	.	1	
KEDOUGOU	1	1	.	.	.	2	
KENTUCKY	5	25	10	7	5	1	2	5	11	71	
KIAMBU	5	2	9	4	9	2	6	.	3	40	
KILWA	3	3	
KINGABWA	2	2	
KINONDONI	1	1	
KINSHASA	.	.	1	1	1	3	
KINTAMBO	1	1	2	.	1	.	.	2	1	8	
KIRKEE	1	1	
KIVU	.	.	2	2	
KOKOMLEMLE	.	1	1	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
KOTTBUS	.	.	1	.	.	2	.	.	1	1	5
KRALENDYK	.	.	2	1	.	3
KRALINGEN	1	1
KREFELD	1	.	.	.	1
KUA	.	.	1	.	.	1	2
LANDWASSER	.	.	1	.	.	1	2
LANKA	.	.	1	1
LANSING	1	1
LAROCHELLE	2	4
LICHTENBERG	.	1	1
LILLE	1	1
LINDENBURG	.	1	.	.	.	1	.	1	1	.	5
LITCHFIELD	13	27	20	9	19	5	21	5	16	135	
LIVERPOOL	.	.	2	2
LIVINGSTONE	.	1	1	.	2	.	4
LOCKLEAZE	.	1	1
LOHBRUEGGE	.	1	1	.	.	.	2
LOMALINDA	2	2	.	.	1	3	8
LONDON	5	9	6	9	2	2	4	3	3	41	
LUANSHYA	1	1

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
LUCIANA	.	1	.	.	.	1	1	.	.	3	6
MADELIA	1	.	2	2	4	.	2	.	.	1	12
MANCHESTER	1	.	.	1
MANGO	1	1
MANHATTAN	5	7	14	10	10	3	7	.	22	.	78
MANILA	1	.	1
MARINA	13	8	8	2	6	3	1	.	3	.	44
MATADI	1	.	1	.	.	.	2
MBANDAKA	6	13	24	10	21	6	10	26	115	.	231
MELEAGRIDS	3	1	.	2	.	.	1	.	7	.	14
MEMPHIS	.	1	1
MENDEN	.	1	1
MENDOZA	1	1
MGULANI	.	2	2
MIAMI	19	8	5	2	54	1	1	1	4	.	95
MICHIGAN	2	.	2
MIKAWASIMA	1	.	3	.	4
MINNESOTA	.	.	1	1	5	.	5	6	5	.	23
MISSION	1	1
MISSISSIPPI	7	11	5	4	104	51	60	1	5	.	248

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
MONSCHAUI	.	.	3	.	2	5
MONTEVIDEO	56	129	140	72	131	32	86	75	130	851	
MOUNDOU	.	1	1	
MUENCHEN	49	72	195	77	227	47	99	138	428	1332	
MUENSTER	10	14	8	9	9	2	4	3	6	65	
NAPOLI	.	.	.	1	1	2	
NEUBRUNSWICK	.	.	3	.	.	1	2	6	11	23	
NEWINGTON	.	.	3	.	4	.	1	14	1	23	
NEWPORT	101	203	213	156	953	158	492	196	146	2618	
NEWROCHELLE	1	.	.	.	1	
NEWYORK	.	.	1	1	
NGOR	2	.	.	2	
NIMA	1	1	
NITRA	.	.	.	1	1	
NORDENHAM	1	.	.	.	1	
NORWICH	.	5	5	19	13	14	15	2	1	74	
OAKLAND	.	.	1	1	
OHIO	9	10	9	6	12	5	1	8	17	77	
OLDENBURG	.	.	1	1	
ONDERSTEEPOORT	1	1	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
ONTARIO	.	1	1
ORANIENBURG	49	44	95	29	61	18	89	128	103	616	
ORIENTALIS	2	2	
OSLO	3	5	7	4	2	.	.	2	5	28	
OTHMARSCHEN	.	12	.	.	2	1	1	4	.	20	
OVERSCHIE	1	1	2	
PAKISTAN	.	.	1	.	1	.	.	.	4	6	
PANAMA	1	23	17	10	12	4	13	21	31	132	
PARATYPHI A	6	15	4	5	10	1	18	3	15	77	
PARATYPHI B	6	20	36	5	7	3	20	15	60	172	
PARATYPHI C	1	1	
PARERA	1	.	1	2	
PENSACOLA	7	1	.	.	.	8	
PHARR	.	1	1	
PHOENIX	6	.	.	6	
POMONA	3	11	.	.	1	.	.	.	13	28	
POONA	14	35	45	10	27	5	28	50	35	249	
PORTSMOUTH	.	1	1	
POTSDAM	1	1	1	.	2	.	.	.	4	9	
PRAHA	1	1	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
PULLORUM	1	1
PUTTEN	.	.	1	.	2	3
RAMATGAN	1	1
RAUS	.	.	3	3
READING	14	25	8	8	15	2	3	5	17		97
REMO	.	1	1	.	.	.	2
RICHMOND	1	1	2
RIDGE	1	1
RIOGRANDE	1	.	.	.	1
RISSEN	.	2	.	1	1	1	.	.	1	1	6
ROMANBY	.	.	2	.	.	.	1	.	3		6
ROODEPOORT	.	.	.	1	1		2
ROTTNEST	1	1
RUBISLAW	1	1	4	2	51	7	30	.	1		97
RUIRU	1		1
RUZIZI	1	.	.	.	1
SAINTPAUL	24	84	62	28	84	14	27	31	118		472
SALINATIS	1		1
SANDIEGO	4	33	13	6	10	6	9	11	12		104
SANJUAN	.	.	1	1		2

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION											TOTAL			
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific						
SANTIAGO	•	•	•	•	1	•	•	•	•	•	•	•	•	•	1
SAPHRA	•	•	•	•	2	1	7	2	1	•	•	•	•	•	13
SCHLEISSHEIM	•	•	•	•	•	6	•	•	•	•	•	•	•	•	6
SCHWARZENGRUND	12	20	15	7	66	5	4	4	22	•	•	•	•	•	155
SENDAI	•	•	•	•	•	•	•	•	•	•	•	•	•	•	1
SENFENBERG	5	16	18	11	19	4	13	13	21	•	•	•	•	•	120
SHUBRA	•	1	•	•	•	1	2	•	2	•	•	•	•	•	7
SINGAPORE	•	•	•	2	•	•	•	•	2	•	•	•	•	•	4
SINSTORF	•	•	•	2	1	•	•	•	•	•	•	•	•	•	3
SOERENGA	•	•	•	•	1	•	•	•	•	•	•	•	•	•	2
SOMONE	•	•	•	•	1	•	•	•	•	•	•	•	•	•	1
SOUTHAMPTON	•	•	•	•	•	•	•	•	•	•	•	•	•	•	1
STACHUS	•	1	•	•	1	•	•	•	•	•	•	•	•	•	2
STANLEY	11	34	38	14	25	•	3	14	33	•	•	•	•	•	172
STANLEYVILLE	•	8	1	1	•	•	•	•	•	•	•	•	•	•	10
STOCKHOLM	•	•	•	•	4	•	•	•	•	•	•	•	•	•	4
SUBSPECIES I	2	16	2	4	46	•	2	5	4	•	•	•	•	•	81
SUBSPECIES II	•	•	•	2	•	1	2	1	•	•	•	•	•	•	6
SUBSPECIES III	1	1	•	•	•	•	•	•	•	•	•	•	•	•	3
SUBSPECIES IIIA	•	1	•	3	6	1	•	5	•	•	•	•	•	•	16

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
SUBSPECIES IIIA/IIIB	.	.	.	2	10	.	1	1	.	14	
SUBSPECIES IIIB	.	.	.	1	6	1	.	.	1	9	
SUBSPECIES IV	5	.	.	1	16	4	.	.	.	26	
SUELLDORF	.	1	1	
SUNDSVALL	1	.	.	3	4	
TAFO	1	.	.	.	1	
TAKORADI	4	4	
TALLAHASSEE	1	1	.	.	1	1	.	1	.	5	
TAMBACOUNDA	.	1	1	
TAMPICO	2	2	
TELELKEBIR	.	1	3	3	2	1	1	.	4	15	
TENNESSEE	.	4	5	4	4	2	1	4	5	29	
THIES	1	1	
THOMASVILLE	1	.	.	1	2	4	
THOMPSON	50	75	93	53	60	33	55	35	148	602	
TILENE	1	.	.	.	1	
TRAVIS	1	.	1	
TUCSON	.	1	1	
TYPHI	29	90	41	8	48	1	26	8	101	352	
TYPHIMURIUM	566	1196	1186	653	1296	371	495	541	821	7125	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
TYPHIMURIUM VAR COPE	153	116	41	34	230	36	.	144	172	926	
UCCLE	.	.	.	1	1	
UGANDA	1	11	7	2	12	2	4	.	19	58	
UNKNOWN	25	33	71	25	36	24	126	4	55	399	
URBANA	4	7	12	2	11	2	12	.	6	56	
UZARAMO	1	1	
VEJLE	1	.	1	
VIRCHOW	4	10	10	7	10	5	5	.	19	70	
VIRGINIA	1	3	.	.	.	4	.	1	1	10	
WA	1	.	.	1	
WANDSWORTH	.	4	1	.	3	.	.	.	1	9	
WASHINGTON	.	1	1	
WASSENAAR	.	2	6	.	1	.	1	.	1	11	
WAYCROSS	.	1	1	.	2	
WELIKADE	1	1	
WELTEVREDEN	1	5	4	1	6	.	1	3	33	54	
WESLACO	.	1	1	
WESTERSTEDE	.	1	1	
WESTHAMPTON	1	.	1	2	
WIDEMARSH	1	1	

(Continued)

TABLE 5
SALMONELLA ISOLATIONS FROM HUMAN SOURCES
BY SEROTYPE AND GEOGRAPHIC REGIONS, 1999

Serotype	REGION										TOTAL
	New England	Mid Atlantic	East North Central	West North Central	South Atlantic	East South Central	West South Central	Mountain	Pacific		
WIEN	1	.	.	1
WIL	.	1	1
WINNEBA	.	1	1
WORTHINGTON	.	5	3	2	5	.	4	4	5	.	28
YOVOKOME	.	.	.	1	1
ZANZIBAR	1	.	1
TOTAL	2250	5280	4690	2410	6489	1481	2807	2615	4760		32782