


Biology and Management of Horseweed

- Mark Loux, *Ohio State University*
- Jeff Stachler, *Ohio State University*
- Bill Johnson, *Purdue University*
- Glenn Nice, *Purdue University*
- Vince Davis, *Purdue University*
- Dawn Nordby, *University of Illinois*

Purdue Extension

Knowledge to Go

1-888-EXT-INFO





**INDIANA
SOYBEAN
BOARD**



**Illinois Soybean Association
Illinois Soybean Checkoff Board**

This publication is sponsored by the Indiana Soybean Board, Illinois Soybean Association/Illinois Soybean Checkoff Board, and U.S. Department of Agriculture Critical and Emerging Pest Competitive Grants Program, and is a collaboration of Ohio State University Extension, Purdue University Extension, and University of Illinois Extension.

Biology and Management of Horseweed

Heavy reliance on glyphosate for weed control in soybeans has resulted in a serious concern for the long-term viability of this valuable weed management tool. Weed resistance to glyphosate is not a new phenomenon. Glyphosate-resistant field bindweed was reported in Indiana during the mid-1980s in areas that had received repeated glyphosate applications (Degennaro and Weller, 1984). But, weed resistance to glyphosate did not become a major issue in U.S. agronomic crops until several years after the 1996 release of Roundup Ready soybeans. Of greatest concern recently, has been horseweed. Since the initial report of glyphosate-resistant horseweed in 2000 (VanGessel, 2001), this weed management issue has been reported in over nine states. Common to all known cases of glyphosate-resistant horseweed is the frequent use of glyphosate for control of all weeds, little or no use of alternative herbicides that control horseweed, and long-term no-tillage crop production practices.



Figure 1. This photo shows the flower structure of a mature horseweed plant.

The purpose of this publication is two-fold. First, we will discuss some of horseweed's biological characteristics that make it particularly troublesome to control in agronomic crops. Second, we will provide management strategies using technologies now available that will hopefully slow the spread of glyphosate-resistant horseweed.

Identification

Horseweed, also known as marestalk, is native to North America. This annual weed can follow a winter or summer annual life cycle. After emergence in the fall, horseweed forms a basal rosette for winter survival. In a winter annual life cycle, the rosette bolts in the spring, growing to a height of 1.5 to 6 feet. Horseweed leaves are alternate, linear, and simple with entirely or slightly toothed margins. Mature plants have leaves with no petioles (sessile). Leaves get progressively smaller in size toward the top of the plant. Stems are erect and tend to be unbranched unless damaged by herbicides, mowing, or animal or insect feeding. Flowers are arranged in a panicle with numerous white ray flowers (1/16 to 2/16 inches long) and 20 to 40 yellow disk flowers (see Figure 1). The seeds are small achenes

Biology and Management of Horseweed

(1/16 to 1/4 inches long), with a pappus of tan to white bristles. A pappus is a structure that allows the seed to be dispersed by the wind, similar to that of common dandelion.

Horseweed (see Figure 2) is often misidentified as whitlowgrass, mouseear chickweed, corn or Persian speedwell, shepherd's-purse, or several of the fleabane species, especially annual fleabane (see Figure 3). The two most common misidentifications are whitlowgrass and fleabane species. Whitlowgrass has shorter and narrower leaves and an entirely to slightly smooth margin compared to horseweed. Around the leaf margin, annual fleabane



Figure 2. This photo shows a horseweed seedling in rosette stage, just beginning to bolt.



Figure 3. The photos above show weeds commonly confused with horseweed: whitlowgrass (top left), mouseear chickweed (middle left), annual fleabane (bottom left) corn or Persian speedwell (top right), and shepherd's-purse (bottom right).

tends to have purple spots at the tip of each tooth while horseweed does not have these spots. Also, the distance between nodes on all fleabane species after bolting are greater than the internode distance on horseweed. Fleabanes will produce more branches than horseweed. Mouseear chickweed, and corn and Persian speedwell can easily be separated from horseweed at a young stage by the presence of opposite leaves at all early stage nodes. Horseweed also will always have alternating leaves. The first node of shepherd's-purse tends to have opposite leaves while subsequent nodes have alternate leaves, although all nodes may be alternating. The hairs on shepherd's-purse leaves are shorter and tend to stay on the upper leaf surface compared to horseweed leaf hairs that are larger and tend to be present on leaf margins. In later development stages, shepherd's-purse leaves will be deeply lobed compared to the toothed or slightly lobed leaves of horseweed.

Distribution and Emergence

Horseweed seed germinates readily as soon as it falls off a mature plant. Horseweed plants usually germinate in the fall or spring, but they also can germinate in midsummer if growing conditions are adequate. In northern regions of Indiana, Illinois, and Ohio most horseweed germinates in the fall, overwinters as a rosette, and begins to bolt in the spring. Typically, 14 to 84 percent of fall emerging plants survive until spring. This broad survival range can be attributed to weather conditions and size of the horseweed rosette. The larger the rosette is prior to winter, the greater the chance of survival in the spring. Spring germination is more predominant in southern regions of Indiana, Illinois, and Ohio. In southeast Indiana and southwest Ohio, spring germinating horseweed is one of the most problematic summer weeds. Spring horseweed generally remains a rosette for a relatively short period prior to bolting. Horseweeds germinating in July and August tend to remain as rosettes until the following spring, with only a few plants bolting and producing flowers in the fall.

Growth and Development

Following rosette formation, horseweed plants begin to bolt in mid-April and start flowering at the end of July. Horseweed is self-compatible and releases pollen before the capitula are fully opened, supporting the idea that horseweed is primarily self-pollinating; however, outcrossing within a population has been cited to range from 1.2 to 14.5 percent. Horseweed's wind dissemination and relatively high seed production suggests that the dispersal of resistant horseweed plants across an agricultural landscape could be very rapid. Regehr and Bazzaz (1979) reported that horseweed seed input into a corn field ranged from 12,500 seeds per square yard at 20 feet from the seed source, and more than 125 seeds per square yard at 400 feet from the seed source. Taller plants produce more seed than shorter plants, suggesting that seed height on plants might influence the distance wind can transport seed.




Figure 4. This graph shows the relative life cycles of horseweed throughout the year for plants that emerge in the spring and fall.

Interference and Competition

In southeast Indiana and southwest Ohio, horseweed quite often behaves like a summer annual. In these situations, horseweed begins to emerge from late March into late June. Southeast Indiana soils characteristically have a shallow layer of silt over a thick clay layer. These soils have relatively poor internal drainage and water holding capacity and require frequent light rains to maintain optimal crop growth. Horseweed tolerates drought conditions well and continues to grow and produce biomass and seed under conditions stressful for crop growth. Horseweed can produce up to 200,000 seeds per plant and approximately 80 percent of the seed will germinate right off the plant. Because of the unique soils in this region and widespread adoption of no-till practices, horseweed has emerged as the number one weed problem in this area.

Bruce and Kells (1990) showed in management studies conducted in Michigan that soybean yields could be reduced up to 83 percent by horseweed in untreated check treatments. In general, we feel horseweed is much less competitive than most of the other summer annual weeds. However, there have been no traditional competition studies published to date that have evaluated the effect of density or emergence time on its yield loss potential in soybean or corn. We do know that horseweed can affect crop production in other ways besides direct competition for light, water and nutrients. Horseweed can be a host for the tarnished plant bug, an alfalfa pest. It can also be a host for aster yellows, a viral disease that can be transmitted by aster leaf hoppers to a wide variety of plants.

Increased Prevalence

Three factors are commonly mentioned as the causes of the increased horseweed prevalence: crop rotation, tillage, and herbicide resistance.

Crop Rotation

Crop rotation appears to have a minimal impact on horseweed prevalence. In field surveys conducted in areas where horseweed was most problematic in Indiana, horseweed was found in 63 percent of double crop soybean fields compared to 51 percent in continuous soybean fields and 47 percent in fields in a corn-soybean rotation.

Tillage

In the same survey mentioned above, increases in tillage intensity reduced horseweed prevalence by 50 percent or more. Horseweed was found in 61 percent of the no-till fields compared to 24 percent in reduced tillage and 8 percent in conventional tillage fields.

Herbicide Resistance

The first case of horseweed herbicide resistance was reported by Japanese researchers in 1980, who found biotypes resistant to Gramoxone[®] (paraquat). More Gramoxone[®]-resistant biotypes have since been found in Mississippi. The second case of resistance was reported in 1989 by researchers in Belgium, who found biotypes resistant to atrazine in corn, nurseries, and roadsides. Additional atrazine-resistant biotypes have since been found in Michigan. ALS-resistant horseweed biotypes were first found

Biology and Management of Horseweed

in the United States in Indiana and Ohio during the 1999 growing season, and today are believed to be present in all Ohio counties west of Interstate 71 and U.S. Route 23.

Biotypes resistant to glyphosate (which is sold under the trade names Roundup[®], Touchdown[®], and others) were first reported in Delaware in 2000. Since 2000, glyphosate-resistant biotypes have been found in nine additional states including Ohio, Indiana, and Kentucky (but not in Illinois). In Indiana, glyphosate-resistant biotypes have been found in 19 counties, mostly in the southeast. However, horseweed has been found in Indiana in Wells County in the northeast and Montgomery County in the west-central area. In Ohio, glyphosate-resistant biotypes have been found in 18 southwest counties, stretching to Union and Franklin counties.

Glyphosate-resistant horseweed populations in Ohio and Indiana were first reported in areas where the following production practices are common: soybeans grown in the same field for consecutive years (up to 14 years in some fields), use of only glyphosate for weed control, and little or no tillage.

In horseweed, resistance to more than one herbicide has been reported in Israel with a biotype resistant to atrazine and Glean[®] (but not glyphosate); in Michigan with a biotype resistant to atrazine, simazine, and diuron; and in Ohio with biotypes resistant to glyphosate and FirstRate[®]. The glyphosate and FirstRate[®] resistant biotype in Ohio has been confirmed in at least two fields, one each in Montgomery and Miami counties.

In Indiana, horseweed populations resistant to glyphosate and FirstRate[®] have been found in Bartholomew, Jefferson, Scott, and Washington counties; populations resistant to glyphosate and Classic[®] have been found in Bartholomew, Scott, and Wells counties; and populations resistant to glyphosate, FirstRate[®], and Classic[®] have been found in Bartholomew and Scott counties.

Control

The primary goal of a horseweed management program in no-till soybeans should be effective control of emerged horseweed plants prior to planting. Soybeans planted before early to mid-May also will require a residual




Figure 5. Glyphosate-resistant horseweed populations have been confirmed in 37 Indiana and Ohio counties (highlighted in red). To date, no resistant populations have been found in Illinois.




Figure 6. Ohio horseweed populations showing resistance to ALS and glyphosate herbicides (top row), resistance to glyphosate herbicides only (second row from top), resistance to ALS herbicides only (third row from top), and no herbicide resistance (bottom row).

herbicide to control later emerging plants. This strategy will reduce the need for postemergence herbicide treatments, which can be limited in effectiveness and exert further selection for herbicide resistance in the population. The following principles are important in horseweed control programs:

- 2,4-D ester should be included in herbicide treatments when possible;
- herbicides should be applied before horseweed plants are 4 to 6 inches tall;
- herbicides applied in the fall will control emerged horseweed, but may not adequately control spring-emerging plants; and
- spring applications prior to early May should include a residual herbicide to control later-emerging plants.

Control of emerged plants before planting

The most effective treatments for controlling horseweed plants up to about 6 inches tall are ranked roughly in order of effectiveness below. Treatments containing fewer than three herbicides may be less effective on an ALS- or glyphosate-resistant population.

- A combination of glyphosate and 2,4-D ester; plus one of the following: Canopy[®]EX or FirstRate[®]/Amplify[®] or Gangster[®].
- A combination of glyphosate and 2,4-D ester.
- A combination of Sencor[®], Gramxone[®], and 2,4-D ester.
- A combination of glyphosate plus one of the following: FirstRate[®]/Amplify[®] or Gangster[®], or Canopy[®]EX.

Several other treatments can be effective when plants are less than two inches tall, including: Sencor[®] plus 2,4-D ester; Sencor[®] plus Gramoxone[®]; and 2,4-D ester alone. A combination of 2,4-D ester plus Canopy[®]EX or FirstRate[®]/Amplify[®] or Gangster[®] can also be used on small plants, but effectiveness will be reduced in ALS-resistant populations.

Residual horseweed control (spring application)

In fields where horseweed may be ALS-resistant, the most effective herbicides include Gangster[®], metribuzin (at least 0.4 lb ai/A), sulfentrazone, and Valor[®]. Canopy[®]EX, FirstRate[®]/Amplify[®] or Gangster[®], or Python[®] can be used for residual control in fields where the horseweed is not ALS-resistant.

Spring herbicide recommendations based on horseweed size

Fields treated with herbicide the previous fall

Fields should be free of overwintering horseweed in the spring, as long as 2,4-D ester was a component of the fall treatment, but additional horseweed emergence is likely. It is possible that residual herbicides applied in the fall, such as Canopy[®]EX, Valor[®], and Sencor[®], can control horseweed through early June. This is most likely to occur in sparser populations that are not ALS-resistant. Regardless of the herbicide(s) applied in fall, fields should be scouted before planting. Apply herbicide as needed prior to soybean planting to control emerged horseweed, and include residual herbicides if the field is planted before mid-May.



Figure 7. Horseweed plants protruding through soybean canopy in late summer.

Horseweed in the seedling or rosette stage (April)

Controlling horseweed in the seedling or rosette stage can be extremely effective, since small plants are easily controlled and residual herbicides applied at these stages can provide control through early June. Emerged plants should be adequately controlled by 2,4-D ester (1 lb ai/A). When the 2,4-D rate is limited to 0.5 lb ai/A, combine with glyphosate, Sencor[®], or Sencor[®] plus Gramoxone[®]. Sencor plus Gramoxone[®] (without 2,4-D) can effectively control seedlings or small rosettes.

Horseweed stem elongated but not more than 4 to 6 inches tall (May)

During this stage, the most effective treatment is glyphosate, plus 2,4-D ester, plus either Canopy[®]EX or FirstRate[®]/Amplify[®]. Glyphosate plus 2,4-D ester is effective where glyphosate resistance is not an issue; and glyphosate, plus either Canopy[®]EX or FirstRate[®]/Amplify[®] is effective where ALS resistance and ALS plus glyphosate resistance is not an issue. Use a glyphosate rate of at least 0.75 lb acid equivalent per acre, unless ALS and glyphosate resistance is present, then use a minimum rate of 1.5 lbs acid equivalent/A. Somewhat more variable but generally effective on this size plant is the combination of Sencor[®] plus Gramoxone[®] (minimum of 1.7 pt/A) plus 2,4-D ester.

Horseweed more than 6 inches tall (mid to late May)

Horseweed of this size will likely be difficult to control. Anything less than a three-way mixture of glyphosate, plus 2,4-D ester, plus Canopy[®]EX or FirstRate[®]/Amplify[®] or Gangster[®] are not recommended. Use a glyphosate rate of at least 1.5 lbs acid equivalent/A. Resistance to glyphosate and/or ALS inhibitors can result in situations where effective control is not possible.

Conclusion

Weed resistance is not new to agriculture; we have been dealing with it for some time. Awareness and prevention are the first steps to dealing with this problem. As with other weed resistance issues, dealing with resistant horseweed requires adjusting management strategies. Having knowledge of herbicide groups and using combinations of herbicides with different modes of action will help prevent additional resistance problems from arising.

References

- Barnes, J., Johnson, B., Gibson, K., and Weller, S. 2004. Crop rotation and tillage system influence late-season incidence of giant ragweed and horseweed in Indiana soybean. Online. Crop Management doi:10.1094/CM-2004-0923-02-BR.
- Bruce, J.A. and J.J. Kells. 1990. Horseweed (*Conyza canadensis*) control in no-tillage soybeans (*Glycine max*) with preplant and preemergence herbicides. *Weed Technology*. 4:642-647.
- DeGennaro, F.P. and S.C. Weller. 1984. Differential susceptibility of field bindweed (*Convolvulus arvensis*) biotypes to glyphosate [Indiana]. *Weed Science*. 32:472-476.
- Johnson, B., Barnes, J., Gibson, K., and Weller, S. 2004. Late season weed escapes in Indiana soybean fields. Online. Crop Management doi:10.1094/CM-2004-0923-01-BR.
- Regehr, D.L. and F.A. Bazzaz. 1979. The population dynamics of *Erigeron canadensis*, a successional winter annual. *Journal of Ecology*. 67: 923-933.
- VanGessel, M.J. 2001. Glyphosate-resistant horseweed from Delaware. *Weed Science*. 49:703-705.

Photo Credits

All photos and illustrations by Mark Loux and Jeff Stachler, Ohio State University; Bill Johnson, Glenn Nice, and Vince Davis, Purdue University; and Dawn Nordby, University of Illinois.

11/04

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.



ID-323

Purdue Extension

Knowledge to Go

1-888-EXT-INFO