

Annual Summary

2004

Shigella



Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Infectious Diseases
Division of Bacterial and Mycotic Diseases
Foodborne and Diarrheal Diseases Branch
Atlanta, Georgia 30333







Eric Mintz, M.D., M.P.H.
Chief, Diarrheal Diseases Epidemiology Section
Foodborne and Diarrheal Diseases Branch



Nancy Strockbine, Ph.D.
Chief, National Reference Lab for *E. coli* and *Shigella*
Foodborne Diseases Laboratory Section
Foodborne and Diarrheal Diseases Branch



Brian D. Plikaytis, Ph.D.
Chief, Biostatistics Office



Robert V. Tauxe, M.D., MPH
Chief, Foodborne and Diarrheal Diseases Branch

Division of Bacterial and Mycotic Diseases

National Center for Infectious Diseases

Centers for Disease Control and Prevention

Recommended Reference Citation:

Centers for Disease Control and Prevention. *Shigella Surveillance: Annual Summary, 2004*. Atlanta, Georgia: US Department of Health and Human Services, November 2005

Single copies of *Shigella Surveillance: Annual Summary 2004* are available from:

Centers for Disease Control and Prevention
Foodborne and Diarrheal Diseases Branch
Mail Stop: A38
1600 Clifton Road
Atlanta, Georgia 30333
Telephone: 404-639-2206
<http://www.cdc.gov/ncidod/dbmd/foodborne/index.htm>

The Adobe Acrobat (PDF) version of this document can be viewed on the world-wide web at <http://www.cdc.gov/ncidod/dbmd/phlisdata/shigella.htm>. Further information concerning data described in this report can be obtained by contacting the Foodborne and Diarrheal Diseases Branch at telephone number (404) 639-2206. For further information concerning PHLIS please contact the Biostatistics Office at telephone number (404) 639-1364.

All material in this report is in the public domain and may be used and reprinted without permission; citation of source is appreciated.

TABLE OF CONTENTS

Introduction	i
Annual Highlights for 2004	ii
Acknowledgements	iii
References	iv
TABLE 1	1
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species in 2004	
TABLE 2	2
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species and Serotype in 2004	
TABLE 3	3
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species, Age Group and Sex, 2004	
TABLE 4 / FIGURE 1	5
Median Age of persons from whom laboratory confirmed <i>Shigella</i> isolates were reported to the CDC by Species and Year for 1990-2004	
TABLE 5 / FIGURE 2	6
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species and Year for 1990-2004	
TABLE 6	7
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species, Serotype and Year for 1990-2004	
TABLE 7	9
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species, Serotype and Month for 2004	
TABLE 8	10
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species, Serotype and Month for 1990-2004	
TABLE 9	12
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species, Geographic Region and Year for 1990-2004	
TABLE 10	14
Laboratory confirmed <i>Shigella</i> isolates reported to the CDC by Species, State and Year for 1990-2004	
FIGURE 3	21
Laboratory confirmed <i>S. sonnei</i> isolates reported to the CDC by Geographical Region and Year for 1990-2004	

Laboratory-Confirmed *Shigella* Surveillance Annual Summary, 2004

The Annual Summary contains surveillance data on reported laboratory-confirmed *Shigella* isolates in the United States. The National *Shigella* Surveillance System collects reports of isolates of *Shigella* from every state in the United States. This information is reported electronically through the Public Health Laboratory Information System (PHLIS) by the State Public Health Laboratory Directors and State and Territorial Epidemiologists to the Foodborne and Diarrheal Diseases Branch (FDDDB) and the Biostatistics Office (BSO) of the Division of Bacterial and Mycotic Diseases in the National Center for Infectious Diseases.

The National *Shigella* Surveillance System is based on data collected by state and territorial public health laboratories. *Shigella* isolates are submitted to the state public health laboratory by clinical diagnostic laboratories. The state and territorial laboratories confirm the isolates as *Shigella*, perform subtyping, and submit the data for reporting through PHLIS. Unusual or untypable isolates are forwarded to the National *Shigella* Reference Laboratory at the Centers for Disease Control and Prevention for further characterization or confirmation. These results are reported back to the state laboratory, where they are reported to CDC through PHLIS.

The capture of isolates in the National *Shigella* Surveillance System is considered to be consistent. However, some *Shigella* isolates may not be forwarded or reported to state public health laboratories and therefore are not captured. In addition, irrespective of the surveillance system, many cases of *Shigella* illness are not reported because the ill person does not seek medical care, the health-care provider does not obtain a specimen for diagnosis or the laboratory does not perform culture for *Shigella*. The results of surveillance reported herein are therefore substantial underestimates of the true number of infections.

The number of isolates reported by state represents the state where laboratory confirmation and subtyping were performed. In some instances, the reporting state is not the same as the state of residence of the person from whom the isolate was obtained. For the Annual Summaries, duplicate records were deleted. All isolates reported herein were from infected humans.

There are 4 major subgroups of *Shigella*, designated A, B, C and D, and 44 recognized serotypes (Table A). Subgroups A, B, C and D have historically been treated as species: subgroup A for *Shigella dysenteriae*; subgroup B for *Shigella flexneri*; subgroup C for *Shigella boydii* and subgroup D for *Shigella sonnei*. These subgroups and serotypes are differentiated from one another by their biochemical traits (ability to ferment D-mannitol) and antigenic properties. The most recently recognized serotype belongs to subgroup C (*S. boydii*) (1).

Table A. Classification of *Shigella* Subgroups

Subgroup	Species	Number of serotypes	Fermentation of D-mannitol	Subgroup B group antigens
A	<i>S. dysenteriae</i>	15	-	-
B	<i>S. flexneri</i>	8 ^a	+	+
C	<i>S. boydii</i>	20	+	-
D	<i>S. sonnei</i>	1	+	-

^a = Serotypes 1-5 are subdivided into 11 subserotypes.

The Statistical Outbreak Detection Algorithm (SODA), developed by BSO and FDDB, is a statistical algorithm performed on the National Surveillance Data to detect unusual clusters of *Shigella* infection. SODA compares current *Shigella* isolates reported through PHLIS by subgroup or serotype with a 5 year historical baseline for that subgroup or serotype for the specified time period to detect unusual increases from the baseline. Analyses can be conducted at state, regional, or national levels. Since 1996, SODA has been implemented at CDC and selected state health departments. If you would like more information on SODA, please call the PHLIS Helpdesk (404) 639-3365.

Annual Highlights for 2004

A total of 9,343 *Shigella* isolates were reported from public health laboratories in 50 states in 2003 (Table 1). This represents a 50% decrease compared with 1994 and a 41% decrease from 2003. This is the lowest number of *Shigella* isolates reported since 1969. The national rate of reported *Shigella* isolates in 2004 was 3.2 per 100,000 population based on 2004 census population estimate figures for the United States.

Similar to previous years, *Shigella* was isolated frequently from children under 5 years of age, who accounted for 30.2% of all isolates. About 31.6% of all isolates came from persons aged 5-19 years, and 28.8% from persons aged 20-59, with declining numbers thereafter. The median age of patients by species is shown in Table 4. The overall distribution of *Shigella* isolates between the sexes was similar, with females accounting for 48.4% of persons from whom *Shigella* was isolated. Gender differences were most notable for a preponderance of females in four age groups, 10-19 (51.8%), 20-29 (64.4%), 30-39 (53.4%), and 70-79 (53.8%) and for a relative paucity of females in two age groups 40-49 (44.2%) and 60-69 (43.6%). These gender differences reflect similar findings among reported isolates of *Shigella sonnei*. Among reported isolates of *Shigella flexneri*, a male predominance is seen, particularly in the age groups 30-39 (64.6%) and 40-49 (65.6%). These estimates, however, are not complete since the District of Columbia did not report the age of persons from whom *Shigella* isolates were obtained. In addition, gender information was not reported for 7.1% of all isolates and age information was not reported for 3.1% of isolates.

The frequency of reported species, and the frequency of reported serotypes within these groups for all *Shigella* isolates are shown in Tables 1 and 2. Of the 9,343 isolates, 8,242 (88.2%) were subgrouped. Trends of subgroups remained constant, with subgroup D (*S. sonnei*) accounting for the largest percentage of isolates (68.9%), followed by subgroup B (*S. flexneri*, 17.2%), subgroup C (*S. boydii*, 1.8%) and subgroup A (*S. dysenteriae*, 0.4%). *Shigella* isolate serotype trends by year are shown in Table 5 and in Figure 2. Over the past decade, the numbers of reported *Shigella* isolates in subgroups A, B and C, and the proportions of all reported *Shigella* isolates due to these three subgroups have declined. Slight increases in the number of *S. boydii* isolates was observed in both 2003 and 2004. The number (1101) and the proportion (11.8%) of all reported *Shigella* isolates that were not identified as belonging to a specific subgroup also increased. The highest numbers and proportions of all reported *Shigella* isolates that were not identified as belonging to a specific subgroup were reported by North Dakota (10,100%), Nebraska (32,68%) and Nevada (24,42%).

Shigella transmission occurs via the fecal-oral route. The majority of subgroup D (*S. sonnei*) infections in the United States occur in young children and in association with crowding and poor personal hygiene. Daycare centers have been implicated in many large *S. sonnei* outbreaks that can last many months and affect many persons (2, 4). *S. sonnei* has also been transmitted through unchlorinated wading pools (5), interactive water fountains (6), food items such as parsley (7) and bean dip (8), and men who have sex

with men (MSM) (9). Until recently, the dominant subgroup causing illness among MSM was subgroup B (*S. flexneri*) (10, 11). However, in a large outbreaks among MSM in San Francisco, the dominant serotype was subgroup D (*S. sonnei*) (9). Recent trends in shigellosis in the United States are reviewed in a publication by Dr. Amita Gupta and co-authors (12).

Geographic trends by region for subgroup D (*S. sonnei*) isolates from 1990 to 2004 are illustrated in Figure 3. All nine regions registered decreases in subgroup D (*S. sonnei*) isolates from 2003 to 2004.

Acknowledgements

Thanks to Richard Bishop and Elizabeth Blanton for assembling this summary and to all the State Public Health laboratories and epidemiologists who participate in this surveillance

References

1. Kalluri P, Cummings K, Abbott S, et al. (Mintz ED). Epidemiological features of a newly described serotype of *Shigella boydii*. *Epidemiology and Infection*. 2004; 132:579-583.
2. Mohle-Boetani JC, Stapleton M, Finger R, Bean N, Poundstone J, Blake P, Griffin PM. Communitywide Shigellosis: Control of an outbreak and risk factors in child day-care centers. *Am J Public Health* 1995;85:812-816.
3. Shane AL, Tucker NA, Crump JA, Mintz ED, Painter JA. Sharing Shigella: Risk Factors for a Multicommunity outbreak of Shigellosis. *Arch Pediatr Adolesc Med* Vol 157: 601-603 June 2003.
4. CDC. Multistate Outbreak of Rhamnose-negative *Shigella sonnei* – Eastern United States, March 2003. *MMWR*; 2004 53:60-63.
5. CDC. Shigellosis outbreak associated with an unchlorinated fill-and-drain wading pool -- Iowa, 2001. *Morbidity and Mortality Weekly Report* 2001;50:797-800.
6. CDC. Outbreak of gastroenteritis associated with an interactive water fountain at a beachside park -- Florida, 1999. *MMWR* 2000;49:565-8.
7. CDC. Outbreaks of *Shigella sonnei* infection associated with eating fresh parsley--United States and Canada, July-August 1998. *MMWR* 1999;48:285-9.
8. Kimura AC, Johnson K, Palumbo MS et al. Multi-state outbreak of drug-resistant *Shigella sonnei* associated with consuming a commercially prepared five-layered dip. *Emerg Infect Dis* 2004;10:1147-9).
9. CDC. *Shigella sonnei* outbreak among men who have sex with men--San Francisco, California, 2000-2001. *MMWR* 2001;50; 922-6.
10. Tauxe RV, McDonald RC, Hargrett-Bean N, Blake PA. The persistence of *Shigella flexneri* in the United States: increasing role of adult males. *Am J Public Health* 1998;78:1432-5.
11. CDC. *Shigella flexneri* Serotype 3 infections among men who have sex with men –Chicago, Illinois, 2003-2004. *MMWR* 2005; 54; 820-2.
12. Gupta A, Polyak CS, Bishop RD, Sobel J, Mintz ED. Laboratory-confirmed shigellosis in the United States, 1989-2002: epidemiologic trends and patterns. *Clinical Infectious Diseases*. 2004;38:1372-1377.