


STATE INJURY INDICATORS REPORT

Third Edition – 2004 Data



State Injury Indicators Report

Third Edition — 2004 Data

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Injury Prevention and Control

July 2007

Editors

Renee L. Johnson, RPT, MSPH
Division of Injury Response
National Center for Injury Prevention and Control

Robert G. Thomas, Jr., MSIM
Office of Statistics and Programming
National Center for Injury Prevention and Control

Karen E. Thomas, MPH
Division of Injury Response
National Center for Injury Prevention and Control

Nimeshkumar Patel, MS
Office of Statistics and Programming
National Center for Injury Prevention and Control

Kelly Sarmiento, MPH
Division of Injury Response
National Center for Injury Prevention and Control

Acknowledgements

The editors thank the State and Territorial Injury Prevention Directors Association, the Council of State and Territorial Epidemiologists, and their respective members. These partners have facilitated the ongoing advancement and success of the development of the injury indicators. The editors also thank Kevin Webb from the Office of Statistics and Programming and Angela Marr from the Division of Injury Response, National Center for Injury Prevention and Control, for their consultation and guidance.

The *State Injury Indicators Report, Third Edition—2004 Data* is a publication of the National Center for Injury Prevention and Control, Centers for Disease Control and Prevention.

Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH
Director

National Center for Injury Prevention and Control

Ileana Arias, PhD
Director

Suggested Citation: Johnson RL, Thomas RG, Thomas KE, Patel N, Sarmiento K. *State Injury Indicators Report, Third Edition — 2004 Data*. Atlanta (GA): Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2007.

Contributors

Arizona Department of Health Services

Jennifer Jung, MSPH
Tomi St. Mars, RN, BSN, CEN

Arkansas Department of Health and Human Services

J. Edward Carson, MHA
Paul W C Johnson, MSPH
Elaine Anthony Jones
Gayla Nooner, MA
Thomas S. Rainer, BBA

California Department of Health Services

Jennifer Harper, MSPH

Colorado Department of Public Health and Environment

Kirk A. Bol, MSPH
Holly B. Hedegaard, MD, MSPH

Connecticut Department of Public Health

Susan J. Hewes, MPH
Jennifer Morin, MPH
Justin Peng, MPH

Florida Department of Health

Michael Lo, MSPH

Georgia Department of Human Resources

Laura Fehrs, MD
Bill Van Fleit, MA
Manxia Wu, MD, MPH
Denise Yeager

Hawaii State Department of Health

Daniel Galanis, PhD

Kansas Department of Health and Environment

Greg Crawford
Lori Haskett, BA
Donald Owen, MA
Ghazala Perveen, MBBS, PhD, MPH
Ismaila Ramon, MPH

Kentucky Cabinet for Health and Family Services

Mike D. Singleton, MS
Lei Yu, MS

Louisiana Department of Health and Hospitals

Mona Doshani, MD, MPH
Shirley Kirkconnell, MSW, MPH

Maine Department of Health and Human Services

Ann Farmer, MS
Cindy Mervis, MPH

Maryland Department of Health and Mental Hygiene

Brenna C. Hogan, MPH
Becky L. Roosevelt, MA
Tracey Serpi, PhD
Chris Tkach, PhD

Massachusetts Department of Public Health

Beth Hume, MPH
Maria McKenna, MPH
Loreta McKeown, MPH

Michigan Department of Community Health

Thomas W. Largo, MPH

Minnesota Department of Health

D. R. Hagel
Mark Kinde, MPH
Jon Roesler, MS

Nebraska Health and Human Services System

Xiaojuan Mi, MS
Peg Prusa-Ogea, MA
Ming Qu, PhD
Lei Zhang, MS

Nevada Department of Health and Human Services

Kelly Langdon, MPH
William Lee, BS
Andrea R. Rivers, BA
Wei Yang, PhD

New Hampshire Department of Health and Human Services

David L. Reichel, MPH, DC

New Mexico Department of Health

Barbara Chatterjee, MS
Glenda Hubbard, MPH
Ajoy Kumar, MBBS, MPH
Leona Woelk, MA

New York State Department of Health

Michael Bauer, MS
Susan Hardman
Erin Shortt, MPH

North Carolina Department of Health and Human Services

Pedro LunaOreo, PhD
Catherine (Kay) Sanford, MSPH
Jean Slosek, MS

Ohio Department of Health

David Engler, PhD (Ohio Hospital Association)
Edward Socie, MS

Oklahoma State Department of Health

Tracy Wendling, MPH

Oregon Department of Human Services

Janice D. Alexander, PhD
Mel Kohn, MD, MPH
Lisa Millet, MSH

Pennsylvania Department of Health

Nathan James, BA
Robert Keyton
Gerald Miller, BS
Tom Truong, MS

Rhode Island Department of Health

Edward Donnelly, RN, MPH
Janice Fontes
Beatriz Perez, MPH

South Carolina Department of Health and Environmental Control

Georgette Demian, MPH
Elizabeth Hall, BS
Byron Kirby, BS
Randy Rambo

Tennessee Department of Health

Tom Spillman, BA

Utah Department of Health

Catherine Groseclose, MS
Gary Mower, MPH
Albert Wang, MPH

Vermont Department of Health

Caroline W. Dawson, BS, MS, MPA
Edith Munene, BA, MA, MA
Patricia Worcester, PhD

Virginia Department of Health

Michelle White, MSW

Washington State Department of Health

Jennifer Sabel, PhD
John Sabel, BS

Wisconsin Department of Health and Family Services

Randall L. Glysch, MS

Table of Contents

Foreword	vii	4. Fire-Related Indicators	31
Abbreviations	viii	4a. Unintentional Fire-Related Fatalities (Overall), 2004	32
Introduction	1	4b. Unintentional Fire-Related Fatalities by Sex, 2004	33
Discussion of Indicators	2	4c. Unintentional Fire-Related Fatalities by Age, 2004	34
Methods	2	4d. Unintentional Fire-Related Hospitalizations (Overall), 2004	35
Hospital Discharge Data Quality.	3	4e. Unintentional Fire-Related Hospitalizations by Sex, 2004	36
Highlights	3	4f. Unintentional Fire-Related Hospitalizations by Age, 2004	37
Table A. Factors Affecting Representativeness of 2004 Hospital Discharge Data	4	5. Firearm-Related Indicators	39
Limitations and Future Efforts	8	5a. Firearm-Related Fatalities (Overall), 2004	40
References	9	5b. Firearm-Related Fatalities by Sex, 2004	41
1. All-Injury Indicators	11	5c. Firearm-Related Fatalities by Age, 2004	42
1a. Hospitalizations for All Injuries (Overall), 2004	12	5d. Firearm-Related Hospitalizations (Overall), 2004.	43
1b. Hospitalizations for All Injuries by Sex, 2004	13	5e. Firearm-Related Hospitalizations by Sex, 2004.	44
1c. Hospitalizations for All Injuries by Age, 2004	14	5f. Firearm-Related Hospitalizations by Age, 2004.	45
2. Traumatic Brain Injury Indicators (TBI)	15	5g. Percentage of Adults Living in Homes with Loaded and Unlocked Firearms, 2004, Behavioral Risk Factor Surveillance System	46
2a. TBI Fatalities (Overall), 2004	16	6. Homicide Indicators	47
2b. TBI Fatalities by Sex, 2004	17	6a. Homicide (Overall), 2004	48
2c. TBI Fatalities by Age, 2004	18	6b. Homicide by Sex, 2004	49
2d. TBI Hospitalizations (Overall), 2004	19	6c. Homicide by Age, 2004	50
2e. TBI Hospitalizations by Sex, 2004	20	7. Suicide Indicators	51
2f. TBI Hospitalizations by Age, 2004	21	7a. Suicide (Overall), 2004	52
3. Drowning Indicators	23	7b. Suicide by Sex, 2004	53
3a. Unintentional Drowning Fatalities (Overall), 2004	24	7c. Suicide by Age, 2004	54
3b. Unintentional Drowning Fatalities by Sex, 2004	25	7d. Suicide Attempt Hospitalizations (Overall), 2004	55
3c. Unintentional Drowning Fatalities by Age, 2004	26	7e. Suicide Attempt Hospitalizations by Sex, 2004	56
3d. Drowning-Related Hospitalizations (Overall), 2004.	27	7f. Suicide Attempt Hospitalizations by Age, 2004	57
3e. Drowning-Related Hospitalizations by Sex, 2004.	28	8. Motor Vehicle Indicators	59
3f. Drowning-Related Hospitalizations by Age, 2004.	29	8a. Unintentional Motor Vehicle Traffic Fatalities Overall), 2004	60

8b. Unintentional Motor Vehicle Traffic Fatalities, by Sex, 2004	61
8c. Unintentional Motor Vehicle Traffic Fatalities, by Age, 2004	62
8d. Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations (Overall), 2004.	63
8e. Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations by Sex, 2004.	64
8f. Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations by Age, 2004.	65
8g. Of the Adults Reporting Drinking at Least One Alcoholic Beverage in the Past Month, the Percentage Reporting Driving after Perhaps Having Too Much to Drink in the Past Month, 2004, Behavioral Risk Factor Surveillance System	66
8h. Alcohol-Related Crash Deaths, 2004, Fatality Analysis Reporting System	67
9. Poisoning Indicators	69
9a. Poisoning Fatalities (Overall), 2004.	70
9b. Poisoning Fatalities by Sex, 2004.	71
9c. Poisoning Fatalities by Age, 2004.	72
9d. Poisoning Hospitalizations (Overall), 2004	73
9e. Poisoning Hospitalizations by Sex, 2004	74
9f. Poisoning Hospitalizations by Age, 2004	75

Foreword

The Centers for Disease Control and Prevention's (CDC) National Center for Injury Prevention and Control (NCIPC) is pleased to provide this third edition of the *State Injury Indicators Report* which presents 2004 data. We hope state public health officials and others will find the injury surveillance data presented in this report useful in making decisions about prevention efforts to reduce the burden of injury in the United States.

The methods used to prepare these data are consistent with those used in previous cycles of injury indicator data collection. The methods are based on recommendations presented in the *Consensus Recommendations for Using Hospital Discharge Data for Injury Surveillance* and from the National Public Health Surveillance System (NPHSS) indicators developed by the State and Territorial Injury Prevention Directors Association (STIPDA) and the Council of State and Territorial Epidemiologists (CSTE). With partner feedback and updated consensus recommendations, CDC continuously modifies and updates the instructions and methods used to prepare these data.

Thirty-four states participated in data submission either as a part of CDC Program Announcement 05027 or through voluntary submission. As more states and U.S. territories participate in this surveillance effort, a broader picture of the burden of injuries can be presented and priorities for prevention can be targeted. NCIPC looks forward to continuing to work with partners to advance and improve injury surveillance using state-based data on fatal and nonfatal injuries.

Abbreviations

BAC	Blood alcohol concentration
BRFSS	Behavioral Risk Factor Surveillance System
CDC	Centers for Disease Control and Prevention
CSTE	Council of State and Territorial Epidemiologists
FARS	Fatality Analysis Reporting System
HDD	Hospital discharge data
ICD-10	International Classification of Diseases – Tenth Revision
ICD-9-CM	International Classification of Diseases – Ninth Revision – Clinical Modification
NCIPC	National Center for Injury Prevention and Control
NPHSS	National Public Health Surveillance System
NVSS	National Vital Statistics System
SAVIR	Society for Advancement of Violence and Injury Research
STIPDA	State and Territorial Injury Prevention Directors Association
TBI	Traumatic brain injury

Introduction

Surveillance is one of the most important and basic elements of injury prevention and control. It helps determine the magnitude of injury morbidity and mortality, the leading causes of injury, and the population groups and behaviors associated with the greatest risk of injury. Surveillance data are also fundamental to determining program and prevention priorities. Furthermore, these data are crucial for evaluating the effectiveness of program activities and for identifying problems that need further investigation.

Injury continues to be the leading cause of death and disability among children and young adults.¹ In 2004, more than 167,000 people died from injuries in the U.S. Of the deaths, 27% were from motor-vehicle crashes, 19% were from suicide, and 10% were from homicide.¹ Additionally, in 2004, more than 29.6 million people were treated for injuries in U.S. emergency departments.¹ The economic cost of injuries is also significant. The total cost of the 50 million medically treated injuries sustained in 2000 is estimated to be \$406 billion in medical expenses and productivity losses.²

The mission of public health includes prevention, mitigation, assurance of access to treatment, and reduction of injury-related disability and death.³ The scope of public health encompasses injuries involving any mechanism (e.g., firearm, motor vehicle, or burn) and includes both intentional and unintentional injuries. An important part of the public health mission is to emphasize that injuries are preventable and to dispel the misconception that injuries are unavoidable.

Recognizing the need for more comprehensive injury surveillance data, the State and Territorial Injury Prevention Directors Association (STIPDA) produced *Consensus Recommendations for Injury Surveillance in State Health Departments* in 1999.⁴ These recommendations were developed by a working group representing STIPDA, the Council of State and Territorial Epidemiologists (CSTE), the Centers for Disease Control and Prevention (CDC) and its National Center for Injury Prevention and Control (NCIPC), and the Society for Advancement of Violence and Injury Research (SAVIR).

The *State Health Department Consensus Recommendations* identified 14 specific injuries and injury risk factors to be placed under surveillance by all states and 11 data sets to monitor these injuries and risk factors.⁴ The goal is to improve state-based injury surveillance to better support injury prevention programs and policies. By enhancing and standardizing methods used to track fatal and nonfatal injuries at the state level, its integration with overall public health surveillance as part of the National Public Health Surveillance System (NPHSS) will be much easier.⁵ With the *State Health Department Consensus Recommendations*, CSTE and STIPDA developed injury indicators that were formally adopted for inclusion in NPHSS.^{6,7} The NPHSS injury indicators add to other indicators developed by CSTE for chronic diseases and other areas.⁶

The *Consensus Recommendations for Using Hospital Discharge Data for Injury Surveillance*, published in 2003, has provided clear and specific recommendations about the evaluation and use of hospital discharge data.⁸ It presents important considerations for the evaluation of data quality and outlines the methods for developing an injury hospitalization data set.

The collection and dissemination of injury indicators in this report is the implementation of the foundation laid by the publication of the STIPDA and CSTE documents. It is the next step in the collaborative process of building strong state-based injury surveillance that is comparable between jurisdictions.

Discussion of Indicators

Methods

The data presented in this report are drawn from a combination of state vital statistics records, state hospital discharge data (HDD), the National Vital Statistics System (NVSS), the Behavioral Risk Factor Surveillance System (BRFSS), and the Fatality Analysis Reporting System (FARS). The methods used to prepare this report were fully delineated in the companion publication, *State Injury Indicators: Instructions for Preparing 2004 Data*.⁹ In brief, injury hospitalization cases were identified from state-based HDD using the principal diagnosis (or first-listed diagnosis code) along with other selective factors. Injury deaths were identified from vital statistics data using the underlying-cause-of-death codes. For the fatal TBI indicator, all multiple-cause-of-death fields were searched. For comparison purposes, national rates for the fatality indicators were generated using the NVSS.

What is an Injury Indicator?

An injury indicator describes a health outcome of an injury, such as hospitalization or death, or a factor known to be associated with an injury.

The following *International Classification of Diseases – Tenth Revision* (ICD-10)¹⁰ and *International Classification of Diseases – Ninth Revision – Clinical Modification* (ICD-9-CM)¹¹ code groupings were used to identify cases of injury-related death and hospitalization, respectively.

Hospitalizations for all injuries

Diagnosis codes: 800–909.2, 909.4, 909.9–994.9, 995.5–995.59, 995.80–995.85

Traumatic brain injury fatalities

Diagnosis codes: S01.0–S01.9, S02.0, S02.1, S02.3, S02.7–S02.9, S06.0–S06.9, S07.0, S07.1, S07.8, S07.9, S09.7–S09.9, T01.0, T02.0, T04.0, T06.0, T90.1, T90.2, T90.4, T90.5, T90.8, T90.9

Traumatic brain injury hospitalizations

Diagnosis codes: 800.0–801.9, 803.0–804.9, 850–854, 959.01

Unintentional drowning fatalities

W65–W74, V90, V92

Nonfatal drowning hospitalizations

Diagnosis code: 994.1
E830, E832, E910, E954, E964, E984

Unintentional fire-related fatalities

X00–X09

Unintentional fire-related hospitalizations

E890–E899

Firearm-related fatalities

W32–W34, X72–X74, X93–X95, Y22–Y24, Y35.0

Firearm-related hospitalizations

E922.0–E922.3, E922.8, E922.9, E955.0–E955.4, E965.0–E965.4, E985.0–E985.4, E970

Homicide

X85–Y09, Y87.1

Suicide

X60–X84, Y87.0

Suicide attempt hospitalizations

E950–E959

Unintentional motor vehicle traffic fatalities

V02–V04 (.1, .9), V09.2, V12–V14 (.3–.9), V19 (.4–.6), V20–V28 (.3–.9), V29–V79 (.4–.9), V80 (.3–.5), V81.1, V82.1, V83–V86 (.0–.3), V87 (.0–.8), V89.2

Unintentional motor vehicle traffic and non-traffic hospitalizations

E810–E825

Poisoning fatalities

X40–X49, X60–X69, X85–X90, Y10–Y19, Y35.2

Poisoning hospitalizations

E850–E858, E860–E869, E950–E952, E962, E972, E980–E982

For the BRFSS data, weighted percentages were generated by state by using the 2004 public-use data file for the questions of interest.¹² Alcohol-related crash death rates were calculated by state using the number of alcohol-related crash deaths from FARS data as the numerator and state-provided population totals for the denominator.¹³ An alcohol-related crash death is defined as a death due to a motor vehicle traffic crash where either the driver or a non-occupant (e.g., pedestrian) had a blood alcohol concentration (BAC) ≥ 0.01 g/dL.


Hospital Discharge Data Quality

The quality and completeness of hospital discharge data is affected by factors including the completeness of external-cause-of-injury coding and the inclusion or exclusion of residents treated in out-of-state hospitals. Both incomplete external-cause-of-injury coding and the loss of state residents treated in out-of-state hospitals decrease the number of cases that can be identified and result in reporting artificially lower rates. Readers are encouraged to consider the representativeness of data from individual state hospital discharge data systems when interpreting rates based on hospital discharge data.

For the 2004 data, hospital discharge data with external cause coding ranged from 66% to 100% (Table A). Ninety-four percent of the reporting states have $>80\%$ of their hospitalization data external-cause-of-injury coded; this is an increase from 56% in the 2nd edition. A parallel increase is seen in the number of states with $>90\%$ of hospitalization data external-cause-of-injury coded; 66% have $>90\%$ coded in this edition while only 35% had $>90\%$ coded in the second edition.¹⁴

Highlights

The injury indicator data presented here provides a unique state-specific look at the magnitude of nine injury topic areas. With 34 states participating, an increase from 26 states in the second edition, the national representativeness of the data continues to increase. By using state-specific datasets and an agreed upon methodology, states are able to compare findings both within indicators and among states. Within a state, comparisons can be made relative to the magnitude and impact of the various causes of injury. Also, as the completeness of reporting and external-cause-of-injury data continues to improve, interstate comparisons become possible.



The thirty-four states are geographically diverse and both urban and rural areas are represented. These states, when considered together, provide an overview of injury in each of the indicator areas.

All-Injury Indicators

The individual state rate of Hospitalizations for All Injuries ranged from 321.0 to 812.3 per 100,000 persons (Figure 1a). The median rate for Hospitalizations for All Injuries across the reporting states was 555.4 per 100,000 persons.

The individual state rate of Hospitalizations for All Injuries ranged from 333.0 to 824.1 per 100,000 persons among males and from 296.5 to 784.2 per 100,000 persons among females (Figure 1b). The median rate for Hospitalizations for All Injuries was 585.7 per 100,000 persons among males and 509.9 per 100,000 persons among females.

In each of the reporting states, the highest Hospitalizations for All Injuries rate was among people 85 years and older, with a range of 2,557.7 to 6,845.1 per 100,000 persons (Figure 1c).

Table A
Factors Affecting Representativeness of 2004 Hospital Discharge Data

	External Cause Percentage*	External Cause Percentage*	Inclusion of Readmissions and Transfers	Cross-Border Hospitalization [†]	Complete Hospital Participation
Arizona		92.8%	Yes	No	Yes
Arkansas		100.0%	Yes	Yes	Yes
California		100.0%	Yes	No	Yes
Colorado		98.8%	Yes	No	Yes
Connecticut		94.7%	Yes	No	No
Florida		85.7%	Yes	No	Yes
Georgia		95.5%	Yes	Unknown	Yes
Hawaii		86.4%	Yes	No	Yes
Kansas		81.5%	Yes	Yes	No
Kentucky		78.6%	Yes	Yes	No
Louisiana		‡	‡	‡	‡
Maine		86.2%	Yes	No	Yes
Maryland		95.6%	Yes	Yes	Yes
Massachusetts		98.9%	Yes	No	Yes
Michigan		86.2%	Yes	No	Yes
Minnesota		84.7%	Yes	No	No
Nebraska		97.0%	Yes	Yes	No
Nevada		66.0%	Yes	No	Yes
New Hampshire		99.8%	Yes	Yes	Yes
New Mexico		‡	‡	‡	‡
New York		99.1%	Yes	No	Yes
North Carolina		90.2%	Yes	No	Yes
Ohio		99.0%	Yes	Yes	Yes
Oklahoma		88.2%	Yes	No	Yes
Oregon		89.0%	Yes	No	Yes
Pennsylvania		97.7%	Yes	Unknown	Yes
Rhode Island		96.7%	Yes	No	Yes
South Carolina		93.0%	Yes	Yes	Yes
Tennessee		96.3%	Yes	No	Yes
Utah		92.0%	Yes	No	Yes
Vermont		98.4%	Yes	No	Yes
Virginia		82.7%	Yes	No	Yes
Washington		98.4%	Yes	Yes	Yes
Wisconsin		100.0%	Yes	No	Yes

0 10 20 30 40 50 60 70 80 90 100

* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.

[†] Subjective assessment by health department staff whether a substantial proportion of state residents injured in-state who require hospitalization are hospitalized in a neighboring state.

‡ No data available.

Traumatic Brain Injury (TBI) Indicators

The individual state rate of TBI Fatalities ranged from 7.1 to 25.2 per 100,000 persons (Figure 2a). The national rate for TBI Fatalities was 17.9 per 100,000 persons.

The individual state rate of TBI Fatalities ranged from 10.9 to 40.9 per 100,000 persons among males and from 3.8 to 12.9 per 100,000 persons among females (Figure 2b). The national rate for TBI Fatalities was 27.8 per 100,000 persons among males and 9.1 per 100,000 persons among females.

In each of the reporting states, the highest TBI Fatality rate was among people 65 years and older, with a range of 24.2 to 61.9 per 100,000 persons (Figure 2c).

The individual state rate of TBI Hospitalizations ranged from 37.2 to 122.2 per 100,000 persons (Figure 2d). The median rate for TBI Hospitalizations across all reporting states was 74.2 per 100,000 persons.

The individual state rate of TBI Hospitalizations ranged from 50.1 to 164.4 per 100,000 persons among males and from 26.0 to 82.2 per 100,000 persons among females (Figure 2e). The median rate for TBI Hospitalizations was 96.7 per 100,000 persons among males and was 49.1 per 100,000 persons among females.

In each of the reporting states, the highest TBI Hospitalization rate was among people 85 years and older, with a range of 228.4 to 678.7 per 100,000 persons (Figure 2f).

Drowning Indicators

The individual state rate of Unintentional Drowning Fatalities ranged from too few cases to report rates to 2.6 per 100,000 persons (Figure 3a). The national rate for Unintentional Drowning Fatalities was 1.3 per 100,000 persons.

The individual state rate of Unintentional Drowning Fatalities ranged from too few cases to report rates to 4.5 per 100,000 persons among males, and from too few cases to report rates to 1.2 per 100,000 persons among females (Figure 3b). The national rate for Unintentional Drowning Fatalities was 2.0 per 100,000 persons among males and 0.5 per 100,000 persons among females.

The individual state rate of Drowning-Related Hospitalizations ranged from too few cases to report rates to 4.4 per 100,000 persons (Figure 3d). The median rate for Drowning-Related Hospitalizations across reporting states with case counts large enough to present rates was 0.7 per 100,000 persons.

The individual state rate of Drowning-Related Hospitalizations ranged from too few cases to report rates to 6.4 per 100,000 persons among males, and from too few cases to report rates to 1.7 per 100,000 persons among females (Figure 3e). Median rates for male and female Drowning-Related Hospitalizations were not calculated due to low case counts.

Fire-Related Indicators

The individual state rate of Unintentional Fire-Related Fatalities ranged from too few cases to report rates to 2.5 per 100,000 persons (Figure 4a). The national rate for Unintentional Fire-Related Fatalities was 1.1 per 100,000 persons.

The individual state rate of Unintentional Fire-Related Fatalities ranged from too few cases to report rates to 3.6 per 100,000 persons among males, and from too few cases to report rates to 1.7 per 100,000 persons among females (Figure 4b). Among males, the national rate for Unintentional Fire-Related Fatalities was 1.4 per 100,000 persons and among females the national rate was 0.8 per 100,000 persons.

The individual state rate of Unintentional Fire-Related Hospitalizations ranged from too few cases to report rate to 6.5 per 100,000 persons (Figure 4d). The median rate for Unintentional Fire-Related Hospitalizations across reporting states with case counts large enough to present rates was 3.3 per 100,000 persons.

The individual state rate of Unintentional Fire-Related Hospitalizations ranged from too few cases to report rates to 10.2 per 100,000 persons among males, and from too few cases to report rates to 3.4 per 100,000 persons among females (Figure 4e). The median rate for Unintentional Fire-Related Hospitalizations was 5.3 per 100,000 persons among males and 2.1 per 100,000 persons among females.

Firearm–Related Indicators

The individual state rate of Firearm-Related Fatalities ranged from 3.1 to 19.6 per 100,000 persons (Figure 5a). The national rate for Firearm-Related Fatalities was 10.0 per 100,000 persons.

The individual state rate of Firearm-Related Fatalities ranged from 5.5 to 35.0 per 100,000 persons among males, and from too few cases to report rates to 5.3 per 100,000 persons among females (Figure 5b). The national rate for Firearm-Related Fatalities was 17.7 per 100,000 persons among males and 2.7 per 100,000 persons among females.

In the reporting states, the highest Firearm-Related Fatality rate was most commonly found among persons in the 15-24 year age group (Figure 5c). However, in some states the highest rates were in persons 25-44 years, 65 years and older, or 45-64 years.

The individual state rate of Firearm-Related Hospitalizations ranged from too few cases to report rates to 15.8 per 100,000 persons (Figure 5d). The median rate for Firearm-Related Hospitalizations across reporting states with case counts large enough to present rates was 7.5 per 100,000 persons.

The individual state rate of Firearm-Related Hospitalizations ranged from too few cases to report rates to 29.2 per 100,000 persons among males, and from too few cases to report rates to 5.3 per 100,000 persons among females (Figure 5e). The median rate for Firearm-Related Hospitalizations was 13.3 per 100,000 persons among males and 2.1 per 100,000 persons among females.

In most of the reporting states, the highest Firearm-Related Hospitalization rate was among person 15-24 years (Figure 5f).

Based on results from the Behavioral Risk Factor Surveillance System, 0.6% to 10.3% of adults lived in a home with a loaded and unlocked firearm for the 34 states reporting indicator data (Figure 5g). The median percent was 3.8%.

Homicide Indicators

The individual state rate of Homicides ranged from too few cases to report rates to 13.0 per 100,000 persons (Figure 6a). The national rate for Homicides was 5.9 per 100,000 persons.

The individual state rate of Homicides ranged from too few cases to report rates to 21.2 per 100,000 persons among males, and from too few cases to report rates to 5.7 per 100,000 persons among females (Figure 6b). The national rate for Homicides was 9.2 per 100,000 persons among males and 2.6 per 100,000 persons among females.

In the reporting states, the highest Homicide rate was most commonly found in the 15-24 year age group (Figure 6c). However, in several states the highest rate was among person 25-44 years.

Suicide Indicators

The individual state rate of Suicides ranged from 6.3 to 18.9 per 100,000 persons (Figure 7a). The national rate for Suicides was 10.9 per 100,000 persons.

The individual state rate of Suicides ranged from 10.6 to 31.7 per 100,000 persons among males, and from too few cases to report rates to 8.0 per 100,000 persons among females (Figure 7b). The national rate for Suicides was 18.0 per 100,000 persons among males and 4.5 per 100,000 persons among females.

In the reporting states, the highest Suicide rate was most commonly found in the 45-64 year age group (Figure 7c). However, in some states the highest rates were for persons 25-44 years, 65 years and older, or 15-24 years.

The individual state rate of Suicide Attempt Hospitalizations ranged from 29.0 to 87.6 per 100,000 persons (Figure 7d). The median rate for Suicide Attempt Hospitalizations across all reporting states was 47.7 per 100,000 persons.

The individual state rate of Suicide Attempt Hospitalizations ranged from 21.1 to 64.4 per 100,000 persons among males and from 33.5 to 115.5 per 100,000 persons among females (Figure 7e). The median rate for Suicide Attempt Hospitalizations was 36.2 per 100,000 persons among males and 56.9 per 100,000 persons among females.

In the reporting states, the highest Suicide Attempt Hospitalization rate was most commonly found in the 15-24 year age group (Figure 7f). However, in approximately a third of the reporting states, the highest rate was for person 25-44 years.

Motor Vehicle Indicators

The individual state rate of Unintentional Motor Vehicle Traffic Fatalities ranged from 6.6 to 27.4 per 100,000 persons (Figure 8a). The national rate for Unintentional Motor Vehicle Traffic Fatalities was 14.7 per 100,000 persons.

The individual state rate of Unintentional Motor Vehicle Traffic Fatalities ranged from 10.7 to 36.5 per 100,000 persons among males, and from too few cases to report rates to 18.8 per 100,000 persons among females (Figure 8b). The national rate for Unintentional Motor Vehicle Traffic Fatalities was 20.6 per 100,000 persons among males and 9.0 per 100,000 persons among females.

In the reporting states, the highest Unintentional Motor Vehicle Traffic Fatality rate was most commonly found in the 15-24 years age group (Figure 8c). However, in a few states, the highest rate was among those 65 years and older.

The individual state rate of Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations ranged from 35.0 to 131.0 per 100,000 persons (Figure 8d). The median rate for Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations across all reporting states was 88.0 per 100,000 persons.

The individual state rate of Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations ranged from 42.8 to 164.3 per 100,000 persons among males and from 27.7 to 96.5 per 100,000 persons among females (Figure 8e). The median rate for Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations was 114.2 per 100,000 persons among males and 63.6 per 100,000 persons among females.

In the reporting states, the highest Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalization rate was most commonly found in the 15-24 years age group (Figure 8f). However, in a few states, the highest rate was among persons 75-84 years or persons 85 years and older.

Based on results from the Behavioral Risk Factor Surveillance System from the 34 states reporting indicator data, 2.2% to 6.4% of the adults who reported drinking at least one alcoholic beverage in the past month also reported driving after perhaps having too much to drink (Figure 8g). The median percent was 3.4%.

Based on data from the Fatality Analysis Reporting System, alcohol-related crash death rates ranged from 2.9 to 11.1 per 100,000 (Figure 8h). The alcohol-related crash death rate for the United States was 5.7 per 100,000.

Poisoning Indicators

The individual state rate of Poisoning Fatalities ranged from 3.1 to 18.6 per 100,000 persons (Figure 9a). The national rate for Poisoning Fatalities was 10.3 per 100,000 persons.

The individual state rate of Poisoning Fatalities ranged from 3.9 to 24.0 per 100,000 persons among males, and from too few cases to report rates to 14.0 per 100,000 persons among females (Figure 9b). The national rate for Poisoning Fatalities was 13.3 per 100,000 persons among males and 7.4 per 100,000 persons among females.

In the reporting states, the highest Poisoning Fatality rate was most commonly found in the 25-44 year age group (Figure 9c). However, in approximately a third of the reporting states, the highest rate was for person 45-64 years.

The individual state rate of Poisoning Hospitalizations ranged from 40.6 to 105.2 per 100,000 persons (Figure 9d). The median rate for Poisoning Hospitalizations across all reporting states was 74.1 per 100,000 persons.

The individual state rate of Poisoning Hospitalizations ranged from 32.6 to 99.5 per 100,000 persons among males and 48.7 to 126.9 per 100,000 persons among females (Figure 9e). The median rate for Poisoning Hospitalizations was 62.4 per 100,000 persons among males and 81.1 per 100,000 persons among females.

In the reporting states, the highest Poisoning Hospitalization rate was most commonly found in the 35-44 year age group (Figure 9f). However, in some states the highest rates were for persons 15-24 years, 25-34 years, or 45-54 years.

Limitations and Future Efforts

While this report provides important information for understanding injury, the following limitations should be considered:

- Only injuries severe enough to lead to either hospitalization or death are reported. Injuries that lead to emergency department visits, physician office visits, or remain untreated need to be accounted for if the total burden of injury is to be understood.
- Not all causes of injury have been included in this report. The absence of fall-related indicators is noted.
- Representation from all states and territories has not yet been achieved.
- Quality and completeness of external-cause-of-injury coding, while improving, continues to be incomplete.

Future editions of the *State Injury Indicators Report* will strive to address these limitations. NCIPC, STIPDA, and CSTE, in conjunction with participating states, will continue to refine current indicators and define new ones for inclusion in future editions of this report. In keeping with the consensus building process and the publication of *Consensus Recommendations for Surveillance of Falls and Fall-Related Injuries*¹⁵ the next edition will include the recommended indicators for fall-related injuries. The incorporation of emergency department data into future editions is under exploration and consideration. Finally, submission of data from additional states and territories is encouraged and welcomed.


References

1. Centers for Disease Control and Prevention. Web-based injury statistics query and reporting system (WISQARS). Atlanta (GA); 2005. [cited 2007 Mar 28]. Available from: www.cdc.gov/ncipc/wisqars.
2. Finkelstein EA, Corso PS, Miller TR, Associates. Incidence and economic burden of injuries in the United States. New York: Oxford University Press; 2006.
3. Institute of Medicine (US). Reducing the burden of injury, advancing prevention and treatment. Washington (DC): National Academy Press; 1999.
4. State and Territorial Injury Prevention Directors Association (STIP-DA). Consensus recommendations for injury surveillance in state health departments. Marietta (GA): STIPDA; 1999.
5. Meriwether RA. Blueprint for a national public health surveillance system for the 21st century. *J Public Health Manag Pract*. 1996;216-23.
6. Council of State and Territorial Epidemiologists. Injury control and prevention position statement. [cited 2007 Mar 29]. Available from: <http://www.cste.org/PositionStatement.asp>.
7. State and Territorial Injury Prevention Directors Association (STIP-DA). Resolutions, October 1999. [cited 2007 Mar 29]. Available from: <http://www.stipda.org/displaycommon.cfm?an=1&subarticlenbr=16>.
8. Injury Surveillance Workgroup. Consensus recommendations for using hospital discharge data for injury surveillance. Marietta (GA): State and Territorial Injury Prevention Directors Association; 2003.
9. Johnson RL, Sarmiento K. State injury indicators: Instructions for preparing 2004 data. Atlanta (GA): Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2006.
10. International Classification of Diseases 10th Revision. [cited 2007 Mar 29]. Available from: <http://www.who.int/classifications/icd/en/index.html>.
11. International Classification of Diseases 9th Revision Clinical Modification. [cited 2007 Mar 29]. Available from: www.cdc.gov/nchs/about/otheract/icd9/abtcd9.htm.
12. Centers for Disease Control and Prevention. Behavioral risk factor surveillance system: technical documents and survey data. [cited 2007 Mar 28]. Available from: http://www.cdc.gov/brfss/technical_infodata/surveydata/2004.htm.
13. National Highway Traffic Safety Administration. Traffic safety facts 2004. [cited 2007 Mar 28]. Available from: <http://www-nrd.nhtsa.dot.gov/pdf/nrd-30/NCSA/TSFAnn/2004HTML/TSF2004.htm>.
14. Thomas C, Butler J, Davies M, Johnson R. State injury indicators report, second edition – 1999 data. Atlanta (GA): Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2004.
15. Injury Surveillance Workgroup on Falls. Consensus recommendations for surveillance of falls and fall-related injuries. Atlanta (GA): State and Territorial Injury Prevention Directors Association; 2006.

All-Injury Indicators

- 1a. Hospitalizations for All Injuries (Overall), 2004
- 1b. Hospitalizations for All Injuries by Sex, 2004
- 1c. Hospitalizations for All Injuries by Age, 2004

Figure 1a
Hospitalizations for All Injuries (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 1b
Hospitalizations for All Injuries by Sex, 2004**



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 1c
Hospitalizations for All Injuries by Age[#], 2004**

State	0-4		5-14		15-24		25-34		35-44		45-54		55-64		65-74		75-84		85+		
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	
Arizona	1,582	367.4	2,242	257.7	5,393	649.9	4,371	515.9	4,361	497.2	4,045	564.0	3,021	599.1	3,232	784.7	5,181	1,942.3	4,143	5,348.0	
Arkansas	285	153.6	563	150.3	2,034	513.3	1,877	521.3	2,146	557.4	2,014	535.9	1,602	544.4	1,920	935.6	3,131	2,445.3	2,708	5,660.3	
California	6,956	264.1	11,071	205.4	25,778	498.8	19,493	371.1	21,402	386.3	21,020	432.5	15,764	488.8	15,440	791.8	26,594	1,957.1	21,619	4,205.9	
Colorado	492	146.4	1,022	161.2	3,284	488.9	2,701	402.3	2,933	404.8	3,138	452.9	2,245	512.1	2,167	828.5	3,591	2,197.5	3,073	5,582.8	
Connecticut	506	237.5	854	178.4	1,980	432.8	1,527	373.0	1,977	350.2	1,922	361.0	1,509	403.3	1,565	722.5	2,990	1,708.4	2,894	3,526.0	
Florida	2,860	262.1	4,697	211.3	11,721	523.6	10,027	468.1	12,267	484.9	12,651	531.6	9,791	524.9	11,573	784.6	21,184	1,974.2	17,691	4,660.8	
Georgia	1,307	192.5	2,000	156.8	5,195	405.9	4,591	336.7	5,276	378.9	5,036	423.5	3,884	483.7	3,978	834.7	5,962	2,166.1	4,520	4,744.3	
Hawaii	226	254.6	369	229.5	944	538.3	664	412.0	706	391.8	791	424.8	561	405.5	526	664.9	976	1,444.7	913	3,602.4	
Kansas	470	126.4	693	359.7	2,056	491.2	1,495	417.0	1,703	439.4	1,811	462.2	1,328	510.6	1,563	918.7	3,346	2,586.0	3,198	5,808.7	
Kentucky	395	148.2	824	150.7	2,628	453.5	2,617	460.7	3,006	483.7	2,870	474.4	2,363	537.9	2,700	959.8	4,210	2,352.0	3,005	5,091.1	
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	166	245.5	305	192.4	1,030	570.9	698	479.1	857	418.5	895	414.7	732	473.3	849	889.9	1,421	2,051.7	1,331	5,304.7	
Maryland	548	146.3	1,132	145.3	4,532	594.4	3,339	469.7	4,174	466.6	3,775	455.6	2,803	488.9	2,410	739.2	4,374	1,935.6	3,338	4,033.7	
Massachusetts	756	191.1	1,416	173.7	3,920	460.4	3,013	342.2	4,089	396.8	4,006	427.5	3,383	518.9	3,571	901.6	7,063	2,192.6	6,619	4,862.4	
Michigan	1,681	258.7	2,775	192.8	6,951	482.3	5,795	443.8	7,250	478.8	7,663	512.2	5,271	516.8	5,109	835.0	9,054	1,969.7	7,453	4,257.2	
Minnesota	745	224.4	1,430	208.7	3,768	499.8	2,712	399.7	3,350	422.4	3,573	473.2	2,563	525.3	2,617	872.0	4,822	2,223.7	4,576	4,659.2	
Nebraska	131	107.3	261	110.1	666	250.1	477	208.0	565	231.2	690	276.8	601	359.5	777	697.8	1,539	1,820.4	1,505	4,191.0	
Nevada	327	190.8	563	167.9	1,399	406.1	1,276	363.4	1,438	397.9	1,430	437.3	1,156	469.9	1,142	729.5	1,566	1,762.0	970	4,397.1	
New Hampshire	93	125.3	223	127.7	663	371.9	516	341.6	774	359.7	679	327.0	566	407.4	582	703.3	1,069	1,941.0	956	4,478.7	
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	4,035	323.8	6,031	237.0	13,292	509.9	10,522	399.6	13,376	446.9	12,387	452.0	9,368	475.5	9,636	785.9	17,047	1,867.7	15,107	4,268.9	
North Carolina	968	167.5	1,804	156.3	4,966	402.5	4,685	390.0	5,693	439.0	5,406	453.6	4,379	504.1	4,993	914.6	7,996	2,290.6	5,706	4,682.6	
Ohio	478	65.5	1,203	77.1	4,659	288.3	4,114	281.0	4,862	288.7	4,683	275.1	3,160	268.5	3,387	453.5	6,371	1,118.3	5,331	2,557.7	
Oklahoma	863	356.3	1,275	271.2	2,940	551.7	2,187	465.7	2,678	548.3	2,509	508.8	1,954	538.8	2,330	936.8	3,730	2,310.9	3,150	5,800.6	
Oregon	509	223.0	876	174.9	2,203	442.0	1,812	366.2	2,142	387.1	2,557	480.3	1,835	575.0	1,711	749.9	3,055	1,822.6	2,715	4,556.2	
Pennsylvania	2,116	289.7	3,982	247.5	10,390	590.5	8,044	544.9	10,334	572.2	10,306	571.4	7,754	585.2	8,677	910.9	16,751	2,429.9	13,749	5,401.5	
Rhode Island	95	154.4	133	95.8	363	232.9	308	221.5	419	251.9	420	264.8	361	329.0	334	517.5	776	1,298.5	801	3,046.8	
South Carolina	546	194.8	1,245	219.4	2,312	381.7	2,140	376.8	2,570	420.5	2,501	420.8	1,965	436.1	2,032	711.2	2,900	1,654.9	2,069	3,480.2	
Tennessee	763	195.5	1,406	174.5	4,414	546.2	4,243	522.5	5,162	585.7	4,868	575.2	3,851	621.4	4,000	1,010.1	6,147	2,482.4	4,575	5,088.1	
Utah	510	208.8	768	187.8	2,441	574.5	1,862	445.4	1,909	620.5	2,006	719.0	1,692	947.6	1,595	1,456.1	2,381	3,294.9	1,788	6,845.1	
Vermont	42	125.8	130	172.0	394	439.9	235	337.7	312	328.2	366	352.6	267	362.5	309	756.8	506	1,778.4	512	4,460.7	
Virginia	852	171.0	1,355	128.4	4,033	397.3	3,384	288.4	3,951	361.4	4,131	413.2	3,231	415.2	3,710	818.5	6,111	2,099.4	4,740	4,621.6	
Washington	777	193.7	1,450	169.3	4,391	497.4	3,747	449.5	4,847	514.5	5,573	599.5	4,256	680.2	4,020	1,143.9	6,765	2,748.5	5,852	5,912.0	
Wisconsin	851	250.2	1,437	196.0	3,642	444.2	2,682	383.5	3,523	417.4	3,516	423.4	2,639	477.6	2,959	843.5	5,258	2,045.5	5,116	4,582.8	

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.


¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Traumatic Brain Injury (TBI) Indicators

- 2a. TBI Fatalities (Overall), 2004
- 2b. TBI Fatalities by Sex, 2004
- 2c. TBI Fatalities by Age, 2004
- 2d. TBI Hospitalizations (Overall), 2004
- 2e. TBI Hospitalizations by Sex, 2004
- 2f. TBI Hospitalizations by Age, 2004

Figure 2a
TBI Fatalities (Overall), 2004




‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 2b
TBI Fatalities by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 2c
TBI Fatalities by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	48	3.7	239	28.8	340	19.7	288	23.6	367	48.5
Arkansas	43	7.7	121	30.5	195	26.2	174	26.0	152	39.9
California	183	2.3	816	15.8	1,137	10.5	1,118	13.8	1,499	39.2
Colorado	46	4.7	176	26.2	287	20.6	273	24.1	233	48.5
Connecticut	8		55	12.0	78	8.0	84	9.3	155	32.7
Florida	140	4.2	521	23.3	872	18.7	925	21.8	1,161	39.7
Georgia	103	5.3	328	25.6	562	20.4	445	22.3	400	47.2
Hawaii	5		22	12.5	40	11.7	40	12.3	68	39.5
Kansas	28	5.0	102	24.4	154	20.6	133	20.4	210	59.2
Kentucky	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Louisiana	63	6.5	235	33.3	336	27.1	253	23.4	190	36.0
Maine	7		37	20.5	64	18.3	53	14.3	63	33.2
Maryland	22	1.9	103	13.5	171	10.7	150	10.7	220	34.7
Massachusetts	16		71	8.3	87	4.6	105	6.6	207	24.2
Michigan	67	3.2	219	15.2	384	13.6	356	14.1	476	38.2
Minnesota	37	3.6	110	14.6	182	12.4	167	13.4	260	42.3
Nebraska	16		78	29.3	68	14.4	78	18.7	137	59.1
Nevada	12		45	13.1	123	17.3	125	21.8	132	49.3
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	26	6.4	96	32.7	128	25.5	116	24.5	107	46.6
New York	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
North Carolina	86	4.9	346	29.5	569	22.5	403	19.8	542	52.5
Ohio	70	3.1	355	22.0	514	16.3	523	18.2	632	41.4
Oklahoma	43	6.0	161	30.2	234	24.4	211	24.7	229	49.3
Oregon	32	4.4	106	21.3	155	14.8	205	24.1	227	49.8
Pennsylvania	59	2.5	308	17.5	527	16.1	505	16.1	785	41.4
Rhode Island	¶		31	19.9	24	7.9	29	10.8	64	42.5
South Carolina	31	3.7	195	32.2	355	30.1	268	25.6	213	40.9
Tennessee	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Utah	20	3.1	75	17.7	125	17.2	102	22.3	86	41.4
Vermont	¶		21	23.4	36	21.9	32	18.0	50	61.9
Virginia	42	2.7	258	25.4	380	16.8	315	17.7	407	48.1
Washington	39	3.1	212	24.0	308	17.3	303	19.5	419	60.2
Wisconsin	32	3.0	204	24.9	234	15.2	240	17.4	366	50.9
United States	2,224	3.7	9,106	21.8	13,822	16.4	12,541	17.7	15,425	42.5

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 2d
TBI Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 2e
TBI Hospitalizations by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 2f
TBI Hospitalizations by Age#, 2004

State	0-4		5-14		15-24		25-34		35-44		45-54		55-64		65-74		75-84		85+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	474	110.1	629	72.3	1,480	178.4	842	99.4	863	98.4	713	99.4	470	93.2	420	102.0	592	221.9	411	530.5
Arkansas	63	34.0	101	27.0	332	83.8	252	70.0	234	60.8	246	65.5	189	64.2	201	97.9	265	207.0	216	451.5
California	1,683	63.9	2,374	44.0	6,174	119.5	3,642	69.3	3,661	66.1	3,571	73.5	2,438	75.6	2,324	119.2	3,720	273.8	2,378	462.6
Colorado	153	45.5	245	38.6	887	132.1	560	83.4	528	72.9	522	75.3	361	82.3	291	111.3	387	236.8	270	490.5
Connecticut	145	68.1	143	29.9	414	90.5	196	47.9	237	42.0	239	44.9	189	50.5	209	96.5	315	180.0	270	329.0
Florida	726	66.5	992	44.6	2,571	114.9	1,618	75.5	1,735	68.6	1,749	73.5	1,203	64.5	1,463	99.2	2,352	219.2	1,825	480.8
Georgia	369	54.3	481	37.7	1,188	92.8	731	53.6	705	50.6	698	58.7	506	63.0	447	93.8	643	233.6	386	405.2
Hawaii	58	65.3	106	65.9	280	159.7	149	92.4	169	93.8	155	83.2	106	76.6	99	125.1	186	275.3	172	678.7
Kansas	85	22.9	163	84.6	440	105.1	210	58.6	193	49.8	213	54.4	141	54.2	109	64.1	176	136.0	170	308.8
Kentucky	109	40.9	167	30.5	492	84.9	348	61.3	344	55.4	322	53.2	232	52.8	245	87.1	369	206.2	201	340.5
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	34	50.3	56	35.3	227	125.8	120	82.4	120	58.6	109	50.5	87	56.3	98	102.7	129	186.3	87	346.7
Maryland	129	34.4	228	29.3	1,185	155.4	699	98.3	726	81.2	638	77.0	405	70.6	347	106.4	554	245.2	335	404.8
Massachusetts	216	54.6	183	22.4	740	86.9	397	45.1	478	46.4	416	44.4	397	60.9	389	98.2	710	220.4	578	424.6
Michigan	433	66.6	632	43.9	1,532	106.3	922	70.6	1,023	67.6	1,098	73.4	727	71.3	609	99.5	1,059	230.4	730	417.0
Minnesota	210	63.2	306	44.7	811	107.6	431	63.5	481	60.6	479	63.4	298	61.1	326	108.6	509	234.7	373	379.8
Nebraska	38	31.1	50	21.1	136	51.1	61	26.6	64	26.2	58	23.3	65	38.9	74	66.5	123	145.5	104	289.6
Nevada	88	51.3	146	43.5	301	87.4	196	55.8	222	61.4	220	67.3	131	53.2	151	96.5	153	172.1	74	335.4
New Hampshire	17		47	26.7	78	43.6	50	32.8	57	26.3	57	27.3	48	34.6	56	67.9	85	155.2	56	260.4
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	868	69.7	1,309	51.4	2,614	100.3	1,595	60.6	1,775	59.3	1,583	57.8	1,265	64.2	1,326	108.1	2,005	219.7	1,540	435.2
North Carolina	204	35.3	289	25.0	855	69.3	521	43.4	606	46.7	562	47.2	440	50.7	457	83.7	675	193.4	413	338.9
Ohio	144	19.7	297	19.0	1,164	72.0	820	56.0	883	52.4	785	46.1	526	44.7	447	59.9	717	125.9	476	228.4
Oklahoma	218	90.0	299	63.6	641	120.3	337	71.8	304	62.2	337	68.3	234	64.5	270	108.6	395	244.7	293	539.5
Oregon	134	58.7	142	28.4	405	81.3	197	39.8	224	40.5	300	56.4	178	55.8	151	66.2	230	137.2	194	325.6
Pennsylvania	610	83.5	1,053	65.5	2,827	160.7	1,611	109.1	1,822	100.9	1,766	97.9	1,225	92.4	1,267	133.0	2,162	313.6	1,533	602.3
Rhode Island	27	43.9	24	17.3	51	32.7	34	24.4	28	16.8	37	23.3	30	27.3	37	57.3	101	169.0	71	270.1
South Carolina	124	44.2	237	41.8	556	91.8	377	66.4	386	63.2	360	60.6	219	48.6	205	71.8	269	153.5	176	296.0
Tennessee	169	43.3	259	32.1	832	103.0	494	60.8	586	66.5	476	56.2	371	59.9	338	85.4	526	212.4	328	364.8
Utah	143	58.5	206	50.4	448	105.4	233	55.7	193	62.7	198	71.0	178	99.7	131	119.6	203	280.9	148	566.6
Vermont	5		21	27.8	83	92.7	33	47.4	38	40.0	37	35.6	22	29.9	38	93.1	44	154.6	32	278.8
Virginia	218	43.7	232	22.0	804	79.2	488	41.6	516	47.2	498	49.8	357	45.9	396	87.4	600	206.1	389	379.3
Washington	187	46.6	295	34.5	744	84.3	458	54.9	403	42.8	464	49.9	339	54.2	346	98.5	588	238.9	427	431.4
Wisconsin	219	64.4	306	41.7	736	89.8	414	59.2	470	55.7	500	60.2	339	61.3	389	110.9	637	247.8	550	492.7

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.


¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Drowning Indicators

- 3a. Unintentional Drowning Fatalities (Overall), 2004
- 3b. Unintentional Drowning Fatalities by Sex, 2004
- 3c. Unintentional Drowning Fatalities by Age, 2004
- 3d. Drowning-Related Hospitalizations (Overall), 2004
- 3e. Drowning-Related Hospitalizations by Sex, 2004
- 3f. Drowning-Related Hospitalizations by Age, 2004

Figure 3a
Unintentional Drowning Fatalities (Overall), 2004




‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

† Case counts are suppressed if fewer than 5 cases were reported.

Figure 3b
Unintentional Drowning Fatalities by Sex, 2004



‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 3c
Unintentional Drowning Fatalities by Age#, 2004

State	0-14		15-44		45+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	25	1.9	36	1.4	38	1.9
Arkansas	12		23	2.0	16	
California	89	1.1	160	1.0	151	1.3
Colorado	5		27	1.3	26	1.6
Connecticut	6		7		15	
Florida	82	2.5	138	2.0	170	2.4
Georgia	30	1.5	62	1.5	37	1.3
Hawaii	¶		11		17	
Kansas	10		11		14	
Kentucky	13		29	1.6	22	1.4
Louisiana	20	2.1	57	2.9	40	2.5
Maine	¶		10		9	
Maryland	7		23	1.0	26	1.3
Massachusetts	6		20	0.7	24	1.0
Michigan	16		32	0.8	41	1.1
Minnesota	11		23	1.0	22	1.2
Nebraska	6		11		10	
Nevada	¶		8		6	
New Hampshire	‡	‡	‡	‡	‡	‡
New Mexico	7		11		11	
New York	24	0.6	46	0.6	36	0.5
North Carolina	10		55	1.5	30	1.0
Ohio	31	1.4	32	0.7	21	0.5
Oklahoma	19		26	1.7	20	1.5
Oregon	11		38	2.5	22	1.7
Pennsylvania	19		35	0.7	32	0.6
Rhode Island	¶		5		6	
South Carolina	15		34	1.9	17	
Tennessee	25	2.1	49	2.0	40	1.8
Utah	10		13		6	
Vermont	¶		¶		¶	
Virginia	21	1.4	29	0.9	32	1.2
Washington	12		64	2.4	42	1.9
Wisconsin	14		27	1.1	16	
United States	775	1.3	1,600	1.3	1,340	1.3

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 3d
Drowning-Related Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 3e
Drowning-Related Hospitalizations by Sex, 2004**



‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 3f
Drowning-Related Hospitalizations by Age[#], 2004**

State	0-14		15-44		45+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	70	5.4	15		7	
Arkansas	10		12		5	
California	336	4.2	96	0.6	50	0.4
Colorado	13		¶		6	
Connecticut	5		6		10	
Florida	218	6.6	83	1.2	69	1.0
Georgia	44	2.3	12		11	
Hawaii	21	8.4	23	4.5	11	
Kansas	14		¶		¶	
Kentucky	6		¶		¶	
Louisiana	‡	‡	‡	‡	‡	‡
Maine	¶		¶		¶	
Maryland	12		7		7	
Massachusetts	15		13		12	
Michigan	44	2.1	13		8	
Minnesota	21	2.1	9		¶	
Nebraska	5		¶		¶	
Nevada	17		5		¶	
New Hampshire	5		7		¶	
New Mexico	‡	‡	‡	‡	‡	‡
New York	69	1.8	30	0.4	21	0.3
North Carolina	22	1.3	9		¶	
Ohio	17		53	1.1	22	0.5
Oklahoma	29	4.1	6		8	
Oregon	12		12		¶	
Pennsylvania	‡	‡	‡	‡	‡	‡
Rhode Island	¶		¶		¶	
South Carolina	20	2.4	8		¶	
Tennessee	22	1.8	9		¶	
Utah	19		7		¶	
Vermont	¶		¶		¶	
Virginia	19		8		7	
Washington	21	1.7	25	0.9	17	
Wisconsin	26	2.4	10		¶	

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.


¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Fire-Related Indicators

- 4a. Unintentional Fire-Related Fatalities (Overall), 2004
- 4b. Unintentional Fire-Related Fatalities by Sex, 2004
- 4c. Unintentional Fire-Related Fatalities by Age, 2004
- 4d. Unintentional Fire-Related Hospitalizations (Overall), 2004
- 4e. Unintentional Fire-Related Hospitalizations by Sex, 2004
- 4f. Unintentional Fire-Related Hospitalizations by Age, 2004

Figure 4a
Unintentional Fire-Related Fatalities (Overall), 2004




‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 4b
Unintentional Fire-Related Fatalities by Sex, 2004**



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 4c
Unintentional Fire-Related Fatalities by Age#, 2004

State	0-14		15-44		45+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	¶		6		24	1.2
Arkansas	10		27	2.4	32	3.0
California	10		37	0.2	169	1.4
Colorado	¶		¶		10	
Connecticut	¶		7		21	1.5
Florida	14		29	0.4	96	1.3
Georgia	34	1.7	31	0.8	77	2.7
Hawaii	¶		¶		¶	
Kansas	¶		11		22	2.2
Kentucky	11		15		34	2.2
Louisiana	27	2.8	20	1.0	48	3.0
Maine	¶		¶		¶	
Maryland	17		11		42	2.1
Massachusetts	¶		7		22	0.9
Michigan	16		28	0.7	74	2.0
Minnesota	6		6		14	
Nebraska	¶		¶		7	
Nevada	¶		¶		8	
New Hampshire	‡	‡	‡	‡	‡	‡
New Mexico	¶		¶		18	
New York	23	0.6	38	0.5	134	1.9
North Carolina	17		28	0.8	79	2.6
Ohio	23	1.0	22	0.5	60	1.4
Oklahoma	10		18		41	3.1
Oregon	¶		6		23	1.8
Pennsylvania	20	0.9	40	0.8	102	2.0
Rhode Island	¶		¶		¶	
South Carolina	10		23	1.3	55	3.5
Tennessee	17		19		74	3.4
Utah	¶		¶		7	
Vermont	¶		¶		¶	
Virginia	19		26	0.8	71	2.7
Washington	¶		10		34	1.5
Wisconsin	6		11		35	1.7
United States	504	0.8	739	0.6	1,983	1.9

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 4d
Unintentional Fire-Related Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 4e
Unintentional Fire-Related Hospitalizations by Sex, 2004



‡ No data available.
 § Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
 ¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 4f
Unintentional Fire-Related Hospitalizations by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	59	4.5	57	6.9	102	5.9	76	6.2	33	4.4
Arkansas	19		24	6.1	61	8.2	31	4.6	42	11.0
California	159	2.0	144	2.8	333	3.1	281	3.5	162	4.2
Colorado	10		20	3.0	45	3.2	40	3.5	22	4.6
Connecticut	13		11		39	4.0	20	2.2	14	
Florida	82	2.5	71	3.2	166	3.6	113	2.7	56	1.9
Georgia	77	3.9	83	6.5	161	5.8	152	7.6	60	7.1
Hawaii	6		¶		7		7		¶	
Kansas	23	4.1	21	5.0	27	3.6	20	3.1	23	6.5
Kentucky	19		24	4.1	77	6.5	44	4.2	23	4.4
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	8		10		14		18		8	
Maryland	27	2.3	32	4.2	57	3.6	52	3.7	38	6.0
Massachusetts	¶		18		37	1.9	56	3.5	37	4.3
Michigan	70	3.4	73	5.1	119	4.2	123	4.9	58	4.7
Minnesota	30	2.9	41	5.4	45	3.1	35	2.8	20	3.3
Nebraska	5		5		9		17		13	
Nevada	¶		9		24	3.4	17		7	
New Hampshire	¶		¶		10		13		¶	
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	113	3.0	112	4.3	230	4.1	181	3.8	167	6.7
North Carolina	39	2.3	43	3.5	90	3.6	85	4.1	56	5.5
Ohio	16		19		45	1.4	46	1.6	26	1.7
Oklahoma	15		20	3.8	37	3.9	19		26	5.6
Oregon	14		18		24	2.3	29	3.4	13	
Pennsylvania	‡	‡	110	6.3	187	5.7	185	5.9	131	6.9
Rhode Island	¶		¶		6		8		¶	
South Carolina	17		¶		20	1.7	24	2.3	15	
Tennessee	41	3.4	39	4.8	125	7.4	93	6.3	64	8.7
Utah	7		11		19		16		15	
Vermont	¶		5		¶		7		7	
Virginia	22	1.4	17		63	2.8	73	4.1	21	2.5
Washington	29	2.3	49	5.6	90	5.1	80	5.1	41	5.9
Wisconsin	47	4.4	34	4.1	73	4.7	60	4.3	56	7.8

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.


¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Firearm–Related Indicators

- 5a. Firearm-Related Fatalities (Overall), 2004
- 5b. Firearm-Related Fatalities by Sex, 2004
- 5c. Firearm-Related Fatalities by Age, 2004
- 5d. Firearm-Related Hospitalizations (Overall), 2004
- 5e. Firearm-Related Hospitalizations by Sex, 2004
- 5f. Firearm-Related Hospitalizations by Age, 2004
- 5g. Percentage of Adults Living in Homes with Loaded and Unlocked Firearms, 2004, Behavioral Risk Factor Surveillance System

Figure 5a
Firearm-Related Fatalities (Overall), 2004




‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 5b
Firearm-Related Fatalities by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
‡ Case counts are suppressed if fewer than 5 cases were reported.

Figure 5c
Firearm-Related Fatalities by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	12		197	23.7	327	19.0	190	15.6	143	18.9
Arkansas	5		51	12.9	147	19.7	131	19.5	71	18.6
California	46	0.6	986	19.1	1,179	10.9	658	8.1	444	11.6
Colorado	5		95	14.1	190	13.6	177	15.6	74	15.4
Connecticut	¶		39	8.5	61	6.3	53	5.8	18	
Florida	18		301	13.4	648	13.9	545	12.8	368	12.6
Georgia	17		198	15.5	425	15.4	279	14.0	138	16.3
Hawaii	¶		¶		14		17		6	
Kansas	7		58	13.9	104	13.9	79	12.1	43	12.1
Kentucky	¶		88	15.2	189	15.9	163	15.6	75	14.4
Louisiana	8		266	37.7	364	29.4	178	16.5	80	15.2
Maine	¶		23	12.7	26	7.4	35	9.4	23	12.1
Maryland	7		202	26.5	269	16.8	113	8.1	62	9.8
Massachusetts	¶		75	8.8	62	3.2	35	2.2	31	3.6
Michigan	16		220	15.3	452	16.0	249	9.9	108	8.7
Minnesota	¶		86	11.4	138	9.4	90	7.2	44	7.2
Nebraska	¶		31	11.6	34	7.2	27	6.5	26	11.2
Nevada	¶		58	16.8	129	18.1	115	20.1	72	26.9
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	¶		69	23.5	95	18.9	71	15.0	41	17.9
New York	6		257	9.9	368	6.5	216	4.6	89	3.6
North Carolina	16		202	17.2	437	17.3	254	12.5	140	13.6
Ohio	8		201	12.4	359	11.4	280	9.7	156	10.2
Oklahoma	6		80	15.0	156	16.3	128	15.0	84	18.1
Oregon	¶		56	11.2	105	10.0	126	14.8	89	19.5
Pennsylvania	15		282	16.0	487	14.8	318	10.2	161	8.5
Rhode Island	¶		14		7		13		¶	
South Carolina	10		102	16.8	217	18.4	152	14.5	79	15.2
Tennessee	10		155	19.2	332	19.6	244	16.6	126	17.2
Utah	¶		43	10.1	79	10.9	67	14.6	30	14.4
Vermont	¶		¶		22	13.4	23	13.0	12	
Virginia	16		165	16.3	279	12.3	208	11.7	127	15.0
Washington	6		99	11.2	187	10.5	165	10.6	110	15.8
Wisconsin	¶		101	12.3	138	8.9	116	8.4	50	6.9
United States	358	0.6	6,540	15.7	11,007	13.1	7,447	10.5	4,190	11.5

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 5d
Firearm-Related Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 5e
Firearm-Related Hospitalizations by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 5f
Firearm-Related Hospitalizations by Age#, 2004**

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	19		356	42.9	355	20.6	60	4.9	19	
Arkansas	12		100	25.2	178	23.9	58	8.7	15	
California	115	1.4	2,194	42.5	1,565	14.5	277	3.4	54	1.4
Colorado	8		131	19.5	109	7.8	34	3.0	¶	
Connecticut	¶		102	22.3	80	8.2	14		¶	
Florida	17		544	24.3	592	12.7	183	4.3	52	1.8
Georgia	23	1.2	344	26.9	415	15.1	125	6.3	33	3.9
Hawaii	¶		8		14		6		¶	
Kansas	7		74	17.7	76	10.2	14		5	
Kentucky	6		100	17.3	154	12.9	52	5.0	8	
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	¶		9		14		5		¶	
Maryland	13		440	57.7	341	21.2	54	3.9	14	
Massachusetts	9		170	20.0	108	5.7	15		¶	
Michigan	31	1.5	445	30.9	554	19.6	137	5.4	14	
Minnesota	26	2.6	118	15.7	95	6.5	30	2.4	8	
Nebraska	8		33	12.4	13		¶		¶	
Nevada	9		86	25.0	91	12.8	25	4.4	5	
New Hampshire	¶		6		9		¶		¶	
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	44	1.2	781	30.0	721	12.8	103	2.2	12	
North Carolina	25	1.4	344	27.9	399	16.0	107	5.2	26	2.6
Ohio	16		350	21.7	372	11.8	95	3.3	12	
Oklahoma	18		122	22.9	145	15.1	54	6.3	¶	
Oregon	¶		57	11.4	68	6.5	24	2.8	9	
Pennsylvania	‡	‡	670	38.1	608	18.5	126	4.0	‡	‡
Rhode Island	¶		18		16		¶		¶	
South Carolina	19		181	29.9	181	15.3	65	6.2	15	
Tennessee	13		239	29.6	380	22.4	113	7.7	17	
Utah	¶		33	7.8	55	7.6	12		¶	
Vermont	¶		5		¶		7		¶	
Virginia	16		257	25.3	212	9.4	49	2.8	22	2.6
Washington	7		140	15.9	145	8.2	58	3.7	17	
Wisconsin	20	1.9	138	16.8	98	6.3	45	3.3	8	

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 5g
Percentage of Adults Reporting Living in a Home with a Loaded and Unlocked Firearm, 2004,
Behavioral Risk Factor Surveillance System




† No data available.

Homicide Indicators

- 6a. Homicide (Overall), 2004
- 6b. Homicide by Sex, 2004
- 6c. Homicide by Age, 2004


Figure 6a
Homicide (Overall), 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 6b
Homicide by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 6c
Homicide by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	23	1.8	131	15.8	223	12.9	76	6.2	27	3.6
Arkansas	13		39	9.8	87	11.7	52	7.8	19	
California	98	1.2	939	18.2	995	9.2	365	4.5	87	2.3
Colorado	21	2.2	57	8.5	82	5.9	47	4.2	13	
Connecticut	7		36	7.9	45	4.6	12		¶	
Florida	64	1.9	235	10.5	438	9.4	224	5.3	62	2.1
Georgia	53	2.7	148	11.6	290	10.5	123	6.2	43	5.1
Hawaii	¶		¶		11		12		¶	
Kansas	11		30	7.2	49	6.6	18		10	
Kentucky	10		52	9.0	91	7.7	38	3.6	14	
Louisiana	17		221	31.3	266	21.5	83	7.7	16	
Maine	¶		¶		9		5		¶	
Maryland	25	2.2	203	26.6	240	14.9	52	3.7	15	
Massachusetts	10		80	9.4	60	3.1	20	1.3	5	
Michigan	40	1.9	161	11.2	337	12.0	105	4.2	29	2.3
Minnesota	13		42	5.6	49	3.3	21	1.7	5	
Nebraska	6		11		12		6		¶	
Nevada	10		50	14.5	81	11.4	28	4.9	11	
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	16		40	13.6	71	14.2	33	7.0	15	
New York	41	1.1	285	10.9	361	6.4	148	3.1	35	1.4
North Carolina	37	2.1	141	12.0	280	11.1	117	5.8	33	3.2
Ohio	33	1.4	150	9.3	226	7.2	101	3.5	34	2.2
Oklahoma	17		49	9.2	92	9.6	48	5.6	13	
Oregon	10		25	5.0	39	3.7	23	2.7	10	
Pennsylvania	41	1.8	214	12.2	303	9.2	101	3.2	19	
Rhode Island	¶		10		7		6		¶	
South Carolina	23	2.7	71	11.7	148	12.6	54	5.2	17	
Tennessee	19		103	12.7	188	11.1	79	5.4	26	3.5
Utah	5		10		16		6		5	
Vermont	¶		¶		¶		¶		¶	
Virginia	26	1.7	124	12.2	165	7.3	63	3.5	22	2.6
Washington	25	2.0	53	6.0	88	5.0	38	2.4	12	
Wisconsin	14		48	5.9	59	3.8	24	1.7	¶	
United States	1,031	1.7	5,085	12.2	7,479	8.9	2,887	4.1	839	2.3

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Suicide Indicators

7a. Suicide (Overall), 2004

7b. Suicide by Sex, 2004


7c. Suicide by Age, 2004

7d. Suicide Attempt Hospitalizations (Overall), 2004

7e. Suicide Attempt Hospitalizations by Sex, 2004

7f. Suicide Attempt Hospitalizations by Age, 2004


Figure 7a
Suicide (Overall), 2004



† No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 7b
Suicide by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 7c
Suicide by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	¶		137	16.5	298	17.3	263	21.5	154	20.4
Arkansas	¶		38	9.6	120	16.1	142	21.2	57	15.0
California	25	0.3	423	8.2	1,134	10.5	1,148	14.2	632	16.5
Colorado	11		101	15.0	299	21.4	292	25.8	89	18.5
Connecticut	5		34	7.4	93	9.5	113	12.5	43	9.1
Florida	16		207	9.2	754	16.1	887	20.9	516	17.6
Georgia	9		131	10.2	364	13.2	313	15.7	149	17.6
Hawaii	¶		14		39	11.4	41	12.6	18	
Kansas	¶		56	13.4	136	18.2	118	18.1	55	15.5
Kentucky	¶		69	11.9	201	16.9	176	16.9	78	15.0
Louisiana	6		85	12.1	209	16.9	159	14.7	74	14.0
Maine	¶		28	15.5	50	14.3	63	17.0	28	14.8
Maryland	6		57	7.5	170	10.6	177	12.6	81	12.8
Massachusetts	¶		51	6.0	165	8.6	141	8.9	69	8.1
Michigan	11		138	9.6	430	15.2	381	15.1	136	10.9
Minnesota	8		84	11.1	209	14.2	165	13.3	56	9.1
Nebraska	¶		33	12.4	54	11.4	50	12.0	26	11.2
Nevada	¶		38	11.0	127	17.8	178	31.1	86	32.2
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	7		76	25.9	129	25.7	97	20.5	49	21.4
New York	12		174	6.7	425	7.6	423	9.0	208	8.3
North Carolina	9		128	10.9	427	16.9	299	14.7	158	15.3
Ohio	19		176	10.9	450	14.3	440	15.3	195	12.8
Oklahoma	¶		74	13.9	186	19.4	151	17.6	88	18.9
Oregon	¶		66	13.2	167	15.9	205	24.1	109	23.9
Pennsylvania	14		161	9.1	499	15.2	513	16.4	208	11.0
Rhode Island	¶		11		27	8.8	33	12.3	8	
South Carolina	¶		63	10.4	158	13.4	170	16.3	79	15.2
Tennessee	¶		99	12.3	292	17.2	281	19.2	116	15.8
Utah	5		62	14.6	155	21.4	117	25.6	37	17.8
Vermont	¶		7		31	18.8	35	19.7	17	
Virginia	5		92	9.1	291	12.8	276	15.5	140	16.5
Washington	¶		102	11.6	301	16.9	278	17.9	139	20.0
Wisconsin	8		101	12.3	253	16.4	213	15.4	81	11.3
United States	285	0.5	4,316	10.4	11,712	13.9	10,917	15.4	5,198	14.3

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 7d
Suicide Attempt Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 7e
Suicide Attempt Hospitalizations by Sex, 2004**



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 7f
Suicide Attempt Hospitalizations by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	80	6.2	691	83.3	1,428	82.8	586	48.0	118	15.6
Arkansas	67	12.0	444	112.0	993	133.3	409	61.0	73	19.2
California	348	4.3	2,821	54.6	4,561	42.3	2,771	34.3	641	16.8
Colorado	58	6.0	507	75.5	894	64.0	477	42.2	85	17.7
Connecticut	36	5.2	290	63.4	515	52.9	278	30.7	60	12.7
Florida	218	6.6	1,702	76.0	3,511	75.2	2,152	50.7	554	18.9
Georgia	79	4.0	752	58.8	1,633	59.3	816	41.0	101	11.9
Hawaii	14		109	62.2	157	46.0	123	37.9	24	14.0
Kansas	37	6.6	402	96.0	614	82.3	255	39.1	37	10.4
Kentucky	43	5.3	467	80.6	1,012	85.1	447	42.8	49	9.4
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	16		197	109.2	350	99.9	190	51.3	25	13.2
Maryland	19		375	49.2	801	49.9	439	31.3	68	10.7
Massachusetts	94	7.8	638	74.9	1,386	72.5	732	46.1	115	13.5
Michigan	148	7.1	1,250	86.7	2,345	83.2	1,206	47.9	177	14.2
Minnesota	107	10.5	760	100.8	1,061	72.1	460	37.0	54	8.8
Nebraska	18		137	51.4	210	44.3	125	30.0	16	
Nevada	32	6.3	186	54.0	364	51.1	207	36.1	42	15.7
New Hampshire	20	8.0	152	85.2	329	89.9	143	41.4	23	14.5
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	241	6.4	1,846	70.8	3,070	54.6	1,682	35.7	372	14.9
North Carolina	96	5.5	721	58.4	1,844	73.8	864	41.9	112	11.0
Ohio	81	3.5	1,039	64.3	2,254	71.6	1,100	38.2	139	9.1
Oklahoma	62	8.7	499	93.6	972	101.5	422	49.3	42	9.0
Oregon	52	7.1	438	87.9	887	84.6	513	60.2	69	15.2
Pennsylvania	131	5.6	1,514	86.0	3,106	94.6	1,636	52.3	256	13.5
Rhode Island	10		95	60.9	180	58.9	97	36.1	15	
South Carolina	70	8.3	299	49.4	834	70.7	421	40.3	45	8.6
Tennessee	64	5.4	682	84.4	1,798	106.2	829	56.5	92	12.5
Utah	56	8.6	561	132.0	785	108.2	351	76.7	37	17.8
Vermont	9		80	89.3	131	79.6	80	45.1	14	
Virginia	100	6.4	686	67.6	1,428	63.0	665	37.4	113	13.3
Washington	100	8.0	830	94.0	1,552	87.4	783	50.3	98	14.1
Wisconsin	289	26.9	1,538	187.6	2,085	135.1	839	60.7	112	15.6

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.


¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Motor Vehicle Indicators

- 8a. Unintentional Motor Vehicle Traffic Fatalities (Overall), 2004
- 8b. Unintentional Motor Vehicle Traffic Fatalities by Sex, 2004
- 8c. Unintentional Motor Vehicle Traffic Fatalities by Age, 2004
- 8d. Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations (Overall), 2004
- 8e. Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations by Sex, 2004
- 8f. Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations by Age, 2004
- 8g. Of the Adults Reporting Drinking at Least One Alcoholic Beverage in the Past Month, the Percentage Reporting Driving after Perhaps Having Too Much to Drink in the Past Month, 2004, Behavioral Risk Factor Surveillance System
- 8h. Alcohol-Related Crash Deaths, 2004, Fatality Analysis Reporting System


Figure 8a
Unintentional Motor Vehicle Traffic Fatalities (Overall), 2004



‡ No data available.
 § Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
 ¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 8b
Unintentional Motor Vehicle Traffic Fatalities by Sex, 2004



‡ No data available.
 § Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
 ¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 8c
Unintentional Motor Vehicle Traffic Fatalities by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	49	3.8	247	29.8	299	17.3	233	19.1	180	23.8
Arkansas	44	7.9	192	48.5	253	34.0	160	23.9	106	27.8
California	217	2.7	924	17.9	1,199	11.1	858	10.6	622	16.3
Colorado	30	3.1	181	26.9	208	14.9	162	14.3	94	19.6
Connecticut	11		83	18.1	96	9.9	71	7.8	58	12.2
Florida	147	4.4	720	32.2	931	19.9	797	18.8	603	20.6
Georgia	77	3.9	319	24.9	449	16.3	352	17.7	230	27.2
Hawaii	¶		31	17.7	55	16.1	22	6.8	22	12.8
Kansas	29	5.1	116	27.7	126	16.9	118	18.1	87	24.5
Kentucky	35	4.3	232	40.0	309	26.0	179	17.1	109	21.0
Louisiana	60	6.2	249	35.3	322	26.0	232	21.5	107	20.3
Maine	8		45	24.9	49	14.0	29	7.8	34	17.9
Maryland	38	3.3	172	22.6	183	11.4	152	10.8	107	16.9
Massachusetts	19		128	15.0	146	7.6	119	7.5	96	11.2
Michigan	61	2.9	308	21.4	340	12.1	283	11.2	233	18.7
Minnesota	35	3.4	147	19.5	154	10.5	148	11.9	118	19.2
Nebraska	11		82	30.8	61	12.9	62	14.9	53	22.9
Nevada	19		72	20.9	126	17.7	103	18.0	62	23.2
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	31	7.7	114	38.8	145	28.9	97	20.5	57	24.8
New York	82	2.2	333	12.8	397	7.1	304	6.5	348	14.0
North Carolina	88	5.0	422	35.9	527	20.8	335	16.5	263	25.5
Ohio	50	2.2	336	20.8	325	10.3	269	9.3	197	12.9
Oklahoma	36	5.1	196	36.8	217	22.6	160	18.7	123	26.5
Oregon	29	4.0	109	21.9	121	11.5	117	13.7	86	18.9
Pennsylvania	54	2.3	354	20.1	413	12.6	349	11.2	324	17.1
Rhode Island	¶		24	15.4	20	6.5	15		16	
South Carolina	28	3.3	244	40.3	356	30.2	241	23.1	117	22.5
Tennessee	53	4.4	326	40.3	416	24.6	356	24.3	193	26.3
Utah	20	3.1	69	16.2	99	13.6	61	13.3	49	23.6
Vermont	¶		22	24.6	21	12.8	20	11.3	15	
Virginia	29	1.9	230	22.7	276	12.2	184	10.3	166	19.6
Washington	24	1.9	184	20.8	180	10.1	156	10.0	103	14.8
Wisconsin	26	2.4	207	25.2	208	13.5	157	11.4	153	21.3
United States	2,165	3.6	10,757	25.8	13,285	15.8	10,024	14.2	7,175	19.7

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 8d
Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

|| Rates are suppressed if fewer than 20 cases were reported.

§ Rate per 100,000 population.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 8e
Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations by Sex, 2004



‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 8f
Unintentional Motor Vehicle Traffic and Non-Traffic Hospitalizations by Age#, 2004

State	0-4		5-14		15-24		25-34		35-44		45-54		55-64		65-74		75-84		85+		
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	
Arizona	209	48.5	666	76.6	2,041	246.0	1,290	152.3	1,136	129.5	906	126.3	556	110.3	384	93.2	331	124.1	114	147.2	
Arkansas	31	16.7	162	43.3	784	197.8	516	143.3	532	138.2	444	118.1	318	108.1	245	119.4	186	145.3	63	131.7	
California	650	24.7	2,402	44.6	7,975	154.3	5,257	100.1	4,942	89.2	4,291	88.3	2,400	74.4	1,577	80.9	1,499	110.3	597	116.1	
Colorado	55	16.4	252	39.7	1,129	168.1	755	112.5	707	97.6	703	101.5	390	89.0	231	88.3	248	151.8	91	165.3	
Connecticut	30	14.1	162	33.8	719	157.2	419	102.3	394	69.8	319	59.9	195	52.1	132	60.9	141	80.6	72	87.7	
Florida	282	25.8	1,174	52.8	3,818	170.6	2,546	118.9	2,457	97.1	2,255	94.7	1,378	73.9	1,069	72.5	979	91.2	372	98.0	
Georgia	186	27.4	612	48.0	2,056	160.6	1,403	102.9	1,248	89.6	1,140	95.9	685	85.3	474	99.5	391	142.1	94	98.7	
Hawaii	22	24.8	87	54.1	355	202.4	183	113.5	199	110.4	172	92.4	108	78.1	72	91.0	58	85.9	35	138.1	
Kansas	32	8.6	125	64.9	438	104.6	240	66.9	223	57.5	233	59.5	165	63.4	85	50.0	77	59.5	41	74.5	
Kentucky	59	22.1	238	43.5	1,030	177.7	702	123.6	667	107.3	555	91.7	393	89.5	262	93.1	228	127.4	71	120.3	
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	12		68	42.9	365	202.3	185	127.0	185	90.4	143	66.3	88	56.9	84	88.0	69	99.6	28	111.6	
Maryland	37	9.9	258	33.1	1,703	223.3	1,004	141.2	957	107.0	773	93.3	470	82.0	281	86.2	229	101.3	93	112.4	
Massachusetts	26	6.6	253	31.0	1,289	151.4	704	80.0	778	75.5	615	65.6	405	62.1	296	74.7	349	108.3	132	97.0	
Michigan	161	24.8	668	46.4	2,070	143.6	1,359	104.1	1,462	96.5	1,268	84.8	764	74.9	462	75.5	524	114.0	165	94.2	
Minnesota	45	13.6	216	31.5	988	131.0	572	84.3	609	76.8	549	72.7	342	70.1	203	67.6	238	109.8	93	94.7	
Nebraska	15		55	23.2	210	78.9	113	49.3	130	53.2	120	48.1	83	49.6	67	60.2	74	87.5	29	80.8	
Nevada	25	14.6	141	42.1	401	116.4	288	82.0	295	81.6	265	81.0	159	64.6	92	58.8	65	73.1	28	126.9	
New Hampshire	10		53	30.2	241	135.2	146	96.7	174	80.9	146	70.2	91	65.6	59	71.6	56	102.3	31	146.1	
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	168	13.5	1,306	51.3	3,482	133.6	2,270	86.2	2,402	80.3	1,969	71.8	1,356	68.8	1,091	89.0	1,108	121.4	389	109.9	
North Carolina	128	22.1	481	41.7	1,827	148.1	1,333	111.0	1,311	101.1	1,062	89.1	657	75.6	432	79.1	443	126.9	130	106.7	
Ohio	69	9.5	305	19.5	1,618	100.1	1,095	74.8	1,128	67.0	981	57.6	564	47.9	335	44.9	398	69.9	126	60.5	
Oklahoma	97	40.0	332	70.6	964	180.9	530	112.8	519	106.3	440	89.2	304	83.8	210	84.4	184	114.0	74	136.3	
Oregon	56	24.5	164	32.7	658	132.0	400	80.8	393	71.0	391	73.4	222	69.6	153	67.1	131	78.2	59	99.0	
Pennsylvania	203	27.8	1,002	62.3	3,523	200.2	2,149	145.6	2,179	120.7	1,850	102.6	1,217	91.8	929	97.5	1,027	149.0	368	144.6	
Rhode Island	6		28	20.2	83	53.2	62	44.6	62	37.3	47	29.6	30	27.3	27	41.8	26	43.5	18		
South Carolina	60	21.4	359	63.3	1,040	171.7	767	135.0	737	120.6	631	106.2	394	87.4	216	75.6	203	115.8	43	72.3	
Tennessee	99	25.4	434	53.9	1,815	224.6	1,253	154.3	1,241	140.8	991	117.1	604	97.5	391	98.7	353	142.6	116	129.0	
Utah	46	18.8	162	39.6	612	144.0	334	79.9	346	112.5	269	96.4	178	99.7	101	92.2	91	125.9	37	141.6	
Vermont	7		29	38.4	157	175.3	68	97.7	89	93.6	73	70.3	46	62.5	31	75.9	20	70.3	13		
Virginia	52	10.4	176	16.7	903	89.0	597	50.9	594	54.3	500	50.0	351	45.1	215	47.4	186	63.9	45	43.9	
Washington	53	13.2	277	32.4	1,142	129.4	694	83.2	700	74.3	712	76.6	403	64.4	247	70.3	270	109.7	95	96.0	
Wisconsin	51	15.0	186	25.4	807	98.4	514	73.5	660	78.2	495	59.6	369	66.8	267	76.1	296	115.2	119	106.6	

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.


Age in years.

Figure 8g
**Of the Adults Reporting Drinking at Least One Alcoholic Beverage in the Past Month,
 the Percentage Reporting Driving after Perhaps Having Too Much to Drink in the Past Month, 2004,
 Behavioral Risk Factor Surveillance System**



‡ No data available.

Figure 8h
**Alcohol-Related Crash Deaths, 2004,
 Fatality Analysis Reporting System**




§ Rate per 100,000 population.

Poisoning Indicators

- 9a. Poisoning Fatalities (Overall), 2004
- 9b. Poisoning Fatalities by Sex, 2004
- 9c. Poisoning Fatalities by Age, 2004
- 9d. Poisoning Hospitalizations (Overall), 2004
- 9e. Poisoning Hospitalizations by Sex, 2004
- 9f. Poisoning Hospitalizations by Age, 2004

Figure 9a
Poisoning Fatalities (Overall), 2004




‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 9b
Poisoning Fatalities by Sex, 2004



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

Figure 9c
Poisoning Fatalities by Age#, 2004

State	0-14		15-24		25-44		45-64		65+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	5		77	9.3	304	17.6	316	25.9	42	5.6
Arkansas	¶		31	7.8	116	15.6	79	11.8	17	
California	12		211	4.1	1,407	13.0	1,564	19.3	168	4.4
Colorado	¶		49	7.3	271	19.4	256	22.6	31	6.5
Connecticut	¶		43	9.4	195	20.0	102	11.3	24	5.1
Florida	12		251	11.2	1,072	22.9	937	22.1	155	5.3
Georgia	6		54	4.2	370	13.4	280	14.1	28	3.3
Hawaii	¶		6		42	12.3	58	17.9	6	
Kansas	¶		20	4.8	124	16.6	96	14.7	15	
Kentucky	8		63	10.9	275	23.1	157	15.0	22	4.2
Louisiana	¶		84	11.9	290	23.4	155	14.4	22	4.2
Maine	¶		23	12.7	78	22.3	37	10.0	8	
Maryland	¶		43	5.6	367	22.9	264	18.8	21	3.3
Massachusetts	¶		83	9.7	390	20.4	237	14.9	21	2.5
Michigan	¶		85	5.9	474	16.8	373	14.8	51	4.1
Minnesota	¶		18		147	10.0	139	11.2	13	
Nebraska	¶		8		45	9.5	33	7.9	¶	
Nevada	¶		39	11.3	179	25.1	169	29.5	12	
New Hampshire	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New Mexico	¶		27	9.2	162	32.3	147	31.0	10	
New York	9		54	2.1	255	4.5	218	4.6	63	2.5
North Carolina	¶		105	8.9	457	18.1	321	15.8	49	4.7
Ohio	9		129	8.0	554	17.6	494	17.2	72	4.7
Oklahoma	¶		62	11.6	245	25.6	191	22.3	25	5.4
Oregon	¶		28	5.6	159	15.2	171	20.1	29	6.4
Pennsylvania	5		218	12.4	799	24.3	548	17.5	84	4.4
Rhode Island	¶		10		49	16.0	41	15.3	¶	
South Carolina	¶		29	4.8	165	14.0	161	15.4	17	
Tennessee	¶		70	8.7	379	22.4	307	20.9	40	5.5
Utah	¶		53	12.5	207	28.5	132	28.9	10	
Vermont	¶		6		25	15.2	21	11.8	¶	
Virginia	¶		63	6.2	310	13.7	198	11.1	28	3.3
Washington	¶		58	6.6	366	20.6	398	25.6	40	5.7
Wisconsin	¶		46	5.6	225	14.6	199	14.4	33	4.6
United States	174	0.3	2,969	7.1	14,190	16.9	11,444	16.2	1,521	4.2

‡ No data available.


§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

Figure 9d
Poisoning Hospitalizations (Overall), 2004



* Percentage of hospital discharge data injury hospitalizations with external cause coding. Incompleteness may lead to bias.


‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 9e
Poisoning Hospitalizations by Sex, 2004**



‡ No data available.
§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.
¶ Case counts are suppressed if fewer than 5 cases were reported.

**Figure 9f
Poisoning Hospitalizations by Age#, 2004**

State	0-4		5-14		15-24		25-34		35-44		45-54		55-64		65-74		75-84		85+	
	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]	N	Rate [§]
Arizona	152	35.3	139	16.0	889	107.1	919	108.5	946	107.9	819	114.2	392	77.7	209	50.7	167	62.6	67	86.5
Arkansas	50	26.9	63	16.8	302	76.2	303	84.2	318	82.6	260	69.2	146	49.6	85	41.4	61	47.6	29	60.6
California	395	15.0	498	9.2	3,567	69.0	2,742	52.2	3,709	66.9	3,582	73.7	1,750	54.3	843	43.2	691	50.9	235	45.7
Colorado	75	22.3	78	12.3	640	95.3	603	89.8	659	91.0	599	86.5	239	54.5	131	50.1	120	73.4	48	87.2
Connecticut	74	34.7	61	12.7	377	82.4	344	84.0	462	81.8	370	69.5	164	43.8	93	42.9	91	52.0	56	68.2
Florida	384	35.2	338	15.2	2,479	110.7	2,262	105.6	2,941	116.3	2,623	110.2	1,160	62.2	634	43.0	591	55.1	224	59.0
Georgia	164	24.2	127	10.0	1,005	78.5	1,003	73.6	1,396	100.3	1,123	94.4	533	66.4	301	63.2	161	58.5	61	64.0
Hawaii	26	29.3	16		130	74.1	103	63.9	100	55.5	151	81.1	54	39.0	35	44.2	27	40.0	15	
Kansas	93	25.0	56	29.1	462	110.4	362	101.0	419	108.1	346	88.3	110	42.3	75	44.1	57	44.1	31	56.3
Kentucky	51	19.1	60	11.0	627	108.2	723	127.3	770	123.9	586	96.9	210	47.8	112	39.8	91	50.8	28	47.4
Louisiana	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
Maine	28	41.4	28	17.7	268	148.6	206	141.4	244	119.2	215	99.6	84	54.3	35	36.7	43	62.1	13	
Maryland	93	24.8	46	5.9	570	74.8	568	79.9	945	105.6	784	94.6	358	62.4	164	50.3	154	68.1	62	74.9
Massachusetts	75	19.0	117	14.4	868	101.9	858	97.5	1,163	112.9	948	101.2	380	58.3	195	49.2	201	62.4	105	77.1
Michigan	284	43.7	227	15.8	1,642	113.9	1,479	113.3	1,956	129.2	1,693	113.2	611	59.9	328	53.6	289	62.9	122	69.7
Minnesota	95	28.6	149	21.7	957	126.9	688	101.4	787	99.2	638	84.5	229	46.9	115	38.3	106	48.9	37	37.7
Nebraska	15		27	11.4	158	59.3	122	53.2	130	53.2	137	55.0	59	35.3	26	23.3	24	28.4	15	
Nevada	29	16.9	37	11.0	233	67.6	240	68.4	293	81.1	247	75.5	119	48.4	56	35.8	30	33.8	15	
New Hampshire	16		28	15.9	200	112.2	184	122.1	253	117.4	159	76.8	71	51.4	32	38.7	30	54.6	22	103.1
New Mexico	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡	‡
New York	533	42.8	431	16.9	2,672	102.5	2,301	87.4	3,656	122.2	2,967	108.3	1,237	62.8	618	50.4	553	60.6	244	68.9
North Carolina	143	24.7	128	11.1	928	75.2	1,129	94.0	1,557	120.1	1,181	99.1	531	61.1	334	61.2	239	68.5	68	55.8
Ohio	64	8.8	128	8.2	1,337	82.7	1,368	93.5	1,625	96.5	1,349	79.2	484	41.1	260	34.8	203	35.6	85	40.8
Oklahoma	171	70.6	121	25.7	743	139.4	627	133.5	795	162.8	590	119.7	275	75.8	137	55.1	133	82.4	47	86.5
Oregon	66	28.9	74	14.8	531	106.5	525	106.1	623	112.6	599	112.5	219	68.6	99	43.4	89	53.1	44	73.8
Pennsylvania	293	40.1	258	16.0	2,308	131.2	2,136	144.7	3,021	167.3	2,490	138.1	901	68.0	458	48.1	433	62.8	186	73.1
Rhode Island	14		14		109	69.9	90	64.7	129	77.6	106	66.8	48	43.7	17		21	35.1	13	
South Carolina	106	37.8	106	18.7	392	64.7	517	91.0	604	98.8	536	90.2	226	50.2	150	52.5	104	59.3	37	62.2
Tennessee	70	17.9	110	13.7	942	116.6	1,177	144.9	1,500	170.2	1,221	144.3	518	83.6	260	65.7	173	69.9	58	64.5
Utah	63	25.8	53	13.0	555	130.6	482	115.3	451	146.6	380	136.2	214	119.9	92	84.0	78	107.9	28	107.2
Vermont	7		9		105	117.2	66	94.9	91	95.7	78	75.1	32	43.4	23	56.3	21	73.8	5	
Virginia	67	13.4	122	11.6	862	84.9	837	71.3	972	88.9	790	79.0	302	38.8	142	31.3	92	31.6	43	41.9
Washington	72	17.9	125	14.6	868	98.3	820	98.4	1,026	108.9	1,046	112.5	482	77.0	252	71.7	242	98.3	105	106.1
Wisconsin	166	48.8	246	33.6	1,341	163.6	968	138.4	1,262	149.5	918	110.5	382	69.1	251	71.6	233	90.6	104	93.2

‡ No data available.

§ Rate per 100,000 population.

|| Rates are suppressed if fewer than 20 cases were reported.

¶ Case counts are suppressed if fewer than 5 cases were reported.

Age in years.

U.S. Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Injury Prevention and Control
www.cdc.gov/injury