



ENSO Cycle: Recent Evolution, Current Status and Predictions

**Update prepared by
Climate Prediction Center / NCEP
12 January 2009**



Outline

- Overview
- Recent Evolution and Current Conditions
- Oceanic Niño Index (ONI) – **“Revised December 2008”**
- Pacific SST Outlook
- U.S. Seasonal Precipitation and Temperature Outlooks
- Summary




Summary

- **Atmospheric and oceanic conditions reflect La Niña.**
- **Recently, negative equatorial SST anomalies have strengthened across portions of the central and eastern Pacific Ocean.**
- **Based on recent trends in the observations and model forecasts, La Niña conditions are likely through early 2009.**



Recent Evolution of Equatorial Pacific SST Departures (°C)



Since October 2008, negative sea surface temperature anomalies have strengthened in portions of the central and eastern equatorial Pacific Ocean.





Niño Region SST Departures (°C)

Recent Evolution

The latest weekly SST departures are:


- Niño 4 -0.8°C
- Niño 3.4 -1.1°C
- Niño 3 -0.9°C
- Niño 1+2 -0.6°C





SST Departures (°C) in the Tropical Pacific During the Last 4 Weeks

During the last 4-weeks, SSTs were at least 0.5°C below-average throughout the central and eastern equatorial Pacific Ocean, with anomalies more than 1°C below-average between 170°W and 110°W.






Global SST Departures (°C)

Average SST Anomalies

14 DEC 2008 – 10 JAN 2009








During the last four weeks, equatorial SSTs were below-average across the central and eastern Pacific Ocean, and above-average in the extreme western Pacific and Atlantic Oceans. Positive anomalies covered much of the North Atlantic and west-central South Pacific Oceans, while negative anomalies were evident in a region extending from the west coast of North America to the Gulf of Alaska and Bering Sea.




Weekly SST Departures (°C) for the Last Four Weeks

- During the last four weeks, negative SST anomalies strengthened and became more widespread in the central and eastern equatorial Pacific.

Weekly SST Anomalies (DEG C)




Change in Weekly SST Anoms (°C)
07JAN2009 minus 10DEC2008





Upper-Ocean Conditions in the Eq. Pacific

Cold Episodes ↓
Warm Episodes ↑




- The basin-wide equatorial upper ocean (0-300 m) heat content is **greatest** prior to and during the early stages of a Pacific **warm** (El Niño) episode (compare top 2 panels) and **least** prior to and during the early stages of a **cold** (La Niña) episode.
- The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.
- Current values of the upper-ocean heat anomalies (negative) and the thermocline slope index (positive) indicate La Niña.

The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).



Central & Eastern Pacific Upper-Ocean (0-300 m) Weekly Heat Content Anomalies


EQ. Upper-Ocean Heat Anoms. (deg C) for 180-100W



The upper ocean heat content was below-average across the eastern half of the equatorial Pacific Ocean between January 2007 and April 2008 and again since mid-August 2008. The negative heat content anomalies have strengthened since mid-October 2008.




Sub-Surface Temperature Departures (°C) in the Equatorial Pacific



- During mid-November 2008 through early January 2009, negative sub-surface temperature anomalies at thermocline depth (dashed black line) strengthened, while positive anomalies in the western Pacific expanded eastward.

- The most recent period (below) shows negative subsurface temperature anomalies in the central and eastern Pacific, and positive anomalies west of 165°W.



Most recent pentad analysis

Time
↓

Longitude




Tropical OLR and Wind Anomalies During the Last 30 Days

Positive OLR anomalies (suppressed convection and precipitation, red shading) were observed between 140°E and 150°W, while negative anomalies (enhanced convection, blue shading) were present over most of Indonesia, Malaysia, the Philippines, and northern Australia.


Low-level (850-hPa) easterly wind anomalies persisted over the western and central tropical Pacific Ocean.

Upper-level (200-hPa) westerly wind anomalies continued over much of the equatorial Pacific. An anomalous cyclonic couplet is evident in the subtropics consistent with La Niña.


OLR Anomalies
14 DEC 2008 to 08 JAN 2009



CDAS 850-hPa Wind Anoms
10 DEC 2008-08 JAN 2009




CDAS 200-hPa Wind Anoms
10 DEC 2008-08 JAN 2009




Atmospheric Circulation over the North Pacific & North America During the Last 60 Days

200-hPa Wind


10 NOV 2008–24 NOV 2008




25 NOV 2008–09 DEC 2008



10 DEC 2008–24 DEC 2008




25 DEC 2008–08 JAN 2009




500-hPa Height & Anoms.


10 NOV 2008–24 NOV 2008




25 NOV 2008–09 DEC 2008



10 DEC 2008–24 DEC 2008




25 DEC 2008–08 JAN 2009




925-hPa Temp. Anoms. (°C)


10 NOV 2008–24 NOV 2008




25 NOV 2008–09 DEC 2008



10 DEC 2008–24 DEC 2008



25 DEC 2008–08 JAN 2009




From mid-November through early December, the atmospheric circulation featured an anomalous ridge (red oval in central panels) and above-normal temperatures over western North America with a trough and below-average temperatures in the East. These features retrograded by mid-December, with an anomalously strong trough and below-normal temperatures covering Canada and the northern tier of the United States. The trough retrograded to Alaska and western Canada by the end of the year, resulting in much below-average temperatures in that region.




U.S. Temperature and Precipitation Departures During the Last 30 and 90 Days

Last 30 Days

30-day (ending 11 Jan 2009) % of average precipitation




30-day (ending 10 Jan 2009) temperature departures (degree C)




Last 90 Days

90-day (ending 11 Jan 2009) % of average precipitation



90-day (ending 10 Jan 2009) temperature departures (degree C)






Intraseasonal Variability

- **Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.**
- **Related to this activity**
 - **significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.**
 - **Several Kelvin waves have occurred during the last year (see next slide).**



Weekly Heat Content Evolution in the Equatorial Pacific



- During January - July 2008, subsurface temperature anomalies across the central and eastern equatorial Pacific increased consistent with the transition from La Niña to ENSO-neutral conditions.


- The upper-ocean heat content was occasionally affected by weak oceanic Kelvin wave activity during the period (dashed lines).

- From September 2008 – January 2009, negative heat content anomalies strengthened in the central and eastern equatorial Pacific, as La Niña conditions redeveloped.

- Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Down-welling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.



Low-level (850-hPa) Zonal (east-west) Wind Anomalies (m s^{-1})



Westerly wind anomalies
(orange/red shading).

Easterly wind anomalies (blue
shading).


Low-level (850-hPa) easterly wind
anomalies have persisted since
January 2007 over the equatorial
Pacific between 150°E and 150°W.

However, in late September- early
October 2008, intraseasonal (MJO)
activity briefly weakened the easterly
anomalies across the central
equatorial Pacific (dashed red oval in
figure).

Since early October, strong easterly
anomalies have dominated the central
equatorial Pacific.



200-hPa Velocity Potential Anomalies (5°N-5°S)



Positive anomalies (brown shading) indicate unfavorable conditions for precipitation.

Negative anomalies (green shading) indicate favorable conditions for precipitation.


Positive velocity potential anomalies (associated with suppressed convection) have been dominant near the Date Line.

The MJO was active during mid-May through mid-July 2008, and again beginning in September 2008.

Recently, MJO activity has been weak.



Outgoing Longwave Radiation (OLR) Anomalies



Drier-than-average conditions (orange/red shading)

Wetter-than-average conditions (blue shading)

Since February 2007, convection has been suppressed across the central equatorial Pacific Ocean.

Convection has occasionally been enhanced over the western equatorial Pacific and central Indian Ocean. Since November 2008, convection has become more persistent over Indonesia.



Oceanic Niño Index (ONI)

- The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.
- Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST – **ERSST.v3b**). The SST reconstruction methodology is described in Smith et al., 2008, *J. Climate*, vol. 21, 2283-2296.)
- Used to place current events into a historical perspective
- NOAA's operational definitions of El Niño and La Niña are keyed to the ONI index.



NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a *positive* ONI greater than or equal to $+0.5^{\circ}\text{C}$.

La Niña: characterized by a *negative* ONI less than or equal to -0.5°C .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 SST departures meet or exceed $\pm 0.5^{\circ}\text{C}$ along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.



ONI (°C): Evolution since 1950

The most recent ONI value (October–December 2008) is -0.3°C .





Historical El Niño and La Niña Episodes

Based on the ONI computed using ERSST.v3b

	Highest		Lowest
<u>El Niño</u>	<u>ONI Value</u>	<u>La Nina</u>	<u>ONI Value</u>
JAS 1951 - NDJ 1951/52	0.8	ASO 1949 – FMA 1951	-1.7
MAM 1957 – MJJ 1958	1.7	MAM 1954 – DJF 1956/57	-2.1
JJA 1963 – DJF 1963/64	1.0	ASO 1962 – DJF 1962/63	-0.8
MJJ 1965 – MAM 1966	1.6	MAM 1964 – DJF 1964/65	-1.1
OND 1968 – MJJ 1969	1.0	NDJ 1967/68 – MAM 1968	-0.9
ASO 1969 – DJF 1969/70	0.8	JJA 1970 – DJF 1971/72	-1.3
AMJ 1972 – FMA 1973	2.1	AMJ 1973 – MAM 1976	-2.0
ASO 1976 – JFM 1977	0.8	SON 1984 – ASO 1985	-1.0
ASO 1977 - DJF 1977/78	0.8	AMJ 1988 – AMJ 1989	-1.9
AMJ 1982 – MJJ 1983	2.3	ASO 1995 – FMA 1996	-0.7
JAS 1986 – JFM 1988	1.6	JJA 1998 – MJJ 2000	-1.6
AMJ 1991 – JJA 1992	1.8	SON 2000 – JFM 2001	-0.7
AMJ 1994 – FMA 1995	1.3	ASO 2007 – AMJ 2008	-1.4
AMJ 1997 – AMJ 1998	2.5		
AMJ 2002 – FMA 2003	1.5		
MJJ 2004 – JFM 2005	0.9		
JAS 2006 - DJF 2006/07	1.1		

NOTE:

After updating the ocean analysis to ERSST.v3b, a new La Niña episode was classified (ASO 1962-DJF 1962/63) and two previous La Niña episodes were combined into one single episode (AMJ 1973- MAM 1976).



Historical Pacific warm (red) and cold (blue) episodes based on a threshold of +/- 0.5 °C for the Oceanic Nino Index (ONI) [3 month running mean of ERSST.v3b SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)], calculated with respect to the 1971-2000 base period. For historical purposes El Niño and La Niña episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons.

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
1950	-1.7	-1.5	-1.3	-1.4	-1.3	-1.1	-0.8	-0.8	-0.8	-0.9	-0.9	-1.0
1951	-1.0	-0.9	-0.6	-0.3	-0.2	0.2	0.4	0.7	0.7	0.8	0.7	0.6
1952	0.3	0.1	0.1	0.2	0.1	-0.1	-0.3	-0.3	-0.2	-0.2	-0.1	0.0
1953	0.2	0.4	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4
1954	0.5	0.3	-0.1	-0.5	-0.7	-0.7	-0.8	-1.0	-1.2	-1.1	-1.1	-1.1
1955	-1.0	-0.9	-0.9	-1.0	-1.0	-1.0	-1.0	-1.0	-1.4	-1.8	-2.0	-1.9
1956	-1.3	-0.9	-0.7	-0.6	-0.6	-0.6	-0.7	-0.8	-0.8	-0.9	-0.9	-0.8
1957	-0.5	-0.1	0.3	0.6	0.7	0.9	0.9	0.9	0.9	1.0	1.2	1.5
1958	1.7	1.5	1.2	0.8	0.6	0.5	0.3	0.1	0.0	0.0	0.2	0.4
1959	0.4	0.5	0.4	0.2	0.0	-0.2	-0.4	-0.5	-0.4	-0.3	-0.2	-0.2
1960	-0.3	-0.3	-0.3	-0.2	-0.2	-0.2	-0.1	0.0	-0.1	-0.2	-0.2	-0.2
1961	-0.2	-0.2	-0.2	-0.1	0.1	0.2	0.0	-0.3	-0.6	-0.6	-0.5	-0.4
1962	-0.4	-0.4	-0.4	-0.5	-0.4	-0.4	-0.3	-0.3	-0.5	-0.6	-0.7	-0.7
1963	-0.6	-0.3	0.0	0.1	0.1	0.3	0.6	0.8	0.9	0.9	1.0	1.0
1964	0.8	0.4	-0.1	-0.5	-0.8	-0.8	-0.9	-1.0	-1.1	-1.2	-1.2	-1.0
1965	-0.8	-0.4	-0.2	0.0	0.3	0.6	1.0	1.2	1.4	1.5	1.6	1.5
1966	1.2	1.0	0.8	0.5	0.2	0.2	0.2	0.0	-0.2	-0.2	-0.3	-0.3
1967	-0.4	-0.4	-0.6	-0.5	-0.3	0.0	0.0	-0.2	-0.4	-0.5	-0.4	-0.5
1968	-0.7	-0.9	-0.8	-0.7	-0.3	0.0	0.3	0.4	0.3	0.4	0.7	0.9
1969	1.0	1.0	0.9	0.7	0.6	0.5	0.4	0.4	0.6	0.7	0.8	0.7
1970	0.5	0.3	0.2	0.1	0.0	-0.3	-0.6	-0.8	-0.9	-0.8	-0.9	-1.1
1971	-1.3	-1.3	-1.1	-0.9	-0.8	-0.8	-0.8	-0.8	-0.8	-0.9	-1.0	-0.9
1972	-0.7	-0.4	0.0	0.2	0.5	0.8	1.0	1.3	1.5	1.8	2.0	2.1
1973	1.8	1.2	0.5	-0.1	-0.6	-0.9	-1.1	-1.3	-1.4	-1.7	-2.0	-2.1
1974	-1.9	-1.7	-1.3	-1.1	-0.9	-0.8	-0.6	-0.5	-0.5	-0.7	-0.9	-0.7
1975	-0.6	-0.6	-0.7	-0.8	-0.9	-1.1	-1.2	-1.3	-1.5	-1.6	-1.7	-1.7



Historical Pacific warm (red) and cold (blue) episodes based on a threshold of +/- 0.5 °C for the Oceanic Nino Index (ONI) [3 month running mean of ERSST.v3b SST anomalies in the Nino 3.4 region (5N-5S, 120-170W)], calculated with respect to the 1971-2000 base period. For historical purposes El Niño and La Niña episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons.

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
1976	-1.6	-1.2	-0.8	-0.6	-0.5	-0.2	0.1	0.3	0.5	0.7	0.8	0.7
1977	0.6	0.5	0.2	0.2	0.2	0.4	0.4	0.4	0.5	0.6	0.7	0.7
1978	0.7	0.4	0.0	-0.3	-0.4	-0.4	-0.4	-0.4	-0.4	-0.3	-0.2	-0.1
1979	-0.1	0.0	0.1	0.1	0.1	-0.1	0.0	0.1	0.3	0.4	0.5	0.5
1980	0.5	0.3	0.2	0.2	0.3	0.3	0.2	0.0	-0.1	-0.1	0.0	-0.1
1981	-0.3	-0.5	-0.5	-0.4	-0.3	-0.3	-0.4	-0.4	-0.3	-0.2	-0.1	-0.1
1982	0.0	0.1	0.1	0.3	0.6	0.7	0.7	1.0	1.5	1.9	2.2	2.3
1983	2.3	2.0	1.5	1.2	1.0	0.6	0.2	-0.2	-0.6	-0.8	-0.9	-0.7
1984	-0.4	-0.2	-0.2	-0.3	-0.5	-0.4	-0.3	-0.2	-0.3	-0.6	-0.9	-1.1
1985	-0.9	-0.8	-0.7	-0.7	-0.7	-0.6	-0.5	-0.5	-0.5	-0.4	-0.3	-0.4
1986	-0.5	-0.4	-0.2	-0.2	-0.1	0.0	0.3	0.5	0.7	0.9	1.1	1.2
1987	1.2	1.3	1.2	1.1	1.0	1.2	1.4	1.6	1.6	1.5	1.3	1.1
1988	0.7	0.5	0.1	-0.2	-0.7	-1.2	-1.3	-1.2	-1.3	-1.6	-1.9	-1.9
1989	-1.7	-1.5	-1.1	-0.8	-0.6	-0.4	-0.3	-0.3	-0.3	-0.3	-0.2	-0.1
1990	0.1	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.4
1991	0.4	0.3	0.3	0.4	0.6	0.8	1.0	0.9	0.9	1.0	1.4	1.6
1992	1.8	1.6	1.5	1.4	1.2	0.8	0.5	0.2	0.0	-0.1	0.0	0.2
1993	0.3	0.4	0.6	0.7	0.8	0.7	0.4	0.4	0.4	0.4	0.3	0.2
1994	0.2	0.2	0.3	0.4	0.5	0.5	0.6	0.6	0.7	0.9	1.2	1.3
1995	1.2	0.9	0.7	0.4	0.3	0.2	0.0	-0.2	-0.5	-0.6	-0.7	-0.7
1996	-0.7	-0.7	-0.5	-0.3	-0.1	-0.1	0.0	-0.1	-0.1	-0.2	-0.3	-0.4
1997	-0.4	-0.3	0.0	0.4	0.8	1.3	1.7	2.0	2.2	2.4	2.5	2.5
1998	2.3	1.9	1.5	1.0	0.5	0.0	-0.5	-0.8	-1.0	-1.1	-1.3	-1.4
1999	-1.4	-1.2	-0.9	-0.8	-0.8	-0.8	-0.9	-0.9	-1.0	-1.1	-1.3	-1.6
2000	-1.6	-1.4	-1.0	-0.8	-0.6	-0.5	-0.4	-0.4	-0.4	-0.5	-0.6	-0.7
2001	-0.6	-0.5	-0.4	-0.2	-0.1	0.1	0.2	0.2	0.1	0.0	-0.1	-0.1



Pacific Niño 3.4 SST Outlook

Nearly all ENSO forecasts indicate below-average SSTs in the central equatorial Pacific through Northern Hemisphere Summer 2009. Many models suggest La Niña conditions through April 2009.





Figure provided by the International Research Institute (IRI) for Climate and Society (updated 17 December 2008).




SST Outlook: NCEP CFS Forecast Issued 12 January 2009



The CFS ensemble mean (heavy blue line) indicates La Niña conditions through Spring 2009.


PDF correction: Forecast *Nino3.4* SST anomalies from CFS






U. S. Seasonal Outlooks January – March 2009

Temperature



Precipitation



These seasonal outlooks combine long-term trends and some aspects of La Niña.



Summary

- **Atmospheric and oceanic conditions reflect La Niña.**
- **Recently, negative equatorial SST anomalies have strengthened across portions of the central and eastern Pacific Ocean.**
- **Based on recent trends in the observations and model forecasts, La Niña conditions are likely through early 2009.**