

NOBANIS – Invasive Alien Species Fact Sheet

Dreissena polymorpha

Author of this fact sheet: Christina Birnbaum, Estonian University of Life Sciences, Institute of Agricultural and Environmental Sciences, Kreutzwaldi 64 Tartu 51014, phone, e-mail: chbirnbaum@yahoo.com

Bibliographical reference – how to cite this fact sheet:

Birnbaum, C. (2006): NOBANIS – Invasive Alien Species Fact Sheet – *Dreissena polymorpha*. – From: Online Database of the North European and Baltic Network on Invasive Alien Species – NOBANIS www.nobanis.org, Date of access x/x/200x.

Species description

Scientific name: *Dreissena polymorpha* (Pallas, 1771), Dreissenidae.

Synonyms: *Mytilus polymorphus* (Pallas), *Mytilus hagenii*, *Tichogonia chemnitzii* (Rossm.)

Common names: Zebra mussel, Wandering mussel (GB), Zebramuschel, Wandermuschel, Dreikantmuschel, Dreiecksmuschel, Schafklaumuschel (DE), Vandremusling (DK), Tavaline ehk muutlik rändkarp (EE), Vaeltajasimpukka (FI), Svītrainā gliemene (LV), Dreisena (LT), Racicznica zmienna (PL), Vandringsmussla (SE).



Fig. 1. *Dreissena polymorpha*, photo by Viktoras Didziulis.



Fig. 2. *Dreissena polymorpha*, photo by Jonne Kotta, source: [D. polymorpha fact-sheet](#), produced by Estonian Marine Institute.

Species identification

The shell of *D. polymorpha* is triangular (height makes 40-60 % of length) or triangular with sharply pointed shell hinge ends (umbos). The maximum size of *D. polymorpha* can be 3-5 cm (Mackie *et al.* 1989). The prominent dark and light banding pattern on the shell is the most obvious characteristic of the *D. polymorpha*. Its specific name, "polymorpha", derives from the many variations in shell colour, pattern and shape depending on substrate, depth, and density of aggregation. Greenish, brownish-yellowish with clear dark and light coloured ("zebra") zig-zag banding. The outer covering of the shell (the periostracum) is generally well polished, light tan in colour with a distinct series of broad, dark, transverse colour bands which may be either smooth or zigzag in shape.

Native range

Dreissena polymorpha is native to the drainage basins of the Black, Caspian and Aral Seas (Gollasch and Leppäkoski 1999). *D. polymorpha* has been found as fossil in Central and Western Europe. Its recent reappearance is therefore regarded as a return migration. However, the return migration was only possible via shipping canals and therefore *D. polymorpha* is an alien species in the "new" habitats (Dr. Stefan Nehring, pers. comm.).

Alien distribution

History of introduction and geographical spread.

Prior to the 19th century, *Dreissena polymorpha* was found in the Black, Caspian, and Azov Seas (Stanczykowska 1977). Between 1800 and 1900, *D. polymorpha* more than doubled its range in Europe (Schloesser 1995). *D. polymorpha* is now introduced to north-west Russia, central and western Europe, southern Scandinavia, Britain, Ireland and North-America (Gollasch *et al.* 1999,

Minchin *et al.* 2002). It is now found throughout most of Europe and extending east into the western Asia and south into Turkey and Ireland in west (Mackie *et al.* 1989).

It is largely unclear how the initial expansion from the Black Sea and Aralo-Caspian Sea took place. *D. polymorpha* may have penetrated from the Black Sea via Dnepr, the Oginskij Canal (completed in 1804) to the River Neman and further to the Curonian Lagoon, SE Baltic Sea. However, it may have come via canals from the Caspian region using the Volga and its tributaries and Lakes Onega and Ladoga. In 1990 it was reported from brackish water in the eastern part of the Gulf of Finland after being present for 150-years in the nearby freshwater Lake Ladoga (Gollasch and Leppäkoski 1999, Minchin *et al.* 2002).

D. polymorpha was presumably introduced into Germany in the course of the extension of the inland waterway network at the beginning of 1800s. It reached the Netherlands by 1826, being found in the Rhine at Rotterdam, most probably carried with timber imports from the Baltic (Kearney and Morton 1970, Minchin *et al.* 2002).

The first introduction of the species in Estonia was into the estuary of Põlula Brook in the Gulf of Finland and Pärnu Bay in the Gulf of Riga in the mid-1800s. A second introduction took place through Lake Peipsi to the easternmost part of the Gulf of Finland in the 1930s.

In Poland, *D. polymorpha* were first identified in the end of 1800s. The mussel is now found mainly in the northern half of the country territory but single sites are known from the upper drainage basins of the Odra and Vistula. (Dr. Katarzyna Zajac, pers. comm.)

Pathways of introduction

The main pathways of the expansion in the range of *D. polymorpha* are through inland navigation, particularly since the opening of new waterways between eastern and central Europe at the beginning of the 1800s (Martens 1865, Rebhan 1984, Kinzelbach 1992, Dreyer 1995, Reinhold and Tittizer 1997, Nehring and Leuchs 1999, Gollasch 1996, Orlova 2002, Nehring 2002), the transfer of animals (including crayfish) for stocking in farms (*e.g.* Thienemann 1950) and the introduction into lakes of mussels attached to boat hulls (*e.g.* Jungbluth 1996), also due to the increase of recreational boating during the post-war period.

Alien status in region

The numbers of *Dreissena* have started to decrease in the Baltic Sea due to the high water salinity and cold winters. Still generally the spread of *Dreissena* is increasing, especially in the warmer Liivi Bay and Peipsi Lake. The massive dispersal could be also caused by global climate changes (Kull 2005).

The species has been established in a few localities of the easternmost Gulf of Finland (Valovirta and Porkka 1996, Kotta *et al.* 1998) and in less saline parts of the Gulf of Riga (Kotta *et al.* 1998). *D. polymorpha* has not been found in the Väinameri.

In German freshwater the most important molluscan invader is *D. polymorpha* (see also table 1). This invasive species lives in lakes, dam reservoirs and large rivers, as well as in brackish water bodies connected to the North and Baltic Sea. Increasing water pollution in the mid-20th century strongly reduced *Dreissena* populations. The continuous improvement of water quality since the 1980s has allowed the populations to recover. Today they have again attained densities of up to 40,000 individuals per m² (Böhmer *et al.* 2001, Nehring 2005).

Between 1900-1925 *D. polymorpha* was rather seldom recorded in water bodies of Poland. During the subsequent 25 years, *D. polymorpha* records started to increase rapidly. Since 1975 the increase in number of new records has decreased, and at the same time in some of the species sites, the population number has also decreased (Katarzyna Zajac, pers. comm.).

It is not found in Iceland (Mandahl-Barth 1938).




Fig. 3. Crosses show the recent distribution area of *D. polymorpha* in the Estonian coastal sea. Map by Jonne Kotta, Source: [D. polymorpha fact-sheet](#), produced by Estonian Marine Institute.

Country	Not found	Not established	Rare	Local	Common	Very common	Not known
Denmark				X			
Estonia					X		
European part of Russia					X		
Finland				X			
Faroe Islands	X						
Germany						X	
Greenland	X						
Iceland	X						
Latvia					X		
Lithuania					X		
Norway	X						
Poland					X		
Sweden				X			

Table 1. The frequency and establishment of *Dreissena polymorpha*, please refer also to the information provided for this species at www.nobanis.org/search.asp. Legend for this table: **Not found** –The species is not found in the country; **Not established** - The species has not formed self-reproducing populations (but is found as a casual or incidental species); **Rare** - Few sites where it is found in the country; **Local** - Locally abundant, many individuals in some areas of the country; **Common** - Many sites in the country; **Very common** - Many sites and many individuals; **Not known** – No information was available.

Ecology

Habitat description

D. polymorpha colonizes lakes, rivers and brackish lagoons. Their preferred habitats include the calm waters upstream of dams *e.g.* on the central section of the Weser River (Busch *et al.* 1995). The mussels require suitable substrate for attachment. They are most abundant on hard surfaces, particularly rocky surfaces, and on macrophytes. (Dr. Katarzyna Zajac comm.) According to Kinzelbach (unpubl.) the mass occurrences of the species show that it is a typical r-strategist, i.e. short-lived but with great reproductive capacity.

Reproduction and life cycle

D. polymorpha usually spawns in May-July, and the fertilized egg give rise to a veliger larva up to 100 µm in size. The larva lives planktonically for 2 to 4 weeks and is capable of swimming rapidly by means of its velum (Einsle and Walz 1972, Einsle 1973, Orlova 2002). The final larval stage lasts for about one week, after which the mussel attaches to hard substrates by secreting byssal threads (termed "spatfall"; cf. Jantz 1996). According to Jantz and Schöll (1998), spatfall usually occurs in early summer, but it may also continue through the summer and into September. The spat settle on stones, deadwood, walls of embankments, the hulls of ships, and on buoys, and they may also attach themselves to crustaceans, snails and the shells of other bivalves (epoecious commensalisms, Kinzelbach 1969, cf. Wagner 1936), often forming dense clusters on these substrates. They are also capable of releasing old byssus threads and forming new ones.

In contrast to lake populations, the recruitment of *D. polymorpha* in rivers depends on populations located further upstream, because the larvae invariably drift downstream during their 2 to 4 week pelagic phase. The age and size composition of river populations vary greatly (Jantz *et al.* 1998, Jantz and Schöll 1998).

Dispersal and spread

The high reproductive output and ability to extend their planktonic stage enables *D. polymorpha* to disperse rapidly. Larvae and adults may be distributed in ballast water or as fouling on ship and boat hulls, navigation buoys, fishing vessel wells, as well as by transport of timber or river gravel, fish stocking water and fishing equipment. The overland transport of *D. polymorpha* by small trailed boats has been repeatedly implicated in inner-lake dispersal (Gollasch and Leppäkoski 1999).

Impact

Affected habitats and indigenous organisms

Owing to its large filtration capacity and mass occurrence *D. polymorpha* outcompetes the native species of similar feeding type. The species slows down the eutrophication processes, indirectly favours the blooms of blue green algae, increases water transparency and ameliorate the conditions for benthic macro-vegetation. Through biodeposition *D. polymorpha* increases the density of benthic deposit feeders. *Dreissena* consumes a lot of plankton in the water. In Lake Esrom, Denmark, 18 % of the phytoplankton production was consumed by *D. polymorpha* (Hamburger *et al.* 1990). This may therefore also cause a decline in the populations of some fish.

Native mussels belonging to the genera *Unio* and *Anodonta*, of which some are listed in national Red Lists in the region, are co-opted by zebra mussels as hard substrate for successful settlement. Those mussels that serve as "hosts" for *Dreissena* are effectively starved, because undisturbed filter feeding is no longer possible (Böhmer *et al.* 2001).

Due mainly to predation by aquatic birds, well-established *Dreissena* populations are nowadays a “non-problematic element of the ecosystem” (Siessegger, pers. comm.), while expanding stocks seem to repeatedly cause severe impacts, both ecological and economic, in their new areas (see below).

Genetic effects

There might be a risk of hybridization between *D. polymorpha* and *D. bugensis*. *D. polymorpha* and *D. bugensis* mussel hybrids were created by pooling gametes collected after exposure to serotonin in the laboratory, indicating that interspecies fertilization may be feasible (Mills *et al.*, 1996). There is, however, evidence for species-specific sperm attractants suggesting that interspecific fertilization may be rare in nature (hybrids are very rare both in North America and in the Volga River basin (M. Orlova, pers. comm.). If hybridization does occur; these hybrids will constitute a very small proportion of the dreissenid community (Mills *et al.* 1996). *D. bugensis* must have arrived more recently than *D. polymorpha* based on differences in size classes, and therefore it seems plausible that *D. bugensis* is still in the process of expanding its non-indigenous range (May and Marsden 1992, MacIsaac 1994). The absence of *D. bugensis* from areas where *D. polymorpha* are present may be related to the timing and location of introduction rather than physiological tolerances (MacIsaac 1994).

The genus *Dreissena* is highly polymorphic and has a high potential for rapid adaptation to extreme environmental conditions by the evolution of allelic frequencies and combinations, possibly leading to significant long-term impacts on North American waters (Mills *et al.* 1996).

Human health effects

Injuries to bathers, resulting from the sharp edges of the shells have been documented (Siessegger 1970, Minchin *et al.* 2002).

Economic and societal effects (positive/negative)

Negative economic impacts caused by *D. polymorpha* are fouling of intake pipes, ship hulls, navigational constructions, cages of aquaculture and reduced angling catches (Gollasch and Leppäkoski 1999, Minchin *et al.* 2002, Table 1).

When the mussels die, their decay causes accelerated corrosion. This problem has also affected supplies of drinking water taken from Lake Constance, and - until recently - the cooling water systems of power plants throughout Germany (Siplingen, Siessegger, pers. comm., see also Bernauer *et al.* 1996).

Another negative effect is the high cost of cleaning the shores from the sharp *Dreissena* shells (when they are massively spread across the shore, it is practically impossible to walk).

On the other hand *Dreissena* is an important food component for some fish, crayfish and some birds. Crushed shells could also be used as a fertilizers and poultry feed.

In conclusion, the impacts of *Dreissena* are many-sided, as summarized in Table 2.

Changes to environment	Changes to industry and leisure
Fouling of firm substrata	Pipe blockages
Increased water clarity	Fouling of craft
Increase of submerged macrophytes	Maintenance of navigation marks
Changes to benthos	Lacerations to bathers
Reduction of unionid populations	Municipal water supply
Changes to food-web dynamics	Disturbance from feeding birds
Fouling of marginal aquatic plants	Fouling of lock gate systems
Changes to phosphorus cycling	Water abstraction

Table 2. Impacts in Ireland following the introduction of zebra mussels (from Minchin *et al.* 2002).

Management approaches

Prevention methods

Prevention is only possible if the repeated introduction via the above mentioned pathways are reduced.

Eradication, control and monitoring efforts

Due to its sensitivity to anthropogenic influences, *Dreissena* is important as a bioindicator and biomonitoring organism (Franz 1992), and quantitative assessments have been conducted regularly since the 1960s in the context of water quality surveys *e.g.* in the Rhine (Schiller 1990).

D. polymorpha is one of the indicator species of the German Federal Government's Environmental Specimen Bank (UBA 1999). Jantz and Schöll (1998) name various monitoring systems that employ *Dreissena* as test organism (cf. ARGE Elbe 1991, Borchering 1992, 1994, Borchering and Volpers 1994, Busch *et al.* 1995, 1998, Franz 1992).

Chemical control of *D. polymorpha* can be achieved by applying chlorine, sodium hydroxide or potassium dichromate (Schalekamp 1971). The larvae suffer total mortality after exposure to ultrasonic vibration (22 to 800 kHz) for 3 minutes (Schalekamp 1971), but the technical effort involved is prohibitive. Exposure to radiation appears equally unfeasible because of the high dosage required (Jungen 1972). Other possible control methods are poisoning with a molluscicide, such as Bayer 73, air exposure, burning, and electrocution.

Fouled pipes for non-potable water and cooling water are cleaned with chemical agents, usually by exposing the mussels to sodium hydroxide (pH 11 to 12) for 24 to 36 h. This dissolves the byssal threads, thus detaching the mussels from the substrate (Siessegger 1971). In cooling water systems, *Dreissena* can be controlled by recirculating the water to raise its temperature to 35 or 40°C for a few hours. This also leads to the detachment of the byssal threads and to gaping of the shells, and the animals fall off after a few days (Siessegger 1971). In those sections of the cooling water system where thermal treatment is impossible, control may be achieved by temporary chlorination during the summer. In drinking water systems, chlorination (at high dosage, 2-10 mg Cl₂ per litre must be conducted at the intake filter, and it must subsequently be neutralized with activated charcoal (Siessegger 1971, 1973).

The timing of the control measures is also important. Ideally, control in non-potable water systems should take place in late fall, because larvae may be present until late October. Control of the larvae is the key, because they are small enough to pass through the sieves. The biomass of the juveniles subsequently increases 20 times in the course of the next two years.

The mussels die after 7 to 14 days of air exposure, but they do not detach themselves from the substrate, so that they need to be scraped off mechanically (Schalekamp 1971).

A combination of chemical and thermal procedures has proved particularly successful. Larvae are killed in a few minutes, and adult mussels in a few hours, at pH 12 and 30°C (Lentz 1993).

Other methods of control include: oxygen deprivation, thermal treatment, exposure and desiccation, radiation, manual scraping, high-pressure jetting, mechanical filtration, removable substrates, molluscicides, ozone, antifouling coatings, electric currents, and sonic vibration. Some industries even build their intake structures and piping at depths too low for zebra mussel colonization.

Biological control so far has proven to be ineffective in controlling *Dreissena* species. Predation by migrating diving ducks, fish species, and crayfish may reduce mussel abundance, though the effects are short-lived (Bially and MacIsaac 2000).

Other biological controls being researched are selectively toxic microbes and parasites that may play a role in management of *Dreissena* populations (Molloy 1998). Another prospective approach is to control *Dreissena* populations by disrupting the reproductive process, by interfering with the

synchronization of spawning by males and females in their release of gametes (Snyder *et al.* 1997). A further approach would be to inhibit the planktonic veliger from settling, since this is the most vulnerable stage in the life cycle (Kennedy 2002).

Information and awareness

In Estonia a booklet has been published which introduces invasive species of local importance including *Dreissena* (“[Invasiived võõrliigid Eestis](#)” 2005). The purpose of this booklet is to make the wider range of people aware of the problems going hand-in-hand with the spread of invasive species. Another purpose of the booklet is to explain and show how the species look and give some simple hints how the spread of species could be controlled. The species has also been included into the [Baltic Sea Alien Species Database](#).

Today the occurrence of *Dreissena polymorpha* in German waters is not well known by the general public, since a purposeful information platform is not installed in Germany yet. Education and awareness raising is needed (Stefan Nehring, pers. comm.).

Knowledge and research

The problems related to *Dreissena* are mainly well known and a lot of control and eradication methods have been worked out (see above). The distribution of the species is partly mapped and described several times (especially in the Baltic region, *e.g.* Gulf of Finland, Gulf of Riga).

Recommendations or comments from experts and local communities

A monitoring program for *D. polymorpha* is recommended. It should focus on the effectiveness of competition with amphipods (*e.g.* *Corophium curvispinum*) and other species (*e.g.* the mussel *Corbicula sp.*). There are no known prevention methods against the further spread of *Dreissena* in Estonia, but the Estonian Marine Institute maps for dispersal of the species provides a basis for future prevention actions.

References and other resources

Contact persons

Jonne Kotta (EE) Estonian Marine Institute, University of Tartu, Mäealuse 10A, EE-12618 Tallinn, Estonia, E-mail: jonne.kotta@sea.ee

Stefan Nehring (DE) AeT umweltplanung, Bismarckstrasse 19, D-56068 Koblenz, Phone: +49 261 1330398, E-mail: nehring@aet-umweltplanung.de

Dr. Katarzyna Zajac (PL) Institute of Nature Conservation, Polish Academy of Sciences, Al. Mickiewicza 33, PL-31-120 Krakow, tel. +48 609 440 104, E-mail: kzajac@iop.krakow.pl

Claus Lindegaard (DK) University of Copenhagen, Freshwater Biological Laboratory, 51 Helsingørsgade, DK-3400 Hillerød, E-mail: CLindegaard@bi.ku.dk

Erkki Leppäkoski (FI) Abo Akademi Universitet, Environmental and Marine Biology, FIN-20500 TURKU/ÅBO, Phone +358-40-595 5488, E-mail eleppako@abo.fi

Viktoras Didziulis (LT) Klaipeda University, CORPI, H. Manto 84, LT- Klaipeda, E-mail: viktoras@ekoinf.net

Gk̄sli Mj̄r Gk̄slason (IS) University of Iceland, Department of Biology, Sturlugata 7, IS-101 Reykjavik, Telephone: (354) 525 4617, E-mail: gmg@hi.is

Vadims Jermakovs (LV) Institute of Aquatic Ecology University of Latvia, Daugavgrivas 8, LV 1049 Riga, E-mail: vadims@hydro.edu.lv

Melanie Josefsson (SE) Swedish Environmental Protection Agency, SE-106 48 Stockholm, Sweden, tel: +48 18 67 31 48, fax: +46 18 67 31 56, E-mail Melanie.Josefsson@snv.slu.se

Links

[D. polymorpha fact-sheet](#), produced by Estonian Marine Institute

[D. polymorpha fact-sheet](#) developed by the University of Southern Mississippi/College of Marine Sciences/Gulf Coast Research Laboratory

Electronical version of booklet “[Invasive species in Estonia](#)” – in estonian, published by Estonian Ministry of Environment.

[Information about zebra mussel](#), provided by Great Lakes Information Network (e.g. maps, news)

[Baltic Sea Alien Species Database](#)

[Aquatic alien species in German inland and coastal waters](#) (database) – in english

Central Fisheries Board – Ireland – [General information on Zebra Mussel](#)

References

- ARGE Elbe [Arbeitsgemeinschaft für die Reinhaltung der Elbe der Länder Hamburg – Niedersachsen – Schleswig-Holstein] (ed., 1991): Biologisches Effektmonitoring mit der Dreikantmuschel *Dreissena polymorpha* in der Meßstation Schnackenburg. – Wassergütestelle Elbe
- Bernauer, D., Kappus, B. and W. Jansen (1996): Neozoen in Kraftwerksproben und Begleituntersuchungen am nördlichen Oberrhein. – In: H. Gebhardt, R. Kinzelbach, S. Schmidt-Fischer (eds.), *Gebietsfremde Tierarten*, 87-96. Landsberg.
- Bially, A. and H. J. MacIsaac. 2000. Fouling mussels (*Dreissena* spp.) colonize soft sediments in Lake Erie and facilitate benthic invertebrates. *Freshwater Biology* 43:85-97.
- Borcherding, J. (1992): A new biomonitor system for the detection of toxic discharges in the aquatic environment based on valve movements of the freshwater mussel *Dreissena polymorpha*. – In: D. Neumann, H. A. Jenner (eds., *Limnologie aktuell*, Band 4 – The zebra mussel *Dreissena polymorpha*: ecology, biological monitoring and first applications in water quality management, 127-146. Stuttgart.
- Borcherding, J. (1994): The „*Dreissena*-Monitor“ – improved evaluation of dynamic limits for the establishment of alarm thresholds during toxicity tests and for continuous water control. – In: I. R. Hill *et al.* (eds.), *Freshwater field tests for hazard assessment of chemicals*, 477-484. Boca Raton.
- Borcherding, J. and M. Volpers (1994): Der *Dreissena*-Monitor – Untersuchungen zum Verhalten der Zebra- und Quagga-Muscheln im Rhein. Störfallbewertung in der kontinuierlichen Gewässerüberwachung. – *Wasser und Boden* 46(8): 25-30.
- Busch, D., Schuchardt, B., Kettler, J. and B. Steinweg (1995): Die Verbreitung der Muschelarten *Dreissena polymorpha* und *Congria leucocephala* in der Weser und ihre Eignung für ein passives Schwermetallbiomonitoring. – In: Gerken and Schirmer (ed.), *Die Weser*, 109-122.
- Busch, D., Lucker, T. and T. Wosniok (1998): Effects of changing salt concentrations and other physical-chemical parameters on bioavailability and bioaccumulation of heavy metals in exposed *Dreissena polymorpha* (Pallas 1771). – *Limnologica* 28(3): 263-274.

- Böhmer, H.J., Heger, T., Trepl, L. (2001): Fallstudien zu gebietsfremden Arten in Deutschland - Case studies on Aliens Species in Germany. – Texte des Umweltbundesamtes 2001 (13), 126pp.
- Dreyer, U. (1995): Potenziale und Strategien der Wiederbesiedlung am Beispiel des Makrozoobenthons in der mittleren Elbe. PhD Thesis University Darmstadt: 102 pp.
- Einsle, U. (1973): Zur Horizontal- und Vertikalverteilung der Larven von *Dreissena polymorpha* im Pelagial des Bodensee-Obersees (1971). – gwf-wasser/abwasser 114(1): 27-30.
- Einsle, U. and N. Walz (1972): Die täglichen Vertikalwanderungen der Larven von *Dreissena polymorpha* Pallas im Bodensee-Obersee. – gwf-wasser/abwasser 113(9): 428-430.
- Franz, H. W. (1992): Der Rhein und seine Besiedlung im Wandel: Schwebstoffzehrende Organismen (Hydrozoa, Kamptozoa und Bryozoa) als Indikatoren für den ökologischen Zustand eines Gewässers. – Pollichia 25. Bad Dürkheim.
- Gollasch, S. (1996): Untersuchungen des Arteintrages durch den internationalen Schiffsverkehr unter besonderer Berücksichtigung nichteinheimischer Arten. – Hamburg.
- Gollasch, S. and Leppäkoski, E (eds.). 1999. Initial Risk Assessment of Alien Species in Nordic Coastal Waters. Nordic Council of Ministers, Copenhagen
- Hamburger K., Dall, P. C. and Jónasson, P. M. (1990): The role of *Dreissena polymorpha* Pallas (Mollusca) in the energy budget of Lake Esrom, Denmark. – Verh. Internat. Verein. Limnol. 24: 621-625.
- Jantz, B. (1996): Wachstum, Reproduktion, Populationsentwicklung und Beeinträchtigung der Zebrauschel (*Dreissena polymorpha*) in einem großen Fließgewässer, dem Rhein. – Diss. Univ. Köln.
- Jantz, B., Jantz, B. and Neumann, D. (1998): Growth and reproductive cycle of the zebra mussel in the River Rhine as studied in a river bypass. – Oecologia 114: 213-225.
- Jantz, B. and F. Schöll (1998): Größenzusammensetzung und Altersstruktur lokaler Bestände einer Zebrauschel-Flußpopulation – Untersuchungen am Rhein zwischen Basel und Emmerich (Rh-km 168-861). – Limnologica 28 (4): 395-413.
- Jungbluth, J.H. (1996): Einwanderer in der Molluskenfauna von Deutschland. I. Der chorologische Befund. In: Gebhardt, H., R. Kinzelbach and S. Schmidt-Fischer (Eds.), Gebietsfremde Tierarten. Ecomed, Landsberg, pp. 105-125
- Jungen, H. (1972): Ermittlung der letalen Strahlendosis für Larven der Wandermuschel *Dreissena polymorpha* Pallas. – Unveröff. Bericht, Wasserwerk Stadt Zürich.
- Jungen, H. (1972): Ermittlung der letalen Strahlendosis für Larven der Wandermuschel *Dreissena polymorpha* Pallas. – Unveröff. Bericht, Wasserwerk Stadt Zürich.
- Kearney, M.P. and Morton, B.S. 1970. The distribution of *Dreissena polymorpha* (Pallas) in Britain. Journal of Conchology 27: 97-100.
- Kennedy, V. S. 2002. Swimming and Settlement Behavior in the Quagga Mussel! Maryland Sea Grant. URL: http://www/mdsg.umd.edu/Research/R_ZM-03.html.
- Kinzelbach, R. (1969): Epökie der Wandermuschel (*Dreissena polymorpha* PALLAS). – Natur und Museum 99 (4): 155-158.
- Kinzelbach, R. (1992): The main features of the phylogeny and dispersal of the Zebra Mussel *Dreissena polymorpha*. – In: The Zebra Mussel *Dreissena polymorpha*. Limnologie aktuell 4: 5-17. Stuttgart.
- Kotta, J., Orav, H. and Kotta, I. 1998. Distribution and filtration activity of Zebra mussel, *Dreissena polymorpha* (Pallas) in the Gulf of Riga and the Gulf of Finland. Proc. Estonian Acad. Sci. Biol. Ecol. 47: 32-41.
- Kull, T. 2005. Invasiivsed võõrliigid Eestis (in estonian). Keskkonnaministeerium. Tallinn.
- Lentz, D. (1993): Möglichkeiten der Bekämpfung der Dreikant- oder Wandermuschel *Dreissena polymorpha* Pallas in wassertechnischen Anlagen. – Unveröff. Diplomarbeit, FB Chemie/Biotechnologie, Technische Fachhochschule Berlin.
- MacIsaac, H. G. 1994. Comparative growth and survival of *Dreissena polymorpha* and *Dreissena bugensis*, exotic mollusks introduced to the Great Lakes. J. Great Lakes Res. 20(4):783-790.
- Mackie, G. L., Gibbons, W. N., Muncaster, B. W., and Gray, I. M. 1989. The zebra mussel, *Dreissena polymorpha*: A synthesis of European experiences and a preview for North America. Toronto: Queens Printer for Ontario.
- Mandahl-Barth, G. 1938. Land- and Freshwater Mollusca. Zoology of Iceland 3 (65): 1-31.
- Martens, E. van (1865): Eine eingewanderte Muschel. Zoologischer Garten 6: 50–59, 89–97
- Wagner, H. (1936): Die Wandermuschel (*Dreissensia*) erobert den Platten-See. – Natur und Volk 66: 37-41.
- May, B. and J. E. Marsden. 1992. Genetic identification and implications of another invasive species of Dreissenid mussel in the Great Lakes. Can. J. Fish. Aquat. Sci. 49:1501-1506.
- Mills, E. L., G. Rosenberg, A. P. Spidle, M. Ludyanskiy, Y. Pligin, and B. May. 1996. A review of the biology and ecology of the quagga mussel (*Dreissena bugensis*), a second species of freshwater Dreissenid introduced to North America. Amer. Zool. 36:271-286.

- Minchin, D., Lucy, F. and Sullivan, M. (2002). Zebra mussel: Impacts and spread. In: Leppäkoski, E., Gollasch, S. and Olenin, S. (eds) Invasive Aquatic Species of Europe. Distribution, Impacts and Management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 135-146.
- Molloy, D. P. 1998. The potential of using biological control technologies in the management of *Dreissena*. Abstracts from the Eighth International Zebra Mussel and Other Nuisance Species Conference. California Sea Grant, Sacramento, California: 69-70.
- Nehring, S. and H. Leuchs (1999): Neozoa (Makrozoobenthos) an der deutschen Nordseeküste. Eine Übersicht. - Bundesanstalt für Gewässerkunde, Bericht BfG-1200. Koblenz.
- Nehring, S. (2002): Biological invasions into German waters: an evaluation of the importance of different human-mediated vectors for nonindigenous macrozoobenthic species. – In: Leppäkoski, E., S. Gollasch and S. Olenin (Eds.), Invasive Aquatic Species of Europe - Distribution, Impacts and Management. Kluwer Academic Publishers, Dordrecht: 373-383.
- Nehring, S. (2005): International shipping - A risk for aquatic biodiversity in Germany. – In: Nentwig, W., Bacher, S., Cock, M.J.W., Dietz, H., Gigon, A. and Wittenberg, R. (Eds.), Biological Invasions - From Ecology to Control. Neobiota 6: 125-143.0
- Orlova, M. I. (2002). *Dreissena (D.) polymorpha* (Pallas, 1771): evolutionary origin and biological peculiarities as prerequisites of invasion success. In: Leppäkoski, E., Gollasch, S. and Olenin, S. (eds) Invasive Aquatic Species of Europe. Distribution, Impacts and Management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 127-134.
- Rebhan, H. (1984): Wandermuschel, Keulenpolyp und Süßwassergarnele – Einwanderer unserer Schifffahrtswege. – Ber. Naturforsch. Ges. Bamberg 59: 37-48.
- Reinhold, M. and T. Tittizer (1997): Zur Rolle von Schiffen als Vektoren beim Faunenaustausch Rhein/Main/Main-Donau-Kanal/Donau. – Dt. Gewässerk. Mitt. 41, 199-205.
- Schalekamp, M. (1971): Warnung vor der Wandermuschel *Dreissena polymorpha* Pallas und Bekämpfung derselben. – GWA 1971/3 des Schweiz. Vereins von Gas- und Wasserfachmännern.
- Schiller, W. (1990): Die Entwicklung der Makrozoobenthosbesiedlung des Rheins in Nordrhein-Westfalen im Zeitraum 1969-1987. – In: R. Kinzelbach, G. Friedrich (ed.), Limnologie aktuell, Band 1, Biologie des Rheins, 259-275. Stuttgart.
- Schloesser, D.W. 1995. Introduced Species, Zebra Mussels In North America. In Encyclopedia Of Environmental Biology, Pp. 337-356. Academic Press, Inc., San Diego, California.
- Siessegger, B. (1970): Schäden durch die Wandermuschel „*Dreissena polymorpha* Pallas“? – Münchner Medizinische Wochenschrift 112(40): 1806-1807.
- Siessegger, B. (1971): Besteht durch die Wandermuschel *Dreissena polymorpha* Pallas eine Gefahr für den Bodensee und seine anliegenden Wasserwerke? – Wasser/Boden/Luft 7: 34-?.
- Siessegger, B. (1973): Untersuchungen zur Bekämpfung der *Dreissena polymorpha*. – Unveröff. Bericht, Langenargen.
- Snyder, F. L., M. B. Hilgendorf, and D. W. Garton. 1997. Zebra Mussels in North America: The invasion and its implications! Ohio Sea Grant, Ohio State University, Columbus, OH.
- Stanczykowska, A. 1977. Ecology of *Dreissena Polymorpha* (Pallas)(Bivalvia) In Lakes. Pol. Arch. Hydrolbiol. 24:461-530.
- Thienemann, A. (1950): Verbreitungsgeschichte der Süßwassertierwelt Europas. In: Thienemann, A. (Ed.), Die Binnengewässer, Bd. XVIII. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 809 pp.
- Umweltbundesamt (ed., 1999): Umweltprobenbank des Bundes. Ergebnisse aus den Jahren 1996 und 1997. – UBA-Texte 61/99. Berlin.
- Valovirta, I. and Porkka, M. 1996. The distribution and abundance of *Dreissena polymorpha* (Pallas) in the eastern Gulf of Finland. Mem. Soc. Fauna Flora Fenn. 72: 63-78.

Date of creation/modification of this species fact sheet – 26-09-2006