

Table 14. Tuberculosis (TB) Cases and Percentages by Reason Tuberculosis Therapy Stopped: United States, 1993–2003

Year	Total Cases ¹	Completed Therapy		Moved		Lost		Refused		Died ²		Unknown ³	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	23741	18042	(76)	1121	(5)	1087	(5)	223	(1)	3051	(13)	217	(1)
1994	23051	17761	(77)	1194	(5)	739	(3)	183	(1)	2743	(12)	431	(2)
1995	21711	17285	(80)	970	(5)	563	(3)	155	(1)	2390	(11)	348	(2)
1996	20297	16510	(81)	783	(4)	520	(3)	156	(1)	1991	(10)	337	(2)
1997	18930	15653	(83)	668	(4)	434	(2)	119	(1)	1755	(9)	301	(2)
1998	17584	14773	(84)	534	(3)	400	(2)	104	(1)	1578	(9)	195	(1)
1999	16863	14238	(84)	457	(3)	356	(2)	104	(1)	1439	(9)	269	(2)
2000	15785	13401	(85)	410	(3)	390	(3)	112	(1)	1294	(8)	178	(1)
2001	15410	13187	(86)	373	(2)	364	(2)	96	(1)	1114	(7)	276	(2)
2002	14556	12415	(85)	333	(2)	366	(3)	86	(1)	1069	(7)	287	(2)
2003	14382	12328	(86)	324	(2)	353	(3)	83	(1)	982	(7)	312	(2)

¹Includes all cases in persons reported as alive at diagnosis and taking one or more TB drugs.

²Died = died of any cause (not only TB).

³Includes cases in persons reporting reason therapy stopped = Other, Missing, or Unknown.

Note: Data for all years are updated through March 29, 2006.

Data complete through 2003 only. See Technical Notes (page 9) for details.