


# Estimated Number of AIDS Cases among Adults and Adolescents, by Transmission Category 1985–2006—United States and Dependent Areas




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.




# Estimated Number and Proportion of AIDS Cases among Adult and Adolescent Injection Drug Users 1985–2006—United States and Dependent Areas




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.



# Estimated Number and Proportion of AIDS Cases, by Sex and Transmission Category among Adults and Adolescents 2006—United States and Dependent Areas



N = 27,694



N = 10,121

- Male-to-male sexual contact
- Injection drug use
- Male-to-male sexual contact and injection drug use
- High-risk heterosexual contact\*
- Other/not identified†



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.



# Estimated Number of AIDS Cases among Male Adult and Adolescent Injection Drug Users, by Race/Ethnicity 2006—50 States and DC

<u>Race/Ethnicity</u>	<u>Cases</u>
White, not Hispanic	739
Black, not Hispanic	1,561
Hispanic	655
Asian/Pacific Islander	34
American Indian/ Alaska Native	14
<b>Total*</b>	<b>3,016</b>

Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Includes 63 males of unknown race or multiple races.



# Estimated Number of AIDS Cases among Female Adult and Adolescent Injection Drug Users, by Race/Ethnicity 2006—50 States and DC


Race/Ethnicity	Cases
White, not Hispanic	559
Black, not Hispanic	1,396
Hispanic	367
Asian/Pacific Islander	16
American Indian/ Alaska Native	17
<b>Total*</b>	<b>2,385</b>

Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Includes 29 females of unknown race or multiple races.



# Proportion of AIDS Cases among Adults and Adolescents, by IDU-related Transmission Category, 2006—United States and Dependent Areas



N = 10,700



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.  
IDU-related cases include: injection drug users (IDUs); men who have sex with men and inject drugs (MSM/IDU); and heterosexual sex partners of IDUs.



# Proportion of AIDS Cases among Adults and Adolescents, by IDU-related Transmission Category and Age Group at Diagnosis 2006—United States and Dependent Areas

N = 10,700

Expanded transmission category	Age group (years), %					Total
	13–19	20–24	25–34	35–44	≥45	
Injection drug use	63	58	60	64	72	67
Male-to-male sexual contact and injection drug use	14	22	21	20	13	17
Heterosexual contact with injection drug user	23	20	19	16	14	16



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

IDU-related cases include: injection drug users (IDUs); men who have sex with men and inject drugs (MSM/IDU); and heterosexual sex partners of IDUs.



# Proportion of AIDS Cases among Adult and Adolescent Injection Drug Users and Heterosexual Sex Partners of Injection Drug Users, by Race/Ethnicity and Age Group at Diagnosis, 2006—United States and Dependent Areas

N = 8,857

Race/Ethnicity	Age group (years), %					Total
	13–19	20–24	25–34	35–44	≥45	
White, not Hispanic	17	14	21	24	21	22
Black, not Hispanic	62	50	47	49	58	53
Hispanic	20	31	28	24	20	23




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.





# Proportion of AIDS Cases among Adult and Adolescent Males, IDU-related Transmission Category and by Race/Ethnicity 2002–2006—50 States and DC

N = 37,286\*



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.


\*Includes 409 males of unknown race or multiple races.

IDU-related cases include: injection drug users (IDUs); men who have sex with men and inject drugs (MSM/IDU); and heterosexual sex partners of IDUs.



# Proportion of AIDS Cases among Adult and Adolescent Females, by IDU-related Transmission Category and Race/Ethnicity 2002–2006—50 States and DC

N = 19,203\*




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Includes 202 females of unknown race or multiple races.

IDU-related cases include: injection drug users (IDUs); and heterosexual sex partners of IDUs.



# Estimated Number of AIDS Cases among Adult and Adolescent Injection Drug Users, by Area of Residence at Diagnosis 2006—United States and Dependent Areas



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.


\*Fewer than 10 AIDS diagnoses among injection drug users.

†Includes 30 persons of unknown state of residence at diagnosis.



# Estimated Number of AIDS Cases among Adult and Adolescent Injection Drug Users and Heterosexual Sex Partners of Injection Drug Users, by Region, 2006—50 States and DC


N = 8,465



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.



# Estimated Number of AIDS Cases among Adult and Adolescent Injection Drug Users, by Region and Race/Ethnicity, 2006—50 States and DC




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Includes 93 persons of unknown race or multiple races.



# Estimated Number of Adult and Adolescent Injection Drug Users Living with AIDS, by Sex, 2006—United States and Dependent Areas

N = 99,443




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

\*Fewer than 10 male (or female) injection drug users living with AIDS.

National and New York totals exclude 1 injection drug user whose sex was not reported.



# Estimated Number of HIV/AIDS Cases among Adults and Adolescents, by Transmission Category 2003–2006—33 states



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.

Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.


†Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.



# Proportion of HIV/AIDS Cases among Adults and Adolescents, by Sex and Transmission Category, 2006—33 States


Male HIV/AIDS cases

N = 25,928



Female HIV/AIDS cases

N = 9,252



- Male-to-male sexual contact
- Injection drug use
- Male-to-male sexual contact and injection drug use
- High-risk heterosexual contact\*
- Other/not identified†

Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

†Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.






# Proportion of HIV/AIDS Cases among Male Adult and Adolescent Injection Drug Users and Male Population by Race/Ethnicity, 2006—33 States


HIV/AIDS diagnoses  
N = 3,016\*

1% ————— <1%



Male population, 33 states  
N = 76,694,617

3% ————— 1%



White, not Hispanic  
 Black, not Hispanic  
 Hispanic  
 Asian/Pacific Islander  
 American Indian/  
Alaska Native



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.



Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.

Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Includes 15 males of unknown race or multiple races.



# Proportion of HIV/AIDS Cases among Female Adult and Adolescent Injection Drug Users and Female Population by Race/Ethnicity, 2006—33 States



White, not Hispanic  
 Black, not Hispanic  
 Hispanic  
 Asian/Pacific Islander  
 American Indian/  
Alaska Native



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.


Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.

Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Includes 8 females of unknown race or multiple races.



# Estimated Number of HIV/AIDS Cases among Adult and Adolescent Injection Drug Users, 2006—33 States



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Fewer than 10 cases among IDU.



# Estimated Number of HIV/AIDS Cases among Male Adult and Adolescent Injection Drug Users by Race/Ethnicity, 2006—33 States

Race/Ethnicity	Cases
White, not Hispanic	739
Black, not Hispanic	1,561
Hispanic	655
Asian/Pacific Islander	34
American Indian/Alaska Native	14
<b>Total*</b>	<b>3,016</b>

Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.

Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Includes 15 males of unknown race or multiple races.



# Estimated Number of HIV/AIDS Cases among Female Adult and Adolescent Injection Drug Users by Race/Ethnicity, 2006—33 States


Race/Ethnicity	Cases
White, not Hispanic	471
Black, not Hispanic	952
Hispanic	257
Asian/Pacific Islander	10
American Indian/Alaska Native	13
<b>Total*</b>	<b>1,712</b>

Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed. Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis. Data from states with confidential named-based HIV infection reporting since at least 2001.

\*Includes 8 females of unknown race or multiple races



# Proportion of HIV/AIDS Cases among Male Adult and Adolescent Injection Drug Users, by Race/Ethnicity 2003–2006—33 States N = 12,591\*




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.  
Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.  
Data from states with confidential named-based HIV infection reporting since at least 2001.  
\*Includes 102 males of unknown race or multiple races.



# Proportion of HIV/AIDS Cases among Female Adult and Adolescent Injection Drug Users, by Race/Ethnicity 2003–2006—33 States

N = 7,315\*




Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.  
Data include persons with a diagnosis of HIV infection regardless of AIDS status at diagnosis.  
Data from states with confidential named-based HIV infection reporting since at least 2001.  
\*Includes 61 females of unknown race or multiple races.



# Estimated Number of Adult and Adolescent Injection Drug Users Living with HIV Infection (not AIDS), by Sex, 2006—33 States

N = 35,343



Note. Data have been adjusted for reporting delays, and cases without risk factor information have been proportionally redistributed.

Data from states with confidential named-based HIV infection reporting since at least 2001.


\*Fewer than 10 male (or female) injection drug users living with HIV infection (not AIDS).

Florida has confidential name-based HIV infection reporting for only diagnoses made during July 1997 or later.





# Proportion of Reported Cases of HIV Infection (not AIDS) among Adults and Adolescents by IDU-related Transmission Category, 2006—50 Areas



Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December, 2006. IDU-related cases include: injection drug users (IDUs); men who have sex with men and inject drugs (MSM/IDU); and heterosexual sex partners of IDUs.



# Proportion of Reported Cases of HIV infection (not AIDS) among Adults and Adolescents, by IDU-Related Transmission Category and Age Group, 2006—50 Areas

N = 8,638

Expanded transmission category	Age group (years), %					Total
	13–19	20–24	25–34	35–44	≥45	
Injection drug use	51	53	59	67	75	65
Male-to-male sexual contact and injection drug use	28	32	29	21	13	23
Heterosexual contact with injection drug user	21	15	12	12	11	12



Note. Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December, 2006. IDU-related cases include: injection drug users (IDUs); men who have sex with men and inject drugs (MSM/IDU); and heterosexual sex partners of IDUs.



# Reported Cases of HIV Infection (not AIDS) Related to Injection Drug Use among Adults and Adolescents, by Race/Ethnicity and Age Group, Cumulative through 2006—50 Areas

N = 58,277\*

Race/Ethnicity	Age group (years), %					Total
	13–19	20–24	25–34	35–44	≥45	
White, not Hispanic	35	39	36	29	21	30
Black, not Hispanic	41	34	40	51	60	47
Hispanic	21	24	23	19	17	20

Note: Data from 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December, 2006.

IDU-related reports include: injection drug users (IDU); men who have sex with men and inject drugs (MSM/IDU); and heterosexual sex partners of IDUs.

\*Includes 485 persons of unknown race or multiple races.

