

PART III

OTHER INFORMATION ON ASBESTOS



This fact sheet answers the most frequently asked health questions about asbestos. For more information, you may call 404-639-6000. This fact sheet is one in a series of summaries about hazardous substances and their health effects. This information is important because this substance may harm you. The effects of exposure to any hazardous substance depend on the dose, the duration, how you are exposed, personal traits and habits, and whether other chemicals are present.

SUMMARY: Exposure to asbestos usually occurs by breathing contaminated air in workplaces that make or use asbestos. Asbestos is also found in the air of buildings containing asbestos that are being torn down or renovated. Asbestos exposure can cause cancer and other serious lung problems. This substance has been found in at least 58 of the 1,430 National Priorities List sites identified by the Environmental Protection Agency (EPA).

What is asbestos?

(Pronounced äs-bēs'təs)

Asbestos is the name that's used for a group of six different fibrous minerals (amosite, chrysotile, crocidolite, and the fibrous varieties of tremolite, actinolite, and anthophyllite) that occur naturally in soil and rocks in some areas. Asbestos fibers vary in length and may be straight or curled.

Asbestos fibers are resistant to heat and most chemicals. Because of this, asbestos fibers are used for a wide range of manufactured goods, mostly roofing shingles, ceiling and floor tiles, paper products, asbestos cement products, friction products (automobile clutch, brake, and transmission parts), textiles, packaging, gaskets, and coatings.

What happens to asbestos when it enters the environment?

- Asbestos can enter the air and water from the weathering of natural deposits and the wearing down of manufactured asbestos products, such as brake pads.
- Small fibers may remain suspended in the air for a long time before settling. Larger fibers tend to settle more quickly.

- Asbestos fibers aren't able to move through soil and they aren't broken down to other compounds in the environment. Therefore, they can remain in the environment for decades or longer.
- Asbestos fibers may build up in animals.

How might I be exposed to asbestos?

- Breathing low levels in air.
- Breathing higher levels in air while working in industries that make or use asbestos products or near a building that contains asbestos products and is being torn down or renovated.
- Breathing higher levels in air near an asbestos-related industry or near an asbestos-containing waste site.
- Drinking water containing asbestos from natural sources or from asbestos-containing cement pipes in drinking water distribution systems.

How can asbestos affect my health?

Asbestos mainly affects the lungs. Changes in the membrane surrounding the lung are quite common in workers exposed to asbestos. These are also sometimes found in people living in areas with high levels of asbestos in the air, but effects on breathing usually aren't serious.

ATSDR Internet home page via WWW is <http://atsdr1.atsdr.cdc.gov:8080/atsdrhome.html>

Breathing very high levels of asbestos may result in a slow buildup of scar-like tissue in the lungs and in the membrane that surrounds the lungs. This disease is called asbestosis, and is usually found in asbestos workers and not in the general public. People with asbestosis have shortness of breath, often along with a cough and sometimes heart enlargement. This is a serious disease and can eventually lead to disability or death.

How likely is asbestos to cause cancer?

The Department of Health and Human Services (DHHS) has determined that asbestos is a known carcinogen.

It is known that asbestos causes cancer in people. There are two types of cancer caused by exposure to high levels of asbestos: cancer of the lung tissue itself and mesothelioma, a cancer of the membrane that surrounds the lung and other internal organs. Both of these are usually fatal. These diseases don't develop immediately, but show up only after many years.

Interactions between cigarette smoke and asbestos increase your chances of getting lung cancer. Studies of workers suggest that breathing asbestos can increase the chances of getting cancer in other parts of the body (stomach, intestines, esophagus, pancreas, kidneys), but this is not certain.

People who are exposed to lower levels of asbestos may also have an increased risk of developing cancer, but the risks are usually small and are difficult to measure.

It is not known whether ingesting asbestos causes cancer. Some people who had been exposed to asbestos fibers in their drinking water had higher-than-average death rates from cancer of the esophagus, stomach, and intestines. However, it isn't known whether this was caused by asbestos or by something else.

Is there a medical test to show whether I've been exposed to asbestos?

Chest X-rays cannot detect asbestos fibers, but can detect early signs of lung disease caused by asbestos. Other tests (lung and CAT scans), are also useful in detecting changes in the lungs.

Tests exist to measure asbestos fibers in urine, feces, mucus, or material rinsed out of the lung. However, low levels of asbestos fibers are found in these body fluids in nearly all people, so higher-than-average levels can only show that you have been exposed to asbestos, not whether you will experience any health effects.

Has the federal government made recommendations to protect human health?

In 1989, the EPA banned all new uses of asbestos; uses established before this date are still allowed. The EPA has established regulations that require school systems to inspect for damaged asbestos and to eliminate or reduce the exposure by removing the asbestos or by covering it up. The EPA has set a limit of 7 million fibers per liter (MFL) as the concentration of long asbestos fibers that may be present in drinking water.

Glossary

Carcinogen: A substance that can cause cancer.

CAS: Chemical Abstract Service.

MFL: Million fibers per liter.

CAT scan: A medical test in which a computer makes a 3-dimensional image of a body organ.

References

Agency for Toxic Substances and Disease Registry (ATSDR). 1995. Toxicological profile for asbestos. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.

Where can I get more information? ATSDR can tell you where to find occupational and environmental health clinics. Their specialists can recognize, evaluate, and treat illnesses resulting from exposure to hazardous substances. You can also contact your community or state health or environmental quality department if you have any more questions or concerns. For more information, contact: Agency for Toxic Substances and Disease Registry, Division of Toxicology, 1600 Clifton Road NE, Mailstop E-29, Atlanta, GA 30333, Phone: 404-639-6000, FAX: 404-639-6315. ATSDR Internet home page via WWW is <http://atsdr1.atsdr.cdc.gov:8080/atsdrhome.html>



CANCER FACTS

National Cancer Institute • National Institutes of Health

Questions and Answers About Asbestos Exposure

1. What is asbestos?

"Asbestos" is the name given to a group of minerals that occur naturally as masses of strong, flexible fibers that can be separated into thin threads and woven. These fibers are not affected by heat or chemicals and do not conduct electricity. For these reasons, asbestos has been widely used in many industries. Four types of asbestos have been commonly used:

- Chrysotile, or white asbestos (curly, flexible white fibers), which accounts for about 90 percent of the asbestos currently used in industry;
- Amosite (straight, brittle fibers that are light gray to pale brown in color);
- Crocidolite, or blue asbestos (straight blue fibers); and
- Anthophyllite (brittle white fibers).

Chrysotile asbestos, with its curly fibers, is in the serpentine family of minerals. The other types of asbestos, which all have needle-like fibers, are known as amphiboles.

Asbestos fiber masses tend to break easily into a dust composed of tiny particles that can float in the air and stick to clothes. The fibers may be easily inhaled or swallowed and can cause serious health problems.

2. How is asbestos used?

Asbestos has been mined and used commercially in North America since the late 1800s, but its use increased greatly during World War II. Since then, it has been used in many industries. For example, the building and construction industry uses it for strengthening cement and plastics as well as for insulation, fireproofing, and sound absorption. The shipbuilding industry has used asbestos to insulate boilers, steampipes, hot water pipes, and nuclear reactors in ships. The automotive industry uses asbestos in vehicle brakeshoes and clutch pads. More than 5,000 products contain or have contained asbestos, some of which are listed below:

- Asbestos cement sheet and pipe products used for water supply and sewage piping, roofing and siding, casings for electrical wires, fire protection material, chemical tanks, electrical switchboards and components, and residential and industrial building materials;
- Friction products, such as clutch facings; brake linings for automobiles, railroad cars, and airplanes; and industrial friction materials;
- Products containing asbestos paper, such as table pads and heat-protective mats, heat and electrical wire insulation, industrial filters for beverages, small appliance components, and underlying material for sheet flooring;
- Asbestos textile products, such as packing components, roofing materials, heat- and fire-resistant clothing, and fireproof draperies; and
- Other products, including ceiling and floor tile; gaskets and packings; paints, coatings, and sealants; caulking and patching tape; and plastics.

In the late 1970s, the U.S. Consumer Product Safety Commission banned the use of asbestos in wallboard patching compounds and gas fireplaces because these products released excessive amounts of asbestos fibers into the environment. In addition, asbestos was voluntarily withdrawn by manufacturers of electric hair dryers. These and other regulatory actions, coupled with widespread public concern about the hazards of asbestos, have resulted in a significant annual decline in U.S. use of asbestos: Domestic use of asbestos amounted to about 560,000 metric tons in 1979, but it had dropped to about 55,000 metric tons by 1989.

3. What are the health hazards of exposure to asbestos?

Exposure to asbestos may increase the risk of several serious diseases:

- Asbestosis—a chronic lung ailment that can produce shortness of breath and permanent lung damage and increase the risk of dangerous lung infections;
- Lung cancer;
- Mesothelioma—a relatively rare cancer of the thin membranes that line the chest and abdomen; and
- Other cancers, such as those of the larynx and of the gastrointestinal tract.

4. Who is at risk?

Since the early 1940s, millions of American workers have been exposed to asbestos dust, including many of the 4.5 million men and women who worked in shipyards during the peak shipbuilding years of World War II. Health hazards from asbestos dust have been recognized in workers exposed in shipbuilding trades, asbestos mining and milling, manufacturing of asbestos textiles and other asbestos products, insulation work in the construction and building trades, brake repair, and a variety of other trades. Demolition workers, drywall removers, and firefighters also may be exposed to asbestos dust. As a result of Government regulations and improved work practices, today's workers (those without previous exposure) are likely to face smaller risks than did those exposed in the past.

Although it is known that the risk to workers increases with heavier exposure and longer exposure time, investigators have found asbestos-related diseases in some shipyard workers exposed to high levels of asbestos fibers for only brief periods (as little as 1 or 2 months). Even workers who may not have worked directly with asbestos but whose jobs were located near contaminated areas have developed asbestosis, mesothelioma, and other cancers associated with asbestos exposure.

Generally, workers who develop asbestos-related diseases show no signs of illness until many years after first exposure. For example, the time between first exposure to asbestos and the appearance of lung cancer is generally 15 years or more; a lag of 30 to 35 years is not unusual. The lag period for development of mesothelioma and asbestosis is even greater, often as long as 40 to 45 years.

There is also some evidence that family members of workers heavily exposed to asbestos face an increased risk of developing mesothelioma and perhaps other asbestos-related diseases. This risk is thought to result from exposure to asbestos dust brought into the home on the shoes, clothing, skin, and hair of workers.

5. How great is the risk?

Not all workers exposed to asbestos will develop diseases related to their exposure. In fact, many will experience no ill effects.

Asbestos that is bonded into finished products such as walls, tiles, and pipes poses no risk to health as long as it is not damaged or disturbed (for example, by sawing or drilling) in such a way as to release fibers into the air. When asbestos particles are set free and inhaled, however, exposed individuals are at risk of developing an asbestos-related disease. Once these nearly indestructible fibers work their way into body tissues, they tend to stay there indefinitely.

The risk of developing asbestos-related diseases varies with the type of industry in which the exposure occurred and with the extent of the exposure. In addition, different

types of asbestos fibers may be associated with different health risks. For example, results of several studies suggest that crocidolite and amosite are more likely than chrysotile to cause lung cancer, asbestosis, and, in particular, mesothelioma. Even so, no fiber type can be considered harmless, and proper safety precautions should always be taken by people working with asbestos.

6. How does smoking affect risk?

Many studies have shown that the combination of smoking and asbestos exposure is particularly hazardous. Cigarette smokers, on the average, are 10 times as likely to develop lung cancer as are nonsmokers. For nonsmokers who work with asbestos, the risk is about five times greater than for those in the general population. By contrast, smokers who also are heavily exposed to asbestos are as much as 90 times more likely to develop lung cancer than are nonexposed individuals who do not smoke. Smoking does not appear to increase the risk of mesothelioma, however.

There is evidence that quitting smoking will reduce the risk of lung cancer among asbestos-exposed workers, perhaps by as much as half or more after at least 5 years without smoking. People who were exposed to asbestos on the job at any time during their life or who suspect they may have been exposed *should not smoke*. If they smoke, they should stop.

7. Who needs to be examined?

Individuals who have been exposed (or suspect they have been exposed) to asbestos dust on the job or at home via a family contact should inform their physician of their exposure history and any symptoms. A thorough physical examination, including a chest x-ray and lung function tests, may be recommended. Interpretation of the chest x-ray may require the help of a specialist who is experienced in reading x-rays for asbestos-related diseases. Other tests also may be necessary.

As noted earlier, the symptoms of asbestos-related diseases may not become apparent for many decades after exposure. If any of the following symptoms develop, a physical examination should be scheduled without delay:

- Shortness of breath;
- A cough or a change in cough pattern;
- Blood in the sputum (fluid) coughed up from the lungs;
- Pain in the chest or abdomen;

- Difficulty in swallowing or prolonged hoarseness; and/or
- Significant weight loss.

8. What are the treatments for asbestos-related diseases?

The key to successful treatment of asbestos-related diseases lies in early detection. The health problems caused by asbestosis are due mainly to lung infections, like pneumonia, that attack weakened lungs. Early medical attention and prompt, aggressive treatment offer the best chance of success in controlling such infections. Depending on the situation, doctors may give a vaccine against influenza or pneumococcal pneumonia as a protective measure.

Treatment of cancer is tailored to the individual patient and may include surgery, anticancer drugs, radiation, or combinations of these therapies. Information about cancer treatment is available from the National Cancer Institute-supported Cancer Information Service, whose toll-free telephone number is 1-800-4-CANCER.

9. How can workers protect themselves?

Employers are required to follow regulations dealing with asbestos exposure on the job that have been issued by the Occupational Safety and Health Administration (OSHA), the Federal agency responsible for health and safety regulations in the workplace. Regulations related to mine safety are enforced by the Mine Safety and Health Administration (MSHA). Workers should use all protective equipment provided by their employers and follow recommended work practices and safety procedures. Workers who are or who have been exposed to asbestos should not smoke cigarettes.

Workers who are concerned about asbestos exposure in the workplace should discuss the situation with other employees, their union, and their employers. If necessary, OSHA can provide more information or make an inspection. Area offices of OSHA are listed in the "United States Government" section of telephone directories' blue pages (under "Department of Labor"). If no listing is found, workers may call or write to one of the OSHA regional offices listed on page 9. Mine workers may contact MSHA's Office of Standards, Variances, and Regulation at Room 627, 4015 Wilson Boulevard, Arlington, VA 22203; the telephone number is 703-235-1910.

The National Institute for Occupational Safety and Health (NIOSH) is another Federal agency that is concerned with asbestos exposure in the workplace. The Institute conducts asbestos-related research, evaluates work sites for possible health hazards, and makes safety recommendations. In addition, NIOSH distributes publications on the health effects of asbestos exposure and can suggest additional sources of information. The address is Office of Information, National Institute of Occupational Safety and

Health, 4676 Columbia Parkway/Mailstop C-19, Cincinnati, OH 45226. The toll-free telephone number is 1-800-35-NIOSH (1-800-356-4674).

10. What should people who have been exposed to asbestos do?

It is important for exposed individuals to:

- Stop smoking;
- Get regular health checkups;
- Get prompt medical attention for any respiratory illness; and
- Use all protective equipment, work practices, and safety procedures designed for working around asbestos.

11. Will the Government provide examinations and treatment or pay for such services? What about insurance coverage?

Medical services related to asbestos exposure are available through the Government only for certain groups of eligible individuals. In general, exposed individuals must pay for their own medical services unless they are covered by private or Government health insurance. Medicare may reimburse people with symptoms of asbestos-related diseases for the costs of diagnosis and treatment (following review of medical procedures for appropriateness). General and specific information about benefits is available from the Medicare office serving each state; for the telephone number of the nearest office, call 1-800-772-1213.

People with asbestos-related diseases also may qualify for financial help, including medical payments, under state workers' compensation laws. Because eligibility requirements vary from state to state, workers should contact the workers' compensation program in the state where the last exposure occurred. (The telephone number may be found in the blue pages of a local telephone directory.)

If exposure occurred during employment with a Federal agency (military or civilian), medical expenses and other compensation may be covered by the Federal Employees' Compensation Act. Workers who are or were employed in a shipyard by a private employer may be covered under the Longshoremen and Harbor Workers' Compensation Act. Information about eligibility or how to file a claim is available from the U.S. Department of Labor, Office of Workers' Compensation Programs, Room S-3229, 200 Constitution Avenue NW, Washington, DC 20210; the telephone number is 202-219-7552.

Retired military personnel and their eligible dependents may receive health care at any Department of Defense medical facility, Department of Veterans Affairs (VA) hospital, or Public Health Service hospital. Where no Federal facility is available, civilian facilities may be used under the Civilian Health and Medical Program for the Uniformed Services. Those over age 65 may be covered by Medicare. Former members of the military who believe they may have a service-related medical problem may inquire about care at a VA facility or telephone the local VA office.

Workers also may wish to contact their international union for information on other sources of medical help and insurance matters. One organization, the Asbestos Victims Special Fund Trust, provides financial assistance to asbestos victims who have not received workers' compensation or compensation through legal avenues. Information is available from the Trust at Suite M-11, 1500 Walnut Street, Philadelphia, PA 19102; the telephone number is 1-800-447-7590.

12. Is there a danger of nonoccupational exposure from products contaminated with asbestos particles?

Asbestos is so widely used that the entire population has been exposed to some degree. Air, beverages, drinking water, food, drug and dental preparations, and a variety of consumer products all may contain small amounts of asbestos. In addition, asbestos fibers are released into the environment from natural deposits in the earth and as a result of wear and deterioration of asbestos products.

The U.S. Environmental Protection Agency (EPA) regulates the general public's exposure to asbestos in buildings, drinking water, and the environment. The EPA's Toxic Substances Control Act (TSCA) Assistance Office can answer questions about toxic substances, including asbestos. Printed material is available on a number of topics, particularly on controlling asbestos exposure in schools and other buildings. The TSCA office can provide information about accredited laboratories for asbestos testing and can refer inquirers to other resources on asbestos. Questions may be directed to the TSCA Assistance Office, U.S. Environmental Protection Agency, 7408 M Street SW, Washington, DC 20024; the telephone number is 202-554-1404.

The Consumer Product Safety Commission (CPSC) is responsible for the regulation of asbestos in consumer products. The CPSC maintains a toll-free information line on the potential hazards of commercial products; the telephone number is 1-800-638-2772. In addition, CPSC provides information about laboratories for asbestos testing, guidelines for repairing and removing asbestos, and general information about asbestos in the home. Publications are available from the Office of Public Affairs, Consumer Product Safety Commission, 4330 East-West Highway, Bethesda, MD 20816; the telephone number is 301-504-0580.

The U.S. Food and Drug Administration is concerned with asbestos contamination of foods, drugs, and cosmetics and will answer questions on these topics. The address is Office of Consumer Affairs, Food and Drug Administration, HFE-88, 5600 Fishers Lane, Rockville, MD 20857; the telephone number is 301-443-3170.

13. What other organizations offer information related to asbestos exposure?

The American Lung Association and the American Cancer Society can provide information about lung disease, cancer, and smoking. Local chapters of these organizations are listed in telephone directories. Material about cancer and how to quit smoking is available by calling the National Cancer Institute-supported Cancer Information Service (CIS). The CIS, a program of the National Cancer Institute, provides a nationwide telephone service for cancer patients and their families, the public, and health care professionals. CIS information specialists have extensive training in providing up-to-date and understandable information about cancer and cancer research. They can answer questions in English and Spanish and can send free printed material. In addition, CIS offices serve specific geographic areas and have information about cancer-related services and resources in their region. The toll-free number of the CIS is 1-800-4-CANCER (1-800-422-6237).

**Occupational Safety and Health Administration (OSHA)
Regional Offices**

Region I (serves Connecticut, Massachusetts, Maine, New Hampshire, Rhode Island, and Vermont)	First Floor 133 Portland Street Boston, MA 02114 617-565-7164
Region II (serves New Jersey, New York, Puerto Rico, and the Virgin Islands)	Room 670 201 Varick Street New York, NY 10014 212-337-2356
Region III (serves the District of Columbia, Delaware, Maryland, Pennsylvania, Virginia, and West Virginia)	Gateway Building, Suite 2100 3535 Market Street Philadelphia, PA 19104 215-596-1201
Region IV (serves Alabama, Florida, Georgia, Kentucky, Mississippi, North Carolina, South Carolina, and Tennessee)	Suite 587 1375 Peachtree Street NE Atlanta, GA 30367 404-347-3573
Region V (serves Illinois, Indiana, Michigan, Minnesota, Ohio, and Wisconsin)	Room 3244 230 South Dearborn Street Chicago, IL 60604 312-353-2220
Region VI (serves Arkansas, Louisiana, New Mexico, Oklahoma, and Texas)	Room 602 525 Griffin Street Dallas, TX 75202 214-767-4731
Region VII (serves Iowa, Kansas, Missouri, and Nebraska)	Room 406 911 Walnut Street Kansas City, MO 64106 816-426-5861
Region VIII (serves Colorado, Montana, North Dakota, South Dakota, Utah, and Wyoming)	Federal Building, Room 1576 1961 Stout Street Denver, CO 80294 303-844-3061

Region IX
(serves American Samoa, Arizona,
California, Guam, Hawaii, Nevada, and
Trust Territories of the Pacific)

Room 415
71 Stevenson Street
San Francisco, CA 94105
415-744-6670

Region X
(serves Alaska, Idaho, Oregon,
and Washington)

Suite 715
1111 Third Avenue
Seattle, WA 98101-3212
206-553-5930

U.S. Department of Labor Program Highlights



Fact Sheet No. OSHA 93-06

BETTER PROTECTION AGAINST ASBESTOS IN THE WORKPLACE

What Is Asbestos?

Asbestos is a widely used, mineral-based material that is resistant to heat and corrosive chemicals. Typically, asbestos appears as a whitish, fibrous material which may release fibers that range in texture from coarse to silky; however, airborne fibers that can cause health damage may be too small to see with the naked eye.

Who Is Exposed?

An estimated 1.3 million employees in construction and general industry face significant asbestos exposure on the job. Heaviest exposures occur in the construction industry, particularly during the removal of asbestos during renovation or demolition. Employees are also likely to be exposed during the manufacture of asbestos products (such as textiles, friction products, insulation, and other building materials) and during automotive brake and clutch repair work.

What Are the Dangers of Asbestos Exposure?

Exposure to asbestos can cause asbestosis (scarring of the lungs resulting in loss of lung function that often progresses to disability and to death); mesothelioma (cancer affecting the membranes lining the lungs and abdomen); lung cancer; and cancers of the esophagus, stomach, colon, and rectum.

What Protections Are Mandatory?

The U.S. Occupational Safety and Health Administration (OSHA) has issued revised regulations covering asbestos exposure in general industry and construction. Both standards set a maximum exposure limit and include provisions for engineering controls and respirators, protective clothing, exposure monitoring, hygiene facilities and practices, warning signs, labeling, recordkeeping, and medical exams.

Nonasbestiform tremolite, anthophyllite, and actinolite were excluded from coverage under the asbestos standard in May 1992.

Here are some of the highlights of the revised rules, published in the Federal Register June 20, 1986; and on Sept. 14, 1988:

— Permissible Exposure Limit: In both general industry and construction, workplace exposure must be limited to 0.2 fibers per cubic centimeter of air (0.2 f/cc), averaged over an eight-hour work shift. The excursion or short-term limit is one fiber per cubic centimeter of air (1 f/cc) averaged over a sampling period of 30 minutes.

— Exposure Monitoring: In general industry, employers must do initial monitoring for workers who may be exposed above the "action level" of 0.1 f/cc. Subsequent monitoring must be conducted at reasonable intervals, in no case longer than six months for employees exposed above the action level.

In construction, daily monitoring must be continued until exposure drops below the action level (0.1 f/cc). Daily monitoring is not required where employees are using supplied-air respirators operated in the positive pressure mode.

— Methods of Compliance: In both general industry and construction, employers must control exposures using engineering controls, to the extent feasible. Where engineering controls are not feasible to meet the exposure limit, they must be used to reduce employee exposures to the lowest levels attainable and must be supplemented by the use of respiratory protection.

— Respirators: In general industry and construction, the level of exposure determines what type of respirator is required; the standards specify the respirator to be used.

— Regulated Areas: In general industry and construction, regulated areas must be established where the 8-hour TWA or 30-minute excursion values for airborne asbestos exceed the prescribed permissible exposure limits. Only authorized persons wearing appropriate respirators can enter a regulated area. In regulated areas, eating, smoking, drinking, chewing tobacco or gum, and applying cosmetics are prohibited.

Warning signs must be displayed at each regulated area and must be posted at all approaches to regulated areas.

This is one of a series of fact sheets highlighting U.S. Department of Labor programs. It is intended as a general description only and does not carry the force of legal opinion. This information will be made available to sensory impaired individuals upon request. Voice phone: (202) 219-8151. TDD message referral phone: 1-800-326-2577.

— **Labels:** Caution labels must be placed on all raw materials, mixtures, scrap, waste, debris, and other products containing asbestos fibers.

— **Recordkeeping:** The employer must keep an accurate record of all measurements taken to monitor employee exposure to asbestos. This record is to include: the date of measurement, operation involving exposure, sampling and analytical methods used, and evidence of their accuracy; number, duration, and results of samples taken; type of respiratory protective devices worn; name, social security number, and the results of all employee exposure measurements. This record must be kept for 30 years.

— **Protective Clothing:** For any employee exposed to airborne concentrations of asbestos that exceed the PEL, the employer must provide and require the use of protective clothing such as coveralls or similar full-body clothing, head coverings, gloves, and foot covering. Wherever the possibility of eye irritation exists, face shields, vented goggles, or other appropriate protective equipment must be provided and worn.

In construction, there are special regulated-area requirements for asbestos removal, renovation, and demolition operations. These provisions include a negative pressure area, decontamination procedures for workers, and a "competent person" with the authority to identify and control asbestos hazards. The standard includes an exemption from the negative pressure enclosure requirements for certain small scale, short duration operations provided special work practices prescribed in an appendix to the standard are followed.

— **Hygiene Facilities and Practices:** Clean change rooms must be furnished by employers for employees who work in areas where exposure is above the TWA and/or excursion limit. Two lockers or storage facilities must be furnished and separated to prevent contamination of the employee's street clothes from protective work clothing and equipment. Showers must be furnished so that employees may shower at the end of the work shift. Employees must enter and exit the regulated area through the decontamination area.

The equipment room must be supplied with impermeable, labeled bags and containers for the containment and disposal of contaminated protective clothing and equipment.

Lunchroom facilities for those employees must have a positive pressure, filtered air supply and be readily accessible to employees. Employees must wash their hands and face prior to eating, drinking or smoking. The employer must ensure that employees do not enter lunchroom facilities with protective work clothing or equipment unless surface fibers

have been removed from the clothing or equipment.

Employees may not smoke in work areas where they are occupationally exposed to asbestos.

— **Medical Exams:** In general industry, exposed employees must have a preplacement physical examination before being assigned to an occupation exposed to airborne concentrations of asbestos at or above the action level or the excursion level. The physical examination must include chest X-ray, medical and work history, and pulmonary function tests. Subsequent exams must be given annually and upon termination of employment, though chest X-rays are required annually only for 45 and over workers whose first asbestos exposure occurred more than 10 years ago.

In construction, examinations must be made available annually for workers exposed above the action level or excursion limit for 30 or more days per year or who are required to wear negative pressure respirators; chest X-rays are at the discretion of the physician.

Where Can I Get More Information?

Copies of the general industry asbestos standard Part II, Health Standards, Stock Number 869-017-00110-4, \$16.00) and the construction industry standard (Stock Number 869-017-00112-1, \$14.00) are available from the Superintendent of Documents, Government Printing Office, Washington, DC 20402-9325, or telephone 202-783-3238. These standards are also available on CD-ROM (Stock Number 729-013-00000-5, \$88.00) by subscription for four updates per year or a single disk for \$28.00.

Two pamphlets summarizing the rule are also available: (in single copies) "Asbestos Standard for General Industry" and "Asbestos Standard for Construction Industry," and can be obtained by sending a self-addressed mailing label to the OSHA Publications Office, Room N-3101, Washington, D.C. 20210, telephone 202-219-4667 or from any local OSHA office.

Questions about the standards can be answered by any local OSHA office or by OSHA regional offices located in Boston, New York, Philadelphia, Atlanta, Chicago, Dallas, Kansas City, Denver, San Francisco, and Seattle.

All local OSHA offices have available for loan slide programs on the general industry and construction asbestos standards. Training on asbestos and other safety and health hazards is conducted at the OSHA Training Institute, 1555 Times Dr., Des Plaines, IL 60018, telephone 708-297-4810; tuition is charged.

#

NTIS Web Site — <http://www.ntis.gov>

SHIP TO ADDRESS (please print or type)

CUSTOMER MASTER NUMBER (IF KNOWN)		DATE
ATTENTION NAME		
ORGANIZATION	DIVISION / ROOM NUMBER	
STREET ADDRESS		
CITY	STATE	ZIP CODE
PROVINCE / TERRITORY	INTERNATIONAL POSTAL CODE	
COUNTRY		
PHONE NUMBER ()	FAX NUMBER ()	
CONTACT NAME	INTERNET E-MAIL ADDRESS	

ORDER BY PHONE (ELIMINATE MAIL TIME)
 8:00 a.m. - 8:00 p.m., Eastern Time, M - F.
 Sales Desk: 1-800-553-NTIS (6847)
 (703) 605-6000

TDD: (703) 605-6043

CUSTOMER SERVICE: (703) 605-6050

ORDER BY FAX
 24 hours/7 days a week: (703) 321-8547
 To verify receipt of fax call: (703) 605-6090
 7:00 a.m. - 5:00 p.m., Eastern Time, M - F.

ORDER BY MAIL
 National Technical Information Service
 5285 Port Royal Road
 Springfield, VA 22161

RUSH SERVICE: available for an additional fee.

NTIS ORDERNOW™ ONLINE
 Order the most recent additions to the NTIS collection at NTIS Web site <http://www.ntis.gov/ordernow>.

ORDER VIA E-MAIL
 Order via E-mail 24 hours a day.
orders@ntis.fedworld.gov
 If concerned about Internet security, you may register your credit card at NTIS. Simply call (703) 605-6070.

BILL ME
 (U.S., Canada, and Mexico only)
 NTIS will gladly bill your order, for an additional fee of \$10.00. A request to be billed must be on a purchase order or company letterhead. An authorizing signature, contact name, and telephone number should be included with this request. Requests may be mailed or faxed.

METHOD OF PAYMENT (please print or type)

VISA MasterCard American Express Discover

CREDIT CARD NUMBER EXPIRATION DATE

CARDHOLDER'S NAME

NTIS Deposit Account Number:

Check / Money Order enclosed for \$ (PAYABLE TO NTIS IN U.S. DOLLARS)

PRODUCT SELECTION (please print or type)

NTIS PRODUCT NUMBER	INTERNAL CUSTOMER ROUTING (OPTIONAL) UP TO 8 CHARACTERS	UNIT PRICE	QUANTITY						INTERNATIONAL AIRMAIL FEES (SEE BELOW)	TOTAL PRICE
			PAPER COPY	MICRO-FICHE	MAGNETIC TAPE *	DISKETTE	CD-ROM	OTHER		
		\$							\$	\$
		\$							\$	\$
		\$							\$	\$
		\$							\$	\$
		\$							\$	\$
		\$							\$	\$
		\$							\$	\$
		\$							\$	\$

* CIRCLE REQUIREMENTS	3480 CARTRIDGE	1600 BPI	6250 BPI	LABELING		FORMAT	
				STANDARD	NONLABELED	EBCDIC	ASCII

TOTAL	\$
HANDLING FEE PER TOTAL ORDER	\$ 4.00
Outside North America-\$8	\$
GRAND TOTAL	\$

PLEASE NOTE

Unless microfiche or other is specified, paper copy will be sent.
 Please call the Sales Desk at 1-800-553-NTIS (6847) for information on multiple copy discounts available for certain documents.
Out-Of-Print Surcharge
 A 25% out-of-print surcharge will be added to titles acquired by NTIS more than three years prior to the current calendar year.
International Airmail Fees
 All regular prepaid orders are shipped "air-to-surface" unless airmail is requested. Airmail service is available for an additional fee. Canada and Mexico add \$4 per item. Other countries add \$8 per item.

Thank you for your order!

