401	PHARMACIST - DETACH HERE AND GIVE INSTRUCTIONS TO PATIENT
402 403	
404	<b>PATIENT INFORMATION</b>
405	
406	TYKERB (TIE-curb)
407	(lapatinib) tablets
408	
409	Read this leaflet before you start taking TYKERB® and each time you get a refill. There may be
410	new information. This information does not take the place of talking with your doctor about your
411	medical condition or treatment.
412 413	What is TYKERB?
413	TYKERB is used with the medicine capecitabine for the treatment of patients with advanced or
415	metastatic breast cancer that is HER2 positive, and who have already had certain other breast
416	cancer treatments.
417	cancer deatherts.
418	Before you start taking TYKERB, tell your doctor about all of your medical conditions,
419	including if you:
420	have heart problems.
421	• have liver problems. You may need a lower dose of TYKERB.
422	• are pregnant or may become pregnant. TYKERB may harm an unborn baby. If you become
423	pregnant during treatment with TYKERB, tell your doctor as soon as possible.
424	• are breastfeeding. It is not known if TYKERB passes into your breast milk or if it can harm
425	your baby. If you are a woman who has or will have a baby, talk with your doctor about the
426	best way to feed your baby.
427	
428	Tell your doctor about all the medicines you take, including prescription and nonprescription
429	medicines and herbal and dietary supplements. TYKERB and many other medicines may interact
430	with each other. Your doctor needs to know what medicines you take so he or she can choose the
431	right dose of TYKERB for you.
432	
433	Especially tell your doctor if you take:
434	<ul> <li>antibiotics and anti-fungals (drugs used to treat infections)</li> </ul>
435	• HIV (AIDS) treatments
436	<ul> <li>anticonvulsant drugs (drugs used to treat seizures)</li> </ul>
437	• calcium channel blockers (drugs used to treat certain heart disorders or high blood pressure)
438	<ul> <li>antidepressants</li> </ul>
439	drugs used for stomach ulcers
440	• St. John's Wort or other herbal supplements

441	
442	Know the medicines you take. Keep a list of your medicines with you to show your doctor. Do
443	not take other medicines during treatment with TYKERB without first checking with your
444	doctor.
445	
446	Because TYKERB is given with another drug called capecitabine, you should also discuss with
447	your doctor or pharmacist any medicines that should be avoided when taking capecitabine.
448	
449	How should I take TYKERB?
450	• Take TYKERB exactly as your doctor has told you. TYKERB and capecitabine are taken in
451	21 day cycles. The usual dose of TYKERB is 1,250 mg (5 tablets) taken by mouth, <b>one time</b>
452	a day on days 1 to 21. Your doctor will tell you the dose of capecitabine you should take
453	and when you should take it.
454	• TYKERB should be taken at least one hour before, or at least one hour after food.
455	Do not eat or drink grapefruit products while taking TYKERB.      TYKERB.
456	Your doctor may adjust your dose of TYKERB depending on how you tolerate the
457	treatment.
458	• If you forget to take your dose of TYKERB, take it as soon as you remember that day. If
459 460	you miss a day, do not double your dose the next day. Just skip the missed dose.
461	What are the possible side effects of TYKERB?
462	Serious side effects include:
463	heart problems
464	<ul> <li>decreased pumping of blood from the heart</li> </ul>
465	abnormal heartbeat
466	• liver problems, which may cause itching, yellow eyes or skin, dark urine, or pain or
467	discomfort in the right upper area of the belly.
468	Your doctor should do blood tests to check your liver before you start taking TYKERB
469	and during treatment.
470	• lung problems
471	• severe diarrhea, which may lead to you becoming dehydrated
472	
473	Call your doctor right away if you have palpitations, persistent cough, shortness of breath,
474	or severe diarrhea.
475	
476	Common side effects of TYKERB in combination with capecitabine include:
477	<ul> <li>diarrhea</li> </ul>

red, painful hands and feet

nausea

rash

478 479

480

- 481 vomiting
- 482 tiredness
- mouth sores
- loss of appetite
- indigestion

486

Tell your doctor about any side effect that gets serious or that does not go away.

488

These are not all the side effects with TYKERB. Ask your doctor or pharmacist for more information.

491

492 **You may also get side effects from capecitabine.** Talk to your doctor about possible side effects with capecitabine.

494 495

496

497

498

499

## How should I store TYKERB tablets?

- Store TYKERB tablets at room temperature between 59° and 86°F (15° to 30°C). Keep the container closed tightly.
  - Do not keep medicine that is out of date or that you no longer need. Be sure that if you throw any medicine away, it is out of the reach of children.
- Keep TYKERB and all medicines out of the reach of children.

500501502

503

504

505

## **General information about TYKERB**

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use TYKERB for any other condition for which it was not prescribed. Do not give TYKERB to other people, even if they have the same condition that you have. It may harm them.

506507

This leaflet summarizes the most important information about TYKERB. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about TYKERB that is written for health professionals. For more information you can call toll-free 1-888-825-5249.

512513

## What are the ingredients in TYKERB?

- 514 **Active Ingredient:** Lapatinib.
- 515 **Inactive Ingredients: Tablet Core:** Magnesium stearate, microcrystalline cellulose, povidone,
- sodium starch glycolate. **Coating:** Orange film-coat: FD&C yellow #6/sunset yellow FCF
- aluminum lake, hypromellose, macrogol/PEG 400, polysorbate 80, titanium dioxide.

518

519 TYKERB tablets are oval, biconvex, orange, film-coated with GS XJG printed on one side.

520



521522523

TYKERB is a registered trademark of GlaxoSmithKline.

524

## gsk GlaxoSmithKline

525

- 526 GlaxoSmithKline
- 527 Research Triangle Park, NC 27709

528

529 ©YEAR, GlaxoSmithKline. All rights reserved.

530

- Revised: Month YEAR
- 532 TKB:XPIL

533