

# Actifed Cold & Allergy, 100 tablets

Combination Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active Ingredients (in each tablet) Purpose</b>	
Pseudoephedrine HCl 60 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine
<b>Uses</b> ■ temporarily relieves these symptoms of hay fever:	
<ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ itchy, watery eyes</li> <li>■ itching of the nose or throat</li> <li>■ sneezing</li> <li>■ nasal congestion</li> </ul>	
■ temporarily relieves nasal congestion due to the common cold	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> <li>■ high blood pressure</li> <li>■ glaucoma</li> <li>■ heart disease</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ a breathing problem such as emphysema or chronic bronchitis</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul>	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	

<b>Drug Facts (continued)</b>	
When using this product	
<ul style="list-style-type: none"> <li>■ do not use more than directed</li> <li>■ avoid alcoholic drinks</li> <li>■ drowsiness may occur</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> <li>■ excitability may occur, especially in children</li> </ul>	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ new symptoms occur</li> <li>■ symptoms do not get better within 7 days or occur with a fever</li> </ul>	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

<b>Drug Facts (continued)</b>	
<b>Directions</b>	
adults and children 12 years and over	take 1 tablet every 4 to 6 hours; not more than 4 tablets in 24 hours
children 6 years to under 12 years	take 1/2 tablet every 4 to 6 hours; not more than 2 tablets in 24 hours
children under 6 years	ask a doctor
<b>Inactive ingredients</b> corn starch, hydroxypropyl methylcellulose, lactose, magnesium stearate, polyethylene glycol, potato starch, povidone, sucrose, titanium dioxide	

\* Note: 14 point Helvetica Condensed Bold Italic Title  
8 point Helvetica Condensed Bold Italic Headings  
6 point Helvetica Condensed Bold Subheadings  
6 point Helvetica Condensed Text  
6.5 point Leading

98N-0337

BKGI

Actifed Cold & Allergy, 20 caplets  
 Combination Product Using Standard Labeling Format\*  
 [Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 500 mg	Pain reliever/fever reducer
Pseudoephedrine HCl 30 mg	Nasal decongestant
Triprolidine HCl 1.25 mg	Antihistamine
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ temporarily relieves these symptoms of hay fever:</li> <li>■ runny nose    ■ itchy, watery eyes    ■ nasal congestion</li> <li>■ itching of the nose or throat    ■ sneezing</li> <li>■ temporarily relieves these symptoms due to the common cold:</li> <li>■ nasal congestion    ■ headache    ■ minor aches and pains</li> <li>■ runny nose    ■ sneezing</li> <li>■ temporarily relieves nasal congestion due to sinusitis</li> <li>■ temporarily reduces fever</li> </ul>	
<b>Warnings</b>	
<p>Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.</p> <p>Do not use you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>■ high blood pressure    ■ heart disease    ■ thyroid disease    ■ glaucoma</li> <li>■ a breathing problem such as emphysema or chronic bronchitis    ■ diabetes</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul> <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</p> <p>When using this product    ■ do not use more than directed</p> <ul style="list-style-type: none"> <li>■ drowsiness may occur    ■ avoid alcoholic drinks</li> <li>■ excitability may occur, especially in children</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> </ul> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless    ■ new symptoms occur</li> <li>■ redness or swelling is present    ■ symptoms do not get better</li> <li>■ you need to use more than 7 days</li> <li>■ fever gets worse or lasts more than 3 days</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>	
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>Drug Facts (continued)</b></p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 6 caplets in 24 hours</li> <li>■ children under 12 years: ask a doctor</li> </ul>	

<b>Drug Facts (continued)</b>
<b>Inactive ingredients:</b> carnauba wax, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

\* Note: 14 point Helvetica Condensed Bold Italic Title  
 8 point Helvetica Condensed Bold Italic Headings  
 6 point Helvetica Condensed Bold Subheadings  
 6 point Helvetica Condensed Text  
 6.5 point Leading

Amitone, 100 tablets

Product Using Standard Labeling Format\*  
[Bottle Label, No Outer Carton]

<b><i>Drug Facts</i></b>	
<b><i>Active ingredient (in each tablet)</i></b>	<b><i>Purpose</i></b>
Calcium carbonate 350 mg	Antacid
<b><i>Use relieves:</i></b> ■ heartburn ■ acid indigestion ■ sour stomach ■ upset stomach due to these symptoms	
<b><i>Warnings</i></b> Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs. Stop use and ask a doctor if symptoms last more than 2 weeks. Keep out of reach of children.	
<b><i>Directions</i></b> ■ chew 2 tablets. Repeat hourly if symptoms return. ■ do not take more than 22 tablets in 24 hours or use the maximum dosage for more than 2 weeks	
<b><i>Inactive ingredients</i></b> magnesium stearate, mineral oil, powdered peppermint, sodium hexametaphosphate, starch, stearic acid, sucrose, talc	

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

# Arthritis Pain Formula, Maximum Strength, 40 caplets

Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredient (in each caplet)</b>	<b>Purpose</b>
Aspirin 500 mg	Pain reliever
<b>Use</b> temporarily relieves minor aches and pains due to: ■ minor arthritis pain ■ headache ■ backache ■ a cold ■ muscular aches ■ toothache ■ premenstrual cramps and menstrual cramps	
<b>Warnings</b> Reye's syndrome: Children and teenagers should not use this drug for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported with aspirin. Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock	

<b>Drug Facts (continued)</b>
<b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.
<b>Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer</b>
<b>Ask a doctor before use if you have</b> ■ asthma ■ ulcers ■ bleeding problems ■ stomach problems that last or come back, such as heartburn, upset stomach, or pain
<b>Ask a doctor or pharmacist before use if you are taking a prescription drug for:</b> ■ anticoagulation (blood thinning) ■ gout ■ diabetes ■ arthritis
<b>Stop use and ask a doctor if</b> ■ allergic reaction occurs. Seek medical help right away. ■ pain gets worse or lasts more than 10 days ■ new symptoms occur ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present

<b>Drug Facts (continued)</b>
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b> ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor
<b>Inactive ingredients</b> aluminum hydroxide, hydrogenated vegetable oil, magnesium hydroxide, microcrystalline cellulose, sodium lauryl sulfate, starch

- \* Note: 14 point Helvetica Condensed Bold Italic Title  
8 point Helvetica Condensed Bold Italic Headings  
6 point Helvetica Condensed Bold Subheadings  
6 point Helvetica Condensed Text  
6.5 point Leading

# Ascriptin Regular Strength, 100 tablets

Product Using Standard Labeling Format\*

[Outer Carton]

<b>Drug Facts</b>	
<b>Active Ingredient (in each tablet)</b>	<b>Purpose</b>
Aspirin 325 mg	Pain reliever
<b>Use</b> temporary relieves minor aches and pains due to: ■ headaches ■ muscular aches ■ menstrual cramps ■ toothache ■ minor arthritis pain	
<b>Warnings</b> Reye's syndrome: Children and teenagers should not use this drug for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported with aspirin. Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding. Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer Ask a doctor before use if you have ■ asthma ■ ulcers ■ bleeding problems ■ stomach problems that last or come back, such as heartburn, upset stomach, or pain	

<b>Drug Facts (continued)</b>
Ask a doctor or pharmacist before use if you are taking a prescription drug for: ■ diabetes ■ gout ■ arthritis ■ anticoagulation (blood thinning)
Stop use and ask a doctor if ■ allergic reaction occurs. Seek medical help right away. ■ redness or swelling is present ■ ringing in the ears or loss of hearing occurs ■ new symptoms occur ■ pain gets worse or lasts more than 10 days
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b> ■ drink a full glass of water with each dose ■ adults and children 12 years and older: take 2 tablets every 4 hours while symptoms last; not more than 12 tablets in 24 hours ■ children under 12 years: ask a doctor
<b>Inactive Ingredients</b> aluminum hydroxide gel, calcium carbonate, hydroxypropyl methylcellulose, magnesium hydroxide, magnesium stearate, microcrystalline cellulose, propylene glycol, starch, talc, titanium dioxide

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

AsthmaHaler Mist Bronchodilator refill, 1/2 fl. oz.

Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredient (in each inhalation)</b>	<b>Purpose</b>
Epinephrine 0.22 mg	Bronchodilator
<b>Uses</b> ■ eases breathing for asthma patients by reducing spasms of bronchial muscles ■ temporarily relieves shortness of breath, lightness of chest, and wheezing due to bronchial asthma	
<b>Warnings</b>	
Asthma alert: Do not continue to use and get medical help right away if you ■ are not better in 20 minutes ■ get worse	
Do not use ■ unless a doctor has said you have asthma	

<b>Drug Facts (continued)</b>
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ ever been hospitalized for asthma ■ trouble urinating due to an enlarged prostate gland
Ask a doctor or pharmacist before use if you are taking a prescription drug for asthma
When using this product ■ over use may cause nervousness, rapid heartbeat, and heart problems ■ do not puncture or throw into incinerator. Contents under pressure. ■ do not use or store near open flame or heat above 120° F (49° C). May cause bursting.

<b>Drug Facts (continued)</b>
Contains CFC-12 and CFC-114, substances which harm public health and environment by destroying ozone in the upper atmosphere
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b> ■ do not use more often or at higher doses unless directed by a doctor ■ supervise children using this product ■ adults and children 4 years and over: start with 1 puff, then wait at least 1 minute. If not relieved, use once more. Do not use again for at least 3 hours. ■ children under 4 years: ask a doctor
<b>Inactive ingredients</b> alcohol 34%, ascorbic acid, fluorocarbons (propellant), water

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

Benadryl Spray Extra Strength, 2 fl. oz.  
 Combination Product Using Section 201.66(d)(10) Modified Format\*  
 [Immediate Container Label, No Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredients</b>	<b>Purpose</b>
Diphenhydramine HCl 2 %	External analgesic
Zinc acetate 0.1 %	Skin protectant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ dries the oozing and weeping of poison:               <ul style="list-style-type: none"> <li>■ ivy ■ oak ■ sumac</li> </ul> </li> <li>■ temporarily relieves pain and itching that occur with:               <ul style="list-style-type: none"> <li>■ minor skin irritations ■ insect bites</li> <li>■ rashes due to poison ivy, oak, sumac</li> </ul> </li> </ul>	
<b>Warnings</b>	
For external use only	
Flammable: Keep away from fire or flame	
When using this product	
<ul style="list-style-type: none"> <li>■ do not use in or near the eyes</li> <li>■ do not use on chicken pox, measles, blisters, or on extensive areas of skin, unless directed by a doctor</li> </ul>	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> <li>■ condition worsens</li> <li>■ symptoms last more than 7 days or clear up and occur again within a few days</li> </ul>	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>■ adults and children 12 years and over: apply to affected area not more than 3 to 4 times daily</li> <li>■ children under 12 years: ask a doctor</li> </ul>	
<b>Inactive ingredients</b> alcohol 73.5% v/v, glycerin, povidone, purified water, tromethamine	

\* Note: 9 point Helvetica Narrow Bold Italic Title  
 8 point Helvetica Narrow Bold Italic Headings  
 6 point Helvetica Narrow Bold Subheadings  
 6 point Helvetica Narrow Text  
 6.5 point Leading

Benylin Multi-Symptom, 4 fl. oz.

Combination Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active Ingredients</b> <i>(in each 5 mL)</i>	<b>Purpose</b>
Dextromethorphan HBr 5 mg.....	Antitussive
Guafenesin 100 mg.....	Expectorant
Pseudoephedrine HCl 15 mg.....	Nasal decongestant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ temporarily relieves:                             <ul style="list-style-type: none"> <li>■ cough due to minor throat and bronchial irritation from inhaled irritants</li> <li>■ nasal congestion due to common cold</li> <li>■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive</li> </ul> </li> </ul>	
<b>Warnings</b>	
<p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>	
<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>■ heart disease      ■ high blood pressure</li> <li>■ thyroid disease    ■ diabetes</li> <li>■ trouble urinating due to an enlarged prostate gland</li> <li>■ cough that occurs with too much phlegm (mucus)</li> <li>■ chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> </ul>	
<p>When using this product</p> <ul style="list-style-type: none"> <li>■ do not use more than directed</li> </ul>	
<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ symptoms do not get better within 7 days or occur with a fever</li> <li>■ cough lasts more than 7 days, comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.</li> </ul>	

<b>Drug Facts (continued)</b>	
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>■ do not take more than 4 doses in 24 hours</li> </ul>	
adults and children 12 years and over	take 20 mL every 4 hours
children 6 years to under 12 years	take 10 mL every 4 hours
children 2 years to under 6 years	take 5 mL every 4 hours
children under 2 years	ask a doctor
<b>Inactive Ingredients</b> caramel, citric acid, D&C red no. 33, edetate disodium, FD&C red no. 40, poloxamer 407, polyethylene glycol 1450, propylgallate, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution	

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 7 point Leading



BFI Antiseptic First-Aid Powder, 1/4 oz.  
Product Using Section 201.66(d)(10) Modified Format\*  
[Immediate Container]

<b>Drug Facts</b>	
<b>Active ingredient</b>	<b>Purpose</b>
Bismuth-formic-iodide 16%	First-aid antiseptic
<b>Use</b> helps reduce the risk of infection in minor: ■ cuts ■ scrapes ■ burns	
<b>Warnings</b> For external use only Do not use if you have ■ deep or puncture wounds ■ animal bites ■ serious burns When using this product ■ do not get in the eyes ■ do not apply over large areas of the body, especially raw skin, in infants and children Stop use and ask a doctor if condition lasts more than 1 week or worsens Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ clean the affected area ■ sprinkle freely to cover the area	
<b>Inactive ingredients</b> bismuth subgallate, boric acid, eucalyptol, magnesium carbonate, menthol, potassium alum, talc, thymol, zinc phenol sulfonate	

Ingredient Not In OTC Drug Review And Has No Compendial Monograph.

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6.25 point Leading

Caladryl Clear, 6 fl. oz.

Combination Product Using Standard Labeling Format\*  
[Immediate Container]

<b>Drug Facts</b>	
<b>Active ingredients</b>	<b>Purpose</b>
Catamine 8%	Skin protectant
Pramoxine HCl 1%	External analgesic
<b>Uses</b>	
■ temporarily relieves pain and itching due to: ■ insect bites ■ minor skin irritations ■ rashes due to poison ivy, poison oak, poison sumac ■ dries the oozing and weeping of: ■ poison ivy ■ poison oak ■ poison sumac	
<b>Warnings</b>	
For external use only	
When using this product do not use in or near the eyes	
Stop use and ask a doctor if	
■ condition worsens ■ symptoms last more than 7 days or clear up and occur again within a few days	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ shake well	
■ adults and children 2 years and over: apply to affected area not more than 3 to 4 times daily	
■ children under 2 years: ask a doctor	
<b>Inactive ingredients</b> SD alcohol 388, camphor, citric acid, diazolidinyl urea, glycerin, hydroxypropyl methylcellulose, methylparaben, polysorbate 40, propylene glycol, propylparaben, purified water, sodium citrate	

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

Caladryl Lotion, 6 fl. oz.

Combination Product Using Standard Labeling Format\*  
[Immediate Container]

<b>Drug Facts</b>	
<b>Active Ingredients</b>	<b>Purpose</b>
Calamine 8%	Skin protectant
Pramoxine HCl 1%	External analgesic
<b>Uses</b>	
■ temporarily relieves pain and itching due to:	
■ insect bites ■ minor skin irritations	
■ rashes due to poison ivy, poison oak, poison sumac	
■ dries the oozing and weeping of:	
■ poison ivy ■ poison oak ■ poison sumac	
<b>Warnings</b>	
For external use only	
When using this product do not use in or near the eyes	
Stop use and ask a doctor if	
■ condition worsens	
■ symptoms last more than 7 days or clear up and occur again within a few days	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away	
<b>Directions</b> ■ shake well	
■ adults and children 2 years and over: apply to affected area not more than 3 to 4 times daily	
■ children under 2 years: ask a doctor	
<b>Inactive Ingredients</b> SD alcohol 38B, camphor, diazolidinyl urea, hydroxypropyl methylcellulose, polysorbate 80, propylene glycol, propylparaben, purified water, xanthan gum	

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

# Konsyl

## Product Using Section 201.66(d)(10) Modified Format\* [Immediate Container]

<b>Drug Facts</b>	
<b>Active ingredient (in each tablet)</b>	<b>Purpose</b>
Calcium polycarbophil 625 mg	Bulk laxative
<b>Use</b> ■ relieves occasional constipation (irregularity) ■ usually causes bowel movement in 12 to 72 hours	
<b>Warnings</b>	
Choking: Taking this product without adequate liquid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.	
Ask a doctor before use if you have ■ stomach pain, nausea, or vomiting ■ a sudden change in bowel habits that lasts over 2 weeks	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Calcium may interact with certain prescription drugs.	
Stop use and ask a doctor if ■ you have no bowel movement within 72 hours or rectal bleeding. These could be signs of a serious condition. ■ you need to use a laxative more than 1 week if pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ taking this product without enough liquid may cause choking. See warnings. ■ drink a full glass (8 oz) of liquid with each dose.	
adults and children 12 years and over	2 tablets 1 to 4 times a day
children 6 to under 12 years	1 tablet 1 to 3 times a day
children under 6 years	ask a doctor
<b>Inactive ingredients</b> caramel crospovidone, ethyl cellulose, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, silicon dioxide	

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6.25 point Leading

Children's Kaopectate  
 Product Using Standard Labeling Format\*  
 [Immediate Container]

<b>Drug Facts</b>	
<b>Active ingredient (in each 15 mL) Purpose</b>	
Attapulgite 600 mg <span style="float: right;">Antidiarrheal</span>	
<b>Use</b>	
<ul style="list-style-type: none"> <li>■ reduces the number of bowel movements in diarrhea</li> </ul>	
<b>Warnings</b>	
<ul style="list-style-type: none"> <li>Do not use</li> <li>■ if you have fever</li> <li>■ in infants or children under 3 years of age</li> </ul>	
<ul style="list-style-type: none"> <li>Stop use and ask a doctor if</li> <li>■ you need to use an antidiarrheal product more than 2 days</li> </ul>	
<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>■ do not exceed 7 doses in 24 hours</li> </ul>	
<ul style="list-style-type: none"> <li>adults and children 12 years and over</li> </ul>	<ul style="list-style-type: none"> <li>30 mL after the first bowel movement; 30 mL after each subsequent bowel movement</li> </ul>
<ul style="list-style-type: none"> <li>children 6 to under 12 years</li> </ul>	<ul style="list-style-type: none"> <li>15 mL after the first bowel movement; 15 mL after each subsequent bowel movement</li> </ul>
<ul style="list-style-type: none"> <li>children 3 to under 6 years</li> </ul>	<ul style="list-style-type: none"> <li>7.5 mL after the first bowel movement; 7.5 mL after each subsequent bowel movement</li> </ul>
<b>Inactive Ingredients</b> FD&C red no. 40, glucono-delta-lactone, magnesium aluminum silicate, methylparaben, purified water, sorbic acid, sucrose, titanium dioxide, and xanthan gum	

\* Note: 14 point Helvetica Condensed Bold Italic Title  
 8 point Helvetica Condensed Bold Italic Headings  
 6 point Helvetica Condensed Bold Subheadings  
 6 point Helvetica Condensed Text  
 6.5 point Leading

Clear Eyes, 0.5 oz.

Combination Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active Ingredients</b>	<b>Purpose</b>
Glycerin 0.2%	Demolucenk
Naphazoline HCl 0.012%	Redness reliever
<b>Uses</b>	
■ temporarily relieves redness due to minor eye irritation	
■ for protection against further irritation or dryness of the eye due to:	
■ smoke ■ sun glare ■ swimming	
■ smog ■ contact lenses	
<b>Warnings</b>	
Ask a doctor before use if you have glaucoma	
<b>When using this product</b>	
■ do not touch tip of container to any surface to avoid contamination	
■ replace cap after each use	
■ do not use if liquid changes color or becomes cloudy	
■ overuse may cause more eye redness	
<b>Stop use and ask a doctor if</b>	
■ you feel eye pain ■ changes in vision occur	
■ redness or irritation of the eyes lasts	
■ condition worsens or lasts more than 72 hours	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
■ put 1 or 2 drops in the affected eye(s) up to 4 times daily	
<b>Inactive Ingredients</b> benzalkonium chloride, boric acid, edetate disodium, purified water, sodium borate	

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

Childrens Sudafed Nasal Decongestant, 4 fl. oz.

Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredient (in each 5 mL)</b>	<b>Purpose</b>
Pseudoephedrine HCl 15 mg	Nasal decongestant
<b>Use</b> temporarily relieves nasal congestion due to: <ul style="list-style-type: none"> <li>■ common cold</li> <li>■ hay fever</li> <li>■ upper respiratory allergies</li> <li>■ sinusitis</li> </ul>	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have: <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul>	
<b>When using this product</b>	
■ do not take more than directed	
<b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ symptoms do not get better within 7 days or occur with a fever</li> </ul>	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
adults and children 12 years and over	take 20 mL every 4 to 6 hours; not more than 80 mL in 24 hours
children 6 years to under 12 years	take 10 mL every 4 to 6 hours; not more than 40 mL in 24 hours
children 2 years to under 6 years	take 5 mL every 4 to 6 hours; not more than 20 mL in 24 hours
children under 2 years	ask a doctor

<b>Drug Facts (continued)</b>
<b>Inactive ingredients</b> citric acid, edetate disodium, FD&C red no. 40, FD&C blue no. 1, glycerin, poloxamer 407, polyethylene glycol 1450, povidone K-90, purified water, saccharin sodium, sodium benzoate, sodium citrate, sorbitol solution

\* Note: 14 point Helvetica Condensed Bold Italic Title  
 8 point Helvetica Condensed Bold Italic Headings  
 6 point Helvetica Condensed Bold Subheadings  
 6 point Helvetica Condensed Text  
 7 point Leading

# Congestac, 12 caplets

Combination Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Guafenesin 400 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal decongestant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial tubes of bothersome mucus, drain bronchial tubes, or make coughs more productive</li> <li>■ temporarily relieves nasal congestion due to the common cold</li> </ul>	
<b>Warnings</b>	
<p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ trouble urinating due to an enlarged prostate gland</li> <li>■ chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> </ul> <p>When using this product</p> <ul style="list-style-type: none"> <li>■ do not use more than directed</li> </ul> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ cough lasts more than 7 days, comes back, or occur with fever, rash, or headache that lasts. These could be signs of a serious condition.</li> <li>■ symptoms do not get better within 7 days or occur with a fever</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
<b>Directions</b>	
adults and children 12 years or over	take 1 caplet every 4 to 6 hours; not more than 4 caplets every 24 hours
children 6 years to under 12 years	take 1/2 caplet every 4 to 6 hours; not more than 2 caplets every 24 hours
children under 6 years	ask a doctor

**Drug Facts (continued)**

**Inactive ingredients** croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, silicon dioxide, starch

\* Note: 14 point Helvetica Condensed Bold Italic Title  
8 point Helvetica Condensed Bold Italic Headings  
6 point Helvetica Condensed Bold Subheadings  
6 point Helvetica Condensed Text  
7 point Leading



Coricidin Nighttime Cough and Cold, 6 fl. oz.  
 Combination Product Using Standard Labeling Format\*  
 [Outer Carton]

**Drug Facts**

---

**Active ingredients (in each 15 mL) Purpose**

Acetaminophen 325 mg.....Pain reliever  
 Diphenhydramine HCl 12.5 mg.....Antihistamine/cough suppressant

---

**Uses** temporarily relieves these cold and flu symptoms:

- headache
- minor aches and pains
- minor sore throat
- runny nose
- sneezing
- cough due to minor throat irritation

---

**Warnings**

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

**Sore throat warning:** Severe or persistent sore throat or sore throat that occurs with high fever, headache, nausea, and vomiting may be serious. Ask a doctor right away. Do not use more than 2 days or give to children under 3 years of age unless directed by a doctor.

---

**Ask a doctor before use if you have**

- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- glaucoma

---

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers**

---

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Drug Facts (continued)**

---

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- redness or swelling is present
- new symptoms occur
- pain or cough gets worse, lasts more than 5 days (children) or 7 days (adults)
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

---

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.**

---

**Directions**

- do not use more than directed

adults and children 12 years and older	take 30 mL every 6 hours; not more than 180 mL in 24 hours
children 6 years to under 12 years	take 15 mL every 4 hours; not more than 60 mL in 24 hours
children under 6 years	ask a doctor

---

**Inactive ingredients** citric acid, FD&C blue no. 1, FD&C red no. 40, glycerin, polyethylene glycol, propylene glycol, sodium citrate, sodium saccharin, sugar, water

\* Note: 14 point Helvetica Bold Italic Title  
 8 point Helvetica Bold Italic Headings  
 6 point Helvetica Bold Subheadings  
 6 point Helvetica Regular Text  
 7 point Leading

Correctol, 10 tablets

Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active Ingredient (in each tablet)</b> .....	<b>Purpose</b>
Bisacodyl 5 mg	Laxative
<b>Use</b>	
<ul style="list-style-type: none"> <li>■ relieves occasional constipation (irregularity)</li> <li>■ this product usually causes bowel movement in 6 to 12 hours</li> </ul>	
<b>Warnings</b>	
Do not use if you cannot swallow without chewing	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> <li>■ stomach pain, nausea, or vomiting</li> <li>■ a sudden change in bowel habits that lasts over 2 weeks</li> </ul>	
<b>When using this product</b>	
<ul style="list-style-type: none"> <li>■ do not chew or crush tablet</li> <li>■ do not use within 1 hour after taking an antacid or milk</li> <li>■ you may have stomach discomfort, faintness, or cramps</li> </ul>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>■ you no bowel movement within 12 hours or rectal bleeding. These could be signs of a serious condition.</li> <li>■ you need to use a laxative more than 1 week</li> </ul>	
if pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	
adults and children 12 years and over	take 1 to 3 tablets (usually 2) daily
children 6 to under 12 years	take 1 tablet daily
children under 6 years	ask a doctor
<b>Other information</b> ■ store at 20-25°C (68-77°F)	

**Drug Facts (continued)**  
**Inactive ingredients** acetylated monoglycerides, calcium sulfate, carnauba wax, coin starch, D&C red no. 7 calcium lake, gelatin, hydroxypropyl methylcellulose phthalate, lactose, magnesium stearate, povidone, sugar, tlc, titanium dioxide, white wax

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

# Drixoral Allergy Sinus, 12 extended release tablets

Combination Product Using Standard Labeling Format\*

[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Acetaminophen 500 mg	Pain reliever
Dexbrompheniramine maleate 3 mg	Antihistamine
Pseudoephedrine sulfate 60 mg	Nasal decongestant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ temporarily relieves these hay fever or other upper respiratory allergy symptoms:                             <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ itchy watery eyes</li> <li>■ itching of nose and throat</li> <li>■ sneezing</li> </ul> </li> <li>■ temporarily relieves nasal congestion due to a cold and sinusitis:                             <ul style="list-style-type: none"> <li>■ shrinks swollen membranes</li> <li>■ restores freer breathing through the nose</li> </ul> </li> <li>■ helps to decongest sinus openings and sinus passages</li> <li>■ reduces swelling of nasal passages</li> <li>■ temporarily relieves minor aches and pains and headaches</li> </ul>	
<b>Warnings</b>	
<p><b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.</p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ glaucoma</li> <li>■ a breathing problem such as emphysema or chronic bronchitis</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul> <p>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives</p> <p>When using this product</p> <ul style="list-style-type: none"> <li>■ do not use more than directed</li> <li>■ drowsiness may occur</li> <li>■ excitability may occur, especially in children</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> </ul> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ new symptoms occur</li> <li>■ redness or swelling is present</li> <li>■ symptoms do not get better</li> <li>■ you need to use more than 7 days</li> <li>■ fever occurs and lasts more than 3 days</li> </ul>	
<b>Drug Facts (continued)</b>	
<b>Other information</b>	
<ul style="list-style-type: none"> <li>■ store at 20-25°C (68-77°F)</li> <li>■ protect from excessive moisture</li> </ul> <p><b>Inactive ingredients</b> calcium phosphate, carnauba wax, D&amp;C yellow no. 10 aluminum lake, FD&amp;C yellow no. 6 aluminum lake, hydroxypropyl methylcellulose, magnesium stearate, methylparaben, PEG, propylparaben, stearic acid</p>	

**Drug Facts (continued)**

If you are pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** ■ children under 12 years: ask a doctor  
 ■ adults and children 12 years or over: take 2 tablets every 12 hours; not more than 4 tablets in 24 hours

\* Note: 14 point Helvetica Condensed Bold Italic Title

8 point Helvetica Condensed Bold Italic Headings

6 point Helvetica Condensed Bold Subheadings

6 point Helvetica Condensed Text

6.5 point-Leading

Drixoral Cold & Flu, 2 extended release tablets  
 Combination Product Using Standard Labeling Format\*  
 [Outer Carton]

<b>Drug Facts</b>	
<b>Active Ingredients (in each tablet)</b>	<b>Purpose</b>
Acetaminophen 500 mg	Pain reliever/fever reducer
Dexbrompheniramine maleate 3 mg	Antihistamine
Pseudoephedrine sulfate 60 mg	Nasal decongestant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ temporarily relieves these cold symptoms:               <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ sneezing</li> </ul> </li> <li>■ temporarily relieves nasal congestion due to a cold and sinusitis:               <ul style="list-style-type: none"> <li>■ shrinks swollen membranes</li> <li>■ restores freer breathing through the nose</li> </ul> </li> <li>■ helps to decongest sinus openings and sinus passages</li> <li>■ reduces swelling of nasal passages</li> <li>■ temporarily relieves minor aches and pains and headaches</li> <li>■ temporarily reduces fever</li> </ul>	
<b>Warnings</b>	
<p><b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.</p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ glaucoma</li> <li>■ a breathing problem such as emphysema or chronic bronchitis</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul> <p>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives</p> <p>When using this product</p> <ul style="list-style-type: none"> <li>■ do not use more than directed</li> <li>■ drowsiness may occur</li> <li>■ excitability may occur, especially in children</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> </ul> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ new symptoms occur</li> <li>■ redness or swelling is present</li> <li>■ symptoms do not get better</li> <li>■ you need to use more than 7 days</li> <li>■ fever gets worse or lasts more than 3 days</li> </ul>	

**Drug Facts (continued)**

If you are pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions** ■ children under 12 years: ask a doctor  
 ■ adults and children 12 years or over: take 2 tablets every 12 hours; not more than 4 tablets in 24 hours

<b>Drug Facts (continued)</b>
<b>Other information</b>
<ul style="list-style-type: none"> <li>■ store at 20-25°C (68-77°F)</li> <li>■ protect from excessive moisture</li> </ul>
<b>Inactive Ingredients</b> calcium phosphate, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C yellow no. 6 aluminum lake, hydroxypropyl methylcellulose, magnesium stearate, methylparaben, PEG, propylparaben, stearic acid

\* Note: 14 point Helvetica Condensed Bold Italic Title  
 8 point Helvetica Condensed Bold Italic Headings  
 6 point Helvetica Condensed Bold Subheadings  
 6 point Helvetica Condensed Text  
 6.5 point Leading

Ducolax Laxative, 10 count  
 Product Using Section 201.66(d)(10) Modified Format\*  
 [Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredient (in each tablet)</b> Bisacodyl 5 mg	<b>Purpose</b> Laxative
<b>Use</b> ■ relieves occasional constipation (irregularity) ■ this product usually causes bowel movement in 6 to 12 hours	
<b>Warnings</b> Do not use if you cannot swallow without chewing Ask a doctor before use if you have ■ stomach pain, nausea, or vomiting ■ a sudden change in bowel habits that lasts over 2 weeks When using this product ■ you may have stomach discomfort, flatness, or cramps ■ do not chew or crush tablet ■ do not use within 1 hour after taking an antacid or milk Stop use and ask a doctor if ■ you have no bowel movement within 12 hours or rectal bleeding. These could be signs of a serious condition. ■ you need to use a laxative more than 1 week If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	adults and children 12 years and over take 1 to 3 tablets (usually 2) daily children 6 to under 12 years take 1 tablet daily children under 6 years ask a doctor
<b>Other information</b> store at 20-25°C (68-77°F)	
<b>Inactive ingredients</b> acacia, acetylated monoglyceride, carnauba wax, cellulose acetate phthalate, corn starch, D&C red no. 30 aluminum lake, D&C yellow no. 10 aluminum lake, dibutyl phthalate, docusate sodium, gelatin, glycerin, iron oxides, kaolin, lactose, magnesium stearate, methylparaben, pharmaceutical glaze, polyethylene glycol, povidone, propylparaben, sodium benzoate, sorbitan monooleate, sucrose, talc, titanium dioxide, white wax	

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6.25 point Leading

## Goody Headache Powder

Combination Product Using Section 201.66(d)(10) Modified Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredients (in each powder)</b>	<b>Purpose</b>
Aspirin 500mg	Pain reliever
Acetaminophen 260mg	Pain reliever
Caffeine 32.5mg	Pain reliever aid
<b>Use</b> temporarily relieves minor aches and pains due to: <input type="checkbox"/> colds <input type="checkbox"/> headache <input type="checkbox"/> minor arthritis pain	
<b>Warnings</b>	
<p><b>Reye's syndrome:</b> Children and teenagers should not use this drug for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported with aspirin.</p> <p><b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen, aspirin or other pain relievers/fever reducers. Acetaminophen and aspirin may cause liver damage and stomach bleeding.</p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include:  <input type="checkbox"/> hives   <input type="checkbox"/> facial swelling   <input type="checkbox"/> asthma (wheezing)   <input type="checkbox"/> shock</p> <p>Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.</p> <p>Ask a doctor before use if you have   <input type="checkbox"/> asthma   <input type="checkbox"/> ulcers   <input type="checkbox"/> bleeding problems   <input type="checkbox"/> stomach problems that last or come back, such as heartburn, upset stomach, or pain.</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for:  <input type="checkbox"/> diabetes   <input type="checkbox"/> gout   <input type="checkbox"/> arthritis   <input type="checkbox"/> anticoagulation (blood thinning)</p> <p>Stop use and ask a doctor if   <input type="checkbox"/> allergic reaction occurs. Seek medical help right away.   <input type="checkbox"/> pain gets worse or lasts for more than 10 days.  <input type="checkbox"/> redness or swelling is present   <input type="checkbox"/> new symptoms occur  <input type="checkbox"/> ringing in the ears or loss of hearing occurs</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	
<b>Drug Facts (continued)</b>	
<p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b>   <input type="checkbox"/> do not take more than directed  <input type="checkbox"/> adults and children 12 years and over: place 1 powder on tongue every 4 to 6 hours. Follow with liquid. May stir powder into glass of water or other liquid and drink; not more than 4 powders in 24 hours.  <input type="checkbox"/> children under 12 years: ask a doctor</p>	
<b>Inactive ingredients</b> lactose, potassium chloride	

grey is actual size of product

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6 point Leading

# Goody Headache Powder

Combination Product Using Section 201.66(d)(10) Modified Format\*  
[Individual Envelope]

Front

Back

PDP Space

---

**Drug Facts**

Active ingredients (in each powder)	Purpose
Aspirin 500mg	Pain reliever
Acetaminophen 260mg	Pain reliever
Caffeine 32.5mg	Pain reliever aid

Use temporarily relieves minor aches and pains due to:  
 colds    headache    minor arthritis pain

**Drug Facts (continued)**

**Warnings**

**Reye's syndrome:** Children and teenagers should not use this drug for chicken pox or flu symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported with aspirin.

**Alcohol warning:** If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen, aspirin or other pain relievers/fever reducers. Acetaminophen and aspirin may cause liver damage and stomach bleeding.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  hives    facial swelling    asthma (wheezing)    shock

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.

Ask a doctor before use if you have  asthma    ulcers    bleeding problems    stomach problems that last or come back, such as heartburn, upset stomach, or pain.

Ask a doctor or pharmacist before use if you are taking a prescription drug for:  diabetes    gout    arthritis    anticoagulation (blood thinning)

Stop use and ask a doctor if  allergic reaction occurs. Seek medical help right away.  pain gets worse or lasts for more than 10 days    redness or swelling is present    new symptoms occur    ringing in the ears or loss of hearing occurs.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

**Directions**    do not take more than directed

- adults and children 12 years and over: place 1 powder on tongue every 4 to 6 hours. Follow with liquid. May stir powder into glass of water or other liquid and drink not more than 4 powders in 24 hours.
- children under 12 years: ask a doctor

**Inactive ingredients:** lactose, potassium chloride

grey line is actual size of product

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6 point Leading

# Phillips Milk of Magnesium

Product Using Section 201.66(d)(10) Modified Format\*  
 [Immediate Container Label, No Outer Carton]

<b>Drug Facts</b>		
<b>Active ingredient (in each 5 mL)</b>	<b>Purpose</b>	
Magnesium hydroxide 400 mg	Saline laxative/antacid	
<b>Uses</b>		
laxative ■ relieves occasional constipation (irregularity)		
■ this product usually causes bowel movement in 1/2 to 6 hours		
antacid ■ relieves; ■ heartburn ■ acid indigestion ■ sour stomach		
■ upset stomach due to these symptoms		
<b>Warnings</b>		
Ask a doctor before use if you have		
■ kidney disease ■ stomach pain, nausea, or vomiting		
■ a sudden change in bowel habits that lasts over 2 weeks		
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.		
Stop use and ask a doctor if		
laxative ■ you have no bowel movement within 6 hours or rectal bleeding.		
These could be signs of a serious condition.		
■ you need to use a laxative for more than 1 week		
antacid ■ heartburn symptoms last for more than 2 weeks		
■ product has a laxative effect		
If pregnant or breast-feeding, ask a health professional before use.		
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.		
<b>Directions</b> ■ drink a full glass (8 oz) of liquid with each dose		
■ do not take more than recommended daily dosage in a 24 hour period		
■ shake well		
	<b>Laxative</b>	<b>Antacid</b>
adults and children 12 years and over	30 to 60 mL once a day	5 to 15 mL, up to four times a day or as directed by a doctor
children 6 to 11 years	15 to 30 mL once a day	ask a doctor
children 2 to 5 years	5 to 15 mL once a day	ask a doctor
children under 2 years	ask a doctor	ask a doctor
<b>Inactive ingredients</b> purified water		

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6 point Leading



# Nupercainal Ointment, 1 oz.

Product Using Standard Labeling Format\*  
[Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredient</b> Dibucaine 1%	<b>Purpose</b> Hemorrhoidal/topical analgesic ointment
<b>Uses</b> temporarily relieves pain and itching due to: ■ hemorrhoids or other anorectal disorders ■ sunburn ■ minor burns ■ minor cuts ■ minor scrapes ■ insect bites ■ minor skin irritations	
<b>Warnings</b> For external use only When using this product ■ do not use in or near the eyes ■ do not put directly in rectum by using fingers or any mechanical device ■ do not use in large quantities, particularly over raw surfaces or blistered areas	

<b>Drug Facts (continued)</b>		
Stop use and ask a doctor if ■ allergic reaction occurs ■ rectal bleeding occurs ■ condition worsens or does not improve within 7 days ■ symptoms clear up and return within a few days ■ redness, irritation, swelling, pain, or other symptoms begin or increase		
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.		
<b>Directions</b> ■ when practical, clean area with mild soap and warm water and rinse thoroughly ■ gently dry by patting or blotting with toilet tissue or soft cloth before applying	adults and children 12 years and over children 2 years to under 12 years children under 2 years	apply externally to the affected area up to 3 or 4 times daily ask a doctor before use do not use

<b>Drug Facts (continued)</b>	
<b>Inactive ingredients</b> acetone sodium bisulfite, lanolin, light mineral oil, purified water, white petrolatum	

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

# Opcon-A Eyedrops, 0.5 fl. oz.

Combination Product Using Standard Labeling Format\*  
[Outer Carton]

<p><b>Drug Facts</b></p> <p><b>Active Ingredients Purpose</b> Naphazoline HCl 0.02675%...Redness reliever Pheniramine maleate 0.315%...Antihistamine</p> <p><b>Use</b> temporarily relieves itchy, red eyes caused by: ■ pollen ■ ragweed ■ animal hair ■ grass ■ dander</p> <p><b>Warnings</b> Ask a doctor before use if you have ■ high blood pressure ■ heart disease ■ narrow angle glaucoma ■ trouble urinating due to an enlarged prostate gland</p> <p><b>When using this product</b> ■ do not touch tip of container to any surface to avoid contamination ■ replace cap after each use ■ do not use if the liquid changes color or becomes cloudy ■ remove contact lenses before using ■ overuse may cause more eye redness</p> <p><b>Stop use and ask a doctor if</b> ■ you feel eye pain ■ changes in vision occur ■ redness or irritation of the eyes lasts ■ condition worsens or lasts more than 72 hours</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b> ■ adults and children 6 years and over: put 1 or 2 drops in affected eye(s) up to 4 times daily ■ children under 6 years: ask a doctor</p> <p><b>Other Information</b> ■ store at 20-25°C (68-77°F) ■ protect from light</p> <p><b>Inactive Ingredients</b> benzalkonium chloride, edetate disodium, hydroxypropyl methylcellulose, purified water with a boric acid, sodium borate buffer system, and sodium chloride</p>
--	--

- \* Note: 14 point Helvetica Condensed Bold Italic Title
- 8 point Helvetica Condensed Bold Italic Headings
- 6 point Helvetica Condensed Bold Subheadings
- 6 point Helvetica Condensed Text
- 6.5 point Leading

Ornex Maximum Strength, 24 caplets  
 Combination Product Using Standard Labeling Format\*  
 [Outer Carton]

<b>Drug Facts</b>	
<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 500 mg	Pain reliever/fever reducer
Pseudoephedrine hydrochloride 30 mg	Nasal decongestant
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ temporarily relieves nasal congestion due to the common cold               <ul style="list-style-type: none"> <li>■ promotes nasal and/or sinus drainage</li> <li>■ relieves sinus congestion and pressure</li> </ul> </li> <li>■ temporarily relieves               <ul style="list-style-type: none"> <li>■ minor aches and pains</li> <li>■ minor sore throat pain</li> </ul> </li> <li>■ temporarily reduces fever due to colds or flu</li> </ul>	
<b>Warnings</b>	
<p><b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.</p> <p><b>Sore throat warning:</b> Severe or persistent sore throat or sore throat that occurs with high fever, headache, nausea, and vomiting may be serious. Ask a doctor right away. Do not use more than 2 days or give to children under 3 years of age unless directed by a doctor.</p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul>	
<b>When using this product</b>	
<ul style="list-style-type: none"> <li>■ do not use more than directed</li> </ul>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ redness or swelling is present</li> <li>■ you need to use more than 7 days</li> <li>■ new symptoms occur</li> <li>■ symptoms do not get better</li> <li>■ fever gets worse or lasts more than 3 days</li> </ul>	
<p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>	

**Drug Facts (continued)**

**Directions**

- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets every 24 hours
- children under 12 years: ask a doctor

**Inactive ingredients** croscarmellose, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyacrylate 40, povidone, starch, titanium dioxide

\* Note: 14 point Helvetica Condensed Bold Italic Title  
 8 point Helvetica Condensed Bold Italic Headings  
 6 point Helvetica Condensed Bold Subheadings  
 6 point Helvetica Condensed Text  
 6.5 point Leading

## Sucrets Oral Anesthetic

Product Using Section 201.66(d)(10) Modified Format\*  
[Immediate Container]

<b>Drug Facts</b>	
<b>Active Ingredient (in each lozenge)</b>	<b>Purpose</b>
Hexylresorcinol 2.4 mg	Oral anesthetic
<b>Use</b> temporarily relieves occasional: ■ minor irritation ■ pain ■ sore mouth ■ sore throat	
<b>Warnings</b> Sore throat warning: Severe or persistent sore throat or sore throat that occurs with high fever, headache, nausea, and vomiting may be serious. Ask a doctor right away. Do not use more than 2 days or give to children under 3 years of age unless directed by a doctor.	
Stop use and ask a doctor if ■ sore throat is severe or irritation, pain or redness lasts and worsens ■ sore mouth does not improve in 7 days	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ adults and children 3 years and older: dissolve 1 lozenge slowly in the mouth. May be repeated every 2 hours as needed or as directed by a doctor. ■ children under 3 years: ask a doctor	
<b>Inactive Ingredients</b> corn syrup, D&C blue 1, D&C yellow 10, mineral oil, silicon dioxide, sucrose	

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6 point Leading

Selsun Blue 4 oz.

Product Using Section 201.66(d)(10) Modified Format\*  
[Irregular Shape Bottle Label, No Outer Carton]

<i>Drug Facts</i>	
<i>Active ingredient</i>	<i>Purpose</i>
Selenium sulfide 1%	Antidandruff seborrheic dermatitis shampoo
<i>Use</i> controls scalp itching and flaking due to dandruff and seborrheic dermatitis	
<i>Warnings</i> For external use only	
Ask a doctor before use if you have	
■ seborrheic dermatitis that covers a large area of the body	
When using this product	
■ do not get into eyes. If contact occurs, rinse eyes thoroughly with water.	
Stop use and ask a doctor if	
■ condition worsens or does not improve after regular use	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
<i>Directions</i> ■ shake well	
■ for best results, use at least 2 times a week.	
<i>Inactive ingredients</i> ammonium lauryl sulfate, ammonium lauryl sulfate, citric acid, cocamide DEA, cocamidopropyl betaine, DMDM hydantoin, D&C red no. 33, FD&C blue no. 1, hydroxypropyl methylcellulose, magnesium aluminum silicate, menthol, purified water, sodium chloride, TEA-lauryl sulfate	

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6 point Leading

Tylenol Extra Strength Caplets 10 count  
Product Using Section 201.66(d)(10) Modified Format\*  
[Tube With Wraparound Label]

<b>Drug Facts</b>	
<b>Active ingredient (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 500mg	Pain reliever/fever reducer
<b>Uses</b> temporarily relieves minor aches and pains relieves fever	
<b>Warnings</b>	
Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.	
Stop use and ask a doctor if new symptoms occur	
redness or swelling is present pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days	
if pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults/children even if you do not notice any signs/symptoms.	
<b>Directions</b> do not take more than directed adults and children over 12 years: take 2 caplets every 6 hours as needed; not more than 8 caplets in 24 hours children under 12 years: do not use	
<b>Inactive ingredients</b> cellulose, corn starch, FD&C red no. 40, hydroxypropyl methylcellulose, magnesium stearate, polyethylene glycol, sodium starch glycolate	

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6.25 point Leading

# Vicks Nyquil Liquicaps Multi-symptom Cold/Flu, 12 softgel

Combination Product Using Standard Labeling Format\*

[Outer Carton]

<p><b>Drug Facts (continued)</b></p> <p><b>Inactive Ingredients:</b> D&amp;C yellow no. 10, FD&amp;C blue no. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special</p>	<p><b>Drug Facts</b></p> <table border="1"> <thead> <tr> <th>Active Ingredients (in each softgel)</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Acetaminophen 500 mg</td> <td>Pain reliever/fever reducer</td> </tr> <tr> <td>Dextromethorphan HBr 15 mg</td> <td>Cough suppressant</td> </tr> <tr> <td>Doxylamine succinate 6.25 mg</td> <td>Antihistamine</td> </tr> <tr> <td>Pseudoephedrine HCl 30 mg</td> <td>Nasal decongestant</td> </tr> </tbody> </table> <p><b>Uses</b> ■ temporarily relieves these cold and flu symptoms:</p> <ul style="list-style-type: none"> <li>■ cough</li> <li>■ sore throat pain</li> <li>■ minor aches and pains</li> <li>■ headache</li> <li>■ stuffy nose</li> <li>■ muscular aches</li> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Alcohol warning:</b> If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.</p> <p><b>Sore throat warning:</b> Severe or persistent sore throat or sore throat that occurs with high fever, headache, nausea, and vomiting may be serious. Ask a doctor right away. Do not use more than 2 days or give to children under 3 years of age unless directed by a doctor.</p> <p><b>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI)</b> (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ glaucoma</li> <li>■ diabetes</li> <li>■ thyroid disease</li> <li>■ cough that occurs with too much phlegm (mucus)</li> <li>■ high blood pressure</li> <li>■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> <li>■ trouble urinating due to an enlarged prostate gland</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</b></p> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ do not use more than directed</li> <li>■ excitability may occur, especially in children</li> <li>■ marked drowsiness may occur</li> <li>■ avoid alcoholic drinks</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ redness or swelling is present</li> <li>■ new symptoms occur</li> <li>■ you need to use more than 7 days</li> <li>■ symptoms do not get better</li> <li>■ fever gets worse or lasts more than 3 days</li> <li>■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.</li> </ul>	Active Ingredients (in each softgel)	Purpose	Acetaminophen 500 mg	Pain reliever/fever reducer	Dextromethorphan HBr 15 mg	Cough suppressant	Doxylamine succinate 6.25 mg	Antihistamine	Pseudoephedrine HCl 30 mg	Nasal decongestant	<p><b>Drug Facts (continued)</b></p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>
Active Ingredients (in each softgel)	Purpose											
Acetaminophen 500 mg	Pain reliever/fever reducer											
Dextromethorphan HBr 15 mg	Cough suppressant											
Doxylamine succinate 6.25 mg	Antihistamine											
Pseudoephedrine HCl 30 mg	Nasal decongestant											
<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ adults and children 12 years and over: take 2 softgels with water every 6 hours; not more than 4 doses daily.</li> <li>■ children under 12 years: do not use</li> </ul>												

- \* Note: 14 point Helvetica Condensed Bold Italic Title  
 8 point Helvetica Condensed Bold Italic Headings  
 6 point Helvetica Condensed Bold Subheadings  
 6 point Helvetica Condensed Text  
 6.5 point Leading

# Coppertone Kids Sunscreen

Product Using Section 201.66(d)(10) Modified Format\*  
 [Immediate Container Label, No Outer Carton]

**Drug Facts**

Active Ingredients	Purpose
Methoxycinnamate 7%	Sunscreen
2-Ethylhexyl Salicylate 7%	
Oxybenzone 7%	
Homosalate 7%	

**Use** to help prevent sunburn

**Warnings**  
 For external use only  
 When using this product do not use in or near the eyes  
 if product gets into eyes, rinse thoroughly with water  
 Stop use and ask a doctor if irritation or rash appears and lasts  
 Keep out of reach of children. If swallowed, contact a Poison Control Center or get medical help right away.

**Directions** in children under 2 years of age should use sunscreen products with a minimum SPF of 4  
 in adults and children 6 months and over: apply liberally 15 minutes before sun or water exposure; reapply after 80 minutes of swimming or excessive sweating or anytime after towel drying  
 in children under 6 months: ask a doctor

**Other Information** Recommended Sunscreen Product Guide

Sunburn and tanning history	Recommended sun protection product
Always burns easily; rarely tans	SPF 20 to 30
Always burns easily; tans minimally	SPF 12 to under 20
Burns moderately; tans gradually	SPF 8 to under 12
Burns minimally; always tans well	SPF 4 to under 8
Rarely burns; tans profusely	SPF 2 to under 4

Sun alert: The sun causes skin damage. Regular use of sunscreens over the years may reduce the chance of skin damage, some types of skin cancer, and other harmful effects due to the sun.

**Inactive ingredients** water, sorbitan isostearate, squalene, polyglyceryl-3 distearate, octadecene MA, copolymer, triethanolamine, stearic acid, barium sulfate, benzyl alcohol fragrance, dimethicone, aloe extract, jojoba oil, methylparaben, tocopherol (vitamin E), propylparaben, carbomer, disodium EDTA, imidazolidinyl urea

grey line is actual size of product

- \* Note: 9 point Helvetica Narrow Bold Italic Title
- 8 point Helvetica Narrow Bold Italic Headings
- 6 point Helvetica Narrow Bold Subheadings
- 6 point Helvetica Narrow Text
- 6 point Leading