



Complete Summary

GUIDELINE TITLE

Screening for congenital hypothyroidism: U.S. Preventive Services Task Force reaffirmation recommendation statement.

BIBLIOGRAPHIC SOURCE(S)

U.S. Preventive Services Task Force (USPSTF). Screening for congenital hypothyroidism: U.S. Preventive Services Task Force reaffirmation recommendation statement. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2008 Mar. 6 p. [5 references]

US Preventive Services Task Force. Screening for congenital hypothyroidism: US Preventive Services Task Force reaffirmation recommendation. Ann Fam Med 2008 Mar-Apr;6(2):166. [PubMed](#)

GUIDELINE STATUS

This is the current release of the guideline.

This release updates a previously published guideline: U.S. Preventive Services Task Force. Guide to clinical preventive services. 2nd ed. Baltimore (MD): Williams & Wilkins; 1996. Chapter 45, Screening for congenital hypothyroidism. p. 503-507. [39 references]

COMPLETE SUMMARY CONTENT

SCOPE
METHODOLOGY - including Rating Scheme and Cost Analysis
RECOMMENDATIONS
EVIDENCE SUPPORTING THE RECOMMENDATIONS
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
QUALIFYING STATEMENTS
IMPLEMENTATION OF THE GUIDELINE
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
CATEGORIES
IDENTIFYING INFORMATION AND AVAILABILITY
DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Congenital hypothyroidism

GUIDELINE CATEGORY

Prevention
Screening

CLINICAL SPECIALTY

Family Practice
Medical Genetics
Nursing
Obstetrics and Gynecology
Pediatrics
Preventive Medicine

INTENDED USERS

Advanced Practice Nurses
Allied Health Personnel
Health Care Providers
Nurses
Physician Assistants
Physicians

GUIDELINE OBJECTIVE(S)

- To summarize the U.S. Preventive Services Task Force (USPSTF) recommendations and supporting scientific evidence on screening for congenital hypothyroidism
- To reaffirm the 1996 USPSTF recommendations on screening for congenital hypothyroidism

TARGET POPULATION

All newborn infants

INTERVENTIONS AND PRACTICES CONSIDERED

Screening for congenital hypothyroidism by serum thyroxine (T₄) and/or thyroid-stimulating hormone (TSH) performed on capillary blood collected from a heel stick and adsorbed onto filter paper

MAJOR OUTCOMES CONSIDERED

- Benefits of screening for congenital hypothyroidism
- Potential harms of screening for congenital hypothyroidism

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Primary Sources)
Hand-searches of Published Literature (Secondary Sources)
Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Note from the National Guideline Clearinghouse (NGC): A targeted review of the literature was prepared by Agency for Healthcare Research and Quality (AHRQ) staff for use by the U.S. Preventive Services Task Force (USPSTF) (see the "Availability of Companion Documents" field).

In 2006, the USPSTF decided to update its recommendation statement on screening for congenital hypothyroidism. Noting that the 1996 recommendation was made on a strong evidence base and that it would take large, high-quality studies or evidence of substantial harms to overturn the current recommendation, the USPSTF chose to perform a reaffirmation update for this topic. The USPSTF performs reaffirmation updates for older recommendation statements that remain USPSTF priorities, are within the scope of the USPSTF, and for which there is compelling reason for the USPSTF to have a current recommendation statement.

To assist the USPSTF in updating its 1996 recommendation on screening for congenital hypothyroidism, staff at AHRQ performed a targeted search of the medical literature from January 1, 1995, to September 14, 2006, and consulted with subject area experts. The goal of this targeted review was to find new, high-quality evidence regarding the benefits and potential harms of screening for congenital hypothyroidism. Thirty-eight studies were initially identified. No studies on benefits and three studies relating to potential harms were found to meet review inclusion criteria.

NUMBER OF SOURCE DOCUMENTS

Three studies relating to potential harms were found to meet review inclusion criteria.

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Expert Consensus

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Not applicable

METHODS USED TO ANALYZE THE EVIDENCE

Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

The U.S. Preventive Services Task Force (USPSTF) systematically reviews the evidence concerning both the benefits and harms of widespread implementation of a preventive service. It then assesses the certainty of the evidence and the magnitude of the benefits and harms. On the basis of this assessment, the USPSTF assigns a letter grade to each preventive service signifying its recommendation about provision of the service (see Table below). An important, but often challenging, step is determining the balance between benefits and harms to estimate "net benefit" (that is, benefits minus harms).

Table 1. U.S. Preventive Services Task Force Recommendation Grid*

Certainty of Net Benefit	Magnitude of Net Benefit			
	Substantial	Moderate	Small	Zero/Negative
High	A	B	C	D
Moderate	B	B	C	D
Low	Insufficient			

*A, B, C, D, and *Insufficient* represent the letter grades of recommendation or statement of insufficient evidence assigned by the U.S. Preventive Services Task Force after assessing certainty and magnitude of net benefit of the service (see the "Rating Scheme for the Strength of the Recommendations" field).

The overarching question that the Task Force seeks to answer for every preventive service is whether evidence suggests that provision of the service would improve health outcomes if implemented in a general primary care population. For screening topics, this standard could be met by a large randomized, controlled trial (RCT) in a representative asymptomatic population with follow-up of all members of both the group "invited for screening" and the group "not invited for screening."

Direct RCT evidence about screening is often unavailable so when the Task Force undertakes a full systematic review, it considers indirect evidence. To guide its selection of indirect evidence, the Task Force constructs a "chain of evidence" within an analytic framework. Each arrow in the framework defines a key question, and each key question represents a link in the chain of evidence. Rectangles in the framework represent the intermediate outcomes (rounded corners) or the health outcomes (square corners); ovals represent harms. To form an unbroken chain, evidence must support each link in the chain, thereby connecting the target population (far left side of the framework) to the improved health outcome (far right side of the framework). For each key question, the body of pertinent literature is critically appraised, focusing on the following 6 questions:

1. Do the studies have the appropriate research design to answer the key question(s)?

2. To what extent are the existing studies of high quality? (i.e., what is the internal validity?)
3. To what extent are the results of the studies generalizable to the general U.S. primary care population and situation? (i.e., what is the external validity?)
4. How many studies have been conducted that address the key question(s)? How large are the studies? (i.e., what is the precision of the evidence?)
5. How consistent are the results of the studies?
6. Are there additional factors that assist us in drawing conclusions (e.g., presence or absence of dose–response effects, fit within a biologic model)?

The next step in the Task Force process is to use the evidence from the key questions to assess whether there would be net benefit if the service were implemented. In 2001, the USPSTF published an article that documented its systematic processes of evidence evaluation and recommendation development. At that time, the Task Force's overall assessment of evidence was described as good, fair, or poor. The Task Force realized that this rating seemed to apply only to how well studies were conducted and did not fully capture all of the issues that go into an overall assessment of the evidence about net benefit. To avoid confusion, the USPSTF has changed its terminology. Whereas individual study quality will continue to be characterized as good, fair, or poor, the term *certainty* will now be used to describe the Task Force's assessment of the overall body of evidence about net benefit of a preventive service and the likelihood that the assessment is correct. Certainty will be determined by considering all 6 questions listed above; the judgment about certainty will be described as high, moderate, or low.

In making its assessment of certainty about net benefit, the evaluation of the evidence from each key question plays a primary role. It is important to note that the Task Force makes recommendations for real-world medical practice in the United States and must determine to what extent the evidence for each key question—even evidence from screening RCTs or treatment RCTs—can be applied to the general primary care population. Frequently, studies are conducted in highly selected populations under special conditions. The Task Force must consider differences between the general primary care population and the populations studied in RCTs and make judgments about the likelihood of observing the same effect in actual practice.

It is also important to note that 1 of the key questions in the analytic framework refers to the potential harms of the preventive service. The Task Force considers the evidence about the benefits and harms of preventive services separately and equally. Data about harms are often obtained from observational studies because harms observed in RCTs may not be representative of those found in usual practice and because some harms are not completely measured and reported in RCTs.

Putting the body of evidence for all key questions together as a chain, the Task Force assesses the certainty of net benefit of a preventive service by asking the 6 major questions listed above. The Task Force would rate a body of convincing evidence about the benefits of a service that, for example, derives from several RCTs of screening in which the estimate of benefits can be generalized to the general primary care population as "high" certainty (see the "Rating Scheme for the Strength of Recommendations" field). The Task Force would rate a body of

evidence that was not clearly applicable to general practice or has other defects in quality, research design, or consistency of studies as "moderate" certainty. Certainty is "low" when, for example, there are gaps in the evidence linking parts of the analytic framework, when evidence to determine the harms of treatment is unavailable, or when evidence about the benefits of treatment is insufficient. Table 4 in the methodology document listed below (see "Availability of Companion Documents" field) summarizes the current terminology used by the Task Force to describe the critical assessment of evidence at all 3 levels: individual studies, key questions, and overall certainty of net benefit of the preventive service.

Some clinical preventive services have a strong, well-established evidence base and are a routine part of clinical practice. Because it is unlikely that new evidence will change USPSTF recommendations for such services, the USPSTF reviews the evidence for them in an expedited manner by conducting literature searches that address benefits and harms and consulting experts. The reaffirmation recommendation statement on Screening for Congenital Hypothyroidism was developed on the basis of such an expedited literature review.

Sources

1. Sawaya GF et al., Update on the methods of the U.S. Preventive Services Task Force: estimating certainty and magnitude of net benefit. *Ann Intern Med.* 2007;147:871-875. [5 references].
2. Guirguis-Blake J, et al. Current processes of the U.S. Preventive Services Task Force: refining evidence-based recommendation development. *Ann Intern Med.* 2007;147:117-122. [2 references]

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

What the United States Preventive Services Task Force (USPSTF) Grades Mean and Suggestions for Practice

Grade	Grade Definitions	Suggestions for Practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
B	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
C	The USPSTF recommends against routinely providing the service. There may be considerations that support providing the service in an individual patient. There is moderate or high certainty that the net benefit is small.	Offer/provide this service only if there are other considerations in support of the offering/providing the service in an individual patient.
D	The USPSTF recommends against the service. There is moderate or high certainty that the service has	Discourage the use of this service.

Grade	Grade Definitions	Suggestions for Practice
	no net benefit or that the harms outweigh the benefits.	
I Statement	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality or conflicting, and the balance of benefits and harms cannot be determined.	Read "Clinical Considerations" section of USPSTF Recommendation Statement (see "Major Recommendations" field). If offered, patients should understand the uncertainty about the balance of benefits and harms.

USPSTF Levels of Certainty Regarding Net Benefit

Definition: The U.S. Preventive Services Task Force defines certainty as "likelihood that the USPSTF assessment of the net benefit of a preventive service is correct." The net benefit is defined as benefit minus harm of the preventive service as implemented in a general, primary care population. The USPSTF assigns a certainty level based on the nature of the overall evidence available to assess the net benefit of a preventive service.

Level of Certainty	Description
High	The available evidence usually includes consistent results from well-designed, well-conducted studies in representative primary care populations. These studies assess the effects of the preventive service on health outcomes. This conclusion is therefore unlikely to be strongly affected by the results of future studies.
Moderate	<p>The available evidence is sufficient to determine the effects of the preventive service on health outcomes, but confidence in the estimate is constrained by factors such as:</p> <ul style="list-style-type: none"> • The number, size, or quality of individual studies • Inconsistency of findings across individual studies • Limited generalizability of findings to routine primary care practice • Lack of coherence in the chain of evidence <p>As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion.</p>
Low	<p>The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of:</p> <ul style="list-style-type: none"> • The limited number or size of studies • Important flaws in study design or methods • Inconsistency of findings across individual studies • Gaps in the chain of evidence • Findings not generalizable to routine primary care practice • A lack of information on important health outcomes

Level of Certainty	Description
	More information may allow an estimation of effects on health outcomes.

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Comparison with Guidelines from Other Groups
 External Peer Review
 Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Peer Review. Before the U.S. Preventive Services Task Force makes its final determinations about recommendations on a given preventive service, the Evidence-Based Practice Center and the Agency for Healthcare Research and Quality send a draft evidence review to 4 to 6 external experts and to federal agencies and professional and disease-based health organizations with interests in the topic. They ask the experts to examine the review critically for accuracy and completeness and to respond to a series of specific questions about the document. After assembling these external review comments and documenting the proposed response to key comments, the topic team presents this information to the Task Force in memo form. In this way, the Task Force can consider these external comments before it votes on its recommendations about the service. Draft recommendation statements are then circulated for comment from reviewers representing professional societies, voluntary organizations and Federal agencies. These comments are discussed before the final recommendations are confirmed.

Comparison with Guidelines from Other Groups. Recommendations for screening from the following groups were discussed: the European Society of Paediatric Endocrinology, the American Academy of Pediatrics, the American Academy of Family Physicians, the American Thyroid Association, and the Lawson Wilkins Pediatric Endocrine Society.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

The US Preventive Services Task Force (USPSTF) grades its recommendations (A, B, C, D, or I) and identifies the Levels of Certainty regarding Net Benefit (High, Moderate, and Low). The definitions of these grades can be found at the end of the "Major Recommendations" field.

Summary of Recommendation and Evidence

The U. S. Preventive Services Task Force (USPSTF) recommends screening for congenital hypothyroidism (CH) in newborns. **This is a grade A recommendation.**

Clinical Considerations

Patient Population under Consideration

This recommendation applies all infants born in the U.S. Premature, very low birth weight and ill infants may benefit from additional screening because these conditions are associated with decreased sensitivity and specificity of screening tests.

Screening Tests

Screening for CH is mandated in all 50 states and the District of Columbia, though methods of screening vary. There are two main methods used in the U.S.: Primary thyroid-stimulating hormone (TSH) with backup serum thyroxine (T_4); and primary T_4 with backup TSH. A few states use both tests in initial screening. Clinicians should become familiar with the tests used in their area and the limitations of the employed screening strategy. For example, a primary TSH method may be falsely negative in low and very low birth weight infants with CH because of delayed elevation in TSH. Additionally, few states currently screen for centrally-mediated congenital hypothyroidism. Families should be provided with appropriate information about newborn screening tests, including the benefits and harms of screening. They should be aware of the potential of a false positive test, and the process required for definitive testing. Nationally, only 1 in 25 positive screening tests are confirmed to be CH. Normal newborn screening results for CH should not preclude appropriate evaluation of infants presenting with clinical signs and symptoms suggestive of hypothyroidism.

Timing of Screening

Infants should be tested between 2 and 4 days of age. Infants discharged from hospitals before 48 hours of life should be tested immediately before discharge. Specimens obtained in the first 24-48 hours of age may be falsely elevated for TSH regardless of the screening method used.

Treatment

Primary care clinicians should ensure that infants with abnormal screens receive confirmatory testing and begin appropriate treatment with thyroid hormone replacement within 2 weeks after birth. Children with positive confirmatory testing in whom no permanent cause of CH is found (such as lack of thyroid tissue on thyroid ultrasound or thyroid scan), should, at some time point after the age of 3 years, undergo a 30-day trial of reduced or discontinued thyroid hormone replacement therapy to determine if the hypothyroidism is permanent or transient.

Definitions:

What the United States Preventive Services Task Force (USPSTF) Grades Mean and Suggestions for Practice

Grade	Grade Definitions	Suggestions for Practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
B	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
C	The USPSTF recommends against routinely providing the service. There may be considerations that support providing the service in an individual patient. There is moderate or high certainty that the net benefit is small.	Offer/provide this service only if there are other considerations in support of the offering/providing the service in an individual patient.
D	The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.
I Statement	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality or conflicting, and the balance of benefits and harms cannot be determined.	Read "Clinical Considerations" section of USPSTF Recommendation Statement (see "Major Recommendations" field). If offered, patients should understand the uncertainty about the balance of benefits and harms.

USPSTF Levels of Certainty Regarding Net Benefit

Definition: The U.S. Preventive Services Task Force defines certainty as "likelihood that the USPSTF assessment of the net benefit of a preventive service is correct." The net benefit is defined as benefit minus harm of the preventive service as implemented in a general, primary care population. The USPSTF assigns a certainty level based on the nature of the overall evidence available to assess the net benefit of a preventive service.

Level of Certainty	Description
High	The available evidence usually includes consistent results from well-designed, well-conducted studies in representative primary care populations. These studies assess the effects of the preventive service on health outcomes. This conclusion is therefore unlikely to be strongly affected by the results of future studies.
Moderate	The available evidence is sufficient to determine the effects of the preventive service on health outcomes, but confidence in the estimate is

Level of Certainty	Description
	<p>constrained by factors such as:</p> <ul style="list-style-type: none"> • The number, size, or quality of individual studies • Inconsistency of findings across individual studies • Limited generalizability of findings to routine primary care practice • Lack of coherence in the chain of evidence <p>As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion.</p>
Low	<p>The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of:</p> <ul style="list-style-type: none"> • The limited number or size of studies • Important flaws in study design or methods • Inconsistency of findings across individual studies • Gaps in the chain of evidence • Findings not generalizable to routine primary care practice • A lack of information on important health outcomes <p>More information may allow an estimation of effects on health outcomes.</p>

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is not specifically stated for each recommendation.

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Benefits of Detection and Early Intervention

Early detection of congenital hypothyroidism (CH) by neonatal screening and appropriate treatment substantially improves neurodevelopmental outcomes for affected persons.

POTENTIAL HARMS

Harms of Detection and Early Treatment

Positive test results, whether true positive or false positive, cause anxiety in parents. For some parents, this anxiety may be considerable.

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

- The U.S. Preventive Services Task Force (USPSTF) makes recommendations about preventive care services for patients without recognized signs or symptoms of the target condition.
- Recommendations are based on a systematic review of the evidence of the benefits and harms and an assessment of the net benefit of the service.
- The USPSTF recognizes that clinical or policy decisions involve more considerations than this body of evidence alone. Clinicians and policy-makers should understand the evidence but individualize decision making to the specific patient or situation.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

The experiences of the first and second U.S. Preventive Services Task Force (USPSTF), as well as that of other evidence-based guideline efforts, have highlighted the importance of identifying effective ways to implement clinical recommendations. Practice guidelines are relatively weak tools for changing clinical practice when used in isolation. To effect change, guidelines must be coupled with strategies to improve their acceptance and feasibility. Such strategies include enlisting the support of local opinion leaders, using reminder systems for clinicians and patients, adopting standing orders, and audit and feedback of information to clinicians about their compliance with recommended practice.

In the case of preventive services guidelines, implementation needs to go beyond traditional dissemination and promotion efforts to recognize the added patient and clinician barriers that affect preventive care. These include clinicians' ambivalence about whether preventive medicine is part of their job, the psychological and practical challenges that patients face in changing behaviors, lack of access to health care or of insurance coverage for preventive services for some patients, competing pressures within the context of shorter office visits, and the lack of organized systems in most practices to ensure the delivery of recommended preventive care.

Dissemination strategies have changed dramatically in this age of electronic information. While recognizing the continuing value of journals and other print formats for dissemination, the Agency for Healthcare Research and Quality will make all U.S. Preventive Services Task Force (USPSTF) products available through its [Web site](#). The combination of electronic access and extensive material in the public domain should make it easier for a broad audience of users to access U.S. Preventive Services Task Force materials and adapt them for their local needs. Online access to U.S. Preventive Services Task Force products also opens up new

possibilities for the appearance of the annual, pocket-size *Guide to Clinical Preventive Services*.

To be successful, approaches for implementing prevention have to be tailored to the local level and deal with the specific barriers at a given site, typically requiring the redesign of systems of care. Such a systems approach to prevention has had notable success in established staff-model health maintenance organizations, by addressing organization of care, emphasizing a philosophy of prevention, and altering the training and incentives for clinicians. Staff-model plans also benefit from integrated information systems that can track the use of needed services and generate automatic reminders aimed at patients and clinicians, some of the most consistently successful interventions. Information systems remain a major challenge for individual clinicians' offices, however, as well as for looser affiliations of practices in network-model managed care and independent practice associations, where data on patient visits, referrals, and test results are not always centralized.

IMPLEMENTATION TOOLS

Foreign Language Translations
Patient Resources
Personal Digital Assistant (PDA) Downloads
Pocket Guide/Reference Cards
Tool Kits

For information about [availability](#), see the "Availability of Companion Documents" and "Patient Resources" fields below.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Staying Healthy

IOM DOMAIN

Effectiveness
Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

U.S. Preventive Services Task Force (USPSTF). Screening for congenital hypothyroidism: U.S. Preventive Services Task Force reaffirmation recommendation statement. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2008 Mar. 6 p. [5 references]

US Preventive Services Task Force. Screening for congenital hypothyroidism: US Preventive Services Task Force reaffirmation recommendation. *Ann Fam Med* 2008 Mar-Apr;6(2):166. [PubMed](#)

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

1996 (revised 2008 Mar)

GUIDELINE DEVELOPER(S)

United States Preventive Services Task Force - Independent Expert Panel

GUIDELINE DEVELOPER COMMENT

The U.S. Preventive Services Task Force (USPSTF) is a federally-appointed panel of independent experts. Conclusions of the U.S. Preventive Services Task Force do not necessarily reflect policy of the U.S. Department of Health and Human Services (DHHS) or its agencies.

SOURCE(S) OF FUNDING

United States Government

GUIDELINE COMMITTEE

U.S. Preventive Services Task Force

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

*Task Force Members**: Ned Calonge, MD, MPH, Chair, USPSTF (Chief Medical Officer and State Epidemiologist, Colorado Department of Public Health and Environment, Denver, CO); Diana B. Petitti, MD, MPH, Vice-chair, USPSTF (Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Sierra Madre, CA); Thomas G. DeWitt, MD (Carl Weihl Professor of Pediatrics and Director of the Division of General and Community Pediatrics, Department of Pediatrics, Children's Hospital Medical Center, Cincinnati, OH); Leon Gordis, MD, MPH, DrPH (Professor, Epidemiology Department, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD); Kimberly D. Gregory, MD, MPH (Director, Women's Health Services Research and Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Cedars-Sinai Medical Center, Los Angeles, CA); Russell Harris, MD, MPH (Professor of Medicine, Sheps Center for Health Services Research, University of North Carolina School of Medicine, Chapel Hill, NC); George Isham, MD, MS (Medical Director and Chief Health Officer, HealthPartners, Minneapolis, MN); Michael L. LeFevre, MD, MSPH (Professor, Department of Family and Community Medicine, University of Missouri School of Medicine, Columbia, MO); Carol Loveland-Cherry, PhD, RN (Executive Associate Dean, Office of Academic Affairs, University of Michigan School of

Nursing, Ann Arbor, MI); Lucy N. Marion, PhD, RN (Dean and Professor, School of Nursing, Medical College of Georgia, Augusta, GA); Virginia A. Moyer, MD, MPH (Professor, Department of Pediatrics, University of Texas Health Science Center, Houston, TX); Judith K. Ockene, PhD (Professor of Medicine and Chief of Division of Preventive and Behavioral Medicine, University of Massachusetts Medical School, Worcester, MA); George F. Sawaya, MD (Associate Professor, Department of Obstetrics, Gynecology, and Reproductive Sciences and Department of Epidemiology and Biostatistics, University of California, San Francisco, CA); Albert L. Siu, MD, MSPH (Professor and Chairman, Brookdale Department of Geriatrics and Adult Development, Mount Sinai Medical Center, New York, NY); Steven M. Teutsch, MD, MPH (Executive Director, Outcomes Research and Management, Merck & Company, Inc., West Point, PA); and Barbara P. Yawn, MD, MSPH, MSc (Director of Research, Olmstead Medical Center, Rochester, MN)

**Members of the Task Force at the time this recommendation was finalized. For a list of current Task Force members, go to www.ahrq.gov/clinic/uspstfab.htm.*

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

The U.S. Preventive Services Task Force has an explicit policy concerning conflict of interest. All members disclose at each meeting if they have a significant financial, professional/business, or intellectual conflict for each topic being discussed. Task Force members with conflicts may be recused from discussing or voting on recommendations about the topic in question.

GUIDELINE STATUS

This is the current release of the guideline.

This release updates a previously published guideline: U.S. Preventive Services Task Force. Guide to clinical preventive services. 2nd ed. Baltimore (MD): Williams & Wilkins; 1996. Chapter 45, Screening for congenital hypothyroidism. p. 503-507. [39 references]

GUIDELINE AVAILABILITY

Electronic copies: Available from the [U.S. Preventive Services Task Force \(USPSTF\) Web site](http://www.ahrq.gov/clinic/uspstf.htm).

Print copies: Available from the Agency for Healthcare Research and Quality (AHRQ) Publications Clearinghouse. For more information, go to <http://www.ahrq.gov/news/pubsix.htm> or call 1-800-358-9295 (U.S. only).

AVAILABILITY OF COMPANION DOCUMENTS

The following are available:

Evidence Reviews:

- Meyers D, Haering S. Screening for congenital hypothyroidism in newborns: a literature update for the U.S. Preventive Services Task Force. AHRQ

Publication No. 08-05109-EF-1. Rockville, Maryland: Agency for Healthcare Research and Quality, March 2008. Electronic copies: Available from the [U.S. Preventive Services Task Force \(USPSTF\) Web site](#).

- Screening for congenital hypothyroidism: clinical summary of U.S. Preventive Services Task Force recommendations. 2008. Electronic copies: Available in Portable Document Format (PDF) from the U.S. Preventive Services Task Force (USPSTF) Web site. Electronic copies: Available from the [U.S. Preventive Services Task Force \(USPSTF\) Web site](#).

Background Articles:

- Barton M et al. How to read the new recommendation statement: methods update from the U.S. Preventive Services Task Force. *Ann Intern Med*. 2007;147:123-127.
- Guirguis-Blake J et al. Current processes of the U.S. Preventive Services Task Force: refining evidence-based recommendation development. *Ann Intern Med*. 2007;147:117-122. [2 references]
- Sawaya GF et al., Update on the methods of the U.S. Preventive Services Task Force: estimating certainty and magnitude of net benefit. *Ann Intern Med*. 2007;147:871-875. [5 references].

Electronic copies: Available from [U.S. Preventive Services Task Force \(USPSTF\) Web site](#).

The following is also available:

- The guide to clinical preventive services, 2007. Recommendations of the U.S. Preventive Services Task Force. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ), 2007. 228 p. Electronic copies available from the [AHRQ Web site](#).
- A step-by-step guide to delivering clinical preventive services: a systems approach. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ), 2002 May. 189 p. Electronic copies available from the [AHRQ Web site](#). See the related QualityTool summary on the [Health Care Innovations Exchange Web site](#).

Print copies: Available from the Agency for Healthcare Research and Quality Publications Clearinghouse. For more information, go to <http://www.ahrq.gov/news/pubsix.htm> or call 1-800-358-9295 (U.S. only).

The [Electronic Preventive Services Selector \(ePSS\)](#), available as a PDA application and a web-based tool, is a quick hands-on tool designed to help primary care clinicians identify the screening, counseling, and preventive medication services that are appropriate for their patients. It is based on current recommendations of the USPSTF and can be searched by specific patient characteristics such as age, sex, and selected behavioral risk factors.

PATIENT RESOURCES

The following are available:

- Men: Stay Healthy at Any Age – Checklist for Your Next Checkup. Rockville (MD): Agency for Healthcare Research and Quality. AHRQ Pub. No. 07-IP006-A. February 2007. Electronic copies: Available from the [USPSTF Web site](#).
- Women: Stay Healthy at Any Age – Checklist for Your Next Checkup. Rockville (MD): Agency for Healthcare Research and Quality. AHRQ Pub. No. 07-IP005-A. February 2007. Electronic copies: Available from the [USPSTF Web site](#).

Print copies: Available from the Agency for Healthcare Research and Quality (AHRQ) Publications Clearinghouse. For more information, go to <http://www.ahrq.gov/news/pubsix.htm> or call 1-800-358-9295 (U.S. only).

Please note: This patient information is intended to provide health professionals with information to share with their patients to help them better understand their health and their diagnosed disorders. By providing access to this patient information, it is not the intention of NGC to provide specific medical advice for particular patients. Rather we urge patients and their representatives to review this material and then to consult with a licensed health professional for evaluation of treatment options suitable for them as well as for diagnosis and answers to their personal medical questions. This patient information has been derived and prepared from a guideline for health care professionals included on NGC by the authors or publishers of that original guideline. The patient information is not reviewed by NGC to establish whether or not it accurately reflects the original guideline's content.

NGC STATUS

This summary was completed by ECRI on June 30, 1998. The information was verified by the guideline developer on December 1, 1998. This NGC summary was updated by ECRI Institute on March 5, 2008. The updated information was verified by the guideline developer on March 11, 2008.

COPYRIGHT STATEMENT

Requests regarding copyright should be sent to: Randie A. Siegel, Electronic Dissemination Advisor, Division of Print and Electronic Publishing, Agency for Healthcare Research and Quality (formerly the Agency for Health Care Policy and Research), 540 Gaither Road, Rockville, MD 20850. Facsimile: 301-427-1873. E-mail: Randie.siegel@ahrq.hhs.gov.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC

Inclusion Criteria which may be found at
<http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 11/3/2008

