



Complete Summary

GUIDELINE TITLE

Management of community-acquired pneumonia in the home: an American College of Chest Physicians clinical position statement.

BIBLIOGRAPHIC SOURCE(S)

Ramsdell J, Narsavage GL, Fink JB. Management of community-acquired pneumonia in the home: an American College of Chest Physicians clinical position statement. Chest 2005 May;127(5):1752-63. [51 references] [PubMed](#)

GUIDELINE STATUS

This is the current release of the guideline.

COMPLETE SUMMARY CONTENT

SCOPE
METHODOLOGY - including Rating Scheme and Cost Analysis
RECOMMENDATIONS
EVIDENCE SUPPORTING THE RECOMMENDATIONS
BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
QUALIFYING STATEMENTS
IMPLEMENTATION OF THE GUIDELINE
INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
CATEGORIES
IDENTIFYING INFORMATION AND AVAILABILITY
DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Community-acquired pneumonia (CAP)

GUIDELINE CATEGORY

Diagnosis
Evaluation
Management
Risk Assessment
Treatment

CLINICAL SPECIALTY

Emergency Medicine
Family Practice
Infectious Diseases
Internal Medicine
Pulmonary Medicine

INTENDED USERS

Advanced Practice Nurses
Health Plans
Managed Care Organizations
Physician Assistants
Physicians

GUIDELINE OBJECTIVE(S)

To address the requirements for successfully managing patients with community-acquired pneumonia (CAP) in the home environment

TARGET POPULATION

Immunocompetent adults with community-acquired pneumonia (CAP) who are living at home or in unskilled residential facilities:

- Previously healthy individuals and chronically ill individuals who may or may not be home-bound but choose not to go into the hospital
- Hospitalized patients with CAP who are completing a hospital discharge plan

Note: These recommendations are not intended to be applied to pediatric patients, patients with human immunodeficiency virus (HIV) or other forms of immune compromise, or patients living in skilled nursing facilities.

INTERVENTIONS AND PRACTICES CONSIDERED

1. Initial evaluation and diagnosis performed in person by a qualified provider (i.e., physician, nurse practitioner, or physician assistant)
2. Determination of site of care
3. Administration of first dose of antibiotic, oxygenation, and hydration within 8 hours of presentation
4. Treatment with antibiotic regimen based on empiric therapy guidelines of the American Thoracic Society (ATS) and Infectious Disease Society of America (IDSA)
5. Transfer to acute care facility if patient care contract cannot be agreed to or is violated
6. Repeat assessment within 24 hours for high-risk patients
7. Closure visit
 - Evaluation of risk of recurrence
 - Discussion of preventive measures
 - Assessment of functional status
8. Chest radiography (a minimum of 8 weeks following diagnosis)

MAJOR OUTCOMES CONSIDERED

- Risk of infection with pathogenic bacteria
- Mortality
- Response to treatment
- Antibiotic resistance

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Hand-searches of Published Literature (Secondary Sources)
Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

This clinical position statement is based on a review of the guideline literature and on a consensus panel meeting of the American Academy of Home Care Physicians (AAHCP) Community-Acquired Pneumonia Working Group (see Appendix 2 of the original guideline document for a list of participants), which was held on October 20, 2000, in Coolfont, WV. The AAHCP panel review process included reviews of guidelines from the American Thoracic Society (ATS) (1993), the Infectious Diseases Society of America (IDSA) (2000), the Canadian Infectious Diseases Society and the Canadian Thoracic Society (2000), the Centers for Disease Control and Prevention (2000), the European Respiratory Society (1998), and the British Thoracic Society (BTS) (1993). The AAHCP recommendations were reconsidered by the American College of Chest Physicians (ACCP) working group and were updated with a MEDLINE search of the literature from October 2000 to July 2003. This review included the 2001 update of the American Thoracic Society recommendations and the 2003 update of the IDSA recommendations.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Expert Consensus (Committee)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Not applicable

METHODS USED TO ANALYZE THE EVIDENCE

Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

This clinical position statement was developed by a working group of the American College of Chest Physicians (ACCP) Home Care Network, which includes representatives of the American Thoracic Society (ATS) and the American Association of Respiratory Care.

The working group concluded that the current literature did not provide the scientific evidence to support a clinical practice guideline as defined by the ACCP (i.e., a statement that is based on scientific evidence, that explicitly documents the process used to develop the statement, and that grades the strength of the evidence used in making clinical recommendations). Because the issue of appropriate resources for the home treatment of community-acquired pneumonia (CAP) is of immediate importance, the working group proceeded to develop this clinical position statement. None of the current guidelines that address the management of patients with community-acquired pneumonia specifically focuses on practice recommendations for the home care patient, but the recommendations in this position statement are compatible with each of these existing guidelines.

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

External Peer Review
Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Each chapter of this guideline was reviewed and approved by the American College of Chest Physicians (ACCP) Health and Science Policy Committee prior to submission, approval by the ACCP Board of Regents, and approval by the American Academy of Home Care Physicians (AAHCP). The guideline was then forwarded to other external organizations for endorsement.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

Recommendations for the evaluation of care provided to the immunocompetent adult patient with community-acquired pneumonia (CAP) who is being treated at home or in an unskilled residential facility are summarized below. Future research should be focused on evidence for appropriate assessment and in-home treatment, such as diagnostic tests, home care interventions, and the exploration of the efficacy of telephone triage in selected groups of patients who are at low risk and are well-known to the provider. Finally, the approach to the in-home management of CAP that has been outlined herein, and others, should be studied to assess their impact on patient outcomes (especially in the elderly) and to ensure that in-home management can achieve the same level of quality and patient outcomes as at any other treatment site for appropriate patient subsets.

Recommendations for the Home Treatment of Patients with CAP

Initial Patient Evaluation and Diagnosis in the Home Environment

1. A qualified provider includes a physician, nurse practitioner, or physician assistant.
2. The initial evaluation should be performed in person by a qualified provider or by a visiting home nurse who is in contact with a qualified provider at the time of the evaluation. In this case, the qualified provider must evaluate the patient within 24 hours of the initial diagnosis. Telephone triage alone is not acceptable.
3. If a qualified provider does not meet with the patient at the time of the initial evaluation, the provider must see the patient sometime between presentation and closure.

Determination of Site of Care

4. Home care should be an option if it can provide the same level of quality and achieve the same level of recovery and functional status, consistent with the patient's wishes and overall treatment goals, as would be possible at any other site of care.
5. Care should be provided in a timely fashion. If these goals cannot be achieved in the home care setting, transfer to an acute care facility should be considered.
 - The first dose of antibiotic should be administered within 8 hours of presentation.
 - Oxygenation should be optimized within 8 hours of presentation.
 - Hydration should be initiated within 8 hours of presentation.
6. If a patient care contract cannot be agreed to or is violated, transfer to an acute care facility should occur.

Management of CAP at Home

7. The patient should be treated with antibiotics based using the empiric therapy guidelines of the American Thoracic Society (ATS) and Infectious Disease Society of America (IDSA).

Monitoring and Goals for Nurses Providing Interim Home Care

8. Repeat assessment should be performed within 24 hours for the high-risk patient.
9. If a provider chooses not to refer a patient to a home care agency, the standards for interim care must still be met.

Closure

10. There should be a closure visit for each patient during which the risk of recurrence is evaluated, preventive measures are discussed, and functional status is assessed.
11. A chest radiograph should be obtained to confirm the resolution of the illness a minimum of 8 weeks following diagnosis.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of evidence supporting the recommendations is not specifically stated. The recommendations are based on a review of the guideline literature and on a consensus panel meeting of the American Academy of Home Care Physicians (AAHCP) Community-Acquired Pneumonia Working Group.

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Successful management of patients with community-acquired pneumonia in the home environment would provide the same level of quality and achieve the same level of recovery and functional status as would be possible in any other site of care.

POTENTIAL HARMS

Not stated

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

- The information provided in this statement should be used in conjunction with clinical judgment. These recommendations may not apply to every individual patient; therefore, it is important for the physician to take into consideration the role of patient preferences and the availability of local resources.
- The American College of Chest Physicians (ACCP) is sensitive to concerns that nationally and/or internationally developed position statements are not always

applicable in local settings. Further, recommendations are just that: recommendations, not dictates. In treating patients, individual circumstances, preferences, and resources do play a role in the course of treatment at every decision level. These recommendations are intended to guide healthcare decisions and may be adapted to be applicable at various levels.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better

IOM DOMAIN

Effectiveness
Patient-centeredness
Timeliness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Ramsdell J, Narsavage GL, Fink JB. Management of community-acquired pneumonia in the home: an American College of Chest Physicians clinical position statement. Chest 2005 May;127(5):1752-63. [51 references] [PubMed](#)

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

2005 May

GUIDELINE DEVELOPER(S)

American Academy of Home Care Physicians - Professional Association
American College of Chest Physicians - Medical Specialty Society

SOURCE(S) OF FUNDING

American College of Chest Physicians

GUIDELINE COMMITTEE

American College of Chest Physicians' Home Care Network Working Group

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Primary Authors: Joe Ramsdell, MD, FCCP, Department of Medicine, University of California, San Diego, Medical Center, San Diego, CA; Georgia L. Narsavage, PhD, RN, CS, Francis Payne Bolton School of Nursing, Case Western Reserve University, Cleveland, OH; James B. Fink, MS, RRT, Aerogen

Work Group Members: Joe Ramsdell, MD, FCCP; Ann Cathcart, MSW, RSW; Georgia L. Narsavage, PhD, RN, CS; Nicholas S. Hill, MD, FCCP; James B. Fink, MS, RRT; Joe Lewarski, RRT; Barry Make, MD, FCCP

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

It is essential that the American College of Chest Physicians (ACCP) have full disclosure of outside interests from those individuals serving on policy development committees, including liaison representatives from outside organizations. Both real and potential conflicts of interest may actually affect impartial or objective decisions or may appear to.

No conflicts are reported in the original consensus document.

GUIDELINE STATUS

This is the current release of the guideline.

GUIDELINE AVAILABILITY

Electronic copies: Available to subscribers of [Chest - The Cardiopulmonary and Critical Care Journal](#).

Print copies: Available from the American College of Chest Physicians, Products and Registration Division, 3300 Dundee Road, Northbrook IL 60062-2348.

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI on July 26, 2005. The information was verified by the guideline developer on July 26, 2005.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which is subject to the guideline developer's copyright restrictions.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 11/3/2008

