5 *1	metrizamide, The Sanofi-Aventis Group.
	disulfiram, Odyssey Pharmaceuticals, Inc.
7	
3	Medication Guide
)	Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal
	Thoughts or Actions
	PARNATE® (PAR-nate) (tranylcypromine sulfate) Tablets
R	ead the Medication Guide that comes with you or your family member's antidepressant
	nedicine. This Medication Guide is only about the risk of suicidal thoughts and actions with
	ntidepressant medicines. Talk to your, or your family member's, healthcare provider
	bout:
•	All risks and benefits of treatment with antidepressant medicines
•	All treatment choices for depression or other serious mental illness
	The committee of the property of the committee of the com
V	What is the most important information I should know about antidepressant medicines,
	epression and other serious mental illnesses, and suicidal thoughts or actions?
1	Antidepressant medicines may increase suicidal thoughts or actions in some children,
	teenagers, and young adults within the first few months of treatment.
2	. Depression and other serious mental illnesses are the most important causes of suicidal
	thoughts and actions. Some people may have a particularly high risk of having suicidal
	thoughts or actions. These include people who have (or have a family history of) bipolar
	illness (also called manic-depressive illness) or suicidal thoughts or actions.
3	. How can I watch for and try to prevent suicidal thoughts and actions in myself or a
	family member?
	 Pay close attention to any changes, especially sudden changes, in mood, behaviors,
	thoughts, or feelings. This is very important when an antidepressant medicine is started or
	when the dose is changed.
	 Call the healthcare provider right away to report new or sudden changes in mood,
	behavior, thoughts, or feelings.
	• Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare
	provider between visits as needed, especially if you have concerns about symptoms.
	all a healthcare provider right away if you or your family member has any of the
Ī	ollowing symptoms, especially if they are new, worse, or worry you:
•	Thoughts about suicide or dying
•	r or
•	New or worse depression
•	New or worse anxiety

435 • Feeling very agitated or restless 436 Panic attacks 437 • Trouble sleeping (insomnia) 438 • New or worse irritability 439 • Acting aggressive, being angry, or violent 440 Acting on dangerous impulses 441 An extreme increase in activity and talking (mania) 442 Other unusual changes in behavior or mood 443 444 What else do I need to know about antidepressant medicines? 445 Never stop an antidepressant medicine without first talking to a healthcare provider. 446 Stopping an antidepressant medicine suddenly can cause other symptoms. 447 Antidepressants are medicines used to treat depression and other illnesses. It is 448 important to discuss all the risks of treating depression and also the risks of not treating it. 449 Patients and their families or other caregivers should discuss all treatment choices with the 450 healthcare provider, not just the use of antidepressants. 451 **Antidepressant medicines have other side effects.** Talk to the healthcare provider about the 452 side effects of the medicine prescribed for you or your family member. 453 **Antidepressant medicines can interact with other medicines.** Know all of the medicines 454 that you or your family member takes. Keep a list of all medicines to show the healthcare 455 provider. Do not start new medicines without first checking with your healthcare provider. 456 Not all antidepressant medicines prescribed for children are FDA approved for use in 457 **children.** Talk to your child's healthcare provider for more information. 458 459 This Medication Guide has been approved by the U.S. Food and Drug Administration for all 460 antidepressants. 461 462 May 2007 PRT:2MG 463 GlaxoSmithKline 464 465 GlaxoSmithKline

469 May 2007 PRT:71PI

466

467

468

470

Research Triangle Park, NC 27709

©2007, GlaxoSmithKline. All rights reserved.