# **Example of Fictional Highlights of Prescribing Information** (Based on Proposed Physician Labeling Rule)

OCRACEPHALOSE® [fictional drug] Rx (spurilous hypothetic chloride) Tablets or Capsules for oral use [fictional drug]	Discontinue Ocraco depression occurs ( • Hypoglycemia can alcohol (5.5, 6.2)
	Most Common Adverse  somnolence, dry m
, , , ,	sommorence, ary m
Adjunct therapy with a sulfonylurea to lower blood glucose in patients with Type 2 diabetes whose hyperglycemia cannot be controlled by diet and exercise (1.1)	To report SUSPECTE #) or FD
Initial dose is 100 mg once every morning and may be titrated up to 300 mg (2.1)	<ul><li>Domecattus reduc</li><li>Alcohol: increases</li></ul>
HOW SUPPLIED	<ul> <li>Hepatic impairmen impairment (4, 7.6</li> </ul>
Tablets: 100 mg (3) Capsules: 100 mg (3)	See P for PATII
CONTRAINDICATIONS	
Hepatic impairment (4)	These highlights do not Ocracephalose safely a prescribing information
Hepatic dysfunction leading to acute liver failure may occur, typically within 3 months of initiation (5.2)	
<ul> <li>Evaluate liver function prior to initiating Ocracephalose and monitor weekly for 3 months. Discontinue if LFTs increase &gt; 3 times upper limit of normal (5.2)</li> </ul>	

1

- Severe depression with suicidal ideation occurred in 2% of patients. ephalose or initiate antidepressant therapy if
- occur with insufficient caloric intake and use of

### e Reactions (> 5%) (8)

nouth, nightmares, and sexual disorders

### ED SERIOUS ADRs, call (manufacturer) at (phone A's MedWatch at 1-800-FDA-1088

## -DRUG INTERACTIONS--

- e domecattus dose by one-half (5.4, 6.1)
- incidence of hypoglycemia (6.2)

### IN SPECIFIC POPULATIONS-

nt: Contraindicated in patients with hepatic

### ENT COUNSELING INFORMATION and --phalose's approved patient labeling

include all the information needed to prescribe nd effectively. See Ocracephalose's comprehensive n provided below.

Revised: 12/2003