

U.S. Vaccines

Vaccine	Name	Manufacturer	Type	Route	Comments
Anthrax	BioThrax	BioPort	Inactivated Bacterial	SC	
DTaP	Daptacel	sanofi	Inactivated Bacterial	IM	Tetanus & diphtheria toxoids and pertussis vaccine. Not licensed for 5 th dose.
	Infanrix	GlaxoSmithKline	Inactivated Bacterial	IM	Tetanus & diphtheria toxoids and pertussis vaccine.
	Tripedia	sanofi	Inactivated Bacterial	IM	Tetanus & diphtheria toxoids and pertussis vaccine.
DT	(Generic)	sanofi	Inactivated Bacterial Toxoids	IM	Pediatric formulation
DTaP/Hib	TriHIBit	sanofi	Inactivated Bacterial	IM	ActHIB reconstituted with Tripedia. Licensed for 4 th dose of DTaP & Hib series (not primary series).
DTaP/IPV	Kinrix	GlaxoSmithKline	Inactivated Bacterial & Viral	IM	Approved for 5 th (DTaP) and 4 th (IPV) booster at 4-6 years
DTaP/IPV/HepB	Pediarix	GlaxoSmithKline	Inactivated Bacterial & Viral	IM	Approved for doses at 2, 4, 6 months (through 6 years of age). Not licensed for boosters.
DTaP/IPV/Hib	Pentacel	sanofi	Inactivated Bacterial & Viral	IM	Approved for 4 doses at 2, 4, 6, and 15-18 months.
<i>Haemophilus influenzae</i> type b (Hib)	PedvaxHIB	Merck	Inactivated Bacterial	IM	PRP-OMP. Polysaccharide conjugate (mening. protein carrier). 2-dose schedule.
	ActHIB	sanofi	Inactivated Bacterial	IM	PRP-T. Polysaccharide conjugate (tetanus toxoid carrier). 3-dose schedule.
Hepatitis A	Havrix	GlaxoSmithKline	Inactivated Viral	IM	Pediatric (≤ 18) and adult formulations. Pediatric = 720 EL.U., 0.5mL Adult = 1,140 EL.U., 1.0mL Minimum age = 1 year.
	Vaqa	Merck	Inactivated Viral	IM	Pediatric (≤ 18) and adult formulations. Pediatric = 25 U, 0.5mL Adult = 50 U, 1.0mL Minimum age = 1 year.
Hepatitis B	Engerix-B	GlaxoSmithKline	Inactivated Viral (recombinant)	IM	Pediatric (≤ 19) and adult formulations. Pediatric formulation is not licensed for adults.
	Recombivax HB	Merck	Inactivated Viral (recombinant)	IM	Pediatric (≤ 19), adult, and dialysis formulations. Two pediatric doses may be substituted for an adult dose.

Vaccine	Name	Manufacturer	Type	Route	Comments
HepA/HepB	Twinrix	GlaxoSmithKline	Inactivated Viral	IM	Pediatric dose of HepA + adult dose of HepB. Minimum age = 18 years. 3-dose series.
HepB/Hib	Comvax	Merck	Inactivated Bacterial & Viral	IM	Should not be used for HepB birth dose.
Human Papillomavirus (HPV)	Gardasil	Merck	Inactivated Viral (recombinant)	IM	Quadrivalent, Types 6, 11, 16 & 18. Licensed for females 9-26 years.
Influenza	Fluarix	GlaxoSmithKline	Inactivated Viral	IM	Trivalent Types A & B. Minimum age = 18 years.
	Fluvirin	Chiron	Inactivated Viral	IM	Trivalent Types A & B Purified surface antigen. Minimum age = 4 years.
	Fluzone	sanofi	Inactivated Viral	IM	Trivalent Types A & B Subvirion. Minimum age multidose vial = 6 months. Age range 0.25ml prefilled syringe = 6-35 months. Minimum age 0.5ml prefilled syringe = 3 years.
	FluLaval	GlaxoSmithKline	Inactivated Viral	IM	Trivalent Types A & B Minimum age = 18 years.
	Afluria	CSL	Inactivated Viral	IM	Trivalent Types A & B Minimum age = 18 years.
	FluMist	Medimmune	Live attenuated viral	Intra-nasal	Trivalent Types A & B. Age range 5-49 years.
Japanese Encephalitis	JE-Vax	sanofi	Inactivated viral	SC	
MMR	M-M-R II	Merck	Live attenuated viral	SC	Measles, mumps, rubella.
MMRV	ProQuad	Merck	Live attenuated viral	SC	Measles, mumps, rubella, varicella.
Measles	Attenuvax	Merck	Live attenuated viral	SC	Edmonston-Enders strain
Mumps	Mumpsvax	Merck	Live attenuated viral	SC	Jeryl Lynn strain
Rubella	Meruvax II	Merck	Live attenuated viral	SC	RA 27/3 strain
Meningococcal	Menomune	sanofi	Inactivated bacterial	SC	Polysaccharide, containing serogroups A, C, Y, & W-135.
	Menactra	sanofi	Inactivated bacterial	IM	Polysaccharide conjugate (diphtheria toxoid carrier), containing serogroups A, C, Y, & W-135. Age range 2-49.

Vaccine	Name	Manufacturer	Type	Route	Comments
Pneumococcal	Pneumovax 23	Merck	Inactivated bacterial	SC or IM	Polysaccharide. Contains 23 strains. Minimum age = 2 yrs.
	Prennar	Wyeth	Inactivated bacterial, conjugate	IM	Polysaccharide conjugate (diphtheria protein carrier). Contains 7 strains. Routine age range = 2-59 months.
Polio	Ipol	sanofi	Inactivated viral	SC or IM	Trivalent, Types 1, 2, & 3.
Rabies	Imovax Rabies	sanofi	Inactivated viral	IM	
	RabAvert	Chiron	Inactivated viral	IM	
Rotavirus	RotaTeq	Merck	Live viral	Oral	Pentavalent. Age range 6-32 weeks. First dose between 6 & 12 weeks; complete 3-dose series by 32 weeks.
	Rotarix	GlaxoSmithKline	Live viral	Oral	Monovalent. Age range 6-24 weeks. First dose beginning at 6 weeks; at least 4 weeks between doses; complete 2-dose series by 24 weeks.
Td	Decavac	sanofi	Inactivated bacterial toxoids	IM	Tetanus/diphtheria toxoids. Adult formulation
	(Generic)	Massachusetts Biological Labs	Inactivated bacterial toxoids	IM	Tetanus/diphtheria toxoids. Adult formulation
Tdap	Boostrix	GlaxoSmithKline	Inactivated bacterial	IM	Tetanus & diphtheria toxoids & pertussis vaccine. Licensed for ages 10-18.
	Adacel	sanofi	Inactivated bacterial	IM	Tetanus & diphtheria toxoids & pertussis vaccine. Licensed for ages 11-64.
TT	(Generic)	sanofi	Inactivated bacterial toxoid	IM	Tetanus toxoid. May be used for adults or children.
Typhoid	Typhim Vi	sanofi	Inactivated bacterial	IM	Polysaccharide.
	Vivotif Berna	Berna	Live bacterial	Oral	Ty21a strain.
Varicella	Varivax	Merck	Live viral	SC	
Vaccinia (Smallpox)	ACAM2000	Acambis	Live Viral	Percutaneous	
Yellow Fever	YF-Vax	sanofi	Live viral	SC	
Zoster (Shingles)	Zostavax	Merck	Live viral	SC	Licensed for age 60 and older.

June 2008