

Mailstop K30
(770) 488-5292
January 8, 1998

Dear Colleague:

Information related to the 1996 BRFSS summary prevalence report of selected risk factors and preventive health measures from the core and modules is attached.

This report consists of four parts:

An introduction that contains:

An overview of the 1996 summary prevalence report.

An explanation of the tables and figures for 1996 Behavioral Risk Factor Surveillance System (BRFSS) prevalence data.

Definitions of BRFSS risk factors and preventive health measures. When appropriate, Year 2000 Objectives are identified.

Analysis of selected questions from the 1996 BRFSS questionnaire. The set of analyses is divided into three sections:

- Section I - selected risk factors and preventive health measures in the BRFSS core.
- Section II - selected risk factors and preventive health measures from modules used by at least 10 states
- Section III - preventive health measures that concern women only.

As soon as your schedule permits, please examine the enclosed materials closely.

Sincerely yours,

David E. Nelson, M.D., M.P.H.
Chief, Behavioral Surveillance Branch
Division of Adult and Community Health
National Center for Chronic Disease
Prevention and Health Promotion

Enclosures

1996 BRFSS SUMMARY PREVALENCE REPORT

BEHAVIORAL SURVEILLANCE BRANCH
DIVISION OF ADULT AND COMMUNITY HEALTH
NATIONAL CENTER FOR CHRONIC DISEASE PREVENTION AND HEALTH PROMOTION
CENTERS FOR DISEASE CONTROL AND PREVENTION

TABLE OF CONTENTS

Overview of the 1996 Summary Report.....	vii
Explanation of Tables and Figures.....	viii
Definitions of Risk Factors & Preventive Health Measures.....	xvi

SECTION I

PREVALENCE REPORT FOR SELECTED CORE QUESTIONS

General Health Status

1. Self reported health status: Fair or poor	1
----------------------------------------------------	---

Quality of Life

2. Physical health: Mean number of days physical health not good.....	5
3. Mental health: Mean number of days mental health not good.....	9
4. Activities limited: Mean number of days activities limited.....	13

Health Insurance

5. No health insurance, ages 18 and over.....	17
6. No health insurance, ages 18-64	21

Diabetes

7. Diabetes Awareness: Ever told you have diabetes.	25
----------------------------------------------------------	----

Smoking

8. Ever Smoked.....	29
9. Current Smoker (Year 2000 Objective 3.4).....	33

Overweight

10. Based on body mass index (BMI) (Year 2000 Objective 2.3)	36
11. Based on percent of median	40

AIDS Knowledge/Attitudes

12. Encourage teen condom use, ages 18-64	44
13. Condom effectiveness, ages 18-64	48
14. HIV blood test, ages 18-64	52
15. Chances of getting HIV, ages 18-64	56

Physical Activity

16. No leisure time physical activity
(Year 2000 Objective 1.5)60

17. Regular and sustained physical activity
(Year 2000 Objective 1.3)64

18. Regular and vigorous physical activity
(Year 2000 Objective 1.4)68

Fruits and Vegetables

19. Consuming five or more servings of fruits and
vegetables a day (Year 2000 Objective 2.6)72

Weight Control

20. Eating fewer calories and/or less fat combined
with increased physical activity76

SECTION II

PREVALENCE REPORT FOR SELECTED MODULES

Smokeless tobacco

21. Ever use.....80

22. Current use82

Diabetes

23. Mean age at onset of diabetes.....84

24. Now taking insulin.....86

25. Mean times seen health professional (past year).....88

Health Care Utilization

26. One or more than one particular place for health care
(Year 2000 Objective 21.3).....90

Oral Health (Year 2000 Objectives 13.3,13.5)

27. Dental visit in last year92

28. Loss of permanent teeth94

Hypertension

29. Awareness: Ever told blood pressure high96

30. Screening: Blood pressure checked within past two years..98

Cholesterol Screening/Awareness

31. Screening: Ever checked100

32. Screening: Checked within past five years
(Year 2000 Objective 15.14).....102

33. Awareness: Ever told cholesterol high.....104

Immunization

34. Had a flu shot within past 12 months, ages 65 and older
(Year 2000 Objective 20.11)106

35. Ever had pneumonia vaccination, ages 65 and older
(Year 2000 Objective 20.11)108

Lack of safety belt usage

36. Does not always wear a safety belt
(Year 2000 Objective 9.12).....110

Injury Control and Child Safety

37. Use of child safety belt
(Approximates Year 2000 Objective 9.12).....112

38. Use of child safety seat
(Approximates Year 2000 Objective 9.12a).....114

39. Use of child bicycle helmet116

Drinking

40. Current Drinkers118

41. Binge Drinking(Acute Drinking)120

42. Chronic Drinking122

43. Drinking and Driving124

SECTION III

PREVALENCE REPORT FOR WOMEN'S HEALTH ISSUES

Mammography

44. Ever had a mammogram, women age 18 and older.....126

45. Ever had a mammogram, women age 40 and older.....129

46. Had a mammogram, women age 50 and older
(within past two years).....132

Breast Exam

- 47. Ever had a breast exam by doctor,
women age 18 and older.....135
- 48. Ever had a breast exam by doctor,
women age 40 and older.....138
- 49. Had a breast exam, women age 50 and older
(within past two years).....141

Mammogram and Breast Exam (Year 2000 Objective 16.11)

- 50. Ever had a mammogram and breast exam,
women age 40 and older.....144
- 51. Had a mammogram and breast exam,
women age 50 and older
(within past two years).....147

Cervical Cancer

- 52. Ever had Pap smear (all women).....150
- 53. Ever had Pap smear, women with uterine cervix
(Year 2000 Objective 16.12).....153
- 54. Had Pap smear, women with uterine cervix
(within past three years)
(Year 2000 Objective 16.12).....156

OVERVIEW OF THE 1996 SUMMARY REPORT

The report is divided into three sections:

Section I displays tables and figures for selected variables in the core. In Section I, there are three tables and a graphical map for each risk factor or preventive measure. Estimated prevalence and relevant statistics are provided for the total population sampled and by gender. Prevalence estimates for both 1995 and 1996, or where appropriate, 1994 and 1996, and the percentage point difference between them are given for variables in the core. A map of the United States displays values at or above the median value versus below the median value.

Section II shows tables for selected variables in modules used by at least 10 states, with the exception of the preventive counseling module, which was new in 1996. Section II contains two tables for each risk factor or preventive measure of interest in the modules. These tables show the estimated prevalence and relevant statistics for the total sampled population and both sex groups who responded to selected questions in the optional modules.

Section III contains tables for preventive health measures that are related to women's health issues. Estimated prevalence and relevant statistics are provided for the total population sampled. Prevalence estimates for both 1995 and 1996 and the percentage point difference between them are given. A map of the United States displays values at or above versus below the median.

The next section describes each of the tables in detail.

EXPLANATION OF TABLES AND FIGURES

Three sample tables (Tables A-C) and Figure A from **Section I** will be used as examples to explain the information contained in this report. Each table or figure describes a behavioral risk factor or preventive health measure from the Behavioral Risk Factor Surveillance System (BRFSS) instrument.

Table A shows the estimated prevalence and relevant statistics for each participant. (Participants refer to all states, and the District of Columbia and territories of the U.S.). The sample table shows the estimated prevalence and relevant statistics for "no health insurance" in the population surveyed by each BRFSS participant. This prevalence estimate excludes respondents with unknown information on insurance coverage.

The columns in Table A provide the following information:

<u>COLUMN HEADING</u>	<u>CONTENTS</u>
<u>Participant</u>	Name of the participant. Participants refers to all states including the District of Columbia and territories of the United States.
<u>Sample size</u>	The number of individuals who responded to the specific BRFSS question. These are unweighted numbers. These numbers <u>exclude</u> individuals who reported "not sure, don't know", those who refused to answer the question, or had no response coded (missing).
<u>Percent/Mean</u>	An estimate of the percent (or prevalence or mean number) of residents who report the behavioral risk factor of interest. This participant's population to estimate the true prevalence or mean in the population. is weighted to the
<u>Standard error</u>	A measure of the amount of variability that exists in the calculation of the prevalence estimate.
<u>95% confidence interval</u>	A range which contains the true population prevalence estimate 95% of the time, if repeated sampling of the population is performed.

SUMMARY STATISTICS are given at the bottom of the table:

<u>Number of participants</u>	The total number of participants who asked question(s) regarding the respective "risk factor".
<u>Median</u>	The middle value of all estimated prevalences.
<u>Range</u>	The lowest prevalence and the highest prevalence among all participants.

For example, in **Table A**, there were 2135 individuals in the state of Alabama who responded to the "no health insurance" question. Of those responding, the estimated prevalence of Alabama's population who reported that they had no health care plan is 15.98 percent. This prevalence estimate has a standard error of 0.91. The 95% confidence interval for this estimate is (14.20, 17.76). Ninety-five percent (95%) of confidence intervals generated under repeated sampling will contain the true population prevalence of Alabama's population who "had no health care plan".

Figure A is a map of the United States that displays state participants in one of two categories: at or above the median or below the median prevalence value. The following United States map shows BRFSS participants in either of these two categories for the "no health insurance" prevalence value.

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2135	15.98	0.91	(14.20, 17.76)
ALASKA	1530	20.76	1.57	(17.68, 23.84)
ARIZONA	1943	17.00	1.23	(14.59, 19.41)
ARKANSAS	1796	17.93	1.07	(15.83, 20.03)
CALIFORNIA	3986	19.51	0.76	(18.02, 21.00)
COLORADO	1803	13.46	0.99	(11.52, 15.40)
CONNECTICUT	1829	9.34	0.84	(7.69, 10.99)
DELAWARE	2122	10.78	0.91	(9.00, 12.56)
DISTRICT OF CO	1196	13.39	1.17	(11.10, 15.68)
FLORIDA	3571	17.55	0.75	(16.08, 19.02)
GEORGIA	2379	9.76	0.72	(8.35, 11.17)
IDAHO	2728	15.00	0.77	(13.49, 16.51)
ILLINOIS	2879	11.84	0.71	(10.45, 13.23)
INDIANA	2205	11.42	0.77	(9.91, 12.93)
IOWA	3597	9.16	0.56	(8.06, 10.26)
KANSAS	2004	10.09	0.79	(8.54, 11.64)
KENTUCKY	3609	14.42	0.72	(13.01, 15.83)
LOUISIANA	1646	22.83	1.23	(20.42, 25.24)
MAINE	1696	14.22	1.02	(12.22, 16.22)
MARYLAND	4474	9.74	0.60	(8.56, 10.92)
MASSACHUSETTS	1780	10.19	0.88	(8.47, 11.91)
MICHIGAN	2481	9.56	0.66	(8.27, 10.85)
MINNESOTA	4395	7.21	0.44	(6.35, 8.07)
MISSISSIPPI	1586	14.21	1.03	(12.19, 16.23)
MISSOURI	1549	12.93	0.98	(11.01, 14.85)
MONTANA	1797	15.61	0.96	(13.73, 17.49)
NEBRASKA	2091	8.57	0.71	(7.18, 9.96)
NEVADA	1795	16.43	1.30	(13.88, 18.98)
NEW HAMPSHIRE	1500	11.88	0.99	(9.94, 13.82)
NEW JERSEY	3137	11.14	0.74	(9.69, 12.59)
NEW MEXICO	1092	21.02	1.60	(17.88, 24.16)
NEW YORK	4300	12.43	0.60	(11.25, 13.61)
NORTH CAROLINA	2789	12.94	0.78	(11.41, 14.47)
NORTH DAKOTA	1805	11.76	0.91	(9.98, 13.54)
OHIO	1552	11.37	1.04	(9.33, 13.41)
OKLAHOMA	1923	18.47	1.12	(16.27, 20.67)
OREGON	2919	13.20	0.77	(11.69, 14.71)
PENNSYLVANIA	3578	11.07	0.62	(9.85, 12.29)
RHODE ISLAND	1845	10.49	0.87	(8.78, 12.20)
SOUTH CAROLINA	1868	16.31	1.21	(13.94, 18.68)
SOUTH DAKOTA	2100	10.12	0.75	(8.65, 11.59)
TENNESSEE	3052	11.98	0.67	(10.67, 13.29)
TEXAS	1694	24.60	1.20	(22.25, 26.95)
UTAH	2873	12.18	0.76	(10.69, 13.67)
VERMONT	2497	12.98	0.80	(11.41, 14.55)
VIRGINIA	1891	14.57	0.99	(12.63, 16.51)
WASHINGTON	3582	11.94	0.62	(10.72, 13.16)
WEST VIRGINIA	2412	17.69	0.90	(15.93, 19.45)
WISCONSIN	2226	9.30	0.85	(7.63, 10.97)
WYOMING	2522	18.33	0.87	(16.62, 20.04)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 12.94 RANGE = 7.21-24.60
* HAVING NO HEALTH CARE PLAN AGES 18 AND OVER

November 19, 1997

Table B shows the estimated prevalence and relevant statistics by sex for each state participant. This example table shows the estimated prevalence and relevant statistics for "no health insurance" among both sex groups. The same statistics are displayed in Table B as in Table A, except that Table B shows values for males and females separately. An explanation of the terms was provided earlier. This side by side display of the tables permits each participant to compare estimates for males and females in their respective state.

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*				*	-----FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	779	15.97	1.51	(13.01, 18.93)	*	1356	15.99	1.09	(13.85, 18.13)
ALASKA	710	24.09	2.48	(19.23, 28.95)	*	820	17.09	1.77	(13.62, 20.56)
ARIZONA	809	18.77	1.98	(14.89, 22.65)	*	1134	15.31	1.54	(12.29, 18.33)
ARKANSAS	647	18.08	1.66	(14.83, 21.33)	*	1149	17.80	1.33	(15.19, 20.41)
CALIFORNIA	1722	22.24	1.21	(19.87, 24.61)	*	2264	16.83	0.93	(15.01, 18.65)
COLORADO	793	14.83	1.56	(11.77, 17.89)	*	1010	12.14	1.22	(9.75, 14.53)
CONNECTICUT	729	9.49	1.24	(7.06, 11.92)	*	1100	9.20	1.15	(6.95, 11.45)
DELAWARE	828	11.87	1.50	(8.93, 14.81)	*	1294	9.79	0.95	(7.93, 11.65)
DISTRICT OF CO	459	13.91	1.97	(10.05, 17.77)	*	737	12.95	1.43	(10.15, 15.75)
FLORIDA	1440	18.39	1.17	(16.10, 20.68)	*	2131	16.78	0.93	(14.96, 18.60)
GEORGIA	1018	10.83	1.20	(8.48, 13.18)	*	1361	8.79	0.87	(7.08, 10.50)
IDAHO	1114	15.58	1.22	(13.19, 17.97)	*	1614	14.44	0.94	(12.60, 16.28)
ILLINOIS	1152	12.83	1.15	(10.58, 15.08)	*	1727	10.93	0.85	(9.26, 12.60)
INDIANA	937	11.11	1.15	(8.86, 13.36)	*	1268	11.69	1.00	(9.73, 13.65)
IOWA	1477	10.94	0.91	(9.16, 12.72)	*	2120	7.55	0.65	(6.28, 8.82)
KANSAS	858	10.06	1.22	(7.67, 12.45)	*	1146	10.12	1.03	(8.10, 12.14)
KENTUCKY	1317	14.43	1.13	(12.22, 16.64)	*	2292	14.41	0.88	(12.69, 16.13)
LOUISIANA	652	23.93	1.98	(20.05, 27.81)	*	994	21.86	1.51	(18.90, 24.82)
MAINE	708	13.69	1.44	(10.87, 16.51)	*	988	14.72	1.38	(12.02, 17.42)
MARYLAND	1835	10.98	1.03	(8.96, 13.00)	*	2639	8.61	0.69	(7.26, 9.96)
MASSACHUSETTS	739	12.15	1.43	(9.35, 14.95)	*	1041	8.43	1.07	(6.33, 10.53)
MICHIGAN	1043	10.67	1.05	(8.61, 12.73)	*	1438	8.56	0.81	(6.97, 10.15)
MINNESOTA	1954	8.28	0.67	(6.97, 9.59)	*	2441	6.21	0.58	(5.07, 7.35)
MISSISSIPPI	587	13.87	1.72	(10.50, 17.24)	*	999	14.51	1.26	(12.04, 16.98)
MISSOURI	620	13.78	1.63	(10.59, 16.97)	*	929	12.17	1.13	(9.96, 14.38)
MONTANA	769	14.11	1.35	(11.46, 16.76)	*	1028	17.03	1.32	(14.44, 19.62)
NEBRASKA	834	10.02	1.17	(7.73, 12.31)	*	1257	7.25	0.85	(5.58, 8.92)
NEVADA	756	18.54	2.09	(14.44, 22.64)	*	1039	14.30	1.50	(11.36, 17.24)
NEW HAMPSHIRE	647	13.39	1.50	(10.45, 16.33)	*	853	10.47	1.26	(8.00, 12.94)
NEW JERSEY	1329	13.52	1.30	(10.97, 16.07)	*	1808	8.99	0.77	(7.48, 10.50)
NEW MEXICO	432	22.78	2.51	(17.86, 27.70)	*	660	19.39	2.02	(15.43, 23.35)
NEW YORK	1740	15.19	1.00	(13.23, 17.15)	*	2560	9.99	0.67	(8.68, 11.30)
NORTH CAROLINA	1142	12.91	1.22	(10.52, 15.30)	*	1647	12.98	0.99	(11.04, 14.92)
NORTH DAKOTA	736	12.65	1.41	(9.89, 15.41)	*	1069	10.90	1.10	(8.74, 13.06)
OHIO	625	13.40	1.76	(9.95, 16.85)	*	927	9.54	1.12	(7.34, 11.74)
OKLAHOMA	854	18.35	1.76	(14.90, 21.80)	*	1069	18.57	1.44	(15.75, 21.39)
OREGON	1229	15.05	1.27	(12.56, 17.54)	*	1690	11.46	0.90	(9.70, 13.22)
PENNSYLVANIA	1516	13.14	1.01	(11.16, 15.12)	*	2062	9.24	0.75	(7.77, 10.71)
RHODE ISLAND	792	11.91	1.23	(9.50, 14.32)	*	1053	9.24	1.25	(6.79, 11.69)
SOUTH CAROLINA	740	15.42	1.96	(11.58, 19.26)	*	1128	17.11	1.47	(14.23, 19.99)
SOUTH DAKOTA	892	10.67	1.18	(8.36, 12.98)	*	1208	9.60	0.92	(7.80, 11.40)
TENNESSEE	1173	13.82	1.11	(11.64, 16.00)	*	1879	10.33	0.80	(8.76, 11.90)
TEXAS	661	23.47	1.86	(19.82, 27.12)	*	1033	25.66	1.53	(22.66, 28.66)
UTAH	1249	11.33	1.06	(9.25, 13.41)	*	1624	12.98	1.08	(10.86, 15.10)
VERMONT	1071	14.35	1.25	(11.90, 16.80)	*	1426	11.70	1.02	(9.70, 13.70)
VIRGINIA	732	13.61	1.59	(10.49, 16.73)	*	1159	15.47	1.23	(13.06, 17.88)
WASHINGTON	1518	14.65	1.03	(12.63, 16.67)	*	2064	9.34	0.70	(7.97, 10.71)
WEST VIRGINIA	900	19.45	1.44	(16.63, 22.27)	*	1512	16.14	1.11	(13.96, 18.32)
WISCONSIN	946	10.37	1.32	(7.78, 12.96)	*	1280	8.29	1.10	(6.13, 10.45)
WYOMING	1046	19.83	1.33	(17.22, 22.44)	*	1476	16.86	1.06	(14.78, 18.94)
SUMMARY STATISTICS:		NUMBER OF PARTICIPANTS =	50			NUMBER OF PARTICIPANTS =	50		
		MEDIAN	= 13.80			MEDIAN	= 11.92		
		RANGE	= 8.28-24.09			RANGE	= 6.21-25.66		

* HAVING NO HEALTH CARE PLAN AGES 18 AND OVER

Table C shows the percent difference between the prevalence estimates, by participant, for the 1995 and 1996 survey years for having no health insurance. A negative change indicates a decrease in prevalence from 1995 to 1996, while a positive change indicates an increase in prevalence between the two years. This difference gives state health officials a quick method to examine changes that may have occurred in their respective state. Some of these changes may have resulted from legislative actions (or inactions) in the state, programs funded, or programs that came into existence because of results that you may have shown from the BRFSS. Since changes between two years may be attributable to many factors, conclusions drawn from this table should be interpreted with extreme caution.

Sample Size. Sample sizes vary by question. Some estimates may be based on responses from fewer than 50 respondents. Interpreting estimates that are based on a small number of respondents can mislead the reader into believing that a given finding is much more precise than it actually is. The BRFSS recommends not interpreting percentages where the denominator is based upon fewer than 50 unweighted respondents (sample size).

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.26	15.98	1.72
ALASKA	15.33	20.76	5.43
ARIZONA	15.20	17.00	1.80
ARKANSAS	15.40	17.93	2.53
CALIFORNIA	19.52	19.51	-0.01
COLORADO	14.19	13.46	-0.73
CONNECTICUT	9.75	9.34	-0.41
DELAWARE	11.35	10.78	-0.57
DISTRICT OF CO	N/A	13.39	N/A
FLORIDA	14.98	17.55	2.57
GEORGIA	10.18	9.76	-0.42
IDAHO	14.30	15.00	0.70
ILLINOIS	11.17	11.84	0.67
INDIANA	10.21	11.42	1.21
IOWA	9.16	9.16	0.00
KANSAS	11.09	10.09	-1.00
KENTUCKY	14.28	14.42	0.14
LOUISIANA	20.96	22.83	1.87
MAINE	16.35	14.22	-2.13
MARYLAND	9.44	9.74	0.30
MASSACHUSETTS	10.47	10.19	-0.28
MICHIGAN	8.55	9.56	1.01
MINNESOTA	8.18	7.21	-0.97
MISSISSIPPI	13.98	14.21	0.23
MISSOURI	15.03	12.93	-2.10
MONTANA	16.31	15.61	-0.70
NEBRASKA	7.82	8.57	0.75
NEVADA	13.11	16.43	3.32
NEW HAMPSHIRE	12.22	11.88	-0.34
NEW JERSEY	8.08	11.14	3.06
NEW MEXICO	18.57	21.02	2.45
NEW YORK	12.25	12.43	0.18
NORTH CAROLINA	12.55	12.94	0.39
NORTH DAKOTA	10.57	11.76	1.19
OHIO	11.03	11.37	0.34
OKLAHOMA	14.92	18.47	3.55
OREGON	12.96	13.20	0.24
PENNSYLVANIA	9.28	11.07	1.79
RHODE ISLAND	10.67	10.49	-0.18
SOUTH CAROLINA	12.06	16.31	4.25
SOUTH DAKOTA	8.96	10.12	1.16
TENNESSEE	11.37	11.98	0.61
TEXAS	17.26	24.60	7.34
UTAH	10.84	12.18	1.34
VERMONT	11.87	12.98	1.11
VIRGINIA	11.41	14.57	3.16
WASHINGTON	11.49	11.94	0.45
WEST VIRGINIA	15.96	17.69	1.73
WISCONSIN	7.77	9.30	1.53
WYOMING	16.25	18.33	2.08

* HAVING NO HEALTH CARE PLAN
 N/A - NOT AVAILABLE

November 19, 1997

DEFINITIONS OF BRFSS RISK FACTORS AND PREVENTIVE HEALTH MEASURES

ACTIVITY LIMITATIONS

ANY DISABILITY - Respondents, age 18-64, who report that an impairment or health problem is responsible for any of the following: (1) keeping them from working at a job or business, (2) limiting the kind or amount of work they can do, (3) keeping them from doing housework, or (4) limiting the kind or amount of housework they can do. This measures Year 2000 Objective 17.2 - Target $\leq 8\%$.

ANY LIMITATIONS (1) - Respondents, age 18-64, who report they are limited in any way in any activities because of any impairment or health problem. This measures Year 2000 Objective 17.2 - Target $\leq 8\%$.

ANY LIMITATIONS (2) - Respondents, age 65 and older, who report they are limited in any way in any activities because of any impairment or health problem. This measures Year 2000 Objective 17.2 - Target $\leq 8\%$.

AIDS KNOWLEDGE/ATTITUDES

HIV BLOOD TEST - Respondents, age 18-64, who have ever had a blood test for HIV infection.

CHANCES OF GETTING HIV - Respondents, age 18-64, who believe that their chances of getting infected with HIV, the virus that causes AIDS are medium or high.

CONDOM EFFECTIVENESS - Respondents, age 18-64, who believe that condoms are very effective in preventing AIDS.

ENCOURAGE TEEN CONDOM USE - Respondents, age 18-64, who would encourage sexually active teenagers to use a condom.

BREAST EXAM

EVER HAD A BREAST EXAM - Female respondents, age 18 and older, who report that they have ever had a breast exam by a doctor.

EVER HAD A BREAST EXAM - Female respondents, age 40 and older, who report that they have ever had a breast exam by a doctor.

HAD A BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they had a breast exam within the past two years.

CERVICAL CANCER

EVER HAD A PAP SMEAR - Female respondents who report that they have ever had a pap smear.

EVER HAD A PAP SMEAR, WITH UTERINE CERVIX - Female respondents, with uterine cervix, who report that they have ever had a pap smear. This measures Year 2000 Objective 16.12. - Target $\geq 95\%$.

HAD A PAP SMEAR WITHIN 3 YEARS - Female respondents, with uterine cervix, who report that they had a pap smear within the last three years. This measures Year 2000 Objective 16.12. - Target $\geq 85\%$.

CHOLESTEROL SCREENING/AWARENESS

CHOLESTEROL SCREENING (1) - Respondents who have ever had their cholesterol checked.

CHOLESTEROL SCREENING (2) - Respondents who have had their cholesterol checked within the past five years. Year 2000 Objective 15.14 - Target $\geq 75\%$.

CHOLESTEROL AWARENESS - Respondents who report that they have been told their cholesterol is high by health professional.

CLINICAL BREAST EXAM See BREAST EXAM

DIABETES

AGE AT ONSET - Mean age at onset of diabetes: age when respondent was told that they have diabetes, among those who report they were told by a doctor that they have diabetes.

DIABETES AWARENESS - Respondents who report they were told by a doctor that they have diabetes.

NOW TAKING INSULIN - Respondents who are now taking insulin, among those who report they were told by a doctor that they have diabetes.

NUMBER OF TIMES SEEN HEALTH PROFESSIONAL - Mean number of times respondents report they have seen a doctor, nurse, or

other health professional for their diabetes in the past year, among those who report they were told by a doctor that they have diabetes.

DRINKING STATUS

CURRENT DRINKERS - Respondents who report that they have had any alcoholic beverages in the past month.

BINGE DRINKING (also called ACUTE DRINKING) - Respondents who report having five or more drinks on an occasion, one or more times in the past month.

CHRONIC DRINKING - Respondents who report an average of two or more drinks per day i.e., 60 or more alcoholic drinks a month. This is based on the total number of drinks per month.

DRINKING and DRIVING - Respondents who report having driven after having too much to drink, one or more times in the past month.

FRUITS AND VEGETABLES

FRUITS AND VEGETABLES - Respondents who report they are consuming five or more servings of fruits and vegetables a day. This measures Year 2000 Objective 2.6 - Target is not specified.

HEALTH CARE UTILIZATION

PARTICULAR PLACE FOR HEALTH CARE - Respondents who report they have one or more than one particular clinic, health center, doctor's office, or other place that they usually go to if they are sick or need advice about their health. This measures Year 2000 Objective 21.3 - Target $\geq 95\%$.

HEALTH INSURANCE

NO HEALTH INSURANCE - Respondents who report they have no health care plan.

NO HEALTH INSURANCE - Respondents age 18-64 years who report they have no health care plan.

HEALTH STATUS

ACTIVITIES LIMITED - Mean number of days in past 30 days activities were limited as reported by respondents.

MENTAL HEALTH NOT GOOD - Mean number of days in past 30 days mental health was not good as reported by respondents.

PHYSICAL HEALTH NOT GOOD - Mean number of days in past 30 days physical health was not good as reported by respondents.

SELF REPORTED HEALTH STATUS - Respondents who report having a general health status of fair or poor.

HYPERTENSION

HYPERTENSION [also called HYPERTENSION AWARENESS] - Respondents who report that they have ever been told they have high blood pressure.

HYPERTENSION SCREENING - Respondents who report they have had their blood pressure checked within the past two years.

IMMUNIZATION

FLU SHOT - Respondents, age 65 and older, who report that they have had a flu shot within past 12 months. This measures Year 2000 Objective 20.11 - Target $\geq 80\%$.

PNEUMONIA VACCINATION - Respondents, age 65 and older, who have ever had a pneumonia vaccination. This measures Year 2000 Objective 20.11 - Target $\geq 80\%$.

INJURY CONTROL AND CHILD SAFETY

USE OF CHILD SAFETY BELTS FOR OLDEST CHILD AGE 5-15 - Respondents who report their oldest child is between age 5 and 15 and "always" or "nearly always" uses a safety belt. This approximates Year 2000 Objective 9.12 - Target $\geq 85\%$.

USE OF CHILD SAFETY SEATS FOR OLDEST CHILD UNDER 5 - Respondents who report their oldest child is under age 5 and "always" or "nearly always" uses a car safety seat. This approximates Year 2000 Objective 9.12a - Target $\geq 95\%$.

USE OF BICYCLE HELMET - Respondents who report their oldest child is under age 15 and child age 5-15 always wears a bicycle helmet when riding a bicycle.

MAMMOGRAPHY

EVER HAD A MAMMOGRAM (1) - Female respondents age 18 and older, who report that they have ever had a mammogram.

EVER HAD A MAMMOGRAM (2) - Female respondents, age 40 and older, who report that they have ever had a mammogram.

HAD A MAMMOGRAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they had a mammogram within the past two years.

MAMMOGRAPHY AND BREAST EXAM

MAMMOGRAM & BREAST EXAM - Female respondents, age 40 and older, who report that they have ever had a mammogram and a breast exam. This measures Year 2000 Objective 16.11 - Target $\geq 80\%$.

MAMMOGRAM & BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they have had a mammogram and a breast exam within the last two years. This measures Year 2000 Objective 16.11 - Target $\geq 60\%$.

OVERWEIGHT

OVERWEIGHT: BY MEDIAN - Respondent at or above 120% of their ideal weight. Ideal weight is defined as the mid-value of a median frame person from the 1959 Metropolitan Life Insurance Corporation height-weight tables.

OVERWEIGHT: BY BODY MASS INDEX - Females with body mass index(BMI) ≥ 27.3 and males with BMI ≥ 27.8 . BMI is defined as weight in kilograms divided by height in meters squared (w/h^2). This measures Year 2000 Objective 2.3 - Target $\leq 20\%$. This should be used with extreme caution since people tend to underreport their weight. BRFSS may underestimate the prevalence of overweight.

ORAL HEALTH

DENTAL VISIT - Respondents who reported visiting a dentist or dental clinic in the last year.

TOOTH LOSS - Respondents who report a loss of 6 or more permanent teeth due to tooth decay or gum disease.

PAP SMEAR See CERVICAL CANCER

PHYSICAL INACTIVITY

NO LEISURE TIME PHYSICAL ACTIVITY - Respondents who report no leisure-time physical activity during the past month. This measures Year 2000 Objective 1.5 - Target \leq 15%.

REGULAR AND SUSTAINED PHYSICAL ACTIVITY - Respondents who report physical activity 5 or more sessions per week, 30 minutes or more per session, regardless of intensity. This measures Year 2000 Objective 1.3 - Target \geq 30%.

REGULAR AND VIGOROUS PHYSICAL ACTIVITY - Respondents who report physical activity or a pair of activities for 3 or more sessions per week, 20 minutes or more per session, at 50% or more capacity. This measures Year 2000 Objective 1.4 - Target \geq 20%.

SAFETY BELTS

LACK OF SAFETY BELT USAGE - Respondents reporting they "nearly always", "sometimes", "seldom", or "never" use safety belts (i.e., do not always use a safety belt). This measures Year 2000 Objective 9.12 - Target \geq 85%.

SMOKELESS TOBACCO

EVER USED - Respondents who have ever used or tried smokeless tobacco.

CURRENT USE - Respondents who report that they currently use smokeless tobacco among all respondents in the state.

SMOKING STATUS

EVER SMOKED - Respondents who have ever smoked 100 cigarettes in their lifetime.

CURRENT SMOKER - Respondents who have ever smoked 100 cigarettes in their lifetime and reported smoking every day or some days. The wording of this question changed between 1995 and 1996. This measures Year 2000 Objective 3.4 - Target \leq 15%.

WEIGHT CONTROL

CALORIES AND PHYSICAL ACTIVITY - Respondents at risk on BMI who report that they are either eating fewer calories or less fat and using physical activity or exercise to lose or maintain weight.

LIMITATIONS OF THE DATA

This report consolidates the most current data on selected risk factors and preventive health measures for states and the District of Columbia. The information was obtained from the data files of individual states. Therefore, while all participants use telephone surveys to collect the data, the data in this report vary somewhat with respect to data collection, sampling design and sample size.

Some states conduct interviews using a paper questionnaire, but most use a computer-assisted telephone interviewing (CATI) software program. All states use probability-based methods to select their sample. For data year 1996, thirty-one states used the Mitofsky-Waksberg method of sampling, fourteen states used a disproportionate stratified sample (DSS) design, and six states used a variety of other designs. **The BRFSS standard for sample designs is that sample records must be justifiable as a probability sample of all households with telephones in the state.** All but Alaska, California, Nevada and Texas met this criterion in 1996. The deviations from the standard by these states may reduce the comparability of their data with those of other areas. Although the effects of the deviations have not been extensively investigated, departure from the standard may produce bias in prevalence estimates and other survey results, as well as constrict confidence intervals. The sampling practices of the four states are highlighted below:

Alaska used a stratified directory/random sample combination where the sample design is made up of 3 random digit-dialed strata and 4 listed strata. Phone numbers in the listed strata are selected from a list of putative household numbers. Forty-five percent of the completed interviews come from the listed strata. Alaska's sampling frame for the listed strata contains an estimated 50% of all household numbers in these strata.

California used a truncated list-assisted design with sampling from 3+ blocks only. Noncoverage was further affected to an unknown degree by substitution for sample numbers used by any client of the commercial sample provider in the preceding six months. Probabilities of selection varied in proportion to the number of listed household numbers in the 100 blocks associated with a number, the number of known business numbers preceding a number, the number of preceding numbers selected by any client of

the commercial sample provider in the preceding six months, and the measure of size used for each county, which could vary each time a sample was drawn. Because of the unavailability of appropriate information, none of these factors is accounted for in the weights.

Nevada used a truncated list-assisted design with sampling from 5+ blocks only. Noncoverage was further affected to an unknown degree by substitution for sample numbers used by any client of the commercial sample provider in the preceding six months. Probabilities of selection varied in proportion to the number of listed household numbers in the 100 blocks associated with a number, the number of known business numbers preceding a number, the number of preceding numbers selected by any client of the commercial sample provider in the preceding six months, and the measure of size used for each county, which could vary each time a sample was drawn. Because of the unavailability of appropriate information, only the number of listed household numbers in the 100 blocks associated with a number is accounted for in the weights.

Texas used a truncated list-assisted sample which included only phone numbers from 100 blocks with one or more listed household numbers. Such hundred blocks are estimated to contain 97.1% of all household numbers.

In addition to departing from the standard for sample designs, **California** modified the wording of mammography and Papanicolaou (PAP) smear, and are therefore excluded from those tables in the report. California also asked the HIV/AIDS section questions to persons 18-45 years of age rather than those 18-65 years of age, and are therefore excluded from tables reporting data on this topic.

Although states collect data monthly, the number of interviews range from 1096 in New Mexico to 4003 in California. Estimates for some states are therefore based upon smaller sample sizes, and for rare events (e.g. diabetes), may yield unstable estimates.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. People may not remember essential information, a question may not mean the same thing to different respondents, and some individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their impact on the data. The user must make his or her own evaluation on the data. Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. When the number of events is small and the probability of such an event is

small, considerable caution must be observed in interpreting the conditions described by the figures.

Final data for **Hawaii** and **Puerto Rico** were not received in time to be included in this report.

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2134	20.19	1.04	(18.15, 22.23)
ALASKA	1532	8.27	0.93	(6.45, 10.09)
ARIZONA	1954	12.10	0.95	(10.24, 13.96)
ARKANSAS	1785	18.90	1.07	(16.80, 21.00)
CALIFORNIA	4003	15.45	0.68	(14.12, 16.78)
COLORADO	1803	9.24	0.77	(7.73, 10.75)
CONNECTICUT	1833	10.14	0.78	(8.61, 11.67)
DELAWARE	2123	13.29	0.77	(11.78, 14.80)
DISTRICT OF CO	1201	13.03	1.17	(10.74, 15.32)
FLORIDA	3575	15.42	0.66	(14.13, 16.71)
GEORGIA	2382	8.86	0.63	(7.63, 10.09)
IDAHO	2732	11.70	0.69	(10.35, 13.05)
ILLINOIS	2879	13.70	0.74	(12.25, 15.15)
INDIANA	2204	14.05	0.79	(12.50, 15.60)
IOWA	3599	11.55	0.57	(10.43, 12.67)
KANSAS	1994	12.19	0.78	(10.66, 13.72)
KENTUCKY	3613	20.92	0.74	(19.47, 22.37)
LOUISIANA	1654	16.69	1.06	(14.61, 18.77)
MAINE	1698	13.18	0.89	(11.44, 14.92)
MARYLAND	4475	11.81	0.64	(10.56, 13.06)
MASSACHUSETTS	1778	12.76	0.89	(11.02, 14.50)
MICHIGAN	2485	11.95	0.72	(10.54, 13.36)
MINNESOTA	4389	10.24	0.50	(9.26, 11.22)
MISSISSIPPI	1589	21.05	1.09	(18.91, 23.19)
MISSOURI	1547	14.05	0.96	(12.17, 15.93)
MONTANA	1800	11.60	0.80	(10.03, 13.17)
NEBRASKA	2095	13.19	0.81	(11.60, 14.78)
NEVADA	1795	11.36	1.06	(9.28, 13.44)
NEW HAMPSHIRE	1498	10.39	0.79	(8.84, 11.94)
NEW JERSEY	3140	13.74	0.72	(12.33, 15.15)
NEW MEXICO	1094	16.18	1.35	(13.53, 18.83)
NEW YORK	4304	12.67	0.57	(11.55, 13.79)
NORTH CAROLINA	2785	16.28	0.80	(14.71, 17.85)
NORTH DAKOTA	1808	12.54	0.87	(10.83, 14.25)
OHIO	1552	12.57	0.88	(10.85, 14.29)
OKLAHOMA	1921	13.96	0.89	(12.22, 15.70)
OREGON	2929	12.20	0.67	(10.89, 13.51)
PENNSYLVANIA	3589	14.40	0.66	(13.11, 15.69)
RHODE ISLAND	1847	14.83	0.98	(12.91, 16.75)
SOUTH CAROLINA	1876	16.86	1.22	(14.47, 19.25)
SOUTH DAKOTA	2101	11.49	0.75	(10.02, 12.96)
TENNESSEE	3047	19.54	0.80	(17.97, 21.11)
TEXAS	1700	18.43	1.04	(16.39, 20.47)
UTAH	2886	11.29	0.74	(9.84, 12.74)
VERMONT	2491	10.36	0.63	(9.13, 11.59)
VIRGINIA	1884	13.08	0.91	(11.30, 14.86)
WASHINGTON	3574	10.85	0.56	(9.75, 11.95)
WEST VIRGINIA	2416	22.78	1.01	(20.80, 24.76)
WISCONSIN	2225	11.46	0.93	(9.64, 13.28)
WYOMING	2531	11.78	0.68	(10.45, 13.11)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 12.90 RANGE = 8.27-22.78

* HEALTH STATUS IS FAIR OR POOR

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	780	20.19	1.70	(16.86, 23.52)	* 1354	20.19	1.22	(17.80, 22.58)
ALASKA	714	8.68	1.37	(5.99, 11.37)	* 818	7.82	1.23	(5.41, 10.23)
ARIZONA	812	10.89	1.44	(8.07, 13.71)	* 1142	13.23	1.34	(10.60, 15.86)
ARKANSAS	646	17.89	1.70	(14.56, 21.22)	* 1139	19.81	1.33	(17.20, 22.42)
CALIFORNIA	1730	15.11	1.00	(13.15, 17.07)	* 2273	15.78	0.91	(14.00, 17.56)
COLORADO	794	8.64	1.11	(6.46, 10.82)	* 1009	9.81	1.06	(7.73, 11.89)
CONNECTICUT	732	8.80	1.12	(6.60, 11.00)	* 1101	11.36	1.12	(9.16, 13.56)
DELAWARE	828	11.99	1.24	(9.56, 14.42)	* 1295	14.47	1.03	(12.45, 16.49)
DISTRICT OF CO	462	11.73	1.79	(8.22, 15.24)	* 739	14.13	1.46	(11.27, 16.99)
FLORIDA	1442	14.87	0.98	(12.95, 16.79)	* 2133	15.92	0.89	(14.18, 17.66)
GEORGIA	1020	7.44	0.82	(5.83, 9.05)	* 1362	10.16	0.91	(8.38, 11.94)
IDAHO	1119	11.73	1.10	(9.57, 13.89)	* 1613	11.68	0.84	(10.03, 13.33)
ILLINOIS	1152	13.15	1.12	(10.95, 15.35)	* 1727	14.21	0.95	(12.35, 16.07)
INDIANA	937	12.80	1.13	(10.59, 15.01)	* 1267	15.18	1.09	(13.04, 17.32)
IOWA	1477	11.39	0.89	(9.65, 13.13)	* 2122	11.69	0.75	(10.22, 13.16)
KANSAS	856	11.98	1.20	(9.63, 14.33)	* 1138	12.39	1.00	(10.43, 14.35)
KENTUCKY	1318	19.03	1.16	(16.76, 21.30)	* 2295	22.64	0.95	(20.78, 24.50)
LOUISIANA	658	16.60	1.65	(13.37, 19.83)	* 996	16.78	1.33	(14.17, 19.39)
MAINE	709	12.81	1.34	(10.18, 15.44)	* 989	13.51	1.14	(11.28, 15.74)
MARYLAND	1835	11.67	0.95	(9.81, 13.53)	* 2640	11.94	0.82	(10.33, 13.55)
MASSACHUSETTS	738	11.49	1.35	(8.84, 14.14)	* 1040	13.90	1.16	(11.63, 16.17)
MICHIGAN	1045	9.17	0.99	(7.23, 11.11)	* 1440	14.49	1.03	(12.47, 16.51)
MINNESOTA	1956	9.25	0.72	(7.84, 10.66)	* 2433	11.16	0.69	(9.81, 12.51)
MISSISSIPPI	587	18.66	1.68	(15.37, 21.95)	* 1002	23.15	1.48	(20.25, 26.05)
MISSOURI	618	12.42	1.49	(9.50, 15.34)	* 929	15.51	1.21	(13.14, 17.88)
MONTANA	771	11.39	1.20	(9.04, 13.74)	* 1029	11.80	1.03	(9.78, 13.82)
NEBRASKA	835	13.26	1.28	(10.75, 15.77)	* 1260	13.13	1.01	(11.15, 15.11)
NEVADA	759	10.85	1.56	(7.79, 13.91)	* 1036	11.88	1.44	(9.06, 14.70)
NEW HAMPSHIRE	644	10.83	1.24	(8.40, 13.26)	* 854	9.97	1.09	(7.83, 12.11)
NEW JERSEY	1332	12.79	1.05	(10.73, 14.85)	* 1808	14.59	0.98	(12.67, 16.51)
NEW MEXICO	434	15.68	2.09	(11.58, 19.78)	* 660	16.65	1.84	(13.04, 20.26)
NEW YORK	1746	11.57	0.85	(9.90, 13.24)	* 2558	13.64	0.77	(12.13, 15.15)
NORTH CAROLINA	1140	15.75	1.18	(13.44, 18.06)	* 1645	16.76	0.99	(14.82, 18.70)
NORTH DAKOTA	740	12.67	1.33	(10.06, 15.28)	* 1068	12.42	1.07	(10.32, 14.52)
OHIO	625	13.10	1.40	(10.36, 15.84)	* 927	12.09	1.07	(9.99, 14.19)
OKLAHOMA	853	12.41	1.28	(9.90, 14.92)	* 1068	15.39	1.19	(13.06, 17.72)
OREGON	1234	11.11	0.95	(9.25, 12.97)	* 1695	13.23	0.95	(11.37, 15.09)
PENNSYLVANIA	1524	14.59	1.01	(12.61, 16.57)	* 2065	14.24	0.87	(12.53, 15.95)
RHODE ISLAND	795	11.03	1.37	(8.34, 13.72)	* 1052	18.22	1.39	(15.50, 20.94)
SOUTH CAROLINA	748	14.62	1.93	(10.84, 18.40)	* 1128	18.88	1.48	(15.98, 21.78)
SOUTH DAKOTA	896	11.86	1.19	(9.53, 14.19)	* 1205	11.15	0.95	(9.29, 13.01)
TENNESSEE	1173	17.76	1.20	(15.41, 20.11)	* 1874	21.14	1.04	(19.10, 23.18)
TEXAS	666	16.17	1.54	(13.15, 19.19)	* 1034	20.56	1.38	(17.86, 23.26)
UTAH	1254	9.84	1.00	(7.88, 11.80)	* 1632	12.66	1.08	(10.54, 14.78)
VERMONT	1069	10.14	0.94	(8.30, 11.98)	* 1422	10.57	0.85	(8.90, 12.24)
VIRGINIA	729	10.61	1.34	(7.98, 13.24)	* 1155	15.38	1.23	(12.97, 17.79)
WASHINGTON	1514	10.55	0.82	(8.94, 12.16)	* 2060	11.13	0.77	(9.62, 12.64)
WEST VIRGINIA	901	23.82	1.57	(20.74, 26.90)	* 1515	21.86	1.20	(19.51, 24.21)
WISCONSIN	944	10.13	1.20	(7.78, 12.48)	* 1281	12.69	1.39	(9.97, 15.41)
WYOMING	1050	11.89	1.06	(9.81, 13.97)	* 1481	11.69	0.85	(10.02, 13.36)
SUMMARY STATISTICS:		NUMBER OF PARTICIPANTS =	50		NUMBER OF PARTICIPANTS =	50		
		MEDIAN	= 11.94		MEDIAN	= 13.77		
		RANGE	= 7.44-23.82		RANGE	= 7.82-23.15		

* HEALTH STATUS IS FAIR OR POOR

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	18.39	20.19	1.80
ALASKA	8.24	8.27	0.03
ARIZONA	13.44	12.10	-1.34
ARKANSAS	18.61	18.90	0.29
CALIFORNIA	14.40	15.45	1.05
COLORADO	10.73	9.24	-1.49
CONNECTICUT	10.87	10.14	-0.73
DELAWARE	13.40	13.29	-0.11
DISTRICT OF CO	N/A	13.03	N/A
FLORIDA	14.32	15.42	1.10
GEORGIA	12.34	8.86	-3.48
IDAHO	11.21	11.70	0.49
ILLINOIS	13.97	13.70	-0.27
INDIANA	14.71	14.05	-0.66
IOWA	11.34	11.55	0.21
KANSAS	12.29	12.19	-0.10
KENTUCKY	21.07	20.92	-0.15
LOUISIANA	17.30	16.69	-0.61
MAINE	12.90	13.18	0.28
MARYLAND	10.42	11.81	1.39
MASSACHUSETTS	12.31	12.76	0.45
MICHIGAN	14.70	11.95	-2.75
MINNESOTA	9.74	10.24	0.50
MISSISSIPPI	21.46	21.05	-0.41
MISSOURI	12.79	14.05	1.26
MONTANA	11.25	11.60	0.35
NEBRASKA	10.59	13.19	2.60
NEVADA	14.07	11.36	-2.71
NEW HAMPSHIRE	10.23	10.39	0.16
NEW JERSEY	10.93	13.74	2.81
NEW MEXICO	17.18	16.18	-1.00
NEW YORK	11.65	12.67	1.02
NORTH CAROLINA	18.57	16.28	-2.29
NORTH DAKOTA	13.51	12.54	-0.97
OHIO	14.20	12.57	-1.63
OKLAHOMA	13.10	13.96	0.86
OREGON	11.35	12.20	0.85
PENNSYLVANIA	13.21	14.40	1.19
RHODE ISLAND	14.20	14.83	0.63
SOUTH CAROLINA	15.50	16.86	1.36
SOUTH DAKOTA	12.73	11.49	-1.24
TENNESSEE	17.76	19.54	1.78
TEXAS	17.24	18.43	1.19
UTAH	11.80	11.29	-0.51
VERMONT	11.51	10.36	-1.15
VIRGINIA	11.54	13.08	1.54
WASHINGTON	10.85	10.85	0.00
WEST VIRGINIA	22.61	22.78	0.17
WISCONSIN	10.14	11.46	1.32
WYOMING	11.00	11.78	0.78

* HEALTH STATUS IS FAIR OR POOR

N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2082	3.27	0.19	(2.90, 3.64)
ALASKA	1505	2.10	0.21	(1.69, 2.51)
ARIZONA	1949	2.16	0.19	(1.79, 2.53)
ARKANSAS	1773	3.08	0.20	(2.69, 3.47)
CALIFORNIA	3966	3.13	0.14	(2.86, 3.40)
COLORADO	1794	2.90	0.18	(2.55, 3.25)
CONNECTICUT	1807	2.53	0.17	(2.20, 2.86)
DELAWARE	2107	2.76	0.17	(2.43, 3.09)
DISTRICT OF CO	1188	2.58	0.24	(2.11, 3.05)
FLORIDA	3535	3.40	0.14	(3.13, 3.67)
GEORGIA	2371	2.04	0.15	(1.75, 2.33)
IDAHO	2708	2.84	0.15	(2.55, 3.13)
ILLINOIS	2857	3.15	0.14	(2.88, 3.42)
INDIANA	2152	3.42	0.18	(3.07, 3.77)
IOWA	3585	2.82	0.12	(2.58, 3.06)
KANSAS	1980	2.04	0.14	(1.77, 2.31)
KENTUCKY	3607	3.60	0.15	(3.31, 3.89)
LOUISIANA	1626	2.92	0.21	(2.51, 3.33)
MAINE	1675	2.89	0.19	(2.52, 3.26)
MARYLAND	4442	2.49	0.13	(2.24, 2.74)
MASSACHUSETTS	1762	3.19	0.18	(2.84, 3.54)
MICHIGAN	2460	2.82	0.16	(2.51, 3.13)
MINNESOTA	4318	2.95	0.12	(2.71, 3.19)
MISSISSIPPI	1554	3.20	0.21	(2.79, 3.61)
MISSOURI	1520	3.48	0.23	(3.03, 3.93)
MONTANA	1783	2.62	0.17	(2.29, 2.95)
NEBRASKA	2083	2.91	0.16	(2.60, 3.22)
NEVADA	1780	2.99	0.23	(2.54, 3.44)
NEW HAMPSHIRE	1481	2.96	0.21	(2.55, 3.37)
NEW JERSEY	3105	3.00	0.17	(2.67, 3.33)
NEW MEXICO	1087	3.72	0.33	(3.07, 4.37)
NEW YORK	4248	3.13	0.13	(2.88, 3.38)
NORTH CAROLINA	2768	2.67	0.14	(2.40, 2.94)
NORTH DAKOTA	1805	3.56	0.20	(3.17, 3.95)
OHIO	1548	2.56	0.20	(2.17, 2.95)
OKLAHOMA	1909	2.13	0.17	(1.80, 2.46)
OREGON	2899	3.02	0.15	(2.73, 3.31)
PENNSYLVANIA	3522	3.16	0.15	(2.87, 3.45)
RHODE ISLAND	1824	3.51	0.21	(3.10, 3.92)
SOUTH CAROLINA	1850	2.68	0.20	(2.29, 3.07)
SOUTH DAKOTA	2093	2.25	0.15	(1.96, 2.54)
TENNESSEE	3013	3.22	0.15	(2.93, 3.51)
TEXAS	1668	3.32	0.21	(2.91, 3.73)
UTAH	2878	3.07	0.15	(2.78, 3.36)
VERMONT	2471	3.16	0.17	(2.83, 3.49)
VIRGINIA	1824	2.91	0.19	(2.54, 3.28)
WASHINGTON	3512	3.28	0.14	(3.01, 3.55)
WEST VIRGINIA	2402	3.25	0.18	(2.90, 3.60)
WISCONSIN	2204	3.16	0.20	(2.77, 3.55)
WYOMING	2505	2.95	0.15	(2.66, 3.24)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 2.98 RANGE = 2.04- 3.72

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD* --BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	768	3.25	0.31	(2.64, 3.86)	*	1314	3.28	0.22	(2.85, 3.71)
ALASKA	703	1.69	0.28	(1.14, 2.24)	*	802	2.54	0.31	(1.93, 3.15)
ARIZONA	810	1.96	0.29	(1.39, 2.53)	*	1139	2.35	0.27	(1.82, 2.88)
ARKANSAS	643	2.56	0.30	(1.97, 3.15)	*	1130	3.55	0.26	(3.04, 4.06)
CALIFORNIA	1721	2.69	0.19	(2.32, 3.06)	*	2245	3.57	0.19	(3.20, 3.94)
COLORADO	788	2.65	0.27	(2.12, 3.18)	*	1006	3.14	0.24	(2.67, 3.61)
CONNECTICUT	726	2.23	0.27	(1.70, 2.76)	*	1081	2.81	0.23	(2.36, 3.26)
DELAWARE	824	2.24	0.24	(1.77, 2.71)	*	1283	3.22	0.23	(2.77, 3.67)
DISTRICT OF CO	459	2.35	0.39	(1.59, 3.11)	*	729	2.76	0.28	(2.21, 3.31)
FLORIDA	1429	2.95	0.22	(2.52, 3.38)	*	2106	3.81	0.18	(3.46, 4.16)
GEORGIA	1017	1.73	0.22	(1.30, 2.16)	*	1354	2.32	0.20	(1.93, 2.71)
IDAHO	1113	2.27	0.21	(1.86, 2.68)	*	1595	3.38	0.20	(2.99, 3.77)
ILLINOIS	1145	2.78	0.22	(2.35, 3.21)	*	1712	3.49	0.19	(3.12, 3.86)
INDIANA	922	2.95	0.25	(2.46, 3.44)	*	1230	3.85	0.24	(3.38, 4.32)
IOWA	1475	2.50	0.18	(2.15, 2.85)	*	2110	3.10	0.16	(2.79, 3.41)
KANSAS	850	1.48	0.18	(1.13, 1.83)	*	1130	2.57	0.21	(2.16, 2.98)
KENTUCKY	1316	3.22	0.23	(2.77, 3.67)	*	2291	3.95	0.19	(3.58, 4.32)
LOUISIANA	649	2.66	0.30	(2.07, 3.25)	*	977	3.15	0.27	(2.62, 3.68)
MAINE	703	2.37	0.27	(1.84, 2.90)	*	972	3.36	0.27	(2.83, 3.89)
MARYLAND	1830	2.12	0.20	(1.73, 2.51)	*	2612	2.84	0.18	(2.49, 3.19)
MASSACHUSETTS	736	2.62	0.25	(2.13, 3.11)	*	1026	3.71	0.26	(3.20, 4.22)
MICHIGAN	1041	2.14	0.22	(1.71, 2.57)	*	1419	3.44	0.22	(3.01, 3.87)
MINNESOTA	1936	2.70	0.16	(2.39, 3.01)	*	2382	3.18	0.16	(2.87, 3.49)
MISSISSIPPI	582	2.64	0.30	(2.05, 3.23)	*	972	3.71	0.31	(3.10, 4.32)
MISSOURI	613	3.38	0.38	(2.64, 4.12)	*	907	3.56	0.27	(3.03, 4.09)
MONTANA	766	2.11	0.24	(1.64, 2.58)	*	1017	3.10	0.24	(2.63, 3.57)
NEBRASKA	836	2.51	0.23	(2.06, 2.96)	*	1247	3.29	0.23	(2.84, 3.74)
NEVADA	753	2.26	0.30	(1.67, 2.85)	*	1027	3.74	0.33	(3.09, 4.39)
NEW HAMPSHIRE	637	2.59	0.33	(1.94, 3.24)	*	844	3.31	0.28	(2.76, 3.86)
NEW JERSEY	1330	2.82	0.29	(2.25, 3.39)	*	1775	3.17	0.20	(2.78, 3.56)
NEW MEXICO	433	3.31	0.51	(2.31, 4.31)	*	654	4.11	0.42	(3.29, 4.93)
NEW YORK	1722	2.75	0.19	(2.38, 3.12)	*	2526	3.47	0.17	(3.14, 3.80)
NORTH CAROLINA	1132	2.27	0.21	(1.86, 2.68)	*	1636	3.02	0.19	(2.65, 3.39)
NORTH DAKOTA	739	3.17	0.31	(2.56, 3.78)	*	1066	3.93	0.28	(3.38, 4.48)
OHIO	621	2.36	0.30	(1.77, 2.95)	*	927	2.75	0.24	(2.28, 3.22)
OKLAHOMA	851	1.74	0.23	(1.29, 2.19)	*	1058	2.49	0.23	(2.04, 2.94)
OREGON	1222	2.23	0.19	(1.86, 2.60)	*	1677	3.76	0.22	(3.33, 4.19)
PENNSYLVANIA	1501	2.85	0.22	(2.42, 3.28)	*	2021	3.45	0.21	(3.04, 3.86)
RHODE ISLAND	787	2.83	0.28	(2.28, 3.38)	*	1037	4.12	0.31	(3.51, 4.73)
SOUTH CAROLINA	745	2.01	0.29	(1.44, 2.58)	*	1105	3.29	0.28	(2.74, 3.84)
SOUTH DAKOTA	895	2.01	0.22	(1.58, 2.44)	*	1198	2.47	0.20	(2.08, 2.86)
TENNESSEE	1163	2.78	0.22	(2.35, 3.21)	*	1850	3.62	0.22	(3.19, 4.05)
TEXAS	654	2.66	0.31	(2.05, 3.27)	*	1014	3.96	0.27	(3.43, 4.49)
UTAH	1252	2.52	0.21	(2.11, 2.93)	*	1626	3.58	0.21	(3.17, 3.99)
VERMONT	1065	2.73	0.23	(2.28, 3.18)	*	1406	3.56	0.25	(3.07, 4.05)
VIRGINIA	706	2.26	0.27	(1.73, 2.79)	*	1118	3.50	0.26	(2.99, 4.01)
WASHINGTON	1502	2.75	0.19	(2.38, 3.12)	*	2010	3.78	0.20	(3.39, 4.17)
WEST VIRGINIA	898	3.27	0.29	(2.70, 3.84)	*	1504	3.23	0.21	(2.82, 3.64)
WISCONSIN	944	3.12	0.31	(2.51, 3.73)	*	1260	3.19	0.27	(2.66, 3.72)
WYOMING	1046	2.73	0.24	(2.26, 3.20)	*	1459	3.18	0.20	(2.79, 3.57)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 2.61	MEDIAN	= 3.34
	RANGE	= 1.48- 3.38	RANGE	= 2.32- 4.12

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 MEAN DIFFERENCES FROM 1995 TO 1996
 PHYSICAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995	1996	DIFFERENCE
	MEAN	MEAN	
ALABAMA	3.40	3.27	-0.13
ALASKA	2.58	2.10	-0.48
ARIZONA	3.42	2.16	-1.26
ARKANSAS	4.03	3.08	-0.95
CALIFORNIA	3.44	3.13	-0.31
COLORADO	3.17	2.90	-0.27
CONNECTICUT	2.50	2.53	0.03
DELAWARE	2.97	2.76	-0.21
DISTRICT OF CO	N/A	2.58	N/A
FLORIDA	3.42	3.40	-0.02
GEORGIA	2.53	2.04	-0.49
IDAHO	2.78	2.84	0.06
ILLINOIS	2.56	3.15	0.59
INDIANA	3.51	3.42	-0.09
IOWA	3.22	2.82	-0.40
KANSAS	3.06	2.04	-1.02
KENTUCKY	3.71	3.60	-0.11
LOUISIANA	3.43	2.92	-0.51
MAINE	2.50	2.89	0.39
MARYLAND	2.59	2.49	-0.10
MASSACHUSETTS	3.14	3.19	0.05
MICHIGAN	3.03	2.82	-0.21
MINNESOTA	2.78	2.95	0.17
MISSISSIPPI	3.27	3.20	-0.07
MISSOURI	2.88	3.48	0.60
MONTANA	2.74	2.62	-0.12
NEBRASKA	2.87	2.91	0.04
NEVADA	3.13	2.99	-0.14
NEW HAMPSHIRE	2.57	2.96	0.39
NEW JERSEY	3.16	3.00	-0.16
NEW MEXICO	3.74	3.72	-0.02
NEW YORK	2.71	3.13	0.42
NORTH CAROLINA	2.70	2.67	-0.03
NORTH DAKOTA	2.82	3.56	0.74
OHIO	2.69	2.56	-0.13
OKLAHOMA	2.53	2.13	-0.40
OREGON	3.02	3.02	0.00
PENNSYLVANIA	3.36	3.16	-0.20
RHODE ISLAND	3.28	3.51	0.23
SOUTH CAROLINA	2.65	2.68	0.03
SOUTH DAKOTA	2.96	2.25	-0.71
TENNESSEE	3.39	3.22	-0.17
TEXAS	3.30	3.32	0.02
UTAH	3.35	3.07	-0.28
VERMONT	3.18	3.16	-0.02
VIRGINIA	2.84	2.91	0.07
WASHINGTON	3.32	3.28	-0.04
WEST VIRGINIA	3.50	3.25	-0.25
WISCONSIN	3.33	3.16	-0.17
WYOMING	2.53	2.95	0.42

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2086	3.35	0.18	(3.00, 3.70)
ALASKA	1492	3.03	0.23	(2.58, 3.48)
ARIZONA	1949	2.15	0.20	(1.76, 2.54)
ARKANSAS	1771	2.97	0.21	(2.56, 3.38)
CALIFORNIA	3970	2.98	0.12	(2.74, 3.22)
COLORADO	1782	3.31	0.20	(2.92, 3.70)
CONNECTICUT	1813	2.93	0.19	(2.56, 3.30)
DELAWARE	2104	2.30	0.15	(2.01, 2.59)
DISTRICT OF CO	1186	3.02	0.25	(2.53, 3.51)
FLORIDA	3529	3.18	0.13	(2.93, 3.43)
GEORGIA	2364	2.16	0.13	(1.91, 2.41)
IDAHO	2709	2.96	0.14	(2.69, 3.23)
ILLINOIS	2855	3.07	0.13	(2.82, 3.32)
INDIANA	2162	3.28	0.17	(2.95, 3.61)
IOWA	3585	2.86	0.12	(2.62, 3.10)
KANSAS	1985	1.99	0.13	(1.74, 2.24)
KENTUCKY	3573	4.59	0.18	(4.24, 4.94)
LOUISIANA	1625	3.10	0.20	(2.71, 3.49)
MAINE	1685	2.93	0.19	(2.56, 3.30)
MARYLAND	4438	2.33	0.12	(2.09, 2.57)
MASSACHUSETTS	1754	3.34	0.19	(2.97, 3.71)
MICHIGAN	2450	3.25	0.16	(2.94, 3.56)
MINNESOTA	4309	3.19	0.12	(2.95, 3.43)
MISSISSIPPI	1541	2.28	0.17	(1.95, 2.61)
MISSOURI	1531	3.03	0.20	(2.64, 3.42)
MONTANA	1771	2.45	0.15	(2.16, 2.74)
NEBRASKA	2068	2.98	0.37	(2.25, 3.71)
NEVADA	1778	3.31	0.24	(2.84, 3.78)
NEW HAMPSHIRE	1480	2.81	0.18	(2.46, 3.16)
NEW JERSEY	3097	3.05	0.14	(2.78, 3.32)
NEW MEXICO	1078	3.42	0.27	(2.89, 3.95)
NEW YORK	4210	2.95	0.11	(2.73, 3.17)
NORTH CAROLINA	2766	2.01	0.13	(1.76, 2.26)
NORTH DAKOTA	1804	2.77	0.16	(2.46, 3.08)
OHIO	1544	1.85	0.16	(1.54, 2.16)
OKLAHOMA	1913	2.05	0.18	(1.70, 2.40)
OREGON	2891	3.16	0.14	(2.89, 3.43)
PENNSYLVANIA	3531	2.90	0.13	(2.65, 3.15)
RHODE ISLAND	1823	3.12	0.17	(2.79, 3.45)
SOUTH CAROLINA	1843	2.82	0.21	(2.41, 3.23)
SOUTH DAKOTA	2098	1.75	0.13	(1.50, 2.00)
TENNESSEE	3009	2.53	0.13	(2.28, 2.78)
TEXAS	1661	3.29	0.21	(2.88, 3.70)
UTAH	2882	3.63	0.17	(3.30, 3.96)
VERMONT	2462	2.91	0.15	(2.62, 3.20)
VIRGINIA	1835	2.61	0.17	(2.28, 2.94)
WASHINGTON	3506	3.16	0.13	(2.91, 3.41)
WEST VIRGINIA	2394	2.47	0.15	(2.18, 2.76)
WISCONSIN	2202	3.00	0.19	(2.63, 3.37)
WYOMING	2511	2.62	0.13	(2.37, 2.87)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 2.97 RANGE = 1.75- 4.59

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
MEAN DIFFERENCES FROM 1995 TO 1996
MENTAL HEALTH NOT GOOD*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995	1996	DIFFERENCE
	MEAN	MEAN	
ALABAMA	3.23	3.35	0.12
ALASKA	2.89	3.03	0.14
ARIZONA	2.96	2.15	-0.81
ARKANSAS	2.72	2.97	0.25
CALIFORNIA	3.15	2.98	-0.17
COLORADO	3.34	3.31	-0.03
CONNECTICUT	2.92	2.93	0.01
DELAWARE	2.92	2.30	-0.62
DISTRICT OF CO	N/A	3.02	N/A
FLORIDA	3.46	3.18	-0.28
GEORGIA	2.76	2.16	-0.60
IDAHO	3.03	2.96	-0.07
ILLINOIS	2.15	3.07	0.92
INDIANA	3.57	3.28	-0.29
IOWA	3.17	2.86	-0.31
KANSAS	2.95	1.99	-0.96
KENTUCKY	4.63	4.59	-0.04
LOUISIANA	3.18	3.10	-0.08
MAINE	2.35	2.93	0.58
MARYLAND	2.22	2.33	0.11
MASSACHUSETTS	3.32	3.34	0.02
MICHIGAN	3.34	3.25	-0.09
MINNESOTA	3.11	3.19	0.08
MISSISSIPPI	2.57	2.28	-0.29
MISSOURI	3.06	3.03	-0.03
MONTANA	2.47	2.45	-0.02
NEBRASKA	2.75	2.98	0.23
NEVADA	3.46	3.31	-0.15
NEW HAMPSHIRE	2.62	2.81	0.19
NEW JERSEY	2.51	3.05	0.54
NEW MEXICO	2.85	3.42	0.57
NEW YORK	3.17	2.95	-0.22
NORTH CAROLINA	1.94	2.01	0.07
NORTH DAKOTA	2.81	2.77	-0.04
OHIO	2.35	1.85	-0.50
OKLAHOMA	1.81	2.05	0.24
OREGON	3.31	3.16	-0.15
PENNSYLVANIA	3.02	2.90	-0.12
RHODE ISLAND	3.45	3.12	-0.33
SOUTH CAROLINA	2.48	2.82	0.34
SOUTH DAKOTA	2.06	1.75	-0.31
TENNESSEE	2.47	2.53	0.06
TEXAS	3.29	3.29	0.00
UTAH	3.22	3.63	0.41
VERMONT	3.13	2.91	-0.22
VIRGINIA	3.07	2.61	-0.46
WASHINGTON	3.09	3.16	0.07
WEST VIRGINIA	2.71	2.47	-0.24
WISCONSIN	2.77	3.00	0.23
WYOMING	2.68	2.62	-0.06

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1130	3.65	0.26	(3.14, 4.16)
ALASKA	787	2.64	0.32	(2.01, 3.27)
ARIZONA	536	5.51	0.55	(4.43, 6.59)
ARKANSAS	677	5.95	0.46	(5.05, 6.85)
CALIFORNIA	2180	3.57	0.19	(3.20, 3.94)
COLORADO	1010	2.94	0.24	(2.47, 3.41)
CONNECTICUT	897	2.75	0.26	(2.24, 3.26)
DELAWARE	761	4.54	0.33	(3.89, 5.19)
DISTRICT OF CO	544	3.60	0.42	(2.78, 4.42)
FLORIDA	1845	3.86	0.22	(3.43, 4.29)
GEORGIA	987	2.80	0.23	(2.35, 3.25)
IDAHO	1423	3.02	0.21	(2.61, 3.43)
ILLINOIS	1652	3.11	0.19	(2.74, 3.48)
INDIANA	1209	3.17	0.22	(2.74, 3.60)
IOWA	1862	2.92	0.17	(2.59, 3.25)
KANSAS	777	2.54	0.25	(2.05, 3.03)
KENTUCKY	1601	6.48	0.30	(5.89, 7.07)
LOUISIANA	765	4.14	0.34	(3.47, 4.81)
MAINE	736	4.57	0.35	(3.88, 5.26)
MARYLAND	1818	3.21	0.24	(2.74, 3.68)
MASSACHUSETTS	1016	2.98	0.24	(2.51, 3.45)
MICHIGAN	1297	3.19	0.23	(2.74, 3.64)
MINNESOTA	2431	3.54	0.17	(3.21, 3.87)
MISSISSIPPI	616	4.94	0.43	(4.10, 5.78)
MISSOURI	788	3.57	0.32	(2.94, 4.20)
MONTANA	882	2.79	0.25	(2.30, 3.28)
NEBRASKA	1033	4.36	0.58	(3.22, 5.50)
NEVADA	996	3.17	0.33	(2.52, 3.82)
NEW HAMPSHIRE	765	3.49	0.34	(2.82, 4.16)
NEW JERSEY	1544	3.36	0.22	(2.93, 3.79)
NEW MEXICO	576	4.04	0.47	(3.12, 4.96)
NEW YORK	2230	3.39	0.18	(3.04, 3.74)
NORTH CAROLINA	948	5.56	0.33	(4.91, 6.21)
NORTH DAKOTA	903	3.03	0.27	(2.50, 3.56)
OHIO	539	3.50	0.38	(2.76, 4.24)
OKLAHOMA	528	5.11	0.50	(4.13, 6.09)
OREGON	1627	3.30	0.21	(2.89, 3.71)
PENNSYLVANIA	1668	3.36	0.21	(2.95, 3.77)
RHODE ISLAND	970	3.80	0.30	(3.21, 4.39)
SOUTH CAROLINA	759	5.22	0.42	(4.40, 6.04)
SOUTH DAKOTA	753	3.07	0.28	(2.52, 3.62)
TENNESSEE	1287	4.21	0.26	(3.70, 4.72)
TEXAS	916	3.74	0.30	(3.15, 4.33)
UTAH	1663	2.78	0.19	(2.41, 3.15)
VERMONT	1223	3.40	0.25	(2.91, 3.89)
VIRGINIA	931	2.80	0.23	(2.35, 3.25)
WASHINGTON	2025	2.92	0.17	(2.59, 3.25)
WEST VIRGINIA	1002	6.06	0.35	(5.37, 6.75)
WISCONSIN	1261	3.06	0.25	(2.57, 3.55)
WYOMING	1322	2.82	0.20	(2.43, 3.21)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 3.40 RANGE = 2.54- 6.48

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
MEAN DIFFERENCES FROM 1995 TO 1996
ACTIVITIES LIMITED*

PARTICIPANT	DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED		DIFFERENCE
	1995 MEAN	1996 MEAN	
ALABAMA	4.19	3.65	-0.54
ALASKA	2.57	2.64	0.07
ARIZONA	3.06	5.51	2.45
ARKANSAS	4.90	5.95	1.05
CALIFORNIA	3.67	3.57	-0.10
COLORADO	3.18	2.94	-0.24
CONNECTICUT	2.90	2.75	-0.15
DELAWARE	4.59	4.54	-0.05
DISTRICT OF CO	N/A	3.60	N/A
FLORIDA	3.66	3.86	0.20
GEORGIA	3.33	2.80	-0.53
IDAHO	2.72	3.02	0.30
ILLINOIS	3.19	3.11	-0.08
INDIANA	3.19	3.17	-0.02
IOWA	3.22	2.92	-0.30
KANSAS	3.43	2.54	-0.89
KENTUCKY	6.29	6.48	0.19
LOUISIANA	4.26	4.14	-0.12
MAINE	4.26	4.57	0.31
MARYLAND	3.77	3.21	-0.56
MASSACHUSETTS	2.84	2.98	0.14
MICHIGAN	3.12	3.19	0.07
MINNESOTA	3.31	3.54	0.23
MISSISSIPPI	5.38	4.94	-0.44
MISSOURI	3.43	3.57	0.14
MONTANA	3.27	2.79	-0.48
NEBRASKA	3.24	4.36	1.12
NEVADA	3.20	3.17	-0.03
NEW HAMPSHIRE	2.77	3.49	0.72
NEW JERSEY	3.34	3.36	0.02
NEW MEXICO	3.93	4.04	0.11
NEW YORK	3.07	3.39	0.32
NORTH CAROLINA	5.10	5.56	0.46
NORTH DAKOTA	2.75	3.03	0.28
OHIO	2.93	3.50	0.57
OKLAHOMA	4.28	5.11	0.83
OREGON	2.92	3.30	0.38
PENNSYLVANIA	3.51	3.36	-0.15
RHODE ISLAND	3.36	3.80	0.44
SOUTH CAROLINA	4.24	5.22	0.98
SOUTH DAKOTA	3.19	3.07	-0.12
TENNESSEE	4.10	4.21	0.11
TEXAS	3.44	3.74	0.30
UTAH	3.04	2.78	-0.26
VERMONT	3.10	3.40	0.30
VIRGINIA	2.94	2.80	-0.14
WASHINGTON	3.02	2.92	-0.10
WEST VIRGINIA	6.52	6.06	-0.46
WISCONSIN	3.10	3.06	-0.04
WYOMING	2.50	2.82	0.32

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2135	15.98	0.91	(14.20, 17.76)
ALASKA	1530	20.76	1.57	(17.68, 23.84)
ARIZONA	1943	17.00	1.23	(14.59, 19.41)
ARKANSAS	1796	17.93	1.07	(15.83, 20.03)
CALIFORNIA	3986	19.51	0.76	(18.02, 21.00)
COLORADO	1803	13.46	0.99	(11.52, 15.40)
CONNECTICUT	1829	9.34	0.84	(7.69, 10.99)
DELAWARE	2122	10.78	0.91	(9.00, 12.56)
DISTRICT OF CO	1196	13.39	1.17	(11.10, 15.68)
FLORIDA	3571	17.55	0.75	(16.08, 19.02)
GEORGIA	2379	9.76	0.72	(8.35, 11.17)
IDAHO	2728	15.00	0.77	(13.49, 16.51)
ILLINOIS	2879	11.84	0.71	(10.45, 13.23)
INDIANA	2205	11.42	0.77	(9.91, 12.93)
IOWA	3597	9.16	0.56	(8.06, 10.26)
KANSAS	2004	10.09	0.79	(8.54, 11.64)
KENTUCKY	3609	14.42	0.72	(13.01, 15.83)
LOUISIANA	1646	22.83	1.23	(20.42, 25.24)
MAINE	1696	14.22	1.02	(12.22, 16.22)
MARYLAND	4474	9.74	0.60	(8.56, 10.92)
MASSACHUSETTS	1780	10.19	0.88	(8.47, 11.91)
MICHIGAN	2481	9.56	0.66	(8.27, 10.85)
MINNESOTA	4395	7.21	0.44	(6.35, 8.07)
MISSISSIPPI	1586	14.21	1.03	(12.19, 16.23)
MISSOURI	1549	12.93	0.98	(11.01, 14.85)
MONTANA	1797	15.61	0.96	(13.73, 17.49)
NEBRASKA	2091	8.57	0.71	(7.18, 9.96)
NEVADA	1795	16.43	1.30	(13.88, 18.98)
NEW HAMPSHIRE	1500	11.88	0.99	(9.94, 13.82)
NEW JERSEY	3137	11.14	0.74	(9.69, 12.59)
NEW MEXICO	1092	21.02	1.60	(17.88, 24.16)
NEW YORK	4300	12.43	0.60	(11.25, 13.61)
NORTH CAROLINA	2789	12.94	0.78	(11.41, 14.47)
NORTH DAKOTA	1805	11.76	0.91	(9.98, 13.54)
OHIO	1552	11.37	1.04	(9.33, 13.41)
OKLAHOMA	1923	18.47	1.12	(16.27, 20.67)
OREGON	2919	13.20	0.77	(11.69, 14.71)
PENNSYLVANIA	3578	11.07	0.62	(9.85, 12.29)
RHODE ISLAND	1845	10.49	0.87	(8.78, 12.20)
SOUTH CAROLINA	1868	16.31	1.21	(13.94, 18.68)
SOUTH DAKOTA	2100	10.12	0.75	(8.65, 11.59)
TENNESSEE	3052	11.98	0.67	(10.67, 13.29)
TEXAS	1694	24.60	1.20	(22.25, 26.95)
UTAH	2873	12.18	0.76	(10.69, 13.67)
VERMONT	2497	12.98	0.80	(11.41, 14.55)
VIRGINIA	1891	14.57	0.99	(12.63, 16.51)
WASHINGTON	3582	11.94	0.62	(10.72, 13.16)
WEST VIRGINIA	2412	17.69	0.90	(15.93, 19.45)
WISCONSIN	2226	9.30	0.85	(7.63, 10.97)
WYOMING	2522	18.33	0.87	(16.62, 20.04)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 12.94 RANGE = 7.21-24.60
 * HAVING NO HEALTH CARE PLAN AGES 18 AND OVER

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
DENOMINATOR IS PERSONS AGE 18 AND OVER
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	779	15.97	1.51	(13.01, 18.93)	1356	15.99	1.09	(13.85, 18.13)
ALASKA	710	24.09	2.48	(19.23, 28.95)	820	17.09	1.77	(13.62, 20.56)
ARIZONA	809	18.77	1.98	(14.89, 22.65)	1134	15.31	1.54	(12.29, 18.33)
ARKANSAS	647	18.08	1.66	(14.83, 21.33)	1149	17.80	1.33	(15.19, 20.41)
CALIFORNIA	1722	22.24	1.21	(19.87, 24.61)	2264	16.83	0.93	(15.01, 18.65)
COLORADO	793	14.83	1.56	(11.77, 17.89)	1010	12.14	1.22	(9.75, 14.53)
CONNECTICUT	729	9.49	1.24	(7.06, 11.92)	1100	9.20	1.15	(6.95, 11.45)
DELAWARE	828	11.87	1.50	(8.93, 14.81)	1294	9.79	0.95	(7.93, 11.65)
DISTRICT OF CO	459	13.91	1.97	(10.05, 17.77)	737	12.95	1.43	(10.15, 15.75)
FLORIDA	1440	18.39	1.17	(16.10, 20.68)	2131	16.78	0.93	(14.96, 18.60)
GEORGIA	1018	10.83	1.20	(8.48, 13.18)	1361	8.79	0.87	(7.08, 10.50)
IDAHO	1114	15.58	1.22	(13.19, 17.97)	1614	14.44	0.94	(12.60, 16.28)
ILLINOIS	1152	12.83	1.15	(10.58, 15.08)	1727	10.93	0.85	(9.26, 12.60)
INDIANA	937	11.11	1.15	(8.86, 13.36)	1268	11.69	1.00	(9.73, 13.65)
IOWA	1477	10.94	0.91	(9.16, 12.72)	2120	7.55	0.65	(6.28, 8.82)
KANSAS	858	10.06	1.22	(7.67, 12.45)	1146	10.12	1.03	(8.10, 12.14)
KENTUCKY	1317	14.43	1.13	(12.22, 16.64)	2292	14.41	0.88	(12.69, 16.13)
LOUISIANA	652	23.93	1.98	(20.05, 27.81)	994	21.86	1.51	(18.90, 24.82)
MAINE	708	13.69	1.44	(10.87, 16.51)	988	14.72	1.38	(12.02, 17.42)
MARYLAND	1835	10.98	1.03	(8.96, 13.00)	2639	8.61	0.69	(7.26, 9.96)
MASSACHUSETTS	739	12.15	1.43	(9.35, 14.95)	1041	8.43	1.07	(6.33, 10.53)
MICHIGAN	1043	10.67	1.05	(8.61, 12.73)	1438	8.56	0.81	(6.97, 10.15)
MINNESOTA	1954	8.28	0.67	(6.97, 9.59)	2441	6.21	0.58	(5.07, 7.35)
MISSISSIPPI	587	13.87	1.72	(10.50, 17.24)	999	14.51	1.26	(12.04, 16.98)
MISSOURI	620	13.78	1.63	(10.59, 16.97)	929	12.17	1.13	(9.96, 14.38)
MONTANA	769	14.11	1.35	(11.46, 16.76)	1028	17.03	1.32	(14.44, 19.62)
NEBRASKA	834	10.02	1.17	(7.73, 12.31)	1257	7.25	0.85	(5.58, 8.92)
NEVADA	756	18.54	2.09	(14.44, 22.64)	1039	14.30	1.50	(11.36, 17.24)
NEW HAMPSHIRE	647	13.39	1.50	(10.45, 16.33)	853	10.47	1.26	(8.00, 12.94)
NEW JERSEY	1329	13.52	1.30	(10.97, 16.07)	1808	8.99	0.77	(7.48, 10.50)
NEW MEXICO	432	22.78	2.51	(17.86, 27.70)	660	19.39	2.02	(15.43, 23.35)
NEW YORK	1740	15.19	1.00	(13.23, 17.15)	2560	9.99	0.67	(8.68, 11.30)
NORTH CAROLINA	1142	12.91	1.22	(10.52, 15.30)	1647	12.98	0.99	(11.04, 14.92)
NORTH DAKOTA	736	12.65	1.41	(9.89, 15.41)	1069	10.90	1.10	(8.74, 13.06)
OHIO	625	13.40	1.76	(9.95, 16.85)	927	9.54	1.12	(7.34, 11.74)
OKLAHOMA	854	18.35	1.76	(14.90, 21.80)	1069	18.57	1.44	(15.75, 21.39)
OREGON	1229	15.05	1.27	(12.56, 17.54)	1690	11.46	0.90	(9.70, 13.22)
PENNSYLVANIA	1516	13.14	1.01	(11.16, 15.12)	2062	9.24	0.75	(7.77, 10.71)
RHODE ISLAND	792	11.91	1.23	(9.50, 14.32)	1053	9.24	1.25	(6.79, 11.69)
SOUTH CAROLINA	740	15.42	1.96	(11.58, 19.26)	1128	17.11	1.47	(14.23, 19.99)
SOUTH DAKOTA	892	10.67	1.18	(8.36, 12.98)	1208	9.60	0.92	(7.80, 11.40)
TENNESSEE	1173	13.82	1.11	(11.64, 16.00)	1879	10.33	0.80	(8.76, 11.90)
TEXAS	661	23.47	1.86	(19.82, 27.12)	1033	25.66	1.53	(22.66, 28.66)
UTAH	1249	11.33	1.06	(9.25, 13.41)	1624	12.98	1.08	(10.86, 15.10)
VERMONT	1071	14.35	1.25	(11.90, 16.80)	1426	11.70	1.02	(9.70, 13.70)
VIRGINIA	732	13.61	1.59	(10.49, 16.73)	1159	15.47	1.23	(13.06, 17.88)
WASHINGTON	1518	14.65	1.03	(12.63, 16.67)	2064	9.34	0.70	(7.97, 10.71)
WEST VIRGINIA	900	19.45	1.44	(16.63, 22.27)	1512	16.14	1.11	(13.96, 18.32)
WISCONSIN	946	10.37	1.32	(7.78, 12.96)	1280	8.29	1.10	(6.13, 10.45)
WYOMING	1046	19.83	1.33	(17.22, 22.44)	1476	16.86	1.06	(14.78, 18.94)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 13.80	MEDIAN	= 11.92
	RANGE	= 8.28-24.09	RANGE	= 6.21-25.66

* HAVING NO HEALTH CARE PLAN AGES 18 AND OVER November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.26	15.98	1.72
ALASKA	15.33	20.76	5.43
ARIZONA	15.20	17.00	1.80
ARKANSAS	15.40	17.93	2.53
CALIFORNIA	19.52	19.51	-0.01
COLORADO	14.19	13.46	-0.73
CONNECTICUT	9.75	9.34	-0.41
DELAWARE	11.35	10.78	-0.57
DISTRICT OF CO	N/A	13.39	N/A
FLORIDA	14.98	17.55	2.57
GEORGIA	10.18	9.76	-0.42
IDAHO	14.30	15.00	0.70
ILLINOIS	11.17	11.84	0.67
INDIANA	10.21	11.42	1.21
IOWA	9.16	9.16	0.00
KANSAS	11.09	10.09	-1.00
KENTUCKY	14.28	14.42	0.14
LOUISIANA	20.96	22.83	1.87
MAINE	16.35	14.22	-2.13
MARYLAND	9.44	9.74	0.30
MASSACHUSETTS	10.47	10.19	-0.28
MICHIGAN	8.55	9.56	1.01
MINNESOTA	8.18	7.21	-0.97
MISSISSIPPI	13.98	14.21	0.23
MISSOURI	15.03	12.93	-2.10
MONTANA	16.31	15.61	-0.70
NEBRASKA	7.82	8.57	0.75
NEVADA	13.11	16.43	3.32
NEW HAMPSHIRE	12.22	11.88	-0.34
NEW JERSEY	8.08	11.14	3.06
NEW MEXICO	18.57	21.02	2.45
NEW YORK	12.25	12.43	0.18
NORTH CAROLINA	12.55	12.94	0.39
NORTH DAKOTA	10.57	11.76	1.19
OHIO	11.03	11.37	0.34
OKLAHOMA	14.92	18.47	3.55
OREGON	12.96	13.20	0.24
PENNSYLVANIA	9.28	11.07	1.79
RHODE ISLAND	10.67	10.49	-0.18
SOUTH CAROLINA	12.06	16.31	4.25
SOUTH DAKOTA	8.96	10.12	1.16
TENNESSEE	11.37	11.98	0.61
TEXAS	17.26	24.60	7.34
UTAH	10.84	12.18	1.34
VERMONT	11.87	12.98	1.11
VIRGINIA	11.41	14.57	3.16
WASHINGTON	11.49	11.94	0.45
WEST VIRGINIA	15.96	17.69	1.73
WISCONSIN	7.77	9.30	1.53
WYOMING	16.25	18.33	2.08

* HAVING NO HEALTH CARE PLAN
 N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1728	18.80	1.08	(16.68, 20.92)
ALASKA	1388	21.84	1.67	(18.57, 25.11)
ARIZONA	1478	20.47	1.46	(17.61, 23.33)
ARKANSAS	1379	21.70	1.31	(19.13, 24.27)
CALIFORNIA	3302	22.62	0.88	(20.90, 24.34)
COLORADO	1521	15.34	1.11	(13.16, 17.52)
CONNECTICUT	1481	10.97	1.01	(8.99, 12.95)
DELAWARE	1701	13.01	1.08	(10.89, 15.13)
DISTRICT OF CO	972	15.36	1.40	(12.62, 18.10)
FLORIDA	2632	22.91	0.95	(21.05, 24.77)
GEORGIA	1903	11.02	0.83	(9.39, 12.65)
IDAHO	2218	18.15	0.93	(16.33, 19.97)
ILLINOIS	2390	14.13	0.84	(12.48, 15.78)
INDIANA	1759	13.55	0.90	(11.79, 15.31)
IOWA	2729	11.61	0.71	(10.22, 13.00)
KANSAS	1542	12.29	0.97	(10.39, 14.19)
KENTUCKY	2689	17.40	0.86	(15.71, 19.09)
LOUISIANA	1308	26.34	1.44	(23.52, 29.16)
MAINE	1331	17.16	1.22	(14.77, 19.55)
MARYLAND	3600	11.41	0.71	(10.02, 12.80)
MASSACHUSETTS	1476	12.49	1.06	(10.41, 14.57)
MICHIGAN	2093	11.09	0.77	(9.58, 12.60)
MINNESOTA	3619	8.03	0.50	(7.05, 9.01)
MISSISSIPPI	1250	17.25	1.24	(14.82, 19.68)
MISSOURI	1202	15.89	1.19	(13.56, 18.22)
MONTANA	1403	19.18	1.14	(16.95, 21.41)
NEBRASKA	1578	10.67	0.89	(8.93, 12.41)
NEVADA	1493	19.20	1.51	(16.24, 22.16)
NEW HAMPSHIRE	1230	14.13	1.17	(11.84, 16.42)
NEW JERSEY	2553	13.09	0.89	(11.35, 14.83)
NEW MEXICO	892	24.59	1.83	(21.00, 28.18)
NEW YORK	3559	15.01	0.71	(13.62, 16.40)
NORTH CAROLINA	2095	15.03	0.92	(13.23, 16.83)
NORTH DAKOTA	1374	14.43	1.11	(12.25, 16.61)
OHIO	1144	13.47	1.24	(11.04, 15.90)
OKLAHOMA	1291	23.41	1.40	(20.67, 26.15)
OREGON	2349	15.89	0.92	(14.09, 17.69)
PENNSYLVANIA	2788	13.45	0.76	(11.96, 14.94)
RHODE ISLAND	1465	12.98	1.08	(10.86, 15.10)
SOUTH CAROLINA	1521	18.89	1.41	(16.13, 21.65)
SOUTH DAKOTA	1561	12.73	0.94	(10.89, 14.57)
TENNESSEE	2438	13.89	0.80	(12.32, 15.46)
TEXAS	1418	28.33	1.37	(25.64, 31.02)
UTAH	2362	14.04	0.88	(12.32, 15.76)
VERMONT	2040	15.29	0.94	(13.45, 17.13)
VIRGINIA	1595	16.48	1.14	(14.25, 18.71)
WASHINGTON	2983	14.00	0.73	(12.57, 15.43)
WEST VIRGINIA	1837	22.15	1.10	(19.99, 24.31)
WISCONSIN	1821	11.42	1.04	(9.38, 13.46)
WYOMING	2098	21.56	1.02	(19.56, 23.56)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 15.16 RANGE = 8.03-28.33 YEAR 2000 OBJECTIVE 21.4
 * HAVING NO HEALTH CARE PLAN November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	653	18.15	1.77	(14.68, 21.62)	1075	19.41	1.32	(16.82, 22.00)
ALASKA	652	25.07	2.61	(19.95, 30.19)	736	18.16	1.92	(14.40, 21.92)
ARIZONA	632	22.18	2.29	(17.69, 26.67)	846	18.74	1.90	(15.02, 22.46)
ARKANSAS	518	20.98	1.99	(17.08, 24.88)	861	22.40	1.66	(19.15, 25.65)
CALIFORNIA	1461	25.22	1.35	(22.57, 27.87)	1841	19.91	1.09	(17.77, 22.05)
COLORADO	694	16.73	1.75	(13.30, 20.16)	827	13.94	1.35	(11.29, 16.59)
CONNECTICUT	607	11.02	1.45	(8.18, 13.86)	874	10.92	1.42	(8.14, 13.70)
DELAWARE	679	13.81	1.75	(10.38, 17.24)	1022	12.25	1.18	(9.94, 14.56)
DISTRICT OF CO	383	15.43	2.26	(11.00, 19.86)	589	15.30	1.73	(11.91, 18.69)
FLORIDA	1081	23.28	1.44	(20.46, 26.10)	1551	22.55	1.21	(20.18, 24.92)
GEORGIA	814	11.69	1.33	(9.08, 14.30)	1089	10.37	1.03	(8.35, 12.39)
IDAHO	950	18.38	1.42	(15.60, 21.16)	1268	17.92	1.16	(15.65, 20.19)
ILLINOIS	986	14.86	1.31	(12.29, 17.43)	1404	13.41	1.04	(11.37, 15.45)
INDIANA	785	12.79	1.32	(10.20, 15.38)	974	14.28	1.20	(11.93, 16.63)
IOWA	1201	13.22	1.10	(11.06, 15.38)	1528	10.01	0.86	(8.32, 11.70)
KANSAS	701	11.98	1.45	(9.14, 14.82)	841	12.61	1.29	(10.08, 15.14)
KENTUCKY	1044	16.82	1.32	(14.23, 19.41)	1645	17.96	1.09	(15.82, 20.10)
LOUISIANA	534	26.50	2.24	(22.11, 30.89)	774	26.19	1.81	(22.64, 29.74)
MAINE	579	15.93	1.66	(12.68, 19.18)	752	18.37	1.71	(15.02, 21.72)
MARYLAND	1529	12.54	1.18	(10.23, 14.85)	2071	10.31	0.82	(8.70, 11.92)
MASSACHUSETTS	632	14.33	1.67	(11.06, 17.60)	844	10.69	1.34	(8.06, 13.32)
MICHIGAN	923	12.09	1.20	(9.74, 14.44)	1170	10.12	0.98	(8.20, 12.04)
MINNESOTA	1680	9.15	0.74	(7.70, 10.60)	1939	6.91	0.68	(5.58, 8.24)
MISSISSIPPI	484	16.39	2.00	(12.47, 20.31)	766	18.07	1.55	(15.03, 21.11)
MISSOURI	515	16.79	1.93	(13.01, 20.57)	687	15.03	1.41	(12.27, 17.79)
MONTANA	628	16.58	1.60	(13.44, 19.72)	775	21.81	1.62	(18.63, 24.99)
NEBRASKA	663	12.13	1.41	(9.37, 14.89)	915	9.23	1.10	(7.07, 11.39)
NEVADA	625	20.95	2.38	(16.29, 25.61)	868	17.36	1.80	(13.83, 20.89)
NEW HAMPSHIRE	548	15.42	1.71	(12.07, 18.77)	682	12.84	1.53	(9.84, 15.84)
NEW JERSEY	1116	15.36	1.50	(12.42, 18.30)	1437	10.87	0.96	(8.99, 12.75)
NEW MEXICO	366	26.48	2.89	(20.82, 32.14)	526	22.76	2.32	(18.21, 27.31)
NEW YORK	1489	17.83	1.17	(15.54, 20.12)	2070	12.33	0.81	(10.74, 13.92)
NORTH CAROLINA	917	14.34	1.40	(11.60, 17.08)	1178	15.70	1.22	(13.31, 18.09)
NORTH DAKOTA	605	15.24	1.67	(11.97, 18.51)	769	13.60	1.39	(10.88, 16.32)
OHIO	482	15.20	2.04	(11.20, 19.20)	662	11.81	1.39	(9.09, 14.53)
OKLAHOMA	564	22.93	2.16	(18.70, 27.16)	727	23.86	1.82	(20.29, 27.43)
OREGON	1026	17.70	1.47	(14.82, 20.58)	1323	14.07	1.11	(11.89, 16.25)
PENNSYLVANIA	1232	15.28	1.18	(12.97, 17.59)	1556	11.68	0.96	(9.80, 13.56)
RHODE ISLAND	670	14.14	1.45	(11.30, 16.98)	795	11.87	1.59	(8.75, 14.99)
SOUTH CAROLINA	642	17.87	2.25	(13.46, 22.28)	879	19.88	1.75	(16.45, 23.31)
SOUTH DAKOTA	708	12.83	1.42	(10.05, 15.61)	853	12.63	1.20	(10.28, 14.98)
TENNESSEE	972	15.18	1.27	(12.69, 17.67)	1466	12.65	0.98	(10.73, 14.57)
TEXAS	572	26.41	2.07	(22.35, 30.47)	846	30.26	1.77	(26.79, 33.73)
UTAH	1062	12.81	1.22	(10.42, 15.20)	1300	15.24	1.26	(12.77, 17.71)
VERMONT	910	16.21	1.43	(13.41, 19.01)	1130	14.37	1.24	(11.94, 16.80)
VIRGINIA	641	15.09	1.78	(11.60, 18.58)	954	17.87	1.43	(15.07, 20.67)
WASHINGTON	1296	16.56	1.17	(14.27, 18.85)	1687	11.39	0.86	(9.70, 13.08)
WEST VIRGINIA	733	23.42	1.70	(20.09, 26.75)	1104	20.93	1.41	(18.17, 23.69)
WISCONSIN	799	12.29	1.56	(9.23, 15.35)	1022	10.55	1.38	(7.85, 13.25)
WYOMING	900	22.48	1.51	(19.52, 25.44)	1198	20.60	1.26	(18.13, 23.07)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 15.68	MEDIAN	= 14.33
	RANGE	= 9.15-26.50	RANGE	= 6.91-30.26

* HAVING NO HEALTH CARE PLAN

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	16.46	18.80	2.34
ALASKA	16.41	21.84	5.43
ARIZONA	18.59	20.47	1.88
ARKANSAS	18.70	21.70	3.00
CALIFORNIA	22.61	22.62	0.01
COLORADO	16.50	15.34	-1.16
CONNECTICUT	11.49	10.97	-0.52
DELAWARE	13.41	13.01	-0.40
DISTRICT OF CO	N/A	15.36	N/A
FLORIDA	19.22	22.91	3.69
GEORGIA	11.27	11.02	-0.25
IDAHO	16.86	18.15	1.29
ILLINOIS	13.42	14.13	0.71
INDIANA	12.12	13.55	1.43
IOWA	11.58	11.61	0.03
KANSAS	13.40	12.29	-1.11
KENTUCKY	16.90	17.40	0.50
LOUISIANA	24.53	26.34	1.81
MAINE	19.57	17.16	-2.41
MARYLAND	10.74	11.41	0.67
MASSACHUSETTS	12.35	12.49	0.14
MICHIGAN	9.96	11.09	1.13
MINNESOTA	9.52	8.03	-1.49
MISSISSIPPI	16.73	17.25	0.52
MISSOURI	18.23	15.89	-2.34
MONTANA	19.49	19.18	-0.31
NEBRASKA	9.76	10.67	0.91
NEVADA	15.24	19.20	3.96
NEW HAMPSHIRE	14.40	14.13	-0.27
NEW JERSEY	9.62	13.09	3.47
NEW MEXICO	21.58	24.59	3.01
NEW YORK	14.33	15.01	0.68
NORTH CAROLINA	14.67	15.03	0.36
NORTH DAKOTA	12.77	14.43	1.66
OHIO	13.14	13.47	0.33
OKLAHOMA	18.07	23.41	5.34
OREGON	15.72	15.89	0.17
PENNSYLVANIA	11.48	13.45	1.97
RHODE ISLAND	13.00	12.98	-0.02
SOUTH CAROLINA	14.09	18.89	4.80
SOUTH DAKOTA	10.84	12.73	1.89
TENNESSEE	13.21	13.89	0.68
TEXAS	19.93	28.33	8.40
UTAH	12.62	14.04	1.42
VERMONT	13.83	15.29	1.46
VIRGINIA	13.05	16.48	3.43
WASHINGTON	13.55	14.00	0.45
WEST VIRGINIA	19.92	22.15	2.23
WISCONSIN	9.28	11.42	2.14
WYOMING	18.97	21.56	2.59

* HAVING NO HEALTH CARE PLAN
 N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2137	7.32	0.65	(6.05, 8.59)
ALASKA	1535	3.49	0.77	(1.98, 5.00)
ARIZONA	1954	3.55	0.55	(2.47, 4.63)
ARKANSAS	1800	4.61	0.55	(3.53, 5.69)
CALIFORNIA	3993	5.10	0.42	(4.28, 5.92)
COLORADO	1803	3.02	0.43	(2.18, 3.86)
CONNECTICUT	1836	4.38	0.51	(3.38, 5.38)
DELAWARE	2122	5.31	0.50	(4.33, 6.29)
DISTRICT OF CO	1195	5.60	0.78	(4.07, 7.13)
FLORIDA	3573	5.19	0.42	(4.37, 6.01)
GEORGIA	2382	4.13	0.44	(3.27, 4.99)
IDAHO	2735	3.46	0.40	(2.68, 4.24)
ILLINOIS	2875	5.76	0.49	(4.80, 6.72)
INDIANA	2210	5.31	0.53	(4.27, 6.35)
IOWA	3598	4.21	0.34	(3.54, 4.88)
KANSAS	2000	3.61	0.47	(2.69, 4.53)
KENTUCKY	3615	4.49	0.38	(3.75, 5.23)
LOUISIANA	1653	6.56	0.66	(5.27, 7.85)
MAINE	1699	4.24	0.51	(3.24, 5.24)
MARYLAND	4478	5.14	0.42	(4.32, 5.96)
MASSACHUSETTS	1780	5.25	0.61	(4.05, 6.45)
MICHIGAN	2487	5.81	0.53	(4.77, 6.85)
MINNESOTA	4400	4.46	0.34	(3.79, 5.13)
MISSISSIPPI	1591	6.11	0.63	(4.88, 7.34)
MISSOURI	1550	4.36	0.55	(3.28, 5.44)
MONTANA	1802	3.70	0.45	(2.82, 4.58)
NEBRASKA	2100	5.28	0.52	(4.26, 6.30)
NEVADA	1797	4.17	0.61	(2.97, 5.37)
NEW HAMPSHIRE	1502	3.98	0.52	(2.96, 5.00)
NEW JERSEY	3142	4.60	0.44	(3.74, 5.46)
NEW MEXICO	1096	4.38	0.71	(2.99, 5.77)
NEW YORK	4311	3.60	0.33	(2.95, 4.25)
NORTH CAROLINA	2792	4.44	0.42	(3.62, 5.26)
NORTH DAKOTA	1811	3.07	0.42	(2.25, 3.89)
OHIO	1555	4.98	0.57	(3.86, 6.10)
OKLAHOMA	1923	3.78	0.51	(2.78, 4.78)
OREGON	2928	4.46	0.41	(3.66, 5.26)
PENNSYLVANIA	3589	5.65	0.43	(4.81, 6.49)
RHODE ISLAND	1846	4.47	0.53	(3.43, 5.51)
SOUTH CAROLINA	1863	5.43	0.62	(4.21, 6.65)
SOUTH DAKOTA	2106	4.25	0.45	(3.37, 5.13)
TENNESSEE	3056	5.02	0.43	(4.18, 5.86)
TEXAS	1705	3.74	0.49	(2.78, 4.70)
UTAH	2885	3.29	0.39	(2.53, 4.05)
VERMONT	2503	3.90	0.41	(3.10, 4.70)
VIRGINIA	1887	5.48	0.57	(4.36, 6.60)
WASHINGTON	3589	3.42	0.33	(2.77, 4.07)
WEST VIRGINIA	2418	5.85	0.52	(4.83, 6.87)
WISCONSIN	2228	4.72	0.68	(3.39, 6.05)
WYOMING	2530	4.22	0.42	(3.40, 5.04)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 4.46 RANGE = 3.02- 7.32
 * EVER TOLD YOU HAVE DIABETES November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS* --BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	780	6.67	0.97	(4.77, 8.57)	*	1357	7.90	0.83	(6.27, 9.53)
ALASKA	715	3.45	1.10	(1.29, 5.61)	*	820	3.54	1.06	(1.46, 5.62)
ARIZONA	814	4.23	0.91	(2.45, 6.01)	*	1140	2.90	0.65	(1.63, 4.17)
ARKANSAS	649	4.09	0.80	(2.52, 5.66)	*	1151	5.08	0.69	(3.73, 6.43)
CALIFORNIA	1723	5.26	0.64	(4.01, 6.51)	*	2270	4.95	0.55	(3.87, 6.03)
COLORADO	793	3.32	0.69	(1.97, 4.67)	*	1010	2.73	0.51	(1.73, 3.73)
CONNECTICUT	733	3.15	0.66	(1.86, 4.44)	*	1103	5.49	0.77	(3.98, 7.00)
DELAWARE	827	5.94	0.83	(4.31, 7.57)	*	1295	4.74	0.60	(3.56, 5.92)
DISTRICT OF CO	459	5.99	1.28	(3.48, 8.50)	*	736	5.27	0.94	(3.43, 7.11)
FLORIDA	1441	5.70	0.68	(4.37, 7.03)	*	2132	4.72	0.50	(3.74, 5.70)
GEORGIA	1020	3.75	0.61	(2.55, 4.95)	*	1362	4.48	0.63	(3.25, 5.71)
IDAHO	1120	4.03	0.63	(2.80, 5.26)	*	1615	2.91	0.46	(2.01, 3.81)
ILLINOIS	1150	5.66	0.73	(4.23, 7.09)	*	1725	5.85	0.67	(4.54, 7.16)
INDIANA	942	6.09	0.89	(4.35, 7.83)	*	1268	4.61	0.64	(3.36, 5.86)
IOWA	1478	3.76	0.51	(2.76, 4.76)	*	2120	4.61	0.47	(3.69, 5.53)
KANSAS	854	3.21	0.63	(1.98, 4.44)	*	1146	3.98	0.69	(2.63, 5.33)
KENTUCKY	1320	4.23	0.57	(3.11, 5.35)	*	2295	4.72	0.48	(3.78, 5.66)
LOUISIANA	658	5.40	0.89	(3.66, 7.14)	*	995	7.59	1.00	(5.63, 9.55)
MAINE	709	3.68	0.68	(2.35, 5.01)	*	990	4.75	0.72	(3.34, 6.16)
MARYLAND	1836	4.92	0.60	(3.74, 6.10)	*	2642	5.35	0.59	(4.19, 6.51)
MASSACHUSETTS	738	4.65	0.90	(2.89, 6.41)	*	1042	5.79	0.84	(4.14, 7.44)
MICHIGAN	1045	5.28	0.77	(3.77, 6.79)	*	1442	6.29	0.72	(4.88, 7.70)
MINNESOTA	1958	4.35	0.50	(3.37, 5.33)	*	2442	4.57	0.46	(3.67, 5.47)
MISSISSIPPI	589	5.83	0.99	(3.89, 7.77)	*	1002	6.35	0.80	(4.78, 7.92)
MISSOURI	620	3.91	0.80	(2.34, 5.48)	*	930	4.76	0.76	(3.27, 6.25)
MONTANA	771	3.68	0.69	(2.33, 5.03)	*	1031	3.71	0.60	(2.53, 4.89)
NEBRASKA	839	5.57	0.84	(3.92, 7.22)	*	1261	5.01	0.64	(3.76, 6.26)
NEVADA	760	5.77	1.05	(3.71, 7.83)	*	1037	2.54	0.60	(1.36, 3.72)
NEW HAMPSHIRE	647	3.43	0.71	(2.04, 4.82)	*	855	4.51	0.77	(3.00, 6.02)
NEW JERSEY	1331	4.70	0.67	(3.39, 6.01)	*	1811	4.50	0.58	(3.36, 5.64)
NEW MEXICO	435	4.98	1.25	(2.53, 7.43)	*	661	3.82	0.79	(2.27, 5.37)
NEW YORK	1749	3.86	0.53	(2.82, 4.90)	*	2562	3.36	0.41	(2.56, 4.16)
NORTH CAROLINA	1144	4.22	0.63	(2.99, 5.45)	*	1648	4.64	0.55	(3.56, 5.72)
NORTH DAKOTA	741	2.29	0.59	(1.13, 3.45)	*	1070	3.82	0.61	(2.62, 5.02)
OHIO	626	3.86	0.75	(2.39, 5.33)	*	929	5.98	0.83	(4.35, 7.61)
OKLAHOMA	855	3.12	0.65	(1.85, 4.39)	*	1068	4.39	0.75	(2.92, 5.86)
OREGON	1234	4.01	0.59	(2.85, 5.17)	*	1694	4.88	0.59	(3.72, 6.04)
PENNSYLVANIA	1525	4.77	0.59	(3.61, 5.93)	*	2064	6.43	0.62	(5.21, 7.65)
RHODE ISLAND	793	4.76	0.84	(3.11, 6.41)	*	1053	4.21	0.66	(2.92, 5.50)
SOUTH CAROLINA	737	5.18	0.93	(3.36, 7.00)	*	1126	5.65	0.77	(4.14, 7.16)
SOUTH DAKOTA	896	4.11	0.67	(2.80, 5.42)	*	1210	4.39	0.62	(3.17, 5.61)
TENNESSEE	1176	4.20	0.61	(3.00, 5.40)	*	1880	5.76	0.62	(4.54, 6.98)
TEXAS	667	3.40	0.77	(1.89, 4.91)	*	1038	4.06	0.62	(2.84, 5.28)
UTAH	1254	3.65	0.57	(2.53, 4.77)	*	1631	2.94	0.54	(1.88, 4.00)
VERMONT	1073	3.99	0.62	(2.77, 5.21)	*	1430	3.81	0.53	(2.77, 4.85)
VIRGINIA	728	4.03	0.74	(2.58, 5.48)	*	1159	6.83	0.85	(5.16, 8.50)
WASHINGTON	1522	3.02	0.46	(2.12, 3.92)	*	2067	3.80	0.47	(2.88, 4.72)
WEST VIRGINIA	904	5.89	0.80	(4.32, 7.46)	*	1514	5.82	0.66	(4.53, 7.11)
WISCONSIN	945	5.34	1.20	(2.99, 7.69)	*	1283	4.14	0.68	(2.81, 5.47)
WYOMING	1050	3.64	0.60	(2.46, 4.82)	*	1480	4.78	0.60	(3.60, 5.96)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 4.21	MEDIAN	= 4.68
	RANGE	= 2.29- 6.67	RANGE	= 2.54- 7.90

* EVER TOLD YOU HAVE DIABETES

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	4.79	7.32	2.53
ALASKA	2.68	3.49	0.81
ARIZONA	4.81	3.55	-1.26
ARKANSAS	5.18	4.61	-0.57
CALIFORNIA	5.60	5.10	-0.50
COLORADO	3.52	3.02	-0.50
CONNECTICUT	4.31	4.38	0.07
DELAWARE	4.34	5.31	0.97
DISTRICT OF CO	N/A	5.60	N/A
FLORIDA	5.28	5.19	-0.09
GEORGIA	3.66	4.13	0.47
IDAHO	3.55	3.46	-0.09
ILLINOIS	5.42	5.76	0.34
INDIANA	5.01	5.31	0.30
IOWA	5.37	4.21	-1.16
KANSAS	4.84	3.61	-1.23
KENTUCKY	3.55	4.49	0.94
LOUISIANA	6.19	6.56	0.37
MAINE	3.51	4.24	0.73
MARYLAND	4.05	5.14	1.09
MASSACHUSETTS	3.84	5.25	1.41
MICHIGAN	5.34	5.81	0.47
MINNESOTA	3.08	4.46	1.38
MISSISSIPPI	6.37	6.11	-0.26
MISSOURI	4.14	4.36	0.22
MONTANA	2.84	3.70	0.86
NEBRASKA	4.44	5.28	0.84
NEVADA	4.82	4.17	-0.65
NEW HAMPSHIRE	4.46	3.98	-0.48
NEW JERSEY	4.20	4.60	0.40
NEW MEXICO	5.58	4.38	-1.20
NEW YORK	4.16	3.60	-0.56
NORTH CAROLINA	4.51	4.44	-0.07
NORTH DAKOTA	3.71	3.07	-0.64
OHIO	4.18	4.98	0.80
OKLAHOMA	2.94	3.78	0.84
OREGON	4.03	4.46	0.43
PENNSYLVANIA	5.65	5.65	0.00
RHODE ISLAND	4.59	4.47	-0.12
SOUTH CAROLINA	5.04	5.43	0.39
SOUTH DAKOTA	2.94	4.25	1.31
TENNESSEE	5.21	5.02	-0.19
TEXAS	5.23	3.74	-1.49
UTAH	3.70	3.29	-0.41
VERMONT	4.50	3.90	-0.60
VIRGINIA	3.85	5.48	1.63
WASHINGTON	3.08	3.42	0.34
WEST VIRGINIA	4.72	5.85	1.13
WISCONSIN	4.50	4.72	0.22
WYOMING	3.19	4.22	1.03

* EVER TOLD YOU HAVE DIABETES
 N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2137	43.90	1.26	(41.43, 46.37)
ALASKA	1531	53.74	1.95	(49.92, 57.56)
ARIZONA	1952	48.82	1.60	(45.68, 51.96)
ARKANSAS	1795	43.27	1.41	(40.51, 46.03)
CALIFORNIA	3995	43.59	0.89	(41.85, 45.33)
COLORADO	1800	46.94	1.35	(44.29, 49.59)
CONNECTICUT	1832	49.99	1.31	(47.42, 52.56)
DELAWARE	2121	50.99	1.26	(48.52, 53.46)
DISTRICT OF CO	1199	38.31	1.64	(35.10, 41.52)
FLORIDA	3572	52.51	0.92	(50.71, 54.31)
GEORGIA	2378	34.44	1.12	(32.24, 36.64)
IDAHO	2730	43.99	1.09	(41.85, 46.13)
ILLINOIS	2878	48.21	1.08	(46.09, 50.33)
INDIANA	2210	52.41	1.17	(50.12, 54.70)
IOWA	3599	47.41	0.93	(45.59, 49.23)
KANSAS	2002	41.56	1.21	(39.19, 43.93)
KENTUCKY	3612	54.83	0.97	(52.93, 56.73)
LOUISIANA	1651	47.40	1.34	(44.77, 50.03)
MAINE	1696	53.07	1.33	(50.46, 55.68)
MARYLAND	4471	44.34	1.01	(42.36, 46.32)
MASSACHUSETTS	1775	53.79	1.38	(51.09, 56.49)
MICHIGAN	2484	51.54	1.11	(49.36, 53.72)
MINNESOTA	4400	43.65	0.82	(42.04, 45.26)
MISSISSIPPI	1587	44.34	1.41	(41.58, 47.10)
MISSOURI	1547	52.46	1.35	(49.81, 55.11)
MONTANA	1798	48.68	1.31	(46.11, 51.25)
NEBRASKA	2093	45.71	1.37	(43.02, 48.40)
NEVADA	1797	55.00	1.68	(51.71, 58.29)
NEW HAMPSHIRE	1498	53.91	1.44	(51.09, 56.73)
NEW JERSEY	3132	47.46	1.04	(45.42, 49.50)
NEW MEXICO	1092	47.02	1.85	(43.39, 50.65)
NEW YORK	4304	46.93	0.87	(45.22, 48.64)
NORTH CAROLINA	2787	48.82	1.13	(46.61, 51.03)
NORTH DAKOTA	1805	48.85	1.32	(46.26, 51.44)
OHIO	1553	49.83	1.47	(46.95, 52.71)
OKLAHOMA	1923	42.50	1.36	(39.83, 45.17)
OREGON	2920	51.07	1.07	(48.97, 53.17)
PENNSYLVANIA	3583	47.75	0.94	(45.91, 49.59)
RHODE ISLAND	1846	49.65	1.32	(47.06, 52.24)
SOUTH CAROLINA	1860	44.70	1.45	(41.86, 47.54)
SOUTH DAKOTA	2104	45.90	1.19	(43.57, 48.23)
TENNESSEE	3054	47.84	1.03	(45.82, 49.86)
TEXAS	1701	44.14	1.33	(41.53, 46.75)
UTAH	2885	32.33	1.15	(30.08, 34.58)
VERMONT	2496	53.30	1.16	(51.03, 55.57)
VIRGINIA	1886	47.75	1.33	(45.14, 50.36)
WASHINGTON	3576	50.28	0.94	(48.44, 52.12)
WEST VIRGINIA	2416	49.51	1.11	(47.33, 51.69)
WISCONSIN	2224	50.83	1.41	(48.07, 53.59)
WYOMING	2526	49.68	1.08	(47.56, 51.80)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 48.03 RANGE = 32.33-55.00

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	781	53.60	2.04	(49.60, 57.60)	* 1356	35.32	1.41	(32.56, 38.08)
ALASKA	713	58.47	2.81	(52.96, 63.98)	* 818	48.50	2.63	(43.35, 53.65)
ARIZONA	812	54.95	2.47	(50.11, 59.79)	* 1140	43.01	1.97	(39.15, 46.87)
ARKANSAS	646	51.94	2.43	(47.18, 56.70)	* 1149	35.54	1.63	(32.35, 38.73)
CALIFORNIA	1725	50.81	1.37	(48.12, 53.50)	* 2270	36.52	1.12	(34.32, 38.72)
COLORADO	791	52.64	1.99	(48.74, 56.54)	* 1009	41.52	1.77	(38.05, 44.99)
CONNECTICUT	730	54.14	2.04	(50.14, 58.14)	* 1102	46.23	1.70	(42.90, 49.56)
DELAWARE	828	55.13	1.97	(51.27, 58.99)	* 1293	47.22	1.59	(44.10, 50.34)
DISTRICT OF CO	461	42.13	2.53	(37.17, 47.09)	* 738	35.07	2.07	(31.01, 39.13)
FLORIDA	1441	59.46	1.39	(56.74, 62.18)	* 2131	46.20	1.19	(43.87, 48.53)
GEORGIA	1018	41.04	1.82	(37.47, 44.61)	* 1360	28.40	1.35	(25.75, 31.05)
IDAHO	1118	49.82	1.67	(46.55, 53.09)	* 1612	38.40	1.33	(35.79, 41.01)
ILLINOIS	1152	54.64	1.63	(51.45, 57.83)	* 1726	42.33	1.36	(39.66, 45.00)
INDIANA	942	61.95	1.68	(58.66, 65.24)	* 1268	43.73	1.50	(40.79, 46.67)
IOWA	1478	55.69	1.43	(52.89, 58.49)	* 2121	39.89	1.19	(37.56, 42.22)
KANSAS	856	50.28	1.88	(46.60, 53.96)	* 1146	33.44	1.50	(30.50, 36.38)
KENTUCKY	1319	63.59	1.48	(60.69, 66.49)	* 2293	46.88	1.16	(44.61, 49.15)
LOUISIANA	656	58.23	2.06	(54.19, 62.27)	* 995	37.82	1.63	(34.63, 41.01)
MAINE	708	60.37	1.96	(56.53, 64.21)	* 988	46.37	1.77	(42.90, 49.84)
MARYLAND	1836	49.20	1.53	(46.20, 52.20)	* 2635	39.90	1.27	(37.41, 42.39)
MASSACHUSETTS	736	57.22	2.21	(52.89, 61.55)	* 1039	50.73	1.73	(47.34, 54.12)
MICHIGAN	1045	56.67	1.68	(53.38, 59.96)	* 1439	46.86	1.45	(44.02, 49.70)
MINNESOTA	1961	48.37	1.24	(45.94, 50.80)	* 2439	39.23	1.07	(37.13, 41.33)
MISSISSIPPI	585	57.45	2.32	(52.90, 62.00)	* 1002	32.84	1.60	(29.70, 35.98)
MISSOURI	618	58.50	2.19	(54.21, 62.79)	* 929	47.07	1.69	(43.76, 50.38)
MONTANA	770	52.57	1.86	(48.92, 56.22)	* 1028	44.98	1.74	(41.57, 48.39)
NEBRASKA	836	54.59	2.13	(50.42, 58.76)	* 1257	37.54	1.57	(34.46, 40.62)
NEVADA	759	59.79	2.53	(54.83, 64.75)	* 1038	50.12	2.20	(45.81, 54.43)
NEW HAMPSHIRE	646	56.32	2.19	(52.03, 60.61)	* 852	51.66	1.83	(48.07, 55.25)
NEW JERSEY	1328	53.15	1.61	(49.99, 56.31)	* 1804	42.31	1.33	(39.70, 44.92)
NEW MEXICO	431	55.60	2.80	(50.11, 61.09)	* 661	39.00	2.33	(34.43, 43.57)
NEW YORK	1745	49.49	1.33	(46.88, 52.10)	* 2559	44.65	1.10	(42.49, 46.81)
NORTH CAROLINA	1140	59.05	1.66	(55.80, 62.30)	* 1647	39.51	1.41	(36.75, 42.27)
NORTH DAKOTA	738	56.00	1.92	(52.24, 59.76)	* 1067	41.93	1.71	(38.58, 45.28)
OHIO	625	57.67	2.17	(53.42, 61.92)	* 928	42.81	1.80	(39.28, 46.34)
OKLAHOMA	854	48.31	2.07	(44.25, 52.37)	* 1069	37.18	1.75	(33.75, 40.61)
OREGON	1231	57.55	1.65	(54.32, 60.78)	* 1689	45.00	1.33	(42.39, 47.61)
PENNSYLVANIA	1521	51.85	1.44	(49.03, 54.67)	* 2062	44.12	1.23	(41.71, 46.53)
RHODE ISLAND	794	55.85	1.98	(51.97, 59.73)	* 1052	44.14	1.72	(40.77, 47.51)
SOUTH CAROLINA	738	51.81	2.25	(47.40, 56.22)	* 1122	38.31	1.76	(34.86, 41.76)
SOUTH DAKOTA	895	52.59	1.81	(49.04, 56.14)	* 1209	39.64	1.56	(36.58, 42.70)
TENNESSEE	1174	56.30	1.56	(53.24, 59.36)	* 1880	40.28	1.28	(37.77, 42.79)
TEXAS	665	53.31	2.11	(49.17, 57.45)	* 1036	35.49	1.60	(32.35, 38.63)
UTAH	1254	40.16	1.83	(36.57, 43.75)	* 1631	24.90	1.35	(22.25, 27.55)
VERMONT	1072	59.56	1.76	(56.11, 63.01)	* 1424	47.46	1.49	(44.54, 50.38)
VIRGINIA	731	54.28	2.06	(50.24, 58.32)	* 1155	41.64	1.67	(38.37, 44.91)
WASHINGTON	1516	55.41	1.42	(52.63, 58.19)	* 2060	45.38	1.23	(42.97, 47.79)
WEST VIRGINIA	902	56.91	1.79	(53.40, 60.42)	* 1514	42.96	1.41	(40.20, 45.72)
WISCONSIN	944	58.53	2.07	(54.47, 62.59)	* 1280	43.67	1.86	(40.02, 47.32)
WYOMING	1050	54.64	1.63	(51.45, 57.83)	* 1476	44.82	1.47	(41.94, 47.70)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 55.04	MEDIAN	= 42.32
	RANGE	= 40.16-63.59	RANGE	= 24.90-51.66

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	45.59	43.90	-1.69
ALASKA	52.25	53.74	1.49
ARIZONA	48.47	48.82	0.35
ARKANSAS	48.66	43.27	-5.39
CALIFORNIA	43.58	43.59	0.01
COLORADO	49.13	46.94	-2.19
CONNECTICUT	49.49	49.99	0.50
DELAWARE	51.20	50.99	-0.21
DISTRICT OF CO	N/A	38.31	N/A
FLORIDA	51.63	52.51	0.88
GEORGIA	38.50	34.44	-4.06
IDAHO	43.87	43.99	0.12
ILLINOIS	46.75	48.21	1.46
INDIANA	50.20	52.41	2.21
IOWA	47.01	47.41	0.40
KANSAS	46.22	41.56	-4.66
KENTUCKY	50.06	54.83	4.77
LOUISIANA	47.39	47.40	0.01
MAINE	52.07	53.07	1.00
MARYLAND	41.81	44.34	2.53
MASSACHUSETTS	51.69	53.79	2.10
MICHIGAN	53.92	51.54	-2.38
MINNESOTA	44.02	43.65	-0.37
MISSISSIPPI	44.68	44.34	-0.34
MISSOURI	51.42	52.46	1.04
MONTANA	50.37	48.68	-1.69
NEBRASKA	47.56	45.71	-1.85
NEVADA	52.80	55.00	2.20
NEW HAMPSHIRE	54.01	53.91	-0.10
NEW JERSEY	44.39	47.46	3.07
NEW MEXICO	48.62	47.02	-1.60
NEW YORK	46.70	46.93	0.23
NORTH CAROLINA	48.50	48.82	0.32
NORTH DAKOTA	46.85	48.85	2.00
OHIO	49.88	49.83	-0.05
OKLAHOMA	38.98	42.50	3.52
OREGON	50.16	51.07	0.91
PENNSYLVANIA	52.25	47.75	-4.50
RHODE ISLAND	53.97	49.65	-4.32
SOUTH CAROLINA	44.13	44.70	0.57
SOUTH DAKOTA	45.79	45.90	0.11
TENNESSEE	46.12	47.84	1.72
TEXAS	47.33	44.14	-3.19
UTAH	31.19	32.33	1.14
VERMONT	53.57	53.30	-0.27
VIRGINIA	46.90	47.75	0.85
WASHINGTON	49.96	50.28	0.32
WEST VIRGINIA	48.93	49.51	0.58
WISCONSIN	49.39	50.83	1.44
WYOMING	48.80	49.68	0.88

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME
 N/A - NOT AVAILABLE

November 19, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2137	22.42	1.05	(20.36, 24.48)
ALASKA	1530	27.68	1.73	(24.29, 31.07)
ARIZONA	1951	23.71	1.28	(21.20, 26.22)
ARKANSAS	1795	25.29	1.23	(22.88, 27.70)
CALIFORNIA	3995	18.60	0.70	(17.23, 19.97)
COLORADO	1800	22.79	1.14	(20.56, 25.02)
CONNECTICUT	1831	21.80	1.14	(19.57, 24.03)
DELAWARE	2121	24.22	1.10	(22.06, 26.38)
DISTRICT OF CO	1199	20.50	1.33	(17.89, 23.11)
FLORIDA	3571	21.81	0.78	(20.28, 23.34)
GEORGIA	2378	20.27	0.97	(18.37, 22.17)
IDAHO	2730	21.14	0.86	(19.45, 22.83)
ILLINOIS	2876	24.80	0.91	(23.02, 26.58)
INDIANA	2209	28.63	1.06	(26.55, 30.71)
IOWA	3598	23.59	0.80	(22.02, 25.16)
KANSAS	2002	22.06	1.03	(20.04, 24.08)
KENTUCKY	3611	31.66	0.93	(29.84, 33.48)
LOUISIANA	1649	25.88	1.24	(23.45, 28.31)
MAINE	1696	25.31	1.20	(22.96, 27.66)
MARYLAND	4471	20.90	0.79	(19.35, 22.45)
MASSACHUSETTS	1774	23.35	1.16	(21.08, 25.62)
MICHIGAN	2483	25.63	0.95	(23.77, 27.49)
MINNESOTA	4400	20.58	0.68	(19.25, 21.91)
MISSISSIPPI	1587	23.24	1.22	(20.85, 25.63)
MISSOURI	1547	27.78	1.25	(25.33, 30.23)
MONTANA	1798	21.69	1.10	(19.53, 23.85)
NEBRASKA	2093	22.09	1.31	(19.52, 24.66)
NEVADA	1797	28.16	1.54	(25.14, 31.18)
NEW HAMPSHIRE	1498	24.84	1.40	(22.10, 27.58)
NEW JERSEY	3131	22.72	0.92	(20.92, 24.52)
NEW MEXICO	1092	22.83	1.57	(19.75, 25.91)
NEW YORK	4303	23.25	0.73	(21.82, 24.68)
NORTH CAROLINA	2787	25.69	1.02	(23.69, 27.69)
NORTH DAKOTA	1805	23.38	1.16	(21.11, 25.65)
OHIO	1553	28.40	1.35	(25.75, 31.05)
OKLAHOMA	1923	24.08	1.20	(21.73, 26.43)
OREGON	2920	23.41	0.88	(21.69, 25.13)
PENNSYLVANIA	3581	24.54	0.82	(22.93, 26.15)
RHODE ISLAND	1845	22.51	1.10	(20.35, 24.67)
SOUTH CAROLINA	1859	24.46	1.29	(21.93, 26.99)
SOUTH DAKOTA	2104	20.66	0.96	(18.78, 22.54)
TENNESSEE	3054	27.98	0.94	(26.14, 29.82)
TEXAS	1701	22.91	1.14	(20.68, 25.14)
UTAH	2885	15.91	0.88	(14.19, 17.63)
VERMONT	2496	24.07	1.14	(21.84, 26.30)
VIRGINIA	1885	24.79	1.17	(22.50, 27.08)
WASHINGTON	3575	23.44	0.81	(21.85, 25.03)
WEST VIRGINIA	2416	26.63	1.00	(24.67, 28.59)
WISCONSIN	2223	24.86	1.17	(22.57, 27.15)
WYOMING	2526	24.56	0.98	(22.64, 26.48)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 23.51 RANGE = 15.91-31.66 YEAR 2000 OBJ. 3.4
 * EVER SMOKED 100 CIGARETTES AND A CURRENT SMOKER (EVERYDAY & SOME DAYS) December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2066	33.89	1.20	(31.54, 36.24)
ALASKA	1484	30.03	1.81	(26.48, 33.58)
ARIZONA	1846	26.28	1.52	(23.30, 29.26)
ARKANSAS	1697	31.83	1.31	(29.26, 34.40)
CALIFORNIA	3866	27.73	0.82	(26.12, 29.34)
COLORADO	1765	22.32	1.11	(20.14, 24.50)
CONNECTICUT	1730	27.16	1.21	(24.79, 29.53)
DELAWARE	2043	31.78	1.19	(29.45, 34.11)
DISTRICT OF CO	1163	29.15	1.64	(25.94, 32.36)
FLORIDA	3474	30.54	0.90	(28.78, 32.30)
GEORGIA	2332	24.07	1.02	(22.07, 26.07)
IDAHO	2642	29.28	0.98	(27.36, 31.20)
ILLINOIS	2809	29.84	0.96	(27.96, 31.72)
INDIANA	2137	32.70	1.11	(30.52, 34.88)
IOWA	3524	32.73	0.88	(31.01, 34.45)
KANSAS	1883	26.06	1.11	(23.88, 28.24)
KENTUCKY	3470	32.96	0.92	(31.16, 34.76)
LOUISIANA	1591	33.39	1.36	(30.72, 36.06)
MAINE	1632	28.14	1.18	(25.83, 30.45)
MARYLAND	4264	30.78	0.93	(28.96, 32.60)
MASSACHUSETTS	1703	26.67	1.22	(24.28, 29.06)
MICHIGAN	2407	33.03	1.06	(30.95, 35.11)
MINNESOTA	4296	27.91	0.76	(26.42, 29.40)
MISSISSIPPI	1545	34.52	1.42	(31.74, 37.30)
MISSOURI	1501	33.11	1.38	(30.41, 35.81)
MONTANA	1742	28.84	1.24	(26.41, 31.27)
NEBRASKA	2002	29.00	1.17	(26.71, 31.29)
NEVADA	1743	27.67	1.53	(24.67, 30.67)
NEW HAMPSHIRE	1436	25.99	1.32	(23.40, 28.58)
NEW JERSEY	2987	27.53	0.92	(25.73, 29.33)
NEW MEXICO	1062	26.82	1.73	(23.43, 30.21)
NEW YORK	4093	28.96	0.82	(27.35, 30.57)
NORTH CAROLINA	2632	30.65	1.02	(28.65, 32.65)
NORTH DAKOTA	1773	32.87	1.23	(30.46, 35.28)
OHIO	1442	33.43	1.49	(30.51, 36.35)
OKLAHOMA	1872	29.05	1.26	(26.58, 31.52)
OREGON	2784	29.28	0.99	(27.34, 31.22)
PENNSYLVANIA	3415	31.76	0.90	(30.00, 33.52)
RHODE ISLAND	1767	28.48	1.20	(26.13, 30.83)
SOUTH CAROLINA	1759	34.68	1.43	(31.88, 37.48)
SOUTH DAKOTA	1985	30.10	1.14	(27.87, 32.33)
TENNESSEE	2905	30.77	0.99	(28.83, 32.71)
TEXAS	1636	31.34	1.26	(28.87, 33.81)
UTAH	2811	25.93	1.09	(23.79, 28.07)
VERMONT	2394	27.12	1.06	(25.04, 29.20)
VIRGINIA	1827	28.33	1.22	(25.94, 30.72)
WASHINGTON	3373	27.34	0.87	(25.63, 29.05)
WEST VIRGINIA	2353	33.46	1.05	(31.40, 35.52)
WISCONSIN	2173	30.54	1.31	(27.97, 33.11)
WYOMING	2467	27.89	1.00	(25.93, 29.85)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 29.28 RANGE = 22.32-34.68 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES November 24, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	774	36.21	1.93	(32.43, 39.99)	*	1292	31.75	1.46	(28.89, 34.61)
ALASKA	703	30.42	2.65	(25.23, 35.61)	*	781	29.59	2.44	(24.81, 34.37)
ARIZONA	798	28.46	2.35	(23.85, 33.07)	*	1048	24.06	1.83	(20.47, 27.65)
ARKANSAS	645	33.99	2.08	(29.91, 38.07)	*	1052	29.71	1.58	(26.61, 32.81)
CALIFORNIA	1701	28.03	1.24	(25.60, 30.46)	*	2165	27.43	1.07	(25.33, 29.53)
COLORADO	783	22.70	1.64	(19.49, 25.91)	*	982	21.94	1.48	(19.04, 24.84)
CONNECTICUT	716	32.57	1.97	(28.71, 36.43)	*	1014	21.90	1.45	(19.06, 24.74)
DELAWARE	819	32.50	1.85	(28.87, 36.13)	*	1224	31.10	1.51	(28.14, 34.06)
DISTRICT OF CO	456	23.93	2.43	(19.17, 28.69)	*	707	33.67	2.17	(29.42, 37.92)
FLORIDA	1428	31.26	1.33	(28.65, 33.87)	*	2046	29.85	1.16	(27.58, 32.12)
GEORGIA	1011	27.24	1.65	(24.01, 30.47)	*	1321	21.12	1.22	(18.73, 23.51)
IDAHO	1111	30.68	1.49	(27.76, 33.60)	*	1531	27.88	1.24	(25.45, 30.31)
ILLINOIS	1140	28.42	1.45	(25.58, 31.26)	*	1669	31.17	1.28	(28.66, 33.68)
INDIANA	930	32.92	1.61	(29.76, 36.08)	*	1207	32.50	1.47	(29.62, 35.38)
IOWA	1473	34.09	1.36	(31.42, 36.76)	*	2051	31.45	1.14	(29.22, 33.68)
KANSAS	836	25.80	1.66	(22.55, 29.05)	*	1047	26.31	1.49	(23.39, 29.23)
KENTUCKY	1304	33.80	1.42	(31.02, 36.58)	*	2166	32.16	1.12	(29.96, 34.36)
LOUISIANA	653	33.02	2.10	(28.90, 37.14)	*	938	33.75	1.71	(30.40, 37.10)
MAINE	700	27.71	1.75	(24.28, 31.14)	*	932	28.56	1.57	(25.48, 31.64)
MARYLAND	1800	30.76	1.37	(28.07, 33.45)	*	2464	30.81	1.17	(28.52, 33.10)
MASSACHUSETTS	723	29.17	1.94	(25.37, 32.97)	*	980	24.35	1.52	(21.37, 27.33)
MICHIGAN	1038	33.43	1.60	(30.29, 36.57)	*	1369	32.65	1.40	(29.91, 35.39)
MINNESOTA	1944	29.57	1.13	(27.36, 31.78)	*	2352	26.30	1.01	(24.32, 28.28)
MISSISSIPPI	580	34.61	2.24	(30.22, 39.00)	*	965	34.45	1.66	(31.20, 37.70)
MISSOURI	616	31.07	2.02	(27.11, 35.03)	*	885	35.01	1.81	(31.46, 38.56)
MONTANA	766	31.20	1.87	(27.53, 34.87)	*	976	26.47	1.51	(23.51, 29.43)
NEBRASKA	825	30.20	1.85	(26.57, 33.83)	*	1177	27.83	1.45	(24.99, 30.67)
NEVADA	758	30.61	2.35	(26.00, 35.22)	*	985	24.51	1.91	(20.77, 28.25)
NEW HAMPSHIRE	637	29.21	1.99	(25.31, 33.11)	*	799	22.80	1.66	(19.55, 26.05)
NEW JERSEY	1309	28.93	1.39	(26.21, 31.65)	*	1678	26.19	1.25	(23.74, 28.64)
NEW MEXICO	433	25.48	2.67	(20.25, 30.71)	*	629	28.14	2.05	(24.12, 32.16)
NEW YORK	1725	29.06	1.20	(26.71, 31.41)	*	2368	28.86	1.09	(26.72, 31.00)
NORTH CAROLINA	1118	31.92	1.54	(28.90, 34.94)	*	1514	29.41	1.34	(26.78, 32.04)
NORTH DAKOTA	738	34.11	1.87	(30.44, 37.78)	*	1035	31.62	1.65	(28.39, 34.85)
OHIO	614	33.20	2.20	(28.89, 37.51)	*	828	33.65	1.83	(30.06, 37.24)
OKLAHOMA	847	30.66	1.83	(27.07, 34.25)	*	1025	27.52	1.63	(24.33, 30.71)
OREGON	1230	29.95	1.50	(27.01, 32.89)	*	1554	28.60	1.28	(26.09, 31.11)
PENNSYLVANIA	1502	32.48	1.38	(29.78, 35.18)	*	1913	31.07	1.20	(28.72, 33.42)
RHODE ISLAND	786	29.99	1.81	(26.44, 33.54)	*	981	27.05	1.60	(23.91, 30.19)
SOUTH CAROLINA	717	36.22	2.17	(31.97, 40.47)	*	1042	33.25	1.87	(29.58, 36.92)
SOUTH DAKOTA	882	31.68	1.67	(28.41, 34.95)	*	1103	28.51	1.48	(25.61, 31.41)
TENNESSEE	1153	31.57	1.46	(28.71, 34.43)	*	1752	30.01	1.27	(27.52, 32.50)
TEXAS	655	31.24	1.95	(27.42, 35.06)	*	981	31.44	1.60	(28.30, 34.58)
UTAH	1254	27.61	1.63	(24.42, 30.80)	*	1557	24.25	1.40	(21.51, 26.99)
VERMONT	1060	28.68	1.67	(25.41, 31.95)	*	1334	25.59	1.32	(23.00, 28.18)
VIRGINIA	726	29.59	1.91	(25.85, 33.33)	*	1101	27.10	1.50	(24.16, 30.04)
WASHINGTON	1499	27.14	1.24	(24.71, 29.57)	*	1874	27.54	1.21	(25.17, 29.91)
WEST VIRGINIA	891	32.44	1.68	(29.15, 35.73)	*	1462	34.39	1.33	(31.78, 37.00)
WISCONSIN	940	31.38	1.91	(27.64, 35.12)	*	1233	29.74	1.79	(26.23, 33.25)
WYOMING	1043	30.84	1.56	(27.78, 33.90)	*	1424	24.91	1.25	(22.46, 27.36)
SUMMARY STATISTICS:									
			NUMBER OF PARTICIPANTS =	50				NUMBER OF PARTICIPANTS =	50
			MEDIAN	= 30.72				MEDIAN	= 28.73
			RANGE	= 22.70-36.22				RANGE	= 21.12-35.01
* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES					November 24, 1997				

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 OVERWEIGHT: BY BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	32.52	33.89	1.37
ALASKA	32.11	30.03	-2.08
ARIZONA	25.54	26.28	0.74
ARKANSAS	31.17	31.83	0.66
CALIFORNIA	27.75	27.73	-0.02
COLORADO	22.56	22.32	-0.24
CONNECTICUT	25.31	27.16	1.85
DELAWARE	30.61	31.78	1.17
DISTRICT OF CO	N/A	29.15	N/A
FLORIDA	30.19	30.54	0.35
GEORGIA	28.66	24.07	-4.59
IDAHO	27.92	29.28	1.36
ILLINOIS	30.79	29.84	-0.95
INDIANA	35.26	32.70	-2.56
IOWA	32.05	32.73	0.68
KANSAS	29.03	26.06	-2.97
KENTUCKY	29.86	32.96	3.10
LOUISIANA	31.53	33.39	1.86
MAINE	27.74	28.14	0.40
MARYLAND	29.93	30.78	0.85
MASSACHUSETTS	22.79	26.67	3.88
MICHIGAN	32.25	33.03	0.78
MINNESOTA	29.19	27.91	-1.28
MISSISSIPPI	32.64	34.52	1.88
MISSOURI	33.79	33.11	-0.68
MONTANA	25.69	28.84	3.15
NEBRASKA	30.44	29.00	-1.44
NEVADA	27.70	27.67	-0.03
NEW HAMPSHIRE	26.78	25.99	-0.79
NEW JERSEY	25.47	27.53	2.06
NEW MEXICO	24.58	26.82	2.24
NEW YORK	28.99	28.96	-0.03
NORTH CAROLINA	30.36	30.65	0.29
NORTH DAKOTA	31.26	32.87	1.61
OHIO	33.23	33.43	0.20
OKLAHOMA	25.09	29.05	3.96
OREGON	29.14	29.28	0.14
PENNSYLVANIA	31.05	31.76	0.71
RHODE ISLAND	25.59	28.48	2.89
SOUTH CAROLINA	30.49	34.68	4.19
SOUTH DAKOTA	30.33	30.10	-0.23
TENNESSEE	31.93	30.77	-1.16
TEXAS	29.61	31.34	1.73
UTAH	25.66	25.93	0.27
VERMONT	26.24	27.12	0.88
VIRGINIA	29.99	28.33	-1.66
WASHINGTON	26.32	27.34	1.02
WEST VIRGINIA	32.87	33.46	0.59
WISCONSIN	31.07	30.54	-0.53
WYOMING	27.73	27.89	0.16

* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES
 N/A - NOT AVAILABLE

November 24, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY MEDIAN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2065	38.46	1.24	(36.03, 40.89)
ALASKA	1480	34.29	1.87	(30.62, 37.96)
ARIZONA	1837	30.30	1.56	(27.24, 33.36)
ARKANSAS	1689	36.38	1.37	(33.69, 39.07)
CALIFORNIA	3864	32.78	0.86	(31.09, 34.47)
COLORADO	1763	26.18	1.16	(23.91, 28.45)
CONNECTICUT	1727	32.00	1.28	(29.49, 34.51)
DELAWARE	2043	35.65	1.23	(33.24, 38.06)
DISTRICT OF CO	1163	33.70	1.74	(30.29, 37.11)
FLORIDA	3474	35.37	0.92	(33.57, 37.17)
GEORGIA	2330	29.84	1.08	(27.72, 31.96)
IDAHO	2640	33.97	1.03	(31.95, 35.99)
ILLINOIS	2807	34.33	1.00	(32.37, 36.29)
INDIANA	2137	37.82	1.11	(35.64, 40.00)
IOWA	3518	37.33	0.90	(35.57, 39.09)
KANSAS	1877	31.05	1.18	(28.74, 33.36)
KENTUCKY	3462	37.75	0.97	(35.85, 39.65)
LOUISIANA	1586	38.65	1.37	(35.96, 41.34)
MAINE	1633	34.16	1.29	(31.63, 36.69)
MARYLAND	4258	34.92	0.97	(33.02, 36.82)
MASSACHUSETTS	1703	31.95	1.28	(29.44, 34.46)
MICHIGAN	2406	38.38	1.10	(36.22, 40.54)
MINNESOTA	4291	32.95	0.80	(31.38, 34.52)
MISSISSIPPI	1543	38.32	1.47	(35.44, 41.20)
MISSOURI	1498	37.17	1.43	(34.37, 39.97)
MONTANA	1742	32.92	1.27	(30.43, 35.41)
NEBRASKA	1998	34.38	1.25	(31.93, 36.83)
NEVADA	1741	31.33	1.57	(28.25, 34.41)
NEW HAMPSHIRE	1435	30.16	1.41	(27.40, 32.92)
NEW JERSEY	2984	31.43	0.96	(29.55, 33.31)
NEW MEXICO	1060	32.12	1.81	(28.57, 35.67)
NEW YORK	4089	34.13	0.86	(32.44, 35.82)
NORTH CAROLINA	2628	35.58	1.09	(33.44, 37.72)
NORTH DAKOTA	1769	37.65	1.28	(35.14, 40.16)
OHIO	1442	37.91	1.52	(34.93, 40.89)
OKLAHOMA	1870	33.93	1.31	(31.36, 36.50)
OREGON	2781	33.93	1.02	(31.93, 35.93)
PENNSYLVANIA	3410	36.44	0.93	(34.62, 38.26)
RHODE ISLAND	1767	32.47	1.25	(30.02, 34.92)
SOUTH CAROLINA	1753	39.76	1.45	(36.92, 42.60)
SOUTH DAKOTA	1981	34.43	1.20	(32.08, 36.78)
TENNESSEE	2902	34.68	1.01	(32.70, 36.66)
TEXAS	1634	35.61	1.30	(33.06, 38.16)
UTAH	2807	29.85	1.15	(27.60, 32.10)
VERMONT	2394	31.40	1.12	(29.20, 33.60)
VIRGINIA	1825	33.24	1.28	(30.73, 35.75)
WASHINGTON	3370	31.82	0.90	(30.06, 33.58)
WEST VIRGINIA	2351	38.30	1.11	(36.12, 40.48)
WISCONSIN	2170	35.38	1.35	(32.73, 38.03)
WYOMING	2466	32.13	1.03	(30.11, 34.15)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 34.23 RANGE = 26.18-39.76
 * WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN November 24, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY MEDIAN* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-MALES-				-FEMALES-			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	773	40.80	1.99	(36.90, 44.70)	1292	36.32	1.51	(33.36, 39.28)
ALASKA	702	34.70	2.74	(29.33, 40.07)	778	33.84	2.52	(28.90, 38.78)
ARIZONA	795	32.71	2.39	(28.03, 37.39)	1042	27.84	1.89	(24.14, 31.54)
ARKANSAS	642	37.78	2.27	(33.33, 42.23)	1047	34.99	1.60	(31.85, 38.13)
CALIFORNIA	1700	33.14	1.30	(30.59, 35.69)	2164	32.43	1.13	(30.22, 34.64)
COLORADO	782	26.42	1.72	(23.05, 29.79)	981	25.96	1.57	(22.88, 29.04)
CONNECTICUT	715	37.15	2.02	(33.19, 41.11)	1012	27.00	1.59	(23.88, 30.12)
DELAWARE	819	35.71	1.91	(31.97, 39.45)	1224	35.60	1.54	(32.58, 38.62)
DISTRICT OF CO	456	28.84	2.60	(23.74, 33.94)	707	37.91	2.25	(33.50, 42.32)
FLORIDA	1426	36.62	1.40	(33.88, 39.36)	2048	34.20	1.18	(31.89, 36.51)
GEORGIA	1011	33.12	1.73	(29.73, 36.51)	1319	26.76	1.31	(24.19, 29.33)
IDAHO	1111	34.97	1.56	(31.91, 38.03)	1529	32.96	1.29	(30.43, 35.49)
ILLINOIS	1138	33.32	1.53	(30.32, 36.32)	1669	35.28	1.32	(32.69, 37.87)
INDIANA	930	37.76	1.63	(34.57, 40.95)	1207	37.87	1.52	(34.89, 40.85)
IOWA	1468	37.94	1.39	(35.22, 40.66)	2050	36.75	1.17	(34.46, 39.04)
KANSAS	831	30.95	1.76	(27.50, 34.40)	1046	31.15	1.57	(28.07, 34.23)
KENTUCKY	1302	37.98	1.46	(35.12, 40.84)	2160	37.53	1.18	(35.22, 39.84)
LOUISIANA	650	38.65	2.13	(34.48, 42.82)	936	38.65	1.75	(35.22, 42.08)
MAINE	702	33.35	1.85	(29.72, 36.98)	931	34.96	1.74	(31.55, 38.37)
MARYLAND	1798	34.61	1.45	(31.77, 37.45)	2460	35.22	1.23	(32.81, 37.63)
MASSACHUSETTS	723	34.93	2.04	(30.93, 38.93)	980	29.19	1.61	(26.03, 32.35)
MICHIGAN	1037	38.80	1.66	(35.55, 42.05)	1369	37.98	1.45	(35.14, 40.82)
MINNESOTA	1940	34.84	1.17	(32.55, 37.13)	2351	31.12	1.06	(29.04, 33.20)
MISSISSIPPI	580	37.63	2.31	(33.10, 42.16)	963	38.94	1.71	(35.59, 42.29)
MISSOURI	613	35.00	2.13	(30.83, 39.17)	885	39.18	1.85	(35.55, 42.81)
MONTANA	766	34.93	1.91	(31.19, 38.67)	976	30.91	1.55	(27.87, 33.95)
NEBRASKA	823	35.08	1.97	(31.22, 38.94)	1175	33.70	1.56	(30.64, 36.76)
NEVADA	757	34.40	2.40	(29.70, 39.10)	984	28.05	1.98	(24.17, 31.93)
NEW HAMPSHIRE	637	33.28	2.10	(29.16, 37.40)	798	27.04	1.73	(23.65, 30.43)
NEW JERSEY	1307	32.75	1.45	(29.91, 35.59)	1677	30.17	1.29	(27.64, 32.70)
NEW MEXICO	433	29.76	2.74	(24.39, 35.13)	627	34.46	2.25	(30.05, 38.87)
NEW YORK	1724	34.35	1.25	(31.90, 36.80)	2365	33.92	1.11	(31.74, 36.10)
NORTH CAROLINA	1117	36.61	1.60	(33.47, 39.75)	1511	34.56	1.42	(31.78, 37.34)
NORTH DAKOTA	734	38.47	1.95	(34.65, 42.29)	1035	36.83	1.72	(33.46, 40.20)
OHIO	614	37.81	2.26	(33.38, 42.24)	828	38.01	1.85	(34.38, 41.64)
OKLAHOMA	845	34.66	1.89	(30.96, 38.36)	1025	33.25	1.72	(29.88, 36.62)
OREGON	1227	33.97	1.55	(30.93, 37.01)	1554	33.88	1.33	(31.27, 36.49)
PENNSYLVANIA	1501	36.41	1.41	(33.65, 39.17)	1909	36.47	1.24	(34.04, 38.90)
RHODE ISLAND	786	34.58	1.89	(30.88, 38.28)	981	30.46	1.66	(27.21, 33.71)
SOUTH CAROLINA	715	41.62	2.25	(37.21, 46.03)	1038	38.01	1.87	(34.34, 41.68)
SOUTH DAKOTA	881	35.08	1.72	(31.71, 38.45)	1100	33.77	1.57	(30.69, 36.85)
TENNESSEE	1152	35.21	1.49	(32.29, 38.13)	1750	34.19	1.31	(31.62, 36.76)
TEXAS	654	34.55	2.01	(30.61, 38.49)	980	36.65	1.66	(33.40, 39.90)
UTAH	1252	31.51	1.70	(28.18, 34.84)	1555	28.19	1.50	(25.25, 31.13)
VERMONT	1060	33.19	1.75	(29.76, 36.62)	1334	29.65	1.40	(26.91, 32.39)
VIRGINIA	727	34.54	2.00	(30.62, 38.46)	1098	31.97	1.59	(28.85, 35.09)
WASHINGTON	1497	31.03	1.30	(28.48, 33.58)	1873	32.63	1.25	(30.18, 35.08)
WEST VIRGINIA	891	36.78	1.77	(33.31, 40.25)	1460	39.68	1.38	(36.98, 42.38)
WISCONSIN	937	35.93	1.98	(32.05, 39.81)	1233	34.84	1.84	(31.23, 38.45)
WYOMING	1043	35.00	1.59	(31.88, 38.12)	1423	29.23	1.30	(26.68, 31.78)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 50 NUMBER OF PARTICIPANTS = 50
 MEDIAN = 34.93 MEDIAN = 34.06
 RANGE = 26.42-41.62 RANGE = 25.96-39.68
 * WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN November 24, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 OVERWEIGHT: BY MEDIAN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	36.72	38.46	1.74
ALASKA	37.96	34.29	-3.67
ARIZONA	30.23	30.30	0.07
ARKANSAS	36.01	36.38	0.37
CALIFORNIA	32.29	32.78	0.49
COLORADO	27.20	26.18	-1.02
CONNECTICUT	30.16	32.00	1.84
DELAWARE	35.06	35.65	0.59
DISTRICT OF CO	N/A	33.70	N/A
FLORIDA	34.02	35.37	1.35
GEORGIA	35.38	29.84	-5.54
IDAHO	32.25	33.97	1.72
ILLINOIS	35.53	34.33	-1.20
INDIANA	39.54	37.82	-1.72
IOWA	37.02	37.33	0.31
KANSAS	33.56	31.05	-2.51
KENTUCKY	34.29	37.75	3.46
LOUISIANA	35.90	38.65	2.75
MAINE	33.06	34.16	1.10
MARYLAND	34.88	34.92	0.04
MASSACHUSETTS	27.28	31.95	4.67
MICHIGAN	37.42	38.38	0.96
MINNESOTA	33.61	32.95	-0.66
MISSISSIPPI	37.86	38.32	0.46
MISSOURI	37.76	37.17	-0.59
MONTANA	31.71	32.92	1.21
NEBRASKA	34.41	34.38	-0.03
NEVADA	31.71	31.33	-0.38
NEW HAMPSHIRE	31.71	30.16	-1.55
NEW JERSEY	31.10	31.43	0.33
NEW MEXICO	29.54	32.12	2.58
NEW YORK	33.99	34.13	0.14
NORTH CAROLINA	34.52	35.58	1.06
NORTH DAKOTA	36.84	37.65	0.81
OHIO	38.27	37.91	-0.36
OKLAHOMA	29.59	33.93	4.34
OREGON	34.11	33.93	-0.18
PENNSYLVANIA	36.43	36.44	0.01
RHODE ISLAND	31.62	32.47	0.85
SOUTH CAROLINA	34.62	39.76	5.14
SOUTH DAKOTA	34.37	34.43	0.06
TENNESSEE	36.52	34.68	-1.84
TEXAS	33.48	35.61	2.13
UTAH	30.64	29.85	-0.79
VERMONT	30.25	31.40	1.15
VIRGINIA	34.30	33.24	-1.06
WASHINGTON	31.30	31.82	0.52
WEST VIRGINIA	37.71	38.30	0.59
WISCONSIN	36.41	35.38	-1.03
WYOMING	32.51	32.13	-0.38

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN
 N/A - NOT AVAILABLE

November 24, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR AIDS: ENCOURAGE TEEN CONDOM USE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1679	91.03	0.76	(89.54, 92.52)
ALASKA	1326	91.44	1.26	(88.97, 93.91)
ARIZONA	1429	91.69	0.97	(89.79, 93.59)
ARKANSAS	1314	78.15	1.43	(75.35, 80.95)
COLORADO	1439	90.15	0.99	(88.21, 92.09)
CONNECTICUT	1410	92.13	0.90	(90.37, 93.89)
DELAWARE	1678	90.08	0.81	(88.49, 91.67)
DISTRICT OF CO	968	93.84	0.95	(91.98, 95.70)
FLORIDA	2581	89.09	0.69	(87.74, 90.44)
GEORGIA	1880	92.57	0.67	(91.26, 93.88)
IDAHO	2179	87.42	0.78	(85.89, 88.95)
ILLINOIS	2361	90.95	0.63	(89.72, 92.18)
INDIANA	1730	87.84	0.87	(86.13, 89.55)
IOWA	2652	92.56	0.56	(91.46, 93.66)
KANSAS	1465	87.04	0.96	(85.16, 88.92)
KENTUCKY	2600	87.58	0.68	(86.25, 88.91)
LOUISIANA	1297	87.25	1.03	(85.23, 89.27)
MAINE	1296	92.38	0.73	(90.95, 93.81)
MARYLAND	3495	87.17	0.79	(85.62, 88.72)
MASSACHUSETTS	1444	93.99	0.70	(92.62, 95.36)
MICHIGAN	2060	88.84	0.77	(87.33, 90.35)
MINNESOTA	3503	80.03	0.76	(78.54, 81.52)
MISSISSIPPI	1225	88.36	1.03	(86.34, 90.38)
MISSOURI	1178	88.53	1.07	(86.43, 90.63)
MONTANA	1375	84.85	1.12	(82.65, 87.05)
NEBRASKA	1522	90.06	0.84	(88.41, 91.71)
NEVADA	1471	90.19	1.05	(88.13, 92.25)
NEW HAMPSHIRE	1184	93.07	0.77	(91.56, 94.58)
NEW JERSEY	2454	91.54	0.66	(90.25, 92.83)
NEW MEXICO	871	88.75	1.26	(86.28, 91.22)
NEW YORK	3479	89.07	0.61	(87.87, 90.27)
NORTH CAROLINA	2061	85.25	0.86	(83.56, 86.94)
NORTH DAKOTA	1346	89.98	0.88	(88.26, 91.70)
OHIO	1119	89.22	1.03	(87.20, 91.24)
OKLAHOMA	1279	86.30	1.07	(84.20, 88.40)
OREGON	2320	91.38	0.63	(90.15, 92.61)
PENNSYLVANIA	2687	89.68	0.66	(88.39, 90.97)
RHODE ISLAND	1426	93.01	0.75	(91.54, 94.48)
SOUTH CAROLINA	1444	88.07	1.10	(85.91, 90.23)
SOUTH DAKOTA	1494	82.93	1.05	(80.87, 84.99)
TENNESSEE	2272	81.84	0.91	(80.06, 83.62)
TEXAS	1394	89.74	0.89	(88.00, 91.48)
UTAH	2308	86.99	0.99	(85.05, 88.93)
VERMONT	1990	93.38	0.65	(92.11, 94.65)
VIRGINIA	1555	89.85	0.94	(88.01, 91.69)
WASHINGTON	2893	90.18	0.62	(88.96, 91.40)
WEST VIRGINIA	1810	87.69	0.86	(86.00, 89.38)
WISCONSIN	1796	91.34	0.80	(89.77, 92.91)
WYOMING	2077	91.60	0.68	(90.27, 92.93)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 49 MEDIAN = 89.74 RANGE = 78.15-93.99

* WOULD ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR AIDS-ENCOURAGE TEEN CONDOM USE* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	629	90.93	1.18	(88.62, 93.24)	* 1050	91.12	0.95	(89.26, 92.98)	
ALASKA	616	89.88	2.04	(85.88, 93.88)	* 710	93.16	1.40	(90.42, 95.90)	
ARIZONA	616	91.00	1.48	(88.10, 93.90)	* 813	92.39	1.33	(89.78, 95.00)	
ARKANSAS	492	75.38	2.44	(70.60, 80.16)	* 822	80.82	1.47	(77.94, 83.70)	
COLORADO	661	91.47	1.17	(89.18, 93.76)	* 778	88.81	1.58	(85.71, 91.91)	
CONNECTICUT	580	91.64	1.21	(89.27, 94.01)	* 830	92.63	1.33	(90.02, 95.24)	
DELAWARE	664	89.40	1.24	(86.97, 91.83)	* 1014	90.72	1.03	(88.70, 92.74)	
DISTRICT OF CO	382	92.84	1.61	(89.68, 96.00)	* 586	94.75	1.06	(92.67, 96.83)	
FLORIDA	1064	88.30	1.04	(86.26, 90.34)	* 1517	89.87	0.87	(88.16, 91.58)	
GEORGIA	804	92.35	1.02	(90.35, 94.35)	* 1076	92.79	0.86	(91.10, 94.48)	
IDAHO	930	86.18	1.19	(83.85, 88.51)	* 1249	88.66	1.00	(86.70, 90.62)	
ILLINOIS	970	90.89	0.95	(89.03, 92.75)	* 1391	91.01	0.83	(89.38, 92.64)	
INDIANA	772	85.47	1.36	(82.80, 88.14)	* 958	90.11	1.07	(88.01, 92.21)	
IOWA	1173	91.74	0.87	(90.03, 93.45)	* 1479	93.38	0.71	(91.99, 94.77)	
KANSAS	670	88.49	1.35	(85.84, 91.14)	* 795	85.54	1.36	(82.87, 88.21)	
KENTUCKY	994	86.70	1.10	(84.54, 88.86)	* 1606	88.42	0.81	(86.83, 90.01)	
LOUISIANA	529	87.39	1.57	(84.31, 90.47)	* 768	87.11	1.29	(84.58, 89.64)	
MAINE	564	92.86	1.08	(90.74, 94.98)	* 732	91.90	1.03	(89.88, 93.92)	
MARYLAND	1489	85.70	1.22	(83.31, 88.09)	* 2006	88.60	0.96	(86.72, 90.48)	
MASSACHUSETTS	619	92.85	1.12	(90.65, 95.05)	* 825	95.10	0.84	(93.45, 96.75)	
MICHIGAN	907	87.57	1.22	(85.18, 89.96)	* 1153	90.07	0.95	(88.21, 91.93)	
MINNESOTA	1615	78.25	1.12	(76.05, 80.45)	* 1888	81.77	0.99	(79.83, 83.71)	
MISSISSIPPI	474	86.99	1.65	(83.76, 90.22)	* 751	89.66	1.17	(87.37, 91.95)	
MISSOURI	501	88.83	1.49	(85.91, 91.75)	* 677	88.25	1.44	(85.43, 91.07)	
MONTANA	614	82.74	1.76	(79.29, 86.19)	* 761	86.99	1.31	(84.42, 89.56)	
NEBRASKA	641	90.15	1.30	(87.60, 92.70)	* 881	89.97	1.06	(87.89, 92.05)	
NEVADA	614	90.84	1.46	(87.98, 93.70)	* 857	89.50	1.51	(86.54, 92.46)	
NEW HAMPSHIRE	530	91.83	1.28	(89.32, 94.34)	* 654	94.34	0.91	(92.56, 96.12)	
NEW JERSEY	1084	91.19	1.02	(89.19, 93.19)	* 1370	91.89	0.84	(90.24, 93.54)	
NEW MEXICO	358	88.74	1.87	(85.07, 92.41)	* 513	88.76	1.65	(85.53, 91.99)	
NEW YORK	1451	87.34	0.98	(85.42, 89.26)	* 2028	90.71	0.73	(89.28, 92.14)	
NORTH CAROLINA	897	83.32	1.34	(80.69, 85.95)	* 1164	87.13	1.08	(85.01, 89.25)	
NORTH DAKOTA	594	89.29	1.34	(86.66, 91.92)	* 752	90.71	1.13	(88.50, 92.92)	
OHIO	474	90.50	1.52	(87.52, 93.48)	* 645	87.97	1.38	(85.27, 90.67)	
OKLAHOMA	561	86.12	1.65	(82.89, 89.35)	* 718	86.47	1.32	(83.88, 89.06)	
OREGON	1015	91.08	0.96	(89.20, 92.96)	* 1305	91.68	0.81	(90.09, 93.27)	
PENNSYLVANIA	1188	89.47	1.00	(87.51, 91.43)	* 1499	89.87	0.87	(88.16, 91.58)	
RHODE ISLAND	655	93.31	1.05	(91.25, 95.37)	* 771	92.71	1.06	(90.63, 94.79)	
SOUTH CAROLINA	607	85.03	1.84	(81.42, 88.64)	* 837	90.95	1.14	(88.72, 93.18)	
SOUTH DAKOTA	662	83.32	1.59	(80.20, 86.44)	* 832	82.54	1.36	(79.87, 85.21)	
TENNESSEE	920	80.36	1.43	(77.56, 83.16)	* 1352	83.30	1.13	(81.09, 85.51)	
TEXAS	558	89.91	1.38	(87.21, 92.61)	* 836	89.56	1.12	(87.36, 91.76)	
UTAH	1036	85.00	1.63	(81.81, 88.19)	* 1272	88.93	1.11	(86.75, 91.11)	
VERMONT	888	93.83	0.93	(92.01, 95.65)	* 1102	92.93	0.91	(91.15, 94.71)	
VIRGINIA	624	88.75	1.55	(85.71, 91.79)	* 931	90.94	1.05	(88.88, 93.00)	
WASHINGTON	1252	90.00	0.92	(88.20, 91.80)	* 1641	90.38	0.82	(88.77, 91.99)	
WEST VIRGINIA	715	85.68	1.43	(82.88, 88.48)	* 1095	89.58	0.96	(87.70, 91.46)	
WISCONSIN	786	92.25	1.07	(90.15, 94.35)	* 1010	90.44	1.20	(88.09, 92.79)	
WYOMING	888	91.51	0.98	(89.59, 93.43)	* 1189	91.70	0.91	(89.92, 93.48)	
SUMMARY STATISTICS:		NUMBER OF PARTICIPANTS = 49				NUMBER OF PARTICIPANTS = 49			
		MEDIAN = 89.40				MEDIAN = 90.07			
		RANGE = 75.38-93.83				RANGE = 80.82-95.10			
* WOULD ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM						November 13, 1997			

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 AIDS:ENCOURAGE TEEN CONDOM USE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	88.65	91.03	2.38
ALASKA	91.04	91.44	0.40
ARIZONA	87.27	91.69	4.42
ARKANSAS	86.71	78.15	-8.56
COLORADO	89.75	90.15	0.40
CONNECTICUT	93.70	92.13	-1.57
DELAWARE	88.70	90.08	1.38
DISTRICT OF CO	N/A	93.84	N/A
FLORIDA	89.87	89.09	-0.78
GEORGIA	93.27	92.57	-0.70
IDAHO	88.39	87.42	-0.97
ILLINOIS	90.42	90.95	0.53
INDIANA	89.16	87.84	-1.32
IOWA	93.96	92.56	-1.40
KANSAS	86.46	87.04	0.58
KENTUCKY	88.98	87.58	-1.40
LOUISIANA	88.04	87.25	-0.79
MAINE	91.14	92.38	1.24
MARYLAND	90.09	87.17	-2.92
MASSACHUSETTS	93.16	93.99	0.83
MICHIGAN	92.93	88.84	-4.09
MINNESOTA	81.30	80.03	-1.27
MISSISSIPPI	87.39	88.36	0.97
MISSOURI	89.07	88.53	-0.54
MONTANA	84.09	84.85	0.76
NEBRASKA	88.48	90.06	1.58
NEVADA	91.18	90.19	-0.99
NEW HAMPSHIRE	93.27	93.07	-0.20
NEW JERSEY	86.12	91.54	5.42
NEW MEXICO	92.54	88.75	-3.79
NEW YORK	88.74	89.07	0.33
NORTH CAROLINA	88.12	85.25	-2.87
NORTH DAKOTA	87.76	89.98	2.22
OHIO	86.88	89.22	2.34
OKLAHOMA	89.67	86.30	-3.37
OREGON	90.32	91.38	1.06
PENNSYLVANIA	91.36	89.68	-1.68
RHODE ISLAND	94.57	93.01	-1.56
SOUTH CAROLINA	91.30	88.07	-3.23
SOUTH DAKOTA	82.16	82.93	0.77
TENNESSEE	83.92	81.84	-2.08
TEXAS	87.97	89.74	1.77
UTAH	87.74	86.99	-0.75
VERMONT	94.58	93.38	-1.20
VIRGINIA	90.95	89.85	-1.10
WASHINGTON	90.38	90.18	-0.20
WEST VIRGINIA	86.33	87.69	1.36
WISCONSIN	90.68	91.34	0.66
WYOMING	90.64	91.60	0.96

* WOULD ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM
 N/A - NOT AVAILABLE

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR AIDS: CONDOM EFFECTIVENESS*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1693	27.94	1.31	(25.37, 30.51)
ALASKA	1354	35.70	1.96	(31.86, 39.54)
ARIZONA	1475	40.60	1.74	(37.19, 44.01)
ARKANSAS	1354	23.08	1.35	(20.43, 25.73)
COLORADO	1455	38.02	1.44	(35.20, 40.84)
CONNECTICUT	1425	37.91	1.47	(35.03, 40.79)
DELAWARE	1699	41.86	1.34	(39.23, 44.49)
DISTRICT OF CO	970	45.74	1.96	(41.90, 49.58)
FLORIDA	2604	34.72	1.05	(32.66, 36.78)
GEORGIA	1893	31.45	1.20	(29.10, 33.80)
IDAHO	2216	28.05	1.04	(26.01, 30.09)
ILLINOIS	2373	37.82	1.11	(35.64, 40.00)
INDIANA	1752	30.68	1.19	(28.35, 33.01)
IOWA	2686	33.88	1.03	(31.86, 35.90)
KANSAS	1475	28.34	1.32	(25.75, 30.93)
KENTUCKY	2646	29.77	1.03	(27.75, 31.79)
LOUISIANA	1303	31.79	1.44	(28.97, 34.61)
MAINE	1321	44.41	1.47	(41.53, 47.29)
MARYLAND	3544	36.55	1.08	(34.43, 38.67)
MASSACHUSETTS	1466	43.23	1.53	(40.23, 46.23)
MICHIGAN	2081	39.69	1.19	(37.36, 42.02)
MINNESOTA	3580	29.50	0.84	(27.85, 31.15)
MISSISSIPPI	1241	29.89	1.49	(26.97, 32.81)
MISSOURI	1192	31.18	1.43	(28.38, 33.98)
MONTANA	1397	31.13	1.38	(28.43, 33.83)
NEBRASKA	1548	35.95	1.57	(32.87, 39.03)
NEVADA	1488	36.42	1.77	(32.95, 39.89)
NEW HAMPSHIRE	1185	40.29	1.67	(37.02, 43.56)
NEW JERSEY	2482	38.31	1.13	(36.10, 40.52)
NEW MEXICO	882	34.24	2.04	(30.24, 38.24)
NEW YORK	3546	38.71	0.95	(36.85, 40.57)
NORTH CAROLINA	2077	30.54	1.14	(28.31, 32.77)
NORTH DAKOTA	1374	24.48	1.31	(21.91, 27.05)
OHIO	1136	30.19	1.54	(27.17, 33.21)
OKLAHOMA	1284	21.97	1.33	(19.36, 24.58)
OREGON	2348	37.94	1.11	(35.76, 40.12)
PENNSYLVANIA	2696	36.98	1.06	(34.90, 39.06)
RHODE ISLAND	1452	38.20	1.46	(35.34, 41.06)
SOUTH CAROLINA	1460	27.68	1.39	(24.96, 30.40)
SOUTH DAKOTA	1545	35.35	1.27	(32.86, 37.84)
TENNESSEE	2359	24.44	1.01	(22.46, 26.42)
TEXAS	1423	31.95	1.37	(29.26, 34.64)
UTAH	2354	29.60	1.24	(27.17, 32.03)
VERMONT	2011	40.28	1.34	(37.65, 42.91)
VIRGINIA	1581	36.63	1.40	(33.89, 39.37)
WASHINGTON	2955	36.72	1.00	(34.76, 38.68)
WEST VIRGINIA	1824	28.91	1.20	(26.56, 31.26)
WISCONSIN	1767	33.24	1.54	(30.22, 36.26)
WYOMING	2098	28.39	1.05	(26.33, 30.45)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 49 MEDIAN = 34.24 RANGE = 21.97-45.74

* BELIEVES CONDOMS VERY EFFECTIVE IN PREVENTING AIDS

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR AIDS: CONDOM EFFECTIVENESS* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	638	29.79	2.11	(25.65, 33.93)	1055	26.19	1.49	(23.27, 29.11)
ALASKA	636	39.82	2.89	(34.16, 45.48)	718	31.05	2.61	(25.93, 36.17)
ARIZONA	629	44.26	2.77	(38.83, 49.69)	846	36.94	2.35	(32.33, 41.55)
ARKANSAS	511	22.60	2.15	(18.39, 26.81)	843	23.55	1.67	(20.28, 26.82)
COLORADO	670	39.99	2.13	(35.82, 44.16)	785	36.01	1.94	(32.21, 39.81)
CONNECTICUT	585	41.49	2.20	(37.18, 45.80)	840	34.37	1.94	(30.57, 38.17)
DELAWARE	677	43.89	2.07	(39.83, 47.95)	1022	39.92	1.68	(36.63, 43.21)
DISTRICT OF CO	384	51.81	3.04	(45.85, 57.77)	586	40.14	2.45	(35.34, 44.94)
FLORIDA	1070	39.11	1.60	(35.97, 42.25)	1534	30.44	1.33	(27.83, 33.05)
GEORGIA	811	35.11	1.89	(31.41, 38.81)	1082	27.88	1.52	(24.90, 30.86)
IDAHO	951	28.10	1.61	(24.94, 31.26)	1265	28.00	1.38	(25.30, 30.70)
ILLINOIS	983	39.64	1.66	(36.39, 42.89)	1390	36.02	1.40	(33.28, 38.76)
INDIANA	784	34.97	1.86	(31.32, 38.62)	968	26.54	1.54	(23.52, 29.56)
IOWA	1185	36.65	1.54	(33.63, 39.67)	1501	31.12	1.31	(28.55, 33.69)
KANSAS	674	30.08	1.99	(26.18, 33.98)	801	26.57	1.71	(23.22, 29.92)
KENTUCKY	1020	33.27	1.60	(30.13, 36.41)	1626	26.43	1.22	(24.04, 28.82)
LOUISIANA	534	34.92	2.38	(30.26, 39.58)	769	28.81	1.80	(25.28, 32.34)
MAINE	576	50.42	2.19	(46.13, 54.71)	745	38.45	1.94	(34.65, 42.25)
MARYLAND	1515	37.26	1.62	(34.08, 40.44)	2029	35.84	1.37	(33.15, 38.53)
MASSACHUSETTS	628	45.04	2.41	(40.32, 49.76)	838	41.46	1.89	(37.76, 45.16)
MICHIGAN	917	42.18	1.78	(38.69, 45.67)	1164	37.25	1.58	(34.15, 40.35)
MINNESOTA	1657	31.38	1.24	(28.95, 33.81)	1923	27.64	1.11	(25.46, 29.82)
MISSISSIPPI	480	32.29	2.36	(27.66, 36.92)	761	27.63	1.82	(24.06, 31.20)
MISSOURI	512	34.58	2.27	(30.13, 39.03)	680	27.96	1.72	(24.59, 31.33)
MONTANA	627	33.09	2.04	(29.09, 37.09)	770	29.14	1.73	(25.75, 32.53)
NEBRASKA	651	37.22	2.32	(32.67, 41.77)	897	34.70	2.14	(30.51, 38.89)
NEVADA	624	38.44	2.69	(33.17, 43.71)	864	34.28	2.26	(29.85, 38.71)
NEW HAMPSHIRE	531	42.10	2.45	(37.30, 46.90)	654	38.46	2.18	(34.19, 42.73)
NEW JERSEY	1097	42.79	1.74	(39.38, 46.20)	1385	33.82	1.50	(30.88, 36.76)
NEW MEXICO	363	39.53	3.15	(33.36, 45.70)	519	29.02	2.44	(24.24, 33.80)
NEW YORK	1486	42.50	1.45	(39.66, 45.34)	2060	35.11	1.20	(32.76, 37.46)
NORTH CAROLINA	910	31.70	1.79	(28.19, 35.21)	1167	29.40	1.52	(26.42, 32.38)
NORTH DAKOTA	608	24.25	1.92	(20.49, 28.01)	766	24.73	1.66	(21.48, 27.98)
OHIO	481	31.26	2.37	(26.61, 35.91)	655	29.15	1.95	(25.33, 32.97)
OKLAHOMA	561	25.11	2.05	(21.09, 29.13)	723	18.97	1.60	(15.83, 22.11)
OREGON	1026	41.88	1.69	(38.57, 45.19)	1322	33.99	1.42	(31.21, 36.77)
PENNSYLVANIA	1196	38.28	1.52	(35.30, 41.26)	1500	35.72	1.43	(32.92, 38.52)
RHODE ISLAND	665	41.96	2.13	(37.79, 46.13)	787	34.57	2.00	(30.65, 38.49)
SOUTH CAROLINA	620	29.08	2.21	(24.75, 33.41)	840	26.33	1.69	(23.02, 29.64)
SOUTH DAKOTA	700	36.29	1.89	(32.59, 39.99)	845	34.41	1.69	(31.10, 37.72)
TENNESSEE	951	26.92	1.56	(23.86, 29.98)	1408	22.01	1.26	(19.54, 24.48)
TEXAS	572	35.74	2.19	(31.45, 40.03)	851	28.18	1.64	(24.97, 31.39)
UTAH	1056	31.81	1.75	(28.38, 35.24)	1298	27.44	1.62	(24.26, 30.62)
VERMONT	901	43.23	2.13	(39.06, 47.40)	1110	37.34	1.64	(34.13, 40.55)
VIRGINIA	637	39.64	2.16	(35.41, 43.87)	944	33.64	1.77	(30.17, 37.11)
WASHINGTON	1285	38.45	1.48	(35.55, 41.35)	1670	34.95	1.34	(32.32, 37.58)
WEST VIRGINIA	724	32.98	1.94	(29.18, 36.78)	1100	25.05	1.39	(22.33, 27.77)
WISCONSIN	767	36.36	2.24	(31.97, 40.75)	1000	30.16	2.09	(26.06, 34.26)
WYOMING	899	31.27	1.65	(28.04, 34.50)	1199	25.42	1.32	(22.83, 28.01)
SUMMARY STATISTICS:		NUMBER OF PARTICIPANTS = 49			NUMBER OF PARTICIPANTS = 49			
		MEDIAN = 36.65			MEDIAN = 30.44			
		RANGE = 22.60-51.81			RANGE = 18.97-41.46			
* BELIEVES CONDOMS VERY EFFECTIVE IN PREVENTING AIDS				November 13, 1997				

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 AIDS: CONDOM EFFECTIVENESS*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	29.11	27.94	-1.17
ALASKA	32.90	35.70	2.80
ARIZONA	32.25	40.60	8.35
ARKANSAS	27.96	23.08	-4.88
COLORADO	36.19	38.02	1.83
CONNECTICUT	43.87	37.91	-5.96
DELAWARE	39.50	41.86	2.36
DISTRICT OF CO	N/A	45.74	N/A
FLORIDA	31.70	34.72	3.02
GEORGIA	34.08	31.45	-2.63
IDAHO	28.37	28.05	-0.32
ILLINOIS	38.84	37.82	-1.02
INDIANA	29.58	30.68	1.10
IOWA	29.61	33.88	4.27
KANSAS	26.84	28.34	1.50
KENTUCKY	29.56	29.77	0.21
LOUISIANA	27.27	31.79	4.52
MAINE	46.78	44.41	-2.37
MARYLAND	42.06	36.55	-5.51
MASSACHUSETTS	43.05	43.23	0.18
MICHIGAN	37.64	39.69	2.05
MINNESOTA	31.48	29.50	-1.98
MISSISSIPPI	27.67	29.89	2.22
MISSOURI	33.16	31.18	-1.98
MONTANA	29.43	31.13	1.70
NEBRASKA	31.86	35.95	4.09
NEVADA	37.73	36.42	-1.31
NEW HAMPSHIRE	43.32	40.29	-3.03
NEW JERSEY	37.68	38.31	0.63
NEW MEXICO	34.58	34.24	-0.34
NEW YORK	41.63	38.71	-2.92
NORTH CAROLINA	28.39	30.54	2.15
NORTH DAKOTA	26.76	24.48	-2.28
OHIO	34.96	30.19	-4.77
OKLAHOMA	20.50	21.97	1.47
OREGON	36.05	37.94	1.89
PENNSYLVANIA	35.36	36.98	1.62
RHODE ISLAND	40.29	38.20	-2.09
SOUTH CAROLINA	29.41	27.68	-1.73
SOUTH DAKOTA	29.79	35.35	5.56
TENNESSEE	26.85	24.44	-2.41
TEXAS	29.53	31.95	2.42
UTAH	26.90	29.60	2.70
VERMONT	40.92	40.28	-0.64
VIRGINIA	38.97	36.63	-2.34
WASHINGTON	36.87	36.72	-0.15
WEST VIRGINIA	26.54	28.91	2.37
WISCONSIN	27.75	33.24	5.49
WYOMING	30.47	28.39	-2.08

* BELIEVES CONDOMS VERY EFFECTIVE IN PREVENTING AIDS
 N/A - NOT AVAILABLE

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR AIDS BLOOD TEST*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1641	46.50	1.46	(43.64, 49.36)
ALASKA	1333	48.56	2.05	(44.54, 52.58)
ARIZONA	1430	44.38	1.85	(40.75, 48.01)
ARKANSAS	1314	44.63	1.60	(41.49, 47.77)
COLORADO	1426	45.89	1.51	(42.93, 48.85)
CONNECTICUT	1388	41.64	1.52	(38.66, 44.62)
DELAWARE	1676	46.05	1.39	(43.33, 48.77)
DISTRICT OF CO	958	61.29	1.89	(57.59, 64.99)
FLORIDA	2564	53.64	1.11	(51.46, 55.82)
GEORGIA	1876	33.63	1.23	(31.22, 36.04)
IDAHO	2163	42.59	1.20	(40.24, 44.94)
ILLINOIS	2350	41.17	1.15	(38.92, 43.42)
INDIANA	1687	39.37	1.34	(36.74, 42.00)
IOWA	2617	32.94	1.04	(30.90, 34.98)
KANSAS	1452	30.12	1.33	(27.51, 32.73)
KENTUCKY	2529	34.59	1.09	(32.45, 36.73)
LOUISIANA	1289	49.04	1.60	(45.90, 52.18)
MAINE	1299	34.41	1.47	(31.53, 37.29)
MARYLAND	3450	50.44	1.12	(48.24, 52.64)
MASSACHUSETTS	1431	38.05	1.48	(35.15, 40.95)
MICHIGAN	2042	43.04	1.21	(40.67, 45.41)
MINNESOTA	3520	31.43	0.87	(29.72, 33.14)
MISSISSIPPI	1208	47.67	1.68	(44.38, 50.96)
MISSOURI	1152	39.68	1.66	(36.43, 42.93)
MONTANA	1342	35.44	1.45	(32.60, 38.28)
NEBRASKA	1506	35.01	1.58	(31.91, 38.11)
NEVADA	1466	54.68	1.86	(51.03, 58.33)
NEW HAMPSHIRE	1169	36.86	1.62	(33.68, 40.04)
NEW JERSEY	2420	45.67	1.21	(43.30, 48.04)
NEW MEXICO	870	50.23	2.17	(45.98, 54.48)
NEW YORK	3489	41.05	0.96	(39.17, 42.93)
NORTH CAROLINA	2056	38.69	1.26	(36.22, 41.16)
NORTH DAKOTA	1359	28.97	1.39	(26.25, 31.69)
OHIO	1115	28.10	1.42	(25.32, 30.88)
OKLAHOMA	1268	38.58	1.59	(35.46, 41.70)
OREGON	2275	42.47	1.15	(40.22, 44.72)
PENNSYLVANIA	2637	41.63	1.11	(39.45, 43.81)
RHODE ISLAND	1414	47.61	1.51	(44.65, 50.57)
SOUTH CAROLINA	1443	43.03	1.63	(39.84, 46.22)
SOUTH DAKOTA	1504	27.31	1.27	(24.82, 29.80)
TENNESSEE	2295	40.03	1.16	(37.76, 42.30)
TEXAS	1385	51.12	1.49	(48.20, 54.04)
UTAH	2278	38.98	1.31	(36.41, 41.55)
VERMONT	1957	39.81	1.32	(37.22, 42.40)
VIRGINIA	1533	53.23	1.49	(50.31, 56.15)
WASHINGTON	2825	44.14	1.06	(42.06, 46.22)
WEST VIRGINIA	1792	30.72	1.18	(28.41, 33.03)
WISCONSIN	1765	41.26	1.56	(38.20, 44.32)
WYOMING	2053	39.37	1.19	(37.04, 41.70)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 49 MEDIAN = 41.26 RANGE = 27.31-61.29

* EVER HAD BLOOD TESTED FOR AIDS VIRUS INFECTION

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR AIDS BLOOD TEST* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	621	50.50	2.21	(46.17, 54.83)	1020	42.68	1.75	(39.25, 46.11)	
ALASKA	627	49.40	2.98	(43.56, 55.24)	706	47.63	2.80	(42.14, 53.12)	
ARIZONA	609	45.71	2.74	(40.34, 51.08)	821	43.05	2.37	(38.40, 47.70)	
ARKANSAS	501	45.27	2.74	(39.90, 50.64)	813	43.98	1.95	(40.16, 47.80)	
COLORADO	655	47.43	2.21	(43.10, 51.76)	771	44.31	2.05	(40.29, 48.33)	
CONNECTICUT	571	45.27	2.29	(40.78, 49.76)	817	38.06	2.00	(34.14, 41.98)	
DELAWARE	668	46.33	2.20	(42.02, 50.64)	1008	45.78	1.77	(42.31, 49.25)	
DISTRICT OF CO	378	63.49	2.80	(58.00, 68.98)	580	59.28	2.54	(54.30, 64.26)	
FLORIDA	1058	55.90	1.68	(52.61, 59.19)	1506	51.42	1.45	(48.58, 54.26)	
GEORGIA	806	34.20	1.90	(30.48, 37.92)	1070	33.07	1.60	(29.93, 36.21)	
IDAHO	934	46.51	1.83	(42.92, 50.10)	1229	38.58	1.51	(35.62, 41.54)	
ILLINOIS	966	44.07	1.76	(40.62, 47.52)	1384	38.34	1.45	(35.50, 41.18)	
INDIANA	754	43.91	1.92	(40.15, 47.67)	933	34.97	1.74	(31.56, 38.38)	
IOWA	1158	37.30	1.59	(34.18, 40.42)	1459	28.57	1.31	(26.00, 31.14)	
KANSAS	660	32.08	2.03	(28.10, 36.06)	792	28.14	1.70	(24.81, 31.47)	
KENTUCKY	978	38.14	1.72	(34.77, 41.51)	1551	31.16	1.31	(28.59, 33.73)	
LOUISIANA	527	51.91	2.42	(47.17, 56.65)	762	46.31	2.03	(42.33, 50.29)	
MAINE	566	38.94	2.26	(34.51, 43.37)	733	29.92	1.86	(26.27, 33.57)	
MARYLAND	1470	51.85	1.68	(48.56, 55.14)	1980	49.05	1.41	(46.29, 51.81)	
MASSACHUSETTS	610	41.73	2.35	(37.12, 46.34)	821	34.46	1.85	(30.83, 38.09)	
MICHIGAN	901	45.54	1.81	(41.99, 49.09)	1141	40.60	1.59	(37.48, 43.72)	
MINNESOTA	1624	29.65	1.24	(27.22, 32.08)	1896	33.16	1.17	(30.87, 35.45)	
MISSISSIPPI	467	47.25	2.60	(42.15, 52.35)	741	48.07	2.09	(43.97, 52.17)	
MISSOURI	491	42.48	2.46	(37.66, 47.30)	661	37.05	2.11	(32.91, 41.19)	
MONTANA	597	39.57	2.18	(35.30, 43.84)	745	31.35	1.79	(27.84, 34.86)	
NEBRASKA	630	39.33	2.40	(34.63, 44.03)	876	30.79	2.17	(26.54, 35.04)	
NEVADA	617	54.61	2.82	(49.08, 60.14)	849	54.75	2.40	(50.05, 59.45)	
NEW HAMPSHIRE	523	37.76	2.41	(33.04, 42.48)	646	35.95	2.13	(31.78, 40.12)	
NEW JERSEY	1062	48.40	1.80	(44.87, 51.93)	1358	42.97	1.61	(39.81, 46.13)	
NEW MEXICO	357	52.14	3.21	(45.85, 58.43)	513	48.36	2.76	(42.95, 53.77)	
NEW YORK	1464	45.91	1.47	(43.03, 48.79)	2025	36.39	1.21	(34.02, 38.76)	
NORTH CAROLINA	897	39.05	1.92	(35.29, 42.81)	1159	38.33	1.63	(35.14, 41.52)	
NORTH DAKOTA	601	34.37	2.13	(30.20, 38.54)	758	23.35	1.64	(20.14, 26.56)	
OHIO	472	30.56	2.26	(26.13, 34.99)	643	25.70	1.92	(21.94, 29.46)	
OKLAHOMA	552	37.75	2.31	(33.22, 42.28)	716	39.37	2.09	(35.27, 43.47)	
OREGON	996	46.02	1.74	(42.61, 49.43)	1279	38.89	1.50	(35.95, 41.83)	
PENNSYLVANIA	1172	44.65	1.63	(41.46, 47.84)	1465	38.69	1.46	(35.83, 41.55)	
RHODE ISLAND	650	51.32	2.23	(46.95, 55.69)	764	44.01	2.05	(39.99, 48.03)	
SOUTH CAROLINA	613	43.78	2.59	(38.70, 48.86)	830	42.30	2.07	(38.24, 46.36)	
SOUTH DAKOTA	681	29.48	1.89	(25.78, 33.18)	823	25.13	1.60	(21.99, 28.27)	
TENNESSEE	931	39.96	1.72	(36.59, 43.33)	1364	40.10	1.47	(37.22, 42.98)	
TEXAS	557	51.22	2.32	(46.67, 55.77)	828	51.02	1.88	(47.34, 54.70)	
UTAH	1027	41.85	1.92	(38.09, 45.61)	1251	36.14	1.83	(32.55, 39.73)	
VERMONT	872	44.28	2.07	(40.22, 48.34)	1085	35.38	1.68	(32.09, 38.67)	
VIRGINIA	619	54.83	2.25	(50.42, 59.24)	914	51.62	1.93	(47.84, 55.40)	
WASHINGTON	1223	44.63	1.58	(41.53, 47.73)	1602	43.64	1.41	(40.88, 46.40)	
WEST VIRGINIA	712	32.18	1.87	(28.51, 35.85)	1080	29.34	1.55	(26.30, 32.38)	
WISCONSIN	773	41.99	2.22	(37.64, 46.34)	992	40.54	2.20	(36.23, 44.85)	
WYOMING	883	40.45	1.81	(36.90, 44.00)	1170	38.25	1.55	(35.21, 41.29)	
SUMMARY STATISTICS:		NUMBER OF PARTICIPANTS = 49				NUMBER OF PARTICIPANTS = 49			
		MEDIAN = 44.28				MEDIAN = 38.69			
		RANGE = 29.48-63.49				RANGE = 23.35-59.28			
* EVER HAD BLOOD TESTED FOR AIDS VIRUS INFECTION						November 13, 1997			

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 AIDS BLOOD TEST*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	40.00	46.50	6.50
ALASKA	50.39	48.56	-1.83
ARIZONA	45.00	44.38	-0.62
ARKANSAS	42.33	44.63	2.30
COLORADO	45.70	45.89	0.19
CONNECTICUT	39.69	41.64	1.95
DELAWARE	41.29	46.05	4.76
DISTRICT OF CO	N/A	61.29	N/A
FLORIDA	49.98	53.64	3.66
GEORGIA	36.71	33.63	-3.08
IDAHO	40.60	42.59	1.99
ILLINOIS	38.12	41.17	3.05
INDIANA	33.72	39.37	5.65
IOWA	28.06	32.94	4.88
KANSAS	37.08	30.12	-6.96
KENTUCKY	32.70	34.59	1.89
LOUISIANA	46.95	49.04	2.09
MAINE	27.21	34.41	7.20
MARYLAND	49.48	50.44	0.96
MASSACHUSETTS	36.94	38.05	1.11
MICHIGAN	41.77	43.04	1.27
MINNESOTA	28.94	31.43	2.49
MISSISSIPPI	40.24	47.67	7.43
MISSOURI	37.68	39.68	2.00
MONTANA	33.24	35.44	2.20
NEBRASKA	30.08	35.01	4.93
NEVADA	57.04	54.68	-2.36
NEW HAMPSHIRE	37.19	36.86	-0.33
NEW JERSEY	40.81	45.67	4.86
NEW MEXICO	44.64	50.23	5.59
NEW YORK	36.67	41.05	4.38
NORTH CAROLINA	35.66	38.69	3.03
NORTH DAKOTA	25.62	28.97	3.35
OHIO	28.13	28.10	-0.03
OKLAHOMA	32.89	38.58	5.69
OREGON	37.31	42.47	5.16
PENNSYLVANIA	38.70	41.63	2.93
RHODE ISLAND	44.10	47.61	3.51
SOUTH CAROLINA	31.98	43.03	11.05
SOUTH DAKOTA	26.36	27.31	0.95
TENNESSEE	38.57	40.03	1.46
TEXAS	49.48	51.12	1.64
UTAH	35.84	38.98	3.14
VERMONT	35.05	39.81	4.76
VIRGINIA	52.36	53.23	0.87
WASHINGTON	43.82	44.14	0.32
WEST VIRGINIA	27.59	30.72	3.13
WISCONSIN	35.00	41.26	6.26
WYOMING	38.91	39.37	0.46

* EVER HAD BLOOD TESTED FOR AIDS VIRUS INFECTION
 N/A - NOT AVAILABLE

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING AIDS MEDIUM/HIGH*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1682	6.20	0.65	(4.93, 7.47)
ALASKA	1351	5.16	0.90	(3.40, 6.92)
ARIZONA	1440	6.74	0.93	(4.92, 8.56)
ARKANSAS	1316	7.84	0.84	(6.19, 9.49)
COLORADO	1457	6.14	0.72	(4.73, 7.55)
CONNECTICUT	1418	6.92	0.78	(5.39, 8.45)
DELAWARE	1694	5.54	0.66	(4.25, 6.83)
DISTRICT OF CO	969	11.11	1.13	(8.90, 13.32)
FLORIDA	2566	7.49	0.59	(6.33, 8.65)
GEORGIA	1882	3.86	0.52	(2.84, 4.88)
IDAHO	2207	5.14	0.55	(4.06, 6.22)
ILLINOIS	2357	7.62	0.68	(6.29, 8.95)
INDIANA	1740	5.76	0.61	(4.56, 6.96)
IOWA	2686	7.28	0.59	(6.12, 8.44)
KANSAS	1471	7.60	0.83	(5.97, 9.23)
KENTUCKY	2621	5.54	0.51	(4.54, 6.54)
LOUISIANA	1300	6.65	0.85	(4.98, 8.32)
MAINE	1316	6.49	0.78	(4.96, 8.02)
MARYLAND	3522	4.55	0.44	(3.69, 5.41)
MASSACHUSETTS	1441	5.41	1.06	(3.33, 7.49)
MICHIGAN	2081	6.13	0.60	(4.95, 7.31)
MINNESOTA	3561	7.49	0.53	(6.45, 8.53)
MISSISSIPPI	1226	6.93	0.89	(5.19, 8.67)
MISSOURI	1184	6.73	0.82	(5.12, 8.34)
MONTANA	1383	4.69	0.63	(3.46, 5.92)
NEBRASKA	1537	10.00	1.50	(7.06, 12.94)
NEVADA	1475	8.42	1.02	(6.42, 10.42)
NEW HAMPSHIRE	1185	5.45	0.71	(4.06, 6.84)
NEW JERSEY	2451	7.60	0.67	(6.29, 8.91)
NEW MEXICO	868	6.15	1.06	(4.07, 8.23)
NEW YORK	3463	6.83	0.48	(5.89, 7.77)
NORTH CAROLINA	2067	7.19	0.65	(5.92, 8.46)
NORTH DAKOTA	1370	6.72	0.78	(5.19, 8.25)
OHIO	1133	5.04	0.76	(3.55, 6.53)
OKLAHOMA	1281	6.63	0.91	(4.85, 8.41)
OREGON	2333	5.28	0.56	(4.18, 6.38)
PENNSYLVANIA	2687	7.19	0.60	(6.01, 8.37)
RHODE ISLAND	1429	5.80	0.83	(4.17, 7.43)
SOUTH CAROLINA	1454	5.68	0.65	(4.41, 6.95)
SOUTH DAKOTA	1526	4.44	0.56	(3.34, 5.54)
TENNESSEE	2325	9.84	0.70	(8.47, 11.21)
TEXAS	1402	9.22	0.89	(7.48, 10.96)
UTAH	2355	5.51	0.62	(4.29, 6.73)
VERMONT	1998	6.22	0.73	(4.79, 7.65)
VIRGINIA	1564	6.30	0.72	(4.89, 7.71)
WASHINGTON	2929	5.44	0.51	(4.44, 6.44)
WEST VIRGINIA	1799	14.01	0.92	(12.21, 15.81)
WISCONSIN	1801	6.46	0.81	(4.87, 8.05)
WYOMING	2094	6.38	0.63	(5.15, 7.61)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 49 MEDIAN = 6.46 RANGE = 3.86-14.01
 * CHANCES OF GETTING MEDIUM/HIGH January 7, 1998

MAP

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING AIDS MEDIUM/HIGH* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	638	5.43	0.92	(3.63, 7.23)	*	1044	6.93	0.85	(5.26, 8.60)
ALASKA	639	5.66	1.43	(2.86, 8.46)	*	712	4.59	1.04	(2.55, 6.63)
ARIZONA	613	5.60	1.17	(3.31, 7.89)	*	827	7.89	1.42	(5.11, 10.67)
ARKANSAS	496	8.98	1.39	(6.26, 11.70)	*	820	6.73	0.92	(4.93, 8.53)
COLORADO	669	7.84	1.17	(5.55, 10.13)	*	788	4.42	0.84	(2.77, 6.07)
CONNECTICUT	584	7.61	1.21	(5.24, 9.98)	*	834	6.23	0.96	(4.35, 8.11)
DELAWARE	675	6.06	1.10	(3.90, 8.22)	*	1019	5.05	0.77	(3.54, 6.56)
DISTRICT OF CO	382	13.40	1.93	(9.62, 17.18)	*	587	9.01	1.24	(6.58, 11.44)
FLORIDA	1060	9.55	0.97	(7.65, 11.45)	*	1506	5.47	0.67	(4.16, 6.78)
GEORGIA	807	4.75	0.88	(3.03, 6.47)	*	1075	2.99	0.58	(1.85, 4.13)
IDAHO	944	5.73	0.91	(3.95, 7.51)	*	1263	4.55	0.63	(3.32, 5.78)
ILLINOIS	970	7.91	0.99	(5.97, 9.85)	*	1387	7.34	0.81	(5.75, 8.93)
INDIANA	780	5.69	0.93	(3.87, 7.51)	*	960	5.82	0.79	(4.27, 7.37)
IOWA	1190	7.61	0.90	(5.85, 9.37)	*	1496	6.95	0.74	(5.50, 8.40)
KANSAS	676	9.12	1.36	(6.45, 11.79)	*	795	6.04	0.92	(4.24, 7.84)
KENTUCKY	1007	6.15	0.85	(4.48, 7.82)	*	1614	4.96	0.58	(3.82, 6.10)
LOUISIANA	533	8.98	1.52	(6.00, 11.96)	*	767	4.42	0.84	(2.77, 6.07)
MAINE	574	6.55	1.08	(4.43, 8.67)	*	742	6.42	1.16	(4.15, 8.69)
MARYLAND	1508	4.88	0.72	(3.47, 6.29)	*	2014	4.23	0.54	(3.17, 5.29)
MASSACHUSETTS	619	6.42	1.98	(2.54, 10.30)	*	822	4.42	0.75	(2.95, 5.89)
MICHIGAN	916	6.69	0.92	(4.89, 8.49)	*	1165	5.58	0.79	(4.03, 7.13)
MINNESOTA	1645	8.22	0.78	(6.69, 9.75)	*	1916	6.77	0.70	(5.40, 8.14)
MISSISSIPPI	478	8.57	1.56	(5.51, 11.63)	*	748	5.37	0.84	(3.72, 7.02)
MISSOURI	508	7.36	1.23	(4.95, 9.77)	*	676	6.15	1.02	(4.15, 8.15)
MONTANA	616	5.29	0.95	(3.43, 7.15)	*	767	4.08	0.78	(2.55, 5.61)
NEBRASKA	642	12.15	2.80	(6.66, 17.64)	*	895	7.89	0.99	(5.95, 9.83)
NEVADA	617	9.86	1.68	(6.57, 13.15)	*	858	6.94	1.14	(4.71, 9.17)
NEW HAMPSHIRE	532	5.21	0.99	(3.27, 7.15)	*	653	5.68	1.05	(3.62, 7.74)
NEW JERSEY	1087	8.69	1.05	(6.63, 10.75)	*	1364	6.51	0.81	(4.92, 8.10)
NEW MEXICO	356	8.83	1.92	(5.07, 12.59)	*	512	3.52	0.89	(1.78, 5.26)
NEW YORK	1451	8.33	0.78	(6.80, 9.86)	*	2012	5.40	0.56	(4.30, 6.50)
NORTH CAROLINA	903	7.95	1.02	(5.95, 9.95)	*	1164	6.45	0.85	(4.78, 8.12)
NORTH DAKOTA	606	7.10	1.17	(4.81, 9.39)	*	764	6.33	1.04	(4.29, 8.37)
OHIO	479	5.26	1.24	(2.83, 7.69)	*	654	4.82	0.90	(3.06, 6.58)
OKLAHOMA	558	8.24	1.52	(5.26, 11.22)	*	723	5.11	1.01	(3.13, 7.09)
OREGON	1019	6.20	0.96	(4.32, 8.08)	*	1314	4.36	0.58	(3.22, 5.50)
PENNSYLVANIA	1192	8.00	0.94	(6.16, 9.84)	*	1495	6.40	0.75	(4.93, 7.87)
RHODE ISLAND	662	6.58	1.43	(3.78, 9.38)	*	767	5.04	0.86	(3.35, 6.73)
SOUTH CAROLINA	615	5.88	1.00	(3.92, 7.84)	*	839	5.49	0.85	(3.82, 7.16)
SOUTH DAKOTA	689	3.62	0.78	(2.09, 5.15)	*	837	5.27	0.82	(3.66, 6.88)
TENNESSEE	937	9.90	1.08	(7.78, 12.02)	*	1388	9.77	0.88	(8.05, 11.49)
TEXAS	568	9.26	1.37	(6.57, 11.95)	*	834	9.18	1.15	(6.93, 11.43)
UTAH	1058	7.02	1.02	(5.02, 9.02)	*	1297	4.03	0.69	(2.68, 5.38)
VERMONT	890	7.97	1.16	(5.70, 10.24)	*	1108	4.49	0.87	(2.78, 6.20)
VIRGINIA	626	6.92	1.18	(4.61, 9.23)	*	938	5.69	0.84	(4.04, 7.34)
WASHINGTON	1274	5.68	0.78	(4.15, 7.21)	*	1655	5.19	0.65	(3.92, 6.46)
WEST VIRGINIA	706	11.53	1.36	(8.86, 14.20)	*	1093	16.33	1.29	(13.80, 18.86)
WISCONSIN	789	7.51	1.33	(4.90, 10.12)	*	1012	5.42	0.91	(3.64, 7.20)
WYOMING	895	7.23	1.01	(5.25, 9.21)	*	1199	5.50	0.71	(4.11, 6.89)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	49	NUMBER OF PARTICIPANTS =	49
	MEDIAN	= 7.36	MEDIAN	= 5.50
	RANGE	= 3.62-13.40	RANGE	= 2.99-16.33

* CHANCES OF GETTING MEDIUM/HIGH
 January 7, 1998

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 CHANCE OF GETTING AIDS MEDIUM/HIGH*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	6.49	6.20	-0.29
ALASKA	5.34	5.16	-0.18
ARIZONA	10.24	6.74	-3.50
ARKANSAS	6.14	7.84	1.70
COLORADO	6.52	6.14	-0.38
CONNECTICUT	6.11	6.92	0.81
DELAWARE	6.26	5.54	-0.72
DISTRICT OF CO	N/A	11.11	N/A
FLORIDA	6.58	7.49	0.91
GEORGIA	6.31	3.86	-2.45
IDAHO	6.11	5.14	-0.97
ILLINOIS	8.28	7.62	-0.66
INDIANA	5.12	5.76	0.64
IOWA	7.19	7.28	0.09
KANSAS	6.92	7.60	0.68
KENTUCKY	6.36	5.54	-0.82
LOUISIANA	6.05	6.65	0.60
MAINE	6.10	6.49	0.39
MARYLAND	5.88	4.55	-1.33
MASSACHUSETTS	9.29	5.41	-3.88
MICHIGAN	6.37	6.13	-0.24
MINNESOTA	7.47	7.49	0.02
MISSISSIPPI	6.44	6.93	0.49
MISSOURI	6.83	6.73	-0.10
MONTANA	5.61	4.69	-0.92
NEBRASKA	8.78	10.00	1.22
NEVADA	7.34	8.42	1.08
NEW HAMPSHIRE	5.42	5.45	0.03
NEW JERSEY	7.18	7.60	0.42
NEW MEXICO	7.31	6.15	-1.16
NEW YORK	6.55	6.83	0.28
NORTH CAROLINA	7.52	7.19	-0.33
NORTH DAKOTA	7.41	6.72	-0.69
OHIO	6.12	5.04	-1.08
OKLAHOMA	4.74	6.63	1.89
OREGON	6.01	5.28	-0.73
PENNSYLVANIA	7.86	7.19	-0.67
RHODE ISLAND	5.12	5.80	0.68
SOUTH CAROLINA	7.16	5.68	-1.48
SOUTH DAKOTA	5.78	4.44	-1.34
TENNESSEE	8.41	9.84	1.43
TEXAS	8.69	9.22	0.53
UTAH	5.23	5.51	0.28
VERMONT	6.70	6.22	-0.48
VIRGINIA	7.33	6.30	-1.03
WASHINGTON	4.67	5.44	0.77
WEST VIRGINIA	9.01	14.01	5.00
WISCONSIN	6.73	6.46	-0.27
WYOMING	6.08	6.38	0.30

* CHANCES OF GETTING MEDIUM/HIGH
 N/A - NOT AVAILABLE

January 7, 1998

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO LEISURE TIME PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2137	32.42	1.15	(30.17, 34.67)
ALASKA	1533	25.41	1.70	(22.08, 28.74)
ARIZONA	1955	33.28	1.49	(30.36, 36.20)
ARKANSAS	1797	37.37	1.31	(34.80, 39.94)
CALIFORNIA	4001	23.59	0.79	(22.04, 25.14)
COLORADO	1806	20.16	1.08	(18.04, 22.28)
CONNECTICUT	1837	25.57	1.14	(23.34, 27.80)
DELAWARE	2125	36.12	1.25	(33.67, 38.57)
DISTRICT OF CO	1201	30.25	1.48	(27.35, 33.15)
FLORIDA	3576	27.10	0.84	(25.45, 28.75)
GEORGIA	2382	51.35	1.20	(49.00, 53.70)
IDAHO	2735	20.61	0.90	(18.85, 22.37)
ILLINOIS	1440	24.85	1.25	(22.40, 27.30)
INDIANA	2210	30.97	1.08	(28.85, 33.09)
IOWA	3600	26.91	0.82	(25.30, 28.52)
KANSAS	2007	36.38	1.18	(34.07, 38.69)
KENTUCKY	3618	45.56	0.98	(43.64, 47.48)
LOUISIANA	1653	34.92	1.34	(32.29, 37.55)
MAINE	1697	34.00	1.30	(31.45, 36.55)
MARYLAND	4478	33.82	0.96	(31.94, 35.70)
MASSACHUSETTS	1781	23.02	1.14	(20.79, 25.25)
MICHIGAN	2488	23.30	0.94	(21.46, 25.14)
MINNESOTA	4401	23.63	0.71	(22.24, 25.02)
MISSISSIPPI	1592	39.52	1.47	(36.64, 42.40)
MISSOURI	1548	30.19	1.30	(27.64, 32.74)
MONTANA	1800	21.20	1.04	(19.16, 23.24)
NEBRASKA	2100	22.90	1.05	(20.84, 24.96)
NEVADA	1800	22.74	1.39	(20.02, 25.46)
NEW HAMPSHIRE	1502	25.49	1.33	(22.88, 28.10)
NEW JERSEY	3148	26.27	0.92	(24.47, 28.07)
NEW MEXICO	1096	27.69	1.78	(24.20, 31.18)
NEW YORK	4311	30.40	0.80	(28.83, 31.97)
NORTH CAROLINA	2794	40.68	1.08	(38.56, 42.80)
NORTH DAKOTA	1811	33.85	1.23	(31.44, 36.26)
OHIO	1555	42.61	1.43	(39.81, 45.41)
OKLAHOMA	1925	38.20	1.27	(35.71, 40.69)
OREGON	2928	19.58	0.79	(18.03, 21.13)
PENNSYLVANIA	3591	26.33	0.82	(24.72, 27.94)
RHODE ISLAND	1849	26.67	1.15	(24.42, 28.92)
SOUTH CAROLINA	1863	29.72	1.31	(27.15, 32.29)
SOUTH DAKOTA	2109	34.80	1.15	(32.55, 37.05)
TENNESSEE	3056	40.77	1.01	(38.79, 42.75)
TEXAS	1705	27.86	1.20	(25.51, 30.21)
UTAH	2887	17.12	0.88	(15.40, 18.84)
VERMONT	2503	21.45	0.95	(19.59, 23.31)
VIRGINIA	1890	29.24	1.26	(26.77, 31.71)
WASHINGTON	3587	19.09	0.74	(17.64, 20.54)
WEST VIRGINIA	2420	42.73	1.17	(40.44, 45.02)
WISCONSIN	2230	22.14	1.12	(19.94, 24.34)
WYOMING	2531	20.35	0.86	(18.66, 22.04)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 27.78 RANGE = 17.12-51.35 YEAR 2000 OBJ. 1.5
 * NO LEISURE TIME PHYSICAL ACTIVITY DURING MONTH November 21, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO LEISURE TIME PHYSICAL ACTIVITY* --BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	780	30.77	1.81	(27.22, 34.32)	1357	33.86	1.44	(31.04, 36.68)
ALASKA	713	20.15	2.27	(15.70, 24.60)	820	31.23	2.49	(26.35, 36.11)
ARIZONA	814	30.74	2.17	(26.49, 34.99)	1141	35.68	1.90	(31.96, 39.40)
ARKANSAS	649	36.14	2.19	(31.85, 40.43)	1148	38.47	1.64	(35.26, 41.68)
CALIFORNIA	1728	22.02	1.17	(19.73, 24.31)	2273	25.12	1.05	(23.06, 27.18)
COLORADO	795	18.79	1.64	(15.58, 22.00)	1011	21.47	1.43	(18.67, 24.27)
CONNECTICUT	733	23.77	1.71	(20.42, 27.12)	1104	27.22	1.52	(24.24, 30.20)
DELAWARE	829	33.82	1.88	(30.14, 37.50)	1296	38.21	1.51	(35.25, 41.17)
DISTRICT OF CO	462	25.14	2.20	(20.83, 29.45)	739	34.57	1.96	(30.73, 38.41)
FLORIDA	1442	24.11	1.22	(21.72, 26.50)	2134	29.81	1.12	(27.61, 32.01)
GEORGIA	1020	49.50	1.84	(45.89, 53.11)	1362	53.03	1.56	(49.97, 56.09)
IDAHO	1120	21.78	1.46	(18.92, 24.64)	1615	19.50	1.04	(17.46, 21.54)
ILLINOIS	584	22.99	1.84	(19.38, 26.60)	856	26.65	1.68	(23.36, 29.94)
INDIANA	941	28.01	1.58	(24.91, 31.11)	1269	33.65	1.47	(30.77, 36.53)
IOWA	1478	27.00	1.28	(24.49, 29.51)	2122	26.83	1.03	(24.81, 28.85)
KANSAS	858	36.84	1.79	(33.33, 40.35)	1149	35.96	1.54	(32.94, 38.98)
KENTUCKY	1320	42.25	1.50	(39.31, 45.19)	2298	48.56	1.20	(46.21, 50.91)
LOUISIANA	657	33.15	2.11	(29.01, 37.29)	996	36.49	1.68	(33.20, 39.78)
MAINE	708	34.77	1.93	(30.99, 38.55)	989	33.30	1.68	(30.01, 36.59)
MARYLAND	1838	30.72	1.39	(28.00, 33.44)	2640	36.64	1.24	(34.21, 39.07)
MASSACHUSETTS	738	20.96	1.75	(17.53, 24.39)	1043	24.87	1.50	(21.93, 27.81)
MICHIGAN	1046	20.62	1.38	(17.92, 23.32)	1442	25.74	1.29	(23.21, 28.27)
MINNESOTA	1961	24.23	1.04	(22.19, 26.27)	2440	23.06	0.94	(21.22, 24.90)
MISSISSIPPI	589	40.01	2.31	(35.48, 44.54)	1003	39.09	1.68	(35.80, 42.38)
MISSOURI	619	28.44	2.07	(24.38, 32.50)	929	31.75	1.68	(28.46, 35.04)
MONTANA	771	20.23	1.53	(17.23, 23.23)	1029	22.11	1.37	(19.42, 24.80)
NEBRASKA	838	23.00	1.64	(19.79, 26.21)	1262	22.81	1.32	(20.22, 25.40)
NEVADA	761	20.93	2.05	(16.91, 24.95)	1039	24.57	1.88	(20.89, 28.25)
NEW HAMPSHIRE	647	28.51	2.05	(24.49, 32.53)	855	22.66	1.60	(19.52, 25.80)
NEW JERSEY	1335	22.54	1.29	(20.01, 25.07)	1813	29.65	1.27	(27.16, 32.14)
NEW MEXICO	435	26.15	2.75	(20.76, 31.54)	661	29.13	2.19	(24.84, 33.42)
NEW YORK	1749	26.20	1.19	(23.87, 28.53)	2562	34.14	1.06	(32.06, 36.22)
NORTH CAROLINA	1144	39.31	1.62	(36.13, 42.49)	1650	41.93	1.42	(39.15, 44.71)
NORTH DAKOTA	741	33.61	1.93	(29.83, 37.39)	1070	34.08	1.53	(31.08, 37.08)
OHIO	626	40.79	2.25	(36.38, 45.20)	929	44.24	1.74	(40.83, 47.65)
OKLAHOMA	855	37.82	1.87	(34.15, 41.49)	1070	38.55	1.74	(35.14, 41.96)
OREGON	1235	19.00	1.20	(16.65, 21.35)	1693	20.12	1.05	(18.06, 22.18)
PENNSYLVANIA	1524	22.93	1.18	(20.62, 25.24)	2067	29.33	1.14	(27.10, 31.56)
RHODE ISLAND	795	22.46	1.65	(19.23, 25.69)	1054	30.40	1.60	(27.26, 33.54)
SOUTH CAROLINA	738	24.61	1.97	(20.75, 28.47)	1125	34.31	1.73	(30.92, 37.70)
SOUTH DAKOTA	899	35.84	1.69	(32.53, 39.15)	1210	33.82	1.49	(30.90, 36.74)
TENNESSEE	1175	38.87	1.54	(35.85, 41.89)	1881	42.46	1.29	(39.93, 44.99)
TEXAS	667	25.85	1.83	(22.26, 29.44)	1038	29.76	1.55	(26.72, 32.80)
UTAH	1255	14.84	1.26	(12.37, 17.31)	1632	19.28	1.23	(16.87, 21.69)
VERMONT	1075	21.09	1.44	(18.27, 23.91)	1428	21.79	1.22	(19.40, 24.18)
VIRGINIA	732	26.30	1.91	(22.56, 30.04)	1158	31.98	1.64	(28.77, 35.19)
WASHINGTON	1520	18.40	1.09	(16.26, 20.54)	2067	19.75	1.01	(17.77, 21.73)
WEST VIRGINIA	904	41.28	1.81	(37.73, 44.83)	1516	44.02	1.44	(41.20, 46.84)
WISCONSIN	946	22.07	1.64	(18.86, 25.28)	1284	22.20	1.52	(19.22, 25.18)
WYOMING	1051	21.79	1.33	(19.18, 24.40)	1480	18.93	1.11	(16.75, 21.11)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 50 NUMBER OF PARTICIPANTS = 50
 MEDIAN = 26.00 MEDIAN = 30.82
 RANGE = 14.84-49.50 RANGE = 18.93-53.03
 * NO LEISURE TIME PHYSICAL ACTIVITY DURING MONTH November 21, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1994 TO 1996
 NO LEISURE TIME PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1994 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	45.84	32.42	-13.42
ALASKA	22.84	25.41	2.57
ARIZONA	23.67	33.28	9.61
ARKANSAS	35.12	37.37	2.25
CALIFORNIA	21.75	23.59	1.84
COLORADO	17.16	20.16	3.00
CONNECTICUT	21.96	25.57	3.61
DELAWARE	36.35	36.12	-0.23
DISTRICT OF CO	48.60	30.25	-18.35
FLORIDA	27.93	27.10	-0.83
GEORGIA	32.93	51.35	18.42
IDAHO	21.86	20.61	-1.25
ILLINOIS	33.48	24.85	-8.63
INDIANA	29.63	30.97	1.34
IOWA	33.25	26.91	-6.34
KANSAS	34.45	36.38	1.93
KENTUCKY	45.88	45.56	-0.32
LOUISIANA	33.44	34.92	1.48
MAINE	40.72	34.00	-6.72
MARYLAND	30.25	33.82	3.57
MASSACHUSETTS	24.01	23.02	-0.99
MICHIGAN	23.04	23.30	0.26
MINNESOTA	21.74	23.63	1.89
MISSISSIPPI	38.45	39.52	1.07
MISSOURI	31.90	30.19	-1.71
MONTANA	21.02	21.20	0.18
NEBRASKA	24.25	22.90	-1.35
NEVADA	21.61	22.74	1.13
NEW HAMPSHIRE	25.66	25.49	-0.17
NEW JERSEY	30.90	26.27	-4.63
NEW MEXICO	19.43	27.69	8.26
NEW YORK	37.12	30.40	-6.72
NORTH CAROLINA	42.79	40.68	-2.11
NORTH DAKOTA	31.98	33.85	1.87
OHIO	37.97	42.61	4.64
OKLAHOMA	30.38	38.20	7.82
OREGON	20.77	19.58	-1.19
PENNSYLVANIA	26.47	26.33	-0.14
RHODE ISLAND	N/A	26.67	N/A
SOUTH CAROLINA	31.38	29.72	-1.66
SOUTH DAKOTA	30.69	34.80	4.11
TENNESSEE	39.69	40.77	1.08
TEXAS	27.80	27.86	0.06
UTAH	20.91	17.12	-3.79
VERMONT	23.25	21.45	-1.80
VIRGINIA	22.93	29.24	6.31
WASHINGTON	18.18	19.09	0.91
WEST VIRGINIA	45.30	42.73	-2.57
WISCONSIN	25.89	22.14	-3.75
WYOMING	20.95	20.35	-0.60

* NO LEISURE TIME PHYSICAL ACTIVITY DURING MONTH
 N/A - NOT AVAILABLE

November 21, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR REGULAR AND SUSTAINED PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2137	17.60	0.95	(15.74, 19.46)
ALASKA	1533	23.77	1.71	(20.42, 27.12)
ARIZONA	1955	20.99	1.30	(18.44, 23.54)
ARKANSAS	1797	18.14	1.21	(15.77, 20.51)
CALIFORNIA	4001	21.41	0.72	(20.00, 22.82)
COLORADO	1806	25.08	1.22	(22.69, 27.47)
CONNECTICUT	1837	21.44	1.12	(19.24, 23.64)
DELAWARE	2125	16.13	0.88	(14.41, 17.85)
DISTRICT OF CO	1201	19.97	1.36	(17.30, 22.64)
FLORIDA	3576	21.03	0.75	(19.56, 22.50)
GEORGIA	2382	11.62	0.79	(10.07, 13.17)
IDAHO	2735	25.11	0.91	(23.33, 26.89)
ILLINOIS	1440	21.12	1.21	(18.75, 23.49)
INDIANA	2210	13.48	0.80	(11.91, 15.05)
IOWA	3600	18.88	0.72	(17.47, 20.29)
KANSAS	2007	17.88	0.93	(16.06, 19.70)
KENTUCKY	3618	12.44	0.62	(11.22, 13.66)
LOUISIANA	1653	15.91	1.05	(13.85, 17.97)
MAINE	1697	21.02	1.12	(18.82, 23.22)
MARYLAND	4478	13.28	0.66	(11.99, 14.57)
MASSACHUSETTS	1781	24.18	1.27	(21.69, 26.67)
MICHIGAN	2488	22.80	0.93	(20.98, 24.62)
MINNESOTA	4401	24.06	0.70	(22.69, 25.43)
MISSISSIPPI	1592	16.96	1.15	(14.71, 19.21)
MISSOURI	1548	17.79	1.11	(15.61, 19.97)
MONTANA	1800	24.21	1.13	(22.00, 26.42)
NEBRASKA	2100	20.24	1.07	(18.14, 22.34)
NEVADA	1800	24.67	1.49	(21.75, 27.59)
NEW HAMPSHIRE	1502	22.45	1.27	(19.96, 24.94)
NEW JERSEY	3148	22.36	0.92	(20.56, 24.16)
NEW MEXICO	1096	23.91	1.56	(20.85, 26.97)
NEW YORK	4311	19.97	0.67	(18.66, 21.28)
NORTH CAROLINA	2794	13.57	0.79	(12.02, 15.12)
NORTH DAKOTA	1811	16.78	0.98	(14.86, 18.70)
OHIO	1555	10.23	0.91	(8.45, 12.01)
OKLAHOMA	1925	16.39	0.99	(14.45, 18.33)
OREGON	2928	25.67	0.91	(23.89, 27.45)
PENNSYLVANIA	3591	20.37	0.75	(18.90, 21.84)
RHODE ISLAND	1849	23.38	1.10	(21.22, 25.54)
SOUTH CAROLINA	1863	14.26	0.98	(12.34, 16.18)
SOUTH DAKOTA	2109	18.59	0.98	(16.67, 20.51)
TENNESSEE	3056	14.30	0.70	(12.93, 15.67)
TEXAS	1705	21.51	1.12	(19.31, 23.71)
UTAH	2887	27.17	1.09	(25.03, 29.31)
VERMONT	2503	24.66	1.01	(22.68, 26.64)
VIRGINIA	1890	21.44	1.05	(19.38, 23.50)
WASHINGTON	3587	26.29	0.84	(24.64, 27.94)
WEST VIRGINIA	2420	12.75	0.76	(11.26, 14.24)
WISCONSIN	2230	22.05	1.19	(19.72, 24.38)
WYOMING	2531	28.44	1.00	(26.48, 30.40)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 21.01 RANGE = 10.23-28.44 YEAR 2000 OBJ. 1.3
 * 5+ TIMES/WEEK, 30+ MINUTES/SESSION, REGARDLESS OF INTENSITY November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR REGULAR AND SUSTAINED PHYSICAL ACTIVITY* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	780	18.92	1.54	(15.90, 21.94)	1357	16.44	1.11	(14.26, 18.62)
ALASKA	713	25.45	2.57	(20.41, 30.49)	820	21.90	2.19	(17.61, 26.19)
ARIZONA	814	19.70	1.90	(15.98, 23.42)	1141	22.22	1.61	(19.06, 25.38)
ARKANSAS	649	20.09	2.21	(15.76, 24.42)	1148	16.39	1.17	(14.10, 18.68)
CALIFORNIA	1728	21.59	1.11	(19.41, 23.77)	2273	21.23	0.93	(19.41, 23.05)
COLORADO	795	22.96	1.69	(19.65, 26.27)	1011	27.10	1.74	(23.69, 30.51)
CONNECTICUT	733	22.02	1.70	(18.69, 25.35)	1104	20.92	1.46	(18.06, 23.78)
DELAWARE	829	17.87	1.49	(14.95, 20.79)	1296	14.56	1.00	(12.60, 16.52)
DISTRICT OF CO	462	21.92	2.15	(17.71, 26.13)	739	18.32	1.75	(14.89, 21.75)
FLORIDA	1442	21.94	1.20	(19.59, 24.29)	2134	20.22	0.95	(18.36, 22.08)
GEORGIA	1020	11.83	1.28	(9.32, 14.34)	1362	11.42	1.01	(9.44, 13.40)
IDAHO	1120	22.26	1.34	(19.63, 24.89)	1615	27.84	1.21	(25.47, 30.21)
ILLINOIS	584	20.94	1.84	(17.33, 24.55)	856	21.30	1.54	(18.28, 24.32)
INDIANA	941	14.61	1.26	(12.14, 17.08)	1269	12.45	1.00	(10.49, 14.41)
IOWA	1478	19.03	1.11	(16.85, 21.21)	2122	18.74	0.93	(16.92, 20.56)
KANSAS	858	18.52	1.43	(15.72, 21.32)	1149	17.29	1.20	(14.94, 19.64)
KENTUCKY	1320	14.28	1.06	(12.20, 16.36)	2298	10.78	0.70	(9.41, 12.15)
LOUISIANA	657	15.72	1.60	(12.58, 18.86)	996	16.09	1.32	(13.50, 18.68)
MAINE	708	21.51	1.71	(18.16, 24.86)	989	20.58	1.46	(17.72, 23.44)
MARYLAND	1838	14.94	1.05	(12.88, 17.00)	2640	11.77	0.79	(10.22, 13.32)
MASSACHUSETTS	738	24.79	2.10	(20.67, 28.91)	1043	23.63	1.48	(20.73, 26.53)
MICHIGAN	1046	25.31	1.48	(22.41, 28.21)	1442	20.51	1.14	(18.28, 22.74)
MINNESOTA	1961	24.81	1.07	(22.71, 26.91)	2440	23.36	0.92	(21.56, 25.16)
MISSISSIPPI	589	18.11	1.91	(14.37, 21.85)	1003	15.95	1.30	(13.40, 18.50)
MISSOURI	619	19.65	1.80	(16.12, 23.18)	929	16.13	1.32	(13.54, 18.72)
MONTANA	771	23.46	1.58	(20.36, 26.56)	1029	24.92	1.54	(21.90, 27.94)
NEBRASKA	838	20.96	1.61	(17.80, 24.12)	1262	19.58	1.43	(16.78, 22.38)
NEVADA	761	26.03	2.28	(21.56, 30.50)	1039	23.28	1.91	(19.54, 27.02)
NEW HAMPSHIRE	647	21.98	1.90	(18.26, 25.70)	855	22.89	1.65	(19.66, 26.12)
NEW JERSEY	1335	24.19	1.48	(21.29, 27.09)	1813	20.70	1.11	(18.52, 22.88)
NEW MEXICO	435	23.18	2.31	(18.65, 27.71)	661	24.60	2.08	(20.52, 28.68)
NEW YORK	1749	20.69	1.03	(18.67, 22.71)	2562	19.33	0.86	(17.64, 21.02)
NORTH CAROLINA	1144	15.58	1.29	(13.05, 18.11)	1650	11.73	0.91	(9.95, 13.51)
NORTH DAKOTA	741	15.97	1.48	(13.07, 18.87)	1070	17.56	1.25	(15.11, 20.01)
OHIO	626	10.40	1.46	(7.54, 13.26)	929	10.08	1.11	(7.90, 12.26)
OKLAHOMA	855	16.44	1.48	(13.54, 19.34)	1070	16.34	1.37	(13.65, 19.03)
OREGON	1235	25.49	1.40	(22.75, 28.23)	1693	25.85	1.18	(23.54, 28.16)
PENNSYLVANIA	1524	21.74	1.17	(19.45, 24.03)	2067	19.15	0.98	(17.23, 21.07)
RHODE ISLAND	795	25.20	1.73	(21.81, 28.59)	1054	21.76	1.40	(19.02, 24.50)
SOUTH CAROLINA	738	14.46	1.50	(11.52, 17.40)	1125	14.07	1.28	(11.56, 16.58)
SOUTH DAKOTA	899	18.25	1.42	(15.47, 21.03)	1210	18.90	1.32	(16.31, 21.49)
TENNESSEE	1175	14.86	1.09	(12.72, 17.00)	1881	13.81	0.86	(12.12, 15.50)
TEXAS	667	22.56	1.79	(19.05, 26.07)	1038	20.52	1.37	(17.83, 23.21)
UTAH	1255	27.20	1.64	(23.99, 30.41)	1632	27.14	1.42	(24.36, 29.92)
VERMONT	1075	23.85	1.47	(20.97, 26.73)	1428	25.41	1.35	(22.76, 28.06)
VIRGINIA	732	21.82	1.61	(18.66, 24.98)	1158	21.09	1.36	(18.42, 23.76)
WASHINGTON	1520	26.80	1.30	(24.25, 29.35)	2067	25.81	1.07	(23.71, 27.91)
WEST VIRGINIA	904	12.17	1.20	(9.82, 14.52)	1516	13.26	0.96	(11.38, 15.14)
WISCONSIN	946	22.00	1.83	(18.41, 25.59)	1284	22.09	1.55	(19.05, 25.13)
WYOMING	1051	27.71	1.51	(24.75, 30.67)	1480	29.15	1.28	(26.64, 31.66)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 50 NUMBER OF PARTICIPANTS = 50
 MEDIAN = 21.55 MEDIAN = 20.37
 RANGE = 10.40-27.71 RANGE = 10.08-29.15
 * 5+ TIMES/WEEK, 30+ MINUTES/SESSION, REGARDLESS OF INTENSITY November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1994 TO 1996
 REGULAR AND SUSTAINED PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1994 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	17.08	17.60	0.52
ALASKA	28.27	23.77	-4.50
ARIZONA	17.77	20.99	3.22
ARKANSAS	17.21	18.14	0.93
CALIFORNIA	21.89	21.41	-0.48
COLORADO	26.48	25.08	-1.40
CONNECTICUT	27.00	21.44	-5.56
DELAWARE	17.72	16.13	-1.59
DISTRICT OF CO	11.56	19.97	8.41
FLORIDA	23.85	21.03	-2.82
GEORGIA	18.03	11.62	-6.41
IDAHO	26.32	25.11	-1.21
ILLINOIS	15.68	21.12	5.44
INDIANA	18.78	13.48	-5.30
IOWA	15.92	18.88	2.96
KANSAS	16.84	17.88	1.04
KENTUCKY	13.21	12.44	-0.77
LOUISIANA	16.78	15.91	-0.87
MAINE	12.99	21.02	8.03
MARYLAND	17.65	13.28	-4.37
MASSACHUSETTS	23.20	24.18	0.98
MICHIGAN	21.77	22.80	1.03
MINNESOTA	20.10	24.06	3.96
MISSISSIPPI	14.00	16.96	2.96
MISSOURI	18.01	17.79	-0.22
MONTANA	21.79	24.21	2.42
NEBRASKA	16.73	20.24	3.51
NEVADA	25.36	24.67	-0.69
NEW HAMPSHIRE	21.25	22.45	1.20
NEW JERSEY	20.69	22.36	1.67
NEW MEXICO	25.66	23.91	-1.75
NEW YORK	14.76	19.97	5.21
NORTH CAROLINA	12.72	13.57	0.85
NORTH DAKOTA	20.23	16.78	-3.45
OHIO	15.85	10.23	-5.62
OKLAHOMA	23.00	16.39	-6.61
OREGON	27.33	25.67	-1.66
PENNSYLVANIA	21.24	20.37	-0.87
RHODE ISLAND	N/A	23.38	N/A
SOUTH CAROLINA	15.11	14.26	-0.85
SOUTH DAKOTA	19.44	18.59	-0.85
TENNESSEE	15.00	14.30	-0.70
TEXAS	20.72	21.51	0.79
UTAH	21.62	27.17	5.55
VERMONT	25.70	24.66	-1.04
VIRGINIA	24.60	21.44	-3.16
WASHINGTON	25.70	26.29	0.59
WEST VIRGINIA	14.29	12.75	-1.54
WISCONSIN	22.75	22.05	-0.70
WYOMING	27.95	28.44	0.49

* 5+ TIMES/WEEK, 30+ MINUTES/SESSION, REGARDLESS OF INTENSITY
 N/A - NOT AVAILABLE

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR REGULAR AND VIGOROUS PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2137	10.11	0.73	(8.68, 11.54)
ALASKA	1533	14.06	1.29	(11.53, 16.59)
ARIZONA	1955	15.83	1.07	(13.73, 17.93)
ARKANSAS	1797	11.53	0.98	(9.61, 13.45)
CALIFORNIA	4001	7.90	0.45	(7.02, 8.78)
COLORADO	1806	16.63	0.98	(14.71, 18.55)
CONNECTICUT	1837	14.33	0.89	(12.59, 16.07)
DELAWARE	2125	12.73	0.81	(11.14, 14.32)
DISTRICT OF CO	1201	14.74	1.21	(12.37, 17.11)
FLORIDA	3576	17.62	0.71	(16.23, 19.01)
GEORGIA	2382	6.46	0.55	(5.38, 7.54)
IDAHO	2735	14.19	0.72	(12.78, 15.60)
ILLINOIS	1440	16.76	1.09	(14.62, 18.90)
INDIANA	2210	11.43	0.75	(9.96, 12.90)
IOWA	3600	13.51	0.62	(12.29, 14.73)
KANSAS	2007	15.86	0.88	(14.14, 17.58)
KENTUCKY	3618	10.67	0.60	(9.49, 11.85)
LOUISIANA	1653	11.29	0.86	(9.60, 12.98)
MAINE	1697	14.06	0.93	(12.24, 15.88)
MARYLAND	4478	12.37	0.61	(11.17, 13.57)
MASSACHUSETTS	1781	17.93	1.01	(15.95, 19.91)
MICHIGAN	2488	16.73	0.81	(15.14, 18.32)
MINNESOTA	4401	17.13	0.62	(15.91, 18.35)
MISSISSIPPI	1592	11.22	0.83	(9.59, 12.85)
MISSOURI	1548	11.86	0.89	(10.12, 13.60)
MONTANA	1800	16.50	0.95	(14.64, 18.36)
NEBRASKA	2100	12.51	0.78	(10.98, 14.04)
NEVADA	1800	15.32	1.23	(12.91, 17.73)
NEW HAMPSHIRE	1502	15.64	1.07	(13.54, 17.74)
NEW JERSEY	3148	14.13	0.69	(12.78, 15.48)
NEW MEXICO	1096	14.92	1.23	(12.51, 17.33)
NEW YORK	4311	13.93	0.59	(12.77, 15.09)
NORTH CAROLINA	2794	9.06	0.61	(7.86, 10.26)
NORTH DAKOTA	1811	11.10	0.79	(9.55, 12.65)
OHIO	1555	6.38	0.65	(5.11, 7.65)
OKLAHOMA	1925	12.15	0.80	(10.58, 13.72)
OREGON	2928	17.26	0.75	(15.79, 18.73)
PENNSYLVANIA	3591	12.63	0.62	(11.41, 13.85)
RHODE ISLAND	1849	15.27	0.91	(13.49, 17.05)
SOUTH CAROLINA	1863	12.72	0.99	(10.78, 14.66)
SOUTH DAKOTA	2109	13.48	0.79	(11.93, 15.03)
TENNESSEE	3056	11.17	0.66	(9.88, 12.46)
TEXAS	1705	14.45	0.92	(12.65, 16.25)
UTAH	2887	16.07	0.89	(14.33, 17.81)
VERMONT	2503	14.43	0.78	(12.90, 15.96)
VIRGINIA	1890	15.01	0.92	(13.21, 16.81)
WASHINGTON	3587	16.13	0.69	(14.78, 17.48)
WEST VIRGINIA	2420	10.43	0.70	(9.06, 11.80)
WISCONSIN	2230	14.67	0.96	(12.79, 16.55)
WYOMING	2531	15.65	0.77	(14.14, 17.16)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 14.10 RANGE = 6.38-17.93 YEAR 2000 OBJ. 1.4
 * 3+ TIMES/WEEK, 20+ MINUTES/SESSION, 50%+ OF CAPACITY November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR REGULAR AND VIGOROUS PHYSICAL ACTIVITY* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	780	9.26	1.11	(7.08, 11.44)	* 1357	10.86	0.94	(9.02, 12.70)
ALASKA	713	12.47	1.76	(9.02, 15.92)	* 820	15.82	1.88	(12.14, 19.50)
ARIZONA	814	10.96	1.38	(8.26, 13.66)	* 1141	20.45	1.58	(17.35, 23.55)
ARKANSAS	649	10.78	1.67	(7.51, 14.05)	* 1148	12.20	1.04	(10.16, 14.24)
CALIFORNIA	1728	6.35	0.62	(5.13, 7.57)	* 2273	9.42	0.66	(8.13, 10.71)
COLORADO	795	13.35	1.28	(10.84, 15.86)	* 1011	19.77	1.45	(16.93, 22.61)
CONNECTICUT	733	12.48	1.29	(9.95, 15.01)	* 1104	16.01	1.26	(13.54, 18.48)
DELAWARE	829	13.25	1.26	(10.78, 15.72)	* 1296	12.25	0.99	(10.31, 14.19)
DISTRICT OF CO	462	13.13	1.79	(9.62, 16.64)	* 739	16.10	1.53	(13.10, 19.10)
FLORIDA	1442	16.05	1.03	(14.03, 18.07)	* 2134	19.04	0.95	(17.18, 20.90)
GEORGIA	1020	4.78	0.74	(3.33, 6.23)	* 1362	8.01	0.81	(6.42, 9.60)
IDAHO	1120	12.83	1.09	(10.69, 14.97)	* 1615	15.50	0.92	(13.70, 17.30)
ILLINOIS	584	16.04	1.68	(12.75, 19.33)	* 856	17.46	1.39	(14.74, 20.18)
INDIANA	941	9.43	1.04	(7.39, 11.47)	* 1269	13.25	1.04	(11.21, 15.29)
IOWA	1478	11.26	0.89	(9.52, 13.00)	* 2122	15.56	0.88	(13.84, 17.28)
KANSAS	858	13.80	1.28	(11.29, 16.31)	* 1149	17.78	1.22	(15.39, 20.17)
KENTUCKY	1320	10.24	0.94	(8.40, 12.08)	* 2298	11.06	0.71	(9.67, 12.45)
LOUISIANA	657	11.18	1.34	(8.55, 13.81)	* 996	11.40	1.09	(9.26, 13.54)
MAINE	708	13.09	1.35	(10.44, 15.74)	* 989	14.95	1.21	(12.58, 17.32)
MARYLAND	1838	11.53	0.90	(9.77, 13.29)	* 2640	13.13	0.83	(11.50, 14.76)
MASSACHUSETTS	738	16.26	1.46	(13.40, 19.12)	* 1043	19.42	1.40	(16.68, 22.16)
MICHIGAN	1046	16.59	1.25	(14.14, 19.04)	* 1442	16.85	1.05	(14.79, 18.91)
MINNESOTA	1961	14.79	0.87	(13.08, 16.50)	* 2440	19.33	0.87	(17.62, 21.04)
MISSISSIPPI	589	10.06	1.27	(7.57, 12.55)	* 1003	12.25	1.11	(10.07, 14.43)
MISSOURI	619	12.36	1.40	(9.62, 15.10)	* 929	11.41	1.11	(9.23, 13.59)
MONTANA	771	15.04	1.39	(12.32, 17.76)	* 1029	17.89	1.24	(15.46, 20.32)
NEBRASKA	838	10.86	1.14	(8.63, 13.09)	* 1262	14.03	1.07	(11.93, 16.13)
NEVADA	761	14.31	1.89	(10.61, 18.01)	* 1039	16.34	1.58	(13.24, 19.44)
NEW HAMPSHIRE	647	13.19	1.54	(10.17, 16.21)	* 855	17.93	1.42	(15.15, 20.71)
NEW JERSEY	1335	11.75	0.97	(9.85, 13.65)	* 1813	16.29	0.98	(14.37, 18.21)
NEW MEXICO	435	14.49	1.78	(11.00, 17.98)	* 661	15.33	1.60	(12.19, 18.47)
NEW YORK	1749	13.30	0.90	(11.54, 15.06)	* 2562	14.50	0.76	(13.01, 15.99)
NORTH CAROLINA	1144	7.90	0.84	(6.25, 9.55)	* 1650	10.12	0.81	(8.53, 11.71)
NORTH DAKOTA	741	8.79	1.10	(6.63, 10.95)	* 1070	13.34	1.13	(11.13, 15.55)
OHIO	626	5.95	0.99	(4.01, 7.89)	* 929	6.77	0.86	(5.08, 8.46)
OKLAHOMA	855	11.21	1.21	(8.84, 13.58)	* 1070	13.01	1.08	(10.89, 15.13)
OREGON	1235	14.92	1.06	(12.84, 17.00)	* 1693	19.46	1.05	(17.40, 21.52)
PENNSYLVANIA	1524	11.53	0.89	(9.79, 13.27)	* 2067	13.61	0.87	(11.90, 15.32)
RHODE ISLAND	795	15.42	1.39	(12.70, 18.14)	* 1054	15.13	1.19	(12.80, 17.46)
SOUTH CAROLINA	738	10.27	1.48	(7.37, 13.17)	* 1125	14.92	1.31	(12.35, 17.49)
SOUTH DAKOTA	899	11.86	1.20	(9.51, 14.21)	* 1210	15.01	1.09	(12.87, 17.15)
TENNESSEE	1175	11.37	1.04	(9.33, 13.41)	* 1881	11.00	0.78	(9.47, 12.53)
TEXAS	667	13.77	1.39	(11.05, 16.49)	* 1038	15.09	1.22	(12.70, 17.48)
UTAH	1255	14.35	1.24	(11.92, 16.78)	* 1632	17.70	1.23	(15.29, 20.11)
VERMONT	1075	11.81	1.02	(9.81, 13.81)	* 1428	16.86	1.17	(14.57, 19.15)
VIRGINIA	732	14.73	1.41	(11.97, 17.49)	* 1158	15.28	1.19	(12.95, 17.61)
WASHINGTON	1520	14.59	1.01	(12.61, 16.57)	* 2067	17.60	0.93	(15.78, 19.42)
WEST VIRGINIA	904	10.34	1.12	(8.14, 12.54)	* 1516	10.50	0.87	(8.79, 12.21)
WISCONSIN	946	10.91	1.20	(8.56, 13.26)	* 1284	18.15	1.45	(15.31, 20.99)
WYOMING	1051	13.50	1.10	(11.34, 15.66)	* 1480	17.75	1.10	(15.59, 19.91)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 12.42	MEDIAN	= 15.21
	RANGE	= 4.78-16.59	RANGE	= 6.77-20.45

* 3+ TIMES/WEEK, 20+ MINUTES/SESSION, 50%+ OF CAPACITY

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1994 TO 1996
 REGULAR AND VIGOROUS PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1994 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	11.20	10.11	-1.09
ALASKA	15.13	14.06	-1.07
ARIZONA	17.91	15.83	-2.08
ARKANSAS	10.72	11.53	0.81
CALIFORNIA	15.65	7.90	-7.75
COLORADO	15.85	16.63	0.78
CONNECTICUT	16.98	14.33	-2.65
DELAWARE	14.15	12.73	-1.42
DISTRICT OF CO	8.69	14.74	6.05
FLORIDA	19.99	17.62	-2.37
GEORGIA	13.50	6.46	-7.04
IDAHO	15.73	14.19	-1.54
ILLINOIS	14.62	16.76	2.14
INDIANA	12.98	11.43	-1.55
IOWA	13.29	13.51	0.22
KANSAS	13.93	15.86	1.93
KENTUCKY	11.35	10.67	-0.68
LOUISIANA	11.27	11.29	0.02
MAINE	11.26	14.06	2.80
MARYLAND	14.53	12.37	-2.16
MASSACHUSETTS	17.37	17.93	0.56
MICHIGAN	14.46	16.73	2.27
MINNESOTA	15.39	17.13	1.74
MISSISSIPPI	9.77	11.22	1.45
MISSOURI	10.85	11.86	1.01
MONTANA	14.98	16.50	1.52
NEBRASKA	14.75	12.51	-2.24
NEVADA	14.13	15.32	1.19
NEW HAMPSHIRE	17.01	15.64	-1.37
NEW JERSEY	11.57	14.13	2.56
NEW MEXICO	18.51	14.92	-3.59
NEW YORK	10.55	13.93	3.38
NORTH CAROLINA	9.31	9.06	-0.25
NORTH DAKOTA	13.92	11.10	-2.82
OHIO	12.36	6.38	-5.98
OKLAHOMA	11.11	12.15	1.04
OREGON	18.72	17.26	-1.46
PENNSYLVANIA	14.54	12.63	-1.91
RHODE ISLAND	N/A	15.27	N/A
SOUTH CAROLINA	11.88	12.72	0.84
SOUTH DAKOTA	11.92	13.48	1.56
TENNESSEE	12.72	11.17	-1.55
TEXAS	12.97	14.45	1.48
UTAH	14.34	16.07	1.73
VERMONT	18.43	14.43	-4.00
VIRGINIA	14.63	15.01	0.38
WASHINGTON	16.86	16.13	-0.73
WEST VIRGINIA	9.82	10.43	0.61
WISCONSIN	12.67	14.67	2.00
WYOMING	16.27	15.65	-0.62

* 3+ TIMES/WEEK, 20+ MINUTES/SESSION, 50%+ OF CAPACITY
 N/A - NOT AVAILABLE

November 13, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FRUITS AND VEGETABLES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2138	21.34	1.03	(19.32, 23.36)
ALASKA	1536	26.16	1.77	(22.69, 29.63)
ARIZONA	1957	24.29	1.31	(21.72, 26.86)
ARKANSAS	1800	34.24	1.44	(31.42, 37.06)
CALIFORNIA	4003	31.96	0.84	(30.31, 33.61)
COLORADO	1806	26.18	1.16	(23.91, 28.45)
CONNECTICUT	1838	27.23	1.19	(24.90, 29.56)
DELAWARE	2125	16.08	0.85	(14.41, 17.75)
DISTRICT OF CO	1201	32.36	1.69	(29.05, 35.67)
FLORIDA	3576	21.94	0.77	(20.43, 23.45)
GEORGIA	2382	22.72	1.00	(20.76, 24.68)
IDAHO	2735	23.15	0.89	(21.41, 24.89)
ILLINOIS	1440	21.14	1.20	(18.79, 23.49)
INDIANA	2212	20.84	0.92	(19.04, 22.64)
IOWA	3600	15.08	0.65	(13.81, 16.35)
KANSAS	2008	28.25	1.09	(26.11, 30.39)
KENTUCKY	3618	17.35	0.70	(15.98, 18.72)
LOUISIANA	1656	17.98	1.00	(16.02, 19.94)
MAINE	1699	26.16	1.17	(23.87, 28.45)
MARYLAND	4482	24.67	0.85	(23.00, 26.34)
MASSACHUSETTS	1782	25.82	1.15	(23.57, 28.07)
MICHIGAN	2488	23.43	0.93	(21.61, 25.25)
MINNESOTA	4405	30.23	0.78	(28.70, 31.76)
MISSISSIPPI	1592	18.25	1.07	(16.15, 20.35)
MISSOURI	1550	22.59	1.21	(20.22, 24.96)
MONTANA	1803	23.50	1.04	(21.46, 25.54)
NEBRASKA	2102	22.48	1.27	(19.99, 24.97)
NEVADA	1800	21.92	1.31	(19.35, 24.49)
NEW HAMPSHIRE	1502	26.33	1.27	(23.84, 28.82)
NEW JERSEY	3149	26.84	0.96	(24.96, 28.72)
NEW MEXICO	1096	21.06	1.48	(18.16, 23.96)
NEW YORK	4312	25.43	0.75	(23.96, 26.90)
NORTH CAROLINA	2794	16.52	0.78	(14.99, 18.05)
NORTH DAKOTA	1811	17.84	0.97	(15.94, 19.74)
OHIO	1555	13.91	0.94	(12.07, 15.75)
OKLAHOMA	1925	17.18	0.99	(15.24, 19.12)
OREGON	2932	25.14	0.90	(23.38, 26.90)
PENNSYLVANIA	3595	23.65	0.78	(22.12, 25.18)
RHODE ISLAND	1849	23.74	1.12	(21.54, 25.94)
SOUTH CAROLINA	1879	23.92	1.20	(21.57, 26.27)
SOUTH DAKOTA	2110	23.73	1.00	(21.77, 25.69)
TENNESSEE	3059	25.78	0.87	(24.07, 27.49)
TEXAS	1706	28.14	1.18	(25.83, 30.45)
UTAH	2887	21.46	0.96	(19.58, 23.34)
VERMONT	2505	28.75	1.01	(26.77, 30.73)
VIRGINIA	1892	25.66	1.14	(23.43, 27.89)
WASHINGTON	3591	22.88	0.79	(21.33, 24.43)
WEST VIRGINIA	2420	20.06	0.92	(18.26, 21.86)
WISCONSIN	2231	25.19	1.28	(22.68, 27.70)
WYOMING	2532	22.13	0.86	(20.44, 23.82)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 23.58 RANGE = 13.91-34.24 YEAR 2000 OBJ 2.6
 * CONSUMING FIVE OR MORE FRUITS AND VEGETABLES PER DAY November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FRUITS AND VEGETABLES* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	781	20.67	1.60	(17.53, 23.81)	* 1357	21.94	1.25	(19.49, 24.39)
ALASKA	715	22.58	2.49	(17.70, 27.46)	* 821	30.12	2.50	(25.22, 35.02)
ARIZONA	815	20.35	1.91	(16.61, 24.09)	* 1142	28.02	1.76	(24.57, 31.47)
ARKANSAS	649	29.45	2.32	(24.90, 34.00)	* 1151	38.52	1.63	(35.33, 41.71)
CALIFORNIA	1730	26.32	1.21	(23.95, 28.69)	* 2273	37.49	1.14	(35.26, 39.72)
COLORADO	795	18.75	1.53	(15.75, 21.75)	* 1011	33.28	1.69	(29.97, 36.59)
CONNECTICUT	733	20.59	1.64	(17.38, 23.80)	* 1105	33.27	1.66	(30.02, 36.52)
DELAWARE	829	10.34	1.08	(8.22, 12.46)	* 1296	21.29	1.22	(18.90, 23.68)
DISTRICT OF CO	462	26.43	2.33	(21.86, 31.00)	* 739	37.38	2.23	(33.01, 41.75)
FLORIDA	1442	18.21	1.10	(16.05, 20.37)	* 2134	25.33	1.03	(23.31, 27.35)
GEORGIA	1020	20.97	1.47	(18.09, 23.85)	* 1362	24.31	1.30	(21.76, 26.86)
IDAHO	1120	17.57	1.22	(15.18, 19.96)	* 1615	28.50	1.22	(26.11, 30.89)
ILLINOIS	584	14.91	1.55	(11.87, 17.95)	* 856	27.18	1.67	(23.91, 30.45)
INDIANA	942	16.61	1.36	(13.94, 19.28)	* 1270	24.68	1.30	(22.13, 27.23)
IOWA	1478	10.90	0.86	(9.21, 12.59)	* 2122	18.88	0.94	(17.04, 20.72)
KANSAS	859	26.79	1.62	(23.61, 29.97)	* 1149	29.60	1.47	(26.72, 32.48)
KENTUCKY	1320	12.97	1.02	(10.97, 14.97)	* 2298	21.32	0.98	(19.40, 23.24)
LOUISIANA	660	14.38	1.40	(11.64, 17.12)	* 996	21.20	1.41	(18.44, 23.96)
MAINE	709	22.01	1.72	(18.64, 25.38)	* 990	29.96	1.57	(26.88, 33.04)
MARYLAND	1840	22.26	1.25	(19.81, 24.71)	* 2642	26.87	1.12	(24.67, 29.07)
MASSACHUSETTS	739	20.22	1.63	(17.03, 23.41)	* 1043	30.83	1.59	(27.71, 33.95)
MICHIGAN	1046	17.55	1.30	(15.00, 20.10)	* 1442	28.79	1.30	(26.24, 31.34)
MINNESOTA	1961	26.90	1.14	(24.67, 29.13)	* 2444	33.34	1.07	(31.24, 35.44)
MISSISSIPPI	589	15.47	1.70	(12.14, 18.80)	* 1003	20.70	1.38	(18.00, 23.40)
MISSOURI	620	19.55	1.82	(15.98, 23.12)	* 930	25.31	1.57	(22.23, 28.39)
MONTANA	772	17.80	1.42	(15.02, 20.58)	* 1031	28.93	1.52	(25.95, 31.91)
NEBRASKA	839	17.81	2.26	(13.38, 22.24)	* 1263	26.78	1.39	(24.06, 29.50)
NEVADA	761	18.53	1.83	(14.94, 22.12)	* 1039	25.36	1.85	(21.73, 28.99)
NEW HAMPSHIRE	647	20.42	1.83	(16.83, 24.01)	* 855	31.85	1.80	(28.32, 35.38)
NEW JERSEY	1336	22.39	1.42	(19.61, 25.17)	* 1813	30.86	1.30	(28.31, 33.41)
NEW MEXICO	435	15.21	2.05	(11.19, 19.23)	* 661	26.56	2.09	(22.46, 30.66)
NEW YORK	1749	22.57	1.12	(20.37, 24.77)	* 2563	27.97	1.02	(25.97, 29.97)
NORTH CAROLINA	1144	13.09	1.06	(11.01, 15.17)	* 1650	19.65	1.13	(17.44, 21.86)
NORTH DAKOTA	741	10.15	1.21	(7.78, 12.52)	* 1070	25.29	1.48	(22.39, 28.19)
OHIO	626	10.01	1.27	(7.52, 12.50)	* 929	17.41	1.36	(14.74, 20.08)
OKLAHOMA	855	13.32	1.38	(10.62, 16.02)	* 1070	20.71	1.40	(17.97, 23.45)
OREGON	1236	18.61	1.28	(16.10, 21.12)	* 1696	31.27	1.23	(28.86, 33.68)
PENNSYLVANIA	1527	19.45	1.12	(17.25, 21.65)	* 2068	27.38	1.10	(25.22, 29.54)
RHODE ISLAND	795	20.40	1.65	(17.17, 23.63)	* 1054	26.72	1.53	(23.72, 29.72)
SOUTH CAROLINA	749	23.01	1.83	(19.42, 26.60)	* 1130	24.75	1.49	(21.83, 27.67)
SOUTH DAKOTA	899	19.62	1.42	(16.84, 22.40)	* 1211	27.59	1.38	(24.89, 30.29)
TENNESSEE	1176	23.13	1.30	(20.58, 25.68)	* 1883	28.16	1.13	(25.95, 30.37)
TEXAS	667	24.58	1.78	(21.09, 28.07)	* 1039	31.49	1.56	(28.43, 34.55)
UTAH	1255	17.78	1.37	(15.09, 20.47)	* 1632	24.95	1.36	(22.28, 27.62)
VERMONT	1075	20.93	1.40	(18.19, 23.67)	* 1430	36.01	1.43	(33.21, 38.81)
VIRGINIA	733	20.52	1.63	(17.33, 23.71)	* 1159	30.47	1.57	(27.39, 33.55)
WASHINGTON	1522	17.56	1.13	(15.35, 19.77)	* 2069	27.98	1.10	(25.82, 30.14)
WEST VIRGINIA	904	16.12	1.36	(13.45, 18.79)	* 1516	23.56	1.19	(21.23, 25.89)
WISCONSIN	947	18.92	1.72	(15.55, 22.29)	* 1284	31.01	1.85	(27.38, 34.64)
WYOMING	1051	17.17	1.23	(14.76, 19.58)	* 1481	26.98	1.23	(24.57, 29.39)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	50	NUMBER OF PARTICIPANTS =	50
	MEDIAN	= 19.19	MEDIAN	= 27.49
	RANGE	= 10.01-29.45	RANGE	= 17.41-38.52

* CONSUMING FIVE OR MORE FRUITS AND VEGETABLES PER DAY
 November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1994 TO 1996
 FRUITS AND VEGETABLES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1994 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	25.05	21.34	-3.71
ALASKA	18.79	26.16	7.37
ARIZONA	23.65	24.29	0.64
ARKANSAS	25.33	34.24	8.91
CALIFORNIA	29.01	31.96	2.95
COLORADO	21.63	26.18	4.55
CONNECTICUT	33.52	27.23	-6.29
DELAWARE	16.75	16.08	-0.67
DISTRICT OF CO	31.10	32.36	1.26
FLORIDA	25.40	21.94	-3.46
GEORGIA	26.37	22.72	-3.65
IDAHO	21.72	23.15	1.43
ILLINOIS	22.26	21.14	-1.12
INDIANA	19.71	20.84	1.13
IOWA	18.18	15.08	-3.10
KANSAS	30.95	28.25	-2.70
KENTUCKY	17.35	17.35	0.00
LOUISIANA	19.31	17.98	-1.33
MAINE	21.05	26.16	5.11
MARYLAND	26.15	24.67	-1.48
MASSACHUSETTS	30.32	25.82	-4.50
MICHIGAN	30.57	23.43	-7.14
MINNESOTA	19.50	30.23	10.73
MISSISSIPPI	13.87	18.25	4.38
MISSOURI	20.99	22.59	1.60
MONTANA	24.42	23.50	-0.92
NEBRASKA	19.76	22.48	2.72
NEVADA	20.93	21.92	0.99
NEW HAMPSHIRE	26.86	26.33	-0.53
NEW JERSEY	30.50	26.84	-3.66
NEW MEXICO	23.29	21.06	-2.23
NEW YORK	19.86	25.43	5.57
NORTH CAROLINA	18.94	16.52	-2.42
NORTH DAKOTA	17.72	17.84	0.12
OHIO	21.59	13.91	-7.68
OKLAHOMA	22.62	17.18	-5.44
OREGON	21.20	25.14	3.94
PENNSYLVANIA	24.75	23.65	-1.10
RHODE ISLAND	N/A	23.74	N/A
SOUTH CAROLINA	27.80	23.92	-3.88
SOUTH DAKOTA	20.92	23.73	2.81
TENNESSEE	25.71	25.78	0.07
TEXAS	23.07	28.14	5.07
UTAH	17.96	21.46	3.50
VERMONT	28.35	28.75	0.40
VIRGINIA	27.84	25.66	-2.18
WASHINGTON	21.36	22.88	1.52
WEST VIRGINIA	22.67	20.06	-2.61
WISCONSIN	21.16	25.19	4.03
WYOMING	21.37	22.13	0.76

* CONSUMING FIVE OR MORE FRUITS AND VEGETABLES PER DAY
 N/A - NOT AVAILABLE

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR WEIGHT CONTROL*
 DENOMINATOR IS PEOPLE WHO ARE OVERWEIGHT BASED ON BMI
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	688	36.64	2.00	(32.72, 40.56)
ALASKA	451	46.75	3.69	(39.52, 53.98)
ARIZONA	464	37.06	2.96	(31.26, 42.86)
ARKANSAS	549	30.89	2.27	(26.44, 35.34)
CALIFORNIA	1065	46.94	1.73	(43.55, 50.33)
COLORADO	394	46.55	2.78	(41.10, 52.00)
CONNECTICUT	459	46.22	2.61	(41.10, 51.34)
DELAWARE	627	28.85	2.04	(24.85, 32.85)
DISTRICT OF CO	341	39.20	2.95	(33.42, 44.98)
FLORIDA	1012	41.32	1.68	(38.03, 44.61)
GEORGIA	554	27.10	2.15	(22.89, 31.31)
IDAHO	765	44.23	1.93	(40.45, 48.01)
ILLINOIS	412	41.84	2.59	(36.76, 46.92)
INDIANA	710	38.80	1.96	(34.96, 42.64)
IOWA	1150	38.50	1.58	(35.40, 41.60)
KANSAS	491	37.08	2.46	(32.26, 41.90)
KENTUCKY	1120	25.33	1.45	(22.49, 28.17)
MAINE	468	36.35	2.42	(31.61, 41.09)
MARYLAND	1322	39.14	1.78	(35.65, 42.63)
MASSACHUSETTS	455	45.76	2.65	(40.57, 50.95)
MICHIGAN	785	42.89	1.95	(39.07, 46.71)
MINNESOTA	1198	42.52	1.59	(39.40, 45.64)
MISSISSIPPI	542	29.45	2.34	(24.86, 34.04)
MISSOURI	495	42.22	2.51	(37.30, 47.14)
MONTANA	485	41.98	2.35	(37.37, 46.59)
NEBRASKA	606	46.60	2.24	(42.21, 50.99)
NEVADA	462	44.28	3.22	(37.97, 50.59)
NEW HAMPSHIRE	392	44.96	2.89	(39.30, 50.62)
NEW JERSEY	814	41.33	1.95	(37.51, 45.15)
NEW MEXICO	288	39.18	3.59	(32.14, 46.22)
NEW YORK	1158	40.21	1.63	(37.02, 43.40)
NORTH CAROLINA	802	27.76	1.78	(24.27, 31.25)
NORTH DAKOTA	571	35.34	2.25	(30.93, 39.75)
OHIO	474	42.31	2.63	(37.16, 47.46)
OKLAHOMA	514	36.28	2.52	(31.34, 41.22)
OREGON	781	46.64	1.99	(42.74, 50.54)
PENNSYLVANIA	1050	42.22	1.79	(38.71, 45.73)
RHODE ISLAND	505	44.40	2.47	(39.56, 49.24)
SOUTH CAROLINA	586	38.42	2.57	(33.38, 43.46)
SOUTH DAKOTA	592	38.38	2.20	(34.07, 42.69)
TENNESSEE	878	35.01	1.80	(31.48, 38.54)
TEXAS	517	42.80	2.40	(38.10, 47.50)
UTAH	722	42.76	2.38	(38.10, 47.42)
VERMONT	650	44.85	2.25	(40.44, 49.26)
VIRGINIA	507	42.38	2.50	(37.48, 47.28)
WASHINGTON	919	43.79	1.85	(40.16, 47.42)
WEST VIRGINIA	789	27.20	1.77	(23.73, 30.67)
WISCONSIN	692	50.36	2.60	(45.26, 55.46)
WYOMING	663	45.45	2.13	(41.28, 49.62)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 49 MEDIAN = 41.84 RANGE = 25.33-50.36

* EATING FEWER CALORIES AND/OR LESS FAT AND EXERCISING TO LOSE WEIGHT December 8, 1997

Note: Louisiana is omitted from this table due to missing data on fewer calories

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR WEIGHT CONTROL* -BY SEX
 DENOMINATOR IS PEOPLE WHO ARE OVERWEIGHT BASED ON BMI
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	274	33.81	2.95	(28.03, 39.59)	*	414	39.57	2.73	(34.22, 44.92)
ALASKA	211	51.80	5.36	(41.29, 62.31)	*	240	41.05	4.76	(31.72, 50.38)
ARIZONA	212	36.45	4.45	(27.73, 45.17)	*	252	37.79	3.83	(30.28, 45.30)
ARKANSAS	235	25.62	3.10	(19.54, 31.70)	*	314	36.86	3.21	(30.57, 43.15)
CALIFORNIA	473	45.08	2.58	(40.02, 50.14)	*	592	48.86	2.29	(44.37, 53.35)
COLORADO	175	45.13	4.07	(37.15, 53.11)	*	219	47.98	3.78	(40.57, 55.39)
CONNECTICUT	227	43.61	3.60	(36.55, 50.67)	*	232	50.01	3.71	(42.74, 57.28)
DELAWARE	264	27.51	3.01	(21.61, 33.41)	*	363	30.19	2.65	(25.00, 35.38)
DISTRICT OF CO	105	40.36	4.97	(30.62, 50.10)	*	236	38.47	3.64	(31.34, 45.60)
FLORIDA	432	41.77	2.54	(36.79, 46.75)	*	580	40.88	2.28	(36.41, 45.35)
GEORGIA	266	23.00	3.10	(16.92, 29.08)	*	288	32.07	2.97	(26.25, 37.89)
IDAHO	336	39.77	2.82	(34.24, 45.30)	*	429	49.10	2.64	(43.93, 54.27)
ILLINOIS	159	38.24	4.09	(30.22, 46.26)	*	253	45.04	3.48	(38.22, 51.86)
INDIANA	307	37.86	2.95	(32.08, 43.64)	*	403	39.70	2.61	(34.58, 44.82)
IOWA	500	34.20	2.27	(29.75, 38.65)	*	650	42.86	2.16	(38.63, 47.09)
KANSAS	216	31.85	3.65	(24.70, 39.00)	*	275	42.08	3.30	(35.61, 48.55)
KENTUCKY	436	23.65	2.22	(19.30, 28.00)	*	684	27.01	1.89	(23.31, 30.71)
MAINE	197	27.99	3.44	(21.25, 34.73)	*	271	44.35	3.30	(37.88, 50.82)
MARYLAND	567	38.28	2.58	(33.22, 43.34)	*	755	39.97	2.37	(35.32, 44.62)
MASSACHUSETTS	211	43.69	3.85	(36.14, 51.24)	*	244	48.06	3.59	(41.02, 55.10)
MICHIGAN	341	43.28	2.93	(37.54, 49.02)	*	444	42.52	2.58	(37.46, 47.58)
MINNESOTA	574	38.92	2.26	(34.49, 43.35)	*	624	46.40	2.16	(42.17, 50.63)
MISSISSIPPI	201	21.74	3.37	(15.13, 28.35)	*	341	36.47	2.94	(30.71, 42.23)
MISSOURI	190	42.03	4.10	(33.99, 50.07)	*	305	42.37	3.07	(36.35, 48.39)
MONTANA	229	37.91	3.29	(31.46, 44.36)	*	256	46.77	3.39	(40.13, 53.41)
NEBRASKA	257	43.35	3.36	(36.76, 49.94)	*	349	50.04	2.89	(44.38, 55.70)
NEVADA	221	40.12	4.51	(31.28, 48.96)	*	241	49.81	4.46	(41.07, 58.55)
NEW HAMPSHIRE	193	36.60	3.86	(29.03, 44.17)	*	199	55.60	3.94	(47.88, 63.32)
NEW JERSEY	385	40.98	2.84	(35.41, 46.55)	*	429	41.69	2.69	(36.42, 46.96)
NEW MEXICO	100	31.82	5.43	(21.18, 42.46)	*	188	45.91	4.14	(37.80, 54.02)
NEW YORK	493	41.27	2.49	(36.39, 46.15)	*	665	39.19	2.17	(34.94, 43.44)
NORTH CAROLINA	354	25.95	2.50	(21.05, 30.85)	*	448	29.65	2.49	(24.77, 34.53)
NORTH DAKOTA	248	32.17	3.25	(25.80, 38.54)	*	323	38.74	3.07	(32.72, 44.76)
OHIO	192	41.03	3.96	(33.27, 48.79)	*	282	43.55	3.44	(36.81, 50.29)
OKLAHOMA	238	28.33	3.40	(21.67, 34.99)	*	276	44.66	3.54	(37.72, 51.60)
OREGON	347	45.34	2.92	(39.62, 51.06)	*	434	48.01	2.68	(42.76, 53.26)
PENNSYLVANIA	466	41.36	2.63	(36.21, 46.51)	*	584	43.07	2.35	(38.46, 47.68)
RHODE ISLAND	244	47.24	3.53	(40.32, 54.16)	*	261	41.39	3.46	(34.61, 48.17)
SOUTH CAROLINA	246	41.62	3.86	(34.05, 49.19)	*	340	35.17	3.35	(28.60, 41.74)
SOUTH DAKOTA	271	37.12	3.24	(30.77, 43.47)	*	321	39.80	2.92	(34.08, 45.52)
TENNESSEE	359	31.22	2.59	(26.14, 36.30)	*	519	38.77	2.37	(34.12, 43.42)
TEXAS	209	41.26	3.74	(33.93, 48.59)	*	308	44.32	3.03	(38.38, 50.26)
UTAH	327	39.96	3.24	(33.61, 46.31)	*	395	45.93	3.17	(39.72, 52.14)
VERMONT	299	39.54	3.24	(33.19, 45.89)	*	351	50.57	2.96	(44.77, 56.37)
VIRGINIA	206	38.29	3.74	(30.96, 45.62)	*	301	46.77	3.21	(40.48, 53.06)
WASHINGTON	417	39.57	2.57	(34.53, 44.61)	*	502	48.07	2.64	(42.90, 53.24)
WEST VIRGINIA	292	23.61	2.68	(18.36, 28.86)	*	497	30.29	2.40	(25.59, 34.99)
WISCONSIN	301	44.25	3.67	(37.06, 51.44)	*	391	56.50	3.74	(49.17, 63.83)
WYOMING	307	41.40	3.06	(35.40, 47.40)	*	356	50.52	2.89	(44.86, 56.18)
SUMMARY STATISTICS:		NUMBER OF PARTICIPANTS =	49			NUMBER OF PARTICIPANTS =	49		
		MEDIAN	= 39.54			MEDIAN	= 42.86		
		RANGE	= 21.74-51.80			RANGE	= 27.01-56.50		

* EATING FEWER CALORIES AND/OR LESS FAT AND EXERCISING TO LOSE WEIGHT

December 8, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1994 TO 1996
 WEIGHT CONTROL*
 DENOMINATOR IS PEOPLE WHO ARE OVERWEIGHT BASED ON BMI
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1994 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	36.15	36.64	0.49
ALASKA	44.56	46.75	2.19
ARIZONA	35.44	37.06	1.62
ARKANSAS	36.36	30.89	-5.47
CALIFORNIA	44.16	46.94	2.78
COLORADO	53.36	46.55	-6.81
CONNECTICUT	36.47	46.22	9.75
DELAWARE	31.39	28.85	-2.54
DISTRICT OF CO	27.62	39.20	11.58
FLORIDA	40.03	41.32	1.29
GEORGIA	38.47	27.10	-11.37
IDAHO	40.87	44.23	3.36
ILLINOIS	39.77	41.84	2.07
INDIANA	39.69	38.80	-0.89
IOWA	35.04	38.50	3.46
KANSAS	27.20	37.08	9.88
KENTUCKY	24.07	25.33	1.26
MAINE	30.49	36.35	5.86
MARYLAND	41.43	39.14	-2.29
MASSACHUSETTS	45.95	45.76	-0.19
MICHIGAN	48.68	42.89	-5.79
MINNESOTA	42.31	42.52	0.21
MISSISSIPPI	33.00	29.45	-3.55
MISSOURI	40.65	42.22	1.57
MONTANA	42.94	41.98	-0.96
NEBRASKA	43.42	46.60	3.18
NEVADA	38.49	44.28	5.79
NEW HAMPSHIRE	42.43	44.96	2.53
NEW JERSEY	35.47	41.33	5.86
NEW MEXICO	45.96	39.18	-6.78
NEW YORK	39.28	40.21	0.93
NORTH CAROLINA	26.40	27.76	1.36
NORTH DAKOTA	37.20	35.34	-1.86
OHIO	35.16	42.31	7.15
OKLAHOMA	35.49	36.28	0.79
OREGON	44.60	46.64	2.04
PENNSYLVANIA	44.42	42.22	-2.20
RHODE ISLAND	N/A	44.40	N/A
SOUTH CAROLINA	32.47	38.42	5.95
SOUTH DAKOTA	33.91	38.38	4.47
TENNESSEE	37.77	35.01	-2.76
TEXAS	38.56	42.80	4.24
UTAH	45.07	42.76	-2.31
VERMONT	44.99	44.85	-0.14
VIRGINIA	45.63	42.38	-3.25
WASHINGTON	42.25	43.79	1.54
WEST VIRGINIA	26.79	27.20	0.41
WISCONSIN	44.98	50.36	5.38
WYOMING	43.64	45.45	1.81

* EATING FEWER CALORIES AND/OR LESS FAT AND EXERCISING TO LOSE WEIGHT
 N/A - NOT AVAILABLE
 December 8, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2083	17.38	1.00	(15.42, 19.34)
ALASKA	1507	25.18	1.68	(21.89, 28.47)
GEORGIA	2374	15.36	0.94	(13.52, 17.20)
INDIANA	2194	18.37	0.96	(16.49, 20.25)
KANSAS	1934	15.12	0.94	(13.28, 16.96)
KENTUCKY	1809	15.36	1.07	(13.26, 17.46)
MONTANA	1802	23.11	1.14	(20.88, 25.34)
NEBRASKA	2066	17.51	1.37	(14.82, 20.20)
NEW HAMPSHIRE	1453	12.55	1.03	(10.53, 14.57)
NORTH CAROLINA	2785	15.72	0.81	(14.13, 17.31)
NORTH DAKOTA	1796	21.12	1.18	(18.81, 23.43)
OHIO	1535	9.95	0.99	(8.01, 11.89)
OKLAHOMA	1922	12.51	0.93	(10.69, 14.33)
SOUTH CAROLINA	1493	10.74	1.01	(8.76, 12.72)
SOUTH DAKOTA	2100	15.65	0.94	(13.81, 17.49)
TEXAS	1704	19.32	1.14	(17.09, 21.55)
VIRGINIA	1888	13.66	0.98	(11.74, 15.58)
WEST VIRGINIA	2410	22.17	1.00	(20.21, 24.13)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 15.69 RANGE = 9.95-25.18
 * EVER USED OR TRIED SMOKELESS TOBACCO December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	761	31.62	1.85	(27.99, 35.25)	1322	4.82	0.73	(3.39, 6.25)
ALASKA	699	40.97	2.77	(35.54, 46.40)	808	7.88	1.34	(5.25, 10.51)
GEORGIA	1020	27.20	1.71	(23.85, 30.55)	1354	4.46	0.62	(3.24, 5.68)
INDIANA	937	35.99	1.70	(32.66, 39.32)	1257	2.28	0.50	(1.30, 3.26)
KANSAS	832	28.71	1.74	(25.30, 32.12)	1102	2.39	0.49	(1.43, 3.35)
KENTUCKY	669	30.15	1.98	(26.27, 34.03)	1140	1.59	0.41	(0.79, 2.39)
MONTANA	771	42.95	1.98	(39.07, 46.83)	1031	4.26	0.69	(2.91, 5.61)
NEBRASKA	827	32.61	2.32	(28.06, 37.16)	1239	3.62	1.09	(1.48, 5.76)
NEW HAMPSHIRE	630	24.41	1.91	(20.67, 28.15)	823	1.27	0.46	(0.37, 2.17)
NORTH CAROLINA	1141	26.45	1.49	(23.53, 29.37)	1644	5.92	0.58	(4.78, 7.06)
NORTH DAKOTA	738	38.34	2.05	(34.32, 42.36)	1058	4.30	0.71	(2.91, 5.69)
OHIO	621	20.17	1.92	(16.41, 23.93)	914	0.71	0.28	(0.16, 1.26)
OKLAHOMA	854	24.57	1.75	(21.14, 28.00)	1068	1.44	0.42	(0.62, 2.26)
SOUTH CAROLINA	605	18.67	1.91	(14.93, 22.41)	888	3.35	0.67	(2.04, 4.66)
SOUTH DAKOTA	894	29.68	1.71	(26.33, 33.03)	1206	2.52	0.49	(1.56, 3.48)
TEXAS	666	34.85	2.03	(30.87, 38.83)	1038	4.66	0.79	(3.11, 6.21)
VIRGINIA	730	26.85	1.86	(23.20, 30.50)	1158	1.40	0.36	(0.69, 2.11)
WEST VIRGINIA	896	44.74	1.78	(41.25, 48.23)	1514	2.29	0.40	(1.51, 3.07)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	18	NUMBER OF PARTICIPANTS =	18
	MEDIAN	= 29.92	MEDIAN	= 2.93
	RANGE	= 18.67-44.74	RANGE	= 0.71- 7.88

* EVER USED OR TRIED SMOKELESS TOBACCO
 December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2083	5.43	0.59	(4.27, 6.59)
ALASKA	1506	4.08	0.65	(2.81, 5.35)
GEORGIA	2374	5.52	0.62	(4.30, 6.74)
INDIANA	2193	3.95	0.49	(2.99, 4.91)
KANSAS	1934	4.26	0.53	(3.22, 5.30)
KENTUCKY	1809	5.06	0.66	(3.77, 6.35)
MONTANA	1802	7.96	0.74	(6.51, 9.41)
NEBRASKA	2066	5.94	1.24	(3.51, 8.37)
NEW HAMPSHIRE	1453	1.27	0.39	(0.51, 2.03)
NORTH CAROLINA	2783	5.03	0.47	(4.11, 5.95)
NORTH DAKOTA	1796	4.98	0.65	(3.71, 6.25)
OHIO	1535	2.71	0.55	(1.63, 3.79)
OKLAHOMA	1922	3.85	0.55	(2.77, 4.93)
SOUTH CAROLINA	1493	3.35	0.57	(2.23, 4.47)
SOUTH DAKOTA	2100	5.42	0.60	(4.24, 6.60)
TEXAS	1704	4.08	0.59	(2.92, 5.24)
VIRGINIA	1888	3.80	0.56	(2.70, 4.90)
WEST VIRGINIA	2410	8.48	0.68	(7.15, 9.81)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 4.62 RANGE = 1.27- 8.48
 * CURRENT USERS OF SMOKELESS TOBACCO December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	761	10.51	1.20	(8.16, 12.86)	1322	0.96	0.24	(0.49, 1.43)
ALASKA	699	6.82	1.21	(4.45, 9.19)	807	1.07	0.27	(0.54, 1.60)
GEORGIA	1020	10.05	1.21	(7.68, 12.42)	1354	1.33	0.33	(0.68, 1.98)
INDIANA	936	7.86	0.96	(5.98, 9.74)	1257	0.37	0.28	(-0.18, 0.92)
KANSAS	832	8.72	1.06	(6.64, 10.80)	1102	0.09	0.09	(-0.09, 0.27)
KENTUCKY	669	10.18	1.34	(7.55, 12.81)	1140	0.29	0.17	(-0.04, 0.62)
MONTANA	771	15.92	1.46	(13.06, 18.78)	1031	0.40	0.21	(-0.01, 0.81)
NEBRASKA	827	10.86	2.25	(6.45, 15.27)	1239	1.41	1.01	(-0.57, 3.39)
NEW HAMPSHIRE	630	2.60	0.80	(1.03, 4.17)	823	0.00	0.00	(0.00, 0.00)
NORTH CAROLINA	1140	7.98	0.87	(6.27, 9.69)	1643	2.35	0.36	(1.64, 3.06)
NORTH DAKOTA	738	10.09	1.28	(7.58, 12.60)	1058	0.00	0.00	(0.00, 0.00)
OHIO	621	5.72	1.14	(3.49, 7.95)	914	0.00	0.00	(0.00, 0.00)
OKLAHOMA	854	7.75	1.11	(5.57, 9.93)	1068	0.27	0.19	(-0.10, 0.64)
SOUTH CAROLINA	605	5.68	1.11	(3.50, 7.86)	888	1.18	0.40	(0.40, 1.96)
SOUTH DAKOTA	894	10.81	1.19	(8.48, 13.14)	1206	0.39	0.17	(0.06, 0.72)
TEXAS	666	7.95	1.15	(5.70, 10.20)	1038	0.42	0.30	(-0.17, 1.01)
VIRGINIA	730	7.51	1.13	(5.30, 9.72)	1158	0.34	0.18	(-0.01, 0.69)
WEST VIRGINIA	896	17.61	1.37	(14.92, 20.30)	1514	0.44	0.17	(0.11, 0.77)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	18	NUMBER OF PARTICIPANTS =	18
	MEDIAN	= 8.35	MEDIAN	= 0.40
	RANGE	= 2.60-17.61	RANGE	= 0.00- 2.35

* CURRENT USERS OF SMOKELESS TOBACCO December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET*
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	44	41	4.45	(32.28, 49.72)
ARIZONA	68	46	2.13	(41.83, 50.17)
ARKANSAS	78	46	2.42	(41.26, 50.74)
CALIFORNIA	194	45	1.46	(42.14, 47.86)
COLORADO	58	49	2.60	(43.90, 54.10)
DISTRICT OF CO	60	43	3.14	(36.85, 49.15)
FLORIDA	179	48	1.74	(44.59, 51.41)
IDAHO	86	49	2.11	(44.86, 53.14)
INDIANA	109	50	1.44	(47.18, 52.82)
IOWA	170	50	1.59	(46.88, 53.12)
KANSAS	68	52	1.90	(48.28, 55.72)
KENTUCKY	173	50	1.19	(47.67, 52.33)
MAINE	79	49	1.91	(45.26, 52.74)
MASSACHUSETTS	83	51	1.85	(47.37, 54.63)
MICHIGAN	143	48	1.35	(45.35, 50.65)
MINNESOTA	192	47	1.58	(43.90, 50.10)
MISSISSIPPI	96	50	1.61	(46.84, 53.16)
MISSOURI	69	50	1.84	(46.39, 53.61)
MONTANA	72	49	1.95	(45.18, 52.82)
NEBRASKA	112	48	1.90	(44.28, 51.72)
NEVADA	73	42	2.59	(36.92, 47.08)
NEW HAMPSHIRE	61	51	1.80	(47.47, 54.53)
NEW JERSEY	132	50	1.50	(47.06, 52.94)
NORTH CAROLINA	137	49	1.33	(46.39, 51.61)
NORTH DAKOTA	59	53	2.25	(48.59, 57.41)
OHIO	86	49	2.12	(44.84, 53.16)
OKLAHOMA	78	47	2.02	(43.04, 50.96)
OREGON	127	48	1.41	(45.24, 50.76)
PENNSYLVANIA	200	49	1.55	(45.96, 52.04)
SOUTH CAROLINA	102	48	1.76	(44.55, 51.45)
TENNESSEE	148	45	1.51	(42.04, 47.96)
TEXAS	67	48	2.10	(43.88, 52.12)
UTAH	111	49	2.42	(44.26, 53.74)
VERMONT	102	47	1.66	(43.75, 50.25)
VIRGINIA	101	47	1.80	(43.47, 50.53)
WISCONSIN	104	53	1.77	(49.53, 56.47)
WYOMING	103	47	1.85	(43.37, 50.63)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 37 MEDIAN = 49.00 RANGE = 41.00-53.00
 * MEAN AGE AT ONSET OF DIABETES December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET* -BY SEX
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	19	41	6.90	(27.48, 54.52)	* 25	42	5.50	(31.22, 52.78)
ARIZONA	36	48	2.40	(43.30, 52.70)	* 32	43	3.59	(35.96, 50.04)
ARKANSAS	27	48	3.81	(40.53, 55.47)	* 51	44	2.88	(38.36, 49.64)
CALIFORNIA	92	44	2.24	(39.61, 48.39)	* 102	46	1.79	(42.49, 49.51)
COLORADO	25	53	3.83	(45.49, 60.51)	* 33	45	3.21	(38.71, 51.29)
DISTRICT OF CO	24	38	4.95	(28.30, 47.70)	* 36	48	2.66	(42.79, 53.21)
FLORIDA	79	48	2.72	(42.67, 53.33)	* 100	48	2.12	(43.84, 52.16)
IDAHO	43	51	2.18	(46.73, 55.27)	* 43	46	3.49	(39.16, 52.84)
INDIANA	49	52	2.03	(48.02, 55.98)	* 60	48	1.99	(44.10, 51.90)
IOWA	58	50	2.90	(44.32, 55.68)	* 112	50	1.74	(46.59, 53.41)
KANSAS	29	51	3.02	(45.08, 56.92)	* 39	53	2.43	(48.24, 57.76)
KENTUCKY	61	48	2.09	(43.90, 52.10)	* 112	52	1.24	(49.57, 54.43)
MAINE	31	49	3.04	(43.04, 54.96)	* 48	49	2.41	(44.28, 53.72)
MASSACHUSETTS	31	50	2.75	(44.61, 55.39)	* 52	51	2.41	(46.28, 55.72)
MICHIGAN	53	45	2.08	(40.92, 49.08)	* 90	50	1.69	(46.69, 53.31)
MINNESOTA	83	46	2.40	(41.30, 50.70)	* 109	47	2.19	(42.71, 51.29)
MISSISSIPPI	32	48	2.62	(42.86, 53.14)	* 64	52	1.93	(48.22, 55.78)
MISSOURI	25	55	2.06	(50.96, 59.04)	* 44	47	2.69	(41.73, 52.27)
MONTANA	30	49	2.87	(43.37, 54.63)	* 42	49	2.52	(44.06, 53.94)
NEBRASKA	47	47	3.07	(40.98, 53.02)	* 65	49	2.17	(44.75, 53.25)
NEVADA	45	43	3.01	(37.10, 48.90)	* 28	39	4.55	(30.08, 47.92)
NEW HAMPSHIRE	27	49	3.06	(43.00, 55.00)	* 34	52	2.13	(47.83, 56.17)
NEW JERSEY	58	51	2.38	(46.34, 55.66)	* 74	50	1.88	(46.32, 53.68)
NORTH CAROLINA	51	50	1.76	(46.55, 53.45)	* 86	49	1.83	(45.41, 52.59)
NORTH DAKOTA	19	51	4.98	(41.24, 60.76)	* 40	55	2.09	(50.90, 59.10)
OHIO	29	50	3.37	(43.39, 56.61)	* 57	49	2.66	(43.79, 54.21)
OKLAHOMA	29	47	3.48	(40.18, 53.82)	* 49	47	2.44	(42.22, 51.78)
OREGON	51	48	2.32	(43.45, 52.55)	* 76	48	1.74	(44.59, 51.41)
PENNSYLVANIA	73	48	2.66	(42.79, 53.21)	* 127	49	1.86	(45.35, 52.65)
SOUTH CAROLINA	37	46	2.86	(40.39, 51.61)	* 65	49	2.38	(44.34, 53.66)
TENNESSEE	50	47	2.16	(42.77, 51.23)	* 98	44	2.09	(39.90, 48.10)
TEXAS	22	50	3.49	(43.16, 56.84)	* 45	47	2.47	(42.16, 51.84)
UTAH	59	52	2.67	(46.77, 57.23)	* 52	45	3.87	(37.41, 52.59)
VERMONT	44	48	2.45	(43.20, 52.80)	* 58	47	2.23	(42.63, 51.37)
VIRGINIA	31	43	3.23	(36.67, 49.33)	* 70	48	2.08	(43.92, 52.08)
WISCONSIN	42	53	2.52	(48.06, 57.94)	* 62	53	2.49	(48.12, 57.88)
WYOMING	38	50	2.43	(45.24, 54.76)	* 65	44	2.63	(38.85, 49.15)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	37	NUMBER OF PARTICIPANTS =	37
	MEDIAN	= 48.00	MEDIAN	= 48.00
	RANGE	= 38.00-55.00	RANGE	= 39.00-55.00

* MEAN AGE AT ONSET OF DIABETES December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TAKING INSULIN*
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	46	48.46	11.58	(25.76, 71.16)
ARIZONA	69	38.53	7.60	(23.63, 53.43)
ARKANSAS	89	41.42	5.82	(30.01, 52.83)
CALIFORNIA	198	34.14	3.95	(26.40, 41.88)
COLORADO	58	34.07	6.77	(20.80, 47.34)
DISTRICT OF CO	66	60.59	7.24	(46.40, 74.78)
FLORIDA	179	43.40	4.25	(35.07, 51.73)
IDAHO	89	34.23	5.53	(23.39, 45.07)
INDIANA	114	32.63	4.74	(23.34, 41.92)
IOWA	173	35.65	3.99	(27.83, 43.47)
KANSAS	71	27.38	6.11	(15.40, 39.36)
KENTUCKY	185	34.06	3.95	(26.32, 41.80)
MAINE	84	21.55	4.89	(11.97, 31.13)
MASSACHUSETTS	86	35.45	5.26	(25.14, 45.76)
MICHIGAN	146	35.26	4.61	(26.22, 44.30)
MINNESOTA	195	43.66	3.92	(35.98, 51.34)
MISSISSIPPI	110	40.61	5.46	(29.91, 51.31)
MISSOURI	70	31.61	5.80	(20.24, 42.98)
MONTANA	73	31.08	5.64	(20.03, 42.13)
NEBRASKA	119	43.92	5.04	(34.04, 53.80)
NEVADA	75	36.70	7.04	(22.90, 50.50)
NEW HAMPSHIRE	65	26.34	5.82	(14.93, 37.75)
NEW JERSEY	135	34.62	4.82	(25.17, 44.07)
NORTH CAROLINA	149	35.99	4.53	(27.11, 44.87)
NORTH DAKOTA	61	43.19	7.06	(29.35, 57.03)
OHIO	87	30.92	5.50	(20.14, 41.70)
OKLAHOMA	85	41.49	6.04	(29.65, 53.33)
OREGON	132	32.37	4.32	(23.90, 40.84)
PENNSYLVANIA	214	34.87	3.74	(27.54, 42.20)
SOUTH CAROLINA	106	39.75	5.47	(29.03, 50.47)
TENNESSEE	164	40.20	4.26	(31.85, 48.55)
TEXAS	70	24.92	5.49	(14.16, 35.68)
UTAH	113	34.84	6.02	(23.04, 46.64)
VERMONT	107	36.64	5.08	(26.68, 46.60)
VIRGINIA	111	35.35	5.05	(25.45, 45.25)
WISCONSIN	105	34.98	7.55	(20.18, 49.78)
WYOMING	105	38.28	5.12	(28.24, 48.32)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 37 MEDIAN = 35.35 RANGE = 21.55-60.59
 * NOW TAKING INSULIN

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR HEALTH PROF VISITS FOR DIABETES*
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	39	3	0.53	(1.96, 4.04)
ARIZONA	52	4	0.54	(2.94, 5.06)
ARKANSAS	74	8	1.18	(5.69, 10.31)
CALIFORNIA	152	5	0.41	(4.20, 5.80)
COLORADO	42	4	0.85	(2.33, 5.67)
DISTRICT OF CO	56	5	0.49	(4.04, 5.96)
FLORIDA	155	7	0.66	(5.71, 8.29)
IDAHO	70	6	0.91	(4.22, 7.78)
INDIANA	95	5	0.92	(3.20, 6.80)
IOWA	147	6	0.52	(4.98, 7.02)
KANSAS	53	7	1.61	(3.84, 10.16)
KENTUCKY	154	5	0.30	(4.41, 5.59)
MAINE	68	5	0.53	(3.96, 6.04)
MASSACHUSETTS	67	8	1.55	(4.96, 11.04)
MICHIGAN	115	6	0.51	(5.00, 7.00)
MINNESOTA	175	5	0.38	(4.26, 5.74)
MISSISSIPPI	89	7	0.79	(5.45, 8.55)
MISSOURI	60	5	0.47	(4.08, 5.92)
MONTANA	58	5	0.46	(4.10, 5.90)
NEBRASKA	102	6	0.69	(4.65, 7.35)
NEVADA	66	5	0.77	(3.49, 6.51)
NEW HAMPSHIRE	55	4	0.45	(3.12, 4.88)
NEW JERSEY	122	6	0.46	(5.10, 6.90)
NORTH CAROLINA	123	6	0.77	(4.49, 7.51)
NORTH DAKOTA	59	4	0.46	(3.10, 4.90)
OHIO	77	4	0.41	(3.20, 4.80)
OKLAHOMA	77	6	0.59	(4.84, 7.16)
OREGON	106	6	1.14	(3.77, 8.23)
PENNSYLVANIA	171	6	0.49	(5.04, 6.96)
SOUTH CAROLINA	86	6	0.65	(4.73, 7.27)
TENNESSEE	130	5	0.36	(4.29, 5.71)
TEXAS	62	6	1.01	(4.02, 7.98)
UTAH	93	4	0.52	(2.98, 5.02)
VERMONT	95	4	0.47	(3.08, 4.92)
VIRGINIA	84	6	0.75	(4.53, 7.47)
WISCONSIN	86	4	0.88	(2.28, 5.72)
WYOMING	83	5	0.73	(3.57, 6.43)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 37 MEDIAN = 5.00 RANGE = 3.00- 8.00
 * MEAN TIMES SEEN HEALTH PROF FOR DIABETES IN PAST YEAR December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR HEALTH PROF VISITS FOR DIABETES* -BY SEX
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	18	3	0.45	(2.12, 3.88)	21	4	0.89	(2.26, 5.74)
ARIZONA	26	4	0.56	(2.90, 5.10)	26	5	0.93	(3.18, 6.82)
ARKANSAS	26	6	1.49	(3.08, 8.92)	48	9	1.61	(5.84, 12.16)
CALIFORNIA	68	5	0.65	(3.73, 6.27)	84	6	0.52	(4.98, 7.02)
COLORADO	19	5	1.51	(2.04, 7.96)	23	3	0.58	(1.86, 4.14)
DISTRICT OF CO	22	4	0.49	(3.04, 4.96)	34	6	0.79	(4.45, 7.55)
FLORIDA	68	7	1.01	(5.02, 8.98)	87	7	0.88	(5.28, 8.72)
IDAHO	34	3	0.46	(2.10, 3.90)	36	9	1.73	(5.61, 12.39)
INDIANA	41	4	0.54	(2.94, 5.06)	54	7	1.76	(3.55, 10.45)
IOWA	52	6	0.84	(4.35, 7.65)	95	6	0.64	(4.75, 7.25)
KANSAS	23	4	0.51	(3.00, 5.00)	30	9	2.87	(3.37, 14.63)
KENTUCKY	52	5	0.50	(4.02, 5.98)	102	5	0.37	(4.27, 5.73)
MAINE	26	5	0.85	(3.33, 6.67)	42	4	0.63	(2.77, 5.23)
MASSACHUSETTS	22	13	3.54	(6.06, 19.94)	45	5	0.52	(3.98, 6.02)
MICHIGAN	40	5	0.53	(3.96, 6.04)	75	7	0.76	(5.51, 8.49)
MINNESOTA	76	6	0.65	(4.73, 7.27)	99	5	0.40	(4.22, 5.78)
MISSISSIPPI	28	9	1.60	(5.86, 12.14)	61	6	0.58	(4.86, 7.14)
MISSOURI	21	4	0.63	(2.77, 5.23)	39	6	0.62	(4.78, 7.22)
MONTANA	25	4	0.56	(2.90, 5.10)	33	5	0.70	(3.63, 6.37)
NEBRASKA	40	7	1.10	(4.84, 9.16)	62	6	0.86	(4.31, 7.69)
NEVADA	40	5	0.95	(3.14, 6.86)	26	5	1.33	(2.39, 7.61)
NEW HAMPSHIRE	24	4	0.70	(2.63, 5.37)	31	5	0.60	(3.82, 6.18)
NEW JERSEY	56	6	0.59	(4.84, 7.16)	66	6	0.71	(4.61, 7.39)
NORTH CAROLINA	44	7	1.65	(3.77, 10.23)	79	5	0.53	(3.96, 6.04)
NORTH DAKOTA	18	5	0.69	(3.65, 6.35)	41	4	0.61	(2.80, 5.20)
OHIO	25	5	0.58	(3.86, 6.14)	52	4	0.55	(2.92, 5.08)
OKLAHOMA	29	5	0.67	(3.69, 6.31)	48	6	0.89	(4.26, 7.74)
OREGON	38	4	0.45	(3.12, 4.88)	68	6	1.83	(2.41, 9.59)
PENNSYLVANIA	55	7	1.20	(4.65, 9.35)	116	5	0.35	(4.31, 5.69)
SOUTH CAROLINA	29	6	0.91	(4.22, 7.78)	57	7	0.76	(5.51, 8.49)
TENNESSEE	41	5	0.57	(3.88, 6.12)	89	5	0.46	(4.10, 5.90)
TEXAS	17	5	1.08	(2.88, 7.12)	45	6	1.53	(3.00, 9.00)
UTAH	49	5	0.65	(3.73, 6.27)	44	4	0.87	(2.29, 5.71)
VERMONT	42	4	0.53	(2.96, 5.04)	53	5	0.75	(3.53, 6.47)
VIRGINIA	27	5	1.01	(3.02, 6.98)	57	6	1.01	(4.02, 7.98)
WISCONSIN	37	4	1.43	(1.20, 6.80)	49	4	0.51	(3.00, 5.00)
WYOMING	27	4	0.66	(2.71, 5.29)	56	6	1.09	(3.86, 8.14)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	37	NUMBER OF PARTICIPANTS =	37		
	MEDIAN	=	5.00	MEDIAN	=	6.00
	RANGE	=	3.00-13.00	RANGE	=	3.00- 9.00

* MEAN TIMES SEEN HEALTH PROF FOR DIABETES IN PAST YEAR December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PARTICULAR PRIMARY CARE CENTER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
INDIANA	2195	86.16	0.81	(84.57, 87.75)
KANSAS	1925	90.38	0.80	(88.81, 91.95)
KENTUCKY	1801	89.64	0.85	(87.97, 91.31)
NEBRASKA	2059	91.94	0.73	(90.51, 93.37)
NEW JERSEY	3028	84.73	0.81	(83.14, 86.32)
NEW YORK	4295	90.85	0.50	(89.87, 91.83)
NORTH DAKOTA	1795	93.79	0.62	(92.57, 95.01)
OHIO	1532	96.82	0.51	(95.82, 97.82)
OKLAHOMA	1913	91.03	0.85	(89.36, 92.70)
VIRGINIA	1870	88.09	0.90	(86.33, 89.85)
WASHINGTON	3540	83.41	0.73	(81.98, 84.84)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 11 MEDIAN = 90.38 RANGE = 83.41-96.82
 * ONE OR MORE PARTICULAR PLACE FOR HEALTH CARE December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PARTICULAR PRIMARY CARE CENTER* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----*				*-----FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
INDIANA	936	80.33	1.38	(77.63, 83.03)	* 1259	91.47	0.89	(89.73, 93.21)
KANSAS	830	87.67	1.35	(85.02, 90.32)	* 1095	92.93	0.89	(91.19, 94.67)
KENTUCKY	647	86.72	1.49	(83.80, 89.64)	* 1154	92.21	0.82	(90.60, 93.82)
NEBRASKA	819	88.89	1.33	(86.28, 91.50)	* 1240	94.71	0.67	(93.40, 96.02)
NEW JERSEY	1307	79.34	1.38	(76.64, 82.04)	* 1721	89.80	0.80	(88.23, 91.37)
NEW YORK	1737	87.93	0.86	(86.24, 89.62)	* 2558	93.43	0.54	(92.37, 94.49)
NORTH DAKOTA	737	91.32	1.08	(89.20, 93.44)	* 1058	96.21	0.64	(94.96, 97.46)
OHIO	619	95.51	0.91	(93.73, 97.29)	* 913	97.99	0.51	(96.99, 98.99)
OKLAHOMA	850	88.62	1.40	(85.88, 91.36)	* 1063	93.24	1.01	(91.26, 95.22)
VIRGINIA	725	84.38	1.54	(81.36, 87.40)	* 1145	91.54	0.96	(89.66, 93.42)
WASHINGTON	1505	75.59	1.24	(73.16, 78.02)	* 2035	90.97	0.74	(89.52, 92.42)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	11	NUMBER OF PARTICIPANTS =	11
	MEDIAN	= 87.67	MEDIAN	= 92.93
	RANGE	= 75.59-95.51	RANGE	= 89.80-97.99

* ONE OR MORE PARTICULAR PLACE FOR HEALTH CARE

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR RECENT DENTAL VISIT*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	1866	69.60	1.51	(66.64, 72.56)
CONNECTICUT	1754	77.26	1.18	(74.95, 79.57)
ILLINOIS	1419	67.50	1.39	(64.78, 70.22)
INDIANA	2179	64.89	1.16	(62.62, 67.16)
KANSAS	1875	68.37	1.18	(66.06, 70.68)
KENTUCKY	1788	61.68	1.37	(58.99, 64.37)
LOUISIANA	1608	62.76	1.33	(60.15, 65.37)
MICHIGAN	2471	73.27	0.98	(71.35, 75.19)
MONTANA	1790	69.58	1.26	(67.11, 72.05)
NEBRASKA	2039	70.95	1.36	(68.28, 73.62)
NEVADA	1794	62.54	1.63	(59.35, 65.73)
NEW HAMPSHIRE	1444	74.66	1.27	(72.17, 77.15)
NEW JERSEY	2991	71.40	1.00	(69.44, 73.36)
NEW YORK	979	71.53	1.58	(68.43, 74.63)
OKLAHOMA	1902	60.28	1.32	(57.69, 62.87)
PENNSYLVANIA	3416	69.95	0.89	(68.21, 71.69)
SOUTH DAKOTA	2073	71.71	1.10	(69.55, 73.87)
TEXAS	1668	64.01	1.30	(61.46, 66.56)
UTAH	2879	71.49	1.07	(69.39, 73.59)
VIRGINIA	1854	72.96	1.20	(70.61, 75.31)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 20 MEDIAN = 69.78 RANGE = 60.28-77.26
 * PERSONS VISITING A DENTIST OR DENTAL CLINIC WITHIN THE PAST YEAR December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR RECENT DENTAL VISIT* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	773	65.73	2.45	(60.93, 70.53)	*	1093	73.20	1.77	(69.73, 76.67)
CONNECTICUT	709	77.44	1.73	(74.05, 80.83)	*	1045	77.10	1.61	(73.94, 80.26)
ILLINOIS	575	67.02	2.06	(62.98, 71.06)	*	844	67.96	1.75	(64.53, 71.39)
INDIANA	933	63.26	1.70	(59.93, 66.59)	*	1246	66.39	1.52	(63.41, 69.37)
KANSAS	808	69.11	1.78	(65.62, 72.60)	*	1067	67.67	1.55	(64.63, 70.71)
KENTUCKY	641	60.67	2.14	(56.48, 64.86)	*	1147	62.56	1.65	(59.33, 65.79)
LOUISIANA	640	60.13	2.20	(55.82, 64.44)	*	968	65.08	1.69	(61.77, 68.39)
MICHIGAN	1038	73.70	1.48	(70.80, 76.60)	*	1433	72.89	1.30	(70.34, 75.44)
MONTANA	765	68.20	1.93	(64.42, 71.98)	*	1025	70.90	1.57	(67.82, 73.98)
NEBRASKA	814	68.50	2.33	(63.93, 73.07)	*	1225	73.19	1.43	(70.39, 75.99)
NEVADA	756	60.95	2.49	(56.07, 65.83)	*	1038	64.15	2.09	(60.05, 68.25)
NEW HAMPSHIRE	627	73.09	1.96	(69.25, 76.93)	*	817	76.16	1.62	(72.98, 79.34)
NEW JERSEY	1294	69.26	1.59	(66.14, 72.38)	*	1697	73.42	1.25	(70.97, 75.87)
NEW YORK	390	70.74	2.45	(65.94, 75.54)	*	589	72.19	2.13	(68.02, 76.36)
OKLAHOMA	844	57.69	2.00	(53.77, 61.61)	*	1058	62.65	1.69	(59.34, 65.96)
PENNSYLVANIA	1464	68.70	1.39	(65.98, 71.42)	*	1952	71.07	1.18	(68.76, 73.38)
SOUTH DAKOTA	884	69.76	1.71	(66.41, 73.11)	*	1189	73.53	1.32	(70.94, 76.12)
TEXAS	651	63.10	2.06	(59.06, 67.14)	*	1017	64.88	1.62	(61.70, 68.06)
UTAH	1253	69.93	1.59	(66.81, 73.05)	*	1626	72.97	1.44	(70.15, 75.79)
VIRGINIA	725	73.23	1.90	(69.51, 76.95)	*	1129	72.70	1.49	(69.78, 75.62)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS = 20	NUMBER OF PARTICIPANTS = 20
	MEDIAN = 68.60	MEDIAN = 71.63
	RANGE = 57.69-77.44	RANGE = 62.56-77.10
* PERSONS VISITING A DENTIST OR DENTAL CLINIC WITHIN THE PAST YEAR		December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOOTH LOSS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	1865	15.61	1.05	(13.55, 17.67)
CONNECTICUT	1731	18.15	1.08	(16.03, 20.27)
ILLINOIS	1424	23.09	1.25	(20.64, 25.54)
INDIANA	2161	25.37	1.01	(23.39, 27.35)
KANSAS	1884	19.35	0.96	(17.47, 21.23)
KENTUCKY	1788	32.44	1.29	(29.91, 34.97)
LOUISIANA	1623	23.66	1.20	(21.31, 26.01)
MICHIGAN	2449	20.63	0.91	(18.85, 22.41)
MONTANA	1777	20.37	1.00	(18.41, 22.33)
NEBRASKA	2045	20.31	0.96	(18.43, 22.19)
NEVADA	1780	22.09	1.44	(19.27, 24.91)
NEW HAMPSHIRE	1432	19.94	1.20	(17.59, 22.29)
NEW JERSEY	2981	19.39	0.81	(17.80, 20.98)
NEW YORK	981	22.18	1.64	(18.97, 25.39)
OKLAHOMA	1906	24.76	1.10	(22.60, 26.92)
PENNSYLVANIA	3388	25.07	0.81	(23.48, 26.66)
SOUTH DAKOTA	2071	21.29	0.92	(19.49, 23.09)
TEXAS	1684	17.70	1.01	(15.72, 19.68)
UTAH	2866	13.50	0.81	(11.91, 15.09)
VIRGINIA	1832	19.57	1.07	(17.47, 21.67)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 20 MEDIAN = 20.50 RANGE = 13.50-32.44
 * PERSONS WITH LOSS OF 6 OR MORE TEETH FROM DECAY OR GUM DISEASE December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOLD BLOOD PRESSURE IS HIGH*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2075	28.32	1.08	(26.20, 30.44)
ILLINOIS	1435	23.68	1.30	(21.13, 26.23)
IOWA	3512	23.01	0.75	(21.54, 24.48)
KENTUCKY	1791	25.48	1.15	(23.23, 27.73)
MAINE	1681	23.17	1.09	(21.03, 25.31)
MISSISSIPPI	1576	29.56	1.27	(27.07, 32.05)
NEW YORK	4291	23.36	0.76	(21.87, 24.85)
OKLAHOMA	1913	23.19	1.10	(21.03, 25.35)
PENNSYLVANIA	3427	24.24	0.83	(22.61, 25.87)
SOUTH CAROLINA	1821	25.50	1.20	(23.15, 27.85)
SOUTH DAKOTA	2092	20.22	0.91	(18.44, 22.00)
TENNESSEE	1432	29.33	1.29	(26.80, 31.86)
VIRGINIA	1886	22.93	1.12	(20.73, 25.13)
WEST VIRGINIA	2405	27.38	1.03	(25.36, 29.40)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 14 MEDIAN = 23.96 RANGE = 20.22-29.56

* EVER TOLD BLOOD PRESSURE IS HIGH

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOLD BLOOD PRESSURE IS HIGH* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	757	26.44	1.70	(23.11, 29.77)	1318	29.97	1.42	(27.19, 32.75)
ILLINOIS	566	22.35	1.99	(18.45, 26.25)	869	24.81	1.67	(21.54, 28.08)
IOWA	1458	20.98	1.17	(18.69, 23.27)	2054	24.88	1.02	(22.88, 26.88)
KENTUCKY	655	23.40	1.76	(19.95, 26.85)	1136	27.37	1.47	(24.49, 30.25)
MAINE	703	21.60	1.57	(18.52, 24.68)	978	24.62	1.49	(21.70, 27.54)
MISSISSIPPI	581	27.14	2.05	(23.12, 31.16)	995	31.67	1.48	(28.77, 34.57)
NEW YORK	1733	23.07	1.10	(20.91, 25.23)	2558	23.62	0.97	(21.72, 25.52)
OKLAHOMA	848	23.52	1.70	(20.19, 26.85)	1065	22.88	1.43	(20.08, 25.68)
PENNSYLVANIA	1470	23.29	1.25	(20.84, 25.74)	1957	25.09	1.11	(22.91, 27.27)
SOUTH CAROLINA	723	22.63	1.83	(19.04, 26.22)	1098	28.08	1.58	(24.98, 31.18)
SOUTH DAKOTA	890	18.70	1.37	(16.01, 21.39)	1202	21.64	1.24	(19.21, 24.07)
TENNESSEE	555	28.77	2.03	(24.79, 32.75)	877	29.85	1.65	(26.62, 33.08)
VIRGINIA	731	21.24	1.72	(17.87, 24.61)	1155	24.52	1.44	(21.70, 27.34)
WEST VIRGINIA	893	27.45	1.63	(24.26, 30.64)	1512	27.32	1.26	(24.85, 29.79)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	14	NUMBER OF PARTICIPANTS =	14
	MEDIAN	= 23.18	MEDIAN	= 24.99
	RANGE	= 18.70-28.77	RANGE	= 21.64-31.67

* EVER TOLD BLOOD PRESSURE IS HIGH
 December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BP CHECKED WITHIN LAST 2 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2065	94.62	0.58	(93.48, 95.76)
ILLINOIS	1432	94.58	0.69	(93.23, 95.93)
IOWA	3508	93.66	0.46	(92.76, 94.56)
KENTUCKY	1785	94.55	0.60	(93.37, 95.73)
MAINE	1671	93.49	0.69	(92.14, 94.84)
MISSISSIPPI	1559	95.71	0.63	(94.48, 96.94)
NEW YORK	4254	96.05	0.33	(95.40, 96.70)
OKLAHOMA	1899	95.00	0.64	(93.75, 96.25)
PENNSYLVANIA	3397	95.79	0.37	(95.06, 96.52)
SOUTH CAROLINA	1799	97.29	0.49	(96.33, 98.25)
SOUTH DAKOTA	2074	95.36	0.50	(94.38, 96.34)
TENNESSEE	1419	97.06	0.50	(96.08, 98.04)
VIRGINIA	1866	96.77	0.48	(95.83, 97.71)
WEST VIRGINIA	2390	94.68	0.51	(93.68, 95.68)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 14 MEDIAN = 95.18 RANGE = 93.49-97.29

* BLOOD PRESSURE CHECKED WITHIN THE PAST 2 YEARS December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BP CHECKED WITHIN LAST 2 YEARS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	753	93.26	1.03	(91.24, 95.28)	*	1312	95.81	0.61	(94.61, 97.01)
ILLINOIS	563	91.78	1.28	(89.27, 94.29)	*	869	96.96	0.66	(95.67, 98.25)
IOWA	1454	92.32	0.76	(90.83, 93.81)	*	2054	94.90	0.54	(93.84, 95.96)
KENTUCKY	654	92.66	1.06	(90.58, 94.74)	*	1131	96.28	0.61	(95.08, 97.48)
MAINE	698	92.36	1.14	(90.13, 94.59)	*	973	94.52	0.77	(93.01, 96.03)
MISSISSIPPI	574	93.23	1.22	(90.84, 95.62)	*	985	97.87	0.49	(96.91, 98.83)
NEW YORK	1715	94.74	0.56	(93.64, 95.84)	*	2539	97.21	0.37	(96.48, 97.94)
OKLAHOMA	842	93.38	1.12	(91.18, 95.58)	*	1057	96.47	0.69	(95.12, 97.82)
PENNSYLVANIA	1450	94.58	0.63	(93.35, 95.81)	*	1947	96.87	0.43	(96.03, 97.71)
SOUTH CAROLINA	715	96.48	0.86	(94.79, 98.17)	*	1084	98.02	0.49	(97.06, 98.98)
SOUTH DAKOTA	878	93.41	0.89	(91.67, 95.15)	*	1196	97.15	0.52	(96.13, 98.17)
TENNESSEE	549	95.11	0.95	(93.25, 96.97)	*	870	98.86	0.36	(98.15, 99.57)
VIRGINIA	723	95.57	0.88	(93.85, 97.29)	*	1143	97.89	0.43	(97.05, 98.73)
WEST VIRGINIA	885	94.16	0.80	(92.59, 95.73)	*	1505	95.14	0.64	(93.89, 96.39)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	14	NUMBER OF PARTICIPANTS =	14
	MEDIAN	= 93.40	MEDIAN	= 96.92
	RANGE	= 91.78-96.48	RANGE	= 94.52-98.86

* BLOOD PRESSURE CHECKED WITHIN THE PAST 2 YEARS
 December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1997	69.65	1.22	(67.26, 72.04)
ILLINOIS	1421	72.47	1.35	(69.82, 75.12)
IOWA	3398	71.02	0.91	(69.24, 72.80)
KENTUCKY	1738	68.19	1.25	(65.74, 70.64)
MAINE	1647	73.61	1.28	(71.10, 76.12)
NEW JERSEY	2945	79.58	0.92	(77.78, 81.38)
OKLAHOMA	1893	71.75	1.22	(69.36, 74.14)
PENNSYLVANIA	3324	72.31	0.92	(70.51, 74.11)
SOUTH CAROLINA	1759	75.79	1.39	(73.07, 78.51)
TENNESSEE	1394	74.97	1.28	(72.46, 77.48)
VERMONT	2402	76.35	1.00	(74.39, 78.31)
VIRGINIA	1836	75.69	1.21	(73.32, 78.06)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 12 MEDIAN = 73.04 RANGE = 68.19-79.58
 * EVER HAD CHOLESTEROL CHECKED

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	731	65.58	2.00	(61.66, 69.50)	*	1266	73.28	1.42	(70.50, 76.06)
ILLINOIS	563	68.76	2.13	(64.59, 72.93)	*	858	75.70	1.63	(72.51, 78.89)
IOWA	1407	67.83	1.38	(65.13, 70.53)	*	1991	73.95	1.13	(71.74, 76.16)
KENTUCKY	653	64.44	2.10	(60.32, 68.56)	*	1085	71.77	1.56	(68.71, 74.83)
MAINE	688	68.69	2.01	(64.75, 72.63)	*	959	78.15	1.61	(74.99, 81.31)
NEW JERSEY	1278	77.11	1.50	(74.17, 80.05)	*	1667	81.91	1.09	(79.77, 84.05)
OKLAHOMA	838	71.10	1.89	(67.40, 74.80)	*	1055	72.35	1.56	(69.29, 75.41)
PENNSYLVANIA	1422	69.89	1.39	(67.17, 72.61)	*	1902	74.48	1.22	(72.09, 76.87)
SOUTH CAROLINA	703	74.54	2.26	(70.11, 78.97)	*	1056	76.93	1.64	(73.72, 80.14)
TENNESSEE	540	71.93	2.00	(68.01, 75.85)	*	854	77.76	1.63	(74.57, 80.95)
VERMONT	1031	74.10	1.53	(71.10, 77.10)	*	1371	78.38	1.31	(75.81, 80.95)
VIRGINIA	714	73.20	1.94	(69.40, 77.00)	*	1122	78.02	1.48	(75.12, 80.92)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	12	NUMBER OF PARTICIPANTS =	12
	MEDIAN	= 70.50	MEDIAN	= 76.32
	RANGE	= 64.44-77.11	RANGE	= 71.77-81.91

* EVER HAD CHOLESTEROL CHECKED

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1969	65.34	1.26	(62.87, 67.81)
ILLINOIS	1414	68.95	1.44	(66.13, 71.77)
IOWA	3378	64.48	0.94	(62.64, 66.32)
KENTUCKY	1708	63.15	1.28	(60.64, 65.66)
MAINE	1631	67.22	1.41	(64.46, 69.98)
NEW JERSEY	2904	75.70	0.97	(73.80, 77.60)
OKLAHOMA	1881	68.72	1.27	(66.23, 71.21)
PENNSYLVANIA	3278	67.25	0.96	(65.37, 69.13)
SOUTH CAROLINA	1742	73.66	1.40	(70.92, 76.40)
TENNESSEE	1373	72.34	1.31	(69.77, 74.91)
VERMONT	2366	70.02	1.07	(67.92, 72.12)
VIRGINIA	1808	71.80	1.28	(69.29, 74.31)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 12 MEDIAN = 68.84 RANGE = 63.15-75.70 YEAR 2000 OBJ. 15.14
 * CHOLESTEROL CHECKED WITHIN THE PAST 5 YEARS December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	728	60.64	2.04	(56.64, 64.64)	*	1241	69.58	1.48	(66.68, 72.48)
ILLINOIS	559	65.95	2.21	(61.62, 70.28)	*	855	71.55	1.74	(68.14, 74.96)
IOWA	1399	60.74	1.44	(57.92, 63.56)	*	1979	67.92	1.20	(65.57, 70.27)
KENTUCKY	645	59.55	2.16	(55.32, 63.78)	*	1063	66.61	1.65	(63.38, 69.84)
MAINE	678	62.24	2.23	(57.87, 66.61)	*	953	71.76	1.74	(68.35, 75.17)
NEW JERSEY	1261	73.06	1.55	(70.02, 76.10)	*	1643	78.19	1.18	(75.88, 80.50)
OKLAHOMA	832	68.44	1.93	(64.66, 72.22)	*	1049	68.98	1.63	(65.79, 72.17)
PENNSYLVANIA	1406	65.80	1.46	(62.94, 68.66)	*	1872	68.57	1.28	(66.06, 71.08)
SOUTH CAROLINA	699	72.64	2.26	(68.21, 77.07)	*	1043	74.59	1.68	(71.30, 77.88)
TENNESSEE	530	69.39	2.12	(65.23, 73.55)	*	843	75.04	1.68	(71.75, 78.33)
VERMONT	1019	68.06	1.62	(64.88, 71.24)	*	1347	71.82	1.41	(69.06, 74.58)
VIRGINIA	709	69.11	2.04	(65.11, 73.11)	*	1099	74.38	1.55	(71.34, 77.42)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	12	NUMBER OF PARTICIPANTS =	12
	MEDIAN	= 67.01	MEDIAN	= 71.66
	RANGE	= 59.55-73.06	RANGE	= 66.61-78.19

* CHOLESTEROL CHECKED WITHIN THE PAST 5 YEARS

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1984	22.46	1.01	(20.48, 24.44)
ILLINOIS	1419	21.08	1.22	(18.69, 23.47)
IOWA	3385	22.14	0.81	(20.55, 23.73)
KENTUCKY	1723	22.11	1.08	(19.99, 24.23)
MAINE	1634	23.07	1.15	(20.82, 25.32)
NEW JERSEY	2921	22.41	0.87	(20.70, 24.12)
OKLAHOMA	1886	18.48	0.99	(16.54, 20.42)
PENNSYLVANIA	3293	22.99	0.82	(21.38, 24.60)
SOUTH CAROLINA	1742	21.39	1.21	(19.02, 23.76)
TENNESSEE	1384	23.25	1.29	(20.72, 25.78)
VERMONT	2390	20.99	0.95	(19.13, 22.85)
VIRGINIA	1819	22.55	1.10	(20.39, 24.71)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 12 MEDIAN = 22.27 RANGE = 18.48-23.25

* EVER TOLD CHOLESTEROL IS HIGH BY HEALTH PROFESSIONAL

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL AWARENESS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*				-----FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	726	20.86	1.58	(17.76, 23.96)	1258	23.89	1.32	(21.30, 26.48)
ILLINOIS	562	18.37	1.76	(14.92, 21.82)	857	23.43	1.67	(20.16, 26.70)
IOWA	1402	20.05	1.19	(17.72, 22.38)	1983	24.06	1.07	(21.96, 26.16)
KENTUCKY	648	19.64	1.60	(16.50, 22.78)	1075	24.47	1.43	(21.67, 27.27)
MAINE	682	21.51	1.67	(18.24, 24.78)	952	24.52	1.51	(21.56, 27.48)
NEW JERSEY	1266	21.46	1.27	(18.97, 23.95)	1655	23.32	1.20	(20.97, 25.67)
OKLAHOMA	835	18.27	1.45	(15.43, 21.11)	1051	18.68	1.35	(16.03, 21.33)
PENNSYLVANIA	1410	23.16	1.27	(20.67, 25.65)	1883	22.85	1.08	(20.73, 24.97)
SOUTH CAROLINA	696	18.37	1.87	(14.70, 22.04)	1046	24.11	1.54	(21.09, 27.13)
TENNESSEE	537	19.78	1.90	(16.06, 23.50)	847	26.46	1.71	(23.11, 29.81)
VERMONT	1025	21.51	1.50	(18.57, 24.45)	1365	20.52	1.17	(18.23, 22.81)
VIRGINIA	709	19.88	1.57	(16.80, 22.96)	1110	25.07	1.54	(22.05, 28.09)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	12	NUMBER OF PARTICIPANTS =	12
	MEDIAN	= 19.96	MEDIAN	= 23.98
	RANGE	= 18.27-23.16	RANGE	= 18.68-26.46

* EVER TOLD CHOLESTEROL IS HIGH BY HEALTH PROFESSIONAL

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FLU SHOT*
 DENOMINATOR IS PERSONS AGE 65 AND OLDER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	450	71.08	2.77	(65.65, 76.51)
ILLINOIS	247	53.91	3.46	(47.13, 60.69)
KENTUCKY	462	60.98	2.42	(56.24, 65.72)
LOUISIANA	323	53.08	2.95	(47.30, 58.86)
MARYLAND	813	61.96	2.22	(57.61, 66.31)
MINNESOTA	772	66.81	1.84	(63.20, 70.42)
MONTANA	391	70.46	2.49	(65.58, 75.34)
NEW JERSEY	524	56.01	2.40	(51.31, 60.71)
OKLAHOMA	626	55.65	2.39	(50.97, 60.33)
OREGON	556	67.69	3.14	(61.54, 73.84)
PENNSYLVANIA	732	62.92	1.97	(59.06, 66.78)
SOUTH CAROLINA	311	57.42	3.43	(50.70, 64.14)
TENNESSEE	585	59.50	2.19	(55.21, 63.79)
TEXAS	265	57.03	3.35	(50.46, 63.60)
WYOMING	413	70.86	2.49	(65.98, 75.74)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 15 MEDIAN = 60.98 RANGE = 53.08-71.08 YEAR 2000 OBJ. 20.11
 * HAD FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OVER November 21, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FLU SHOT* -BY SEX
 DENOMINATOR IS PERSONS AGE 65 AND OLDER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	171	67.18	4.72	(57.93, 76.43)	*	279	73.96	3.11	(67.86, 80.06)
ILLINOIS	88	51.58	5.73	(40.35, 62.81)	*	159	55.48	4.60	(46.46, 64.50)
KENTUCKY	137	60.60	4.27	(52.23, 68.97)	*	325	61.25	2.86	(55.64, 66.86)
LOUISIANA	115	52.37	4.74	(43.08, 61.66)	*	208	53.55	3.78	(46.14, 60.96)
MARYLAND	300	64.54	3.49	(57.70, 71.38)	*	513	60.16	2.99	(54.30, 66.02)
MINNESOTA	272	64.98	3.12	(58.86, 71.10)	*	500	68.08	2.27	(63.63, 72.53)
MONTANA	141	70.05	3.93	(62.35, 77.75)	*	250	70.76	3.27	(64.35, 77.17)
NEW JERSEY	202	52.77	3.87	(45.18, 60.36)	*	322	58.27	3.06	(52.27, 64.27)
OKLAHOMA	290	54.65	3.32	(48.14, 61.16)	*	336	56.47	3.33	(49.94, 63.00)
OREGON	201	58.43	5.92	(46.83, 70.03)	*	355	74.39	2.50	(69.49, 79.29)
PENNSYLVANIA	272	66.98	3.10	(60.90, 73.06)	*	460	60.19	2.47	(55.35, 65.03)
SOUTH CAROLINA	87	59.90	6.34	(47.47, 72.33)	*	224	55.77	3.86	(48.20, 63.34)
TENNESSEE	196	65.00	3.61	(57.92, 72.08)	*	389	55.82	2.74	(50.45, 61.19)
TEXAS	86	62.62	5.65	(51.55, 73.69)	*	179	53.17	4.08	(45.17, 61.17)
WYOMING	143	69.61	4.00	(61.77, 77.45)	*	270	71.81	3.05	(65.83, 77.79)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	15	NUMBER OF PARTICIPANTS =	15
	MEDIAN	= 62.62	MEDIAN	= 60.16
	RANGE	= 51.58-70.05	RANGE	= 53.17-74.39
* HAD FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OVER			November 21, 1997	

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PNEUMONIA VACCINATION*
 DENOMINATOR IS PERSONS AGES 65 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	439	54.26	3.23	(47.93, 60.59)
ILLINOIS	243	34.58	3.57	(27.58, 41.58)
KENTUCKY	449	29.19	2.30	(24.68, 33.70)
LOUISIANA	321	25.42	2.56	(20.40, 30.44)
MARYLAND	800	40.48	2.31	(35.95, 45.01)
MINNESOTA	758	43.23	1.97	(39.37, 47.09)
MONTANA	384	45.86	2.66	(40.65, 51.07)
NEW JERSEY	512	30.81	2.21	(26.48, 35.14)
OKLAHOMA	616	29.24	2.12	(25.08, 33.40)
OREGON	540	47.94	2.87	(42.31, 53.57)
PENNSYLVANIA	710	41.66	1.98	(37.78, 45.54)
SOUTH CAROLINA	306	31.94	3.29	(25.49, 38.39)
TENNESSEE	578	41.33	2.22	(36.98, 45.68)
TEXAS	261	37.55	3.28	(31.12, 43.98)
WYOMING	405	48.18	2.74	(42.81, 53.55)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 15 MEDIAN = 40.48 RANGE = 25.42-54.26 YEAR 2000 OBJ. 20.11
 * EVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PNEUMONIA VACCINATION* -BY SEX
 DENOMINATOR IS PERSONS AGE 65 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	164	43.95	5.27	(33.62, 54.28)	275	61.61	3.85	(54.06, 69.16)
ILLINOIS	86	32.80	5.86	(21.31, 44.29)	157	35.77	4.41	(27.13, 44.41)
KENTUCKY	131	27.59	4.02	(19.71, 35.47)	318	30.29	2.83	(24.74, 35.84)
LOUISIANA	113	23.47	4.09	(15.45, 31.49)	208	26.69	3.40	(20.03, 33.35)
MARYLAND	292	38.75	3.67	(31.56, 45.94)	508	41.67	2.91	(35.97, 47.37)
MINNESOTA	266	44.27	3.27	(37.86, 50.68)	492	42.52	2.38	(37.86, 47.18)
MONTANA	138	46.85	4.28	(38.46, 55.24)	246	45.11	3.55	(38.15, 52.07)
NEW JERSEY	193	29.52	3.44	(22.78, 36.26)	319	31.68	2.85	(26.09, 37.27)
OKLAHOMA	282	23.27	2.88	(17.63, 28.91)	334	34.04	2.98	(28.20, 39.88)
OREGON	191	44.73	5.27	(34.40, 55.06)	349	50.19	3.02	(44.27, 56.11)
PENNSYLVANIA	259	42.02	3.32	(35.51, 48.53)	451	41.42	2.49	(36.54, 46.30)
SOUTH CAROLINA	86	25.15	5.83	(13.72, 36.58)	220	36.45	3.78	(29.04, 43.86)
TENNESSEE	195	43.24	3.88	(35.64, 50.84)	383	40.05	2.77	(34.62, 45.48)
TEXAS	85	36.28	5.55	(25.40, 47.16)	176	38.44	3.99	(30.62, 46.26)
WYOMING	141	46.77	4.46	(38.03, 55.51)	264	49.24	3.36	(42.65, 55.83)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	15	NUMBER OF PARTICIPANTS =	15
	MEDIAN	= 38.75	MEDIAN	= 40.05
	RANGE	= 23.27-46.85	RANGE	= 26.69-61.61

* EVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LACK OF SAFETY BELT USAGE*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	1927	24.63	1.52	(21.65, 27.61)
ILLINOIS	1430	35.44	1.54	(32.42, 38.46)
KENTUCKY	1776	34.55	1.32	(31.96, 37.14)
LOUISIANA	1640	22.00	1.20	(19.65, 24.35)
NEW JERSEY	2967	28.67	0.99	(26.73, 30.61)
OKLAHOMA	1919	44.00	1.38	(41.30, 46.70)
PENNSYLVANIA	3385	35.46	0.95	(33.60, 37.32)
SOUTH DAKOTA	2076	54.60	1.22	(52.21, 56.99)
VIRGINIA	1882	26.94	1.23	(24.53, 29.35)
WEST VIRGINIA	2400	31.28	1.07	(29.18, 33.38)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 10 MEDIAN = 32.92 RANGE = 22.00-54.60 YEAR 2000 OBJ. 9.12
 * DOES NOT ALWAYS WEAR A SAFETY BELT October 31, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LACK OF SAFETY BELT USAGE* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	801	30.49	2.32	(25.94, 35.04)	*	1126	19.09	1.69	(15.78, 22.40)
ILLINOIS	566	40.18	2.27	(35.73, 44.63)	*	864	31.34	1.83	(27.75, 34.93)
KENTUCKY	656	41.91	2.20	(37.60, 46.22)	*	1120	27.75	1.47	(24.87, 30.63)
LOUISIANA	653	29.73	2.07	(25.67, 33.79)	*	987	15.09	1.21	(12.72, 17.46)
NEW JERSEY	1278	36.50	1.59	(33.38, 39.62)	*	1689	21.35	1.13	(19.14, 23.56)
OKLAHOMA	854	52.72	2.09	(48.62, 56.82)	*	1065	35.99	1.74	(32.58, 39.40)
PENNSYLVANIA	1454	42.93	1.44	(40.11, 45.75)	*	1931	28.71	1.20	(26.36, 31.06)
SOUTH DAKOTA	884	61.58	1.76	(58.13, 65.03)	*	1192	48.08	1.59	(44.96, 51.20)
VIRGINIA	731	32.99	1.97	(29.13, 36.85)	*	1151	21.27	1.50	(18.33, 24.21)
WEST VIRGINIA	890	38.49	1.77	(35.02, 41.96)	*	1510	24.94	1.28	(22.43, 27.45)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	10	NUMBER OF PARTICIPANTS =	10
	MEDIAN	= 39.34	MEDIAN	= 26.35
	RANGE	= 29.73-61.58	RANGE	= 15.09-48.08

* DOES NOT ALWAYS WEAR A SAFETY BELT

October 31, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY BELT*
 DENOMINATOR IS PERSONS WHO HAVE A CHILD BETWEEN THE AGE OF 5 AND 15
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	427	92.50	1.82	(88.93, 96.07)
ILLINOIS	412	85.71	1.91	(81.97, 89.45)
KENTUCKY	445	89.17	1.72	(85.80, 92.54)
LOUISIANA	493	92.96	1.31	(90.39, 95.53)
NEW JERSEY	839	91.05	1.26	(88.58, 93.52)
OKLAHOMA	431	86.85	1.80	(83.32, 90.38)
PENNSYLVANIA	935	89.15	1.12	(86.95, 91.35)
SOUTH DAKOTA	540	77.81	1.99	(73.91, 81.71)
VIRGINIA	498	88.89	1.62	(85.71, 92.07)
WEST VIRGINIA	594	92.91	1.08	(90.79, 95.03)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 10 MEDIAN = 89.16 RANGE = 77.81-92.96
 * OLDEST CHILD UNDER 16 ALWAYS OR NEARLY ALWAYS USE SAFETY BELT December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY BELT* -BY SEX
 DENOMINATOR IS PERSONS WHO HAVE A CHILD BETWEEN THE AGE OF 5 AND 15
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	156	92.38	3.33	(85.85, 98.91)	271	92.58	2.01	(88.64, 96.52)
ILLINOIS	141	87.08	2.91	(81.38, 92.78)	271	84.64	2.45	(79.84, 89.44)
KENTUCKY	139	86.87	3.38	(80.25, 93.49)	306	90.74	1.73	(87.35, 94.13)
LOUISIANA	167	93.85	2.02	(89.89, 97.81)	326	92.27	1.74	(88.86, 95.68)
NEW JERSEY	325	91.34	2.20	(87.03, 95.65)	514	90.80	1.39	(88.08, 93.52)
OKLAHOMA	154	84.23	3.17	(78.02, 90.44)	277	88.90	2.10	(84.78, 93.02)
PENNSYLVANIA	354	90.25	1.68	(86.96, 93.54)	581	88.29	1.47	(85.41, 91.17)
SOUTH DAKOTA	228	77.99	2.91	(72.29, 83.69)	312	77.63	2.60	(72.53, 82.73)
VIRGINIA	173	90.70	2.60	(85.60, 95.80)	325	87.45	2.05	(83.43, 91.47)
WEST VIRGINIA	208	96.10	1.35	(93.45, 98.75)	386	90.36	1.62	(87.18, 93.54)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	10	NUMBER OF PARTICIPANTS =	10
	MEDIAN	= 90.48	MEDIAN	= 89.63
	RANGE	= 77.99-96.10	RANGE	= 77.63-92.58

* OLDEST CHILD UNDER 16 ALWAYS OR NEARLY ALWAYS USE SAFETY BELT

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY SEAT*
 DENOMINATOR IS PERSONS WHO HAVE A CHILD UNDER THE AGE OF 5
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	115	94.68	2.57	(89.64, 99.72)
ILLINOIS	121	94.97	1.82	(91.40, 98.54)
KENTUCKY	124	96.64	1.51	(93.68, 99.60)
LOUISIANA	118	92.10	2.86	(86.49, 97.71)
NEW JERSEY	226	95.36	1.38	(92.66, 98.06)
OKLAHOMA	95	97.79	1.67	(94.52, 100.0)**
PENNSYLVANIA	213	97.32	1.06	(95.24, 99.40)
SOUTH DAKOTA	121	93.61	2.55	(88.61, 98.61)
VIRGINIA	156	99.02	0.76	(97.53, 100.5)
WEST VIRGINIA	164	96.40	1.35	(93.75, 99.05)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 10 MEDIAN = 95.88 RANGE = 92.10-99.02
 * OLDEST CHILD UNDER 5 ALWAYS OR NEARLY ALWAYS USE SAFETY SEAT December 3, 1997
 **upper bound CI value constrained to 100.0

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY SEAT* -BY SEX
 DENOMINATOR IS PERSONS WHO HAVE A CHILD UNDER THE AGE OF 5
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	46	96.09	3.84	(88.56, 100.0)**	*	69	92.99	3.26	(86.60, 99.38)
ILLINOIS	38	97.47	2.51	(92.55, 100.0)**	*	83	93.51	2.53	(88.55, 98.47)
KENTUCKY	49	99.01	1.00	(97.05, 100.0)**	*	75	94.39	2.76	(88.98, 99.80)
LOUISIANA	52	92.18	4.48	(83.40, 100.0)**	*	66	92.02	3.65	(84.87, 99.17)
NEW JERSEY	104	93.08	2.46	(88.26, 97.90)	*	122	97.74	1.21	(95.37, 100.0)**
OKLAHOMA	37	98.53	1.47	(95.65, 100.0)**	*	58	97.23	2.73	(91.88, 100.0)**
PENNSYLVANIA	85	96.28	1.93	(92.50, 100.0)**	*	128	98.20	1.08	(96.08, 100.0)**
SOUTH DAKOTA	52	93.38	3.80	(85.93, 100.0)**	*	69	93.82	3.42	(87.12, 100.0)**
VIRGINIA	56	98.35	1.64	(95.14, 100.0)**	*	100	99.54	0.46	(98.64, 100.0)**
WEST VIRGINIA	61	98.72	1.27	(96.23, 100.0)**	*	103	94.61	2.17	(90.36, 98.86)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 10
 MEDIAN = 96.88
 RANGE = 92.18-99.01

NUMBER OF PARTICIPANTS = 10
 MEDIAN = 94.50
 RANGE = 92.02-99.54

* OLDEST CHILD UNDER 5 ALWAYS OR NEARLY ALWAYS USE SAFETY SEAT
 **upper bound CI value constrained to 100.0

December 3, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHILD BICYCLE HELMET USE*
 DENOMINATOR IS RESPONDENTS WITH CHILDREN AGE 5-15 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	426	33.20	2.94	(27.44, 38.96)
ILLINOIS	413	18.33	2.18	(14.06, 22.60)
KENTUCKY	445	20.83	2.21	(16.50, 25.16)
LOUISIANA	494	11.51	1.67	(8.24, 14.78)
NEW JERSEY	844	56.80	1.95	(52.98, 60.62)
OKLAHOMA	430	17.94	2.11	(13.80, 22.08)
PENNSYLVANIA	935	46.83	1.83	(43.24, 50.42)
SOUTH DAKOTA	538	12.57	1.42	(9.79, 15.35)
VIRGINIA	494	33.19	2.40	(28.49, 37.89)
WEST VIRGINIA	590	29.17	2.02	(25.21, 33.13)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 10 MEDIAN = 25.00 RANGE = 11.51-56.80
 * OLDEST CHILD AGE 5-15 YEARS ALWAYS WEARS A BICYCLE HELMET

December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHILD BICYCLE HELMET USE* -BY SEX
 DENOMINATOR IS RESPONDENTS WITH CHILDREN AGE 5-15 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	158	37.36	4.96	(27.64, 47.08)	*	268	30.20	3.47	(23.40, 37.00)
ILLINOIS	142	23.46	3.80	(16.01, 30.91)	*	271	14.29	2.33	(9.72, 18.86)
KENTUCKY	140	21.66	3.65	(14.51, 28.81)	*	305	20.25	2.49	(15.37, 25.13)
LOUISIANA	168	9.91	2.57	(4.87, 14.95)	*	326	12.77	2.01	(8.83, 16.71)
NEW JERSEY	325	59.43	3.12	(53.31, 65.55)	*	519	54.65	2.47	(49.81, 59.49)
OKLAHOMA	154	19.03	3.38	(12.41, 25.65)	*	276	17.09	2.66	(11.88, 22.30)
PENNSYLVANIA	356	44.23	2.85	(38.64, 49.82)	*	579	48.87	2.35	(44.26, 53.48)
SOUTH DAKOTA	224	8.85	1.79	(5.34, 12.36)	*	314	16.13	2.17	(11.88, 20.38)
VIRGINIA	172	35.32	4.01	(27.46, 43.18)	*	322	31.50	2.89	(25.84, 37.16)
WEST VIRGINIA	207	28.39	3.07	(22.37, 34.41)	*	383	29.78	2.61	(24.66, 34.90)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	10	NUMBER OF PARTICIPANTS =	10
	MEDIAN	= 25.92	MEDIAN	= 25.01
	RANGE	= 8.85-59.43	RANGE	= 12.77-54.65
* OLDEST CHILD 5-15 YEARS ALWAYS WEARS A BICYCLE HELMET			December 2, 1997	

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT DRINKERS*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2075	40.75	1.27	(38.26, 43.24)
ALASKA	1505	63.20	1.87	(59.53, 66.87)
ARIZONA	1937	43.58	1.57	(40.50, 46.66)
CALIFORNIA	4001	59.06	0.90	(57.30, 60.82)
IDAHO	2730	48.78	1.14	(46.55, 51.01)
ILLINOIS	1440	57.76	1.52	(54.78, 60.74)
KENTUCKY	1804	32.66	1.36	(29.99, 35.33)
MINNESOTA	4391	56.92	0.85	(55.25, 58.59)
NEVADA	1795	62.56	1.59	(59.44, 65.68)
NEW JERSEY	2992	62.23	1.02	(60.23, 64.23)
NEW YORK	4308	53.44	0.87	(51.73, 55.15)
OKLAHOMA	1920	31.04	1.36	(28.37, 33.71)
PENNSYLVANIA	3406	55.65	0.95	(53.79, 57.51)
TENNESSEE	1498	30.14	1.37	(27.45, 32.83)
VERMONT	2414	61.64	1.17	(59.35, 63.93)
WISCONSIN	2228	70.34	1.27	(67.85, 72.83)
WYOMING	2529	60.28	1.11	(58.10, 62.46)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 17 MEDIAN = 56.92 RANGE = 30.14-70.34
 * HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT DRINKERS* -BY SEX
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	758	53.03	2.05	(49.01, 57.05)	*	1317	29.96	1.42	(27.18, 32.74)
ALASKA	698	68.64	2.62	(63.50, 73.78)	*	807	57.21	2.62	(52.07, 62.35)
ARIZONA	806	52.01	2.37	(47.36, 56.66)	*	1131	35.54	1.86	(31.89, 39.19)
CALIFORNIA	1728	66.98	1.31	(64.41, 69.55)	*	2273	51.30	1.18	(48.99, 53.61)
IDAHO	1118	57.10	1.72	(53.73, 60.47)	*	1612	40.80	1.38	(38.10, 43.50)
ILLINOIS	568	68.97	2.09	(64.87, 73.07)	*	872	48.11	1.95	(44.29, 51.93)
KENTUCKY	648	43.10	2.24	(38.71, 47.49)	*	1156	23.49	1.39	(20.77, 26.21)
MINNESOTA	1950	61.43	1.23	(59.02, 63.84)	*	2441	52.72	1.10	(50.56, 54.88)
NEVADA	760	68.21	2.29	(63.72, 72.70)	*	1035	56.80	2.16	(52.57, 61.03)
NEW JERSEY	1286	69.73	1.49	(66.81, 72.65)	*	1706	55.27	1.39	(52.55, 57.99)
NEW YORK	1747	61.82	1.32	(59.23, 64.41)	*	2561	45.98	1.11	(43.80, 48.16)
OKLAHOMA	852	37.98	2.18	(33.71, 42.25)	*	1068	24.70	1.61	(21.54, 27.86)
PENNSYLVANIA	1462	65.85	1.38	(63.15, 68.55)	*	1944	46.43	1.28	(43.92, 48.94)
TENNESSEE	583	37.03	2.23	(32.66, 41.40)	*	915	23.80	1.61	(20.64, 26.96)
VERMONT	1034	68.06	1.70	(64.73, 71.39)	*	1380	55.64	1.52	(52.66, 58.62)
WISCONSIN	945	76.74	1.82	(73.17, 80.31)	*	1283	64.41	1.77	(60.94, 67.88)
WYOMING	1050	65.66	1.56	(62.60, 68.72)	*	1479	55.02	1.45	(52.18, 57.86)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	17	NUMBER OF PARTICIPANTS =	17
	MEDIAN	= 65.66	MEDIAN	= 48.11
	RANGE	= 37.03-76.74	RANGE	= 23.49-64.41

* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH
 December 2, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2056	13.67	0.93	(11.85, 15.49)
ALASKA	1484	19.59	1.60	(16.45, 22.73)
ARIZONA	1913	13.85	1.17	(11.56, 16.14)
CALIFORNIA	3990	14.61	0.65	(13.34, 15.88)
IDAHO	2710	14.65	0.81	(13.06, 16.24)
ILLINOIS	1435	18.19	1.22	(15.80, 20.58)
KENTUCKY	1789	8.77	0.88	(7.05, 10.49)
MINNESOTA	4367	16.11	0.65	(14.84, 17.38)
NEVADA	1777	20.67	1.54	(17.65, 23.69)
NEW JERSEY	2962	14.91	0.79	(13.36, 16.46)
NEW YORK	4275	12.79	0.59	(11.63, 13.95)
OKLAHOMA	1910	8.54	0.83	(6.91, 10.17)
PENNSYLVANIA	3361	16.97	0.75	(15.50, 18.44)
TENNESSEE	1482	7.23	0.73	(5.80, 8.66)
VERMONT	2399	18.39	1.14	(16.16, 20.62)
WISCONSIN	2220	25.95	1.22	(23.56, 28.34)
WYOMING	2507	19.24	0.89	(17.50, 20.98)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 17 MEDIAN = 14.91 RANGE = 7.23-25.95
 * 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH November 26, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKING* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	742	23.09	1.71	(19.74, 26.44)	*	1314	5.54	0.69	(4.19, 6.89)
ALASKA	683	28.12	2.59	(23.04, 33.20)	*	801	10.29	1.62	(7.11, 13.47)
ARIZONA	792	20.62	2.12	(16.46, 24.78)	*	1121	7.42	1.00	(5.46, 9.38)
CALIFORNIA	1724	22.50	1.14	(20.27, 24.73)	*	2266	6.86	0.58	(5.72, 8.00)
IDAHO	1103	22.61	1.43	(19.81, 25.41)	*	1607	7.10	0.70	(5.73, 8.47)
ILLINOIS	565	30.19	2.15	(25.98, 34.40)	*	870	7.90	1.04	(5.86, 9.94)
KENTUCKY	640	14.84	1.70	(11.51, 18.17)	*	1149	3.45	0.66	(2.16, 4.74)
MINNESOTA	1938	22.65	1.06	(20.57, 24.73)	*	2429	10.01	0.70	(8.64, 11.38)
NEVADA	748	28.28	2.50	(23.38, 33.18)	*	1029	12.97	1.65	(9.74, 16.20)
NEW JERSEY	1269	23.67	1.40	(20.93, 26.41)	*	1693	6.84	0.70	(5.47, 8.21)
NEW YORK	1727	20.07	1.06	(17.99, 22.15)	*	2548	6.35	0.53	(5.31, 7.39)
OKLAHOMA	846	13.17	1.49	(10.25, 16.09)	*	1064	4.32	0.81	(2.73, 5.91)
PENNSYLVANIA	1435	25.66	1.31	(23.09, 28.23)	*	1926	9.20	0.70	(7.83, 10.57)
TENNESSEE	574	11.45	1.39	(8.73, 14.17)	*	908	3.37	0.61	(2.17, 4.57)
VERMONT	1025	27.83	1.99	(23.93, 31.73)	*	1374	9.60	0.93	(7.78, 11.42)
WISCONSIN	939	38.02	2.00	(34.10, 41.94)	*	1281	14.80	1.32	(12.21, 17.39)
WYOMING	1038	28.09	1.51	(25.13, 31.05)	*	1469	10.61	0.86	(8.92, 12.30)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	17	NUMBER OF PARTICIPANTS =	17
	MEDIAN	= 23.09	MEDIAN	= 7.42
	RANGE	= 11.45-38.02	RANGE	= 3.37-14.80

* 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH
 November 26, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHRONIC DRINKING*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2048	4.08	0.54	(3.02, 5.14)
ALASKA	1479	3.78	0.68	(2.45, 5.11)
ARIZONA	1901	5.03	0.80	(3.46, 6.60)
IDAHO	2702	3.31	0.46	(2.41, 4.21)
ILLINOIS	1432	3.45	0.54	(2.39, 4.51)
KENTUCKY	1783	2.75	0.45	(1.87, 3.63)
MINNESOTA	4350	2.99	0.31	(2.38, 3.60)
NEVADA	1762	7.14	1.04	(5.10, 9.18)
NEW JERSEY	2931	3.20	0.38	(2.46, 3.94)
NEW YORK	4245	2.75	0.29	(2.18, 3.32)
OKLAHOMA	1906	2.49	0.49	(1.53, 3.45)
PENNSYLVANIA	3329	3.33	0.40	(2.55, 4.11)
TENNESSEE	1466	1.35	0.32	(0.72, 1.98)
VERMONT	2377	4.32	0.94	(2.48, 6.16)
WISCONSIN	2206	6.46	0.69	(5.11, 7.81)
WYOMING	2499	4.22	0.46	(3.32, 5.12)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 16 MEDIAN = 3.39 RANGE = 1.35- 7.14

* 60 OR MORE DRINKS DURING PAST MONTH

December 3, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHRONIC DRINKING* -BY SEX
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	743	8.02	1.11	(5.84, 10.20)	1305	0.64	0.23	(0.19, 1.09)
ALASKA	683	6.82	1.28	(4.31, 9.33)	796	0.46	0.23	(0.01, 0.91)
ARIZONA	787	8.22	1.52	(5.24, 11.20)	1114	1.98	0.59	(0.82, 3.14)
IDAHO	1101	6.22	0.90	(4.46, 7.98)	1601	0.53	0.21	(0.12, 0.94)
ILLINOIS	564	6.64	1.12	(4.44, 8.84)	868	0.71	0.31	(0.10, 1.32)
KENTUCKY	637	4.86	0.89	(3.12, 6.60)	1146	0.92	0.31	(0.31, 1.53)
MINNESOTA	1930	5.41	0.61	(4.21, 6.61)	2420	0.73	0.21	(0.32, 1.14)
NEVADA	742	10.97	1.69	(7.66, 14.28)	1020	3.24	1.18	(0.93, 5.55)
NEW JERSEY	1254	5.99	0.76	(4.50, 7.48)	1677	0.62	0.18	(0.27, 0.97)
NEW YORK	1712	4.13	0.53	(3.09, 5.17)	2533	1.54	0.28	(0.99, 2.09)
OKLAHOMA	847	4.03	0.92	(2.23, 5.83)	1059	1.06	0.39	(0.30, 1.82)
PENNSYLVANIA	1419	5.87	0.80	(4.30, 7.44)	1910	1.06	0.24	(0.59, 1.53)
TENNESSEE	567	2.43	0.64	(1.18, 3.68)	899	0.36	0.19	(-0.01, 0.73)
VERMONT	1018	8.16	1.87	(4.49, 11.83)	1359	0.73	0.21	(0.32, 1.14)
WISCONSIN	936	12.28	1.36	(9.61, 14.95)	1270	1.08	0.29	(0.51, 1.65)
WYOMING	1033	7.67	0.88	(5.95, 9.39)	1466	0.87	0.25	(0.38, 1.36)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	16	NUMBER OF PARTICIPANTS =	16
	MEDIAN	= 6.43	MEDIAN	= 0.80
	RANGE	= 2.43-12.28	RANGE	= 0.36- 3.24

* 60 OR MORE DRINKS DURING PAST MONTH

December 3, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DRINKING AND DRIVING*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2068	3.04	0.47	(2.12, 3.96)
ALASKA	1498	1.71	0.39	(0.95, 2.47)
ARIZONA	1925	3.24	0.69	(1.89, 4.59)
CALIFORNIA	3998	2.53	0.29	(1.96, 3.10)
IDAHO	2726	2.38	0.39	(1.62, 3.14)
ILLINOIS	1440	2.38	0.45	(1.50, 3.26)
KENTUCKY	1799	0.43	0.16	(0.12, 0.74)
MINNESOTA	4372	3.35	0.31	(2.74, 3.96)
NEVADA	1792	3.01	0.60	(1.83, 4.19)
NEW JERSEY	2981	2.23	0.35	(1.54, 2.92)
NEW YORK	4296	1.61	0.22	(1.18, 2.04)
OKLAHOMA	1916	1.56	0.36	(0.85, 2.27)
PENNSYLVANIA	3381	2.91	0.38	(2.17, 3.65)
TENNESSEE	1490	1.39	0.36	(0.68, 2.10)
VERMONT	2408	3.06	0.41	(2.26, 3.86)
WISCONSIN	2227	5.55	0.61	(4.35, 6.75)
WYOMING	2518	3.72	0.43	(2.88, 4.56)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 17 MEDIAN = 2.53 RANGE = 0.43- 5.55
 * DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES November 26, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DRINKING AND DRIVING* -BY SEX
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	752	5.29	0.93	(3.47, 7.11)	*	1316	1.06	0.28	(0.51, 1.61)
ALASKA	692	2.90	0.73	(1.47, 4.33)	*	806	0.41	0.17	(0.08, 0.74)
ARIZONA	800	5.84	1.36	(3.17, 8.51)	*	1125	0.76	0.31	(0.15, 1.37)
CALIFORNIA	1726	3.69	0.49	(2.73, 4.65)	*	2272	1.40	0.29	(0.83, 1.97)
IDAHO	1116	4.46	0.76	(2.97, 5.95)	*	1610	0.38	0.18	(0.03, 0.73)
ILLINOIS	568	3.88	0.86	(2.19, 5.57)	*	872	1.08	0.33	(0.43, 1.73)
KENTUCKY	645	0.66	0.30	(0.07, 1.25)	*	1154	0.23	0.13	(-0.02, 0.48)
MINNESOTA	1940	5.40	0.58	(4.26, 6.54)	*	2432	1.45	0.27	(0.92, 1.98)
NEVADA	759	3.70	0.97	(1.80, 5.60)	*	1033	2.31	0.70	(0.94, 3.68)
NEW JERSEY	1278	3.56	0.64	(2.31, 4.81)	*	1703	0.99	0.32	(0.36, 1.62)
NEW YORK	1739	3.09	0.45	(2.21, 3.97)	*	2557	0.29	0.12	(0.05, 0.53)
OKLAHOMA	848	2.62	0.70	(1.25, 3.99)	*	1068	0.59	0.26	(0.08, 1.10)
PENNSYLVANIA	1448	5.03	0.76	(3.54, 6.52)	*	1933	0.99	0.22	(0.56, 1.42)
TENNESSEE	579	2.20	0.61	(1.00, 3.40)	*	911	0.64	0.39	(-0.12, 1.40)
VERMONT	1032	5.02	0.79	(3.47, 6.57)	*	1376	1.22	0.29	(0.65, 1.79)
WISCONSIN	944	9.02	1.14	(6.79, 11.25)	*	1283	2.33	0.50	(1.35, 3.31)
WYOMING	1044	5.44	0.77	(3.93, 6.95)	*	1474	2.04	0.36	(1.33, 2.75)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	17	NUMBER OF PARTICIPANTS =	17
	MEDIAN	= 3.88	MEDIAN	= 0.99
	RANGE	= 0.66- 9.02	RANGE	= 0.23- 2.33

* DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES
 November 26, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM - FEMALES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1341	60.17	1.47	(57.29, 63.05)
ALASKA	814	56.89	2.64	(51.72, 62.06)
ARIZONA	1135	60.86	2.10	(56.74, 64.98)
ARKANSAS	1149	53.99	1.72	(50.62, 57.36)
COLORADO	988	58.49	1.85	(54.86, 62.12)
CONNECTICUT	1072	65.63	1.67	(62.36, 68.90)
DELAWARE	1296	59.29	1.62	(56.11, 62.47)
DISTRICT OF CO	736	58.60	2.13	(54.43, 62.77)
FLORIDA	2124	64.63	1.13	(62.42, 66.84)
GEORGIA	1357	60.14	1.50	(57.20, 63.08)
IDAHO	1615	53.79	1.40	(51.05, 56.53)
ILLINOIS	1724	57.02	1.31	(54.45, 59.59)
INDIANA	1263	55.70	1.53	(52.70, 58.70)
IOWA	2098	57.01	1.20	(54.66, 59.36)
KANSAS	1117	56.12	1.61	(52.96, 59.28)
KENTUCKY	2269	57.18	1.22	(54.79, 59.57)
LOUISIANA	991	55.76	1.73	(52.37, 59.15)
MAINE	984	59.83	1.77	(56.36, 63.30)
MARYLAND	2610	61.93	1.26	(59.46, 64.40)
MASSACHUSETTS	1035	59.58	1.70	(56.25, 62.91)
MICHIGAN	1439	61.73	1.42	(58.95, 64.51)
MINNESOTA	2441	55.33	1.12	(53.13, 57.53)
MISSISSIPPI	996	54.44	1.72	(51.07, 57.81)
MISSOURI	924	58.17	1.84	(54.56, 61.78)
MONTANA	1029	60.48	1.67	(57.21, 63.75)
NEBRASKA	1255	53.70	1.72	(50.33, 57.07)
NEVADA	1039	59.72	2.18	(55.45, 63.99)
NEW HAMPSHIRE	834	57.08	1.96	(53.24, 60.92)
NEW JERSEY	1770	58.64	1.38	(55.94, 61.34)
NEW MEXICO	658	53.32	2.47	(48.48, 58.16)
NEW YORK	2561	60.23	1.11	(58.05, 62.41)
NORTH CAROLINA	1643	57.56	1.43	(54.76, 60.36)
NORTH DAKOTA	1063	56.26	1.73	(52.87, 59.65)
OHIO	925	57.35	1.83	(53.76, 60.94)
OKLAHOMA	1070	52.74	1.74	(49.33, 56.15)
OREGON	1695	61.97	1.30	(59.42, 64.52)
PENNSYLVANIA	2020	58.99	1.27	(56.50, 61.48)
RHODE ISLAND	1049	62.65	1.72	(59.28, 66.02)
SOUTH CAROLINA	1109	57.83	1.84	(54.22, 61.44)
SOUTH DAKOTA	1204	55.77	1.56	(52.71, 58.83)
TENNESSEE	1862	56.61	1.30	(54.06, 59.16)
TEXAS	1035	52.31	1.69	(49.00, 55.62)
UTAH	1627	49.81	1.65	(46.58, 53.04)
VERMONT	1418	55.70	1.50	(52.76, 58.64)
VIRGINIA	1158	58.95	1.69	(55.64, 62.26)
WASHINGTON	2061	58.26	1.22	(55.87, 60.65)
WEST VIRGINIA	1514	58.57	1.42	(55.79, 61.35)
WISCONSIN	1278	62.23	1.84	(58.62, 65.84)
WYOMING	1481	54.55	1.42	(51.77, 57.33)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 57.83 RANGE = 49.81-65.63
 * EVER HAD A MAMMOGRAM, AGE 18 AND OLDER November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 MAMMOGRAM - FEMALES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	57.80	60.17	2.37
ALASKA	50.72	56.89	6.17
ARIZONA	61.28	60.86	-0.42
ARKANSAS	53.52	53.99	0.47
COLORADO	55.13	58.49	3.36
CONNECTICUT	62.05	65.63	3.58
DELAWARE	58.71	59.29	0.58
DISTRICT OF CO	N/A	58.60	N/A
FLORIDA	62.29	64.63	2.34
GEORGIA	57.40	60.14	2.74
IDAHO	53.10	53.79	0.69
ILLINOIS	56.51	57.02	0.51
INDIANA	55.63	55.70	0.07
IOWA	56.44	57.01	0.57
KANSAS	56.72	56.12	-0.60
KENTUCKY	56.88	57.18	0.30
LOUISIANA	54.33	55.76	1.43
MAINE	56.91	59.83	2.92
MARYLAND	59.57	61.93	2.36
MASSACHUSETTS	61.32	59.58	-1.74
MICHIGAN	59.85	61.73	1.88
MINNESOTA	53.86	55.33	1.47
MISSISSIPPI	53.11	54.44	1.33
MISSOURI	57.57	58.17	0.60
MONTANA	55.10	60.48	5.38
NEBRASKA	52.22	53.70	1.48
NEVADA	55.96	59.72	3.76
NEW HAMPSHIRE	55.27	57.08	1.81
NEW JERSEY	55.58	58.64	3.06
NEW MEXICO	54.43	53.32	-1.11
NEW YORK	57.88	60.23	2.35
NORTH CAROLINA	56.22	57.56	1.34
NORTH DAKOTA	55.90	56.26	0.36
OHIO	57.76	57.35	-0.41
OKLAHOMA	53.31	52.74	-0.57
OREGON	60.56	61.97	1.41
PENNSYLVANIA	58.93	58.99	0.06
RHODE ISLAND	58.09	62.65	4.56
SOUTH CAROLINA	62.63	57.83	-4.80
SOUTH DAKOTA	51.71	55.77	4.06
TENNESSEE	56.46	56.61	0.15
TEXAS	52.60	52.31	-0.29
UTAH	50.46	49.81	-0.65
VERMONT	53.34	55.70	2.36
VIRGINIA	55.97	58.95	2.98
WASHINGTON	57.53	58.26	0.73
WEST VIRGINIA	56.02	58.57	2.55
WISCONSIN	55.57	62.23	6.66
WYOMING	54.16	54.55	0.39

* EVER HAD A MAMMOGRAM, AGE 18 AND OLDER
 N/A - NOT AVAILABLE

November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	763	82.88	1.42	(80.10, 85.66)
ALASKA	449	91.66	1.76	(88.21, 95.11)
ARIZONA	704	84.39	2.00	(80.47, 88.31)
ARKANSAS	744	75.68	1.72	(72.31, 79.05)
COLORADO	583	87.83	1.46	(84.97, 90.69)
CONNECTICUT	648	91.23	1.19	(88.90, 93.56)
DELAWARE	754	86.61	1.41	(83.85, 89.37)
DISTRICT OF CO	398	87.59	1.83	(84.00, 91.18)
FLORIDA	1308	86.99	1.00	(85.03, 88.95)
GEORGIA	763	88.23	1.23	(85.82, 90.64)
IDAHO	929	79.11	1.50	(76.17, 82.05)
ILLINOIS	975	83.59	1.29	(81.06, 86.12)
INDIANA	793	79.25	1.62	(76.07, 82.43)
IOWA	1325	80.66	1.22	(78.27, 83.05)
KANSAS	679	78.99	1.67	(75.72, 82.26)
KENTUCKY	1438	79.26	1.22	(76.87, 81.65)
LOUISIANA	582	76.09	1.93	(72.31, 79.87)
MAINE	634	87.30	1.40	(84.56, 90.04)
MARYLAND	1537	89.11	1.05	(87.05, 91.17)
MASSACHUSETTS	575	89.54	1.37	(86.85, 92.23)
MICHIGAN	838	88.65	1.20	(86.30, 91.00)
MINNESOTA	1438	84.32	1.04	(82.28, 86.36)
MISSISSIPPI	616	76.32	1.87	(72.65, 79.99)
MISSOURI	577	80.95	1.75	(77.52, 84.38)
MONTANA	642	85.91	1.43	(83.11, 88.71)
NEBRASKA	804	78.40	1.59	(75.28, 81.52)
NEVADA	644	83.95	2.10	(79.83, 88.07)
NEW HAMPSHIRE	482	85.94	1.66	(82.69, 89.19)
NEW JERSEY	1051	81.86	1.38	(79.16, 84.56)
NEW MEXICO	421	81.25	2.29	(76.76, 85.74)
NEW YORK	1461	85.58	0.97	(83.68, 87.48)
NORTH CAROLINA	1007	82.01	1.35	(79.36, 84.66)
NORTH DAKOTA	655	84.27	1.54	(81.25, 87.29)
OHIO	583	81.27	1.76	(77.82, 84.72)
OKLAHOMA	710	71.79	2.00	(67.87, 75.71)
OREGON	1021	87.23	1.17	(84.94, 89.52)
PENNSYLVANIA	1246	82.19	1.18	(79.88, 84.50)
RHODE ISLAND	639	88.61	1.35	(85.96, 91.26)
SOUTH CAROLINA	689	84.07	1.72	(80.70, 87.44)
SOUTH DAKOTA	753	81.52	1.53	(78.52, 84.52)
TENNESSEE	1140	80.10	1.26	(77.63, 82.57)
TEXAS	588	77.25	1.90	(73.53, 80.97)
UTAH	895	82.75	1.52	(79.77, 85.73)
VERMONT	891	85.74	1.24	(83.31, 88.17)
VIRGINIA	668	85.57	1.47	(82.69, 88.45)
WASHINGTON	1176	86.35	1.08	(84.23, 88.47)
WEST VIRGINIA	996	80.59	1.35	(77.94, 83.24)
WISCONSIN	749	89.22	1.39	(86.50, 91.94)
WYOMING	874	78.93	1.51	(75.97, 81.89)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 83.95 RANGE = 71.79-91.66
 * EVER HAD A MAMMOGRAM, AGE 40 AND OVER November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 MAMMOGRAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	79.01	82.88	3.87
ALASKA	87.09	91.66	4.57
ARIZONA	86.29	84.39	-1.90
ARKANSAS	73.68	75.68	2.00
COLORADO	84.46	87.83	3.37
CONNECTICUT	87.08	91.23	4.15
DELAWARE	85.56	86.61	1.05
DISTRICT OF CO	N/A	87.59	N/A
FLORIDA	84.43	86.99	2.56
GEORGIA	82.82	88.23	5.41
IDAHO	80.30	79.11	-1.19
ILLINOIS	81.24	83.59	2.35
INDIANA	79.72	79.25	-0.47
IOWA	81.17	80.66	-0.51
KANSAS	80.06	78.99	-1.07
KENTUCKY	80.51	79.26	-1.25
LOUISIANA	74.03	76.09	2.06
MAINE	82.09	87.30	5.21
MARYLAND	86.01	89.11	3.10
MASSACHUSETTS	90.89	89.54	-1.35
MICHIGAN	87.71	88.65	0.94
MINNESOTA	81.81	84.32	2.51
MISSISSIPPI	72.97	76.32	3.35
MISSOURI	81.25	80.95	-0.30
MONTANA	78.29	85.91	7.62
NEBRASKA	76.14	78.40	2.26
NEVADA	82.52	83.95	1.43
NEW HAMPSHIRE	86.32	85.94	-0.38
NEW JERSEY	78.43	81.86	3.43
NEW MEXICO	82.23	81.25	-0.98
NEW YORK	84.36	85.58	1.22
NORTH CAROLINA	81.21	82.01	0.80
NORTH DAKOTA	81.39	84.27	2.88
OHIO	82.24	81.27	-0.97
OKLAHOMA	74.34	71.79	-2.55
OREGON	85.99	87.23	1.24
PENNSYLVANIA	82.79	82.19	-0.60
RHODE ISLAND	86.78	88.61	1.83
SOUTH CAROLINA	82.41	84.07	1.66
SOUTH DAKOTA	76.54	81.52	4.98
TENNESSEE	79.67	80.10	0.43
TEXAS	78.70	77.25	-1.45
UTAH	82.00	82.75	0.75
VERMONT	82.78	85.74	2.96
VIRGINIA	83.58	85.57	1.99
WASHINGTON	86.26	86.35	0.09
WEST VIRGINIA	77.80	80.59	2.79
WISCONSIN	82.44	89.22	6.78
WYOMING	79.09	78.93	-0.16

* EVER HAD A MAMMOGRAM, AGE 40 AND OVER
 N/A - NOT AVAILABLE

November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	502	70.17	2.13	(66.00, 74.34)
ALASKA	219	80.06	4.10	(72.02, 88.10)
ARIZONA	516	79.32	2.09	(75.22, 83.42)
ARKANSAS	529	59.86	2.38	(55.20, 64.52)
COLORADO	358	78.50	2.53	(73.54, 83.46)
CONNECTICUT	397	79.89	2.21	(75.56, 84.22)
DELAWARE	499	79.19	1.98	(75.31, 83.07)
DISTRICT OF CO	259	82.57	2.48	(77.71, 87.43)
FLORIDA	937	80.40	1.39	(77.68, 83.12)
GEORGIA	475	79.75	2.00	(75.83, 83.67)
IDAHO	623	66.35	2.02	(62.39, 70.31)
ILLINOIS	649	69.04	2.05	(65.02, 73.06)
INDIANA	518	64.01	2.38	(59.35, 68.67)
IOWA	947	62.11	1.77	(58.64, 65.58)
KANSAS	447	71.51	2.26	(67.08, 75.94)
KENTUCKY	1044	63.96	1.66	(60.71, 67.21)
LOUISIANA	393	63.71	2.71	(58.40, 69.02)
MAINE	433	76.84	2.18	(72.57, 81.11)
MARYLAND	993	82.15	1.56	(79.09, 85.21)
MASSACHUSETTS	370	80.97	2.22	(76.62, 85.32)
MICHIGAN	529	75.48	2.04	(71.48, 79.48)
MINNESOTA	948	69.92	1.59	(66.80, 73.04)
MISSISSIPPI	427	60.88	2.56	(55.86, 65.90)
MISSOURI	404	65.52	2.60	(60.42, 70.62)
MONTANA	423	68.84	2.47	(64.00, 73.68)
NEBRASKA	574	66.11	2.14	(61.92, 70.30)
NEVADA	402	68.05	3.29	(61.60, 74.50)
NEW HAMPSHIRE	293	74.93	2.63	(69.78, 80.08)
NEW JERSEY	662	69.35	2.04	(65.35, 73.35)
NEW MEXICO	274	67.79	3.50	(60.93, 74.65)
NEW YORK	926	76.67	1.47	(73.79, 79.55)
NORTH CAROLINA	721	72.49	1.84	(68.88, 76.10)
NORTH DAKOTA	468	69.88	2.25	(65.47, 74.29)
OHIO	416	70.80	2.45	(66.00, 75.60)
OKLAHOMA	527	60.54	2.47	(55.70, 65.38)
OREGON	669	78.48	1.73	(75.09, 81.87)
PENNSYLVANIA	831	69.99	1.76	(66.54, 73.44)
RHODE ISLAND	433	77.27	2.22	(72.92, 81.62)
SOUTH CAROLINA	453	75.93	2.32	(71.38, 80.48)
SOUTH DAKOTA	538	67.44	2.15	(63.23, 71.65)
TENNESSEE	761	68.87	1.77	(65.40, 72.34)
TEXAS	361	67.88	2.66	(62.67, 73.09)
UTAH	586	69.69	2.30	(65.18, 74.20)
VERMONT	563	74.66	2.03	(70.68, 78.64)
VIRGINIA	418	72.50	2.36	(67.87, 77.13)
WASHINGTON	719	79.16	1.61	(76.00, 82.32)
WEST VIRGINIA	681	70.35	1.98	(66.47, 74.23)
WISCONSIN	482	74.02	2.48	(69.16, 78.88)
WYOMING	544	64.04	2.19	(59.75, 68.33)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 70.35 RANGE = 59.86-82.57

* HAD A MAMMOGRAM WITHIN TWO YEARS AGE 50 AND OVER

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 MAMMOGRAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	67.66	70.17	2.51
ALASKA	77.71	80.06	2.35
ARIZONA	78.81	79.32	0.51
ARKANSAS	63.18	59.86	-3.32
COLORADO	69.95	78.50	8.55
CONNECTICUT	78.01	79.89	1.88
DELAWARE	74.09	79.19	5.10
DISTRICT OF CO	N/A	82.57	N/A
FLORIDA	77.27	80.40	3.13
GEORGIA	75.77	79.75	3.98
IDAHO	66.91	66.35	-0.56
ILLINOIS	72.93	69.04	-3.89
INDIANA	67.41	64.01	-3.40
IOWA	67.06	62.11	-4.95
KANSAS	68.64	71.51	2.87
KENTUCKY	65.27	63.96	-1.31
LOUISIANA	62.99	63.71	0.72
MAINE	71.87	76.84	4.97
MARYLAND	79.75	82.15	2.40
MASSACHUSETTS	82.01	80.97	-1.04
MICHIGAN	79.80	75.48	-4.32
MINNESOTA	72.55	69.92	-2.63
MISSISSIPPI	55.09	60.88	5.79
MISSOURI	69.86	65.52	-4.34
MONTANA	63.78	68.84	5.06
NEBRASKA	65.79	66.11	0.32
NEVADA	70.31	68.05	-2.26
NEW HAMPSHIRE	77.93	74.93	-3.00
NEW JERSEY	63.27	69.35	6.08
NEW MEXICO	72.09	67.79	-4.30
NEW YORK	73.90	76.67	2.77
NORTH CAROLINA	67.94	72.49	4.55
NORTH DAKOTA	69.71	69.88	0.17
OHIO	68.93	70.80	1.87
OKLAHOMA	62.10	60.54	-1.56
OREGON	77.56	78.48	0.92
PENNSYLVANIA	64.13	69.99	5.86
RHODE ISLAND	71.41	77.27	5.86
SOUTH CAROLINA	73.07	75.93	2.86
SOUTH DAKOTA	64.40	67.44	3.04
TENNESSEE	67.80	68.87	1.07
TEXAS	70.34	67.88	-2.46
UTAH	67.88	69.69	1.81
VERMONT	70.33	74.66	4.33
VIRGINIA	71.96	72.50	0.54
WASHINGTON	76.85	79.16	2.31
WEST VIRGINIA	65.35	70.35	5.00
WISCONSIN	63.85	74.02	10.17
WYOMING	62.43	64.04	1.61

* HAD A MAMMOGRAM WITHIN TWO YEARS AGE 50 AND OVER
 N/A - NOT AVAILABLE

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BREAST EXAM - FEMALES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1335	86.70	1.10	(84.54, 88.86)
ALASKA	812	93.83	1.13	(91.62, 96.04)
ARIZONA	1131	87.40	1.39	(84.68, 90.12)
ARKANSAS	1146	84.85	1.16	(82.58, 87.12)
CALIFORNIA	2270	86.04	0.90	(84.28, 87.80)
COLORADO	986	92.06	0.96	(90.18, 93.94)
CONNECTICUT	1073	87.57	1.15	(85.32, 89.82)
DELAWARE	1293	92.96	0.88	(91.24, 94.68)
DISTRICT OF CO	735	90.57	1.37	(87.88, 93.26)
FLORIDA	2116	88.40	0.82	(86.79, 90.01)
GEORGIA	1356	92.51	0.82	(90.90, 94.12)
IDAHO	1608	91.33	0.79	(89.78, 92.88)
ILLINOIS	1721	89.02	0.88	(87.30, 90.74)
INDIANA	1260	89.52	0.96	(87.64, 91.40)
IOWA	2087	91.70	0.69	(90.35, 93.05)
KANSAS	1108	91.22	0.96	(89.34, 93.10)
KENTUCKY	2265	86.97	0.83	(85.34, 88.60)
LOUISIANA	992	85.73	1.28	(83.22, 88.24)
MAINE	983	91.99	1.10	(89.83, 94.15)
MARYLAND	2592	94.15	0.63	(92.92, 95.38)
MASSACHUSETTS	1034	92.82	0.95	(90.96, 94.68)
MICHIGAN	1437	89.93	0.92	(88.13, 91.73)
MINNESOTA	2429	92.03	0.61	(90.83, 93.23)
MISSISSIPPI	994	87.70	1.19	(85.37, 90.03)
MISSOURI	921	89.43	1.04	(87.39, 91.47)
MONTANA	1028	94.41	0.83	(92.78, 96.04)
NEBRASKA	1251	92.34	0.78	(90.81, 93.87)
NEVADA	1038	89.72	1.30	(87.17, 92.27)
NEW HAMPSHIRE	827	90.70	1.19	(88.37, 93.03)
NEW JERSEY	1757	83.86	1.04	(81.82, 85.90)
NEW MEXICO	656	89.03	1.58	(85.93, 92.13)
NEW YORK	2557	90.22	0.74	(88.77, 91.67)
NORTH CAROLINA	1634	92.46	0.81	(90.87, 94.05)
NORTH DAKOTA	1059	90.50	0.97	(88.60, 92.40)
OHIO	923	90.46	1.12	(88.26, 92.66)
OKLAHOMA	1068	92.28	1.04	(90.24, 94.32)
OREGON	1690	93.31	0.72	(91.90, 94.72)
PENNSYLVANIA	2012	84.10	0.99	(82.16, 86.04)
RHODE ISLAND	1046	90.02	1.09	(87.88, 92.16)
SOUTH CAROLINA	1108	90.02	1.17	(87.73, 92.31)
SOUTH DAKOTA	1193	90.07	0.95	(88.21, 91.93)
TENNESSEE	1854	86.92	0.92	(85.12, 88.72)
TEXAS	1035	82.24	1.37	(79.55, 84.93)
UTAH	1623	89.36	1.07	(87.26, 91.46)
VERMONT	1409	88.30	1.06	(86.22, 90.38)
VIRGINIA	1156	90.26	1.02	(88.26, 92.26)
WASHINGTON	2052	91.36	0.71	(89.97, 92.75)
WEST VIRGINIA	1508	90.49	0.85	(88.82, 92.16)
WISCONSIN	1272	94.63	0.78	(93.10, 96.16)
WYOMING	1476	91.60	0.81	(90.01, 93.19)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 50 MEDIAN = 90.24 RANGE = 82.24-94.63
 * EVER HAD A BREAST EXAM BY A DOCTOR, AGE 18 AND OLDER November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 BREAST EXAM - FEMALES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	89.73	86.70	-3.03
ALASKA	92.89	93.83	0.94
ARIZONA	89.46	87.40	-2.06
ARKANSAS	87.75	84.85	-2.90
CALIFORNIA	86.88	86.04	-0.84
COLORADO	92.26	92.06	-0.20
CONNECTICUT	89.25	87.57	-1.68
DELAWARE	92.18	92.96	0.78
DISTRICT OF CO	N/A	90.57	N/A
FLORIDA	88.63	88.40	-0.23
GEORGIA	93.18	92.51	-0.67
IDAHO	90.85	91.33	0.48
ILLINOIS	86.49	89.02	2.53
INDIANA	89.14	89.52	0.38
IOWA	90.35	91.70	1.35
KANSAS	89.38	91.22	1.84
KENTUCKY	87.69	86.97	-0.72
LOUISIANA	84.37	85.73	1.36
MAINE	92.68	91.99	-0.69
MARYLAND	93.00	94.15	1.15
MASSACHUSETTS	91.09	92.82	1.73
MICHIGAN	88.64	89.93	1.29
MINNESOTA	89.53	92.03	2.50
MISSISSIPPI	86.84	87.70	0.86
MISSOURI	90.97	89.43	-1.54
MONTANA	93.78	94.41	0.63
NEBRASKA	89.46	92.34	2.88
NEVADA	88.46	89.72	1.26
NEW HAMPSHIRE	92.20	90.70	-1.50
NEW JERSEY	91.01	83.86	-7.15
NEW MEXICO	89.74	89.03	-0.71
NEW YORK	89.91	90.22	0.31
NORTH CAROLINA	92.50	92.46	-0.04
NORTH DAKOTA	88.77	90.50	1.73
OHIO	90.77	90.46	-0.31
OKLAHOMA	92.27	92.28	0.01
OREGON	93.79	93.31	-0.48
PENNSYLVANIA	86.92	84.10	-2.82
RHODE ISLAND	91.43	90.02	-1.41
SOUTH CAROLINA	90.86	90.02	-0.84
SOUTH DAKOTA	89.71	90.07	0.36
TENNESSEE	88.42	86.92	-1.50
TEXAS	87.95	82.24	-5.71
UTAH	90.77	89.36	-1.41
VERMONT	89.13	88.30	-0.83
VIRGINIA	89.57	90.26	0.69
WASHINGTON	93.38	91.36	-2.02
WEST VIRGINIA	88.86	90.49	1.63
WISCONSIN	91.49	94.63	3.14
WYOMING	91.58	91.60	0.02

* EVER HAD A BREAST EXAM BY A DOCTOR, AGE 18 AND OLDER
 N/A - NOT AVAILABLE

November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BREAST EXAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	757	84.83	1.46	(81.97, 87.69)
ALASKA	449	95.80	1.16	(93.53, 98.07)
ARIZONA	700	92.46	1.12	(90.26, 94.66)
ARKANSAS	741	84.20	1.41	(81.44, 86.96)
CALIFORNIA	1287	89.87	1.01	(87.89, 91.85)
COLORADO	582	91.74	1.22	(89.35, 94.13)
CONNECTICUT	648	88.26	1.47	(85.38, 91.14)
DELAWARE	752	94.29	0.92	(92.49, 96.09)
DISTRICT OF CO	397	93.48	1.27	(90.99, 95.97)
FLORIDA	1300	90.36	0.91	(88.58, 92.14)
GEORGIA	762	92.69	1.03	(90.67, 94.71)
IDAHO	923	92.12	0.97	(90.22, 94.02)
ILLINOIS	974	89.11	1.18	(86.80, 91.42)
INDIANA	791	89.68	1.16	(87.41, 91.95)
IOWA	1315	90.73	0.85	(89.06, 92.40)
KANSAS	671	90.92	1.21	(88.55, 93.29)
KENTUCKY	1435	87.57	0.94	(85.73, 89.41)
LOUISIANA	583	84.51	1.70	(81.18, 87.84)
MAINE	634	92.52	1.11	(90.34, 94.70)
MARYLAND	1521	94.11	0.86	(92.42, 95.80)
MASSACHUSETTS	574	93.94	1.03	(91.92, 95.96)
MICHIGAN	836	90.00	1.17	(87.71, 92.29)
MINNESOTA	1430	92.86	0.73	(91.43, 94.29)
MISSISSIPPI	614	87.71	1.36	(85.04, 90.38)
MISSOURI	574	89.14	1.35	(86.49, 91.79)
MONTANA	641	96.04	0.75	(94.57, 97.51)
NEBRASKA	801	91.59	0.98	(89.67, 93.51)
NEVADA	643	90.74	1.60	(87.60, 93.88)
NEW HAMPSHIRE	476	89.67	1.57	(86.59, 92.75)
NEW JERSEY	1042	83.02	1.37	(80.33, 85.71)
NEW MEXICO	420	88.71	1.79	(85.20, 92.22)
NEW YORK	1456	92.38	0.84	(90.73, 94.03)
NORTH CAROLINA	1000	93.45	0.84	(91.80, 95.10)
NORTH DAKOTA	651	90.09	1.24	(87.66, 92.52)
OHIO	581	93.08	1.04	(91.04, 95.12)
OKLAHOMA	708	92.68	1.23	(90.27, 95.09)
OREGON	1019	93.58	0.86	(91.89, 95.27)
PENNSYLVANIA	1239	83.03	1.18	(80.72, 85.34)
RHODE ISLAND	637	89.53	1.36	(86.86, 92.20)
SOUTH CAROLINA	688	91.69	1.30	(89.14, 94.24)
SOUTH DAKOTA	746	89.90	1.16	(87.63, 92.17)
TENNESSEE	1133	86.29	1.13	(84.08, 88.50)
TEXAS	588	83.67	1.67	(80.40, 86.94)
UTAH	892	91.30	1.23	(88.89, 93.71)
VERMONT	882	88.95	1.14	(86.72, 91.18)
VIRGINIA	667	91.07	1.15	(88.82, 93.32)
WASHINGTON	1170	90.88	0.97	(88.98, 92.78)
WEST VIRGINIA	990	90.35	1.03	(88.33, 92.37)
WISCONSIN	745	95.62	0.86	(93.93, 97.31)
WYOMING	869	90.57	1.04	(88.53, 92.61)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 50 MEDIAN = 90.74 RANGE = 83.02-96.04

* EVER HAD A BREAST EXAM BY A DOCTOR, AGE 40 AND OVER

November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 BREAST EXAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	89.06	84.83	-4.23
ALASKA	94.07	95.80	1.73
ARIZONA	92.89	92.46	-0.43
ARKANSAS	86.88	84.20	-2.68
CALIFORNIA	88.89	89.87	0.98
COLORADO	90.78	91.74	0.96
CONNECTICUT	89.53	88.26	-1.27
DELAWARE	90.58	94.29	3.71
DISTRICT OF CO	N/A	93.48	N/A
FLORIDA	89.32	90.36	1.04
GEORGIA	93.31	92.69	-0.62
IDAHO	92.32	92.12	-0.20
ILLINOIS	88.26	89.11	0.85
INDIANA	87.98	89.68	1.70
IOWA	91.15	90.73	-0.42
KANSAS	90.21	90.92	0.71
KENTUCKY	86.01	87.57	1.56
LOUISIANA	84.92	84.51	-0.41
MAINE	93.27	92.52	-0.75
MARYLAND	93.53	94.11	0.58
MASSACHUSETTS	93.97	93.94	-0.03
MICHIGAN	90.13	90.00	-0.13
MINNESOTA	90.23	92.86	2.63
MISSISSIPPI	87.11	87.71	0.60
MISSOURI	89.74	89.14	-0.60
MONTANA	94.74	96.04	1.30
NEBRASKA	90.38	91.59	1.21
NEVADA	90.05	90.74	0.69
NEW HAMPSHIRE	90.63	89.67	-0.96
NEW JERSEY	90.68	83.02	-7.66
NEW MEXICO	90.77	88.71	-2.06
NEW YORK	91.19	92.38	1.19
NORTH CAROLINA	92.03	93.45	1.42
NORTH DAKOTA	91.22	90.09	-1.13
OHIO	92.43	93.08	0.65
OKLAHOMA	93.42	92.68	-0.74
OREGON	95.86	93.58	-2.28
PENNSYLVANIA	83.14	83.03	-0.11
RHODE ISLAND	91.09	89.53	-1.56
SOUTH CAROLINA	91.97	91.69	-0.28
SOUTH DAKOTA	89.83	89.90	0.07
TENNESSEE	88.68	86.29	-2.39
TEXAS	88.70	83.67	-5.03
UTAH	92.58	91.30	-1.28
VERMONT	89.98	88.95	-1.03
VIRGINIA	87.88	91.07	3.19
WASHINGTON	94.69	90.88	-3.81
WEST VIRGINIA	89.55	90.35	0.80
WISCONSIN	93.44	95.62	2.18
WYOMING	90.88	90.57	-0.31

* EVER HAD A BREAST EXAM BY A DOCTOR, AGE 40 AND OVER
 N/A - NOT AVAILABLE

November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BREAST EXAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	489	68.39	2.27	(63.94, 72.84)
ALASKA	218	82.55	4.14	(74.44, 90.66)
ARIZONA	507	82.68	1.94	(78.88, 86.48)
ARKANSAS	523	65.34	2.32	(60.79, 69.89)
CALIFORNIA	841	77.76	1.64	(74.55, 80.97)
COLORADO	355	77.15	2.58	(72.09, 82.21)
CONNECTICUT	392	74.67	2.49	(69.79, 79.55)
DELAWARE	499	80.22	1.92	(76.46, 83.98)
DISTRICT OF CO	257	83.41	2.46	(78.59, 88.23)
FLORIDA	929	79.77	1.49	(76.85, 82.69)
GEORGIA	473	87.06	1.66	(83.81, 90.31)
IDAHO	613	69.33	2.01	(65.39, 73.27)
ILLINOIS	645	70.99	2.09	(66.89, 75.09)
INDIANA	511	65.15	2.39	(60.47, 69.83)
IOWA	929	64.27	1.76	(60.82, 67.72)
KANSAS	446	75.72	2.20	(71.41, 80.03)
KENTUCKY	1042	70.18	1.56	(67.12, 73.24)
LOUISIANA	395	66.95	2.70	(61.66, 72.24)
MAINE	429	78.70	2.07	(74.64, 82.76)
MARYLAND	971	83.71	1.56	(80.65, 86.77)
MASSACHUSETTS	367	82.74	2.08	(78.66, 86.82)
MICHIGAN	527	74.01	2.15	(69.80, 78.22)
MINNESOTA	938	75.44	1.53	(72.44, 78.44)
MISSISSIPPI	418	66.17	2.48	(61.31, 71.03)
MISSOURI	396	67.30	2.46	(62.48, 72.12)
MONTANA	420	78.59	2.12	(74.43, 82.75)
NEBRASKA	567	74.42	2.00	(70.50, 78.34)
NEVADA	399	71.90	3.08	(65.86, 77.94)
NEW HAMPSHIRE	288	78.03	2.62	(72.89, 83.17)
NEW JERSEY	658	69.28	2.03	(65.30, 73.26)
NEW MEXICO	272	70.57	3.36	(63.98, 77.16)
NEW YORK	922	81.35	1.43	(78.55, 84.15)
NORTH CAROLINA	707	80.24	1.73	(76.85, 83.63)
NORTH DAKOTA	471	74.72	2.17	(70.47, 78.97)
OHIO	416	79.89	2.00	(75.97, 83.81)
OKLAHOMA	522	77.56	2.02	(73.60, 81.52)
OREGON	655	79.49	1.71	(76.14, 82.84)
PENNSYLVANIA	821	68.76	1.75	(65.33, 72.19)
RHODE ISLAND	431	78.45	2.17	(74.20, 82.70)
SOUTH CAROLINA	448	82.05	2.09	(77.95, 86.15)
SOUTH DAKOTA	534	70.34	2.08	(66.26, 74.42)
TENNESSEE	759	71.75	1.80	(68.22, 75.28)
TEXAS	358	69.53	2.64	(64.36, 74.70)
UTAH	581	74.79	2.18	(70.52, 79.06)
VERMONT	557	74.63	1.98	(70.75, 78.51)
VIRGINIA	420	77.65	2.13	(73.48, 81.82)
WASHINGTON	705	77.35	1.71	(74.00, 80.70)
WEST VIRGINIA	678	74.23	1.85	(70.60, 77.86)
WISCONSIN	476	77.13	2.48	(72.27, 81.99)
WYOMING	532	69.55	2.09	(65.45, 73.65)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 50 MEDIAN = 75.12 RANGE = 64.27-87.06

* HAD A BREAST EXAM WITHIN TWO YEARS AGE 50 AND OVER

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 BREAST EXAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	73.79	68.39	-5.40
ALASKA	77.28	82.55	5.27
ARIZONA	81.81	82.68	0.87
ARKANSAS	69.21	65.34	-3.87
CALIFORNIA	75.89	77.76	1.87
COLORADO	74.70	77.15	2.45
CONNECTICUT	76.97	74.67	-2.30
DELAWARE	75.91	80.22	4.31
DISTRICT OF CO	N/A	83.41	N/A
FLORIDA	78.56	79.77	1.21
GEORGIA	83.54	87.06	3.52
IDAHO	71.36	69.33	-2.03
ILLINOIS	78.54	70.99	-7.55
INDIANA	70.32	65.15	-5.17
IOWA	73.09	64.27	-8.82
KANSAS	72.17	75.72	3.55
KENTUCKY	68.21	70.18	1.97
LOUISIANA	69.95	66.95	-3.00
MAINE	81.32	78.70	-2.62
MARYLAND	83.17	83.71	0.54
MASSACHUSETTS	85.24	82.74	-2.50
MICHIGAN	75.05	74.01	-1.04
MINNESOTA	77.12	75.44	-1.68
MISSISSIPPI	66.12	66.17	0.05
MISSOURI	69.25	67.30	-1.95
MONTANA	69.79	78.59	8.80
NEBRASKA	73.36	74.42	1.06
NEVADA	71.09	71.90	0.81
NEW HAMPSHIRE	79.32	78.03	-1.29
NEW JERSEY	77.83	69.28	-8.55
NEW MEXICO	74.73	70.57	-4.16
NEW YORK	82.51	81.35	-1.16
NORTH CAROLINA	76.99	80.24	3.25
NORTH DAKOTA	74.47	74.72	0.25
OHIO	77.47	79.89	2.42
OKLAHOMA	81.74	77.56	-4.18
OREGON	80.17	79.49	-0.68
PENNSYLVANIA	62.70	68.76	6.06
RHODE ISLAND	78.96	78.45	-0.51
SOUTH CAROLINA	82.43	82.05	-0.38
SOUTH DAKOTA	75.13	70.34	-4.79
TENNESSEE	73.42	71.75	-1.67
TEXAS	71.90	69.53	-2.37
UTAH	74.62	74.79	0.17
VERMONT	75.81	74.63	-1.18
VIRGINIA	75.08	77.65	2.57
WASHINGTON	80.37	77.35	-3.02
WEST VIRGINIA	73.59	74.23	0.64
WISCONSIN	73.68	77.13	3.45
WYOMING	65.63	69.55	3.92

* HAD A BREAST EXAM WITHIN TWO YEARS AGE 50 AND OVER
 N/A - NOT AVAILABLE

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD A MAMMOGRAM AND A BREAST EXAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	755	74.78	1.72	(71.41, 78.15)
ALASKA	447	88.04	2.07	(83.98, 92.10)
ARIZONA	700	80.53	2.11	(76.39, 84.67)
ARKANSAS	740	69.56	1.84	(65.95, 73.17)
COLORADO	582	82.04	1.74	(78.63, 85.45)
CONNECTICUT	646	83.51	1.64	(80.30, 86.72)
DELAWARE	752	84.08	1.48	(81.18, 86.98)
DISTRICT OF CO	397	83.82	1.98	(79.94, 87.70)
FLORIDA	1299	81.01	1.18	(78.70, 83.32)
GEORGIA	762	84.56	1.38	(81.86, 87.26)
IDAHO	923	74.76	1.60	(71.62, 77.90)
ILLINOIS	972	78.08	1.52	(75.10, 81.06)
INDIANA	788	74.49	1.71	(71.14, 77.84)
IOWA	1313	75.42	1.30	(72.87, 77.97)
KANSAS	671	73.77	1.82	(70.20, 77.34)
KENTUCKY	1431	76.25	1.26	(73.78, 78.72)
LOUISIANA	581	69.23	2.08	(65.15, 73.31)
MAINE	633	83.01	1.60	(79.87, 86.15)
MARYLAND	1520	85.49	1.23	(83.08, 87.90)
MASSACHUSETTS	574	85.80	1.54	(82.78, 88.82)
MICHIGAN	835	81.71	1.49	(78.79, 84.63)
MINNESOTA	1429	82.15	1.08	(80.03, 84.27)
MISSISSIPPI	611	71.85	1.99	(67.95, 75.75)
MISSOURI	574	74.99	1.90	(71.27, 78.71)
MONTANA	640	84.36	1.51	(81.40, 87.32)
NEBRASKA	801	74.36	1.68	(71.07, 77.65)
NEVADA	643	78.92	2.29	(74.43, 83.41)
NEW HAMPSHIRE	476	81.45	1.95	(77.63, 85.27)
NEW JERSEY	1042	72.84	1.58	(69.74, 75.94)
NEW MEXICO	419	76.33	2.52	(71.39, 81.27)
NEW YORK	1456	81.93	1.12	(79.73, 84.13)
NORTH CAROLINA	996	80.14	1.41	(77.38, 82.90)
NORTH DAKOTA	650	79.90	1.69	(76.59, 83.21)
OHIO	580	79.09	1.80	(75.56, 82.62)
OKLAHOMA	708	69.94	2.07	(65.88, 74.00)
OREGON	1018	83.65	1.29	(81.12, 86.18)
PENNSYLVANIA	1237	73.43	1.38	(70.73, 76.13)
RHODE ISLAND	637	81.82	1.68	(78.53, 85.11)
SOUTH CAROLINA	685	80.03	1.90	(76.31, 83.75)
SOUTH DAKOTA	744	76.43	1.72	(73.06, 79.80)
TENNESSEE	1132	72.75	1.42	(69.97, 75.53)
TEXAS	586	68.74	2.10	(64.62, 72.86)
UTAH	891	78.10	1.71	(74.75, 81.45)
VERMONT	881	79.05	1.47	(76.17, 81.93)
VIRGINIA	667	80.16	1.66	(76.91, 83.41)
WASHINGTON	1169	81.13	1.25	(78.68, 83.58)
WEST VIRGINIA	990	76.91	1.45	(74.07, 79.75)
WISCONSIN	744	87.09	1.51	(84.13, 90.05)
WYOMING	869	74.78	1.60	(71.64, 77.92)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 79.09 RANGE = 68.74-88.04 YEAR 2000 OBJ 16.11
 * EVER HAD A MAMMOGRAM AND A BREAST EXAM FOR AGE 40 AND OVER November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 HAD A MAMMOGRAM AND A BREAST EXAM - FEMALES*
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	72.82	74.78	1.96
ALASKA	83.75	88.04	4.29
ARIZONA	82.09	80.53	-1.56
ARKANSAS	67.61	69.56	1.95
COLORADO	79.64	82.04	2.40
CONNECTICUT	80.07	83.51	3.44
DELAWARE	81.33	84.08	2.75
DISTRICT OF CO	N/A	83.82	N/A
FLORIDA	78.32	81.01	2.69
GEORGIA	79.99	84.56	4.57
IDAHO	76.32	74.76	-1.56
ILLINOIS	76.02	78.08	2.06
INDIANA	73.23	74.49	1.26
IOWA	77.39	75.42	-1.97
KANSAS	75.38	73.77	-1.61
KENTUCKY	74.48	76.25	1.77
LOUISIANA	68.03	69.23	1.20
MAINE	79.56	83.01	3.45
MARYLAND	82.22	85.49	3.27
MASSACHUSETTS	87.13	85.80	-1.33
MICHIGAN	81.61	81.71	0.10
MINNESOTA	78.44	82.15	3.71
MISSISSIPPI	67.19	71.85	4.66
MISSOURI	74.99	74.99	0.00
MONTANA	76.19	84.36	8.17
NEBRASKA	72.18	74.36	2.18
NEVADA	78.86	78.92	0.06
NEW HAMPSHIRE	80.80	81.45	0.65
NEW JERSEY	74.81	72.84	-1.97
NEW MEXICO	77.77	76.33	-1.44
NEW YORK	80.41	81.93	1.52
NORTH CAROLINA	77.70	80.14	2.44
NORTH DAKOTA	77.24	79.90	2.66
OHIO	79.19	79.09	-0.10
OKLAHOMA	72.32	69.94	-2.38
OREGON	83.92	83.65	-0.27
PENNSYLVANIA	72.44	73.43	0.99
RHODE ISLAND	81.53	81.82	0.29
SOUTH CAROLINA	79.66	80.03	0.37
SOUTH DAKOTA	74.53	76.43	1.90
TENNESSEE	74.57	72.75	-1.82
TEXAS	72.54	68.74	-3.80
UTAH	78.70	78.10	-0.60
VERMONT	78.36	79.05	0.69
VIRGINIA	76.14	80.16	4.02
WASHINGTON	83.15	81.13	-2.02
WEST VIRGINIA	73.55	76.91	3.36
WISCONSIN	78.63	87.09	8.46
WYOMING	74.24	74.78	0.54

* EVER HAD A MAMMOGRAM AND A BREAST EXAM FOR AGE 40 AND OVER

N/A - NOT AVAILABLE

November 20, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD A MAMMOGRAM AND BREAST EXAM *
 DENOMINATOR IS FEMALES AGE 50 AND OVER

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	518	56.01	2.37	(51.36, 60.66)
ALASKA	226	69.40	4.89	(59.82, 78.98)
ARIZONA	527	71.58	2.46	(66.76, 76.40)
ARKANSAS	541	51.76	2.41	(47.04, 56.48)
COLORADO	370	69.23	2.72	(63.90, 74.56)
CONNECTICUT	416	65.65	2.62	(60.51, 70.79)
DELAWARE	502	73.49	2.17	(69.24, 77.74)
DISTRICT OF CO	261	75.46	2.79	(69.99, 80.93)
FLORIDA	953	70.92	1.61	(67.76, 74.08)
GEORGIA	477	74.83	2.09	(70.73, 78.93)
IDAHO	629	56.62	2.14	(52.43, 60.81)
ILLINOIS	654	60.42	2.14	(56.23, 64.61)
INDIANA	529	53.13	2.47	(48.29, 57.97)
IOWA	965	52.38	1.81	(48.83, 55.93)
KANSAS	478	60.67	2.41	(55.95, 65.39)
KENTUCKY	1081	59.35	1.65	(56.12, 62.58)
LOUISIANA	398	53.66	2.75	(48.27, 59.05)
MAINE	439	69.08	2.37	(64.43, 73.73)
MARYLAND	1013	73.09	1.81	(69.54, 76.64)
MASSACHUSETTS	376	74.48	2.43	(69.72, 79.24)
MICHIGAN	532	65.89	2.28	(61.42, 70.36)
MINNESOTA	956	64.94	1.66	(61.69, 68.19)
MISSISSIPPI	429	53.05	2.60	(47.95, 58.15)
MISSOURI	408	55.70	2.63	(50.55, 60.85)
MONTANA	432	63.13	2.55	(58.13, 68.13)
NEBRASKA	580	59.68	2.20	(55.37, 63.99)
NEVADA	404	59.21	3.46	(52.43, 65.99)
NEW HAMPSHIRE	301	67.57	2.80	(62.08, 73.06)
NEW JERSEY	692	57.01	2.14	(52.82, 61.20)
NEW MEXICO	277	58.22	3.72	(50.93, 65.51)
NEW YORK	935	71.84	1.57	(68.76, 74.92)
NORTH CAROLINA	732	67.92	1.96	(64.08, 71.76)
NORTH DAKOTA	482	64.50	2.34	(59.91, 69.09)
OHIO	423	67.48	2.48	(62.62, 72.34)
OKLAHOMA	528	56.84	2.54	(51.86, 61.82)
OREGON	671	70.30	1.93	(66.52, 74.08)
PENNSYLVANIA	865	57.32	1.82	(53.75, 60.89)
RHODE ISLAND	441	68.02	2.46	(63.20, 72.84)
SOUTH CAROLINA	462	68.10	2.50	(63.20, 73.00)
SOUTH DAKOTA	550	59.15	2.23	(54.78, 63.52)
TENNESSEE	782	58.63	1.85	(55.00, 62.26)
TEXAS	366	56.34	2.82	(50.81, 61.87)
UTAH	588	61.51	2.43	(56.75, 66.27)
VERMONT	579	63.59	2.22	(59.24, 67.94)
VIRGINIA	427	63.55	2.55	(58.55, 68.55)
WASHINGTON	732	67.42	1.88	(63.74, 71.10)
WEST VIRGINIA	694	63.09	2.08	(59.01, 67.17)
WISCONSIN	491	66.44	2.68	(61.19, 71.69)
WYOMING	545	55.76	2.26	(51.33, 60.19)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 63.55 RANGE = 51.76-75.46 YEAR 2000 OBJ 16.11
 * HAD A MAMMOGRAM AND A BREAST EXAM WITHIN 2 YEARS, AGE 50 AND OLDER December 3, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 HAD A MAMMOGRAM AND BREAST EXAM *
 DENOMINATOR IS FEMALES AGE 50 AND OVER

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	55.33	56.01	0.68
ALASKA	70.23	69.40	-0.83
ARIZONA	70.79	71.58	0.79
ARKANSAS	55.10	51.76	-3.34
COLORADO	60.83	69.23	8.40
CONNECTICUT	67.57	65.65	-1.92
DELAWARE	65.94	73.49	7.55
DISTRICT OF CO	N/A	75.46	N/A
FLORIDA	69.17	70.92	1.75
GEORGIA	70.00	74.83	4.83
IDAHO	57.89	56.62	-1.27
ILLINOIS	64.53	60.42	-4.11
INDIANA	55.78	53.13	-2.65
IOWA	58.38	52.38	-6.00
KANSAS	57.04	60.67	3.63
KENTUCKY	55.94	59.35	3.41
LOUISIANA	53.95	53.66	-0.29
MAINE	68.50	69.08	0.58
MARYLAND	71.76	73.09	1.33
MASSACHUSETTS	75.71	74.48	-1.23
MICHIGAN	67.60	65.89	-1.71
MINNESOTA	66.07	64.94	-1.13
MISSISSIPPI	47.51	53.05	5.54
MISSOURI	58.65	55.70	-2.95
MONTANA	58.76	63.13	4.37
NEBRASKA	58.15	59.68	1.53
NEVADA	59.19	59.21	0.02
NEW HAMPSHIRE	64.89	67.57	2.68
NEW JERSEY	57.84	57.01	-0.83
NEW MEXICO	64.45	58.22	-6.23
NEW YORK	68.78	71.84	3.06
NORTH CAROLINA	61.69	67.92	6.23
NORTH DAKOTA	62.23	64.50	2.27
OHIO	63.87	67.48	3.61
OKLAHOMA	58.89	56.84	-2.05
OREGON	70.35	70.30	-0.05
PENNSYLVANIA	51.10	57.32	6.22
RHODE ISLAND	62.15	68.02	5.87
SOUTH CAROLINA	68.43	68.10	-0.33
SOUTH DAKOTA	57.11	59.15	2.04
TENNESSEE	58.65	58.63	-0.02
TEXAS	59.67	56.34	-3.33
UTAH	61.35	61.51	0.16
VERMONT	63.17	63.59	0.42
VIRGINIA	58.10	63.55	5.45
WASHINGTON	69.38	67.42	-1.96
WEST VIRGINIA	58.84	63.09	4.25
WISCONSIN	55.11	66.44	11.33
WYOMING	51.31	55.76	4.45

* HAD A MAMMOGRAM AND A BREAST EXAM WITHIN 2 YEARS, AGE 50 AND OLDER

N/A - NOT AVAILABLE

December 3, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PAP SMEAR*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1338	95.69	0.70	(94.32, 97.06)
ALASKA	813	97.18	0.81	(95.59, 98.77)
ARIZONA	1130	93.85	1.05	(91.79, 95.91)
ARKANSAS	1144	92.87	0.91	(91.09, 94.65)
COLORADO	984	96.51	0.67	(95.20, 97.82)
CONNECTICUT	1060	93.57	0.89	(91.83, 95.31)
DELAWARE	1294	94.07	0.89	(92.33, 95.81)
DISTRICT OF CO	735	95.76	0.89	(94.02, 97.50)
FLORIDA	2117	94.44	0.59	(93.28, 95.60)
GEORGIA	1353	96.01	0.61	(94.81, 97.21)
IDAHO	1610	95.88	0.57	(94.76, 97.00)
ILLINOIS	1720	94.00	0.68	(92.67, 95.33)
INDIANA	1258	95.34	0.76	(93.85, 96.83)
IOWA	2092	95.58	0.54	(94.52, 96.64)
KANSAS	1106	95.12	0.76	(93.63, 96.61)
KENTUCKY	2268	92.90	0.64	(91.65, 94.15)
LOUISIANA	985	93.02	0.92	(91.22, 94.82)
MAINE	983	95.66	0.86	(93.97, 97.35)
MARYLAND	2597	95.38	0.58	(94.24, 96.52)
MASSACHUSETTS	1030	92.94	1.02	(90.94, 94.94)
MICHIGAN	1438	96.00	0.59	(94.84, 97.16)
MINNESOTA	2430	93.40	0.59	(92.24, 94.56)
MISSISSIPPI	994	92.68	0.94	(90.84, 94.52)
MISSOURI	921	95.65	0.80	(94.08, 97.22)
MONTANA	1022	96.28	0.72	(94.87, 97.69)
NEBRASKA	1250	94.46	0.72	(93.05, 95.87)
NEVADA	1038	96.18	0.80	(94.61, 97.75)
NEW HAMPSHIRE	831	93.79	1.28	(91.28, 96.30)
NEW JERSEY	1760	91.03	0.91	(89.25, 92.81)
NEW MEXICO	658	90.22	1.82	(86.65, 93.79)
NEW YORK	2554	92.28	0.65	(91.01, 93.55)
NORTH CAROLINA	1642	95.41	0.67	(94.10, 96.72)
NORTH DAKOTA	1059	95.28	0.75	(93.81, 96.75)
OHIO	915	94.82	0.84	(93.17, 96.47)
OKLAHOMA	1070	94.99	0.92	(93.19, 96.79)
OREGON	1684	96.15	0.59	(94.99, 97.31)
PENNSYLVANIA	2008	93.24	0.77	(91.73, 94.75)
RHODE ISLAND	1046	91.78	1.04	(89.74, 93.82)
SOUTH CAROLINA	1107	94.06	1.05	(92.00, 96.12)
SOUTH DAKOTA	1200	94.78	0.76	(93.29, 96.27)
TENNESSEE	1855	93.44	0.70	(92.07, 94.81)
TEXAS	1036	90.80	1.11	(88.62, 92.98)
UTAH	1627	90.70	1.03	(88.68, 92.72)
VERMONT	1411	94.66	0.87	(92.95, 96.37)
VIRGINIA	1156	94.86	0.82	(93.25, 96.47)
WASHINGTON	2053	96.31	0.47	(95.39, 97.23)
WEST VIRGINIA	1512	94.10	0.77	(92.59, 95.61)
WISCONSIN	1270	96.47	0.70	(95.10, 97.84)
WYOMING	1479	96.32	0.61	(95.12, 97.52)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 94.78 RANGE = 90.22-97.18 YEAR 2000 OBJ 16.11
 * EVER HAD A PAP SMEAR November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 PAP SMEAR*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	95.85	95.69	-0.16
ALASKA	95.55	97.18	1.63
ARIZONA	93.86	93.85	-0.01
ARKANSAS	94.17	92.87	-1.30
COLORADO	97.28	96.51	-0.77
CONNECTICUT	93.74	93.57	-0.17
DELAWARE	94.02	94.07	0.05
DISTRICT OF CO	N/A	95.76	N/A
FLORIDA	94.88	94.44	-0.44
GEORGIA	95.58	96.01	0.43
IDAHO	94.31	95.88	1.57
ILLINOIS	88.42	94.00	5.58
INDIANA	94.68	95.34	0.66
IOWA	94.58	95.58	1.00
KANSAS	93.41	95.12	1.71
KENTUCKY	93.47	92.90	-0.57
LOUISIANA	88.94	93.02	4.08
MAINE	93.40	95.66	2.26
MARYLAND	91.70	95.38	3.68
MASSACHUSETTS	93.55	92.94	-0.61
MICHIGAN	94.70	96.00	1.30
MINNESOTA	90.59	93.40	2.81
MISSISSIPPI	93.30	92.68	-0.62
MISSOURI	94.75	95.65	0.90
MONTANA	94.90	96.28	1.38
NEBRASKA	93.02	94.46	1.44
NEVADA	93.45	96.18	2.73
NEW HAMPSHIRE	95.70	93.79	-1.91
NEW JERSEY	90.07	91.03	0.96
NEW MEXICO	94.29	90.22	-4.07
NEW YORK	89.35	92.28	2.93
NORTH CAROLINA	95.84	95.41	-0.43
NORTH DAKOTA	93.44	95.28	1.84
OHIO	94.63	94.82	0.19
OKLAHOMA	95.11	94.99	-0.12
OREGON	96.52	96.15	-0.37
PENNSYLVANIA	94.23	93.24	-0.99
RHODE ISLAND	93.80	91.78	-2.02
SOUTH CAROLINA	95.81	94.06	-1.75
SOUTH DAKOTA	92.80	94.78	1.98
TENNESSEE	92.91	93.44	0.53
TEXAS	93.12	90.80	-2.32
UTAH	91.20	90.70	-0.50
VERMONT	94.81	94.66	-0.15
VIRGINIA	94.39	94.86	0.47
WASHINGTON	96.78	96.31	-0.47
WEST VIRGINIA	91.79	94.10	2.31
WISCONSIN	95.00	96.47	1.47
WYOMING	95.79	96.32	0.53

* EVER HAD A PAP SMEAR
 N/A - NOT AVAILABLE

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER HAD A PAP SMEAR*
 WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	966	94.98	0.87	(93.27, 96.69)
ALASKA	659	96.99	0.98	(95.07, 98.91)
ARIZONA	822	92.30	1.36	(89.63, 94.97)
ARKANSAS	772	93.43	1.09	(91.29, 95.57)
COLORADO	750	96.75	0.77	(95.24, 98.26)
CONNECTICUT	861	94.06	0.99	(92.12, 96.00)
DELAWARE	1067	94.25	1.00	(92.29, 96.21)
DISTRICT OF CO	619	95.66	0.97	(93.76, 97.56)
FLORIDA	1602	93.95	0.72	(92.54, 95.36)
GEORGIA	1048	95.86	0.74	(94.41, 97.31)
IDAHO	1161	95.59	0.72	(94.18, 97.00)
ILLINOIS	1394	94.10	0.76	(92.61, 95.59)
INDIANA	948	94.90	0.95	(93.04, 96.76)
IOWA	1618	95.44	0.65	(94.17, 96.71)
KANSAS	881	95.43	0.86	(93.74, 97.12)
KENTUCKY	1598	92.77	0.80	(91.20, 94.34)
LOUISIANA	689	94.58	1.00	(92.62, 96.54)
MAINE	776	95.31	1.03	(93.29, 97.33)
MARYLAND	2077	95.75	0.61	(94.55, 96.95)
MASSACHUSETTS	883	92.64	1.15	(90.39, 94.89)
MICHIGAN	1129	95.96	0.68	(94.63, 97.29)
MINNESOTA	2013	93.66	0.65	(92.39, 94.93)
MISSISSIPPI	681	92.64	1.21	(90.27, 95.01)
MISSOURI	703	96.39	0.80	(94.82, 97.96)
MONTANA	791	96.20	0.87	(94.49, 97.91)
NEBRASKA	954	94.56	0.85	(92.89, 96.23)
NEVADA	765	96.12	0.83	(94.49, 97.75)
NEW HAMPSHIRE	696	93.79	1.42	(91.01, 96.57)
NEW JERSEY	1512	90.61	1.00	(88.65, 92.57)
NEW MEXICO	483	90.12	2.23	(85.75, 94.49)
NEW YORK	2180	91.74	0.74	(90.29, 93.19)
NORTH CAROLINA	1219	94.65	0.86	(92.96, 96.34)
NORTH DAKOTA	830	94.97	0.89	(93.23, 96.71)
OHIO	694	94.36	1.02	(92.36, 96.36)
OKLAHOMA	698	95.83	1.04	(93.79, 97.87)
OREGON	1233	95.45	0.77	(93.94, 96.96)
PENNSYLVANIA	1608	93.02	0.92	(91.22, 94.82)
RHODE ISLAND	833	91.62	1.20	(89.27, 93.97)
SOUTH CAROLINA	795	93.21	1.32	(90.62, 95.80)
SOUTH DAKOTA	966	95.14	0.86	(93.45, 96.83)
TENNESSEE	1346	93.28	0.87	(91.57, 94.99)
TEXAS	746	89.52	1.38	(86.82, 92.22)
UTAH	1230	90.85	1.20	(88.50, 93.20)
VERMONT	1173	95.11	0.97	(93.21, 97.01)
VIRGINIA	918	94.49	0.98	(92.57, 96.41)
WASHINGTON	1609	95.80	0.57	(94.68, 96.92)
WEST VIRGINIA	1078	94.86	0.82	(93.25, 96.47)
WISCONSIN	1013	96.72	0.75	(95.25, 98.19)
WYOMING	1099	96.60	0.69	(95.25, 97.95)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 94.65 RANGE = 89.52-96.99 YEAR 2000 OBJ 16.12
 * EVER HAD A PAP SMEAR, AGE 18 AND OLDER November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 EVER HAD A PAP SMEAR*
 WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	94.92	94.98	0.06
ALASKA	95.68	96.99	1.31
ARIZONA	93.72	92.30	-1.42
ARKANSAS	93.85	93.43	-0.42
COLORADO	97.44	96.75	-0.69
CONNECTICUT	94.16	94.06	-0.10
DELAWARE	94.58	94.25	-0.33
DISTRICT OF CO	N/A	95.66	N/A
FLORIDA	94.40	93.95	-0.45
GEORGIA	95.16	95.86	0.70
IDAHO	93.59	95.59	2.00
ILLINOIS	89.37	94.10	4.73
INDIANA	93.77	94.90	1.13
IOWA	94.50	95.44	0.94
KANSAS	93.57	95.43	1.86
KENTUCKY	93.32	92.77	-0.55
LOUISIANA	89.84	94.58	4.74
MAINE	93.58	95.31	1.73
MARYLAND	91.72	95.75	4.03
MASSACHUSETTS	93.09	92.64	-0.45
MICHIGAN	94.57	95.96	1.39
MINNESOTA	91.43	93.66	2.23
MISSISSIPPI	92.55	92.64	0.09
MISSOURI	94.93	96.39	1.46
MONTANA	94.49	96.20	1.71
NEBRASKA	92.36	94.56	2.20
NEVADA	93.09	96.12	3.03
NEW HAMPSHIRE	95.67	93.79	-1.88
NEW JERSEY	90.05	90.61	0.56
NEW MEXICO	94.35	90.12	-4.23
NEW YORK	88.85	91.74	2.89
NORTH CAROLINA	95.88	94.65	-1.23
NORTH DAKOTA	93.05	94.97	1.92
OHIO	93.93	94.36	0.43
OKLAHOMA	95.04	95.83	0.79
OREGON	95.87	95.45	-0.42
PENNSYLVANIA	94.28	93.02	-1.26
RHODE ISLAND	93.50	91.62	-1.88
SOUTH CAROLINA	95.84	93.21	-2.63
SOUTH DAKOTA	93.41	95.14	1.73
TENNESSEE	92.60	93.28	0.68
TEXAS	92.38	89.52	-2.86
UTAH	89.92	90.85	0.93
VERMONT	94.94	95.11	0.17
VIRGINIA	94.76	94.49	-0.27
WASHINGTON	96.39	95.80	-0.59
WEST VIRGINIA	91.80	94.86	3.06
WISCONSIN	95.40	96.72	1.32
WYOMING	95.70	96.60	0.90

* EVER HAD A PAP SMEAR, AGE 18 AND OLDER
 N/A - NOT AVAILABLE

November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEIL LANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LAST PAP SMEAR*
 WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	913	88.84	1.14	(86.61, 91.07)
ALASKA	630	95.74	1.04	(93.70, 97.78)
ARIZONA	756	91.80	1.38	(89.10, 94.50)
ARKANSAS	711	89.05	1.22	(86.66, 91.44)
COLORADO	724	92.75	1.15	(90.50, 95.00)
CONNECTICUT	807	89.81	1.29	(87.28, 92.34)
DELAWARE	1021	91.33	0.93	(89.51, 93.15)
DISTRICT OF CO	592	95.12	1.00	(93.16, 97.08)
FLORIDA	1506	89.65	0.89	(87.91, 91.39)
GEORGIA	1001	96.54	0.62	(95.32, 97.76)
IDAHO	1114	87.34	1.00	(85.38, 89.30)
ILLINOIS	1318	90.03	0.93	(88.21, 91.85)
INDIANA	898	86.34	1.23	(83.93, 88.75)
IOWA	1546	84.01	1.04	(81.97, 86.05)
KANSAS	841	91.64	1.00	(89.68, 93.60)
KENTUCKY	1477	86.63	0.92	(84.83, 88.43)
LOUISIANA	651	90.13	1.32	(87.54, 92.72)
MAINE	739	87.81	1.33	(85.20, 90.42)
MARYLAND	1978	92.10	0.78	(90.57, 93.63)
MASSACHUSETTS	824	90.63	1.12	(88.43, 92.83)
MICHIGAN	1081	90.15	1.02	(88.15, 92.15)
MINNESOTA	1893	92.36	0.64	(91.11, 93.61)
MISSISSIPPI	632	88.83	1.31	(86.26, 91.40)
MISSOURI	673	90.64	1.12	(88.44, 92.84)
MONTANA	762	89.98	1.09	(87.84, 92.12)
NEBRASKA	895	87.77	1.20	(85.42, 90.12)
NEVADA	726	88.23	1.65	(85.00, 91.46)
NEW HAMPSHIRE	654	90.16	1.22	(87.77, 92.55)
NEW JERSEY	1388	88.66	0.96	(86.78, 90.54)
NEW MEXICO	450	91.86	1.36	(89.19, 94.53)
NEW YORK	2019	90.44	0.73	(89.01, 91.87)
NORTH CAROLINA	1142	92.18	0.92	(90.38, 93.98)
NORTH DAKOTA	788	86.38	1.29	(83.85, 88.91)
OHIO	654	91.01	1.13	(88.80, 93.22)
OKLAHOMA	666	90.61	1.31	(88.04, 93.18)
OREGON	1187	90.17	0.98	(88.25, 92.09)
PENNSYLVANIA	1499	86.88	0.96	(85.00, 88.76)
RHODE ISLAND	771	90.51	1.13	(88.30, 92.72)
SOUTH CAROLINA	745	92.38	1.08	(90.26, 94.50)
SOUTH DAKOTA	919	88.42	1.14	(86.19, 90.65)
TENNESSEE	637	90.55	1.24	(88.12, 92.98)
TEXAS	675	89.97	1.24	(87.54, 92.40)
UTAH	1126	89.59	1.04	(87.55, 91.63)
VERMONT	1127	91.45	0.85	(89.78, 93.12)
VIRGINIA	865	92.54	0.99	(90.60, 94.48)
WASHINGTON	1533	89.72	0.87	(88.01, 91.43)
WEST VIRGINIA	1020	85.83	1.12	(83.63, 88.03)
WISCONSIN	974	90.00	1.18	(87.69, 92.31)
WYOMING	1064	85.96	1.19	(83.63, 88.29)

SUMMARY STATISTICS NO. OF PARTICIPANTS = 49 MEDIAN = 90.13 RANGE = 84.01-96.54 YEAR 2000 OBJ 16.12
 * LAST PAP SMEAR WITHIN LAST 3 YEARS, AGE 18 AND OLDER November 14, 1997

1996 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1996
 LAST PAP SMEAR*
 WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1995 PERCENT	1996 PERCENT	PERCENT DIFFERENCE
ALABAMA	90.26	88.84	-1.42
ALASKA	95.29	95.74	0.45
ARIZONA	90.20	91.80	1.60
ARKANSAS	87.72	89.05	1.33
COLORADO	90.75	92.75	2.00
CONNECTICUT	92.95	89.81	-3.14
DELAWARE	91.39	91.33	-0.06
DISTRICT OF CO	N/A	95.12	N/A
FLORIDA	90.91	89.65	-1.26
GEORGIA	94.04	96.54	2.50
IDAHO	86.96	87.34	0.38
ILLINOIS	91.80	90.03	-1.77
INDIANA	87.97	86.34	-1.63
IOWA	87.96	84.01	-3.95
KANSAS	88.47	91.64	3.17
KENTUCKY	85.27	86.63	1.36
LOUISIANA	88.73	90.13	1.40
MAINE	94.08	87.81	-6.27
MARYLAND	93.73	92.10	-1.63
MASSACHUSETTS	91.81	90.63	-1.18
MICHIGAN	89.34	90.15	0.81
MINNESOTA	92.62	92.36	-0.26
MISSISSIPPI	91.10	88.83	-2.27
MISSOURI	88.72	90.64	1.92
MONTANA	86.91	89.98	3.07
NEBRASKA	89.80	87.77	-2.03
NEVADA	90.08	88.23	-1.85
NEW HAMPSHIRE	92.37	90.16	-2.21
NEW JERSEY	90.26	88.66	-1.60
NEW MEXICO	92.39	91.86	-0.53
NEW YORK	91.21	90.44	-0.77
NORTH CAROLINA	92.32	92.18	-0.14
NORTH DAKOTA	89.55	86.38	-3.17
OHIO	90.56	91.01	0.45
OKLAHOMA	93.31	90.61	-2.70
OREGON	88.24	90.17	1.93
PENNSYLVANIA	87.41	86.88	-0.53
RHODE ISLAND	88.73	90.51	1.78
SOUTH CAROLINA	93.64	92.38	-1.26
SOUTH DAKOTA	90.48	88.42	-2.06
TENNESSEE	91.66	90.55	-1.11
TEXAS	90.18	89.97	-0.21
UTAH	89.71	89.59	-0.12
VERMONT	91.19	91.45	0.26
VIRGINIA	93.70	92.54	-1.16
WASHINGTON	91.14	89.72	-1.42
WEST VIRGINIA	86.99	85.83	-1.16
WISCONSIN	88.79	90.00	1.21
WYOMING	84.55	85.96	1.41

* LAST PAP SMEAR WITHIN LAST 3 YEARS, AGE 18 AND OLDER
 N/A - NOT AVAILABLE

November 14, 1997

