

Mailstop K30
(770) 488-5292
August 21, 1998

Dear Colleague:

Information related to the 1997 BRFSS summary prevalence report of selected risk factors and preventive health measures from the core and modules is attached.

This report consists of four parts:

An introduction that contains:

An overview of the 1997 summary prevalence report.

An explanation of the tables for 1997 Behavioral Risk Factor Surveillance System (BRFSS) prevalence data.

Definitions of BRFSS risk factors and preventive health measures.
When appropriate, Year 2000 Objectives are identified.

Analysis of selected questions from the 1997 BRFSS questionnaire. The set of analyses is divided into three sections:

Section I - selected risk factors and preventive health measures in the BRFSS core.

Section II - selected risk factors and preventive health measures from modules used by at least 13 states

Section III - preventive health measures that concern women only.

As soon as your schedule permits, please examine the enclosed materials closely.

Sincerely yours,

David E. Nelson, M.D.,M.P.H.
Chief, Behavioral Surveillance Branch
Division of Adult and Community Health
National Center for Chronic Disease
Prevention and Health Promotion

Enclosures

1997 BRFSS SUMMARY PREVALENCE REPORT

BEHAVIORAL SURVEILLANCE BRANCH
DIVISION OF ADULT AND COMMUNITY HEALTH
NATIONAL CENTER FOR CHRONIC DISEASE PREVENTION
AND HEALTH PROMOTION
CENTERS FOR DISEASE CONTROL AND PREVENTION

TABLE OF CONTENTS

Overview of the 1997 Summary Report.....	ix
Explanation of Tables	x
Definitions of Risk Factors & Preventive Health Measures....	xvii

SECTION I

PREVALENCE REPORT FOR CORE QUESTIONS

General Health Status

TABLE 1.1	Self reported health status: Fair or poor
1.2	Self reported health status: Fair or poor-by sex
1.3	Self reported health status: Fair or poor, change 1996-1997

Quality of Life: Physical and Mental Health

TABLE 2.1	Physical health not good
2.2	Physical health not good-by sex
2.3	Physical health not good, change 1996-1997
TABLE 3.1	Mental health not good
3.2	Mental health not good-by sex
3.3	Mental health not good, change 1996-1997
TABLE 4.1	Activities limited
4.2	Activities limited-by sex
4.3	Activities limited, change 1996-1997

Health Insurance

TABLE 5.1	No health insurance, age 18 and over
5.2	No health insurance, age 18 and over-by sex
5.3	No health insurance, age 18 and over, change 1996-1997
TABLE 6.1	No health insurance, age 18-64 years
6.2	No health insurance, age 18-64 year-by sex
6.3	No health insurance, age 18-64 years, change 1996-1997

Diabetes Awareness

TABLE 7.1	Ever told you have diabetes
7.2	Ever told you have diabetes-by sex
7.3	Ever told you have diabetes, change 1996-1997

Hypertension Awareness and Screening

TABLE 8.1	Ever told blood pressure high
8.2	Ever told blood pressure high-by sex
8.3	Ever told blood pressure high, change 1995-1997
TABLE 9.1	Had blood pressure check within past two years
9.2	Had blood pressure check within past two years-by sex
9.3	Had blood pressure check within past two years, change 1995-1997

Cholesterol Screening and Awareness

TABLE 10.1	Ever had cholesterol check
10.2	Ever had cholesterol check-by sex
10.3	Ever had cholesterol check, change 1995-1997

TABLE 11.1	Had cholesterol check within past five years (Year 2000 Objective 15.14)
11.2	Had cholesterol check within past five years-by sex
11.3	Had cholesterol check within past five years, change 1995-1997

TABLE 12.1	Ever told cholesterol high
12.2	Ever told cholesterol high-by sex
12.3	Ever told cholesterol high, change 1995-1997

Smoking Status

TABLE 13.1	Ever smoked
13.2	Ever smoked-by sex
13.3	Ever smoked, change 1996-1997

TABLE 14.1	Current smoker (Year 2000 Objective 3.4)
14.2	Current smoker-by sex
14.3	Current smoker, change 1996-1997

Adult safety belt usage

TABLE 15.1	Does not always wear a safety belt (Approximates Year 2000 Objective 9.12)
15.2	Does not always wear a safety belt-by sex
15.3	Does not always wear a safety belt, change 1995-1997

Injury Control and Child Safety

TABLE 16.1	Child does not always use safety belt (Approximates Year 2000 Objective 9.12)
16.2	Child does not always use safety belt-by sex
16.3	Child does not always use safety belt, change 1995-1997
TABLE 17.1	Child does not always use safety seat (Approximates Year 2000 Objective 9.12a)
17.2	Child does not always use safety seat-by sex
17.3	Child does not always use safety seat, change 1995-1997

TABLE 18.1	Child does not always use bicycle helmet
18.2	Child does not always use bicycle helmet-by sex
18.3	Child does not always use bicycle helmet, change 1995-1997

Alcohol Consumption

TABLE 19.1	Current drinkers
19.2	Current drinkers-by sex
19.3	Current drinkers, change 1995-1997

TABLE 20.1	Binge drinkers (acute drinking)
20.2	Binge drinkers (acute drinking)-by sex
20.3	Binge drinkers (acute drinking), change 1995-1997

TABLE 21.1	Chronic drinkers (acute drinking)
------------	-----------------------------------

21.2	Chronic drinkers (acute drinking)-by sex
21.3	Chronic drinkers (acute drinking), change 1995-1997
TABLE 22.1	Drinking and driving
22.2	Drinking and driving-by sex
22.3	Drinking and driving, change 1995-1997

Immunization

TABLE 23.1	Had a flu shot within past 12 months, age 65 and older (Year 2000 Objective 20.11)
TABLE 23.2	Had a flu shot within past 12 months, age 65 and older-by sex
23.3	Had a flu shot within past 12 months, age 65 and older, change 1995-1997
TABLE 24.1	Ever had pneumonia vaccination, age 65 and older (Year 2000 Objective 20.11)
24.2	Ever had pneumonia vaccination, age 65 and older-by sex
24.3	Ever had pneumonia vaccination, age 65 and older, change 1995-1997

Colorectal Cancer Screening

TABLE 25.1	Ever had sigmoidoscopy, age 50 and older (Year 2000 Objective 16.13)
25.2	Ever had sigmoidoscopy, age 50 and older-by sex
TABLE 26.1	Did not use home blood stool test kit within past two years, age 50 and older (Year 2000 Objective 16.13)
26.2	Did not use home blood stool test kit within past two years, age 50 and older-by sex

Overweight

TABLE 27.1	Based on body mass index (BMI) (Year 2000 Objective 2.3)
27.2	Based on body mass index (BMI)-by sex
27.3	Based on body mass index (BMI), change 1996-1997
TABLE 28.1	Based on percent of median
28.2	Based on percent of median-by sex
28.3	Based on percent of median, change 1996-1997

AIDS Knowledge/Attitudes

TABLE 29.1	Encourage teen condom use, age 18-64
29.2	Encourage teen condom use-by sex
29.3	Encourage teen condom use, change 1996-1997
TABLE 30.1	HIV blood test, age 18-64
30.2	HIV blood test, age 18-64 -by sex
30.3	HIV blood test, change 1996-1997
TABLE 31.1	Chances of getting HIV, age 18-64
31.2	Chances of getting HIV, age 18-64 -by sex
31.3	Chances of getting HIV, change 1996-1997

SECTION II

PREVALENCE REPORT FOR MODULES

Smokeless tobacco

TABLE 32.1 Ever Use
 32.2 Ever Use-by sex

TABLE 33.1 Current Use
 33.2 Current Use-by sex

Diabetes

TABLE 34.1 Age at onset of diabetes
 34.2 Age at onset of diabetes-by sex

TABLE 35.1 Diabetics using insulin
 35.2 Diabetics using insulin-by sex

TABLE 36.1 Health professional visits in past year
 36.2 Health professional visits in past year-by sex

Oral Health

TABLE 37.1 Had dental visit in past year
 (Year 2000 Objective 13.14)
 37.2 Had dental visit in past year-by sex

TABLE 38.1 Loss of permanent teeth
 (Year 2000 Objective 13.3)
 38.2 Loss of permanent teeth-by sex

Sexual Behavior

TABLE 39.1 Condom effectiveness, age 18-49
 39.2 Condom effectiveness, age 18-49 -by sex

SECTION III

PREVALENCE REPORT FOR WOMEN'S HEALTH ISSUES

Mammography

TABLE 40.1 Ever had a mammogram, women age 18 and older
 40.2 Ever had a mammogram, women age 18 and older,
 change 1996-1997

TABLE 41.1 Ever had a mammogram, women age 40 and older
 41.2 Ever had a mammogram, women age 40 and older,
 change 1996-1997

TABLE 42.1 Had a mammogram within the past two years, women
 age 50 and older
 42.2 Had a mammogram within the past two years, women
 age 50 and older, change 1996-1997

Breast Exam

TABLE 43.1 Ever had breast exam, women age 18 and older
 43.2 Ever had breast exam, women age 18 and older,
 change 1996-1997

TABLE 44.1 Ever had breast exam, women age 40 and older
44.2 Ever had breast exam, women age 40 and older,
 change 1996-1997

TABLE 45.1 Had breast exam within the past two years, women
 age 50 and older
45.2 Had breast exam within the past two years, women
 age 50 and older, change 1996-1997

Mammogram and Breast Exam

TABLE 46.1 Ever had a mammogram and breast exam, women
 age 40 and older
(Approximates Year 2000 Objective 16.11)
46.2 Ever had a mammogram and breast exam, women
 age 40 and older, change 1996-1997

TABLE 47.1 Ever had a mammogram and breast exam within the
 past two years, women age 50 and older
(Approximates Year 2000 Objective 16.11)
47.2 Had a mammogram and breast exam within the past
 two years, women age 50 and older, change 1996-1997

Cervical Cancer

TABLE 48.1 Ever had pap smear, women age 18 and older
(Year 2000 Objective 16.12)
48.2 Ever had pap smear, women age 18 and older, change 1996-1997

TABLE 49.1 Ever had pap smear, women with uterine cervix
49.2 Ever had pap smear, women with uterine cervix,
 change 1996-1997

TABLE 50.1 Had pap smear within the past three years, women
 with uterine cervix
50.2 Had pap smear within the past three years, women
 with uterine cervix, change 1996-1997

OVERVIEW OF THE 1997 SUMMARY REPORT

The report is divided into three sections:

Section I displays tables for selected variables in the core. In Section I, there are three tables for each risk factor or preventive measure unless otherwise noted. Estimated prevalence and relevant statistics are provided for the total population sampled and by sex. Prevalence estimates for both 1996 and 1997, or where appropriate 1995 and 1997, and the percentage point difference between the estimates are given for variables in the core.

Section II shows tables for selected variables in modules used by at least 13 states. Section II contains two tables for each risk factor or preventive measure of interest in the modules. These tables show the estimated prevalence and relevant statistics for the total sampled population and both sex groups who responded to selected questions in the optional modules.

Section III contains tables for preventive health measures that are related to women's health issues. Estimated prevalence and relevant statistics are provided for the total population sampled. Prevalence estimates for both 1996 and 1997 and the percentage point difference between them are given.

The next section describes each of the tables in detail.

EXPLANATION OF TABLES AND FIGURES

Three sample tables (Tables A-C) from Section I will be used as examples to explain the information contained in this report.

Each table describes a behavioral risk factor or preventive health measure from the Behavioral Risk Factor Surveillance System (BRFSS) instrument.

Table A shows the estimated prevalence and relevant statistics for each participant. (Participants refer to all states, and the District of Columbia and territories of the U.S.). The sample table shows the estimated prevalence and relevant statistics for "no health insurance" in the population surveyed by each BRFSS participant. This prevalence estimate excludes respondents with unknown information on insurance coverage.

The columns in Table A provide the following information:

<u>COLUMN HEADING</u>	<u>CONTENTS</u>
<u>Participant</u>	Name of the participant. Participants refers to all states including the District of Columbia and territories of the United States.
<u>Sample size</u> specific BRFSS These numbers exclude don't know", those who refused had no response coded (missing or	The number of individuals who responded to the question. These are unweighted numbers. individuals who reported "not sure, to answer the question, or skipped).
<u>Percent/Mean</u> number) of interest. This population to estimate the true	An estimate of the percent (or prevalence or mean residents who report the behavioral risk factor of is weighted to the participant's prevalence or mean in the population.
<u>Standard error</u> the	A measure of the amount of variability that exists in calculation of the prevalence estimate.
<u>95% confidence interval</u> estimate 95% population is performed.	A range which contains the true population prevalence of the time, if repeated sampling of the population is performed.
<u>SUMMARY STATISTICS</u> are given at the bottom of the table:	
<u>Number of participants</u> question(s) regarding	The total number of participants who asked the respective "risk factor".
<u>Median</u>	The middle value of all estimated prevalences.
<u>Range</u> among all	The lowest prevalence and the highest prevalence participants.

For example, in Table A, there were 2171 individuals in the state of Alabama who responded to the "no health insurance" question. Of those responding, the estimated prevalence of Alabama's population who reported that they had no health care plan is 14.4 percent. This prevalence estimate has a standard error of 0.9. The 95% confidence interval for this estimate is (12.7, 16.1). Ninety-five percent (95%) of confidence intervals generated under repeated sampling will contain the true population prevalence of Alabama's population who "had no health care plan".

TABLE A 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2171	14.4	0.9	(12.7, 16.1)
ALASKA	1536	21.3	1.6	(18.2, 24.4)
ARIZONA	1902	14.7	1.2	(12.4, 16.9)
ARKANSAS	1792	17.3	1.2	(15.0, 19.6)
CALIFORNIA	4058	19.3	0.8	(17.8, 20.8)
COLORADO	1824	11.9	0.9	(10.0, 13.7)
CONNECTICUT	2246	9.4	0.9	(7.8, 11.1)
DELAWARE	2554	10.8	0.8	(9.2, 12.3)
DISTRICT OF CO	1504	11.6	1.0	(9.6, 13.6)
FLORIDA	3487	17.6	0.8	(16.1, 19.1)
GEORGIA	2303	12.0	0.9	(10.3, 13.7)
HAWAII	2166	6.4	0.7	(5.1, 7.7)
IDAHO	4911	16.6	0.8	(15.1, 18.1)
ILLINOIS	2883	11.3	0.7	(10.0, 12.7)
INDIANA	2365	12.4	0.8	(10.8, 14.0)
IOWA	3596	9.6	0.6	(8.4, 10.7)
KANSAS	2000	9.5	0.8	(8.0, 10.9)
KENTUCKY	3607	13.8	0.7	(12.4, 15.2)
LOUISIANA	1652	20.4	1.2	(18.0, 22.8)
MAINE	1699	12.0	0.9	(10.3, 13.8)
MARYLAND	4566	10.2	0.6	(9.0, 11.4)
MASSACHUSETTS	1765	9.3	0.8	(7.6, 10.9)
MICHIGAN	2561	9.9	0.7	(8.6, 11.3)
MINNESOTA	4808	8.4	0.5	(7.5, 9.3)
MISSISSIPPI	1595	15.2	1.1	(13.0, 17.4)
MISSOURI	1848	12.2	1.0	(10.3, 14.1)
MONTANA	1800	14.6	0.9	(12.8, 16.4)
NEBRASKA	2692	7.6	0.6	(6.4, 8.8)
NEVADA	2496	14.1	1.4	(11.3, 16.8)
NEW HAMPSHIRE	1511	10.3	1.0	(8.4, 12.2)
NEW JERSEY	2658	11.7	0.8	(10.1, 13.2)
NEW MEXICO	1807	22.2	1.2	(19.9, 24.5)
NEW YORK	3397	14.0	0.7	(12.6, 15.4)
NORTH CAROLINA	3624	14.7	0.7	(13.3, 16.1)
NORTH DAKOTA	1799	11.7	0.9	(9.9, 13.5)
OHIO	3164	10.5	0.7	(9.0, 11.9)
OKLAHOMA	1880	17.0	1.1	(14.9, 19.2)
OREGON	3313	12.5	0.7	(11.1, 14.0)
PENNSYLVANIA	3595	9.4	0.6	(8.3, 10.6)
RHODE ISLAND	1836	10.7	0.9	(8.9, 12.5)
SOUTH CAROLINA	2150	15.0	1.0	(13.1, 17.0)
SOUTH DAKOTA	2196	13.2	0.9	(11.5, 14.9)
TENNESSEE	2973	11.7	0.7	(10.3, 13.2)
TEXAS	2486	24.2	1.0	(22.3, 26.2)
UTAH	2861	11.2	0.8	(9.6, 12.7)
VERMONT	3177	14.4	0.8	(12.9, 15.9)
VIRGINIA	3512	11.3	0.7	(9.9, 12.6)
WASHINGTON	3598	10.9	0.6	(9.7, 12.2)
WEST VIRGINIA	2429	18.1	0.9	(16.3, 19.9)
WISCONSIN	2243	9.3	0.9	(7.6, 11.0)
WYOMING	2406	19.0	1.2	(16.6, 21.3)
PUERTO RICO	2260	10.3	0.8	(8.8, 11.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 12.0 RANGE = 6.4-24.2

* HAVING NO HEALTH CARE PLAN

Table B shows the estimated prevalence and relevant statistics by sex for each state participant. This example table shows the estimated prevalence and relevant statistics for "no health insurance" among both sex groups. The same statistics are displayed in Table B as in Table A, except that Table B shows values for males and females separately. An explanation of the terms was provided earlier. This side by side display of the tables permits each participant to compare estimates for males and females in their respective state.

TABLE B 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE* - BY SEX
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	810	11.5	1.3	(9.0, 14.0)	*	1361	17.0	1.2	(14.6, 19.3)	*
ALASKA	713	21.6	2.4	(17.0, 26.2)	*	823	20.9	2.1	(16.8, 25.1)	*
ARIZONA	889	15.9	1.8	(12.5, 19.4)	*	1013	13.4	1.6	(10.4, 16.5)	*
ARKANSAS	675	18.0	2.0	(14.0, 21.9)	*	1117	16.7	1.3	(14.2, 19.2)	*
CALIFORNIA	1757	22.3	1.2	(19.9, 24.8)	*	2301	16.4	0.9	(14.5, 18.2)	*
COLORADO	817	13.7	1.4	(10.9, 16.5)	*	1007	10.2	1.2	(7.8, 12.5)	*
CONNECTICUT	887	10.7	1.5	(7.8, 13.6)	*	1359	8.3	0.9	(6.5, 10.1)	*
DELAWARE	1012	13.0	1.3	(10.4, 15.5)	*	1542	8.7	0.9	(7.0, 10.5)	*
DISTRICT OF CO	572	13.4	1.7	(10.1, 16.7)	*	932	10.0	1.1	(7.8, 12.3)	*
FLORIDA	1432	17.8	1.2	(15.5, 20.1)	*	2055	17.3	1.0	(15.4, 19.3)	*
GEORGIA	959	12.9	1.4	(10.1, 15.7)	*	1344	11.2	1.1	(9.1, 13.3)	*
HAWAII	946	6.7	0.9	(4.9, 8.4)	*	1220	6.2	0.9	(4.4, 8.0)	*
IDAHO	2051	17.6	1.2	(15.3, 19.9)	*	2860	15.7	1.0	(13.8, 17.7)	*
ILLINOIS	1162	10.8	1.1	(8.7, 12.8)	*	1721	11.9	0.9	(10.1, 13.7)	*
INDIANA	1064	13.1	1.3	(10.5, 15.6)	*	1301	11.8	1.0	(9.8, 13.7)	*
IOWA	1476	10.4	0.9	(8.6, 12.2)	*	2120	8.8	0.7	(7.4, 10.2)	*
KANSAS	785	9.8	1.2	(7.4, 12.2)	*	1215	9.1	0.9	(7.4, 10.8)	*
KENTUCKY	1323	13.8	1.1	(11.6, 15.9)	*	2284	13.9	0.9	(12.1, 15.6)	*
LOUISIANA	626	21.2	2.0	(17.3, 25.2)	*	1026	19.7	1.5	(16.8, 22.6)	*
MAINE	751	13.4	1.4	(10.6, 16.1)	*	948	10.8	1.1	(8.6, 13.0)	*
MARYLAND	1954	10.5	1.0	(8.6, 12.4)	*	2612	10.0	0.8	(8.5, 11.5)	*
MASSACHUSETTS	721	11.1	1.4	(8.4, 13.8)	*	1044	7.6	0.9	(5.7, 9.4)	*
MICHIGAN	1067	11.8	1.1	(9.6, 14.0)	*	1494	8.2	0.8	(6.7, 9.7)	*
MINNESOTA	2201	8.9	0.7	(7.6, 10.2)	*	2607	7.9	0.6	(6.7, 9.1)	*
MISSISSIPPI	590	15.8	1.9	(12.2, 19.5)	*	1005	14.6	1.3	(12.0, 17.2)	*
MISSOURI	698	14.2	1.7	(10.9, 17.4)	*	1150	10.4	1.0	(8.3, 12.4)	*
MONTANA	757	14.4	1.4	(11.7, 17.1)	*	1043	14.7	1.2	(12.4, 17.1)	*
NEBRASKA	1046	7.8	1.0	(5.8, 9.7)	*	1646	7.4	0.8	(5.8, 8.9)	*
NEVADA	1159	13.9	2.2	(9.6, 18.3)	*	1337	14.2	1.7	(10.9, 17.5)	*
NEW HAMPSHIRE	602	10.9	1.6	(7.8, 13.9)	*	909	9.8	1.1	(7.7, 11.9)	*
NEW JERSEY	1057	12.9	1.3	(10.2, 15.5)	*	1601	10.6	0.9	(8.8, 12.4)	*
NEW MEXICO	753	20.8	1.7	(17.4, 24.1)	*	1054	23.6	1.6	(20.5, 26.6)	*
NEW YORK	1352	15.9	1.2	(13.6, 18.2)	*	2045	12.3	0.9	(10.6, 14.0)	*
NORTH CAROLINA	1434	16.3	1.1	(14.1, 18.5)	*	2190	13.3	0.9	(11.6, 14.9)	*
NORTH DAKOTA	799	12.3	1.4	(9.6, 14.9)	*	1000	11.2	1.1	(9.0, 13.4)	*
OHIO	1229	10.8	1.1	(8.6, 13.0)	*	1935	10.2	0.9	(8.3, 12.0)	*
OKLAHOMA	827	15.3	1.7	(11.9, 18.6)	*	1053	18.6	1.5	(15.7, 21.5)	*
OREGON	1398	14.8	1.3	(12.3, 17.2)	*	1915	10.5	0.8	(8.9, 12.1)	*
PENNSYLVANIA	1439	10.8	0.9	(8.9, 12.6)	*	2156	8.2	0.7	(6.9, 9.6)	*
RHODE ISLAND	753	11.5	1.3	(8.9, 14.2)	*	1083	9.9	1.3	(7.4, 12.4)	*
SOUTH CAROLINA	819	15.9	1.6	(12.8, 18.9)	*	1331	14.3	1.2	(12.0, 16.6)	*
SOUTH DAKOTA	953	13.4	1.3	(10.8, 16.0)	*	1243	13.0	1.1	(10.8, 15.2)	*
TENNESSEE	1042	13.6	1.2	(11.2, 16.0)	*	1931	10.1	0.8	(8.5, 11.6)	*
TEXAS	1018	25.8	1.6	(22.7, 28.8)	*	1468	22.8	1.2	(20.3, 25.2)	*
UTAH	1275	12.0	1.1	(9.8, 14.2)	*	1586	10.3	1.1	(8.2, 12.4)	*
VERMONT	1387	16.4	1.2	(14.0, 18.7)	*	1790	12.5	0.9	(10.7, 14.3)	*
VIRGINIA	1409	10.7	1.1	(8.6, 12.7)	*	2103	11.9	0.9	(10.1, 13.7)	*
WASHINGTON	1544	11.7	0.9	(9.9, 13.5)	*	2054	10.2	0.9	(8.5, 11.9)	*
WEST VIRGINIA	959	18.6	1.4	(15.8, 21.4)	*	1470	17.7	1.2	(15.4, 20.0)	*
WISCONSIN	991	10.8	1.4	(8.1, 13.5)	*	1252	7.9	1.1	(5.9, 10.0)	*
WYOMING	1013	20.7	2.0	(16.8, 24.6)	*	1393	17.3	1.1	(15.1, 19.4)	*
PUERTO RICO	936	11.8	1.2	(9.5, 14.1)	*	1324	8.9	1.0	(7.0, 10.8)	*
SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS	=	52			NUMBER OF PARTICIPANTS	=	52		
	MEDIAN	=	13.4			MEDIAN	=	11.2		
	RANGE	=	6.7-25.8			RANGE	=	6.2-23.6		

* HAVING NO HEALTH CARE PLAN

Table C shows the percent difference between the prevalence estimates, by participant, for the 1996 and 1997 survey years for having no health insurance. A negative change indicates a decrease in prevalence from 1996 to 1997, while a positive change indicates an increase in prevalence between the two years. This difference gives state health officials a quick method to examine changes that may have occurred in their respective state. Some of these changes may have resulted from legislative actions (or inactions) in the state, programs funded, or programs that came into existence because of results that you may have shown from the BRFSS. Since changes between two years may be attributable to many factors, conclusions drawn from this table should be interpreted with extreme caution.

Sample Size. Sample sizes vary by question. Some estimates may be based on responses from less than 50 respondents. Interpreting estimates that are based on a small number of respondents can mislead the reader into believing that a given finding is much more precise than it actually is. The BRFSS recommends not interpreting percentages where the denominator is based upon fewer than 50 unweighted respondents (sample size)

TABLE C 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

PREVALENCE DIFFERENCES FROM 1996 TO 1997

NO HEALTH INSURANCE*

DENOMINATOR IS PERSONS AGE 18 AND OVER

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	16.0	14.4	-1.6
ALASKA	20.8	21.3	0.5
ARIZONA	17.0	14.7	-2.4
ARKANSAS	17.9	17.3	-0.6
CALIFORNIA	19.5	19.3	-0.2
COLORADO	13.5	11.9	-1.6
CONNECTICUT	9.3	9.4	0.1
DELAWARE	10.8	10.8	-0.0
DISTRICT OF CO	13.4	11.6	-1.8
FLORIDA	17.6	17.6	0.0
GEORGIA	9.8	12.0	2.3
HAWAII	8.1	6.4	-1.7
IDAHO	15.0	16.6	1.6
ILLINOIS	11.8	11.3	-0.5
INDIANA	11.4	12.4	0.9
IOWA	9.2	9.6	0.4
KANSAS	10.1	9.5	-0.6
KENTUCKY	14.4	13.8	-0.6
LOUISIANA	22.8	20.4	-2.4
MAINE	14.2	12.0	-2.2
MARYLAND	9.7	10.2	0.5
MASSACHUSETTS	10.2	9.3	-0.9
MICHIGAN	9.6	9.9	0.4
MINNESOTA	7.2	8.4	1.2
MISSISSIPPI	14.2	15.2	0.9
MISSOURI	12.9	12.2	-0.8
MONTANA	15.6	14.6	-1.0
NEBRASKA	8.6	7.6	-1.0
NEVADA	16.4	14.1	-2.4
NEW HAMPSHIRE	11.9	10.3	-1.6
NEW JERSEY	11.1	11.7	0.5
NEW MEXICO	21.0	22.2	1.2
NEW YORK	12.4	14.0	1.6
NORTH CAROLINA	12.9	14.7	1.8
NORTH DAKOTA	11.8	11.7	-0.0
OHIO	11.4	10.5	-0.9
OKLAHOMA	18.5	17.0	-1.5
OREGON	13.2	12.5	-0.7
PENNSYLVANIA	11.1	9.4	-1.7
RHODE ISLAND	10.5	10.7	0.2
SOUTH CAROLINA	16.3	15.0	-1.3
SOUTH DAKOTA	10.1	13.2	3.1
TENNESSEE	12.0	11.7	-0.3
TEXAS	24.6	24.2	-0.4
UTAH	12.2	11.2	-1.0
VERMONT	13.0	14.4	1.4
VIRGINIA	14.6	11.3	-3.3
WASHINGTON	11.9	10.9	-1.0
WEST VIRGINIA	17.7	18.1	0.4
WISCONSIN	9.3	9.3	0.0
WYOMING	18.3	19.0	0.6
PUERTO RICO	13.6	10.3	-3.3

* HAVING NO HEALTH CARE PLAN

N/A - NOT AVAILABLE

DEFINITIONS OF BRFSS RISK FACTORS AND PREVENTIVE HEALTH MEASURES

ACTIVITY LIMITATIONS

ANY DISABILITY - Respondents, age 18-64, who report that an impairment or health problem is responsible for any of the following: (1) keeping them from working at a job or business, (2) limiting the kind or amount of work they can do, (3) keeping them from doing housework, or (4) limiting the kind or amount of housework they can do. This measures Year 2000 Objective 17.2 - Target < 8%.

ANY LIMITATIONS (1) - Respondents, age 18-64, who report they are limited in any way in any activities because of any impairment or health problem. This measures Year 2000 Objective 17.2 - Target < 8%.

AIDS KNOWLEDGE/ATTITUDES

HIV BLOOD TEST - Respondents, age 18-64, who have ever had a blood test for HIV infection.

CHANCES OF GETTING HIV - Respondents, age 18-64, who believe that their chances of getting infected with HIV, the virus that causes AIDS are medium or high.

ENCOURAGE TEEN CONDOM USE - Respondents, age 18-64, who would encourage sexually active teenagers to use a condom.

BREAST EXAM

EVER HAD A BREAST EXAM - Female respondents, age 18 and older, who report that they have ever had a breast exam by a doctor.

EVER HAD A BREAST EXAM - Female respondents, age 40 and older, who report that they have ever had a breast exam by a doctor.

HAD A BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they had a breast exam within the past two years.

CERVICAL CANCER

EVER HAD A PAP SMEAR - Female respondents, age 18 and older who report that they have ever had a pap smear. This measures Year 2000 Objective 16.12. - Target > 95%.

EVER HAD A PAP SMEAR, WITH UTERINE CERVIX - Female respondents, with uterine cervix, age 18 and older, who report that they have ever had a pap smear.

HAD A PAP SMEAR WITHIN 3 YEARS - Female respondents, with uterine cervix, age 18 and older, who report that they had a pap smear within the last three years.

CHOLESTEROL SCREENING/AWARENESS

CHOLESTEROL SCREENING (1) - Respondents who have ever had their cholesterol checked.

CHOLESTEROL SCREENING (2) - Respondents who have had their cholesterol checked within the past five years. Year 2000 Objective 15.14 - Target $> 75\%$.

CHOLESTEROL AWARENESS - Respondents who report that their cholesterol has been checked and they have been told their cholesterol is high by health professional.

CLINICAL BREAST EXAM See BREAST EXAM

COLORECTAL CANCER SCREENING

PROCTOSCOPIC EXAM - Respondents, age 50 and older, who report the

BLOOD STOOL TEST - Respondents , age 50 and older, who did not report

DIABETES

AGE AT ONSET - Mean age at onset of diabetes: age when respondent was told that they have diabetes, among those who report they were told by a doctor that they have diabetes.

DIABETES AWARENESS - Respondents who report they were told by a doctor that they have diabetes.

NOW TAKING INSULIN - Respondents who are now taking insulin, among those who report they were told by a doctor that they have diabetes.

NUMBER OF TIMES SEEN HEALTH PROFESSIONAL - Mean number of times respondents report they have seen a doctor, nurse, or other health professional for their diabetes in the past year, among those who report they were told by a doctor that they have diabetes.

DRINKING STATUS

CURRENT DRINKERS - Respondents who report that they have had any alcoholic beverages in the past month.

BINGE DRINKING (also called ACUTE DRINKING) - Respondents who report having five or more drinks on an occasion, one or more times in the past month.

CHRONIC DRINKING - Respondents who report an average of two or more drinks per day i.e., 60 or more alcoholic drinks a month. This is based on the total number of drinks per month.

DRINKING and DRIVING - Respondents who report having driven after having too much to drink, one or more times in the past month.

HEALTH INSURANCE

NO HEALTH INSURANCE - Respondents age 18 and older who report they have no health care plan.

NO HEALTH INSURANCE - Respondents age 18-64 years who report they have no health care plan.

HEALTH STATUS

ACTIVITIES LIMITED - Mean number of days in past 30 days activities were limited as reported by respondents.

MENTAL HEALTH NOT GOOD - Mean number of days in past 30 days mental health was not good as reported by respondents.

PHYSICAL HEALTH NOT GOOD - Mean number of days in past 30 days physical health was not good as reported by respondents.

SELF REPORTED HEALTH STATUS - Respondents who report having a general health status of fair or poor.

HYPERTENSION

HYPERTENSION [also called HYPERTENSION AWARENESS] - Respondents who report that their blood pressure has been checked and they have been told they have high blood pressure.

HYPERTENSION SCREENING - Respondents who report they have had their blood pressure checked within the past two years.

IMMUNIZATION

FLU SHOT - Respondents, age 65 and older, who report that they have had a flu shot within past 12 months. This measures Year 2000 Objective 20.11 - Target > 80%.

PNEUMONIA VACCINATION - Respondents, age 65 and older, who report they have ever had a pneumonia vaccination. This measures Year 2000 Objective 20.11 - Target > 80%.

INJURY CONTROL AND CHILD SAFETY

USE OF CHILD SAFETY BELTS FOR OLDEST CHILD AGE 5-15 - Respondents who report their oldest child is between age 5 and 15 and "always" uses a safety belt when riding in a car. This approximates Year 2000 Objective 9.12 - Target > 85%.

USE OF CHILD SAFETY SEATS FOR OLDEST CHILD UNDER 5 - Respondents who report their oldest child is under age 5 and "always" uses a car safety seat when riding in a car. This approximates Year 2000 Objective 9.12a - Target > 95%.

USE OF BICYCLE HELMET - Respondents who report their oldest child is under age 15 and child age 5-15 always wears a bicycle helmet when riding a bicycle.

MAMMOGRAPHY

EVER HAD A MAMMOGRAM (1) - Female respondents age 18 and older, who report that they have ever had a mammogram.

EVER HAD A MAMMOGRAM (2) - Female respondents, age 40 and older, who report that they have ever had a mammogram.

HAD A MAMMOGRAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they had a mammogram within the past two years.

MAMMOGRAPHY AND BREAST EXAM

MAMMOGRAM & BREAST EXAM - Female respondents, age 40 and older, who report that they have ever had a mammogram and a breast exam. This measures Year 2000 Objective 16.11 - Target > 80%.

MAMMOGRAM & BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they have had a mammogram and a breast exam within the last two years. This measures Year 2000 Objective 16.11 - Target > 60%.

OVERWEIGHT

OVERWEIGHT: BY MEDIAN - Respondents at or above 120% of their ideal weight. Ideal weight is defined as the mid-value of a median frame person from the 1959 Metropolitan Life Insurance Corporation height-weight tables.

OVERWEIGHT: BY BODY MASS INDEX - Females with body mass index(BMI) Year 2000 Objective 2.3 - Target \leq 25 of overweight.

ORAL HEALTH

DENTAL VISIT - Respondents who reported visiting a dentist or dental

TOOTH LOSS - Respondents who report a loss of 6 or more permanent

PAP SMEAR See CERVICAL CANCER

PROCTOSCOPIC EXAM see COLORECTAL CANCER SCREENING

SAFETY BELTS

LACK OF SAFETY BELT USAGE - Respondents reporting they "nearly always", "sometimes", "seldom", or "never" use safety belts when occupants of a car (i.e., do not always use a safety belt). This measures Year 2000 Objective 9.12 - Target \geq 85%.

SEXUAL BEHAVIOR

CONDOM EFFECTIVENESS - Respondents, age 18-49, who believe that condo

SMOKELESS TOBACCO

EVER USED - Respondents who have ever used or tried smokeless tobacco.

CURRENT USE - Respondents who report that they currently use smokeless tobacco.

SMOKING STATUS

EVER SMOKED - Respondents who have ever smoked 100 cigarettes in their lifetime.

CURRENT SMOKER - Respondents who have ever smoked 100 cigarettes in their lifetime and reported smoking every

day or some days. This measures Year 2000 Objective 3.4
- Target < 15%.

LIMITATIONS OF THE DATA

This report consolidates the most current data on selected risk factors and preventive health measures for states and the District of Columbia. The information was obtained from the data files of individual states. Therefore, while all participants use telephone surveys to collect the data, the data in this report vary somewhat with respect to data collection, sampling design and sample size.

Some states conduct interviews using a paper questionnaire, but most use a computer-assisted telephone interviewing (CATI) software program. All states use probability-based methods to select their sample. For data year 1997, twenty-nine states used the Mitofsky-Waksberg method of sampling, seventeen states used a disproportionate stratified sample (DSS) design, and six states used a variety of other designs. **The BRFSS standard for sample designs is that sample records must be justifiable as a probability sample of all households with telephones in the state.** All but Alaska, California, Nevada and Texas met this criterion in 1996. The deviations from the standard by these states may reduce the comparability of their data with those of other areas. Although the effects of the deviations have not been extensively investigated, departure from the standard may produce bias in prevalence estimates and other survey results, as well as constrict confidence intervals. The sampling practices of the four states are highlighted below:

Alaska used a stratified directory/random sample combination where the sample design is made up of 3 random digit-dialed strata and 4 listed strata. Phone numbers in the listed strata are selected from a list of putative household numbers. Forty-five percent of the completed interviews come from the listed strata. Alaska's sampling frame for the listed strata contains an estimated 50% of all household numbers in these strata.

California used a truncated list-assisted design with sampling from 3+ blocks only. Noncoverage was further affected to an unknown degree by substitution for sample numbers used by any client of the commercial sample provider in the preceding six months. Probabilities of selection varied in proportion to the number of listed household numbers in the 100 blocks associated with a number, the number of known business numbers preceding a number, the number of preceding numbers selected by any client of the commercial sample provider in the preceding six months, and the measure of size used for each county, which could vary each

time a sample was drawn. Because of the unavailability of appropriate information, none of these factors is accounted for in the weights.

Nevada used a truncated list-assisted design with sampling from 5+ blocks only. Noncoverage was further affected to an unknown degree by substitution for sample numbers used by any client of the commercial sample provider in the preceding six months. Probabilities of selection varied in proportion to the number of listed household numbers in the 100 blocks associated with a number, the number of known business numbers preceding a number, the number of preceding numbers selected by any client of the commercial sample provider in the preceding six months, and the measure of size used for each county, which could vary each time a sample was drawn. Because of the unavailability of appropriate information, only the number of listed household numbers in the 100 blocks associated with a number is accounted for in the weights.

Texas used a truncated list-assisted sample which included only phone numbers from 100 blocks with one or more listed household numbers. Such hundred blocks are estimated to contain 97.1% of all household numbers.

California modified the wording of mammography, Papanicolaou (PAP) smear, and chronic alcohol use and are therefore excluded from those tables in the report. California also asked the HIV/AIDS section questions to persons 18-45 years of age rather than those 18-65 years of age, and are therefore excluded from tables reporting data on this topic.

Although states collect data monthly, the number of interviews range from 1500 in New Hampshire to 4896 in Idaho. Estimates for some states are therefore based upon smaller sample sizes, and for rare events (e.g. diabetes), may yield unstable estimates.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. People may not remember essential information, a question may not mean the same thing to different respondents, and some individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their impact on the data. The user must make his or her own evaluation on the data. Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. When the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the conditions described.

TABLE 1.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2171	19.0	1.0	(17.1, 20.9)
ALASKA	1538	11.0	1.1	(8.8, 13.2)
ARIZONA	1902	11.4	1.0	(9.4, 13.3)
ARKANSAS	1787	20.9	1.1	(18.8, 23.0)
CALIFORNIA	4059	15.3	0.7	(14.0, 16.6)
COLORADO	1825	10.0	0.7	(8.5, 11.4)
CONNECTICUT	2247	10.2	0.7	(8.8, 11.6)
DELAWARE	2549	13.8	0.8	(12.2, 15.3)
DISTRICT OF CO	1502	9.7	0.9	(7.9, 11.4)
FLORIDA	3484	16.4	0.7	(15.0, 17.7)
GEORGIA	2303	11.4	0.8	(9.8, 12.9)
HAWAII	2162	14.0	0.9	(12.2, 15.8)
IDAHO	4911	12.4	0.6	(11.3, 13.5)
ILLINOIS	2882	14.0	0.8	(12.5, 15.5)
INDIANA	2366	14.1	0.8	(12.5, 15.7)
IOWA	3593	11.3	0.6	(10.2, 12.5)
KANSAS	1997	10.7	0.7	(9.3, 12.1)
KENTUCKY	3611	21.8	0.8	(20.3, 23.3)
LOUISIANA	1647	15.3	1.0	(13.4, 17.2)
MAINE	1698	13.9	0.9	(12.1, 15.6)
MARYLAND	4570	11.9	0.6	(10.8, 13.1)
MASSACHUSETTS	1764	12.0	0.9	(10.1, 13.8)
MICHIGAN	2562	12.7	0.7	(11.3, 14.2)
MINNESOTA	4790	9.5	0.5	(8.6, 10.4)
MISSISSIPPI	1597	21.9	1.2	(19.7, 24.2)
MISSOURI	1842	15.0	1.0	(13.2, 16.9)
MONTANA	1798	10.1	0.8	(8.6, 11.6)
NEBRASKA	2696	11.4	0.7	(10.1, 12.8)
NEVADA	2500	14.4	1.4	(11.7, 17.2)
NEW HAMPSHIRE	1512	9.6	0.8	(8.0, 11.2)
NEW JERSEY	2663	13.8	0.8	(12.3, 15.4)
NEW MEXICO	1812	14.4	0.9	(12.5, 16.2)
NEW YORK	3400	13.3	0.7	(12.0, 14.6)
NORTH CAROLINA	3626	16.1	0.7	(14.7, 17.4)
NORTH DAKOTA	1801	12.1	0.8	(10.5, 13.7)
OHIO	3158	13.7	0.8	(12.1, 15.3)
OKLAHOMA	1881	13.4	0.9	(11.7, 15.2)
OREGON	3317	12.8	0.7	(11.4, 14.2)
PENNSYLVANIA	3595	13.1	0.6	(11.9, 14.4)
RHODE ISLAND	1835	12.6	0.9	(10.9, 14.4)
SOUTH CAROLINA	2154	16.8	0.9	(15.0, 18.6)
SOUTH DAKOTA	2199	11.3	0.8	(9.8, 12.8)
TENNESSEE	2969	18.1	0.8	(16.6, 19.6)
TEXAS	2500	16.9	0.8	(15.2, 18.5)
UTAH	2863	11.2	0.8	(9.6, 12.7)
VERMONT	3187	11.4	0.7	(10.1, 12.8)
VIRGINIA	3505	11.2	0.7	(9.8, 12.6)
WASHINGTON	3596	10.6	0.6	(9.5, 11.7)
WEST VIRGINIA	2435	21.7	0.9	(20.0, 23.5)
WISCONSIN	2245	9.8	0.7	(8.3, 11.2)
WYOMING	2409	11.3	1.0	(9.3, 13.4)
PUERTO RICO	2252	35.0	1.1	(32.8, 37.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 13.0 RANGE = 9.5-35.0

* HEALTH STATUS IS FAIR OR POOR

August 14, 1998

TABLE 1.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	811	16.3	1.5	(13.4, 19.2)	*	1360	21.3	1.2	(18.9, 23.8)	
ALASKA	714	8.5	1.4	(5.8, 11.2)	*	824	13.7	1.8	(10.2, 17.2)	
ARIZONA	890	10.5	1.4	(7.7, 13.2)	*	1012	12.2	1.4	(9.4, 14.9)	
ARKANSAS	672	19.9	1.8	(16.4, 23.4)	*	1115	21.8	1.3	(19.2, 24.4)	
CALIFORNIA	1759	14.7	1.0	(12.7, 16.6)	*	2300	15.9	0.9	(14.2, 17.6)	
COLORADO	818	9.6	1.1	(7.5, 11.8)	*	1007	10.3	1.0	(8.3, 12.3)	
CONNECTICUT	889	9.4	1.1	(7.3, 11.5)	*	1358	11.0	0.9	(9.2, 12.8)	
DELAWARE	1013	13.2	1.2	(10.8, 15.6)	*	1536	14.3	1.0	(12.3, 16.2)	
DISTRICT OF CO	572	9.2	1.3	(6.6, 11.7)	*	930	10.1	1.2	(7.8, 12.5)	
FLORIDA	1431	15.7	1.1	(13.6, 17.7)	*	2053	17.0	0.9	(15.1, 18.8)	
GEORGIA	959	9.1	1.0	(7.1, 11.1)	*	1344	13.5	1.1	(11.2, 15.7)	
HAWAII	947	13.7	1.3	(11.1, 16.2)	*	1215	14.4	1.3	(11.9, 16.8)	
IDAHO	2056	12.3	0.8	(10.7, 14.0)	*	2855	12.5	0.8	(10.9, 14.0)	
ILLINOIS	1161	13.5	1.1	(11.2, 15.7)	*	1721	14.5	1.0	(12.6, 16.4)	
INDIANA	1066	13.1	1.1	(10.9, 15.3)	*	1300	15.0	1.1	(12.8, 17.2)	
IOWA	1471	11.2	0.9	(9.4, 13.0)	*	2122	11.5	0.8	(10.0, 13.0)	
KANSAS	782	10.0	1.1	(7.9, 12.2)	*	1215	11.4	1.0	(9.5, 13.2)	
KENTUCKY	1325	21.6	1.2	(19.2, 24.0)	*	2286	22.0	0.9	(20.2, 23.8)	
LOUISIANA	625	14.3	1.5	(11.3, 17.3)	*	1022	16.3	1.2	(13.9, 18.7)	
MAINE	750	13.5	1.4	(10.8, 16.1)	*	948	14.2	1.2	(11.8, 16.6)	
MARYLAND	1960	11.2	0.9	(9.4, 12.9)	*	2610	12.7	0.9	(10.9, 14.4)	
MASSACHUSETTS	723	11.4	1.3	(8.9, 14.0)	*	1041	12.4	1.3	(9.9, 14.9)	
MICHIGAN	1069	12.3	1.1	(10.1, 14.4)	*	1493	13.2	1.0	(11.3, 15.0)	
MINNESOTA	2196	9.3	0.7	(7.9, 10.6)	*	2594	9.6	0.6	(8.4, 10.8)	
MISSISSIPPI	591	20.9	1.8	(17.4, 24.4)	*	1006	22.9	1.5	(19.9, 25.8)	
MISSOURI	696	13.1	1.4	(10.4, 15.7)	*	1146	16.8	1.3	(14.3, 19.3)	
MONTANA	760	8.8	1.1	(6.7, 10.9)	*	1038	11.3	1.1	(9.3, 13.4)	
NEBRASKA	1046	11.9	1.1	(9.7, 14.1)	*	1650	10.9	0.8	(9.3, 12.6)	
NEVADA	1163	16.0	2.4	(11.4, 20.6)	*	1337	12.8	1.5	(9.9, 15.7)	
NEW HAMPSHIRE	603	8.6	1.2	(6.2, 10.9)	*	909	10.6	1.1	(8.5, 12.7)	
NEW JERSEY	1060	13.3	1.2	(10.9, 15.6)	*	1603	14.4	1.0	(12.4, 16.3)	
NEW MEXICO	754	12.8	1.4	(10.1, 15.5)	*	1058	15.8	1.2	(13.4, 18.2)	
NEW YORK	1355	12.3	1.0	(10.3, 14.3)	*	2045	14.2	0.9	(12.5, 15.9)	
NORTH CAROLINA	1437	14.3	1.0	(12.3, 16.2)	*	2189	17.7	0.9	(15.9, 19.5)	
NORTH DAKOTA	800	11.6	1.2	(9.2, 14.0)	*	1001	12.6	1.0	(10.6, 14.6)	
OHIO	1227	12.3	1.2	(9.9, 14.7)	*	1931	14.9	1.1	(12.8, 17.0)	
OKLAHOMA	830	10.0	1.2	(7.6, 12.4)	*	1051	16.6	1.3	(14.1, 19.1)	
OREGON	1400	12.9	1.1	(10.7, 15.0)	*	1917	12.8	0.9	(11.1, 14.5)	
PENNSYLVANIA	1442	13.0	1.0	(11.0, 14.9)	*	2153	13.2	0.8	(11.6, 14.8)	
RHODE ISLAND	755	11.5	1.3	(9.0, 13.9)	*	1080	13.7	1.2	(11.2, 16.1)	
SOUTH CAROLINA	820	15.5	1.4	(12.7, 18.3)	*	1334	18.0	1.2	(15.7, 20.3)	
SOUTH DAKOTA	957	10.7	1.1	(8.5, 12.9)	*	1242	11.9	1.0	(9.9, 13.8)	
TENNESSEE	1041	14.7	1.2	(12.4, 17.1)	*	1928	21.1	1.0	(19.1, 23.1)	
TEXAS	1027	15.9	1.3	(13.5, 18.4)	*	1473	17.8	1.1	(15.6, 20.0)	
UTAH	1276	11.1	1.1	(8.9, 13.3)	*	1587	11.2	1.1	(9.1, 13.3)	
VERMONT	1393	11.3	0.9	(9.4, 13.1)	*	1794	11.6	1.0	(9.6, 13.5)	
VIRGINIA	1414	10.2	1.0	(8.2, 12.1)	*	2091	12.2	1.0	(10.3, 14.2)	
WASHINGTON	1543	10.6	0.9	(8.9, 12.3)	*	2053	10.6	0.8	(9.2, 12.1)	
WEST VIRGINIA	965	20.0	1.4	(17.3, 22.6)	*	1470	23.3	1.2	(21.0, 25.7)	
WISCONSIN	991	9.5	1.1	(7.4, 11.6)	*	1254	10.0	1.0	(8.1, 11.9)	
WYOMING	1019	11.2	1.8	(7.6, 14.7)	*	1390	11.5	1.1	(9.4, 13.5)	
PUERTO RICO	934	28.3	1.6	(25.2, 31.4)	*	1318	40.9	1.5	(38.0, 43.8)	

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 12.3
RANGE = 8.5-28.3

NUMBER OF PARTICIPANTS = 52
MEDIAN = 13.6
RANGE = 9.6-40.9

August 14, 1998

* HEALTH STATUS IS FAIR OR POOR

TABLE 1.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	20.2	19.0	-1.2
ALASKA	8.3	11.0	2.7
ARIZONA	12.1	11.4	-0.8
ARKANSAS	18.9	20.9	2.0
CALIFORNIA	15.5	15.3	-0.1
COLORADO	9.2	10.0	0.7
CONNECTICUT	10.1	10.2	0.1
DELAWARE	13.3	13.8	0.5
DISTRICT OF CO	13.0	9.7	-3.3
FLORIDA	15.4	16.4	0.9
GEORGIA	8.9	11.4	2.5
HAWAII	12.5	14.0	1.5
IDAHO	11.7	12.4	0.7
ILLINOIS	13.7	14.0	0.3
INDIANA	14.1	14.1	0.0
IOWA	11.6	11.3	-0.2
KANSAS	12.2	10.7	-1.5
KENTUCKY	20.9	21.8	0.9
LOUISIANA	16.7	15.3	-1.4
MAINE	13.2	13.9	0.7
MARYLAND	11.8	11.9	0.1
MASSACHUSETTS	12.8	12.0	-0.8
MICHIGAN	12.0	12.7	0.8
MINNESOTA	10.2	9.5	-0.8
MISSISSIPPI	21.1	21.9	0.9
MISSOURI	14.1	15.0	1.0
MONTANA	11.6	10.1	-1.5
NEBRASKA	13.2	11.4	-1.8
NEVADA	11.4	14.4	3.1
NEW HAMPSHIRE	10.4	9.6	-0.8
NEW JERSEY	13.7	13.8	0.1
NEW MEXICO	16.2	14.4	-1.8
NEW YORK	12.7	13.3	0.6
NORTH CAROLINA	16.3	16.1	-0.2
NORTH DAKOTA	12.5	12.1	-0.4
OHIO	12.6	13.7	1.1
OKLAHOMA	14.0	13.4	-0.5
OREGON	12.2	12.8	0.6
PENNSYLVANIA	14.4	13.1	-1.3
RHODE ISLAND	14.8	12.6	-2.2
SOUTH CAROLINA	16.9	16.8	-0.1
SOUTH DAKOTA	11.5	11.3	-0.2
TENNESSEE	19.5	18.1	-1.5
TEXAS	18.4	16.9	-1.5
UTAH	11.3	11.2	-0.1
VERMONT	10.4	11.4	1.1
VIRGINIA	13.1	11.2	-1.9
WASHINGTON	10.9	10.6	-0.2
WEST VIRGINIA	22.8	21.7	-1.0
WISCONSIN	11.5	9.8	-1.7
WYOMING	11.8	11.3	-0.4
PUERTO RICO	37.4	35.0	-2.4

* HEALTH STATUS IS FAIR OR POOR
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 2.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2088	3.8	0.2	(3.4, 4.2)
ALASKA	1509	2.9	0.3	(2.3, 3.4)
ARIZONA	1904	1.1	0.1	(0.8, 1.3)
ARKANSAS	1775	3.4	0.2	(3.0, 3.8)
CALIFORNIA	4048	3.1	0.1	(2.9, 3.4)
COLORADO	1810	2.8	0.2	(2.5, 3.2)
CONNECTICUT	2211	3.1	0.2	(2.6, 3.5)
DELAWARE	2533	3.7	0.2	(3.4, 4.1)
DISTRICT OF CO	1496	1.5	0.1	(1.2, 1.8)
FLORIDA	3437	3.5	0.2	(3.2, 3.8)
GEORGIA	2260	2.2	0.2	(1.9, 2.5)
HAWAII	2129	3.0	0.2	(2.6, 3.4)
IDAHO	4868	3.4	0.1	(3.2, 3.7)
ILLINOIS	2870	3.3	0.2	(3.0, 3.6)
INDIANA	2322	3.4	0.2	(3.0, 3.7)
IOWA	3578	2.9	0.1	(2.7, 3.2)
KANSAS	1980	2.1	0.1	(1.8, 2.3)
KENTUCKY	3602	4.1	0.2	(3.8, 4.4)
LOUISIANA	1615	3.2	0.2	(2.8, 3.6)
MAINE	1678	3.5	0.2	(3.1, 3.9)
MARYLAND	4551	2.4	0.1	(2.2, 2.7)
MASSACHUSETTS	1748	3.3	0.2	(2.9, 3.7)
MICHIGAN	2540	3.3	0.2	(2.9, 3.6)
MINNESOTA	4714	3.0	0.1	(2.8, 3.2)
MISSISSIPPI	1553	3.1	0.2	(2.7, 3.5)
MISSOURI	1821	3.5	0.2	(3.1, 3.9)
MONTANA	1767	2.7	0.2	(2.4, 3.1)
NEBRASKA	2681	2.9	0.2	(2.6, 3.3)
NEVADA	2467	3.5	0.3	(2.9, 4.1)
NEW HAMPSHIRE	1497	3.1	0.2	(2.7, 3.5)
NEW JERSEY	2625	3.0	0.2	(2.7, 3.4)
NEW MEXICO	1799	3.2	0.2	(2.8, 3.6)
NEW YORK	3347	3.1	0.2	(2.8, 3.4)
NORTH CAROLINA	3568	2.9	0.1	(2.6, 3.1)
NORTH DAKOTA	1798	2.9	0.2	(2.5, 3.2)
OHIO	3152	2.6	0.2	(2.3, 2.9)
OKLAHOMA	1872	2.3	0.2	(1.9, 2.6)
OREGON	3275	3.2	0.1	(2.9, 3.5)
PENNSYLVANIA	3541	3.2	0.2	(2.9, 3.5)
RHODE ISLAND	1804	3.0	0.2	(2.7, 3.4)
SOUTH CAROLINA	2135	2.9	0.2	(2.5, 3.2)
SOUTH DAKOTA	2184	2.6	0.2	(2.3, 3.0)
TENNESSEE	2922	3.4	0.2	(3.1, 3.7)
TEXAS	2451	3.2	0.2	(2.8, 3.5)
UTAH	2852	3.6	0.2	(3.2, 3.9)
VERMONT	3134	2.7	0.1	(2.5, 3.0)
VIRGINIA	3406	2.4	0.1	(2.2, 2.7)
WASHINGTON	3527	3.1	0.1	(2.8, 3.3)
WEST VIRGINIA	2418	3.4	0.2	(3.0, 3.7)
WISCONSIN	2224	2.8	0.2	(2.4, 3.1)
WYOMING	2388	2.6	0.2	(2.3, 2.9)
PUERTO RICO	2243	2.8	0.1	(2.6, 3.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.1 RANGE = 1.1- 4.1

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD August 14, 1998

TABLE 2.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	791	3.2	0.3	(2.5, 3.8)	*	1297	4.3	0.3	(3.8, 4.8)	
ALASKA	703	1.9	0.3	(1.4, 2.4)	*	806	3.9	0.5	(2.9, 4.8)	
ARIZONA	891	0.6	0.1	(0.4, 0.9)	*	1013	1.5	0.3	(1.0, 2.0)	
ARKANSAS	672	3.0	0.3	(2.3, 3.7)	*	1103	3.8	0.3	(3.2, 4.3)	
CALIFORNIA	1752	2.8	0.2	(2.4, 3.1)	*	2296	3.5	0.2	(3.2, 3.9)	
COLORADO	812	2.5	0.3	(2.1, 3.0)	*	998	3.1	0.2	(2.6, 3.6)	
CONNECTICUT	876	2.6	0.4	(1.9, 3.3)	*	1335	3.5	0.2	(3.0, 3.9)	
DELAWARE	1004	3.4	0.3	(2.8, 3.9)	*	1529	4.1	0.3	(3.6, 4.6)	
DISTRICT OF CO	570	1.3	0.2	(1.0, 1.7)	*	926	1.7	0.2	(1.3, 2.0)	
FLORIDA	1421	3.2	0.2	(2.7, 3.6)	*	2016	3.9	0.2	(3.5, 4.2)	
GEORGIA	949	1.7	0.2	(1.3, 2.1)	*	1311	2.7	0.2	(2.3, 3.1)	
HAWAII	937	2.7	0.3	(2.1, 3.2)	*	1192	3.4	0.3	(2.8, 3.9)	
IDAHO	2046	3.1	0.2	(2.7, 3.5)	*	2822	3.8	0.2	(3.4, 4.2)	
ILLINOIS	1157	2.9	0.2	(2.4, 3.3)	*	1713	3.8	0.2	(3.3, 4.2)	
INDIANA	1056	2.8	0.2	(2.3, 3.2)	*	1266	4.0	0.3	(3.5, 4.5)	
IOWA	1473	2.4	0.2	(2.0, 2.7)	*	2105	3.4	0.2	(3.0, 3.7)	
KANSAS	780	1.7	0.2	(1.3, 2.1)	*	1200	2.4	0.2	(2.0, 2.8)	
KENTUCKY	1322	3.8	0.3	(3.3, 4.3)	*	2280	4.4	0.2	(4.0, 4.8)	
LOUISIANA	616	3.1	0.4	(2.4, 3.8)	*	999	3.3	0.3	(2.8, 3.7)	
MAINE	742	3.0	0.3	(2.4, 3.6)	*	936	3.9	0.3	(3.3, 4.5)	
MARYLAND	1952	2.1	0.2	(1.7, 2.4)	*	2599	2.7	0.2	(2.4, 3.1)	
MASSACHUSETTS	716	2.9	0.3	(2.3, 3.4)	*	1032	3.7	0.3	(3.1, 4.2)	
MICHIGAN	1062	2.6	0.2	(2.2, 3.1)	*	1478	3.8	0.2	(3.4, 4.3)	
MINNESOTA	2173	2.6	0.2	(2.3, 2.9)	*	2541	3.4	0.2	(3.1, 3.7)	
MISSISSIPPI	582	2.8	0.3	(2.2, 3.5)	*	971	3.3	0.3	(2.8, 3.8)	
MISSOURI	692	3.0	0.3	(2.4, 3.6)	*	1129	3.9	0.3	(3.4, 4.5)	
MONTANA	748	2.3	0.3	(1.8, 2.8)	*	1019	3.1	0.2	(2.6, 3.6)	
NEBRASKA	1041	2.6	0.3	(2.1, 3.1)	*	1640	3.2	0.2	(2.8, 3.7)	
NEVADA	1152	3.5	0.5	(2.5, 4.5)	*	1315	3.5	0.4	(2.8, 4.3)	
NEW HAMPSHIRE	600	2.4	0.3	(1.8, 3.0)	*	897	3.7	0.3	(3.2, 4.2)	
NEW JERSEY	1048	2.6	0.2	(2.2, 3.1)	*	1577	3.4	0.2	(3.0, 3.8)	
NEW MEXICO	748	2.9	0.3	(2.3, 3.5)	*	1051	3.4	0.3	(2.9, 3.9)	
NEW YORK	1339	2.7	0.2	(2.3, 3.1)	*	2008	3.5	0.2	(3.1, 3.9)	
NORTH CAROLINA	1420	2.4	0.2	(2.1, 2.8)	*	2148	3.3	0.2	(3.0, 3.6)	
NORTH DAKOTA	799	2.5	0.3	(2.0, 3.0)	*	999	3.2	0.3	(2.8, 3.7)	
OHIO	1227	2.0	0.2	(1.6, 2.4)	*	1925	3.1	0.2	(2.6, 3.5)	
OKLAHOMA	827	1.7	0.3	(1.2, 2.2)	*	1045	2.8	0.2	(2.3, 3.2)	
OREGON	1385	2.7	0.2	(2.3, 3.1)	*	1890	3.6	0.2	(3.3, 4.0)	
PENNSYLVANIA	1426	2.8	0.2	(2.4, 3.2)	*	2115	3.5	0.2	(3.1, 3.9)	
RHODE ISLAND	746	2.4	0.3	(1.9, 2.9)	*	1058	3.6	0.3	(3.1, 4.2)	
SOUTH CAROLINA	817	2.8	0.3	(2.2, 3.3)	*	1318	2.9	0.2	(2.5, 3.4)	
SOUTH DAKOTA	953	2.4	0.3	(1.9, 2.9)	*	1231	2.9	0.2	(2.5, 3.3)	
TENNESSEE	1029	2.9	0.3	(2.4, 3.3)	*	1893	3.9	0.2	(3.4, 4.3)	
TEXAS	1006	2.5	0.2	(2.1, 3.0)	*	1445	3.8	0.2	(3.3, 4.3)	
UTAH	1272	2.9	0.2	(2.4, 3.3)	*	1580	4.2	0.3	(3.7, 4.7)	
VERMONT	1375	2.6	0.2	(2.2, 3.0)	*	1759	2.9	0.2	(2.5, 3.2)	
VIRGINIA	1380	2.4	0.2	(2.0, 2.8)	*	2026	2.5	0.2	(2.1, 2.8)	
WASHINGTON	1523	2.9	0.2	(2.5, 3.3)	*	2004	3.2	0.2	(2.9, 3.6)	
WEST VIRGINIA	960	3.0	0.3	(2.5, 3.5)	*	1458	3.6	0.2	(3.2, 4.1)	
WISCONSIN	987	2.4	0.2	(2.0, 2.9)	*	1237	3.0	0.2	(2.6, 3.4)	
WYOMING	1011	2.3	0.2	(1.8, 2.8)	*	1377	3.0	0.2	(2.6, 3.4)	
PUERTO RICO	933	2.3	0.2	(1.9, 2.7)	*	1310	3.3	0.2	(2.9, 3.7)	

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 2.6
 RANGE = 0.6- 3.8

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.4
 RANGE = 1.5- 4.4

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

August 14, 1998

TABLE 2.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

MEAN DIFFERENCES FROM 1996 TO 1997

PHYSICAL HEALTH NOT GOOD*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 MEAN	1997 MEAN	DIFFERENCE
ALABAMA	3.3	3.8	0.5
ALASKA	2.1	2.9	0.8
ARIZONA	2.2	1.1	-1.1
ARKANSAS	3.1	3.4	0.3
CALIFORNIA	3.1	3.1	0.0
COLORADO	2.9	2.8	-0.1
CONNECTICUT	2.5	3.1	0.5
DELAWARE	2.8	3.7	1.0
DISTRICT OF CO	2.6	1.5	-1.1
FLORIDA	3.4	3.5	0.1
GEORGIA	2.0	2.2	0.2
HAWAII	2.5	3.0	0.5
IDAHO	2.8	3.4	0.6
ILLINOIS	3.2	3.3	0.2
INDIANA	3.4	3.4	-0.0
IOWA	2.8	2.9	0.1
KANSAS	2.0	2.1	0.0
KENTUCKY	3.6	4.1	0.5
LOUISIANA	2.9	3.2	0.3
MAINE	2.9	3.5	0.6
MARYLAND	2.5	2.4	-0.1
MASSACHUSETTS	3.2	3.3	0.1
MICHIGAN	2.8	3.3	0.5
MINNESOTA	3.0	3.0	0.0
MISSISSIPPI	3.2	3.1	-0.1
MISSOURI	3.5	3.5	0.0
MONTANA	2.6	2.7	0.1
NEBRASKA	2.9	2.9	0.0
NEVADA	3.0	3.5	0.5
NEW HAMPSHIRE	3.0	3.1	0.1
NEW JERSEY	3.0	3.0	0.0
NEW MEXICO	3.7	3.2	-0.5
NEW YORK	3.1	3.1	-0.0
NORTH CAROLINA	2.7	2.9	0.2
NORTH DAKOTA	3.6	2.9	-0.7
OHIO	2.6	2.6	0.0
OKLAHOMA	2.1	2.3	0.1
OREGON	3.0	3.2	0.2
PENNSYLVANIA	3.2	3.2	0.0
RHODE ISLAND	3.5	3.0	-0.5
SOUTH CAROLINA	2.7	2.9	0.2
SOUTH DAKOTA	2.3	2.6	0.4
TENNESSEE	3.2	3.4	0.2
TEXAS	3.3	3.2	-0.1
UTAH	3.1	3.6	0.5
VERMONT	3.2	2.7	-0.4
VIRGINIA	2.9	2.4	-0.5
WASHINGTON	3.3	3.1	-0.2
WEST VIRGINIA	3.3	3.4	0.1
WISCONSIN	3.2	2.8	-0.4
WYOMING	3.0	2.6	-0.3
PUERTO RICO	3.1	2.8	-0.3

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

N/A - NOT AVAILABLE

August 14, 1998

TABLE 3.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2120	3.3	0.2	(2.9, 3.7)
ALASKA	1506	3.2	0.3	(2.6, 3.7)
ARIZONA	1902	1.5	0.2	(1.1, 1.8)
ARKANSAS	1764	3.0	0.2	(2.5, 3.4)
CALIFORNIA	4043	3.4	0.1	(3.2, 3.7)
COLORADO	1810	3.2	0.2	(2.8, 3.5)
CONNECTICUT	2213	2.8	0.2	(2.4, 3.2)
DELAWARE	2522	3.2	0.2	(2.8, 3.5)
DISTRICT OF CO	1495	2.1	0.2	(1.7, 2.4)
FLORIDA	3446	3.4	0.2	(3.1, 3.7)
GEORGIA	2269	2.8	0.2	(2.5, 3.2)
HAWAII	2147	2.6	0.2	(2.2, 2.9)
IDAHO	4874	3.1	0.1	(2.9, 3.3)
ILLINOIS	2862	3.1	0.1	(2.8, 3.4)
INDIANA	2327	3.6	0.2	(3.3, 3.9)
IOWA	3572	2.9	0.1	(2.6, 3.1)
KANSAS	1975	1.7	0.1	(1.4, 1.9)
KENTUCKY	3579	4.9	0.2	(4.5, 5.3)
LOUISIANA	1623	3.5	0.2	(3.1, 3.9)
MAINE	1679	2.9	0.2	(2.5, 3.2)
MARYLAND	4549	1.8	0.1	(1.6, 2.0)
MASSACHUSETTS	1733	3.1	0.2	(2.7, 3.5)
MICHIGAN	2527	3.3	0.2	(3.0, 3.6)
MINNESOTA	4732	3.0	0.1	(2.8, 3.2)
MISSISSIPPI	1556	2.5	0.2	(2.1, 2.9)
MISSOURI	1813	3.2	0.2	(2.8, 3.6)
MONTANA	1757	2.5	0.2	(2.2, 2.7)
NEBRASKA	2660	2.5	0.1	(2.3, 2.8)
NEVADA	2478	3.7	0.3	(3.1, 4.3)
NEW HAMPSHIRE	1491	3.3	0.2	(2.8, 3.7)
NEW JERSEY	2621	2.9	0.2	(2.6, 3.2)
NEW MEXICO	1796	3.3	0.2	(2.9, 3.6)
NEW YORK	3302	3.0	0.2	(2.7, 3.3)
NORTH CAROLINA	3555	2.5	0.1	(2.2, 2.7)
NORTH DAKOTA	1796	3.0	0.2	(2.7, 3.4)
OHIO	3148	2.2	0.2	(1.9, 2.5)
OKLAHOMA	1873	1.8	0.2	(1.5, 2.1)
OREGON	3278	3.2	0.1	(2.9, 3.5)
PENNSYLVANIA	3541	2.9	0.1	(2.7, 3.2)
RHODE ISLAND	1814	2.7	0.2	(2.4, 3.0)
SOUTH CAROLINA	2146	1.8	0.2	(1.5, 2.1)
SOUTH DAKOTA	2172	2.7	0.2	(2.4, 3.1)
TENNESSEE	2912	3.5	0.2	(3.2, 3.9)
TEXAS	2460	2.9	0.2	(2.6, 3.2)
UTAH	2850	3.5	0.2	(3.1, 3.8)
VERMONT	3141	2.7	0.1	(2.4, 2.9)
VIRGINIA	3426	2.7	0.1	(2.4, 2.9)
WASHINGTON	3533	3.1	0.1	(2.8, 3.4)
WEST VIRGINIA	2409	2.3	0.1	(2.0, 2.6)
WISCONSIN	2225	2.5	0.1	(2.2, 2.8)
WYOMING	2394	2.7	0.2	(2.4, 3.0)
PUERTO RICO	2235	1.7	0.1	(1.4, 1.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 2.9 RANGE = 1.5- 4.9

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD August 14, 1998

TABLE 3.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	799	2.7	0.3	(2.1, 3.2)	*	1321	3.9	0.3	(3.4, 4.4)	
ALASKA	696	2.4	0.4	(1.7, 3.1)	*	810	4.0	0.4	(3.2, 4.8)	
ARIZONA	891	0.6	0.1	(0.3, 0.8)	*	1011	2.4	0.3	(1.7, 3.0)	
ARKANSAS	664	2.6	0.3	(2.0, 3.3)	*	1100	3.3	0.3	(2.7, 3.8)	
CALIFORNIA	1748	2.8	0.2	(2.4, 3.1)	*	2295	4.1	0.2	(3.7, 4.4)	
COLORADO	815	2.5	0.2	(2.1, 3.0)	*	995	3.8	0.3	(3.2, 4.3)	
CONNECTICUT	878	2.4	0.4	(1.6, 3.1)	*	1335	3.2	0.2	(2.7, 3.6)	
DELAWARE	1001	2.5	0.3	(2.0, 3.0)	*	1521	3.8	0.3	(3.3, 4.3)	
DISTRICT OF CO	571	2.1	0.3	(1.5, 2.7)	*	924	2.0	0.2	(1.6, 2.4)	
FLORIDA	1421	3.0	0.2	(2.6, 3.5)	*	2025	3.8	0.2	(3.4, 4.2)	
GEORGIA	950	2.7	0.3	(2.1, 3.3)	*	1319	2.9	0.2	(2.5, 3.3)	
HAWAII	939	2.1	0.2	(1.7, 2.5)	*	1208	3.0	0.3	(2.5, 3.6)	
IDAHO	2045	2.2	0.2	(1.9, 2.5)	*	2829	4.0	0.2	(3.6, 4.3)	
ILLINOIS	1157	2.7	0.2	(2.3, 3.1)	*	1705	3.5	0.2	(3.1, 3.8)	
INDIANA	1055	3.0	0.2	(2.5, 3.4)	*	1272	4.2	0.3	(3.7, 4.7)	
IOWA	1468	2.2	0.2	(1.9, 2.6)	*	2104	3.4	0.2	(3.1, 3.8)	
KANSAS	779	1.0	0.2	(0.7, 1.3)	*	1196	2.3	0.2	(1.9, 2.7)	
KENTUCKY	1313	3.9	0.3	(3.4, 4.5)	*	2266	5.8	0.3	(5.3, 6.3)	
LOUISIANA	620	2.9	0.3	(2.3, 3.6)	*	1003	4.0	0.3	(3.4, 4.6)	
MAINE	743	2.6	0.3	(2.0, 3.2)	*	936	3.1	0.3	(2.6, 3.5)	
MARYLAND	1950	1.4	0.1	(1.1, 1.7)	*	2599	2.2	0.2	(1.9, 2.6)	
MASSACHUSETTS	707	2.9	0.3	(2.2, 3.5)	*	1026	3.4	0.3	(2.9, 3.9)	
MICHIGAN	1057	2.7	0.2	(2.3, 3.1)	*	1470	3.8	0.2	(3.3, 4.2)	
MINNESOTA	2174	2.5	0.2	(2.2, 2.7)	*	2558	3.5	0.2	(3.1, 3.8)	
MISSISSIPPI	581	1.9	0.3	(1.4, 2.4)	*	975	3.0	0.2	(2.5, 3.5)	
MISSOURI	684	2.6	0.3	(2.0, 3.2)	*	1129	3.8	0.3	(3.2, 4.3)	
MONTANA	743	1.8	0.2	(1.4, 2.2)	*	1014	3.1	0.2	(2.7, 3.5)	
NEBRASKA	1037	1.8	0.2	(1.4, 2.1)	*	1623	3.3	0.2	(2.8, 3.7)	
NEVADA	1154	3.6	0.5	(2.6, 4.6)	*	1324	3.8	0.4	(3.1, 4.5)	
NEW HAMPSHIRE	595	3.2	0.4	(2.5, 4.0)	*	896	3.3	0.3	(2.8, 3.8)	
NEW JERSEY	1051	2.5	0.2	(2.0, 3.0)	*	1570	3.3	0.2	(2.9, 3.7)	
NEW MEXICO	751	2.6	0.3	(2.1, 3.1)	*	1045	3.9	0.3	(3.4, 4.4)	
NEW YORK	1326	2.8	0.2	(2.4, 3.3)	*	1976	3.2	0.2	(2.8, 3.6)	
NORTH CAROLINA	1419	2.1	0.2	(1.7, 2.5)	*	2136	2.8	0.2	(2.5, 3.1)	
NORTH DAKOTA	797	2.5	0.2	(2.1, 3.0)	*	999	3.5	0.2	(3.0, 4.0)	
OHIO	1226	1.6	0.2	(1.2, 2.0)	*	1922	2.7	0.2	(2.3, 3.1)	
OKLAHOMA	827	1.2	0.2	(0.8, 1.5)	*	1046	2.3	0.2	(1.9, 2.7)	
OREGON	1389	2.5	0.2	(2.1, 2.9)	*	1889	3.8	0.2	(3.4, 4.2)	
PENNSYLVANIA	1426	2.3	0.2	(1.9, 2.7)	*	2115	3.5	0.2	(3.1, 3.9)	
RHODE ISLAND	749	1.9	0.2	(1.5, 2.3)	*	1065	3.4	0.2	(2.9, 3.9)	
SOUTH CAROLINA	817	1.8	0.3	(1.3, 2.3)	*	1329	1.8	0.2	(1.4, 2.1)	
SOUTH DAKOTA	946	2.2	0.3	(1.7, 2.7)	*	1226	3.2	0.2	(2.8, 3.7)	
TENNESSEE	1031	2.9	0.2	(2.4, 3.3)	*	1881	4.1	0.2	(3.7, 4.6)	
TEXAS	1009	2.1	0.2	(1.7, 2.5)	*	1451	3.6	0.2	(3.2, 4.1)	
UTAH	1269	2.9	0.2	(2.4, 3.3)	*	1581	4.0	0.2	(3.6, 4.5)	
VERMONT	1368	1.9	0.2	(1.5, 2.2)	*	1773	3.4	0.2	(3.0, 3.8)	
VIRGINIA	1384	2.3	0.2	(1.8, 2.7)	*	2042	3.0	0.2	(2.7, 3.4)	
WASHINGTON	1519	2.8	0.2	(2.3, 3.2)	*	2014	3.4	0.2	(3.0, 3.8)	
WEST VIRGINIA	955	1.6	0.2	(1.3, 2.0)	*	1454	3.0	0.2	(2.6, 3.3)	
WISCONSIN	985	1.8	0.2	(1.5, 2.2)	*	1240	3.2	0.2	(2.7, 3.6)	
WYOMING	1015	1.9	0.2	(1.5, 2.3)	*	1379	3.4	0.2	(3.0, 3.9)	
PUERTO RICO	927	1.5	0.2	(1.1, 1.9)	*	1308	1.8	0.2	(1.5, 2.2)	

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 2.4
RANGE = 0.6- 3.9

NUMBER OF PARTICIPANTS = 52
MEDIAN = 3.4
RANGE = 1.8- 5.8

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

August 14, 1998

TABLE 3.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

MEAN DIFFERENCES FROM 1996 TO 1997

MENTAL HEALTH NOT GOOD*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 MEAN	1997 MEAN	DIFFERENCE
ALABAMA	3.4	3.3	-0.1
ALASKA	3.0	3.2	0.1
ARIZONA	2.2	1.5	-0.7
ARKANSAS	3.0	3.0	0.0
CALIFORNIA	3.0	3.4	0.4
COLORADO	3.3	3.2	-0.1
CONNECTICUT	2.9	2.8	-0.2
DELAWARE	2.3	3.2	0.9
DISTRICT OF CO	3.0	2.1	-1.0
FLORIDA	3.2	3.4	0.2
GEORGIA	2.2	2.8	0.7
HAWAII	3.0	2.6	-0.4
IDAHO	3.0	3.1	0.1
ILLINOIS	3.1	3.1	0.0
INDIANA	3.3	3.6	0.3
IOWA	2.9	2.9	0.0
KANSAS	2.0	1.7	-0.3
KENTUCKY	4.6	4.9	0.3
LOUISIANA	3.1	3.5	0.4
MAINE	2.9	2.9	-0.1
MARYLAND	2.3	1.8	-0.5
MASSACHUSETTS	3.3	3.1	-0.2
MICHIGAN	3.3	3.3	0.0
MINNESOTA	3.2	3.0	-0.2
MISSISSIPPI	2.3	2.5	0.2
MISSOURI	3.0	3.2	0.2
MONTANA	2.5	2.5	0.0
NEBRASKA	3.0	2.5	-0.4
NEVADA	3.3	3.7	0.4
NEW HAMPSHIRE	2.8	3.3	0.4
NEW JERSEY	3.1	2.9	-0.1
NEW MEXICO	3.4	3.3	-0.1
NEW YORK	3.0	3.0	0.1
NORTH CAROLINA	2.0	2.5	0.4
NORTH DAKOTA	2.8	3.0	0.3
OHIO	1.9	2.2	0.3
OKLAHOMA	2.1	1.8	-0.3
OREGON	3.2	3.2	0.0
PENNSYLVANIA	2.9	2.9	0.0
RHODE ISLAND	3.1	2.7	-0.4
SOUTH CAROLINA	2.8	1.8	-1.1
SOUTH DAKOTA	1.8	2.7	1.0
TENNESSEE	2.5	3.5	1.0
TEXAS	3.3	2.9	-0.4
UTAH	3.6	3.5	-0.2
VERMONT	2.9	2.7	-0.3
VIRGINIA	2.6	2.7	0.1
WASHINGTON	3.2	3.1	-0.1
WEST VIRGINIA	2.5	2.3	-0.2
WISCONSIN	3.0	2.5	-0.5
WYOMING	2.6	2.7	0.1
PUERTO RICO	2.4	1.7	-0.7

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

N/A - NOT AVAILABLE

August 14, 1998

TABLE 4.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1127	4.2	0.3	(3.6, 4.8)
ALASKA	811	2.3	0.3	(1.8, 2.8)
ARIZONA	335	5.2	0.7	(3.7, 6.6)
ARKANSAS	677	5.9	0.4	(5.0, 6.7)
CALIFORNIA	2144	3.4	0.2	(3.1, 3.7)
COLORADO	993	2.9	0.3	(2.4, 3.4)
CONNECTICUT	1052	3.7	0.5	(2.8, 4.6)
DELAWARE	1148	5.2	0.3	(4.6, 5.9)
DISTRICT OF CO	609	2.0	0.2	(1.5, 2.4)
FLORIDA	1734	4.1	0.2	(3.6, 4.5)
GEORGIA	963	3.2	0.3	(2.7, 3.7)
HAWAII	1064	3.2	0.2	(2.7, 3.6)
IDAHO	2675	3.5	0.2	(3.1, 3.9)
ILLINOIS	1644	3.1	0.2	(2.7, 3.5)
INDIANA	1316	3.1	0.2	(2.6, 3.5)
IOWA	1862	3.0	0.2	(2.7, 3.4)
KANSAS	752	2.3	0.3	(1.8, 2.8)
KENTUCKY	1548	7.2	0.3	(6.5, 7.8)
LOUISIANA	809	4.3	0.4	(3.6, 5.0)
MAINE	799	4.1	0.3	(3.5, 4.7)
MARYLAND	1633	3.5	0.3	(3.1, 4.0)
MASSACHUSETTS	888	3.7	0.3	(3.1, 4.3)
MICHIGAN	1367	3.2	0.2	(2.7, 3.6)
MINNESOTA	2580	3.7	0.2	(3.4, 4.0)
MISSISSIPPI	670	3.6	0.3	(3.0, 4.3)
MISSOURI	963	3.7	0.3	(3.1, 4.4)
MONTANA	863	2.8	0.2	(2.3, 3.3)
NEBRASKA	1335	3.6	0.3	(3.1, 4.1)
NEVADA	1429	3.5	0.4	(2.8, 4.3)
NEW HAMPSHIRE	786	3.5	0.4	(2.8, 4.2)
NEW JERSEY	1276	3.7	0.3	(3.2, 4.2)
NEW MEXICO	957	3.8	0.3	(3.2, 4.3)
NEW YORK	1663	3.8	0.2	(3.3, 4.3)
NORTH CAROLINA	1495	4.0	0.2	(3.6, 4.5)
NORTH DAKOTA	902	3.0	0.3	(2.6, 3.5)
OHIO	1174	3.9	0.3	(3.3, 4.5)
OKLAHOMA	511	3.6	0.4	(2.9, 4.3)
OREGON	1869	2.9	0.2	(2.6, 3.3)
PENNSYLVANIA	1708	3.5	0.2	(3.0, 3.9)
RHODE ISLAND	895	3.9	0.3	(3.3, 4.5)
SOUTH CAROLINA	758	4.9	0.4	(4.1, 5.6)
SOUTH DAKOTA	1062	2.8	0.3	(2.3, 3.3)
TENNESSEE	1409	3.9	0.3	(3.4, 4.4)
TEXAS	1266	3.7	0.3	(3.2, 4.2)
UTAH	1722	3.2	0.2	(2.8, 3.7)
VERMONT	1507	3.0	0.2	(2.6, 3.4)
VIRGINIA	1741	2.8	0.2	(2.5, 3.2)
WASHINGTON	2005	3.0	0.2	(2.6, 3.4)
WEST VIRGINIA	926	5.9	0.4	(5.2, 6.6)
WISCONSIN	1182	2.6	0.2	(2.2, 2.9)
WYOMING	1212	2.7	0.2	(2.2, 3.1)
PUERTO RICO	758	4.0	0.3	(3.3, 4.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.6 RANGE = 2.0- 7.2

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

August 14, 1998

TABLE 4.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*					
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		
ALABAMA	352	4.5	0.6	(3.4,	5.6)	*	775	4.0	0.3	(3.3,	4.6)
ALASKA	327	2.0	0.4	(1.3,	2.8)	*	484	2.5	0.4	(1.7,	3.3)
ARIZONA	108	4.1	0.9	(2.3,	6.0)	*	227	5.7	1.0	(3.8,	7.5)
ARKANSAS	212	6.6	0.8	(5.0,	8.2)	*	465	5.4	0.5	(4.4,	6.3)
CALIFORNIA	815	3.6	0.3	(3.0,	4.2)	*	1329	3.3	0.2	(2.9,	3.6)
COLORADO	392	3.2	0.4	(2.4,	4.0)	*	601	2.7	0.3	(2.0,	3.3)
CONNECTICUT	359	4.3	1.0	(2.4,	6.1)	*	693	3.3	0.3	(2.7,	4.0)
DELAWARE	390	5.7	0.6	(4.6,	6.9)	*	758	4.9	0.4	(4.1,	5.6)
DISTRICT OF CO	222	1.9	0.4	(1.2,	2.6)	*	387	2.0	0.3	(1.5,	2.6)
FLORIDA	627	4.0	0.4	(3.3,	4.7)	*	1107	4.1	0.3	(3.6,	4.7)
GEORGIA	356	2.9	0.4	(2.2,	3.6)	*	607	3.4	0.3	(2.8,	4.1)
HAWAII	418	3.6	0.4	(2.8,	4.4)	*	646	2.8	0.3	(2.3,	3.3)
IDAHO	943	3.5	0.3	(2.9,	4.1)	*	1732	3.4	0.3	(2.9,	3.9)
ILLINOIS	584	3.5	0.4	(2.7,	4.2)	*	1060	2.8	0.2	(2.4,	3.3)
INDIANA	507	3.0	0.4	(2.3,	3.7)	*	809	3.1	0.3	(2.5,	3.6)
IOWA	654	3.1	0.3	(2.5,	3.7)	*	1208	2.9	0.2	(2.5,	3.3)
KANSAS	240	2.4	0.5	(1.5,	3.4)	*	512	2.2	0.3	(1.7,	2.8)
KENTUCKY	484	8.0	0.6	(6.9,	9.1)	*	1064	6.6	0.4	(5.9,	7.3)
LOUISIANA	267	5.1	0.7	(3.8,	6.4)	*	542	3.7	0.4	(3.0,	4.5)
MAINE	311	4.8	0.6	(3.7,	5.8)	*	488	3.6	0.4	(2.8,	4.3)
MARYLAND	620	3.7	0.4	(2.9,	4.5)	*	1013	3.4	0.3	(2.8,	4.1)
MASSACHUSETTS	338	4.1	0.5	(3.0,	5.1)	*	550	3.4	0.3	(2.8,	4.0)
MICHIGAN	501	3.3	0.4	(2.5,	4.0)	*	866	3.1	0.3	(2.6,	3.6)
MINNESOTA	1055	3.9	0.3	(3.3,	4.4)	*	1525	3.6	0.2	(3.2,	4.0)
MISSISSIPPI	211	4.2	0.7	(2.9,	5.5)	*	459	3.3	0.4	(2.5,	4.0)
MISSOURI	314	4.1	0.6	(3.0,	5.2)	*	649	3.5	0.3	(2.8,	4.1)
MONTANA	295	2.7	0.4	(1.8,	3.5)	*	568	2.9	0.3	(2.3,	3.5)
NEBRASKA	460	3.4	0.4	(2.6,	4.2)	*	875	3.8	0.3	(3.1,	4.4)
NEVADA	601	3.2	0.6	(2.1,	4.3)	*	828	3.9	0.5	(2.8,	4.9)
NEW HAMPSHIRE	283	3.3	0.6	(2.1,	4.5)	*	503	3.7	0.4	(2.9,	4.4)
NEW JERSEY	439	3.8	0.4	(3.0,	4.7)	*	837	3.6	0.3	(3.0,	4.2)
NEW MEXICO	335	3.9	0.5	(2.9,	4.8)	*	622	3.7	0.3	(3.1,	4.3)
NEW YORK	601	4.3	0.4	(3.5,	5.1)	*	1062	3.5	0.3	(2.9,	4.0)
NORTH CAROLINA	503	4.1	0.4	(3.3,	4.9)	*	992	4.0	0.3	(3.5,	4.5)
NORTH DAKOTA	359	3.1	0.4	(2.2,	4.0)	*	543	3.0	0.3	(2.4,	3.6)
OHIO	382	4.1	0.5	(3.0,	5.2)	*	792	3.7	0.4	(3.0,	4.5)
OKLAHOMA	150	3.1	0.5	(2.1,	4.2)	*	361	3.8	0.5	(2.9,	4.8)
OREGON	697	2.7	0.3	(2.1,	3.2)	*	1172	3.1	0.2	(2.6,	3.6)
PENNSYLVANIA	584	3.7	0.4	(2.9,	4.4)	*	1124	3.3	0.3	(2.8,	3.8)
RHODE ISLAND	314	3.8	0.5	(2.8,	4.8)	*	581	4.0	0.4	(3.2,	4.8)
SOUTH CAROLINA	255	5.2	0.6	(3.9,	6.4)	*	503	4.6	0.4	(3.8,	5.5)
SOUTH DAKOTA	409	3.3	0.5	(2.4,	4.2)	*	653	2.4	0.3	(1.9,	2.9)
TENNESSEE	411	4.3	0.5	(3.4,	5.2)	*	998	3.7	0.3	(3.1,	4.2)
TEXAS	455	3.3	0.4	(2.5,	4.1)	*	811	4.0	0.3	(3.3,	4.6)
UTAH	674	3.6	0.4	(2.8,	4.3)	*	1048	3.0	0.3	(2.5,	3.5)
VERMONT	592	3.0	0.3	(2.4,	3.7)	*	915	3.0	0.2	(2.5,	3.4)
VIRGINIA	632	3.0	0.3	(2.4,	3.6)	*	1109	2.7	0.2	(2.3,	3.1)
WASHINGTON	782	3.1	0.3	(2.5,	3.8)	*	1223	2.9	0.2	(2.4,	3.3)
WEST VIRGINIA	305	6.6	0.6	(5.4,	7.9)	*	621	5.5	0.4	(4.7,	6.3)
WISCONSIN	450	2.6	0.3	(2.0,	3.2)	*	732	2.6	0.2	(2.1,	3.0)
WYOMING	458	2.5	0.4	(1.8,	3.2)	*	754	2.8	0.3	(2.2,	3.4)
PUERTO RICO	256	4.9	0.6	(3.7,	6.1)	*	502	3.3	0.3	(2.7,	4.0)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.6
 RANGE = 1.9- 8.0

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.4
 RANGE = 2.0- 6.6

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

August 14, 1998

TABLE 4.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

MEAN DIFFERENCES FROM 1996 TO 1997

ACTIVITIES LIMITED*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 MEAN	1997 MEAN	DIFFERENCE
ALABAMA	3.7	4.2	0.5
ALASKA	2.6	2.3	-0.3
ARIZONA	5.5	5.2	-0.4
ARKANSAS	6.0	5.9	-0.1
CALIFORNIA	3.6	3.4	-0.2
COLORADO	2.9	2.9	-0.0
CONNECTICUT	2.8	3.7	1.0
DELAWARE	4.5	5.2	0.7
DISTRICT OF CO	3.6	2.0	-1.6
FLORIDA	3.9	4.1	0.2
GEORGIA	2.8	3.2	0.4
HAWAII	3.0	3.2	0.1
IDAHO	3.0	3.5	0.5
ILLINOIS	3.1	3.1	-0.0
INDIANA	3.2	3.1	-0.1
IOWA	2.9	3.0	0.1
KANSAS	2.5	2.3	-0.2
KENTUCKY	6.5	7.2	0.7
LOUISIANA	4.1	4.3	0.2
MAINE	4.6	4.1	-0.5
MARYLAND	3.2	3.5	0.3
MASSACHUSETTS	3.0	3.7	0.7
MICHIGAN	3.2	3.2	-0.0
MINNESOTA	3.5	3.7	0.2
MISSISSIPPI	4.9	3.6	-1.3
MISSOURI	3.6	3.7	0.2
MONTANA	2.8	2.8	0.0
NEBRASKA	4.4	3.6	-0.8
NEVADA	3.2	3.5	0.4
NEW HAMPSHIRE	3.5	3.5	0.0
NEW JERSEY	3.4	3.7	0.3
NEW MEXICO	4.0	3.8	-0.3
NEW YORK	3.4	3.8	0.4
NORTH CAROLINA	5.6	4.0	-1.5
NORTH DAKOTA	3.0	3.0	0.0
OHIO	3.5	3.9	0.4
OKLAHOMA	5.1	3.6	-1.6
OREGON	3.3	2.9	-0.4
PENNSYLVANIA	3.4	3.5	0.1
RHODE ISLAND	3.8	3.9	0.1
SOUTH CAROLINA	5.2	4.9	-0.4
SOUTH DAKOTA	3.1	2.8	-0.3
TENNESSEE	4.2	3.9	-0.3
TEXAS	3.7	3.7	-0.1
UTAH	2.8	3.2	0.5
VERMONT	3.4	3.0	-0.4
VIRGINIA	2.8	2.8	0.0
WASHINGTON	2.9	3.0	0.1
WEST VIRGINIA	6.1	5.9	-0.1
WISCONSIN	3.1	2.6	-0.5
WYOMING	2.8	2.7	-0.2
PUERTO RICO	4.5	4.0	-0.6

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 5.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2171	14.4	0.9	(12.7, 16.1)
ALASKA	1536	21.3	1.6	(18.2, 24.4)
ARIZONA	1902	14.7	1.2	(12.4, 16.9)
ARKANSAS	1792	17.3	1.2	(15.0, 19.6)
CALIFORNIA	4058	19.3	0.8	(17.8, 20.8)
COLORADO	1824	11.9	0.9	(10.0, 13.7)
CONNECTICUT	2246	9.4	0.9	(7.8, 11.1)
DELAWARE	2554	10.8	0.8	(9.2, 12.3)
DISTRICT OF CO	1504	11.6	1.0	(9.6, 13.6)
FLORIDA	3487	17.6	0.8	(16.1, 19.1)
GEORGIA	2303	12.0	0.9	(10.3, 13.7)
HAWAII	2166	6.4	0.7	(5.1, 7.7)
IDAHO	4911	16.6	0.8	(15.1, 18.1)
ILLINOIS	2883	11.3	0.7	(10.0, 12.7)
INDIANA	2365	12.4	0.8	(10.8, 14.0)
IOWA	3596	9.6	0.6	(8.4, 10.7)
KANSAS	2000	9.5	0.8	(8.0, 10.9)
KENTUCKY	3607	13.8	0.7	(12.4, 15.2)
LOUISIANA	1652	20.4	1.2	(18.0, 22.8)
MAINE	1699	12.0	0.9	(10.3, 13.8)
MARYLAND	4566	10.2	0.6	(9.0, 11.4)
MASSACHUSETTS	1765	9.3	0.8	(7.6, 10.9)
MICHIGAN	2561	9.9	0.7	(8.6, 11.3)
MINNESOTA	4808	8.4	0.5	(7.5, 9.3)
MISSISSIPPI	1595	15.2	1.1	(13.0, 17.4)
MISSOURI	1848	12.2	1.0	(10.3, 14.1)
MONTANA	1800	14.6	0.9	(12.8, 16.4)
NEBRASKA	2692	7.6	0.6	(6.4, 8.8)
NEVADA	2496	14.1	1.4	(11.3, 16.8)
NEW HAMPSHIRE	1511	10.3	1.0	(8.4, 12.2)
NEW JERSEY	2658	11.7	0.8	(10.1, 13.2)
NEW MEXICO	1807	22.2	1.2	(19.9, 24.5)
NEW YORK	3397	14.0	0.7	(12.6, 15.4)
NORTH CAROLINA	3624	14.7	0.7	(13.3, 16.1)
NORTH DAKOTA	1799	11.7	0.9	(9.9, 13.5)
OHIO	3164	10.5	0.7	(9.0, 11.9)
OKLAHOMA	1880	17.0	1.1	(14.9, 19.2)
OREGON	3313	12.5	0.7	(11.1, 14.0)
PENNSYLVANIA	3595	9.4	0.6	(8.3, 10.6)
RHODE ISLAND	1836	10.7	0.9	(8.9, 12.5)
SOUTH CAROLINA	2150	15.0	1.0	(13.1, 17.0)
SOUTH DAKOTA	2196	13.2	0.9	(11.5, 14.9)
TENNESSEE	2973	11.7	0.7	(10.3, 13.2)
TEXAS	2486	24.2	1.0	(22.3, 26.2)
UTAH	2861	11.2	0.8	(9.6, 12.7)
VERMONT	3177	14.4	0.8	(12.9, 15.9)
VIRGINIA	3512	11.3	0.7	(9.9, 12.6)
WASHINGTON	3598	10.9	0.6	(9.7, 12.2)
WEST VIRGINIA	2429	18.1	0.9	(16.3, 19.9)
WISCONSIN	2243	9.3	0.9	(7.6, 11.0)
WYOMING	2406	19.0	1.2	(16.6, 21.3)
PUERTO RICO	2260	10.3	0.8	(8.8, 11.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 12.0 RANGE = 6.4-24.2

* HAVING NO HEALTH CARE PLAN

August 14, 1998

TABLE 5.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE* - BY SEX
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	810	11.5	1.3	(9.0, 14.0)	*	1361	17.0	1.2	(14.6, 19.3)	*
ALASKA	713	21.6	2.4	(17.0, 26.2)	*	823	20.9	2.1	(16.8, 25.1)	*
ARIZONA	889	15.9	1.8	(12.5, 19.4)	*	1013	13.4	1.6	(10.4, 16.5)	*
ARKANSAS	675	18.0	2.0	(14.0, 21.9)	*	1117	16.7	1.3	(14.2, 19.2)	*
CALIFORNIA	1757	22.3	1.2	(19.9, 24.8)	*	2301	16.4	0.9	(14.5, 18.2)	*
COLORADO	817	13.7	1.4	(10.9, 16.5)	*	1007	10.2	1.2	(7.8, 12.5)	*
CONNECTICUT	887	10.7	1.5	(7.8, 13.6)	*	1359	8.3	0.9	(6.5, 10.1)	*
DELAWARE	1012	13.0	1.3	(10.4, 15.5)	*	1542	8.7	0.9	(7.0, 10.5)	*
DISTRICT OF CO	572	13.4	1.7	(10.1, 16.7)	*	932	10.0	1.1	(7.8, 12.3)	*
FLORIDA	1432	17.8	1.2	(15.5, 20.1)	*	2055	17.3	1.0	(15.4, 19.3)	*
GEORGIA	959	12.9	1.4	(10.1, 15.7)	*	1344	11.2	1.1	(9.1, 13.3)	*
HAWAII	946	6.7	0.9	(4.9, 8.4)	*	1220	6.2	0.9	(4.4, 8.0)	*
IDAHO	2051	17.6	1.2	(15.3, 19.9)	*	2860	15.7	1.0	(13.8, 17.7)	*
ILLINOIS	1162	10.8	1.1	(8.7, 12.8)	*	1721	11.9	0.9	(10.1, 13.7)	*
INDIANA	1064	13.1	1.3	(10.5, 15.6)	*	1301	11.8	1.0	(9.8, 13.7)	*
IOWA	1476	10.4	0.9	(8.6, 12.2)	*	2120	8.8	0.7	(7.4, 10.2)	*
KANSAS	785	9.8	1.2	(7.4, 12.2)	*	1215	9.1	0.9	(7.4, 10.8)	*
KENTUCKY	1323	13.8	1.1	(11.6, 15.9)	*	2284	13.9	0.9	(12.1, 15.6)	*
LOUISIANA	626	21.2	2.0	(17.3, 25.2)	*	1026	19.7	1.5	(16.8, 22.6)	*
MAINE	751	13.4	1.4	(10.6, 16.1)	*	948	10.8	1.1	(8.6, 13.0)	*
MARYLAND	1954	10.5	1.0	(8.6, 12.4)	*	2612	10.0	0.8	(8.5, 11.5)	*
MASSACHUSETTS	721	11.1	1.4	(8.4, 13.8)	*	1044	7.6	0.9	(5.7, 9.4)	*
MICHIGAN	1067	11.8	1.1	(9.6, 14.0)	*	1494	8.2	0.8	(6.7, 9.7)	*
MINNESOTA	2201	8.9	0.7	(7.6, 10.2)	*	2607	7.9	0.6	(6.7, 9.1)	*
MISSISSIPPI	590	15.8	1.9	(12.2, 19.5)	*	1005	14.6	1.3	(12.0, 17.2)	*
MISSOURI	698	14.2	1.7	(10.9, 17.4)	*	1150	10.4	1.0	(8.3, 12.4)	*
MONTANA	757	14.4	1.4	(11.7, 17.1)	*	1043	14.7	1.2	(12.4, 17.1)	*
NEBRASKA	1046	7.8	1.0	(5.8, 9.7)	*	1646	7.4	0.8	(5.8, 8.9)	*
NEVADA	1159	13.9	2.2	(9.6, 18.3)	*	1337	14.2	1.7	(10.9, 17.5)	*
NEW HAMPSHIRE	602	10.9	1.6	(7.8, 13.9)	*	909	9.8	1.1	(7.7, 11.9)	*
NEW JERSEY	1057	12.9	1.3	(10.2, 15.5)	*	1601	10.6	0.9	(8.8, 12.4)	*
NEW MEXICO	753	20.8	1.7	(17.4, 24.1)	*	1054	23.6	1.6	(20.5, 26.6)	*
NEW YORK	1352	15.9	1.2	(13.6, 18.2)	*	2045	12.3	0.9	(10.6, 14.0)	*
NORTH CAROLINA	1434	16.3	1.1	(14.1, 18.5)	*	2190	13.3	0.9	(11.6, 14.9)	*
NORTH DAKOTA	799	12.3	1.4	(9.6, 14.9)	*	1000	11.2	1.1	(9.0, 13.4)	*
OHIO	1229	10.8	1.1	(8.6, 13.0)	*	1935	10.2	0.9	(8.3, 12.0)	*
OKLAHOMA	827	15.3	1.7	(11.9, 18.6)	*	1053	18.6	1.5	(15.7, 21.5)	*
OREGON	1398	14.8	1.3	(12.3, 17.2)	*	1915	10.5	0.8	(8.9, 12.1)	*
PENNSYLVANIA	1439	10.8	0.9	(8.9, 12.6)	*	2156	8.2	0.7	(6.9, 9.6)	*
RHODE ISLAND	753	11.5	1.3	(8.9, 14.2)	*	1083	9.9	1.3	(7.4, 12.4)	*
SOUTH CAROLINA	819	15.9	1.6	(12.8, 18.9)	*	1331	14.3	1.2	(12.0, 16.6)	*
SOUTH DAKOTA	953	13.4	1.3	(10.8, 16.0)	*	1243	13.0	1.1	(10.8, 15.2)	*
TENNESSEE	1042	13.6	1.2	(11.2, 16.0)	*	1931	10.1	0.8	(8.5, 11.6)	*
TEXAS	1018	25.8	1.6	(22.7, 28.8)	*	1468	22.8	1.2	(20.3, 25.2)	*
UTAH	1275	12.0	1.1	(9.8, 14.2)	*	1586	10.3	1.1	(8.2, 12.4)	*
VERMONT	1387	16.4	1.2	(14.0, 18.7)	*	1790	12.5	0.9	(10.7, 14.3)	*
VIRGINIA	1409	10.7	1.1	(8.6, 12.7)	*	2103	11.9	0.9	(10.1, 13.7)	*
WASHINGTON	1544	11.7	0.9	(9.9, 13.5)	*	2054	10.2	0.9	(8.5, 11.9)	*
WEST VIRGINIA	959	18.6	1.4	(15.8, 21.4)	*	1470	17.7	1.2	(15.4, 20.0)	*
WISCONSIN	991	10.8	1.4	(8.1, 13.5)	*	1252	7.9	1.1	(5.9, 10.0)	*
WYOMING	1013	20.7	2.0	(16.8, 24.6)	*	1393	17.3	1.1	(15.1, 19.4)	*
PUERTO RICO	936	11.8	1.2	(9.5, 14.1)	*	1324	8.9	1.0	(7.0, 10.8)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 13.4
 RANGE = 6.7-25.8

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 11.2
 RANGE = 6.2-23.6

* HAVING NO HEALTH CARE PLAN

August 14, 1998

TABLE 5.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	16.0	14.4	-1.6
ALASKA	20.8	21.3	0.5
ARIZONA	17.0	14.7	-2.4
ARKANSAS	17.9	17.3	-0.6
CALIFORNIA	19.5	19.3	-0.2
COLORADO	13.5	11.9	-1.6
CONNECTICUT	9.3	9.4	0.1
DELAWARE	10.8	10.8	-0.0
DISTRICT OF CO	13.4	11.6	-1.8
FLORIDA	17.6	17.6	0.0
GEORGIA	9.8	12.0	2.3
HAWAII	8.1	6.4	-1.7
IDAHO	15.0	16.6	1.6
ILLINOIS	11.8	11.3	-0.5
INDIANA	11.4	12.4	0.9
IOWA	9.2	9.6	0.4
KANSAS	10.1	9.5	-0.6
KENTUCKY	14.4	13.8	-0.6
LOUISIANA	22.8	20.4	-2.4
MAINE	14.2	12.0	-2.2
MARYLAND	9.7	10.2	0.5
MASSACHUSETTS	10.2	9.3	-0.9
MICHIGAN	9.6	9.9	0.4
MINNESOTA	7.2	8.4	1.2
MISSISSIPPI	14.2	15.2	0.9
MISSOURI	12.9	12.2	-0.8
MONTANA	15.6	14.6	-1.0
NEBRASKA	8.6	7.6	-1.0
NEVADA	16.4	14.1	-2.4
NEW HAMPSHIRE	11.9	10.3	-1.6
NEW JERSEY	11.1	11.7	0.5
NEW MEXICO	21.0	22.2	1.2
NEW YORK	12.4	14.0	1.6
NORTH CAROLINA	12.9	14.7	1.8
NORTH DAKOTA	11.8	11.7	-0.0
OHIO	11.4	10.5	-0.9
OKLAHOMA	18.5	17.0	-1.5
OREGON	13.2	12.5	-0.7
PENNSYLVANIA	11.1	9.4	-1.7
RHODE ISLAND	10.5	10.7	0.2
SOUTH CAROLINA	16.3	15.0	-1.3
SOUTH DAKOTA	10.1	13.2	3.1
TENNESSEE	12.0	11.7	-0.3
TEXAS	24.6	24.2	-0.4
UTAH	12.2	11.2	-1.0
VERMONT	13.0	14.4	1.4
VIRGINIA	14.6	11.3	-3.3
WASHINGTON	11.9	10.9	-1.0
WEST VIRGINIA	17.7	18.1	0.4
WISCONSIN	9.3	9.3	0.0
WYOMING	18.3	19.0	0.6
PUERTO RICO	13.6	10.3	-3.3

* HAVING NO HEALTH CARE PLAN
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 6.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1710	17.3	1.1	(15.2, 19.3)
ALASKA	1374	22.8	1.7	(19.5, 26.2)
ARIZONA	1423	17.9	1.4	(15.1, 20.6)
ARKANSAS	1374	21.6	1.4	(18.7, 24.4)
CALIFORNIA	3319	22.7	0.9	(21.0, 24.5)
COLORADO	1491	14.0	1.1	(11.8, 16.1)
CONNECTICUT	1777	10.8	1.0	(8.9, 12.8)
DELAWARE	1986	13.0	0.9	(11.2, 14.9)
DISTRICT OF CO	1255	13.2	1.2	(10.9, 15.6)
FLORIDA	2519	22.6	1.0	(20.6, 24.5)
GEORGIA	1948	13.7	1.0	(11.7, 15.6)
HAWAII	1767	7.4	0.8	(5.9, 8.9)
IDAHO	3958	20.1	0.9	(18.3, 22.0)
ILLINOIS	2387	13.5	0.8	(11.9, 15.1)
INDIANA	1932	14.6	1.0	(12.7, 16.5)
IOWA	2725	12.0	0.7	(10.6, 13.5)
KANSAS	1561	11.5	0.9	(9.7, 13.3)
KENTUCKY	2756	16.8	0.9	(15.2, 18.5)
LOUISIANA	1342	24.4	1.4	(21.6, 27.2)
MAINE	1381	14.3	1.1	(12.2, 16.4)
MARYLAND	3677	11.7	0.7	(10.3, 13.2)
MASSACHUSETTS	1420	11.0	1.0	(9.0, 12.9)
MICHIGAN	2138	11.8	0.8	(10.2, 13.4)
MINNESOTA	3887	9.6	0.5	(8.5, 10.6)
MISSISSIPPI	1258	18.3	1.3	(15.7, 20.9)
MISSOURI	1462	15.0	1.2	(12.7, 17.3)
MONTANA	1409	17.9	1.1	(15.7, 20.0)
NEBRASKA	2016	9.5	0.8	(8.0, 11.0)
NEVADA	2122	16.7	1.7	(13.5, 19.9)
NEW HAMPSHIRE	1215	12.3	1.1	(10.1, 14.5)
NEW JERSEY	2102	14.0	1.0	(12.1, 15.8)
NEW MEXICO	1448	25.7	1.4	(23.0, 28.3)
NEW YORK	2786	16.9	0.9	(15.2, 18.6)
NORTH CAROLINA	2839	17.2	0.8	(15.6, 18.8)
NORTH DAKOTA	1351	14.4	1.1	(12.2, 16.6)
OHIO	2317	12.7	0.9	(11.0, 14.5)
OKLAHOMA	1258	20.9	1.4	(18.2, 23.6)
OREGON	2624	15.2	0.9	(13.5, 17.0)
PENNSYLVANIA	2827	11.5	0.7	(10.1, 12.9)
RHODE ISLAND	1427	13.1	1.1	(10.8, 15.3)
SOUTH CAROLINA	1660	17.6	1.2	(15.3, 19.9)
SOUTH DAKOTA	1633	16.5	1.1	(14.3, 18.6)
TENNESSEE	2376	13.9	0.9	(12.2, 15.6)
TEXAS	2106	28.2	1.1	(25.9, 30.4)
UTAH	2421	12.8	0.9	(11.0, 14.5)
VERMONT	2566	16.5	0.9	(14.8, 18.2)
VIRGINIA	2916	12.8	0.8	(11.2, 14.3)
WASHINGTON	2998	13.0	0.8	(11.5, 14.5)
WEST VIRGINIA	1828	22.4	1.1	(20.2, 24.6)
WISCONSIN	1820	11.2	1.0	(9.2, 13.3)
WYOMING	1994	22.4	1.4	(19.7, 25.2)
PUERTO RICO	1845	11.8	0.9	(10.1, 13.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 14.3 RANGE = 7.4-28.2 YEAR 2000 OBJECTIVE 21.4
 * HAVING NO HEALTH CARE PLAN August 14, 1998

TABLE 6.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE* - BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING DON'T KNOW AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	672	13.2	1.5	(10.3, 16.1)	*	1038	21.1	1.5	(18.3, 24.0)	*
ALASKA	637	23.4	2.5	(18.4, 28.3)	*	737	22.2	2.3	(17.8, 26.7)	*
ARIZONA	691	19.0	2.1	(15.0, 23.1)	*	732	16.7	1.9	(12.9, 20.5)	*
ARKANSAS	546	21.7	2.4	(17.0, 26.4)	*	828	21.4	1.6	(18.2, 24.6)	*
CALIFORNIA	1469	25.5	1.4	(22.8, 28.2)	*	1850	19.8	1.1	(17.7, 22.0)	*
COLORADO	693	15.7	1.6	(12.5, 18.8)	*	798	12.3	1.5	(9.4, 15.1)	*
CONNECTICUT	701	11.8	1.7	(8.4, 15.1)	*	1076	9.9	1.0	(7.9, 12.0)	*
DELAWARE	821	15.2	1.5	(12.3, 18.2)	*	1165	10.9	1.1	(8.8, 13.1)	*
DISTRICT OF CO	499	14.8	1.8	(11.2, 18.4)	*	756	11.8	1.4	(9.2, 14.5)	*
FLORIDA	1052	22.3	1.5	(19.3, 25.2)	*	1467	22.9	1.3	(20.4, 25.3)	*
GEORGIA	824	14.6	1.6	(11.4, 17.7)	*	1124	12.8	1.3	(10.4, 15.3)	*
HAWAII	790	7.4	1.0	(5.4, 9.4)	*	977	7.4	1.1	(5.2, 9.6)	*
IDAHO	1740	20.7	1.4	(17.9, 23.4)	*	2218	19.6	1.2	(17.2, 22.0)	*
ILLINOIS	1008	12.4	1.2	(10.0, 14.7)	*	1379	14.6	1.1	(12.4, 16.8)	*
INDIANA	903	14.8	1.5	(11.9, 17.7)	*	1029	14.4	1.3	(11.9, 16.8)	*
IOWA	1215	12.6	1.1	(10.4, 14.8)	*	1510	11.5	0.9	(9.7, 13.3)	*
KANSAS	657	11.7	1.5	(8.8, 14.6)	*	904	11.2	1.1	(9.1, 13.4)	*
KENTUCKY	1071	16.3	1.3	(13.8, 18.7)	*	1685	17.4	1.1	(15.3, 19.5)	*
LOUISIANA	521	24.8	2.3	(20.3, 29.3)	*	821	24.0	1.7	(20.7, 27.4)	*
MAINE	627	15.2	1.7	(11.9, 18.4)	*	754	13.5	1.4	(10.7, 16.2)	*
MARYLAND	1617	11.7	1.1	(9.6, 13.9)	*	2060	11.7	1.0	(9.9, 13.6)	*
MASSACHUSETTS	607	12.7	1.6	(9.6, 15.8)	*	813	9.3	1.2	(7.0, 11.6)	*
MICHIGAN	920	13.8	1.3	(11.3, 16.4)	*	1218	9.9	1.0	(8.0, 11.7)	*
MINNESOTA	1890	10.1	0.8	(8.6, 11.6)	*	1997	9.0	0.7	(7.6, 10.5)	*
MISSISSIPPI	498	18.7	2.2	(14.5, 22.9)	*	760	18.0	1.6	(14.8, 21.1)	*
MISSOURI	567	16.7	2.0	(12.9, 20.5)	*	895	13.4	1.3	(10.8, 15.9)	*
MONTANA	613	16.9	1.6	(13.7, 20.1)	*	796	18.8	1.5	(15.9, 21.7)	*
NEBRASKA	823	9.3	1.2	(7.0, 11.7)	*	1193	9.6	1.0	(7.6, 11.6)	*
NEVADA	1012	16.1	2.6	(11.1, 21.1)	*	1110	17.3	2.1	(13.3, 21.3)	*
NEW HAMPSHIRE	487	12.6	1.8	(9.1, 16.1)	*	728	12.0	1.3	(9.5, 14.5)	*
NEW JERSEY	861	14.8	1.6	(11.7, 17.9)	*	1241	13.2	1.1	(11.0, 15.3)	*
NEW MEXICO	622	24.0	2.0	(20.1, 27.8)	*	826	27.3	1.8	(23.8, 30.9)	*
NEW YORK	1146	18.7	1.4	(16.0, 21.4)	*	1640	15.2	1.0	(13.2, 17.3)	*
NORTH CAROLINA	1191	18.4	1.3	(15.9, 20.8)	*	1648	16.1	1.0	(14.0, 18.1)	*
NORTH DAKOTA	627	14.5	1.6	(11.4, 17.6)	*	724	14.2	1.4	(11.4, 17.0)	*
OHIO	926	12.5	1.3	(9.9, 15.1)	*	1391	12.9	1.2	(10.6, 15.2)	*
OKLAHOMA	515	18.1	2.0	(14.1, 22.1)	*	743	23.7	1.9	(20.1, 27.3)	*
OREGON	1161	17.4	1.5	(14.5, 20.3)	*	1463	13.1	1.0	(11.1, 15.0)	*
PENNSYLVANIA	1170	12.7	1.1	(10.5, 15.0)	*	1657	10.3	0.9	(8.6, 12.1)	*
RHODE ISLAND	612	13.3	1.6	(10.3, 16.4)	*	815	12.8	1.7	(9.6, 16.0)	*
SOUTH CAROLINA	665	17.9	1.8	(14.4, 21.4)	*	995	17.3	1.4	(14.5, 20.0)	*
SOUTH DAKOTA	751	16.0	1.6	(12.9, 19.0)	*	882	17.0	1.4	(14.2, 19.7)	*
TENNESSEE	878	15.5	1.4	(12.7, 18.3)	*	1498	12.3	1.0	(10.4, 14.2)	*
TEXAS	892	29.3	1.8	(25.9, 32.8)	*	1214	27.0	1.5	(24.2, 29.9)	*
UTAH	1103	13.5	1.3	(11.0, 16.0)	*	1318	12.1	1.2	(9.6, 14.5)	*
VERMONT	1165	18.1	1.4	(15.5, 20.8)	*	1401	14.8	1.1	(12.7, 17.0)	*
VIRGINIA	1222	11.9	1.2	(9.6, 14.2)	*	1694	13.6	1.1	(11.5, 15.7)	*
WASHINGTON	1339	13.6	1.1	(11.5, 15.7)	*	1659	12.5	1.1	(10.4, 14.6)	*
WEST VIRGINIA	779	22.0	1.7	(18.7, 25.3)	*	1049	22.8	1.5	(19.9, 25.7)	*
WISCONSIN	858	12.5	1.6	(9.4, 15.7)	*	962	9.9	1.3	(7.4, 12.5)	*
WYOMING	875	23.8	2.3	(19.4, 28.3)	*	1119	21.0	1.3	(18.5, 23.6)	*
PUERTO RICO	776	13.6	1.4	(11.0, 16.2)	*	1069	10.2	1.1	(8.0, 12.4)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 15.2	MEDIAN	= 13.5
	RANGE	= 7.4-29.3	RANGE	= 7.4-27.3

* HAVING NO HEALTH CARE PLAN

August 14, 1998

TABLE 6.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 NO HEALTH INSURANCE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING DON'T KNOW AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	18.8	17.3	-1.5
ALASKA	21.8	22.8	1.0
ARIZONA	20.5	17.9	-2.6
ARKANSAS	21.7	21.6	-0.1
CALIFORNIA	22.6	22.7	0.1
COLORADO	15.3	14.0	-1.4
CONNECTICUT	11.0	10.8	-0.1
DELAWARE	13.0	13.0	0.0
DISTRICT OF CO	15.4	13.2	-2.1
FLORIDA	22.9	22.6	-0.4
GEORGIA	11.0	13.7	2.7
HAWAII	9.5	7.4	-2.1
IDAHO	18.2	20.1	2.0
ILLINOIS	14.1	13.5	-0.6
INDIANA	13.6	14.6	1.0
IOWA	11.6	12.0	0.4
KANSAS	12.3	11.5	-0.8
KENTUCKY	17.4	16.8	-0.6
LOUISIANA	26.3	24.4	-1.9
MAINE	17.2	14.3	-2.9
MARYLAND	11.4	11.7	0.3
MASSACHUSETTS	12.5	11.0	-1.5
MICHIGAN	11.1	11.8	0.7
MINNESOTA	8.0	9.6	1.5
MISSISSIPPI	17.3	18.3	1.1
MISSOURI	15.9	15.0	-0.9
MONTANA	19.2	17.9	-1.3
NEBRASKA	10.7	9.5	-1.2
NEVADA	19.2	16.7	-2.5
NEW HAMPSHIRE	14.1	12.3	-1.8
NEW JERSEY	13.1	14.0	0.9
NEW MEXICO	24.6	25.7	1.1
NEW YORK	15.0	16.9	1.9
NORTH CAROLINA	15.0	17.2	2.2
NORTH DAKOTA	14.4	14.4	-0.1
OHIO	13.5	12.7	-0.8
OKLAHOMA	23.4	20.9	-2.5
OREGON	15.9	15.2	-0.7
PENNSYLVANIA	13.5	11.5	-1.9
RHODE ISLAND	13.0	13.1	0.1
SOUTH CAROLINA	18.9	17.6	-1.3
SOUTH DAKOTA	12.7	16.5	3.7
TENNESSEE	13.9	13.9	-0.0
TEXAS	28.3	28.2	-0.1
UTAH	14.0	12.8	-1.3
VERMONT	15.3	16.5	1.2
VIRGINIA	16.5	12.8	-3.7
WASHINGTON	14.0	13.0	-1.0
WEST VIRGINIA	22.2	22.4	0.3
WISCONSIN	11.4	11.2	-0.2
WYOMING	21.6	22.4	0.9
PUERTO RICO	15.4	11.8	-3.6

* HAVING NO HEALTH CARE PLAN
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 7.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2175	7.0	0.6	(5.8, 8.1)
ALASKA	1544	3.3	0.6	(2.1, 4.6)
ARIZONA	1898	3.1	0.6	(2.0, 4.2)
ARKANSAS	1790	5.1	0.6	(4.0, 6.2)
CALIFORNIA	4063	5.6	0.4	(4.7, 6.4)
COLORADO	1826	3.9	0.5	(2.8, 4.9)
CONNECTICUT	2246	5.2	0.6	(4.0, 6.3)
DELAWARE	2548	6.4	0.5	(5.4, 7.5)
DISTRICT OF CO	1495	4.6	0.6	(3.3, 5.8)
FLORIDA	3488	5.7	0.4	(4.9, 6.6)
GEORGIA	2304	4.1	0.5	(3.2, 5.0)
HAWAII	2165	5.0	0.6	(3.8, 6.1)
IDAHO	4921	4.0	0.3	(3.4, 4.7)
ILLINOIS	2884	7.0	0.6	(5.9, 8.2)
INDIANA	2345	5.3	0.5	(4.2, 6.3)
IOWA	3597	4.6	0.4	(3.9, 5.4)
KANSAS	2004	3.0	0.4	(2.2, 3.8)
KENTUCKY	3612	5.3	0.4	(4.5, 6.0)
LOUISIANA	1653	5.5	0.6	(4.3, 6.7)
MAINE	1698	4.9	0.6	(3.8, 6.0)
MARYLAND	4568	5.9	0.5	(5.0, 6.8)
MASSACHUSETTS	1764	4.7	0.6	(3.5, 5.9)
MICHIGAN	2551	5.8	0.5	(4.9, 6.8)
MINNESOTA	4783	3.9	0.3	(3.3, 4.5)
MISSISSIPPI	1599	6.1	0.7	(4.8, 7.4)
MISSOURI	1849	4.8	0.6	(3.7, 5.9)
MONTANA	1802	3.2	0.4	(2.3, 4.0)
NEBRASKA	2697	4.2	0.4	(3.3, 5.0)
NEVADA	2505	4.0	0.9	(2.3, 5.6)
NEW HAMPSHIRE	1512	3.9	0.5	(2.9, 5.0)
NEW JERSEY	2664	5.3	0.5	(4.3, 6.2)
NEW MEXICO	1809	4.9	0.5	(3.9, 6.0)
NEW YORK	3402	4.8	0.4	(4.0, 5.6)
NORTH CAROLINA	3628	5.0	0.4	(4.2, 5.7)
NORTH DAKOTA	1800	3.5	0.5	(2.6, 4.4)
OHIO	3162	4.7	0.5	(3.9, 5.6)
OKLAHOMA	1884	5.9	0.6	(4.8, 7.1)
OREGON	3316	4.7	0.4	(3.9, 5.5)
PENNSYLVANIA	3595	5.1	0.4	(4.3, 5.9)
RHODE ISLAND	1839	4.9	0.5	(3.8, 5.9)
SOUTH CAROLINA	2144	4.9	0.5	(3.9, 5.9)
SOUTH DAKOTA	2197	3.8	0.5	(2.9, 4.7)
TENNESSEE	2968	4.4	0.4	(3.6, 5.1)
TEXAS	2500	5.9	0.5	(4.8, 6.9)
UTAH	2864	4.1	0.5	(3.1, 5.1)
VERMONT	3184	4.7	0.6	(3.6, 5.8)
VIRGINIA	3519	4.2	0.4	(3.5, 4.9)
WASHINGTON	3600	4.1	0.4	(3.4, 4.8)
WEST VIRGINIA	2438	6.3	0.5	(5.3, 7.3)
WISCONSIN	2245	4.7	0.6	(3.4, 5.9)
WYOMING	2411	3.0	0.4	(2.3, 3.8)
PUERTO RICO	2238	10.5	0.7	(9.1, 11.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 4.8 RANGE = 3.0-10.5

* EVER TOLD YOU HAVE DIABETES

August 14, 1998

TABLE 7.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	813	5.7	0.8	(4.0, 7.3)	*	1362	8.1	0.8	(6.5, 9.7)	
ALASKA	717	2.9	0.9	(1.2, 4.6)	*	827	3.8	0.9	(1.9, 5.6)	
ARIZONA	885	2.3	0.6	(1.2, 3.4)	*	1013	3.9	0.9	(2.0, 5.7)	
ARKANSAS	674	5.7	1.0	(3.9, 7.6)	*	1116	4.6	0.7	(3.3, 5.9)	
CALIFORNIA	1759	5.4	0.6	(4.2, 6.7)	*	2304	5.7	0.6	(4.6, 6.9)	
COLORADO	820	3.9	0.7	(2.4, 5.3)	*	1006	3.9	0.8	(2.3, 5.5)	
CONNECTICUT	886	5.1	0.8	(3.5, 6.7)	*	1360	5.2	0.8	(3.6, 6.8)	
DELAWARE	1008	5.9	0.8	(4.4, 7.4)	*	1540	6.9	0.7	(5.5, 8.4)	
DISTRICT OF CO	563	3.9	0.9	(2.1, 5.7)	*	932	5.1	0.8	(3.5, 6.7)	
FLORIDA	1431	6.8	0.7	(5.4, 8.2)	*	2057	4.8	0.5	(3.8, 5.7)	
GEORGIA	958	2.7	0.5	(1.7, 3.8)	*	1346	5.4	0.7	(3.9, 6.8)	
HAWAII	947	4.3	0.8	(2.7, 5.9)	*	1218	5.6	0.8	(4.1, 7.2)	
IDAHO	2057	4.5	0.5	(3.5, 5.6)	*	2864	3.5	0.4	(2.8, 4.3)	
ILLINOIS	1163	7.0	0.9	(5.2, 8.8)	*	1721	7.1	0.7	(5.6, 8.5)	
INDIANA	1041	5.0	0.8	(3.4, 6.6)	*	1304	5.5	0.7	(4.1, 6.8)	
IOWA	1477	4.3	0.6	(3.2, 5.5)	*	2120	4.9	0.5	(4.0, 5.9)	
KANSAS	787	2.6	0.6	(1.5, 3.8)	*	1217	3.4	0.5	(2.3, 4.4)	
KENTUCKY	1325	5.6	0.7	(4.3, 6.9)	*	2287	5.0	0.5	(4.1, 5.9)	
LOUISIANA	626	5.5	1.0	(3.7, 7.4)	*	1027	5.5	0.8	(3.9, 7.0)	
MAINE	750	4.9	0.8	(3.3, 6.5)	*	948	4.9	0.8	(3.4, 6.5)	
MARYLAND	1954	5.7	0.7	(4.4, 7.1)	*	2614	6.0	0.6	(4.8, 7.2)	
MASSACHUSETTS	719	5.0	0.9	(3.1, 6.8)	*	1045	4.5	0.8	(3.0, 5.9)	
MICHIGAN	1059	5.2	0.7	(3.8, 6.6)	*	1492	6.4	0.7	(5.1, 7.7)	
MINNESOTA	2173	4.0	0.4	(3.1, 4.9)	*	2610	3.8	0.4	(3.1, 4.6)	
MISSISSIPPI	592	5.1	1.0	(3.2, 7.1)	*	1007	6.9	0.9	(5.3, 8.6)	
MISSOURI	698	5.4	0.9	(3.6, 7.2)	*	1151	4.2	0.6	(3.0, 5.4)	
MONTANA	760	3.5	0.7	(2.2, 4.8)	*	1042	2.8	0.5	(1.8, 3.8)	
NEBRASKA	1046	3.7	0.6	(2.4, 5.0)	*	1651	4.6	0.6	(3.4, 5.7)	
NEVADA	1165	5.4	1.6	(2.3, 8.4)	*	1340	2.6	0.7	(1.3, 3.9)	
NEW HAMPSHIRE	604	3.8	0.8	(2.2, 5.4)	*	908	4.1	0.7	(2.7, 5.5)	
NEW JERSEY	1060	5.5	0.7	(4.1, 6.9)	*	1604	5.1	0.7	(3.7, 6.4)	
NEW MEXICO	752	5.3	0.8	(3.7, 6.9)	*	1057	4.6	0.7	(3.2, 5.9)	
NEW YORK	1355	5.2	0.7	(3.8, 6.5)	*	2047	4.5	0.5	(3.4, 5.5)	
NORTH CAROLINA	1436	4.5	0.5	(3.4, 5.5)	*	2192	5.5	0.5	(4.5, 6.5)	
NORTH DAKOTA	799	3.6	0.7	(2.2, 4.9)	*	1001	3.5	0.6	(2.4, 4.6)	
OHIO	1228	3.9	0.6	(2.7, 5.1)	*	1934	5.5	0.7	(4.2, 6.8)	
OKLAHOMA	830	5.5	0.8	(3.8, 7.1)	*	1054	6.4	0.8	(4.7, 8.0)	
OREGON	1397	4.5	0.6	(3.3, 5.7)	*	1919	4.9	0.6	(3.8, 6.0)	
PENNSYLVANIA	1439	5.1	0.6	(3.9, 6.3)	*	2156	5.1	0.5	(4.0, 6.1)	
RHODE ISLAND	754	3.8	0.7	(2.4, 5.2)	*	1085	5.9	0.8	(4.3, 7.4)	
SOUTH CAROLINA	810	4.6	0.8	(3.1, 6.2)	*	1334	5.1	0.6	(4.0, 6.3)	
SOUTH DAKOTA	955	3.5	0.7	(2.1, 4.8)	*	1242	4.2	0.6	(3.0, 5.3)	
TENNESSEE	1032	3.4	0.5	(2.4, 4.5)	*	1936	5.2	0.6	(4.1, 6.3)	
TEXAS	1025	5.0	0.8	(3.5, 6.5)	*	1475	6.7	0.7	(5.3, 8.2)	
UTAH	1276	4.6	0.8	(3.1, 6.1)	*	1588	3.7	0.7	(2.4, 5.0)	
VERMONT	1389	4.9	1.0	(2.9, 6.9)	*	1795	4.5	0.6	(3.4, 5.6)	
VIRGINIA	1416	3.8	0.6	(2.7, 5.0)	*	2103	4.5	0.5	(3.6, 5.5)	
WASHINGTON	1548	3.3	0.5	(2.4, 4.2)	*	2052	4.9	0.5	(3.9, 6.0)	
WEST VIRGINIA	966	7.2	0.9	(5.4, 8.9)	*	1472	5.6	0.6	(4.3, 6.8)	
WISCONSIN	991	5.2	1.1	(3.0, 7.5)	*	1254	4.2	0.6	(2.9, 5.4)	
WYOMING	1019	2.5	0.5	(1.5, 3.5)	*	1392	3.6	0.5	(2.5, 4.6)	
PUERTO RICO	919	9.9	1.0	(8.0, 11.8)	*	1319	11.0	0.9	(9.2, 12.7)	

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 4.9	MEDIAN	= 4.9
	RANGE	= 2.3- 9.9	RANGE	= 2.6-11.0

* EVER TOLD YOU HAVE DIABETES

August 14, 1998

TABLE 7.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	7.3	7.0	-0.4
ALASKA	3.5	3.3	-0.2
ARIZONA	3.6	3.1	-0.4
ARKANSAS	4.6	5.1	0.5
CALIFORNIA	5.1	5.6	0.5
COLORADO	3.0	3.9	0.9
CONNECTICUT	4.4	5.2	0.8
DELAWARE	5.3	6.4	1.1
DISTRICT OF CO	5.6	4.6	-1.0
FLORIDA	5.2	5.7	0.5
GEORGIA	4.1	4.1	-0.0
HAWAII	4.6	5.0	0.3
IDAHO	3.5	4.0	0.6
ILLINOIS	5.8	7.0	1.3
INDIANA	5.3	5.3	-0.1
IOWA	4.2	4.6	0.4
KANSAS	3.6	3.0	-0.6
KENTUCKY	4.5	5.3	0.8
LOUISIANA	6.6	5.5	-1.1
MAINE	4.2	4.9	0.7
MARYLAND	5.1	5.9	0.8
MASSACHUSETTS	5.3	4.7	-0.5
MICHIGAN	5.8	5.8	0.0
MINNESOTA	4.5	3.9	-0.6
MISSISSIPPI	6.1	6.1	-0.0
MISSOURI	4.4	4.8	0.4
MONTANA	3.7	3.2	-0.6
NEBRASKA	5.3	4.2	-1.1
NEVADA	4.2	4.0	-0.2
NEW HAMPSHIRE	4.0	3.9	-0.1
NEW JERSEY	4.6	5.3	0.7
NEW MEXICO	4.4	4.9	0.5
NEW YORK	3.6	4.8	1.2
NORTH CAROLINA	4.4	5.0	0.5
NORTH DAKOTA	3.1	3.5	0.5
OHIO	5.0	4.7	-0.2
OKLAHOMA	3.8	5.9	2.2
OREGON	4.5	4.7	0.3
PENNSYLVANIA	5.7	5.1	-0.6
RHODE ISLAND	4.5	4.9	0.4
SOUTH CAROLINA	5.4	4.9	-0.5
SOUTH DAKOTA	4.3	3.8	-0.4
TENNESSEE	5.0	4.4	-0.6
TEXAS	3.7	5.9	2.1
UTAH	3.3	4.1	0.9
VERMONT	3.9	4.7	0.8
VIRGINIA	5.5	4.2	-1.3
WASHINGTON	3.4	4.1	0.7
WEST VIRGINIA	5.9	6.3	0.5
WISCONSIN	4.7	4.7	-0.0
WYOMING	4.2	3.0	-1.2
PUERTO RICO	10.8	10.5	-0.3

* EVER TOLD YOU HAVE DIABETES
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 8.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOLD BLOOD PRESSURE IS HIGH*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER CHECKED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2170	28.9	1.1	(26.8, 31.1)
ALASKA	1539	22.6	1.6	(19.5, 25.6)
ARIZONA	1867	16.3	1.1	(14.2, 18.5)
ARKANSAS	1784	26.3	1.2	(24.0, 28.7)
CALIFORNIA	4041	21.3	0.7	(19.9, 22.6)
COLORADO	1819	20.4	1.0	(18.4, 22.4)
CONNECTICUT	2235	20.6	1.0	(18.7, 22.5)
DELAWARE	2547	25.5	1.0	(23.6, 27.5)
DISTRICT OF CO	1501	19.4	1.2	(17.0, 21.8)
FLORIDA	3473	26.1	0.8	(24.4, 27.7)
GEORGIA	2307	21.4	1.0	(19.5, 23.3)
HAWAII	2161	24.0	1.1	(21.8, 26.1)
IDAHO	4900	24.1	0.8	(22.5, 25.6)
ILLINOIS	1437	24.3	1.2	(21.9, 26.7)
INDIANA	2364	25.2	1.0	(23.2, 27.1)
IOWA	3593	23.4	0.8	(21.8, 25.0)
KANSAS	1996	20.9	1.0	(18.9, 22.8)
KENTUCKY	3585	27.1	0.8	(25.5, 28.7)
LOUISIANA	1649	25.1	1.2	(22.7, 27.5)
MAINE	1691	22.8	1.1	(20.7, 25.0)
MARYLAND	4566	23.8	0.8	(22.2, 25.4)
MASSACHUSETTS	1757	19.8	1.1	(17.7, 21.9)
MICHIGAN	2561	23.3	0.9	(21.5, 25.1)
MINNESOTA	4793	21.2	0.6	(19.9, 22.4)
MISSISSIPPI	1589	34.4	1.4	(31.7, 37.1)
MISSOURI	1843	27.3	1.2	(25.0, 29.6)
MONTANA	1798	22.9	1.0	(20.9, 24.9)
NEBRASKA	2690	22.4	0.9	(20.6, 24.2)
NEVADA	2494	24.1	1.7	(20.8, 27.4)
NEW HAMPSHIRE	1508	22.6	1.1	(20.4, 24.9)
NEW JERSEY	2650	23.6	1.0	(21.7, 25.5)
NEW MEXICO	1803	21.3	1.1	(19.2, 23.4)
NEW YORK	3379	22.7	0.8	(21.1, 24.3)
NORTH CAROLINA	3615	23.3	0.8	(21.8, 24.8)
NORTH DAKOTA	1799	25.5	1.1	(23.3, 27.6)
OHIO	3155	22.0	1.0	(20.1, 23.9)
OKLAHOMA	1882	21.7	1.1	(19.6, 23.8)
OREGON	3309	22.8	0.9	(21.1, 24.5)
PENNSYLVANIA	3584	21.7	0.8	(20.2, 23.2)
RHODE ISLAND	1834	22.5	1.1	(20.4, 24.5)
SOUTH CAROLINA	2148	26.9	1.0	(24.8, 28.9)
SOUTH DAKOTA	2194	20.6	1.0	(18.7, 22.5)
TENNESSEE	2978	27.8	0.9	(25.9, 29.6)
TEXAS	2483	23.1	0.9	(21.3, 25.0)
UTAH	2860	22.5	1.1	(20.4, 24.7)
VERMONT	3177	20.9	0.9	(19.2, 22.6)
VIRGINIA	3513	24.5	1.0	(22.5, 26.4)
WASHINGTON	3584	23.2	0.8	(21.5, 24.8)
WEST VIRGINIA	2431	28.3	1.0	(26.4, 30.2)
WISCONSIN	2240	23.1	1.1	(21.0, 25.3)
WYOMING	2404	22.1	1.0	(20.2, 24.0)
PUERTO RICO	2235	20.9	0.9	(19.1, 22.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 23.0 RANGE = 16.3-34.4

* EVER TOLD BLOOD PRESSURE IS HIGH

August 14, 1998

TABLE 8.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOLD BLOOD PRESSURE IS HIGH* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER CHECKED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	809	24.8	1.6	(21.6, 28.1)	*	1361	32.5	1.4	(29.7, 35.3)	
ALASKA	715	20.3	2.1	(16.2, 24.4)	*	824	25.1	2.3	(20.6, 29.5)	
ARIZONA	877	15.3	1.4	(12.5, 18.1)	*	990	17.3	1.6	(14.2, 20.5)	
ARKANSAS	665	24.7	1.9	(21.0, 28.5)	*	1119	27.8	1.5	(24.8, 30.7)	
CALIFORNIA	1746	20.5	1.0	(18.5, 22.5)	*	2295	22.0	1.0	(20.1, 23.8)	
COLORADO	812	19.3	1.5	(16.4, 22.1)	*	1007	21.5	1.4	(18.6, 24.3)	
CONNECTICUT	880	20.5	1.5	(17.5, 23.5)	*	1355	20.7	1.3	(18.3, 23.2)	
DELAWARE	1008	24.3	1.6	(21.2, 27.5)	*	1539	26.6	1.3	(24.1, 29.1)	
DISTRICT OF CO	570	16.7	1.7	(13.3, 20.1)	*	931	21.6	1.6	(18.5, 24.8)	
FLORIDA	1422	25.2	1.3	(22.7, 27.7)	*	2051	26.8	1.1	(24.7, 28.9)	
GEORGIA	962	18.4	1.3	(15.8, 20.9)	*	1345	24.2	1.4	(21.4, 27.0)	
HAWAII	947	24.6	1.6	(21.4, 27.8)	*	1214	23.3	1.5	(20.4, 26.2)	
IDAHO	2046	24.5	1.2	(22.1, 26.9)	*	2854	23.7	1.0	(21.6, 25.7)	
ILLINOIS	574	22.7	1.9	(19.1, 26.4)	*	863	25.6	1.6	(22.5, 28.8)	
INDIANA	1061	23.8	1.5	(21.0, 26.7)	*	1303	26.4	1.4	(23.7, 29.0)	
IOWA	1470	22.4	1.2	(20.0, 24.8)	*	2123	24.3	1.0	(22.2, 26.3)	
KANSAS	784	19.0	1.5	(16.1, 22.0)	*	1212	22.6	1.3	(20.1, 25.1)	
KENTUCKY	1308	26.5	1.3	(24.0, 29.0)	*	2277	27.7	1.0	(25.7, 29.6)	
LOUISIANA	626	23.1	1.8	(19.5, 26.6)	*	1023	26.9	1.6	(23.9, 30.0)	
MAINE	746	22.3	1.6	(19.1, 25.5)	*	945	23.3	1.5	(20.4, 26.2)	
MARYLAND	1954	22.9	1.2	(20.5, 25.4)	*	2612	24.6	1.1	(22.4, 26.7)	
MASSACHUSETTS	716	20.1	1.7	(16.9, 23.3)	*	1041	19.5	1.4	(16.7, 22.3)	
MICHIGAN	1068	22.4	1.4	(19.7, 25.2)	*	1493	24.2	1.2	(21.8, 26.5)	
MINNESOTA	2188	20.1	0.9	(18.3, 21.9)	*	2605	22.2	0.9	(20.5, 23.9)	
MISSISSIPPI	586	33.1	2.2	(28.7, 37.5)	*	1003	35.5	1.7	(32.3, 38.8)	
MISSOURI	694	25.0	1.9	(21.3, 28.6)	*	1149	29.4	1.6	(26.3, 32.4)	
MONTANA	758	22.2	1.6	(19.2, 25.3)	*	1040	23.5	1.4	(20.8, 26.2)	
NEBRASKA	1046	21.5	1.5	(18.6, 24.4)	*	1644	23.1	1.2	(20.8, 25.4)	
NEVADA	1157	26.4	2.6	(21.2, 31.5)	*	1337	21.8	2.1	(17.7, 25.9)	
NEW HAMPSHIRE	602	23.4	1.9	(19.8, 27.0)	*	906	21.9	1.5	(19.0, 24.8)	
NEW JERSEY	1051	24.0	1.5	(21.0, 27.0)	*	1599	23.3	1.2	(20.9, 25.7)	
NEW MEXICO	749	22.7	1.6	(19.6, 25.8)	*	1054	20.0	1.3	(17.4, 22.6)	
NEW YORK	1343	21.6	1.3	(19.1, 24.0)	*	2036	23.7	1.0	(21.7, 25.7)	
NORTH CAROLINA	1429	21.0	1.1	(18.7, 23.2)	*	2186	25.5	1.1	(23.4, 27.6)	
NORTH DAKOTA	799	25.6	1.6	(22.4, 28.8)	*	1000	25.4	1.5	(22.4, 28.3)	
OHIO	1223	21.2	1.5	(18.3, 24.1)	*	1932	22.7	1.2	(20.3, 25.1)	
OKLAHOMA	828	20.8	1.6	(17.7, 23.9)	*	1054	22.5	1.4	(19.8, 25.3)	
OREGON	1396	22.3	1.3	(19.8, 24.8)	*	1913	23.3	1.1	(21.1, 25.5)	
PENNSYLVANIA	1434	20.7	1.2	(18.4, 23.1)	*	2150	22.5	1.0	(20.5, 24.5)	
RHODE ISLAND	752	20.4	1.6	(17.3, 23.5)	*	1082	24.3	1.5	(21.4, 27.1)	
SOUTH CAROLINA	815	25.5	1.6	(22.3, 28.7)	*	1333	28.0	1.4	(25.3, 30.7)	
SOUTH DAKOTA	952	18.4	1.4	(15.6, 21.1)	*	1242	22.7	1.3	(20.1, 25.3)	
TENNESSEE	1043	24.6	1.5	(21.7, 27.5)	*	1935	30.6	1.2	(28.3, 32.9)	
TEXAS	1012	20.8	1.4	(18.1, 23.5)	*	1471	25.3	1.3	(22.8, 27.7)	
UTAH	1274	22.3	1.6	(19.2, 25.4)	*	1586	22.8	1.4	(20.0, 25.5)	
VERMONT	1389	21.6	1.4	(18.9, 24.3)	*	1788	20.3	1.1	(18.2, 22.5)	
VIRGINIA	1413	23.9	1.6	(20.8, 27.0)	*	2100	25.0	1.3	(22.6, 27.5)	
WASHINGTON	1541	21.3	1.2	(18.8, 23.7)	*	2043	25.0	1.1	(22.8, 27.1)	
WEST VIRGINIA	960	28.1	1.5	(25.1, 31.1)	*	1471	28.5	1.3	(26.1, 31.0)	
WISCONSIN	986	24.3	1.7	(20.9, 27.7)	*	1254	22.1	1.4	(19.4, 24.8)	
WYOMING	1015	21.1	1.4	(18.3, 23.9)	*	1389	23.1	1.3	(20.5, 25.7)	
PUERTO RICO	920	19.2	1.4	(16.5, 21.9)	*	1315	22.3	1.2	(19.9, 24.7)	

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 22.3
 RANGE = 15.3-33.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 23.6
 RANGE = 17.3-35.5

* EVER TOLD BLOOD PRESSURE IS HIGH

August 14, 1998

TABLE 8.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 TOLD BLOOD PRESSURE IS HIGH*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER CHECKED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	24.8	28.9	4.2
ALASKA	19.3	22.6	3.3
ARIZONA	20.4	16.3	-4.1
ARKANSAS	26.7	26.3	-0.3
CALIFORNIA	22.1	21.3	-0.9
COLORADO	21.5	20.4	-1.1
CONNECTICUT	19.2	20.6	1.4
DELAWARE	21.3	25.5	4.2
DISTRICT OF CO	N/A	19.4	N/A
FLORIDA	23.3	26.1	2.7
GEORGIA	19.7	21.4	1.7
HAWAII	21.1	24.0	2.8
IDAHO	21.6	24.1	2.5
ILLINOIS	22.2	24.3	2.1
INDIANA	26.1	25.2	-1.0
IOWA	23.6	23.4	-0.2
KANSAS	23.4	20.9	-2.5
KENTUCKY	21.7	27.1	5.4
LOUISIANA	24.2	25.1	0.9
MAINE	20.9	22.8	1.9
MARYLAND	20.8	23.8	3.0
MASSACHUSETTS	23.7	19.8	-3.8
MICHIGAN	24.4	23.3	-1.0
MINNESOTA	18.6	21.2	2.6
MISSISSIPPI	29.9	34.4	4.5
MISSOURI	23.1	27.3	4.3
MONTANA	19.5	22.9	3.4
NEBRASKA	21.8	22.4	0.6
NEVADA	21.8	24.1	2.3
NEW HAMPSHIRE	20.3	22.6	2.3
NEW JERSEY	25.9	23.6	-2.2
NEW MEXICO	19.8	21.3	1.5
NEW YORK	22.2	22.7	0.6
NORTH CAROLINA	20.7	23.3	2.6
NORTH DAKOTA	22.3	25.5	3.2
OHIO	23.6	22.0	-1.6
OKLAHOMA	21.6	21.7	0.1
OREGON	24.0	22.8	-1.2
PENNSYLVANIA	24.5	21.7	-2.8
RHODE ISLAND	23.3	22.5	-0.9
SOUTH CAROLINA	23.5	26.9	3.4
SOUTH DAKOTA	20.7	20.6	-0.1
TENNESSEE	26.8	27.8	0.9
TEXAS	24.0	23.1	-0.9
UTAH	19.2	22.5	3.3
VERMONT	22.8	20.9	-1.9
VIRGINIA	23.1	24.5	1.3
WASHINGTON	20.9	23.2	2.2
WEST VIRGINIA	25.5	28.3	2.8
WISCONSIN	22.8	23.1	0.4
WYOMING	21.2	22.1	0.9
PUERTO RICO	N/A	20.9	N/A

* EVER TOLD BLOOD PRESSURE IS HIGH

N/A - NOT AVAILABLE

August 14, 1998

TABLE 9.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BP CHECKED WITHIN LAST 2 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2150	95.4	0.5	(94.5, 96.4)
ALASKA	1531	91.8	1.1	(89.6, 94.1)
ARIZONA	1892	93.5	0.8	(91.9, 95.2)
ARKANSAS	1775	93.3	0.7	(91.9, 94.8)
CALIFORNIA	4039	90.7	0.6	(89.6, 91.8)
COLORADO	1815	93.1	0.7	(91.7, 94.6)
CONNECTICUT	2228	95.1	0.6	(94.0, 96.2)
DELAWARE	2548	94.3	0.6	(93.0, 95.5)
DISTRICT OF CO	1499	97.3	0.5	(96.3, 98.3)
FLORIDA	3483	94.3	0.5	(93.3, 95.2)
GEORGIA	2294	96.4	0.5	(95.5, 97.3)
HAWAII	2152	95.5	0.5	(94.4, 96.5)
IDAHO	4890	91.3	0.6	(90.1, 92.4)
ILLINOIS	1436	94.3	0.7	(93.0, 95.7)
INDIANA	2354	93.1	0.6	(91.9, 94.3)
IOWA	3585	92.7	0.5	(91.7, 93.6)
KANSAS	1981	94.7	0.6	(93.6, 95.8)
KENTUCKY	3567	93.8	0.5	(92.9, 94.7)
LOUISIANA	1643	94.4	0.7	(93.0, 95.7)
MAINE	1692	94.5	0.7	(93.2, 95.8)
MARYLAND	4547	96.4	0.4	(95.7, 97.1)
MASSACHUSETTS	1753	96.0	0.5	(95.0, 97.0)
MICHIGAN	2561	94.9	0.5	(93.9, 95.9)
MINNESOTA	4770	92.5	0.4	(91.7, 93.3)
MISSISSIPPI	1583	95.5	0.6	(94.4, 96.7)
MISSOURI	1835	94.7	0.7	(93.4, 96.1)
MONTANA	1785	92.3	0.7	(90.9, 93.6)
NEBRASKA	2676	93.2	0.6	(91.9, 94.4)
NEVADA	2497	93.3	0.9	(91.5, 95.0)
NEW HAMPSHIRE	1498	94.0	0.7	(92.6, 95.5)
NEW JERSEY	2650	94.4	0.6	(93.3, 95.5)
NEW MEXICO	1802	90.8	0.8	(89.2, 92.4)
NEW YORK	3371	94.8	0.5	(93.9, 95.7)
NORTH CAROLINA	3601	93.9	0.5	(93.0, 94.9)
NORTH DAKOTA	1793	93.0	0.7	(91.7, 94.3)
OHIO	3160	95.5	0.5	(94.5, 96.6)
OKLAHOMA	1881	94.2	0.7	(92.9, 95.5)
OREGON	3292	92.2	0.6	(91.0, 93.4)
PENNSYLVANIA	3564	95.6	0.4	(94.8, 96.3)
RHODE ISLAND	1831	95.7	0.5	(94.7, 96.8)
SOUTH CAROLINA	2143	96.5	0.5	(95.5, 97.4)
SOUTH DAKOTA	2187	92.7	0.6	(91.5, 93.9)
TENNESSEE	2967	95.1	0.5	(94.2, 96.0)
TEXAS	2480	92.4	0.6	(91.1, 93.6)
UTAH	2854	92.1	0.7	(90.8, 93.5)
VERMONT	3160	93.6	0.5	(92.6, 94.5)
VIRGINIA	3466	93.9	0.9	(92.2, 95.6)
WASHINGTON	3579	92.9	0.5	(91.9, 93.9)
WEST VIRGINIA	2408	93.6	0.6	(92.4, 94.8)
WISCONSIN	2232	92.6	0.7	(91.3, 93.9)
WYOMING	2397	90.7	0.6	(89.5, 92.0)
PUERTO RICO	2145	94.9	0.6	(93.8, 95.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 94.0 RANGE = 90.7-97.3

* BLOOD PRESSURE CHECKED WITHIN THE PAST 2 YEARS

August 14, 1998

TABLE 9.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BP CHECKED WITHIN LAST 2 YEARS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	802	94.4	0.9	(92.7, 96.0)	*	1348	96.4	0.6	(95.3, 97.5)	
ALASKA	709	88.5	1.9	(84.8, 92.3)	*	822	95.4	1.1	(93.2, 97.6)	
ARIZONA	885	92.0	1.2	(89.5, 94.4)	*	1007	95.0	1.0	(93.0, 97.1)	
ARKANSAS	666	92.0	1.1	(89.8, 94.3)	*	1109	94.5	0.9	(92.7, 96.3)	
CALIFORNIA	1743	86.5	1.0	(84.6, 88.4)	*	2296	94.7	0.6	(93.6, 95.9)	
COLORADO	815	89.0	1.3	(86.4, 91.5)	*	1000	97.1	0.6	(96.0, 98.3)	
CONNECTICUT	882	94.4	0.8	(92.8, 96.1)	*	1346	95.7	0.7	(94.3, 97.1)	
DELAWARE	1008	90.9	1.3	(88.4, 93.3)	*	1540	97.4	0.4	(96.6, 98.2)	
DISTRICT OF CO	570	95.5	1.0	(93.6, 97.4)	*	929	98.8	0.5	(97.8, 99.8)	
FLORIDA	1428	92.0	0.8	(90.3, 93.6)	*	2055	96.3	0.5	(95.4, 97.2)	
GEORGIA	952	94.1	0.9	(92.4, 95.8)	*	1342	98.5	0.4	(97.7, 99.2)	
HAWAII	940	93.3	0.9	(91.5, 95.0)	*	1212	97.7	0.6	(96.6, 98.8)	
IDAHO	2044	88.6	1.0	(86.6, 90.6)	*	2846	93.8	0.5	(92.8, 94.8)	
ILLINOIS	573	91.1	1.3	(88.5, 93.7)	*	863	97.2	0.6	(96.0, 98.4)	
INDIANA	1059	90.1	1.1	(88.0, 92.1)	*	1295	95.9	0.7	(94.5, 97.2)	
IOWA	1468	89.4	0.9	(87.6, 91.1)	*	2117	95.6	0.5	(94.7, 96.6)	
KANSAS	777	93.4	0.9	(91.6, 95.3)	*	1204	95.9	0.6	(94.7, 97.2)	
KENTUCKY	1297	91.4	0.9	(89.6, 93.1)	*	2270	96.0	0.4	(95.1, 96.9)	
LOUISIANA	627	92.2	1.2	(89.8, 94.6)	*	1016	96.3	0.7	(94.9, 97.7)	
MAINE	746	92.1	1.2	(89.7, 94.5)	*	946	96.7	0.6	(95.5, 97.9)	
MARYLAND	1942	94.6	0.6	(93.4, 95.9)	*	2605	98.0	0.3	(97.3, 98.6)	
MASSACHUSETTS	716	95.1	0.8	(93.5, 96.7)	*	1037	96.8	0.6	(95.5, 98.0)	
MICHIGAN	1068	93.4	0.9	(91.8, 95.1)	*	1493	96.2	0.5	(95.1, 97.3)	
MINNESOTA	2180	89.7	0.7	(88.3, 91.0)	*	2590	95.2	0.5	(94.3, 96.1)	
MISSISSIPPI	584	94.0	1.0	(92.0, 96.0)	*	999	96.9	0.7	(95.6, 98.3)	
MISSOURI	689	93.0	1.2	(90.6, 95.4)	*	1146	96.3	0.7	(94.9, 97.6)	
MONTANA	753	89.5	1.2	(87.0, 91.9)	*	1032	95.0	0.7	(93.5, 96.4)	
NEBRASKA	1036	90.5	1.1	(88.4, 92.6)	*	1640	95.6	0.7	(94.3, 96.9)	
NEVADA	1159	91.2	1.6	(88.2, 94.3)	*	1338	95.3	1.0	(93.4, 97.2)	
NEW HAMPSHIRE	597	91.8	1.3	(89.3, 94.3)	*	901	96.1	0.7	(94.7, 97.5)	
NEW JERSEY	1056	92.2	1.0	(90.2, 94.2)	*	1594	96.4	0.5	(95.4, 97.4)	
NEW MEXICO	752	87.7	1.4	(85.0, 90.4)	*	1050	93.7	0.8	(92.0, 95.3)	
NEW YORK	1336	92.6	0.8	(91.0, 94.1)	*	2035	96.8	0.5	(95.9, 97.7)	
NORTH CAROLINA	1424	90.2	0.9	(88.4, 92.0)	*	2177	97.3	0.4	(96.5, 98.1)	
NORTH DAKOTA	796	89.5	1.2	(87.3, 91.8)	*	997	96.4	0.7	(95.1, 97.7)	
OHIO	1226	93.3	1.0	(91.3, 95.2)	*	1934	97.5	0.5	(96.6, 98.5)	
OKLAHOMA	828	93.0	1.0	(91.0, 95.0)	*	1053	95.3	0.8	(93.6, 96.9)	
OREGON	1384	87.9	1.2	(85.6, 90.2)	*	1908	96.2	0.5	(95.2, 97.1)	
PENNSYLVANIA	1423	94.3	0.6	(93.1, 95.6)	*	2141	96.6	0.5	(95.7, 97.6)	
RHODE ISLAND	751	94.2	1.0	(92.3, 96.1)	*	1080	97.1	0.5	(96.1, 98.1)	
SOUTH CAROLINA	812	94.6	0.9	(92.9, 96.4)	*	1331	98.2	0.4	(97.3, 99.0)	
SOUTH DAKOTA	950	89.3	1.1	(87.1, 91.4)	*	1237	95.9	0.6	(94.6, 97.1)	
TENNESSEE	1037	92.7	0.9	(91.0, 94.4)	*	1930	97.3	0.4	(96.5, 98.1)	
TEXAS	1017	89.0	1.1	(86.8, 91.2)	*	1463	95.6	0.6	(94.3, 96.8)	
UTAH	1269	90.2	1.1	(88.0, 92.3)	*	1585	94.0	0.9	(92.2, 95.7)	
VERMONT	1380	91.3	0.9	(89.7, 93.0)	*	1780	95.6	0.5	(94.6, 96.6)	
VIRGINIA	1388	91.5	1.2	(89.2, 93.8)	*	2078	96.2	1.3	(93.6, 98.8)	
WASHINGTON	1533	89.8	0.9	(88.1, 91.5)	*	2046	95.8	0.5	(94.9, 96.8)	
WEST VIRGINIA	953	91.3	1.1	(89.2, 93.5)	*	1455	95.6	0.6	(94.4, 96.8)	
WISCONSIN	986	89.1	1.2	(86.8, 91.4)	*	1246	95.9	0.6	(94.7, 97.1)	
WYOMING	1010	87.1	1.1	(84.9, 89.3)	*	1387	94.3	0.7	(93.0, 95.6)	
PUERTO RICO	873	93.5	0.9	(91.7, 95.3)	*	1272	96.1	0.6	(94.8, 97.3)	

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 91.7	MEDIAN	= 96.1
	RANGE	= 86.5-95.5	RANGE	= 93.7-98.8

* BLOOD PRESSURE CHECKED WITHIN THE PAST 2 YEARS

August 14, 1998

TABLE 9.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 BP CHECKED WITHIN LAST 2 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	93.4	95.4	2.1
ALASKA	92.0	91.8	-0.1
ARIZONA	94.3	93.5	-0.8
ARKANSAS	94.0	93.3	-0.6
CALIFORNIA	91.3	90.7	-0.6
COLORADO	91.1	93.1	2.0
CONNECTICUT	95.1	95.1	0.0
DELAWARE	96.1	94.3	-1.8
DISTRICT OF CO	N/A	97.3	N/A
FLORIDA	94.0	94.3	0.3
GEORGIA	95.9	96.4	0.5
HAWAII	95.9	95.5	-0.4
IDAHO	91.9	91.3	-0.6
ILLINOIS	93.9	94.3	0.4
INDIANA	93.2	93.1	-0.1
IOWA	93.8	92.7	-1.1
KANSAS	93.0	94.7	1.7
KENTUCKY	94.0	93.8	-0.2
LOUISIANA	95.3	94.4	-1.0
MAINE	94.2	94.5	0.3
MARYLAND	95.2	96.4	1.2
MASSACHUSETTS	94.3	96.0	1.7
MICHIGAN	94.7	94.9	0.2
MINNESOTA	93.2	92.5	-0.6
MISSISSIPPI	92.8	95.5	2.7
MISSOURI	93.6	94.7	1.2
MONTANA	91.8	92.3	0.5
NEBRASKA	93.4	93.2	-0.3
NEVADA	93.4	93.3	-0.1
NEW HAMPSHIRE	94.5	94.0	-0.5
NEW JERSEY	95.2	94.4	-0.8
NEW MEXICO	91.1	90.8	-0.3
NEW YORK	95.1	94.8	-0.3
NORTH CAROLINA	94.0	93.9	-0.1
NORTH DAKOTA	93.7	93.0	-0.7
OHIO	95.7	95.5	-0.2
OKLAHOMA	94.7	94.2	-0.5
OREGON	91.5	92.2	0.7
PENNSYLVANIA	93.0	95.6	2.5
RHODE ISLAND	95.1	95.7	0.6
SOUTH CAROLINA	95.5	96.5	1.0
SOUTH DAKOTA	95.5	92.7	-2.9
TENNESSEE	96.1	95.1	-1.0
TEXAS	93.6	92.4	-1.3
UTAH	92.2	92.1	-0.1
VERMONT	93.8	93.6	-0.3
VIRGINIA	96.2	93.9	-2.3
WASHINGTON	93.2	92.9	-0.3
WEST VIRGINIA	94.4	93.6	-0.8
WISCONSIN	93.6	92.6	-1.0
WYOMING	91.6	90.7	-0.9
PUERTO RICO	N/A	94.9	N/A

* BLOOD PRESSURE CHECKED WITHIN THE PAST 2 YEARS
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 10.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2086	74.8	1.1	(72.6, 76.9)
ALASKA	1525	67.4	1.8	(63.8, 71.0)
ARIZONA	1872	70.9	1.6	(67.7, 74.0)
ARKANSAS	1721	63.5	1.4	(60.8, 66.1)
CALIFORNIA	3994	69.9	0.9	(68.2, 71.6)
COLORADO	1786	75.6	1.2	(73.2, 78.0)
CONNECTICUT	2202	77.9	1.2	(75.6, 80.2)
DELAWARE	2505	74.1	1.1	(72.0, 76.2)
DISTRICT OF CO	1485	81.3	1.3	(78.8, 83.8)
FLORIDA	3433	79.0	0.8	(77.4, 80.6)
GEORGIA	2272	75.8	1.1	(73.7, 78.0)
HAWAII	2116	74.7	1.1	(72.5, 76.9)
IDAHO	4789	72.7	0.8	(71.1, 74.3)
ILLINOIS	1412	71.9	1.4	(69.1, 74.7)
INDIANA	2298	71.9	1.1	(69.8, 74.1)
IOWA	3526	72.9	0.9	(71.1, 74.7)
KANSAS	1968	58.4	1.2	(56.0, 60.9)
KENTUCKY	3483	71.1	0.9	(69.3, 72.9)
LOUISIANA	1612	69.8	1.3	(67.2, 72.4)
MAINE	1666	77.5	1.2	(75.2, 79.8)
MARYLAND	4498	77.4	0.8	(75.7, 79.0)
MASSACHUSETTS	1724	79.1	1.2	(76.8, 81.3)
MICHIGAN	2534	75.6	1.0	(73.7, 77.5)
MINNESOTA	4712	65.9	0.8	(64.3, 67.4)
MISSISSIPPI	1551	66.5	1.5	(63.4, 69.5)
MISSOURI	1775	74.4	1.3	(71.9, 77.0)
MONTANA	1743	71.4	1.3	(68.9, 73.9)
NEBRASKA	2620	73.3	1.2	(71.1, 75.6)
NEVADA	2444	71.7	1.8	(68.1, 75.3)
NEW HAMPSHIRE	1475	77.7	1.3	(75.2, 80.2)
NEW JERSEY	2601	78.1	1.0	(76.1, 80.0)
NEW MEXICO	1759	68.0	1.3	(65.5, 70.6)
NEW YORK	3297	75.9	0.9	(74.2, 77.6)
NORTH CAROLINA	3503	75.7	0.8	(74.0, 77.3)
NORTH DAKOTA	1742	71.8	1.2	(69.4, 74.3)
OHIO	3102	70.0	1.1	(67.8, 72.2)
OKLAHOMA	1873	77.5	1.2	(75.1, 79.8)
OREGON	3229	73.9	1.0	(71.9, 75.8)
PENNSYLVANIA	3526	72.2	0.9	(70.4, 74.0)
RHODE ISLAND	1801	78.3	1.2	(76.1, 80.6)
SOUTH CAROLINA	2101	74.9	1.1	(72.7, 77.2)
SOUTH DAKOTA	2147	68.6	1.2	(66.2, 71.0)
TENNESSEE	2927	73.1	1.0	(71.2, 75.0)
TEXAS	2431	71.0	1.1	(68.9, 73.1)
UTAH	2789	71.9	1.1	(69.7, 74.0)
VERMONT	3131	74.5	0.9	(72.7, 76.3)
VIRGINIA	3401	78.3	1.1	(76.0, 80.5)
WASHINGTON	3498	75.8	0.9	(74.1, 77.6)
WEST VIRGINIA	2378	70.5	1.1	(68.4, 72.6)
WISCONSIN	2180	75.6	1.2	(73.3, 78.0)
WYOMING	2363	75.0	1.1	(72.9, 77.1)
PUERTO RICO	2238	80.9	1.0	(79.0, 82.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 74.0 RANGE = 58.4-81.3

* EVER HAD CHOLESTEROL CHECKED

August 14, 1998

TABLE 10.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	780	69.3	1.8	(65.7, 72.9)	*	1306	79.6	1.3	(77.1, 82.0)	
ALASKA	704	63.0	2.8	(57.6, 68.5)	*	821	72.2	2.3	(67.6, 76.7)	
ARIZONA	876	68.1	2.5	(63.2, 72.9)	*	996	73.6	1.9	(69.8, 77.4)	
ARKANSAS	645	59.0	2.3	(54.5, 63.6)	*	1076	67.4	1.8	(64.0, 70.9)	
CALIFORNIA	1721	64.8	1.4	(62.1, 67.4)	*	2273	74.9	1.1	(72.8, 77.0)	
COLORADO	804	72.3	1.9	(68.7, 75.9)	*	982	78.8	1.6	(75.8, 81.8)	
CONNECTICUT	874	76.8	2.0	(72.9, 80.7)	*	1328	78.9	1.3	(76.4, 81.4)	
DELAWARE	989	72.2	1.7	(68.9, 75.5)	*	1516	75.8	1.4	(73.1, 78.5)	
DISTRICT OF CO	559	77.1	2.2	(72.9, 81.3)	*	926	84.9	1.3	(82.3, 87.5)	
FLORIDA	1412	76.0	1.4	(73.3, 78.6)	*	2021	81.8	1.0	(79.8, 83.7)	
GEORGIA	944	70.6	1.8	(67.1, 74.1)	*	1328	80.6	1.3	(78.1, 83.2)	
HAWAII	926	73.2	1.7	(69.9, 76.5)	*	1190	76.2	1.5	(73.2, 79.2)	
IDAHO	1998	71.6	1.2	(69.2, 74.0)	*	2791	73.8	1.2	(71.5, 76.1)	
ILLINOIS	565	68.0	2.3	(63.5, 72.5)	*	847	75.3	1.7	(72.0, 78.6)	
INDIANA	1038	68.8	1.7	(65.5, 72.0)	*	1260	74.8	1.4	(72.0, 77.6)	
IOWA	1449	68.8	1.4	(66.1, 71.6)	*	2077	76.7	1.1	(74.5, 78.8)	
KANSAS	777	56.4	2.0	(52.6, 60.3)	*	1191	60.3	1.5	(57.3, 63.4)	
KENTUCKY	1270	66.8	1.5	(63.8, 69.8)	*	2213	75.0	1.1	(72.8, 77.1)	
LOUISIANA	607	66.5	2.2	(62.1, 70.8)	*	1005	72.7	1.6	(69.6, 75.9)	
MAINE	733	73.5	1.9	(69.9, 77.2)	*	933	81.1	1.5	(78.3, 83.9)	
MARYLAND	1918	73.7	1.3	(71.2, 76.3)	*	2580	80.6	1.0	(78.6, 82.6)	
MASSACHUSETTS	697	76.4	1.9	(72.6, 80.1)	*	1027	81.4	1.4	(78.8, 84.1)	
MICHIGAN	1061	71.8	1.5	(68.8, 74.8)	*	1473	79.1	1.2	(76.8, 81.5)	
MINNESOTA	2161	59.7	1.1	(57.5, 61.9)	*	2551	71.6	1.0	(69.6, 73.6)	
MISSISSIPPI	577	62.0	2.5	(57.2, 66.9)	*	974	70.4	1.8	(66.9, 73.9)	
MISSOURI	673	71.9	2.1	(67.8, 75.9)	*	1102	76.7	1.6	(73.7, 79.8)	
MONTANA	737	71.4	1.9	(67.7, 75.0)	*	1006	71.4	1.7	(68.1, 74.7)	
NEBRASKA	1014	69.5	1.9	(65.8, 73.1)	*	1606	76.9	1.3	(74.2, 79.5)	
NEVADA	1133	67.2	3.0	(61.4, 73.0)	*	1311	76.2	2.1	(72.0, 80.3)	
NEW HAMPSHIRE	591	76.4	2.1	(72.2, 80.6)	*	884	79.0	1.5	(75.9, 82.0)	
NEW JERSEY	1031	76.3	1.6	(73.1, 79.4)	*	1570	79.7	1.2	(77.3, 82.1)	
NEW MEXICO	740	65.1	1.9	(61.3, 68.9)	*	1019	70.9	1.7	(67.5, 74.2)	
NEW YORK	1319	73.9	1.4	(71.1, 76.6)	*	1978	77.7	1.1	(75.6, 79.8)	
NORTH CAROLINA	1376	71.1	1.4	(68.3, 73.9)	*	2127	79.7	1.0	(77.8, 81.7)	
NORTH DAKOTA	777	68.7	1.9	(65.1, 72.3)	*	965	74.9	1.7	(71.6, 78.1)	
OHIO	1210	68.0	1.8	(64.4, 71.6)	*	1892	71.8	1.4	(69.1, 74.4)	
OKLAHOMA	826	76.1	1.9	(72.5, 79.8)	*	1047	78.7	1.5	(75.7, 81.6)	
OREGON	1369	70.9	1.6	(67.9, 74.0)	*	1860	76.7	1.2	(74.3, 79.0)	
PENNSYLVANIA	1409	69.4	1.4	(66.6, 72.2)	*	2117	74.7	1.2	(72.3, 77.0)	
RHODE ISLAND	743	75.5	1.8	(71.9, 79.1)	*	1058	80.8	1.5	(77.9, 83.7)	
SOUTH CAROLINA	790	70.6	1.9	(66.9, 74.3)	*	1311	78.7	1.3	(76.1, 81.3)	
SOUTH DAKOTA	939	63.5	1.8	(59.9, 67.1)	*	1208	73.5	1.6	(70.4, 76.5)	
TENNESSEE	1027	69.5	1.6	(66.5, 72.5)	*	1900	76.3	1.2	(74.0, 78.6)	
TEXAS	997	68.1	1.7	(64.8, 71.4)	*	1434	73.7	1.3	(71.1, 76.3)	
UTAH	1243	70.6	1.6	(67.4, 73.8)	*	1546	73.1	1.5	(70.1, 76.1)	
VERMONT	1370	72.1	1.4	(69.3, 75.0)	*	1761	76.7	1.1	(74.5, 78.9)	
VIRGINIA	1365	76.9	1.6	(73.9, 80.0)	*	2036	79.5	1.7	(76.3, 82.8)	
WASHINGTON	1508	72.8	1.4	(70.1, 75.5)	*	1990	78.8	1.2	(76.4, 81.2)	
WEST VIRGINIA	949	67.0	1.7	(63.7, 70.3)	*	1429	73.6	1.3	(71.0, 76.2)	
WISCONSIN	957	72.9	1.8	(69.3, 76.5)	*	1223	78.2	1.5	(75.3, 81.1)	
WYOMING	1001	72.0	1.7	(68.7, 75.4)	*	1362	77.9	1.4	(75.3, 80.6)	
PUERTO RICO	922	74.5	1.6	(71.3, 77.7)	*	1316	86.6	1.1	(84.4, 88.7)	

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 70.8
RANGE = 56.4-77.1

NUMBER OF PARTICIPANTS = 52
MEDIAN = 76.7
RANGE = 60.3-86.6

* EVER HAD CHOLESTEROL CHECKED

August 14, 1998

TABLE 10.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 CHOLESTEROL SCREENING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	66.9	74.8	7.9
ALASKA	67.4	67.4	-0.0
ARIZONA	72.3	70.9	-1.4
ARKANSAS	65.8	63.5	-2.4
CALIFORNIA	68.1	69.9	1.8
COLORADO	73.4	75.6	2.2
CONNECTICUT	76.1	77.9	1.8
DELAWARE	70.7	74.1	3.4
DISTRICT OF CO	N/A	81.3	N/A
FLORIDA	75.8	79.0	3.3
GEORGIA	72.4	75.8	3.4
HAWAII	71.0	74.7	3.7
IDAHO	71.3	72.7	1.5
ILLINOIS	68.6	71.9	3.3
INDIANA	69.3	71.9	2.7
IOWA	72.2	72.9	0.8
KANSAS	71.6	58.4	-13.1
KENTUCKY	67.6	71.1	3.5
LOUISIANA	67.9	69.8	1.9
MAINE	68.6	77.5	8.9
MARYLAND	75.4	77.4	1.9
MASSACHUSETTS	78.7	79.1	0.4
MICHIGAN	74.0	75.6	1.6
MINNESOTA	65.3	65.9	0.6
MISSISSIPPI	62.0	66.5	4.4
MISSOURI	70.4	74.4	4.0
MONTANA	71.1	71.4	0.3
NEBRASKA	68.5	73.3	4.8
NEVADA	69.4	71.7	2.3
NEW HAMPSHIRE	77.3	77.7	0.4
NEW JERSEY	75.2	78.1	2.9
NEW MEXICO	67.0	68.0	1.1
NEW YORK	74.6	75.9	1.3
NORTH CAROLINA	71.7	75.7	4.0
NORTH DAKOTA	70.8	71.8	1.0
OHIO	64.5	70.0	5.5
OKLAHOMA	69.5	77.5	8.0
OREGON	72.8	73.9	1.0
PENNSYLVANIA	73.0	72.2	-0.7
RHODE ISLAND	77.3	78.3	1.0
SOUTH CAROLINA	72.0	74.9	2.9
SOUTH DAKOTA	68.5	68.6	0.1
TENNESSEE	70.1	73.1	3.0
TEXAS	73.9	71.0	-2.9
UTAH	66.7	71.9	5.2
VERMONT	73.8	74.5	0.7
VIRGINIA	75.7	78.3	2.6
WASHINGTON	75.6	75.8	0.3
WEST VIRGINIA	69.3	70.5	1.2
WISCONSIN	73.6	75.6	2.0
WYOMING	68.1	75.0	6.9
PUERTO RICO	N/A	80.9	N/A

* EVER HAD CHOLESTEROL CHECKED
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 11.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2041	71.0	1.2	(68.7, 73.3)
ALASKA	1509	62.4	1.9	(58.7, 66.1)
ARIZONA	1862	68.5	1.7	(65.2, 71.7)
ARKANSAS	1680	58.6	1.4	(55.9, 61.4)
CALIFORNIA	3978	65.8	0.9	(64.0, 67.5)
COLORADO	1775	69.7	1.3	(67.2, 72.2)
CONNECTICUT	2172	73.3	1.2	(70.9, 75.7)
DELAWARE	2477	69.3	1.1	(67.1, 71.5)
DISTRICT OF CO	1478	79.4	1.3	(76.8, 81.9)
FLORIDA	3397	75.3	0.9	(73.6, 77.0)
GEORGIA	2236	72.6	1.2	(70.3, 74.8)
HAWAII	2083	69.5	1.2	(67.1, 71.8)
IDAHO	4753	65.0	0.9	(63.1, 66.8)
ILLINOIS	1404	67.2	1.5	(64.3, 70.1)
INDIANA	2262	66.4	1.2	(64.1, 68.7)
IOWA	3493	66.1	1.0	(64.2, 68.0)
KANSAS	1941	54.9	1.3	(52.4, 57.4)
KENTUCKY	3439	65.6	1.0	(63.6, 67.5)
LOUISIANA	1591	66.3	1.4	(63.6, 69.0)
MAINE	1646	71.8	1.3	(69.3, 74.4)
MARYLAND	4458	74.5	0.9	(72.8, 76.2)
MASSACHUSETTS	1702	74.7	1.2	(72.3, 77.1)
MICHIGAN	2522	71.1	1.0	(69.1, 73.1)
MINNESOTA	4651	61.2	0.8	(59.7, 62.8)
MISSISSIPPI	1530	62.5	1.5	(59.5, 65.5)
MISSOURI	1749	70.4	1.3	(67.8, 73.0)
MONTANA	1708	63.2	1.3	(60.6, 65.8)
NEBRASKA	2590	65.5	1.2	(63.1, 67.9)
NEVADA	2431	68.4	1.9	(64.8, 72.1)
NEW HAMPSHIRE	1461	73.0	1.4	(70.4, 75.7)
NEW JERSEY	2579	75.3	1.0	(73.3, 77.3)
NEW MEXICO	1747	62.8	1.3	(60.1, 65.4)
NEW YORK	3256	72.5	0.9	(70.7, 74.3)
NORTH CAROLINA	3451	72.1	0.9	(70.4, 73.8)
NORTH DAKOTA	1729	63.8	1.3	(61.2, 66.4)
OHIO	3087	66.5	1.2	(64.2, 68.7)
OKLAHOMA	1862	74.4	1.2	(72.0, 76.9)
OREGON	3189	68.2	1.0	(66.2, 70.2)
PENNSYLVANIA	3470	68.5	1.0	(66.6, 70.4)
RHODE ISLAND	1778	74.7	1.2	(72.3, 77.1)
SOUTH CAROLINA	2088	72.2	1.2	(69.8, 74.6)
SOUTH DAKOTA	2125	63.2	1.3	(60.7, 65.6)
TENNESSEE	2904	70.3	1.0	(68.3, 72.2)
TEXAS	2405	67.4	1.1	(65.3, 69.6)
UTAH	2778	65.6	1.2	(63.3, 67.9)
VERMONT	3102	68.9	1.0	(67.0, 70.8)
VIRGINIA	3323	73.6	1.2	(71.2, 75.9)
WASHINGTON	3471	69.6	1.0	(67.7, 71.5)
WEST VIRGINIA	2353	67.2	1.1	(65.0, 69.4)
WISCONSIN	2170	70.2	1.2	(67.8, 72.6)
WYOMING	2346	70.2	1.1	(68.0, 72.4)
PUERTO RICO	2196	79.3	1.0	(77.3, 81.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 69.1 RANGE = 54.9-79.4 YEAR 2000 OBJ. 15.14
 * CHOLESTEROL CHECKED WITHIN THE PAST 5 YEARS August 14, 1998

TABLE 11.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL SCREENING* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES-----*					FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	768	65.2	1.9	(61.4, 68.9)	*	1273	76.2	1.3	(73.6, 78.8)	
ALASKA	698	58.9	2.8	(53.4, 64.3)	*	811	66.2	2.5	(61.4, 71.1)	
ARIZONA	868	65.8	2.6	(60.8, 70.8)	*	994	71.0	2.0	(67.1, 74.9)	
ARKANSAS	633	55.2	2.3	(50.6, 59.7)	*	1047	61.8	1.8	(58.2, 65.3)	
CALIFORNIA	1715	60.3	1.4	(57.6, 63.0)	*	2263	71.1	1.1	(68.9, 73.2)	
COLORADO	800	66.1	1.9	(62.3, 69.9)	*	975	73.2	1.7	(69.9, 76.4)	
CONNECTICUT	861	72.4	2.1	(68.3, 76.4)	*	1311	74.2	1.4	(71.5, 76.9)	
DELAWARE	982	68.0	1.8	(64.5, 71.4)	*	1495	70.5	1.5	(67.6, 73.4)	
DISTRICT OF CO	557	75.5	2.2	(71.2, 79.8)	*	921	82.6	1.4	(79.9, 85.4)	
FLORIDA	1395	72.0	1.4	(69.2, 74.7)	*	2002	78.3	1.1	(76.2, 80.4)	
GEORGIA	930	67.5	1.8	(63.9, 71.1)	*	1306	77.2	1.4	(74.4, 80.0)	
HAWAII	912	68.4	1.8	(64.9, 71.9)	*	1171	70.6	1.6	(67.4, 73.8)	
IDAHO	1986	64.1	1.4	(61.4, 66.7)	*	2767	65.9	1.3	(63.4, 68.4)	
ILLINOIS	560	63.4	2.4	(58.6, 68.1)	*	844	70.6	1.8	(67.1, 74.0)	
INDIANA	1028	63.5	1.7	(60.2, 66.9)	*	1234	69.0	1.6	(66.0, 72.0)	
IOWA	1445	61.9	1.5	(59.0, 64.8)	*	2048	70.0	1.2	(67.7, 72.3)	
KANSAS	766	53.0	2.0	(49.1, 56.9)	*	1175	56.7	1.6	(53.6, 59.7)	
KENTUCKY	1259	61.1	1.6	(58.0, 64.3)	*	2180	69.6	1.2	(67.3, 71.8)	
LOUISIANA	603	62.6	2.3	(58.2, 67.1)	*	988	69.6	1.7	(66.3, 72.8)	
MAINE	728	67.9	2.0	(64.0, 71.8)	*	918	75.5	1.6	(72.4, 78.6)	
MARYLAND	1899	70.7	1.4	(68.0, 73.3)	*	2559	78.0	1.1	(75.9, 80.1)	
MASSACHUSETTS	690	72.4	2.0	(68.5, 76.3)	*	1012	76.8	1.5	(73.9, 79.7)	
MICHIGAN	1059	67.0	1.6	(63.8, 70.1)	*	1463	74.9	1.3	(72.4, 77.4)	
MINNESOTA	2135	54.5	1.2	(52.3, 56.8)	*	2516	67.5	1.0	(65.5, 69.6)	
MISSISSIPPI	571	58.1	2.5	(53.2, 63.0)	*	959	66.4	1.8	(62.9, 70.0)	
MISSOURI	668	68.2	2.1	(64.1, 72.3)	*	1081	72.4	1.6	(69.2, 75.6)	
MONTANA	721	63.0	2.0	(59.1, 66.9)	*	987	63.4	1.7	(59.9, 66.8)	
NEBRASKA	1007	61.5	2.0	(57.7, 65.3)	*	1583	69.2	1.5	(66.4, 72.1)	
NEVADA	1127	64.4	3.0	(58.5, 70.2)	*	1304	72.5	2.3	(68.1, 76.9)	
NEW HAMPSHIRE	586	71.1	2.3	(66.6, 75.5)	*	875	74.9	1.6	(71.7, 78.1)	
NEW JERSEY	1021	73.6	1.7	(70.3, 76.8)	*	1558	76.9	1.3	(74.4, 79.4)	
NEW MEXICO	734	59.2	2.0	(55.2, 63.1)	*	1013	66.2	1.7	(62.8, 69.6)	
NEW YORK	1299	70.0	1.5	(67.1, 72.9)	*	1957	74.7	1.1	(72.5, 76.9)	
NORTH CAROLINA	1362	67.0	1.5	(64.1, 69.8)	*	2089	76.7	1.1	(74.6, 78.8)	
NORTH DAKOTA	770	60.3	1.9	(56.5, 64.1)	*	959	67.1	1.8	(63.7, 70.6)	
OHIO	1203	64.1	1.9	(60.4, 67.8)	*	1884	68.6	1.4	(65.9, 71.4)	
OKLAHOMA	823	73.1	2.0	(69.2, 76.9)	*	1039	75.7	1.6	(72.7, 78.8)	
OREGON	1354	65.0	1.6	(61.9, 68.2)	*	1835	71.2	1.3	(68.7, 73.7)	
PENNSYLVANIA	1396	66.3	1.5	(63.4, 69.1)	*	2074	70.5	1.2	(68.1, 73.0)	
RHODE ISLAND	735	71.7	1.9	(67.9, 75.4)	*	1043	77.5	1.5	(74.5, 80.5)	
SOUTH CAROLINA	788	67.8	2.0	(64.0, 71.6)	*	1300	76.1	1.4	(73.4, 78.8)	
SOUTH DAKOTA	933	59.0	1.9	(55.3, 62.7)	*	1192	67.1	1.7	(63.9, 70.3)	
TENNESSEE	1021	65.8	1.6	(62.7, 68.9)	*	1883	74.3	1.2	(71.9, 76.6)	
TEXAS	988	64.1	1.7	(60.8, 67.5)	*	1417	70.5	1.4	(67.9, 73.2)	
UTAH	1240	65.3	1.7	(61.9, 68.7)	*	1538	65.9	1.6	(62.7, 69.0)	
VERMONT	1355	66.6	1.5	(63.6, 69.6)	*	1747	71.1	1.2	(68.7, 73.5)	
VIRGINIA	1339	72.7	1.7	(69.4, 75.9)	*	1984	74.4	1.7	(71.0, 77.8)	
WASHINGTON	1500	66.3	1.5	(63.4, 69.1)	*	1971	72.9	1.3	(70.4, 75.4)	
WEST VIRGINIA	940	63.8	1.8	(60.4, 67.2)	*	1413	70.2	1.4	(67.5, 73.0)	
WISCONSIN	954	68.7	1.9	(65.0, 72.4)	*	1216	71.7	1.6	(68.6, 74.8)	
WYOMING	996	66.3	1.8	(62.8, 69.7)	*	1350	74.1	1.4	(71.3, 76.9)	
PUERTO RICO	902	72.2	1.7	(68.9, 75.5)	*	1294	85.5	1.1	(83.3, 87.7)	

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 65.9
RANGE = 53.0-75.5

NUMBER OF PARTICIPANTS = 52
MEDIAN = 71.2
RANGE = 56.7-85.5

* CHOLESTEROL CHECKED WITHIN THE PAST 5 YEARS

August 14, 1998

TABLE 11.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 CHOLESTEROL SCREENING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	64.1	71.0	6.9
ALASKA	63.2	62.4	-0.8
ARIZONA	69.2	68.5	-0.7
ARKANSAS	63.3	58.6	-4.6
CALIFORNIA	64.9	65.8	0.8
COLORADO	68.1	69.7	1.6
CONNECTICUT	72.6	73.3	0.7
DELAWARE	68.3	69.3	1.0
DISTRICT OF CO	N/A	79.4	N/A
FLORIDA	73.6	75.3	1.7
GEORGIA	70.0	72.6	2.5
HAWAII	68.9	69.5	0.6
IDAHO	66.3	65.0	-1.3
ILLINOIS	66.4	67.2	0.8
INDIANA	64.9	66.4	1.5
IOWA	67.1	66.1	-1.0
KANSAS	68.2	54.9	-13.3
KENTUCKY	64.0	65.6	1.5
LOUISIANA	65.6	66.3	0.7
MAINE	65.1	71.8	6.7
MARYLAND	72.8	74.5	1.7
MASSACHUSETTS	75.7	74.7	-1.0
MICHIGAN	70.8	71.1	0.3
MINNESOTA	62.1	61.2	-0.9
MISSISSIPPI	58.3	62.5	4.2
MISSOURI	65.6	70.4	4.8
MONTANA	64.2	63.2	-1.0
NEBRASKA	62.1	65.5	3.5
NEVADA	66.0	68.4	2.5
NEW HAMPSHIRE	73.5	73.0	-0.4
NEW JERSEY	73.2	75.3	2.2
NEW MEXICO	64.2	62.8	-1.5
NEW YORK	72.7	72.5	-0.2
NORTH CAROLINA	69.0	72.1	3.1
NORTH DAKOTA	65.5	63.8	-1.8
OHIO	62.1	66.5	4.4
OKLAHOMA	67.1	74.4	7.3
OREGON	67.9	68.2	0.3
PENNSYLVANIA	69.1	68.5	-0.5
RHODE ISLAND	74.8	74.7	-0.0
SOUTH CAROLINA	70.3	72.2	1.9
SOUTH DAKOTA	65.5	63.2	-2.3
TENNESSEE	68.2	70.3	2.0
TEXAS	70.0	67.4	-2.6
UTAH	63.2	65.6	2.4
VERMONT	69.2	68.9	-0.3
VIRGINIA	72.8	73.6	0.8
WASHINGTON	70.4	69.6	-0.8
WEST VIRGINIA	66.9	67.2	0.3
WISCONSIN	68.9	70.2	1.4
WYOMING	64.6	70.2	5.6
PUERTO RICO	N/A	79.3	N/A

* CHOLESTEROL CHECKED WITHIN THE PAST 5 YEARS
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 12.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER HAD CHECKED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1602	28.8	1.2	(26.3, 31.2)
ALASKA	1051	25.5	1.9	(21.8, 29.2)
ARIZONA	1341	31.9	1.8	(28.4, 35.4)
ARKANSAS	1146	30.1	1.5	(27.1, 33.0)
CALIFORNIA	2955	30.4	1.0	(28.5, 32.2)
COLORADO	1383	27.9	1.4	(25.2, 30.6)
CONNECTICUT	1721	24.3	1.2	(22.0, 26.7)
DELAWARE	1909	28.9	1.2	(26.6, 31.2)
DISTRICT OF CO	1217	18.2	1.3	(15.7, 20.6)
FLORIDA	2766	31.9	1.0	(29.9, 33.9)
GEORGIA	1752	24.1	1.1	(21.9, 26.3)
HAWAII	1607	31.5	1.4	(28.8, 34.2)
IDAHO	3473	29.8	1.1	(27.7, 31.9)
ILLINOIS	1038	34.3	1.7	(31.0, 37.5)
INDIANA	1670	29.0	1.3	(26.5, 31.5)
IOWA	2655	28.1	1.0	(26.2, 30.0)
KANSAS	1187	28.1	1.4	(25.2, 30.9)
KENTUCKY	2560	30.1	1.0	(28.1, 32.1)
LOUISIANA	1150	26.6	1.5	(23.7, 29.5)
MAINE	1313	32.2	1.4	(29.3, 35.0)
MARYLAND	3533	28.6	1.0	(26.7, 30.5)
MASSACHUSETTS	1361	25.0	1.3	(22.4, 27.6)
MICHIGAN	1954	31.2	1.2	(28.9, 33.4)
MINNESOTA	3161	31.2	0.9	(29.5, 33.0)
MISSISSIPPI	1090	28.8	1.5	(25.9, 31.7)
MISSOURI	1352	30.5	1.5	(27.6, 33.4)
MONTANA	1275	31.0	1.4	(28.2, 33.7)
NEBRASKA	1914	30.0	1.4	(27.2, 32.8)
NEVADA	1793	29.6	2.2	(25.3, 33.9)
NEW HAMPSHIRE	1177	30.9	1.5	(28.0, 33.9)
NEW JERSEY	2055	27.9	1.2	(25.6, 30.2)
NEW MEXICO	1261	28.0	1.4	(25.2, 30.7)
NEW YORK	2515	28.0	1.0	(26.0, 30.1)
NORTH CAROLINA	2704	26.5	0.9	(24.7, 28.2)
NORTH DAKOTA	1289	29.8	1.4	(27.1, 32.5)
OHIO	2299	27.7	1.3	(25.2, 30.2)
OKLAHOMA	1482	22.0	1.3	(19.5, 24.4)
OREGON	2441	32.0	1.1	(29.8, 34.1)
PENNSYLVANIA	2598	25.9	1.0	(24.0, 27.8)
RHODE ISLAND	1431	28.1	1.3	(25.5, 30.7)
SOUTH CAROLINA	1616	24.5	1.2	(22.2, 26.8)
SOUTH DAKOTA	1520	25.7	1.2	(23.3, 28.1)
TENNESSEE	2187	29.5	1.1	(27.3, 31.6)
TEXAS	1779	28.7	1.2	(26.4, 31.0)
UTAH	1991	26.4	1.3	(23.7, 29.0)
VERMONT	2361	25.9	1.1	(23.7, 28.0)
VIRGINIA	2674	29.5	1.3	(26.9, 32.0)
WASHINGTON	2675	25.6	0.9	(23.8, 27.5)
WEST VIRGINIA	1723	32.2	1.2	(29.8, 34.6)
WISCONSIN	1673	26.9	1.3	(24.4, 29.4)
WYOMING	1790	29.9	1.2	(27.6, 32.2)
PUERTO RICO	1851	24.4	1.1	(22.3, 26.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 28.7 RANGE = 18.2-34.3

* EVER TOLD CHOLESTEROL IS HIGH BY HEALTH PROFESSIONAL

August 14, 1998

TABLE 12.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHOLESTEROL AWARENESS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER HAD CHECKED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	552	28.7	2.1	(24.6, 32.8)	*	1050	28.9	1.5	(25.8, 31.9)	*
ALASKA	465	24.8	2.7	(19.5, 30.1)	*	586	26.2	2.7	(21.0, 31.4)	*
ARIZONA	612	30.1	2.5	(25.2, 35.0)	*	729	33.5	2.3	(28.9, 38.0)	*
ARKANSAS	402	32.7	2.5	(27.7, 37.6)	*	744	28.1	1.8	(24.5, 31.6)	*
CALIFORNIA	1201	30.2	1.5	(27.3, 33.1)	*	1754	30.5	1.2	(28.1, 33.0)	*
COLORADO	598	30.0	2.1	(25.9, 34.1)	*	785	26.1	1.8	(22.6, 29.6)	*
CONNECTICUT	685	25.1	1.9	(21.2, 28.9)	*	1036	23.7	1.5	(20.8, 26.5)	*
DELAWARE	733	27.9	1.8	(24.3, 31.5)	*	1176	29.7	1.5	(26.7, 32.8)	*
DISTRICT OF CO	434	16.3	2.0	(12.4, 20.1)	*	783	19.6	1.6	(16.5, 22.7)	*
FLORIDA	1102	30.3	1.5	(27.3, 33.3)	*	1664	33.3	1.3	(30.7, 35.8)	*
GEORGIA	684	20.6	1.7	(17.3, 23.9)	*	1068	26.9	1.5	(24.0, 29.9)	*
HAWAII	691	31.2	2.0	(27.2, 35.2)	*	916	31.8	1.9	(28.1, 35.5)	*
IDAHO	1396	30.3	1.5	(27.3, 33.3)	*	2077	29.3	1.5	(26.4, 32.2)	*
ILLINOIS	390	33.1	2.7	(27.9, 38.3)	*	648	35.2	2.1	(31.1, 39.3)	*
INDIANA	724	26.8	1.9	(23.0, 30.5)	*	946	30.9	1.7	(27.5, 34.3)	*
IOWA	1009	27.4	1.5	(24.5, 30.4)	*	1646	28.6	1.3	(26.2, 31.1)	*
KANSAS	450	27.2	2.3	(22.7, 31.6)	*	737	28.8	1.8	(25.4, 32.3)	*
KENTUCKY	885	28.5	1.6	(25.2, 31.7)	*	1675	31.5	1.2	(29.0, 33.9)	*
LOUISIANA	418	24.1	2.3	(19.7, 28.5)	*	732	28.5	1.8	(25.0, 32.1)	*
MAINE	551	32.7	2.2	(28.4, 37.0)	*	762	31.7	1.9	(28.0, 35.4)	*
MARYLAND	1446	28.6	1.5	(25.6, 31.5)	*	2087	28.7	1.3	(26.2, 31.2)	*
MASSACHUSETTS	538	24.7	2.1	(20.5, 28.8)	*	823	25.2	1.7	(21.9, 28.6)	*
MICHIGAN	777	31.5	1.8	(28.0, 35.1)	*	1177	30.9	1.5	(28.0, 33.7)	*
MINNESOTA	1289	30.4	1.4	(27.7, 33.0)	*	1872	31.9	1.2	(29.6, 34.2)	*
MISSISSIPPI	384	27.6	2.4	(22.9, 32.2)	*	706	29.7	1.9	(26.0, 33.4)	*
MISSOURI	493	30.1	2.3	(25.7, 34.6)	*	859	30.8	1.9	(27.2, 34.5)	*
MONTANA	527	29.9	2.1	(25.8, 33.9)	*	748	32.0	1.9	(28.3, 35.8)	*
NEBRASKA	694	32.5	2.3	(28.0, 36.9)	*	1220	28.0	1.8	(24.6, 31.4)	*
NEVADA	795	29.8	3.6	(22.8, 36.8)	*	998	29.4	2.7	(24.2, 34.6)	*
NEW HAMPSHIRE	463	30.8	2.3	(26.2, 35.3)	*	714	31.1	2.0	(27.2, 34.9)	*
NEW JERSEY	801	26.4	1.9	(22.8, 30.1)	*	1254	29.2	1.5	(26.2, 32.1)	*
NEW MEXICO	514	28.7	2.2	(24.4, 33.0)	*	747	27.3	1.7	(24.0, 30.7)	*
NEW YORK	990	26.2	1.6	(23.1, 29.3)	*	1525	29.6	1.3	(27.1, 32.1)	*
NORTH CAROLINA	1004	25.2	1.4	(22.3, 28.0)	*	1700	27.5	1.2	(25.2, 29.8)	*
NORTH DAKOTA	542	27.8	2.1	(23.7, 32.0)	*	747	31.6	1.8	(28.1, 35.1)	*
OHIO	884	28.4	2.0	(24.6, 32.3)	*	1415	27.0	1.6	(24.0, 30.1)	*
OKLAHOMA	649	20.0	1.9	(16.1, 23.8)	*	833	23.8	1.6	(20.6, 27.0)	*
OREGON	986	31.2	1.7	(27.8, 34.5)	*	1455	32.6	1.4	(29.8, 35.5)	*
PENNSYLVANIA	993	25.5	1.5	(22.4, 28.5)	*	1605	26.2	1.2	(23.8, 28.7)	*
RHODE ISLAND	572	27.1	2.0	(23.2, 31.1)	*	859	28.9	1.7	(25.5, 32.3)	*
SOUTH CAROLINA	569	22.6	1.9	(18.9, 26.4)	*	1047	25.9	1.5	(23.1, 28.8)	*
SOUTH DAKOTA	612	24.9	1.9	(21.2, 28.6)	*	908	26.3	1.6	(23.3, 29.4)	*
TENNESSEE	711	26.1	1.8	(22.7, 29.6)	*	1476	32.2	1.3	(29.6, 34.8)	*
TEXAS	705	26.6	1.8	(23.1, 30.1)	*	1074	30.4	1.5	(27.4, 33.4)	*
UTAH	869	27.5	2.0	(23.6, 31.5)	*	1122	25.3	1.8	(21.9, 28.7)	*
VERMONT	1009	26.1	1.7	(22.7, 29.5)	*	1352	25.7	1.3	(23.1, 28.3)	*
VIRGINIA	1038	32.6	2.2	(28.3, 36.9)	*	1636	26.6	1.5	(23.7, 29.5)	*
WASHINGTON	1096	24.4	1.4	(21.6, 27.1)	*	1579	26.8	1.2	(24.3, 29.2)	*
WEST VIRGINIA	648	31.0	1.9	(27.2, 34.8)	*	1075	33.2	1.6	(30.2, 36.3)	*
WISCONSIN	702	26.6	2.0	(22.8, 30.5)	*	971	27.2	1.7	(23.9, 30.5)	*
WYOMING	726	32.2	1.9	(28.5, 35.9)	*	1064	27.8	1.5	(24.9, 30.8)	*
PUERTO RICO	704	23.2	1.7	(19.8, 26.6)	*	1147	25.3	1.4	(22.5, 28.0)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 27.9	MEDIAN	= 28.8
	RANGE	= 16.3-33.1	RANGE	= 19.6-35.2

* EVER TOLD CHOLESTEROL IS HIGH BY HEALTH PROFESSIONAL

August 14, 1998

TABLE 12.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 CHOLESTEROL AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER HAD CHECKED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	27.3	28.8	1.5
ALASKA	24.9	25.5	0.6
ARIZONA	26.8	31.9	5.1
ARKANSAS	28.8	30.1	1.3
CALIFORNIA	28.0	30.4	2.3
COLORADO	27.9	27.9	-0.0
CONNECTICUT	25.3	24.3	-1.0
DELAWARE	30.0	28.9	-1.1
DISTRICT OF CO	N/A	18.2	N/A
FLORIDA	31.0	31.9	0.9
GEORGIA	22.0	24.1	2.2
HAWAII	27.1	31.5	4.4
IDAHO	28.1	29.8	1.7
ILLINOIS	27.3	34.3	7.0
INDIANA	30.2	29.0	-1.2
IOWA	29.5	28.1	-1.4
KANSAS	31.9	28.1	-3.8
KENTUCKY	29.8	30.1	0.4
LOUISIANA	26.3	26.6	0.3
MAINE	29.6	32.2	2.5
MARYLAND	25.0	28.6	3.7
MASSACHUSETTS	31.3	25.0	-6.3
MICHIGAN	31.5	31.2	-0.3
MINNESOTA	27.1	31.2	4.1
MISSISSIPPI	25.7	28.8	3.1
MISSOURI	28.8	30.5	1.7
MONTANA	28.3	31.0	2.7
NEBRASKA	28.5	30.0	1.5
NEVADA	29.9	29.6	-0.3
NEW HAMPSHIRE	26.6	30.9	4.3
NEW JERSEY	25.5	27.9	2.4
NEW MEXICO	29.9	28.0	-2.0
NEW YORK	25.8	28.0	2.3
NORTH CAROLINA	23.9	26.5	2.6
NORTH DAKOTA	29.4	29.8	0.4
OHIO	28.2	27.7	-0.5
OKLAHOMA	28.1	22.0	-6.1
OREGON	29.5	32.0	2.5
PENNSYLVANIA	30.7	25.9	-4.8
RHODE ISLAND	27.5	28.1	0.6
SOUTH CAROLINA	26.2	24.5	-1.7
SOUTH DAKOTA	25.1	25.7	0.6
TENNESSEE	27.5	29.5	2.0
TEXAS	33.4	28.7	-4.7
UTAH	23.0	26.4	3.3
VERMONT	27.4	25.9	-1.5
VIRGINIA	29.0	29.5	0.5
WASHINGTON	29.2	25.6	-3.5
WEST VIRGINIA	30.4	32.2	1.8
WISCONSIN	29.6	26.9	-2.7
WYOMING	27.9	29.9	2.0
PUERTO RICO	N/A	24.4	N/A

* EVER TOLD CHOLESTEROL IS HIGH BY HEALTH PROFESSIONAL
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 13.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2169	46.7	1.2	(44.2, 49.1)
ALASKA	1543	51.2	1.9	(47.5, 54.9)
ARIZONA	1901	34.7	1.5	(31.8, 37.6)
ARKANSAS	1787	48.6	1.4	(45.8, 51.5)
CALIFORNIA	4059	45.4	0.9	(43.6, 47.1)
COLORADO	1820	48.1	1.3	(45.5, 50.7)
CONNECTICUT	2237	50.7	1.3	(48.1, 53.3)
DELAWARE	2551	53.1	1.2	(50.7, 55.5)
DISTRICT OF CO	1502	34.6	1.5	(31.7, 37.6)
FLORIDA	3489	50.0	1.0	(48.1, 51.9)
GEORGIA	2300	37.7	1.2	(35.3, 40.2)
HAWAII	2165	45.1	1.3	(42.6, 47.6)
IDAHO	4915	44.2	0.9	(42.4, 46.1)
ILLINOIS	2884	49.0	1.1	(46.9, 51.0)
INDIANA	2364	49.8	1.2	(47.5, 52.2)
IOWA	3596	47.2	1.0	(45.3, 49.0)
KANSAS	2003	42.0	1.2	(39.6, 44.4)
KENTUCKY	3603	54.2	1.0	(52.4, 56.1)
LOUISIANA	1654	46.4	1.4	(43.6, 49.1)
MAINE	1697	53.7	1.4	(51.1, 56.3)
MARYLAND	4569	43.3	1.0	(41.3, 45.2)
MASSACHUSETTS	1762	50.2	1.4	(47.5, 52.9)
MICHIGAN	2566	51.1	1.1	(48.9, 53.2)
MINNESOTA	4809	44.3	0.8	(42.7, 45.8)
MISSISSIPPI	1598	44.4	1.4	(41.5, 47.2)
MISSOURI	1847	50.3	1.4	(47.6, 53.0)
MONTANA	1801	47.1	1.3	(44.6, 49.6)
NEBRASKA	2695	44.9	1.2	(42.5, 47.2)
NEVADA	2500	53.4	2.0	(49.5, 57.3)
NEW HAMPSHIRE	1512	56.0	1.5	(53.2, 58.9)
NEW JERSEY	2656	47.6	1.2	(45.3, 49.8)
NEW MEXICO	1810	47.4	1.3	(44.8, 50.0)
NEW YORK	3392	46.8	1.0	(44.9, 48.7)
NORTH CAROLINA	3616	48.2	1.0	(46.3, 50.1)
NORTH DAKOTA	1800	46.7	1.3	(44.1, 49.3)
OHIO	3162	47.9	1.2	(45.6, 50.2)
OKLAHOMA	1885	43.7	1.4	(41.0, 46.4)
OREGON	3308	51.1	1.0	(49.1, 53.1)
PENNSYLVANIA	3589	46.9	1.0	(45.0, 48.8)
RHODE ISLAND	1838	51.0	1.3	(48.3, 53.6)
SOUTH CAROLINA	2154	43.5	1.2	(41.1, 45.8)
SOUTH DAKOTA	2198	50.1	1.2	(47.7, 52.6)
TENNESSEE	2980	48.3	1.1	(46.2, 50.3)
TEXAS	2490	42.9	1.1	(40.7, 45.1)
UTAH	2858	31.0	1.1	(28.8, 33.2)
VERMONT	3179	52.4	1.0	(50.3, 54.4)
VIRGINIA	3506	50.3	1.3	(47.7, 52.8)
WASHINGTON	3590	50.7	1.0	(48.8, 52.7)
WEST VIRGINIA	2437	51.8	1.2	(49.4, 54.1)
WISCONSIN	2240	48.5	1.3	(46.0, 50.9)
WYOMING	2411	49.4	1.3	(46.8, 51.9)
PUERTO RICO	2257	30.1	1.1	(28.0, 32.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 48.0 RANGE = 30.1-56.0

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

August 14, 1998

TABLE 13.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	810	58.2	2.0	(54.3, 62.1)	*	1359	36.5	1.5	(33.6, 39.3)	
ALASKA	716	55.2	2.8	(49.8, 60.6)	*	827	46.8	2.6	(41.8, 51.9)	
ARIZONA	888	36.9	2.2	(32.7, 41.1)	*	1013	32.6	2.2	(28.4, 36.8)	
ARKANSAS	673	59.3	2.3	(54.8, 63.8)	*	1114	39.1	1.7	(35.8, 42.5)	
CALIFORNIA	1758	54.9	1.4	(52.2, 57.5)	*	2301	36.0	1.1	(33.8, 38.2)	
COLORADO	818	54.5	2.0	(50.7, 58.4)	*	1002	42.0	1.8	(38.5, 45.5)	
CONNECTICUT	885	52.1	2.1	(47.9, 56.3)	*	1352	49.5	1.6	(46.3, 52.6)	
DELAWARE	1012	61.2	1.9	(57.5, 64.8)	*	1539	45.7	1.5	(42.9, 48.6)	
DISTRICT OF CO	572	38.7	2.4	(34.0, 43.5)	*	930	31.2	1.8	(27.6, 34.7)	
FLORIDA	1435	56.2	1.5	(53.3, 59.1)	*	2054	44.3	1.2	(41.9, 46.8)	
GEORGIA	960	42.3	1.9	(38.7, 45.9)	*	1340	33.6	1.6	(30.5, 36.6)	
HAWAII	947	53.2	1.9	(49.5, 56.9)	*	1218	36.9	1.6	(33.6, 40.1)	
IDAHO	2057	53.1	1.4	(50.3, 55.8)	*	2858	35.8	1.2	(33.4, 38.1)	
ILLINOIS	1162	55.2	1.6	(52.0, 58.4)	*	1722	43.3	1.3	(40.7, 45.8)	
INDIANA	1063	56.4	1.7	(53.0, 59.8)	*	1301	43.9	1.6	(40.8, 47.0)	
IOWA	1476	56.1	1.4	(53.2, 58.9)	*	2120	39.1	1.2	(36.7, 41.4)	
KANSAS	786	50.5	2.0	(46.6, 54.3)	*	1217	34.2	1.5	(31.3, 37.0)	
KENTUCKY	1321	62.1	1.6	(59.1, 65.2)	*	2282	47.1	1.1	(44.9, 49.3)	
LOUISIANA	629	55.3	2.3	(50.9, 59.7)	*	1025	38.3	1.7	(35.0, 41.6)	
MAINE	751	59.0	2.0	(55.0, 62.9)	*	946	48.9	1.8	(45.3, 52.4)	
MARYLAND	1957	46.4	1.5	(43.6, 49.3)	*	2612	40.3	1.2	(37.9, 42.8)	
MASSACHUSETTS	719	52.8	2.2	(48.6, 57.0)	*	1043	48.0	1.8	(44.5, 51.4)	
MICHIGAN	1071	57.9	1.7	(54.6, 61.1)	*	1495	44.8	1.4	(42.1, 47.6)	
MINNESOTA	2201	48.4	1.2	(46.2, 50.7)	*	2608	40.3	1.0	(38.3, 42.4)	
MISSISSIPPI	592	56.7	2.4	(52.0, 61.4)	*	1006	33.5	1.7	(30.1, 36.8)	
MISSOURI	698	57.0	2.2	(52.7, 61.3)	*	1149	44.2	1.7	(41.0, 47.5)	
MONTANA	759	52.8	1.9	(49.1, 56.6)	*	1042	41.7	1.7	(38.3, 45.0)	
NEBRASKA	1046	52.3	1.9	(48.5, 56.1)	*	1649	38.0	1.6	(34.9, 41.1)	
NEVADA	1163	52.0	3.1	(45.8, 58.1)	*	1337	54.8	2.5	(50.0, 59.7)	
NEW HAMPSHIRE	603	56.2	2.3	(51.7, 60.7)	*	909	55.9	1.8	(52.3, 59.4)	
NEW JERSEY	1056	53.0	1.8	(49.5, 56.5)	*	1600	42.7	1.5	(39.9, 45.6)	
NEW MEXICO	753	53.0	2.0	(49.1, 56.9)	*	1057	42.1	1.6	(38.9, 45.3)	
NEW YORK	1350	51.0	1.6	(47.9, 54.0)	*	2042	43.2	1.2	(40.8, 45.5)	
NORTH CAROLINA	1432	56.4	1.5	(53.4, 59.3)	*	2184	40.9	1.2	(38.5, 43.2)	
NORTH DAKOTA	800	52.7	2.0	(48.9, 56.6)	*	1000	40.9	1.7	(37.5, 44.2)	
OHIO	1229	54.1	1.9	(50.4, 57.7)	*	1933	42.3	1.4	(39.5, 45.2)	
OKLAHOMA	830	48.3	2.2	(44.0, 52.5)	*	1055	39.6	1.7	(36.3, 42.9)	
OREGON	1396	57.2	1.6	(54.1, 60.3)	*	1912	45.4	1.4	(42.7, 48.0)	
PENNSYLVANIA	1439	52.8	1.5	(49.9, 55.8)	*	2150	41.7	1.3	(39.2, 44.1)	
RHODE ISLAND	755	53.9	2.0	(49.9, 57.9)	*	1083	48.3	1.7	(44.9, 51.7)	
SOUTH CAROLINA	820	54.1	1.9	(50.4, 57.8)	*	1334	33.9	1.5	(31.0, 36.8)	
SOUTH DAKOTA	956	59.8	1.8	(56.2, 63.4)	*	1242	41.1	1.6	(38.0, 44.1)	
TENNESSEE	1046	55.4	1.7	(52.0, 58.7)	*	1934	41.9	1.3	(39.4, 44.4)	
TEXAS	1022	52.4	1.7	(49.0, 55.8)	*	1468	34.0	1.4	(31.3, 36.7)	
UTAH	1274	38.6	1.7	(35.2, 41.9)	*	1584	23.7	1.4	(21.0, 26.4)	
VERMONT	1388	57.8	1.5	(54.7, 60.8)	*	1791	47.3	1.4	(44.7, 50.0)	
VIRGINIA	1409	54.1	1.9	(50.3, 57.8)	*	2097	46.7	1.7	(43.4, 50.0)	
WASHINGTON	1540	56.2	1.5	(53.3, 59.1)	*	2050	45.5	1.3	(42.9, 48.0)	
WEST VIRGINIA	966	58.2	1.8	(54.6, 61.8)	*	1471	46.0	1.4	(43.3, 48.7)	

WISCONSIN	989	57.6	1.9	(53.8, 61.3)	*	1251	40.0	1.6	(36.8, 43.2)
WYOMING	1019	55.1	1.9	(51.3, 58.9)	*	1392	43.7	1.6	(40.5, 46.9)
PUERTO RICO	934	44.1	1.8	(40.6, 47.5)	*	1323	17.8	1.1	(15.6, 20.0)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 54.3
 RANGE = 36.9-62.1 NUMBER OF PARTICIPANTS = 52
 MEDIAN = 41.8
 RANGE = 17.8-55.9

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

August 14, 1998

TABLE 13.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	43.9	46.7	2.8
ALASKA	53.7	51.2	-2.5
ARIZONA	48.8	34.7	-14.2
ARKANSAS	43.3	48.6	5.4
CALIFORNIA	43.6	45.4	1.8
COLORADO	46.9	48.1	1.2
CONNECTICUT	50.0	50.7	0.7
DELAWARE	51.0	53.1	2.1
DISTRICT OF CO	38.3	34.6	-3.7
FLORIDA	52.5	50.0	-2.5
GEORGIA	34.4	37.7	3.3
HAWAII	46.3	45.1	-1.2
IDAHO	44.0	44.2	0.2
ILLINOIS	48.2	49.0	0.8
INDIANA	52.4	49.8	-2.6
IOWA	47.4	47.2	-0.3
KANSAS	41.6	42.0	0.5
KENTUCKY	54.8	54.2	-0.6
LOUISIANA	47.4	46.4	-1.0
MAINE	53.1	53.7	0.6
MARYLAND	44.3	43.3	-1.1
MASSACHUSETTS	53.8	50.2	-3.5
MICHIGAN	51.5	51.1	-0.5
MINNESOTA	43.7	44.3	0.6
MISSISSIPPI	44.3	44.4	0.0
MISSOURI	52.5	50.3	-2.2
MONTANA	48.7	47.1	-1.6
NEBRASKA	45.7	44.9	-0.9
NEVADA	55.0	53.4	-1.6
NEW HAMPSHIRE	53.9	56.0	2.1
NEW JERSEY	47.5	47.6	0.1
NEW MEXICO	47.0	47.4	0.4
NEW YORK	46.9	46.8	-0.1
NORTH CAROLINA	48.8	48.2	-0.6
NORTH DAKOTA	48.8	46.7	-2.2
OHIO	49.8	47.9	-2.0
OKLAHOMA	42.5	43.7	1.2
OREGON	51.1	51.1	0.0

PENNSYLVANIA	47.8	46.9	-0.8
RHODE ISLAND	49.6	51.0	1.3
SOUTH CAROLINA	44.7	43.5	-1.2
SOUTH DAKOTA	45.9	50.1	4.2
TENNESSEE	47.8	48.3	0.4
TEXAS	44.1	42.9	-1.2
UTAH	32.3	31.0	-1.4
VERMONT	53.3	52.4	-0.9
VIRGINIA	47.8	50.3	2.5
WASHINGTON	50.3	50.7	0.4
WEST VIRGINIA	49.5	51.8	2.2
WISCONSIN	50.8	48.5	-2.4
WYOMING	49.7	49.4	-0.3
PUERTO RICO	30.7	30.1	-0.6

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

N/A - NOT AVAILABLE

August 14, 1998

TABLE 14.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2169	24.7	1.0	(22.6, 26.7)
ALASKA	1543	26.5	1.7	(23.3, 29.8)
ARIZONA	1901	21.1	1.3	(18.6, 23.6)
ARKANSAS	1787	28.4	1.3	(25.8, 31.1)
CALIFORNIA	4059	18.4	0.7	(17.0, 19.8)
COLORADO	1819	22.6	1.1	(20.3, 24.8)
CONNECTICUT	2235	21.6	1.1	(19.5, 23.7)
DELAWARE	2550	26.6	1.1	(24.4, 28.7)
DISTRICT OF CO	1501	18.8	1.2	(16.4, 21.2)
FLORIDA	3488	23.6	0.8	(22.0, 25.2)
GEORGIA	2300	22.4	1.1	(20.3, 24.5)
HAWAII	2165	18.7	1.0	(16.8, 20.6)
IDAHO	4915	19.9	0.7	(18.5, 21.4)
ILLINOIS	2884	23.2	0.9	(21.5, 24.9)
INDIANA	2364	26.4	1.1	(24.3, 28.5)
IOWA	3596	23.1	0.8	(21.5, 24.7)
KANSAS	2003	22.6	1.0	(20.6, 24.6)
KENTUCKY	3603	30.7	0.9	(29.0, 32.5)
LOUISIANA	1654	24.5	1.2	(22.1, 26.8)
MAINE	1697	22.7	1.1	(20.5, 24.8)
MARYLAND	4568	20.4	0.8	(18.9, 22.0)
MASSACHUSETTS	1758	20.5	1.1	(18.2, 22.7)
MICHIGAN	2566	26.0	0.9	(24.2, 27.9)
MINNESOTA	4809	21.9	0.7	(20.6, 23.1)
MISSISSIPPI	1598	23.1	1.3	(20.6, 25.6)
MISSOURI	1846	28.6	1.3	(26.1, 31.1)
MONTANA	1801	20.5	1.0	(18.5, 22.5)
NEBRASKA	2694	22.1	1.0	(20.1, 24.2)
NEVADA	2500	28.0	1.8	(24.6, 31.5)
NEW HAMPSHIRE	1512	24.7	1.3	(22.2, 27.2)
NEW JERSEY	2654	21.4	1.0	(19.5, 23.2)

NEW MEXICO	1810	22.1	1.1	(19.9, 24.2)
NEW YORK	3391	23.1	0.8	(21.5, 24.7)
NORTH CAROLINA	3615	25.9	0.9	(24.2, 27.5)
NORTH DAKOTA	1800	22.3	1.1	(20.1, 24.4)
OHIO	3161	25.1	1.0	(23.1, 27.1)
OKLAHOMA	1885	24.6	1.2	(22.2, 27.0)
OREGON	3308	20.7	0.9	(19.0, 22.4)
PENNSYLVANIA	3589	24.2	0.8	(22.6, 25.8)
RHODE ISLAND	1838	24.3	1.2	(21.9, 26.7)
SOUTH CAROLINA	2153	23.4	1.0	(21.4, 25.4)
SOUTH DAKOTA	2197	24.3	1.1	(22.2, 26.4)
TENNESSEE	2980	26.9	1.0	(25.1, 28.8)
TEXAS	2489	22.6	1.0	(20.7, 24.4)
UTAH	2858	13.8	0.8	(12.1, 15.4)
VERMONT	3179	23.3	1.0	(21.4, 25.1)
VIRGINIA	3504	24.4	1.1	(22.3, 26.5)
WASHINGTON	3589	23.8	0.9	(22.1, 25.6)
WEST VIRGINIA	2437	27.4	1.0	(25.4, 29.5)
WISCONSIN	2240	23.2	1.1	(21.0, 25.3)
WYOMING	2411	24.0	1.2	(21.6, 26.3)
PUERTO RICO	2257	14.5	0.8	(12.8, 16.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 23.2 RANGE = 13.8-30.7 YEAR 2000 OBJ. 3.4
 * A CURRENT SMOKER (EVERDAY OR SOME DAYS)

August 14, 1998

TABLE 14.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR
	*	*	*	*	*	*	*	95% CONFIDENCE INTERVAL
ALABAMA	810	28.6	1.7	(25.3, 31.9)	*	1359	21.2	1.2 (18.7, 23.6)
ALASKA	716	27.2	2.5	(22.3, 32.1)	*	827	25.8	2.2 (21.5, 30.1)
ARIZONA	888	22.0	2.0	(18.1, 25.9)	*	1013	20.2	1.8 (16.6, 23.7)
ARKANSAS	673	32.0	2.2	(27.6, 36.4)	*	1114	25.2	1.5 (22.2, 28.2)
CALIFORNIA	1758	22.4	1.2	(20.1, 24.7)	*	2301	14.5	0.8 (12.9, 16.1)
COLORADO	818	24.0	1.6	(20.8, 27.2)	*	1001	21.1	1.5 (18.1, 24.1)
CONNECTICUT	884	21.3	1.6	(18.1, 24.4)	*	1351	21.9	1.4 (19.2, 24.5)
DELAWARE	1012	29.2	1.8	(25.8, 32.6)	*	1538	24.2	1.3 (21.7, 26.7)
DISTRICT OF CO	572	22.7	2.0	(18.7, 26.7)	*	929	15.4	1.4 (12.7, 18.2)
FLORIDA	1434	26.0	1.3	(23.4, 28.5)	*	2054	21.4	1.0 (19.5, 23.3)
GEORGIA	960	25.2	1.6	(22.0, 28.4)	*	1340	19.8	1.4 (17.2, 22.5)
HAWAII	947	21.5	1.5	(18.6, 24.4)	*	1218	15.8	1.3 (13.3, 18.3)
IDAHO	2057	21.8	1.1	(19.6, 24.0)	*	2858	18.1	1.0 (16.2, 20.0)
ILLINOIS	1162	25.0	1.4	(22.3, 27.7)	*	1722	21.6	1.1 (19.4, 23.7)
INDIANA	1063	29.2	1.6	(26.1, 32.4)	*	1301	23.8	1.4 (21.1, 26.4)
IOWA	1476	25.5	1.2	(23.1, 27.9)	*	2120	21.0	1.0 (19.0, 23.0)
KANSAS	786	26.7	1.7	(23.4, 30.1)	*	1217	18.8	1.2 (16.5, 21.2)
KENTUCKY	1321	33.0	1.5	(30.1, 35.9)	*	2282	28.7	1.1 (26.6, 30.7)
LOUISIANA	629	29.2	2.1	(25.1, 33.2)	*	1025	20.3	1.4 (17.6, 22.9)
MAINE	751	25.2	1.7	(21.8, 28.5)	*	946	20.4	1.4 (17.6, 23.2)
MARYLAND	1957	21.7	1.2	(19.3, 24.0)	*	2611	19.3	1.0 (17.4, 21.3)
MASSACHUSETTS	716	21.8	1.9	(18.1, 25.5)	*	1042	19.3	1.3 (16.6, 21.9)
MICHIGAN	1071	29.6	1.5	(26.6, 32.6)	*	1495	22.8	1.1 (20.5, 25.0)

MINNESOTA	2201	24.1	1.0	(22.1, 26.0)	*	2608	19.8	0.8	(18.1, 21.4)
MISSISSIPPI	592	28.3	2.1	(24.1, 32.5)	*	1006	18.5	1.4	(15.8, 21.3)
MISSOURI	697	31.6	2.1	(27.5, 35.7)	*	1149	26.0	1.5	(23.1, 28.8)
MONTANA	759	20.8	1.5	(17.8, 23.8)	*	1042	20.2	1.4	(17.6, 22.9)
NEBRASKA	1045	24.3	1.6	(21.2, 27.4)	*	1649	20.1	1.4	(17.4, 22.8)
NEVADA	1163	25.6	2.6	(20.6, 30.6)	*	1337	30.5	2.3	(25.9, 35.1)
NEW HAMPSHIRE	603	25.9	2.1	(21.8, 30.0)	*	909	23.6	1.5	(20.5, 26.6)
NEW JERSEY	1055	23.4	1.6	(20.4, 26.4)	*	1599	19.5	1.2	(17.2, 21.8)
NEW MEXICO	753	21.6	1.7	(18.3, 24.8)	*	1057	22.5	1.4	(19.8, 25.2)
NEW YORK	1350	24.9	1.3	(22.3, 27.5)	*	2041	21.5	1.0	(19.5, 23.5)
NORTH CAROLINA	1432	29.7	1.4	(27.0, 32.3)	*	2183	22.4	1.0	(20.4, 24.4)
NORTH DAKOTA	800	24.3	1.6	(21.1, 27.5)	*	1000	20.3	1.4	(17.5, 23.0)
OHIO	1228	26.3	1.6	(23.2, 29.5)	*	1933	24.0	1.3	(21.5, 26.4)
OKLAHOMA	830	25.2	1.9	(21.5, 28.9)	*	1055	24.0	1.5	(21.0, 27.1)
OREGON	1396	22.1	1.4	(19.4, 24.8)	*	1912	19.3	1.1	(17.2, 21.4)
PENNSYLVANIA	1439	26.2	1.3	(23.5, 28.8)	*	2150	22.4	1.0	(20.4, 24.4)
RHODE ISLAND	755	25.5	1.8	(22.0, 29.1)	*	1083	23.2	1.6	(20.0, 26.4)
SOUTH CAROLINA	819	29.5	1.8	(26.0, 32.9)	*	1334	18.0	1.2	(15.6, 20.3)
SOUTH DAKOTA	955	28.0	1.7	(24.8, 31.3)	*	1242	20.8	1.3	(18.3, 23.4)
TENNESSEE	1046	27.9	1.6	(24.9, 31.0)	*	1934	26.0	1.1	(23.8, 28.2)
TEXAS	1021	28.0	1.6	(24.9, 31.0)	*	1468	17.4	1.1	(15.3, 19.6)
UTAH	1274	16.1	1.3	(13.6, 18.6)	*	1584	11.5	1.0	(9.5, 13.5)
VERMONT	1388	25.2	1.5	(22.3, 28.1)	*	1791	21.5	1.2	(19.1, 23.8)
VIRGINIA	1407	26.0	1.7	(22.7, 29.4)	*	2097	22.9	1.3	(20.3, 25.4)
WASHINGTON	1540	25.0	1.4	(22.3, 27.8)	*	2049	22.7	1.1	(20.5, 24.8)
WEST VIRGINIA	966	27.3	1.6	(24.2, 30.4)	*	1471	27.6	1.3	(25.0, 30.2)
WISCONSIN	989	25.6	1.7	(22.3, 29.0)	*	1251	20.9	1.4	(18.1, 23.7)
WYOMING	1019	23.9	1.9	(20.1, 27.7)	*	1392	24.0	1.5	(21.2, 26.9)
PUERTO RICO	934	21.5	1.5	(18.6, 24.4)	*	1323	8.2	0.8	(6.6, 9.7)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 25.3
 RANGE = 16.1-33.0

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 21.1
 RANGE = 8.2-30.5

* A CURRENT SMOKER (EVERYDAY OR SOME DAYS)

August 14, 1998

TABLE 14.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	22.4	24.7	2.2
ALASKA	27.7	26.5	-1.1
ARIZONA	23.7	21.1	-2.6
ARKANSAS	25.3	28.4	3.2
CALIFORNIA	18.6	18.4	-0.2
COLORADO	22.8	22.6	-0.2
CONNECTICUT	21.8	21.6	-0.2
DELAWARE	24.2	26.6	2.4
DISTRICT OF CO	20.5	18.8	-1.7
FLORIDA	21.8	23.6	1.8
GEORGIA	20.3	22.4	2.1
HAWAII	22.2	18.7	-3.5
IDAHO	21.1	19.9	-1.2

ILLINOIS	24.8	23.2	-1.6
INDIANA	28.6	26.4	-2.3
IOWA	23.6	23.1	-0.5
KANSAS	22.1	22.6	0.6
KENTUCKY	31.7	30.7	-0.9
LOUISIANA	25.9	24.5	-1.4
MAINE	25.3	22.7	-2.6
MARYLAND	20.9	20.4	-0.5
MASSACHUSETTS	23.4	20.5	-2.9
MICHIGAN	25.6	26.0	0.4
MINNESOTA	20.6	21.9	1.3
MISSISSIPPI	23.2	23.1	-0.1
MISSOURI	27.8	28.6	0.8
MONTANA	21.7	20.5	-1.2
NEBRASKA	22.1	22.1	0.0
NEVADA	28.2	28.0	-0.1
NEW HAMPSHIRE	24.8	24.7	-0.1
NEW JERSEY	22.7	21.4	-1.4
NEW MEXICO	22.8	22.1	-0.8
NEW YORK	23.3	23.1	-0.1
NORTH CAROLINA	25.7	25.9	0.2
NORTH DAKOTA	23.4	22.3	-1.1
OHIO	28.4	25.1	-3.3
OKLAHOMA	24.1	24.6	0.5
OREGON	23.4	20.7	-2.7
PENNSYLVANIA	24.5	24.2	-0.3
RHODE ISLAND	22.5	24.3	1.8
SOUTH CAROLINA	24.5	23.4	-1.1
SOUTH DAKOTA	20.7	24.3	3.6
TENNESSEE	28.0	26.9	-1.1
TEXAS	22.9	22.6	-0.4
UTAH	15.9	13.8	-2.2
VERMONT	24.1	23.3	-0.8
VIRGINIA	24.8	24.4	-0.4
WASHINGTON	23.4	23.8	0.4
WEST VIRGINIA	26.6	27.4	0.8
WISCONSIN	24.9	23.2	-1.7
WYOMING	24.6	24.0	-0.6
PUERTO RICO	14.5	14.5	-0.1

* A CURRENT SMOKER (EVERDAY OR SOME DAYS)

N/A - NOT AVAILABLE

August 14, 1998

TABLE 15.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LACK OF SAFETY BELT USAGE*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, OR NEVER RIDE IN CAR

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2166	33.8	1.2	(31.4, 36.2)
ALASKA	1493	34.7	1.8	(31.1, 38.2)
ARIZONA	1894	19.5	1.3	(17.0, 22.0)
ARKANSAS	1725	34.5	1.4	(31.7, 37.3)

CALIFORNIA	4056	12.8	0.6	(11.6, 13.9)
COLORADO	1821	28.6	1.2	(26.2, 31.0)
CONNECTICUT	2237	30.9	1.2	(28.5, 33.2)
DELAWARE	2554	30.1	1.1	(27.9, 32.2)
DISTRICT OF CO	1494	22.0	1.4	(19.3, 24.7)
FLORIDA	3475	23.9	0.9	(22.2, 25.5)
GEORGIA	2300	24.6	1.1	(22.4, 26.8)
HAWAII	2149	12.9	0.9	(11.2, 14.6)
IDAH0	4913	40.4	0.9	(38.6, 42.2)
ILLINOIS	1435	31.9	1.4	(29.3, 34.5)
INDIANA	2365	38.1	1.2	(35.9, 40.4)
IOWA	3593	32.8	0.9	(31.0, 34.6)
KANSAS	1998	46.2	1.2	(43.7, 48.6)
KENTUCKY	3604	34.6	1.0	(32.7, 36.5)
LOUISIANA	1647	25.7	1.3	(23.2, 28.3)
MAINE	1693	30.5	1.3	(28.0, 33.0)
MARYLAND	4555	23.9	0.9	(22.2, 25.6)
MASSACHUSETTS	1753	37.3	1.4	(34.6, 39.9)
MICHIGAN	2559	27.7	1.0	(25.8, 29.7)
MINNESOTA	4806	40.3	0.8	(38.7, 41.8)
MISSISSIPPI	1597	43.5	1.4	(40.7, 46.3)
MISSOURI	1842	38.1	1.4	(35.4, 40.9)
MONTANA	1802	42.4	1.3	(39.8, 45.0)
NEBRASKA	2688	42.3	1.3	(39.8, 44.7)
NEVADA	2498	26.3	1.9	(22.6, 30.0)
NEW HAMPSHIRE	1505	41.8	1.5	(38.8, 44.7)
NEW JERSEY	2647	27.8	1.0	(25.7, 29.8)
NEW MEXICO	1809	16.5	1.0	(14.5, 18.5)
NEW YORK	3308	26.2	0.9	(24.4, 28.0)
NORTH CAROLINA	3621	15.3	0.7	(13.9, 16.6)
NORTH DAKOTA	1792	59.8	1.3	(57.2, 62.4)
OHIO	3155	30.0	1.1	(27.8, 32.2)
OKLAHOMA	1884	36.9	1.4	(34.3, 39.5)
OREGON	3316	16.0	0.8	(14.4, 17.5)
PENNSYLVANIA	3567	32.7	0.9	(30.9, 34.6)
RHODE ISLAND	1836	43.2	1.3	(40.6, 45.8)
SOUTH CAROLINA	2150	19.5	1.0	(17.6, 21.5)
SOUTH DAKOTA	2185	58.0	1.2	(55.5, 60.4)
TENNESSEE	2967	33.6	1.0	(31.6, 35.6)
TEXAS	2485	18.8	0.9	(17.0, 20.5)
UTAH	2862	35.0	1.2	(32.7, 37.3)
VERMONT	3181	26.3	0.9	(24.4, 28.1)
VIRGINIA	3510	28.3	1.1	(26.2, 30.4)
WASHINGTON	3587	24.2	0.9	(22.4, 26.0)
WEST VIRGINIA	2429	29.4	1.0	(27.4, 31.4)
WISCONSIN	2242	38.7	1.3	(36.2, 41.1)
WYOMING	2410	49.8	1.3	(47.3, 52.3)
PUERTO RICO	2242	24.7	1.0	(22.6, 26.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 30.7 RANGE = 12.8-59.8 YEAR 2000 OBJ. 9.12

* DOES NOT ALWAYS WEAR A SAFETY BELT

August 14, 1998

TABLE 15.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LACK OF SAFETY BELT USAGE* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, OR NEVER RIDE IN CAR

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	808	43.5	2.0	(39.7, 47.4)	*	1358	25.2	1.4	(22.5, 27.9)	*
ALASKA	697	39.8	2.7	(34.5, 45.1)	*	796	29.0	2.3	(24.6, 33.4)	*
ARIZONA	888	19.7	1.7	(16.3, 23.1)	*	1006	19.3	1.8	(15.7, 22.8)	*
ARKANSAS	643	42.2	2.4	(37.5, 46.8)	*	1082	27.7	1.6	(24.6, 30.7)	*
CALIFORNIA	1755	14.9	0.9	(13.1, 16.7)	*	2301	10.7	0.7	(9.2, 12.1)	*
COLORADO	818	31.9	1.9	(28.2, 35.5)	*	1003	25.5	1.6	(22.4, 28.5)	*
CONNECTICUT	883	37.1	2.0	(33.1, 41.1)	*	1354	25.2	1.3	(22.6, 27.7)	*
DELAWARE	1013	37.9	1.8	(34.4, 41.4)	*	1541	23.0	1.2	(20.5, 25.4)	*
DISTRICT OF CO	569	27.4	2.5	(22.5, 32.2)	*	925	17.5	1.4	(14.7, 20.2)	*
FLORIDA	1432	31.9	1.4	(29.2, 34.6)	*	2043	16.5	0.9	(14.7, 18.3)	*
GEORGIA	959	31.2	1.8	(27.6, 34.7)	*	1341	18.7	1.3	(16.2, 21.1)	*
HAWAII	943	16.0	1.4	(13.3, 18.7)	*	1206	9.6	1.0	(7.6, 11.6)	*
IDAHO	2054	48.9	1.4	(46.1, 51.7)	*	2859	32.2	1.1	(30.1, 34.3)	*
ILLINOIS	576	37.1	2.2	(32.9, 41.3)	*	859	27.3	1.7	(24.0, 30.6)	*
INDIANA	1063	47.1	1.8	(43.7, 50.5)	*	1302	30.0	1.5	(27.1, 32.9)	*
IOWA	1472	41.1	1.5	(38.2, 43.9)	*	2121	25.3	1.1	(23.1, 27.4)	*
KANSAS	782	53.3	2.0	(49.5, 57.2)	*	1216	39.5	1.5	(36.6, 42.5)	*
KENTUCKY	1321	45.1	1.6	(42.1, 48.2)	*	2283	25.0	1.0	(23.0, 27.1)	*
LOUISIANA	627	33.8	2.2	(29.5, 38.1)	*	1020	18.5	1.4	(15.7, 21.2)	*
MAINE	748	38.3	2.0	(34.4, 42.3)	*	945	23.3	1.5	(20.4, 26.3)	*
MARYLAND	1950	31.3	1.4	(28.6, 33.9)	*	2605	17.2	1.0	(15.3, 19.1)	*
MASSACHUSETTS	720	43.2	2.1	(39.0, 47.3)	*	1033	32.0	1.7	(28.6, 35.3)	*
MICHIGAN	1068	35.0	1.6	(31.8, 38.1)	*	1491	21.1	1.2	(18.9, 23.4)	*
MINNESOTA	2204	48.9	1.2	(46.6, 51.2)	*	2602	32.2	1.0	(30.2, 34.1)	*
MISSISSIPPI	592	52.1	2.3	(47.6, 56.7)	*	1005	35.8	1.7	(32.5, 39.1)	*
MISSOURI	693	45.4	2.2	(41.1, 49.6)	*	1149	31.7	1.7	(28.4, 35.0)	*
MONTANA	760	51.3	2.0	(47.5, 55.2)	*	1042	34.0	1.6	(30.7, 37.2)	*
NEBRASKA	1044	50.6	1.9	(46.8, 54.3)	*	1644	34.6	1.5	(31.6, 37.7)	*
NEVADA	1159	37.6	3.2	(31.4, 43.8)	*	1339	14.8	1.6	(11.7, 17.9)	*
NEW HAMPSHIRE	600	51.9	2.4	(47.3, 56.5)	*	905	32.2	1.8	(28.7, 35.7)	*
NEW JERSEY	1053	34.5	1.7	(31.1, 37.8)	*	1594	21.7	1.2	(19.4, 24.0)	*
NEW MEXICO	754	20.8	1.7	(17.5, 24.0)	*	1055	12.5	1.1	(10.3, 14.7)	*
NEW YORK	1326	33.9	1.5	(30.9, 36.8)	*	1982	19.3	1.0	(17.3, 21.4)	*
NORTH CAROLINA	1435	21.6	1.2	(19.2, 24.0)	*	2186	9.5	0.7	(8.1, 10.8)	*
NORTH DAKOTA	793	71.0	1.8	(67.4, 74.5)	*	999	49.1	1.8	(45.6, 52.5)	*
OHIO	1224	34.4	1.8	(30.8, 37.9)	*	1931	26.1	1.4	(23.4, 28.9)	*
OKLAHOMA	829	43.3	2.1	(39.3, 47.4)	*	1055	31.0	1.7	(27.8, 34.3)	*
OREGON	1398	21.0	1.3	(18.5, 23.6)	*	1918	11.3	0.9	(9.4, 13.1)	*
PENNSYLVANIA	1431	40.6	1.5	(37.7, 43.5)	*	2136	25.7	1.2	(23.4, 28.0)	*
RHODE ISLAND	756	47.7	2.0	(43.7, 51.7)	*	1080	39.1	1.8	(35.7, 42.6)	*
SOUTH CAROLINA	818	26.0	1.7	(22.6, 29.4)	*	1332	13.7	1.1	(11.6, 15.8)	*
SOUTH DAKOTA	954	66.4	1.8	(62.9, 69.8)	*	1231	50.0	1.7	(46.8, 53.2)	*
TENNESSEE	1039	39.7	1.7	(36.4, 42.9)	*	1928	28.2	1.2	(25.9, 30.4)	*
TEXAS	1024	24.5	1.5	(21.6, 27.4)	*	1461	13.3	1.0	(11.4, 15.2)	*
UTAH	1275	43.1	1.8	(39.6, 46.6)	*	1587	27.4	1.4	(24.6, 30.2)	*
VERMONT	1393	33.8	1.4	(31.0, 36.7)	*	1788	19.2	1.2	(16.9, 21.5)	*
VIRGINIA	1414	34.8	1.7	(31.4, 38.2)	*	2096	22.2	1.4	(19.5, 24.9)	*
WASHINGTON	1537	30.8	1.5	(27.9, 33.6)	*	2050	17.9	1.1	(15.7, 20.1)	*
WEST VIRGINIA	963	39.3	1.7	(35.9, 42.7)	*	1466	20.6	1.2	(18.3, 22.9)	*
WISCONSIN	990	49.0	2.0	(45.2, 52.8)	*	1252	29.1	1.5	(26.2, 32.0)	*
WYOMING	1019	56.0	1.9	(52.3, 59.8)	*	1391	43.8	1.7	(40.5, 47.0)	*
PUERTO RICO	933	28.9	1.7	(25.6, 32.2)	*	1309	20.9	1.3	(18.4, 23.4)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 38.1
 RANGE = 14.9-71.0

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 25.2
 RANGE = 9.5-50.0

* DOES NOT ALWAYS WEAR A SAFETY BELT

August 14, 1998

TABLE 15.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 LACK OF SAFETY BELT USAGE*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, OR NEVER RIDE IN CAR

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	32.6	33.8	1.2
ALASKA	33.7	34.7	1.0
ARIZONA	26.0	19.5	-6.5
ARKANSAS	32.8	34.5	1.7
CALIFORNIA	14.9	12.8	-2.1
COLORADO	35.6	28.6	-7.0
CONNECTICUT	31.0	30.9	-0.1
DELAWARE	29.1	30.1	1.0
DISTRICT OF CO	N/A	22.0	N/A
FLORIDA	24.4	23.9	-0.6
GEORGIA	36.2	24.6	-11.5
HAWAII	12.7	12.9	0.2
IDAHO	42.6	40.4	-2.3
ILLINOIS	30.9	31.9	1.0
INDIANA	43.5	38.1	-5.3
IOWA	37.3	32.8	-4.5
KANSAS	42.4	46.2	3.8
KENTUCKY	34.8	34.6	-0.3
LOUISIANA	32.9	25.7	-7.2
MAINE	48.8	30.5	-18.3
MARYLAND	25.5	23.9	-1.6
MASSACHUSETTS	42.0	37.3	-4.7
MICHIGAN	29.8	27.7	-2.0
MINNESOTA	41.3	40.3	-1.0
MISSISSIPPI	41.9	43.5	1.5
MISSOURI	36.2	38.1	1.9
MONTANA	43.4	42.4	-1.0
NEBRASKA	47.0	42.3	-4.8
NEVADA	28.5	26.3	-2.3
NEW HAMPSHIRE	47.1	41.8	-5.3
NEW JERSEY	31.7	27.8	-3.9
NEW MEXICO	15.6	16.5	0.9
NEW YORK	26.6	26.2	-0.4
NORTH CAROLINA	14.1	15.3	1.2
NORTH DAKOTA	58.5	59.8	1.3
OHIO	31.9	30.0	-1.8
OKLAHOMA	46.0	36.9	-9.1
OREGON	17.2	16.0	-1.2
PENNSYLVANIA	39.3	32.7	-6.6
RHODE ISLAND	50.0	43.2	-6.8
SOUTH CAROLINA	23.2	19.5	-3.7
SOUTH DAKOTA	57.0	58.0	1.0
TENNESSEE	38.5	33.6	-4.9
TEXAS	22.4	18.8	-3.6
UTAH	39.8	35.0	-4.8
VERMONT	28.9	26.3	-2.6
VIRGINIA	27.8	28.3	0.5
WASHINGTON	22.1	24.2	2.1
WEST VIRGINIA	30.1	29.4	-0.7
WISCONSIN	43.9	38.7	-5.2
WYOMING	52.3	49.8	-2.5
PUERTO RICO	N/A	24.7	N/A

* DOES NOT ALWAYS WEAR A SAFETY BELT

N/A - NOT AVAILABLE

August 14, 1998

TABLE 16.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY BELT*
 DENOMINATOR IS PERSONS WHO HAVE A CHILD BETWEEN THE AGE OF 5 AND 15
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES IN A CAR

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	594	75.7	2.1	(71.6, 79.8)
ALASKA	555	80.0	2.5	(75.1, 84.9)
ARIZONA	426	93.3	1.4	(90.6, 95.9)
ARKANSAS	490	78.5	2.1	(74.5, 82.6)
CALIFORNIA	1014	91.8	1.1	(89.7, 93.8)
COLORADO	485	82.1	1.9	(78.3, 85.9)
CONNECTICUT	581	88.1	1.6	(85.0, 91.1)
DELAWARE	662	84.8	1.6	(81.7, 87.8)
DISTRICT OF CO	271	89.4	1.9	(85.7, 93.2)
FLORIDA	838	85.1	1.4	(82.4, 87.8)
GEORGIA	718	85.6	1.5	(82.7, 88.6)
HAWAII	612	77.2	2.1	(73.1, 81.3)
IDAHO	1524	66.3	1.6	(63.3, 69.4)
ILLINOIS	419	78.0	2.3	(73.5, 82.5)
INDIANA	692	75.8	1.8	(72.2, 79.4)
IOWA	939	71.6	1.7	(68.3, 75.0)
KANSAS	458	71.2	2.3	(66.7, 75.7)
KENTUCKY	900	82.0	1.4	(79.2, 84.7)
LOUISIANA	493	81.4	2.1	(77.4, 85.4)
MAINE	471	95.3	1.1	(93.2, 97.4)
MARYLAND	1231	87.5	1.2	(85.2, 89.9)
MASSACHUSETTS	440	84.9	2.0	(81.1, 88.8)
MICHIGAN	769	83.1	1.5	(80.3, 86.0)
MINNESOTA	1215	69.0	1.5	(66.1, 71.9)
MISSISSIPPI	508	63.7	2.6	(58.6, 68.8)
MISSOURI	510	72.1	2.5	(67.3, 76.9)
MONTANA	473	69.1	2.3	(64.7, 73.6)
NEBRASKA	696	67.8	2.2	(63.6, 72.0)
NEVADA	715	84.6	2.4	(80.0, 89.2)
NEW HAMPSHIRE	410	88.6	1.8	(85.1, 92.1)
NEW JERSEY	725	84.4	1.7	(81.0, 87.8)
NEW MEXICO	595	88.6	1.5	(85.7, 91.5)
NEW YORK	865	86.3	1.4	(83.6, 89.0)
NORTH CAROLINA	936	88.4	1.1	(86.2, 90.6)
NORTH DAKOTA	425	61.4	2.4	(56.7, 66.2)
OHIO	784	85.3	1.5	(82.3, 88.2)
OKLAHOMA	449	78.9	2.2	(74.7, 83.2)
OREGON	895	91.1	1.2	(88.9, 93.4)
PENNSYLVANIA	925	86.1	1.3	(83.6, 88.6)
RHODE ISLAND	466	88.5	1.7	(85.1, 91.9)
SOUTH CAROLINA	524	89.2	1.5	(86.2, 92.1)
SOUTH DAKOTA	590	54.3	2.3	(49.8, 58.8)
TENNESSEE	723	80.9	1.7	(77.5, 84.2)
TEXAS	794	78.9	1.7	(75.6, 82.2)
UTAH	1038	61.7	2.0	(57.7, 65.6)
VERMONT	899	92.4	1.7	(89.1, 95.7)
VIRGINIA	963	82.8	1.5	(79.9, 85.7)
WASHINGTON	1006	86.1	1.2	(83.6, 88.5)
WEST VIRGINIA	582	84.4	1.6	(81.1, 87.6)

WISCONSIN	625	73.3	2.3	(68.9, 77.8)
WYOMING	747	63.7	2.1	(59.7, 67.8)
PUERTO RICO	716	73.2	1.8	(69.6, 76.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 82.4 RANGE = 54.3-95.3

APPROXIMATES YEAR 2000 OBJ. 9.12

* OLDEST CHILD UNDER 16 ALWAYS USE SAFETY BELT

August 14, 1998

TABLE 16.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY BELT* -BY SEX
 DENOMINATOR IS PERSONS WHO HAVE A CHILD BETWEEN THE AGE OF 5 AND 15
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES IN A CAR

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	200	75.0	3.4	(68.3, 81.7)	*	394	76.3	2.5	(71.4, 81.3)	*
ALASKA	243	79.5	4.0	(71.7, 87.3)	*	312	80.6	2.8	(75.0, 86.2)	*
ARIZONA	190	94.4	1.5	(91.4, 97.4)	*	236	92.3	2.2	(88.0, 96.5)	*
ARKANSAS	186	80.3	3.2	(74.1, 86.5)	*	304	76.8	2.7	(71.5, 82.1)	*
CALIFORNIA	405	92.5	1.6	(89.4, 95.6)	*	609	91.1	1.4	(88.3, 93.8)	*
COLORADO	204	84.5	2.8	(79.1, 89.9)	*	281	79.8	2.7	(74.5, 85.2)	*
CONNECTICUT	191	91.8	2.4	(87.1, 96.4)	*	390	85.2	2.1	(81.2, 89.2)	*
DELAWARE	260	84.0	2.5	(79.0, 88.9)	*	402	85.5	1.9	(81.7, 89.3)	*
DISTRICT OF CO	66	93.5	3.0	(87.6, 99.5)	*	205	87.1	2.5	(82.2, 92.0)	*
FLORIDA	298	82.4	2.4	(77.7, 87.2)	*	540	87.3	1.6	(84.2, 90.3)	*
GEORGIA	276	82.3	2.6	(77.1, 87.4)	*	442	88.5	1.6	(85.3, 91.6)	*
HAWAII	232	75.9	3.3	(69.5, 82.2)	*	380	78.3	2.7	(72.9, 83.6)	*
IDAHO	653	65.8	2.3	(61.2, 70.3)	*	871	66.9	2.0	(63.0, 70.9)	*
ILLINOIS	159	78.3	3.6	(71.3, 85.3)	*	260	77.8	2.9	(72.1, 83.4)	*
INDIANA	293	75.5	2.9	(69.9, 81.1)	*	399	76.1	2.3	(71.6, 80.6)	*
IOWA	397	69.1	2.6	(63.9, 74.3)	*	542	74.0	2.1	(69.9, 78.2)	*
KANSAS	161	72.1	3.8	(64.8, 79.5)	*	297	70.5	2.9	(64.9, 76.1)	*
KENTUCKY	326	82.2	2.3	(77.7, 86.7)	*	574	81.8	1.7	(78.4, 85.2)	*
LOUISIANA	176	80.7	3.4	(73.9, 87.4)	*	317	82.0	2.3	(77.4, 86.5)	*
MAINE	207	95.0	1.6	(91.9, 98.1)	*	264	95.6	1.5	(92.7, 98.4)	*
MARYLAND	488	85.7	2.1	(81.7, 89.8)	*	743	89.1	1.4	(86.2, 91.9)	*
MASSACHUSETTS	158	85.9	3.0	(80.1, 91.7)	*	282	84.3	2.6	(79.2, 89.4)	*
MICHIGAN	286	84.7	2.3	(80.2, 89.2)	*	483	81.9	1.9	(78.2, 85.6)	*
MINNESOTA	522	69.6	2.1	(65.4, 73.7)	*	693	68.6	2.0	(64.7, 72.4)	*
MISSISSIPPI	177	55.1	4.6	(46.2, 64.0)	*	331	70.8	2.7	(65.6, 76.1)	*
MISSOURI	169	71.8	4.0	(64.0, 79.6)	*	341	72.3	2.8	(66.8, 77.8)	*
MONTANA	191	67.9	3.5	(61.0, 74.8)	*	282	70.2	2.9	(64.5, 76.0)	*
NEBRASKA	266	67.8	3.4	(61.2, 74.4)	*	430	67.8	2.8	(62.4, 73.2)	*
NEVADA	311	84.8	3.6	(77.8, 91.8)	*	404	84.4	3.0	(78.4, 90.3)	*
NEW HAMPSHIRE	153	89.5	2.7	(84.2, 94.7)	*	257	87.9	2.2	(83.5, 92.3)	*
NEW JERSEY	263	83.3	3.1	(77.3, 89.2)	*	462	85.4	1.8	(81.8, 88.9)	*
NEW MEXICO	222	89.3	2.2	(85.0, 93.5)	*	373	88.1	2.0	(84.2, 91.9)	*
NEW YORK	294	87.5	2.4	(82.9, 92.1)	*	571	85.4	1.7	(82.2, 88.6)	*
NORTH CAROLINA	318	86.3	2.0	(82.3, 90.3)	*	618	89.9	1.3	(87.4, 92.4)	*
NORTH DAKOTA	176	56.9	3.9	(49.4, 64.5)	*	249	65.3	3.1	(59.3, 71.4)	*
OHIO	280	84.0	2.6	(78.8, 89.2)	*	504	86.3	1.8	(82.8, 89.8)	*
OKLAHOMA	153	77.9	3.6	(70.8, 84.9)	*	296	79.8	2.6	(74.6, 84.9)	*
OREGON	345	90.9	1.7	(87.6, 94.2)	*	550	91.3	1.6	(88.2, 94.4)	*
PENNSYLVANIA	343	82.2	2.3	(77.6, 86.7)	*	582	89.4	1.4	(86.7, 92.1)	*

RHODE ISLAND	167	86.2	2.9	(80.4, 91.9)	*	299	90.5	2.0	(86.6, 94.4)
SOUTH CAROLINA	174	89.6	2.5	(84.7, 94.6)	*	350	88.8	1.9	(85.2, 92.4)
SOUTH DAKOTA	249	51.0	3.4	(44.2, 57.7)	*	341	57.4	3.0	(51.6, 63.2)
TENNESSEE	232	81.7	2.8	(76.2, 87.1)	*	491	80.2	2.1	(76.0, 84.4)
TEXAS	300	77.0	2.8	(71.6, 82.4)	*	494	80.6	2.0	(76.7, 84.5)
UTAH	443	63.3	2.9	(57.7, 68.9)	*	595	60.1	2.8	(54.8, 65.5)
VERMONT	372	90.2	3.2	(83.8, 96.5)	*	527	94.5	1.0	(92.4, 96.5)
VIRGINIA	374	82.5	2.2	(78.3, 86.7)	*	589	83.1	2.1	(79.1, 87.1)
WASHINGTON	405	84.3	2.1	(80.2, 88.3)	*	601	87.8	1.5	(84.9, 90.6)
WEST VIRGINIA	221	80.8	2.8	(75.2, 86.3)	*	361	87.5	1.9	(83.7, 91.3)
WISCONSIN	273	72.4	3.8	(65.0, 79.9)	*	352	74.1	2.7	(68.8, 79.4)
WYOMING	310	66.6	3.0	(60.8, 72.3)	*	437	61.1	2.9	(55.5, 66.7)
PUERTO RICO	288	76.2	2.7	(70.9, 81.6)	*	428	70.5	2.5	(65.7, 75.3)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 82.2
 RANGE = 51.0-95.0 NUMBER OF PARTICIPANTS = 52
 MEDIAN = 81.9
 RANGE = 57.4-95.6

* OLDEST CHILD UNDER 16 ALWAYS USE SAFETY BELT

August 14, 1998

TABLE 16.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 USE OF CHILD SAFETY BELT*
 DENOMINATOR IS PERSONS WHO HAVE A CHILD BETWEEN THE AGE OF 5 AND 15
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES IN A CAR

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	70.3	75.7	5.4
ALASKA	82.7	80.0	-2.7
ARIZONA	80.0	93.3	13.3
ARKANSAS	75.4	78.5	3.1
CALIFORNIA	91.5	91.8	0.3
COLORADO	74.8	82.1	7.2
CONNECTICUT	83.4	88.1	4.7
DELAWARE	85.8	84.8	-1.0
DISTRICT OF CO	N/A	89.4	N/A
FLORIDA	83.0	85.1	2.1
GEORGIA	73.6	85.6	12.0
HAWAII	82.3	77.2	-5.1
IDAHO	65.8	66.3	0.5
ILLINOIS	74.5	78.0	3.5
INDIANA	70.0	75.8	5.8
IOWA	63.0	71.6	8.6
KANSAS	64.4	71.2	6.8
KENTUCKY	79.5	82.0	2.5
LOUISIANA	74.1	81.4	7.2
MAINE	94.9	95.3	0.4
MARYLAND	80.7	87.5	6.8
MASSACHUSETTS	83.0	84.9	2.0
MICHIGAN	77.4	83.1	5.7
MINNESOTA	69.7	69.0	-0.7
MISSISSIPPI	66.8	63.7	-3.1
MISSOURI	72.1	72.1	0.0
MONTANA	64.6	69.1	4.5
NEBRASKA	63.4	67.8	4.4
NEVADA	77.5	84.6	7.1
NEW HAMPSHIRE	88.1	88.6	0.5

NEW JERSEY	85.1	84.4	-0.8
NEW MEXICO	88.5	88.6	0.1
NEW YORK	81.2	86.3	5.1
NORTH CAROLINA	88.6	88.4	-0.2
NORTH DAKOTA	58.2	61.4	3.2
OHIO	73.8	85.3	11.5
OKLAHOMA	69.6	78.9	9.4
OREGON	90.6	91.1	0.5
PENNSYLVANIA	76.6	86.1	9.5
RHODE ISLAND	75.9	88.5	12.6
SOUTH CAROLINA	78.6	89.2	10.6
SOUTH DAKOTA	57.4	54.3	-3.1
TENNESSEE	80.5	80.9	0.4
TEXAS	73.2	78.9	5.7
UTAH	59.2	61.7	2.5
VERMONT	88.6	92.4	3.8
VIRGINIA	81.4	82.8	1.5
WASHINGTON	86.5	86.1	-0.5
WEST VIRGINIA	85.3	84.4	-0.9
WISCONSIN	67.0	73.3	6.3
WYOMING	57.2	63.7	6.5
PUERTO RICO	N/A	73.2	N/A

* OLDEST CHILD UNDER 16 ALWAYS USE SAFETY BELT
N/A - NOT AVAILABLE

August 14, 1998

TABLE 17.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY SEAT*
DENOMINATOR IS PERSONS WHO HAVE A CHILD UNDER THE AGE OF 5
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES IN CAR

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	148	93.9	2.0	(90.0, 97.7)
ALASKA	109	92.8	2.5	(87.9, 97.7)
ARIZONA	96	96.7	1.4	(94.0, 99.4)
ARKANSAS	121	85.9	5.0	(76.1, 95.6)
CALIFORNIA	307	93.3	1.5	(90.4, 96.3)
COLORADO	122	94.9	2.5	(90.1, 99.7)
CONNECTICUT	173	96.3	1.6	(93.2, 99.3)
DELAWARE	151	97.7	1.5	(94.8, 100.5)
DISTRICT OF CO	75	85.0	4.6	(76.0, 93.9)
FLORIDA	242	96.2	1.4	(93.5, 98.8)
GEORGIA	197	99.3	0.5	(98.2, 100.3)
HAWAII	170	90.6	2.5	(85.7, 95.5)
IDAHO	394	88.8	2.0	(84.8, 92.8)
ILLINOIS	126	95.7	1.8	(92.2, 99.2)
INDIANA	198	91.2	2.3	(86.7, 95.7)
IOWA	251	91.9	1.8	(88.4, 95.4)
KANSAS	119	95.2	1.9	(91.4, 98.9)
KENTUCKY	242	98.1	0.9	(96.2, 99.9)
LOUISIANA	111	93.8	2.2	(89.5, 98.1)
MAINE	112	99.6	0.4	(98.8, 100.4)
MARYLAND	312	97.1	1.1	(95.0, 99.3)
MASSACHUSETTS	144	94.7	2.7	(89.4, 99.9)

MICHIGAN	190	90.6	2.1	(86.5, 94.8)
MINNESOTA	304	93.9	1.5	(91.0, 96.8)
MISSISSIPPI	109	93.8	2.5	(88.8, 98.7)
MISSOURI	131	97.7	1.5	(94.8, 100.6)
MONTANA	134	92.0	2.4	(87.3, 96.8)
NEBRASKA	195	94.4	1.9	(90.7, 98.1)
NEVADA	179	94.7	1.9	(90.9, 98.4)
NEW HAMPSHIRE	109	97.6	1.7	(94.2, 100.9)
NEW JERSEY	196	96.0	1.4	(93.2, 98.7)
NEW MEXICO	119	93.8	2.5	(89.0, 98.6)
NEW YORK	247	94.6	1.4	(91.8, 97.4)
NORTH CAROLINA	272	93.5	1.8	(89.9, 97.0)
NORTH DAKOTA	121	96.8	1.7	(93.5, 100.0)
OHIO	193	94.3	2.7	(89.0, 99.6)
OKLAHOMA	113	93.3	2.7	(87.9, 98.6)
OREGON	224	93.0	2.3	(88.5, 97.6)
PENNSYLVANIA	237	94.9	1.7	(91.5, 98.3)
RHODE ISLAND	113	95.8	2.1	(91.8, 99.8)
SOUTH CAROLINA	156	97.4	1.5	(94.4, 100.5)
SOUTH DAKOTA	131	91.0	3.1	(84.8, 97.1)
TENNESSEE	208	97.5	1.5	(94.6, 100.4)
TEXAS	214	85.9	2.6	(80.9, 90.9)
UTAH	286	90.7	1.9	(87.0, 94.4)
VERMONT	202	98.3	0.8	(96.8, 99.9)
VIRGINIA	267	96.1	1.4	(93.3, 99.0)
WASHINGTON	269	87.3	3.1	(81.3, 93.2)
WEST VIRGINIA	144	96.3	1.6	(93.2, 99.4)
WISCONSIN	171	93.6	1.8	(90.0, 97.2)
WYOMING	146	91.5	2.4	(86.9, 96.2)
PUERTO RICO	214	92.3	1.8	(88.8, 95.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 94.4 RANGE = 85.0-99.6

APPROXIMATES YEAR 2000 OBJ. 9.12A

* OLDEST CHILD UNDER 5 ALWAYS USE SAFETY SEAT

August 14, 1998

TABLE 17.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR USE OF CHILD SAFETY SEAT* -BY SEX
DENOMINATOR IS PERSONS WHO HAVE A CHILD UNDER THE AGE OF 5
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES IN CAR

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD	95% CONFIDENCE	SAMPLE SIZE	PERCENT	STANDARD	95% CONFIDENCE	
			ERROR	INTERVAL			ERROR	INTERVAL	
ALABAMA	54	94.3	3.1	(88.1, 100.4)	*	94	93.6	2.5	(88.7, 98.5)
ALASKA	43	94.9	2.5	(90.0, 99.9)	*	66	91.1	3.9	(83.4, 98.8)
ARIZONA	40	94.4	2.9	(88.7, 100.0)	*	56	98.3	1.2	(95.9, 100.8)
ARKANSAS	34	76.0	10.5	(55.3, 96.6)	*	87	92.5	3.2	(86.3, 98.7)
CALIFORNIA	134	94.2	2.0	(90.2, 98.2)	*	173	92.3	2.2	(88.0, 96.6)
COLORADO	57	91.3	4.6	(82.2, 100.3)	*	65	98.5	1.5	(95.6, 101.4)
CONNECTICUT	67	95.1	2.8	(89.7, 100.5)	*	106	97.4	1.6	(94.3, 100.4)
DELAWARE	44	100	0.0	(100.0, 100.0)	*	107	96.5	2.2	(92.2, 100.8)
DISTRICT OF CO	16	90.8	8.7	(73.8, 107.8)	*	59	82.4	5.3	(72.0, 92.8)
FLORIDA	79	96.3	2.1	(92.1, 100.4)	*	163	96.1	1.8	(92.7, 99.6)
GEORGIA	77	100	0.0	(100.0, 100.0)	*	120	98.6	1.0	(96.7, 100.5)
HAWAII	69	87.7	4.1	(79.7, 95.6)	*	101	93.1	3.0	(87.2, 99.0)
IDAHO	163	90.4	2.4	(85.8, 95.0)	*	231	87.5	3.2	(81.3, 93.6)

ILLINOIS	49	97.2	2.1	(93.1, 101.3)	*	77	94.4	2.8	(89.0, 99.9)
INDIANA	80	94.4	2.5	(89.6, 99.3)	*	118	88.6	3.5	(81.7, 95.4)
IOWA	101	93.6	2.5	(88.7, 98.4)	*	150	90.6	2.5	(85.7, 95.5)
KANSAS	45	94.9	3.0	(89.0, 100.8)	*	74	95.4	2.4	(90.7, 100.1)
KENTUCKY	80	97.4	1.8	(93.9, 101.0)	*	162	98.6	0.8	(97.1, 100.1)
LOUISIANA	32	93.7	3.7	(86.4, 100.9)	*	79	93.9	2.7	(88.5, 99.2)
MAINE	40	100	0.0	(100.0, 100.0)	*	72	99.4	0.7	(98.1, 100.6)
MARYLAND	119	96.6	1.8	(93.1, 100.1)	*	193	97.5	1.4	(94.8, 100.2)
MASSACHUSETTS	51	92.9	5.3	(82.6, 103.2)	*	93	96.0	2.5	(91.1, 100.8)
MICHIGAN	83	94.8	2.4	(90.2, 99.5)	*	107	85.9	3.5	(79.0, 92.8)
MINNESOTA	150	94.4	1.9	(90.7, 98.2)	*	154	93.4	2.0	(89.4, 97.3)
MISSISSIPPI	45	91.7	4.3	(83.3, 100.1)	*	64	95.9	2.5	(90.9, 100.8)
MISSOURI	41	96.0	3.1	(89.9, 102.1)	*	90	99.2	0.6	(98.0, 100.4)
MONTANA	54	88.7	4.4	(80.2, 97.3)	*	80	94.7	2.6	(89.7, 99.7)
NEBRASKA	61	95.2	3.3	(88.6, 101.7)	*	134	93.8	2.2	(89.5, 98.1)
NEVADA	78	88.7	4.6	(79.8, 97.6)	*	101	98.4	1.3	(95.9, 100.9)
NEW HAMPSHIRE	33	96.3	3.6	(89.3, 103.4)	*	76	98.3	1.7	(94.9, 101.6)
NEW JERSEY	80	98.1	1.4	(95.4, 100.8)	*	116	94.1	2.3	(89.5, 98.6)
NEW MEXICO	43	95.7	3.0	(89.9, 101.6)	*	76	92.3	3.7	(85.1, 99.5)
NEW YORK	98	97.4	1.5	(94.5, 100.3)	*	149	91.9	2.4	(87.2, 96.6)
NORTH CAROLINA	99	92.5	2.8	(87.1, 97.9)	*	173	94.2	2.4	(89.5, 99.0)
NORTH DAKOTA	50	100	0.0	(100.0, 100.0)	*	71	94.4	2.9	(88.7, 100.0)
OHIO	71	89.8	5.4	(79.1, 100.4)	*	122	98.2	1.5	(95.2, 101.1)
OKLAHOMA	38	92.9	4.6	(83.9, 102.0)	*	75	93.5	3.2	(87.2, 99.8)
OREGON	88	94.8	2.4	(90.2, 99.4)	*	136	91.8	3.6	(84.8, 98.8)
PENNSYLVANIA	86	94.8	3.2	(88.5, 101.1)	*	151	95.0	1.9	(91.3, 98.7)
RHODE ISLAND	42	93.3	3.8	(85.8, 100.7)	*	71	98.0	2.0	(94.0, 101.9)
SOUTH CAROLINA	51	100	0.0	(100.0, 100.0)	*	105	95.5	2.7	(90.2, 100.7)
SOUTH DAKOTA	52	93.7	3.6	(86.5, 100.8)	*	79	89.0	4.7	(79.9, 98.2)
TENNESSEE	66	95.9	3.4	(89.3, 102.5)	*	142	98.6	0.9	(96.9, 100.3)
TEXAS	78	88.2	3.9	(80.6, 95.7)	*	136	84.2	3.4	(77.5, 90.8)
UTAH	134	90.0	2.9	(84.4, 95.6)	*	152	91.4	2.5	(86.6, 96.3)
VERMONT	86	99.4	0.6	(98.2, 100.6)	*	116	97.5	1.3	(94.8, 100.1)
VIRGINIA	112	95.4	2.5	(90.6, 100.2)	*	155	96.8	1.6	(93.7, 99.9)
WASHINGTON	115	89.2	3.1	(83.2, 95.2)	*	154	85.5	5.0	(75.7, 95.3)
WEST VIRGINIA	69	95.8	2.4	(91.1, 100.5)	*	75	97.0	2.0	(93.0, 100.9)
WISCONSIN	76	94.3	2.7	(89.0, 99.6)	*	95	92.8	2.5	(88.0, 97.6)
WYOMING	61	89.3	4.2	(81.0, 97.5)	*	85	93.6	2.4	(88.9, 98.3)
PUERTO RICO	85	92.5	2.8	(87.0, 97.9)	*	129	92.2	2.3	(87.7, 96.6)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 94.4
 RANGE = 76.0- 100

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 94.3
 RANGE = 82.4-99.4

* OLDEST CHILD UNDER 5 ALWAYS USE SAFETY SEAT

August 14, 1998

TABLE 17.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 USE OF CHILD SAFETY SEAT*
 DENOMINATOR IS PERSONS WHO HAVE A CHILD UNDER THE AGE OF 5
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES IN CAR

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	88.8	93.9	5.1
ALASKA	96.3	92.8	-3.5
ARIZONA	90.7	96.7	6.1
ARKANSAS	98.3	85.9	-12.4

CALIFORNIA	89.0	93.3	4.4
COLORADO	92.0	94.9	2.9
CONNECTICUT	90.9	96.3	5.3
DELAWARE	97.6	97.7	0.1
DISTRICT OF CO	N/A	85.0	N/A
FLORIDA	93.2	96.2	3.0
GEORGIA	95.7	99.3	3.6
HAWAII	97.6	90.6	-7.0
IDAHO	86.6	88.8	2.2
ILLINOIS	88.3	95.7	7.4
INDIANA	94.8	91.2	-3.6
IOWA	88.6	91.9	3.4
KANSAS	93.6	95.2	1.5
KENTUCKY	95.7	98.1	2.3
LOUISIANA	87.6	93.8	6.2
MAINE	98.5	99.6	1.1
MARYLAND	96.2	97.1	0.9
MASSACHUSETTS	94.9	94.7	-0.3
MICHIGAN	92.0	90.6	-1.4
MINNESOTA	94.1	93.9	-0.2
MISSISSIPPI	93.8	93.8	0.0
MISSOURI	87.5	97.7	10.2
MONTANA	82.9	92.0	9.1
NEBRASKA	92.6	94.4	1.8
NEVADA	92.0	94.7	2.6
NEW HAMPSHIRE	95.5	97.6	2.1
NEW JERSEY	99.2	96.0	-3.2
NEW MEXICO	92.6	93.8	1.2
NEW YORK	90.8	94.6	3.8
NORTH CAROLINA	98.6	93.5	-5.2
NORTH DAKOTA	94.9	96.8	1.9
OHIO	98.0	94.3	-3.7
OKLAHOMA	97.3	93.3	-4.0
OREGON	88.4	93.0	4.6
PENNSYLVANIA	87.2	94.9	7.7
RHODE ISLAND	97.2	95.8	-1.4
SOUTH CAROLINA	96.5	97.4	1.0
SOUTH DAKOTA	91.0	91.0	-0.0
TENNESSEE	97.1	97.5	0.4
TEXAS	93.8	85.9	-7.9
UTAH	86.5	90.7	4.3
VERMONT	97.9	98.3	0.5
VIRGINIA	92.9	96.1	3.2
WASHINGTON	94.9	87.3	-7.7
WEST VIRGINIA	96.8	96.3	-0.5
WISCONSIN	86.8	93.6	6.8
WYOMING	86.9	91.5	4.6
PUERTO RICO	N/A	92.3	N/A

* OLDEST CHILD UNDER 5 ALWAYS USE SAFETY SEAT

N/A - NOT AVAILABLE

August 14, 1998

TABLE 18.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHILD BICYCLE HELMET USE*
DENOMINATOR IS RESPONDENTS WITH CHILDREN UNDER 16
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES A BIKE

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	438	34.1	2.6	(29.0, 39.3)
ALASKA	503	34.5	3.4	(27.8, 41.2)
ARIZONA	370	39.7	3.5	(32.8, 46.5)
ARKANSAS	375	22.1	2.4	(17.5, 26.8)
CALIFORNIA	772	61.1	2.0	(57.2, 65.1)
COLORADO	425	37.4	2.6	(32.3, 42.5)
CONNECTICUT	493	59.1	2.7	(53.9, 64.3)
DELAWARE	546	59.9	2.4	(55.3, 64.5)
DISTRICT OF CO	218	40.1	4.1	(32.1, 48.1)
FLORIDA	625	47.1	2.3	(42.6, 51.6)
GEORGIA	592	49.7	2.3	(45.1, 54.3)
HAWAII	416	31.2	2.6	(26.1, 36.3)
IDAHO	1325	27.5	1.7	(24.2, 30.8)
ILLINOIS	363	25.8	2.6	(20.8, 30.8)
INDIANA	609	16.3	1.6	(13.1, 19.4)
IOWA	834	20.4	1.5	(17.4, 23.3)
KANSAS	389	23.3	2.3	(18.8, 27.8)
KENTUCKY	673	23.8	1.9	(20.2, 27.5)
LOUISIANA	388	18.3	2.3	(13.8, 22.7)
MAINE	414	49.6	2.7	(44.4, 54.8)
MARYLAND	1044	54.4	2.0	(50.5, 58.4)
MASSACHUSETTS	383	60.7	2.8	(55.3, 66.1)
MICHIGAN	712	20.8	1.6	(17.6, 24.0)
MINNESOTA	1127	20.7	1.3	(18.1, 23.3)
MISSISSIPPI	352	18.9	2.4	(14.2, 23.6)
MISSOURI	409	21.4	2.4	(16.6, 26.2)
MONTANA	424	32.4	2.5	(27.6, 37.3)
NEBRASKA	586	20.2	2.3	(15.7, 24.7)
NEVADA	594	32.2	3.8	(24.9, 39.6)
NEW HAMPSHIRE	376	61.8	2.8	(56.4, 67.2)
NEW JERSEY	623	63.8	2.4	(59.1, 68.4)
NEW MEXICO	466	31.1	2.4	(26.4, 35.8)
NEW YORK	720	69.3	1.9	(65.7, 73.0)
NORTH CAROLINA	801	31.7	1.8	(28.1, 35.2)
NORTH DAKOTA	391	15.4	1.9	(11.6, 19.1)
OHIO	656	32.4	2.3	(28.0, 36.8)
OKLAHOMA	333	25.4	2.5	(20.5, 30.3)
OREGON	783	66.8	1.9	(63.0, 70.5)
PENNSYLVANIA	792	59.2	2.0	(55.4, 63.0)
RHODE ISLAND	408	56.6	2.7	(51.3, 61.9)
SOUTH CAROLINA	450	23.2	2.1	(19.1, 27.3)
SOUTH DAKOTA	507	13.5	1.7	(10.1, 16.8)
TENNESSEE	636	32.2	2.1	(28.1, 36.2)
TEXAS	548	22.5	2.0	(18.6, 26.4)
UTAH	880	19.0	1.6	(15.8, 22.1)
VERMONT	833	62.9	2.1	(58.7, 67.0)
VIRGINIA	783	41.0	2.6	(35.9, 46.1)
WASHINGTON	889	52.4	2.0	(48.5, 56.3)
WEST VIRGINIA	465	55.1	2.6	(50.1, 60.1)
WISCONSIN	588	22.0	1.9	(18.3, 25.7)
WYOMING	675	24.8	2.0	(20.9, 28.7)
PUERTO RICO	502	12.5	1.5	(9.5, 15.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 32.2 RANGE = 12.5-69.3

* OLDEST CHILD UNDER 16 ALWAYS WEARS A BICYCLE HELMET

August 14, 1998

TABLE 18.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHILD BICYCLE HELMET USE* -BY SEX
 DENOMINATOR IS RESPONDENTS WITH CHILDREN UNDER 16
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES A BIKE

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	149	34.3	4.4	(25.6, 42.9)	*	289	34.0	(28.0, 40.0)
ALASKA	231	31.9	4.8	(22.4, 41.4)	*	272	37.6	(28.3, 46.9)
ARIZONA	163	39.7	5.3	(29.4, 50.0)	*	207	39.6	(31.0, 48.3)
ARKANSAS	144	22.5	3.7	(15.3, 29.6)	*	231	21.8	(15.9, 27.7)
CALIFORNIA	312	58.7	3.2	(52.5, 64.9)	*	460	63.5	(58.4, 68.5)
COLORADO	182	35.2	3.9	(27.6, 42.7)	*	243	39.6	(32.7, 46.5)
CONNECTICUT	168	55.4	4.5	(46.6, 64.2)	*	325	62.2	(56.2, 68.1)
DELAWARE	230	58.2	3.7	(50.9, 65.5)	*	316	61.7	(55.7, 67.7)
DISTRICT OF CO	48	50.5	8.2	(34.5, 66.5)	*	170	35.3	(26.6, 43.9)
FLORIDA	224	46.7	3.6	(39.5, 53.8)	*	401	47.5	(41.9, 53.0)
GEORGIA	231	47.7	3.5	(40.8, 54.6)	*	361	51.4	(45.3, 57.6)
HAWAII	163	33.3	4.0	(25.4, 41.2)	*	253	29.3	(22.6, 35.9)
IDAHO	582	25.4	2.3	(20.9, 29.8)	*	743	29.8	(25.0, 34.6)
ILLINOIS	141	23.9	4.0	(15.9, 31.8)	*	222	27.5	(21.1, 33.8)
INDIANA	263	16.7	2.5	(11.9, 21.6)	*	346	15.8	(11.4, 20.2)
IOWA	361	18.2	2.2	(14.0, 22.4)	*	473	22.6	(18.5, 26.6)
KANSAS	137	21.2	3.7	(14.0, 28.4)	*	252	25.0	(19.4, 30.6)
KENTUCKY	248	23.4	2.9	(17.8, 29.0)	*	425	24.2	(19.7, 28.6)
LOUISIANA	143	17.8	3.5	(10.9, 24.7)	*	245	18.8	(13.2, 24.3)
MAINE	184	50.2	3.9	(42.5, 58.0)	*	230	49.1	(42.3, 55.8)
MARYLAND	411	51.4	3.1	(45.3, 57.5)	*	633	56.9	(52.0, 61.9)
MASSACHUSETTS	140	60.9	4.4	(52.2, 69.6)	*	243	60.5	(53.7, 67.3)
MICHIGAN	267	21.6	2.6	(16.5, 26.7)	*	445	20.2	(16.1, 24.2)
MINNESOTA	490	19.1	1.9	(15.4, 22.7)	*	637	22.1	(18.5, 25.7)
MISSISSIPPI	134	16.0	3.4	(9.4, 22.7)	*	218	21.6	(15.4, 27.7)
MISSOURI	151	18.4	3.7	(11.1, 25.7)	*	258	24.0	(18.0, 30.0)
MONTANA	179	33.0	3.8	(25.6, 40.4)	*	245	31.8	(25.7, 37.9)
NEBRASKA	231	20.1	3.0	(14.3, 25.9)	*	355	20.3	(13.5, 27.1)
NEVADA	266	34.6	6.1	(22.6, 46.5)	*	328	29.8	(21.7, 37.8)
NEW HAMPSHIRE	143	62.2	4.1	(54.1, 70.3)	*	233	61.5	(54.5, 68.4)
NEW JERSEY	228	63.2	3.8	(55.9, 70.6)	*	395	64.2	(58.5, 69.9)
NEW MEXICO	173	31.5	3.8	(24.1, 39.0)	*	293	30.7	(25.0, 36.5)
NEW YORK	248	69.1	3.1	(63.1, 75.2)	*	472	69.5	(65.0, 73.9)
NORTH CAROLINA	281	31.2	2.9	(25.5, 36.9)	*	520	32.0	(27.7, 36.4)
NORTH DAKOTA	171	10.6	2.5	(5.7, 15.5)	*	220	19.8	(14.3, 25.3)
OHIO	251	28.0	3.3	(21.4, 34.5)	*	405	36.5	(30.5, 42.6)
OKLAHOMA	119	24.2	4.1	(16.2, 32.1)	*	214	26.5	(20.3, 32.7)
OREGON	315	65.8	3.0	(59.9, 71.7)	*	468	67.6	(62.8, 72.4)
PENNSYLVANIA	305	58.1	3.1	(52.1, 64.2)	*	487	60.1	(55.3, 65.0)
RHODE ISLAND	151	60.3	4.3	(52.0, 68.6)	*	257	53.4	(46.7, 60.1)
SOUTH CAROLINA	151	22.6	3.4	(15.8, 29.3)	*	299	23.7	(18.6, 28.8)
SOUTH DAKOTA	210	12.8	2.7	(7.5, 18.1)	*	297	14.0	(9.9, 18.1)
TENNESSEE	206	28.7	3.5	(21.8, 35.5)	*	430	35.1	(30.4, 39.8)
TEXAS	213	23.8	3.3	(17.4, 30.2)	*	335	21.3	(16.7, 25.9)
UTAH	382	18.6	2.4	(13.9, 23.3)	*	498	19.3	(15.0, 23.7)
VERMONT	347	57.5	3.4	(50.8, 64.2)	*	486	67.9	(63.2, 72.5)
VIRGINIA	309	43.5	4.1	(35.4, 51.5)	*	474	38.9	(32.4, 45.4)
WASHINGTON	364	51.8	3.2	(45.7, 58.0)	*	525	53.0	(48.2, 57.8)
WEST VIRGINIA	179	53.5	4.0	(45.6, 61.4)	*	286	56.6	(50.1, 63.0)
WISCONSIN	256	21.4	2.9	(15.8, 27.0)	*	332	22.4	(17.5, 27.3)
WYOMING	281	23.2	2.7	(17.8, 28.5)	*	394	26.4	(20.8, 32.0)
PUERTO RICO	192	10.7	2.2	(6.4, 15.1)	*	310	14.1	(9.9, 18.2)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 31.7
 RANGE = 10.6-69.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 31.9
 RANGE = 14.0-69.5

* OLDEST CHILD UNDER 16 ALWAYS WEARS A BICYCLE HELMET

August 14, 1998

TABLE 18.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 CHILD BICYCLE HELMET USE*
 DENOMINATOR IS RESPONDENTS WITH CHILDREN UNDER 16
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED, AND NEVER RIDES A BIKE

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	19.8	34.1	14.3
ALASKA	33.1	34.5	1.4
ARIZONA	23.5	39.7	16.2
ARKANSAS	17.8	22.1	4.3
CALIFORNIA	55.6	61.1	5.5
COLORADO	28.3	37.4	9.1
CONNECTICUT	52.5	59.1	6.6
DELAWARE	33.0	59.9	26.9
DISTRICT OF CO	N/A	40.1	N/A
FLORIDA	20.5	47.1	26.6
GEORGIA	41.1	49.7	8.6
HAWAII	20.3	31.2	10.9
IDAHO	24.0	27.5	3.5
ILLINOIS	20.2	25.8	5.7
INDIANA	13.1	16.3	3.1
IOWA	13.0	20.4	7.3
KANSAS	18.0	23.3	5.3
KENTUCKY	18.6	23.8	5.3
LOUISIANA	12.8	18.3	5.5
MAINE	43.6	49.6	6.0
MARYLAND	38.1	54.4	16.4
MASSACHUSETTS	47.5	60.7	13.2
MICHIGAN	16.9	20.8	3.9
MINNESOTA	17.8	20.7	2.9
MISSISSIPPI	9.3	18.9	9.6
MISSOURI	21.0	21.4	0.4
MONTANA	22.7	32.4	9.7
NEBRASKA	11.8	20.2	8.4
NEVADA	23.7	32.2	8.5
NEW HAMPSHIRE	49.7	61.8	12.1
NEW JERSEY	58.7	63.8	5.0
NEW MEXICO	24.5	31.1	6.6
NEW YORK	60.4	69.3	8.9
NORTH CAROLINA	24.3	31.7	7.4
NORTH DAKOTA	12.1	15.4	3.3
OHIO	23.7	32.4	8.7
OKLAHOMA	21.4	25.4	4.0
OREGON	62.8	66.8	3.9
PENNSYLVANIA	42.2	59.2	17.0
RHODE ISLAND	32.7	56.6	24.0
SOUTH CAROLINA	18.2	23.2	5.0
SOUTH DAKOTA	13.0	13.5	0.5
TENNESSEE	27.1	32.2	5.0
TEXAS	17.1	22.5	5.4
UTAH	15.9	19.0	3.1
VERMONT	46.4	62.9	16.5
VIRGINIA	33.1	41.0	7.9
WASHINGTON	50.4	52.4	2.0
WEST VIRGINIA	20.9	55.1	34.2
WISCONSIN	15.7	22.0	6.2
WYOMING	19.3	24.8	5.5

PUERTO RICO N/A 12.5 N/A

* OLDEST CHILD UNDER 16 ALWAYS WEARS A BICYCLE HELMET

N/A - NOT AVAILABLE

August 14, 1998

TABLE 19.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT DRINKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2171	36.8	1.2	(34.4, 39.2)
ALASKA	1542	55.2	1.9	(51.5, 58.9)
ARIZONA	1904	41.1	1.6	(37.9, 44.3)
ARKANSAS	1793	30.6	1.3	(28.0, 33.2)
CALIFORNIA	4063	61.4	0.9	(59.7, 63.1)
COLORADO	1822	62.0	1.3	(59.5, 64.6)
CONNECTICUT	2241	62.1	1.3	(59.6, 64.6)
DELAWARE	2556	54.2	1.2	(51.9, 56.5)
DISTRICT OF CO	1503	43.7	1.6	(40.5, 46.8)
FLORIDA	3490	54.8	1.0	(52.8, 56.7)
GEORGIA	2308	44.6	1.2	(42.2, 47.0)
HAWAII	2167	51.1	1.3	(48.6, 53.6)
IDAHO	4916	47.6	0.9	(45.7, 49.4)
ILLINOIS	1440	54.7	1.5	(51.8, 57.6)
INDIANA	2366	48.5	1.2	(46.1, 50.8)
IOWA	3595	53.9	1.0	(51.9, 55.8)
KANSAS	2000	41.3	1.2	(38.9, 43.7)
KENTUCKY	3605	33.7	1.0	(31.8, 35.6)
LOUISIANA	1653	50.6	1.5	(47.7, 53.4)
MAINE	1697	55.4	1.4	(52.7, 58.0)
MARYLAND	4567	44.4	1.0	(42.4, 46.4)
MASSACHUSETTS	1765	64.8	1.3	(62.3, 67.4)
MICHIGAN	2563	57.4	1.1	(55.3, 59.5)
MINNESOTA	4813	53.5	0.8	(51.9, 55.1)
MISSISSIPPI	1598	35.1	1.4	(32.4, 37.9)
MISSOURI	1849	49.2	1.4	(46.4, 51.9)
MONTANA	1796	57.2	1.3	(54.6, 59.7)
NEBRASKA	2689	58.0	1.2	(55.7, 60.4)
NEVADA	2505	62.9	1.9	(59.1, 66.6)
NEW HAMPSHIRE	1512	61.3	1.4	(58.5, 64.0)
NEW JERSEY	2656	58.3	1.1	(56.1, 60.5)
NEW MEXICO	1809	53.6	1.4	(50.9, 56.3)
NEW YORK	3399	53.5	1.0	(51.6, 55.5)
NORTH CAROLINA	3623	37.6	1.0	(35.7, 39.5)
NORTH DAKOTA	1800	53.7	1.3	(51.2, 56.3)
OHIO	3165	33.7	1.1	(31.5, 35.9)
OKLAHOMA	1885	32.8	1.3	(30.1, 35.4)
OREGON	3318	56.7	1.0	(54.7, 58.7)
PENNSYLVANIA	3588	49.3	1.0	(47.3, 51.2)
RHODE ISLAND	1841	59.4	1.3	(56.9, 61.9)
SOUTH CAROLINA	2154	39.7	1.3	(37.2, 42.2)
SOUTH DAKOTA	2197	55.5	1.2	(53.2, 57.8)
TENNESSEE	2973	28.7	1.0	(26.7, 30.7)
TEXAS	2497	49.4	1.1	(47.3, 51.6)

UTAH	2861	28.3	1.1	(26.1, 30.4)
VERMONT	3190	60.2	1.0	(58.3, 62.2)
VIRGINIA	3517	54.9	1.2	(52.4, 57.3)
WASHINGTON	3594	60.4	1.0	(58.5, 62.3)
WEST VIRGINIA	2435	31.0	1.1	(28.9, 33.2)
WISCONSIN	2242	70.2	1.2	(67.9, 72.5)
WYOMING	2412	52.6	1.3	(50.2, 55.1)
PUERTO RICO	2257	26.7	1.1	(24.6, 28.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 53.5 RANGE = 26.7-70.2

* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH

August 14, 1998

TABLE 19.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT DRINKERS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-MALES-				-FEMALES-				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	811	46.1	2.0	(42.2, 50.0)	*	1360	28.6	1.4	(25.8, 31.3)
ALASKA	715	61.9	2.7	(56.6, 67.1)	*	827	47.9	2.6	(42.9, 53.0)
ARIZONA	891	45.8	2.2	(41.5, 50.1)	*	1013	36.6	2.1	(32.5, 40.8)
ARKANSAS	676	39.3	2.2	(34.9, 43.7)	*	1117	22.8	1.5	(20.0, 25.7)
CALIFORNIA	1759	69.8	1.3	(67.3, 72.2)	*	2304	53.2	1.2	(50.9, 55.5)
COLORADO	815	71.0	1.8	(67.5, 74.4)	*	1007	53.6	1.8	(50.0, 57.1)
CONNECTICUT	885	69.4	1.9	(65.6, 73.1)	*	1356	55.5	1.6	(52.4, 58.7)
DELAWARE	1013	65.2	1.7	(61.9, 68.5)	*	1543	44.2	1.5	(41.4, 47.1)
DISTRICT OF CO	572	49.1	2.4	(44.4, 53.9)	*	931	39.1	1.9	(35.2, 42.9)
FLORIDA	1437	62.0	1.5	(59.1, 64.9)	*	2053	48.1	1.2	(45.7, 50.5)
GEORGIA	962	52.4	1.9	(48.7, 56.0)	*	1346	37.5	1.5	(34.5, 40.6)
HAWAII	948	63.5	1.8	(59.9, 67.1)	*	1219	38.5	1.6	(35.3, 41.7)
IDAHO	2055	55.5	1.4	(52.8, 58.2)	*	2861	40.0	1.2	(37.6, 42.4)
ILLINOIS	576	63.2	2.2	(58.8, 67.5)	*	864	47.2	1.9	(43.6, 50.8)
INDIANA	1064	56.9	1.7	(53.5, 60.2)	*	1302	40.9	1.6	(37.8, 44.0)
IOWA	1476	62.6	1.5	(59.7, 65.4)	*	2119	46.0	1.3	(43.6, 48.5)
KANSAS	785	50.9	2.0	(47.0, 54.7)	*	1215	32.3	1.4	(29.5, 35.1)
KENTUCKY	1320	43.0	1.6	(39.9, 46.1)	*	2285	25.3	1.1	(23.2, 27.4)
LOUISIANA	629	61.0	2.2	(56.7, 65.3)	*	1024	41.2	1.7	(37.8, 44.6)
MAINE	751	64.2	2.0	(60.3, 68.1)	*	946	47.3	1.8	(43.7, 50.8)
MARYLAND	1955	51.8	1.5	(48.8, 54.8)	*	2612	37.6	1.3	(35.1, 40.1)
MASSACHUSETTS	720	71.8	1.9	(68.1, 75.5)	*	1045	58.6	1.7	(55.2, 62.1)
MICHIGAN	1069	67.5	1.6	(64.4, 70.5)	*	1494	48.2	1.4	(45.4, 51.0)
MINNESOTA	2202	58.7	1.2	(56.4, 61.0)	*	2611	48.6	1.1	(46.5, 50.8)
MISSISSIPPI	592	47.1	2.3	(42.6, 51.5)	*	1006	24.6	1.5	(21.6, 27.6)
MISSOURI	698	57.9	2.2	(53.7, 62.1)	*	1151	41.4	1.8	(37.9, 44.8)
MONTANA	757	66.4	1.9	(62.8, 70.1)	*	1039	48.4	1.7	(45.0, 51.7)
NEBRASKA	1043	66.3	1.8	(62.8, 69.7)	*	1646	50.5	1.6	(47.3, 53.6)
NEVADA	1164	70.8	2.7	(65.4, 76.1)	*	1341	54.9	2.5	(50.0, 59.8)
NEW HAMPSHIRE	603	66.6	2.2	(62.2, 70.9)	*	909	56.3	1.8	(52.7, 59.9)
NEW JERSEY	1060	65.6	1.7	(62.2, 68.9)	*	1596	51.7	1.5	(48.8, 54.6)
NEW MEXICO	752	64.0	1.9	(60.2, 67.8)	*	1057	43.8	1.7	(40.4, 47.2)
NEW YORK	1353	61.7	1.5	(58.7, 64.7)	*	2046	46.3	1.3	(43.8, 48.7)
NORTH CAROLINA	1436	47.0	1.5	(44.0, 50.0)	*	2187	29.1	1.1	(26.8, 31.3)
NORTH DAKOTA	799	64.4	1.9	(60.7, 68.0)	*	1001	43.5	1.7	(40.2, 46.8)
OHIO	1230	40.9	1.8	(37.3, 44.4)	*	1935	27.3	1.3	(24.7, 29.8)

OKLAHOMA	830	38.6	2.1	(34.4, 42.7)	*	1055	27.4	1.6	(24.2, 30.6)
OREGON	1400	64.2	1.5	(61.2, 67.1)	*	1918	49.7	1.3	(47.0, 52.3)
PENNSYLVANIA	1440	58.2	1.5	(55.2, 61.1)	*	2148	41.4	1.3	(38.9, 43.8)
RHODE ISLAND	756	66.0	1.9	(62.2, 69.7)	*	1085	53.6	1.7	(50.2, 57.0)
SOUTH CAROLINA	821	51.4	2.0	(47.5, 55.2)	*	1333	29.2	1.4	(26.4, 32.0)
SOUTH DAKOTA	956	63.8	1.8	(60.3, 67.2)	*	1241	47.7	1.6	(44.5, 50.9)
TENNESSEE	1043	37.8	1.7	(34.5, 41.2)	*	1930	20.5	1.0	(18.5, 22.5)
TEXAS	1023	59.4	1.7	(56.1, 62.7)	*	1474	40.1	1.4	(37.4, 42.8)
UTAH	1277	32.6	1.7	(29.3, 35.8)	*	1584	24.2	1.4	(21.5, 26.9)
VERMONT	1395	67.4	1.4	(64.6, 70.2)	*	1795	53.6	1.4	(50.9, 56.2)
VIRGINIA	1415	62.5	1.8	(59.0, 66.1)	*	2102	47.7	1.7	(44.4, 51.0)
WASHINGTON	1546	67.8	1.5	(64.9, 70.6)	*	2048	53.3	1.3	(50.7, 55.9)
WEST VIRGINIA	966	40.7	1.7	(37.4, 44.1)	*	1469	22.4	1.3	(19.8, 24.9)
WISCONSIN	991	80.0	1.6	(76.8, 83.1)	*	1251	61.2	1.6	(58.0, 64.3)
WYOMING	1020	63.0	1.8	(59.5, 66.6)	*	1392	42.5	1.6	(39.4, 45.6)
PUERTO RICO	934	43.7	1.8	(40.2, 47.2)	*	1323	11.6	1.0	(9.7, 13.5)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 61.9
 RANGE = 32.6-80.0

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 43.7
 RANGE = 11.6-61.2

* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH

August 14, 1998

TABLE 19.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 CURRENT DRINKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	40.3	36.8	-3.5
ALASKA	58.8	55.2	-3.6
ARIZONA	49.4	41.1	-8.3
ARKANSAS	32.7	30.6	-2.0
CALIFORNIA	61.6	61.4	-0.2
COLORADO	62.5	62.0	-0.4
CONNECTICUT	65.1	62.1	-2.9
DELAWARE	53.2	54.2	1.1
DISTRICT OF CO	N/A	43.7	N/A
FLORIDA	54.9	54.8	-0.1
GEORGIA	40.3	44.6	4.3
HAWAII	41.7	51.1	9.4
IDAHO	47.8	47.6	-0.2
ILLINOIS	52.0	54.7	2.7
INDIANA	46.8	48.5	1.6
IOWA	57.2	53.9	-3.3
KANSAS	49.6	41.3	-8.3
KENTUCKY	34.5	33.7	-0.8
LOUISIANA	48.2	50.6	2.4
MAINE	49.7	55.4	5.7
MARYLAND	45.1	44.4	-0.7
MASSACHUSETTS	67.0	64.8	-2.2
MICHIGAN	58.2	57.4	-0.8
MINNESOTA	59.6	53.5	-6.1
MISSISSIPPI	36.5	35.1	-1.4
MISSOURI	52.3	49.2	-3.1

MONTANA	56.0	57.2	1.1
NEBRASKA	55.4	58.0	2.7
NEVADA	59.5	62.9	3.3
NEW HAMPSHIRE	65.0	61.3	-3.7
NEW JERSEY	61.7	58.3	-3.4
NEW MEXICO	54.4	53.6	-0.8
NEW YORK	56.0	53.5	-2.5
NORTH CAROLINA	33.8	37.6	3.8
NORTH DAKOTA	52.0	53.7	1.7
OHIO	34.2	33.7	-0.5
OKLAHOMA	27.7	32.8	5.0
OREGON	56.7	56.7	-0.0
PENNSYLVANIA	58.0	49.3	-8.8
RHODE ISLAND	65.4	59.4	-6.0
SOUTH CAROLINA	38.3	39.7	1.4
SOUTH DAKOTA	47.0	55.5	8.5
TENNESSEE	28.3	28.7	0.4
TEXAS	50.6	49.4	-1.2
UTAH	29.0	28.3	-0.7
VERMONT	64.0	60.2	-3.8
VIRGINIA	53.8	54.9	1.0
WASHINGTON	58.6	60.4	1.8
WEST VIRGINIA	27.4	31.0	3.6
WISCONSIN	69.7	70.2	0.5
WYOMING	53.8	52.6	-1.1
PUERTO RICO	N/A	26.7	N/A

* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH

N/A - NOT AVAILABLE

August 14, 1998

TABLE 20.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKING*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	SAMPLE PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2144	11.4	0.8	(9.8, 13.0)
ALASKA	1513	16.5	1.5	(13.6, 19.3)
ARIZONA	1897	8.8	1.0	(6.9, 10.8)
ARKANSAS	1777	9.2	0.9	(7.5, 11.0)
CALIFORNIA	4050	15.2	0.7	(13.9, 16.5)
COLORADO	1808	15.3	1.0	(13.4, 17.3)
CONNECTICUT	2217	15.6	1.2	(13.3, 17.8)
DELAWARE	2539	11.9	0.8	(10.3, 13.6)
DISTRICT OF CO	1492	12.1	1.0	(10.1, 14.1)
FLORIDA	3464	13.1	0.7	(11.7, 14.4)
GEORGIA	2294	9.4	0.7	(7.9, 10.8)
HAWAII	2157	17.1	1.0	(15.2, 19.0)
IDAHO	4886	14.9	0.7	(13.5, 16.4)
ILLINOIS	1431	16.3	1.1	(14.1, 18.4)
INDIANA	2348	12.6	0.8	(11.1, 14.2)
IOWA	3573	17.9	0.8	(16.3, 19.4)
KANSAS	1988	13.3	0.9	(11.5, 15.1)
KENTUCKY	3589	9.4	0.6	(8.2, 10.6)

LOUISIANA	1624	15.2	1.1	(13.1, 17.3)
MAINE	1686	13.8	1.1	(11.6, 16.0)
MARYLAND	4557	6.3	0.5	(5.3, 7.2)
MASSACHUSETTS	1744	17.9	1.2	(15.6, 20.2)
MICHIGAN	2546	18.9	0.9	(17.2, 20.6)
MINNESOTA	4789	15.6	0.6	(14.5, 16.7)
MISSISSIPPI	1580	9.5	0.9	(7.7, 11.4)
MISSOURI	1834	15.1	1.1	(13.0, 17.1)
MONTANA	1772	14.0	0.9	(12.2, 15.9)
NEBRASKA	2668	16.3	1.0	(14.3, 18.3)
NEVADA	2483	19.2	1.9	(15.6, 22.8)
NEW HAMPSHIRE	1501	16.1	1.1	(13.9, 18.3)
NEW JERSEY	2629	13.1	0.8	(11.4, 14.7)
NEW MEXICO	1799	14.6	1.1	(12.5, 16.6)
NEW YORK	3376	9.2	0.6	(8.1, 10.3)
NORTH CAROLINA	3576	9.0	0.6	(7.9, 10.1)
NORTH DAKOTA	1791	18.4	1.0	(16.3, 20.4)
OHIO	3156	8.7	0.7	(7.3, 10.1)
OKLAHOMA	1879	8.8	0.9	(7.0, 10.5)
OREGON	3302	14.3	0.8	(12.7, 16.0)
PENNSYLVANIA	3548	14.6	0.8	(13.1, 16.0)
RHODE ISLAND	1821	14.9	1.1	(12.8, 17.0)
SOUTH CAROLINA	2132	9.7	0.8	(8.2, 11.3)
SOUTH DAKOTA	2182	20.9	1.0	(18.9, 22.8)
TENNESSEE	2950	7.2	0.6	(6.0, 8.3)
TEXAS	2474	17.4	0.9	(15.7, 19.2)
UTAH	2843	7.7	0.6	(6.5, 8.9)
VERMONT	3148	16.1	0.9	(14.3, 17.8)
VIRGINIA	3482	14.5	1.0	(12.6, 16.4)
WASHINGTON	3573	14.5	0.8	(13.0, 16.0)
WEST VIRGINIA	2423	8.4	0.7	(7.1, 9.7)
WISCONSIN	2225	23.3	1.1	(21.1, 25.5)
WYOMING	2395	15.4	1.0	(13.6, 17.3)
PUERTO RICO	2194	10.9	0.8	(9.4, 12.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 14.5 RANGE = 6.3-23.3

* 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH August 14, 1998

TABLE 20.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKING* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES-----*				FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	* SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	789	18.3	1.5	(15.3, 21.3)	*	1355	5.5	0.7 (4.2, 6.7)
ALASKA	694	24.9	2.5	(20.1, 29.7)	*	819	7.5	1.3 (5.0, 10.0)
ARIZONA	888	12.2	1.7	(8.8, 15.6)	*	1009	5.6	1.0 (3.7, 7.5)
ARKANSAS	669	16.1	1.7	(12.7, 19.5)	*	1108	3.1	0.5 (2.0, 4.1)
CALIFORNIA	1753	23.5	1.2	(21.2, 25.8)	*	2297	7.0	0.6 (5.9, 8.2)
COLORADO	802	23.6	1.7	(20.3, 27.0)	*	1006	7.5	1.0 (5.7, 9.4)
CONNECTICUT	871	23.9	2.1	(19.8, 28.0)	*	1346	8.0	1.0 (6.0, 9.9)
DELAWARE	1001	20.8	1.6	(17.7, 23.8)	*	1538	4.0	0.6 (2.8, 5.1)
DISTRICT OF CO	567	15.2	1.8	(11.8, 18.7)	*	925	9.4	1.2 (7.1, 11.8)

FLORIDA	1419	20.3	1.2	(17.9, 22.7)	*	2045	6.5	0.6	(5.4, 7.7)
GEORGIA	952	15.0	1.3	(12.4, 17.6)	*	1342	4.2	0.7	(3.0, 5.5)
HAWAII	942	27.3	1.6	(24.1, 30.5)	*	1215	6.8	0.9	(5.0, 8.5)
IDAHO	2037	22.0	1.2	(19.6, 24.4)	*	2849	8.2	0.9	(6.5, 9.9)
ILLINOIS	570	25.4	1.9	(21.6, 29.1)	*	861	8.3	1.0	(6.3, 10.3)
INDIANA	1053	19.5	1.4	(16.9, 22.2)	*	1295	6.4	0.8	(4.8, 8.0)
IOWA	1463	26.9	1.3	(24.3, 29.5)	*	2110	9.7	0.8	(8.1, 11.2)
KANSAS	778	21.3	1.7	(18.0, 24.5)	*	1210	5.9	0.8	(4.5, 7.4)
KENTUCKY	1308	15.7	1.1	(13.5, 17.9)	*	2281	3.7	0.5	(2.7, 4.7)
LOUISIANA	609	24.2	1.9	(20.4, 28.0)	*	1015	7.4	0.9	(5.6, 9.1)
MAINE	745	22.5	2.0	(18.7, 26.3)	*	941	5.8	0.8	(4.2, 7.4)
MARYLAND	1946	9.8	0.9	(8.1, 11.5)	*	2611	3.1	0.4	(2.3, 3.9)
MASSACHUSETTS	709	27.5	2.1	(23.5, 31.6)	*	1035	9.4	1.0	(7.4, 11.4)
MICHIGAN	1062	29.1	1.5	(26.1, 32.1)	*	1484	9.7	0.8	(8.1, 11.2)
MINNESOTA	2186	22.7	1.0	(20.8, 24.7)	*	2603	8.9	0.6	(7.7, 10.1)
MISSISSIPPI	579	17.4	1.8	(13.9, 20.9)	*	1001	2.7	0.7	(1.3, 4.2)
MISSOURI	688	23.8	1.9	(20.0, 27.6)	*	1146	7.3	1.0	(5.4, 9.1)
MONTANA	736	22.0	1.7	(18.7, 25.2)	*	1036	6.7	0.9	(5.0, 8.4)
NEBRASKA	1028	24.9	1.8	(21.3, 28.4)	*	1640	8.4	0.9	(6.7, 10.1)
NEVADA	1146	29.0	3.2	(22.8, 35.2)	*	1337	9.3	1.4	(6.5, 12.1)
NEW HAMPSHIRE	597	25.1	1.9	(21.2, 28.9)	*	904	7.8	1.0	(5.8, 9.8)
NEW JERSEY	1045	20.6	1.5	(17.7, 23.6)	*	1584	6.2	0.8	(4.8, 7.7)
NEW MEXICO	748	23.1	1.9	(19.4, 26.7)	*	1051	6.6	0.8	(5.0, 8.2)
NEW YORK	1340	13.5	1.1	(11.5, 15.6)	*	2036	5.4	0.6	(4.3, 6.5)
NORTH CAROLINA	1405	15.4	1.1	(13.3, 17.5)	*	2171	3.3	0.5	(2.4, 4.2)
NORTH DAKOTA	791	26.4	1.7	(23.0, 29.8)	*	1000	10.6	1.2	(8.3, 12.9)
OHIO	1224	13.6	1.3	(11.1, 16.1)	*	1932	4.3	0.6	(3.1, 5.6)
OKLAHOMA	825	13.7	1.6	(10.5, 16.9)	*	1054	4.3	0.7	(2.8, 5.7)
OREGON	1390	21.9	1.5	(18.9, 24.8)	*	1912	7.3	0.7	(6.0, 8.6)
PENNSYLVANIA	1418	23.4	1.3	(20.8, 25.9)	*	2130	6.8	0.8	(5.3, 8.3)
RHODE ISLAND	741	21.6	1.8	(18.2, 25.1)	*	1080	9.1	1.3	(6.5, 11.7)
SOUTH CAROLINA	805	16.6	1.5	(13.6, 19.5)	*	1327	3.7	0.6	(2.6, 4.8)
SOUTH DAKOTA	944	32.1	1.7	(28.7, 35.4)	*	1238	10.5	1.0	(8.5, 12.5)
TENNESSEE	1030	11.4	1.1	(9.3, 13.5)	*	1920	3.4	0.5	(2.4, 4.3)
TEXAS	1010	27.9	1.6	(24.9, 31.0)	*	1464	7.6	0.8	(6.1, 9.1)
UTAH	1269	12.3	1.1	(10.1, 14.4)	*	1574	3.4	0.5	(2.4, 4.4)
VERMONT	1370	25.6	1.5	(22.6, 28.6)	*	1778	7.3	0.9	(5.5, 9.0)
VIRGINIA	1390	23.5	1.7	(20.1, 27.0)	*	2092	6.1	0.8	(4.5, 7.7)
WASHINGTON	1536	22.7	1.3	(20.1, 25.3)	*	2037	6.6	0.7	(5.3, 7.9)
WEST VIRGINIA	956	13.8	1.2	(11.4, 16.2)	*	1467	3.6	0.6	(2.4, 4.7)
WISCONSIN	979	36.3	1.9	(32.6, 39.9)	*	1246	11.2	1.2	(8.9, 13.6)
WYOMING	1007	23.5	1.5	(20.5, 26.6)	*	1388	7.6	1.1	(5.4, 9.8)
PUERTO RICO	881	21.1	1.6	(18.0, 24.1)	*	1313	2.4	0.6	(1.3, 3.5)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
MEDIAN = 22.3
RANGE = 9.8-36.3

* 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH

NUMBER OF PARTICIPANTS = 52
MEDIAN = 6.7
RANGE = 2.4-11.2

August 14, 1998

TABLE 20.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1995 TO 1997
BINGE DRINKING*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
-------------	--------------	--------------	--------------------

ALABAMA	13.8	11.4	-2.4
ALASKA	19.5	16.5	-3.0
ARIZONA	13.7	8.8	-4.9
ARKANSAS	8.9	9.2	0.4
CALIFORNIA	15.4	15.2	-0.2
COLORADO	16.4	15.3	-1.1
CONNECTICUT	14.5	15.6	1.0
DELAWARE	8.7	11.9	3.3
DISTRICT OF CO	N/A	12.1	N/A
FLORIDA	13.2	13.1	-0.2
GEORGIA	12.0	9.4	-2.6
HAWAII	12.4	17.1	4.7
IDAHO	13.1	14.9	1.9
ILLINOIS	13.6	16.3	2.6
INDIANA	12.9	12.6	-0.3
IOWA	18.0	17.9	-0.2
KANSAS	14.1	13.3	-0.8
KENTUCKY	9.8	9.4	-0.4
LOUISIANA	14.2	15.2	1.0
MAINE	11.5	13.8	2.3
MARYLAND	8.3	6.3	-2.0
MASSACHUSETTS	18.0	17.9	-0.1
MICHIGAN	18.3	18.9	0.6
MINNESOTA	18.2	15.6	-2.6
MISSISSIPPI	8.8	9.5	0.8
MISSOURI	14.3	15.1	0.8
MONTANA	14.4	14.0	-0.4
NEBRASKA	16.1	16.3	0.2
NEVADA	19.1	19.2	0.1
NEW HAMPSHIRE	16.8	16.1	-0.6
NEW JERSEY	14.1	13.1	-1.0
NEW MEXICO	14.2	14.6	0.4
NEW YORK	12.5	9.2	-3.3
NORTH CAROLINA	5.8	9.0	3.2
NORTH DAKOTA	17.0	18.4	1.3
OHIO	9.9	8.7	-1.2
OKLAHOMA	6.7	8.8	2.1
OREGON	14.0	14.3	0.4
PENNSYLVANIA	19.5	14.6	-4.9
RHODE ISLAND	18.8	14.9	-3.9
SOUTH CAROLINA	9.3	9.7	0.4
SOUTH DAKOTA	14.5	20.9	6.4
TENNESSEE	5.3	7.2	1.9
TEXAS	15.4	17.4	2.0
UTAH	10.0	7.7	-2.3
VERMONT	16.1	16.1	-0.1
VIRGINIA	14.7	14.5	-0.2
WASHINGTON	13.7	14.5	0.8
WEST VIRGINIA	6.0	8.4	2.4
WISCONSIN	23.1	23.3	0.2
WYOMING	15.7	15.4	-0.3
PUERTO RICO	N/A	10.9	N/A

* 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH

N/A - NOT AVAILABLE

August 14, 1998

TABLE 21.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHRONIC DRINKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	SAMPLE PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2114	3.0	0.5	(2.1, 3.8)
ALASKA	1493	4.0	0.8	(2.4, 5.5)
ARIZONA	1892	2.6	0.6	(1.5, 3.7)
ARKANSAS	1768	2.3	0.5	(1.3, 3.3)
COLORADO	1798	2.7	0.4	(1.9, 3.6)
CONNECTICUT	2199	3.2	0.5	(2.2, 4.2)
DELAWARE	2530	3.4	0.5	(2.4, 4.4)
DISTRICT OF CO	1492	3.0	0.5	(1.9, 4.0)
FLORIDA	3460	4.0	0.4	(3.2, 4.8)
GEORGIA	2272	1.8	0.3	(1.2, 2.4)
HAWAII	2141	5.1	0.6	(3.9, 6.2)
IDAHO	4870	3.0	0.3	(2.3, 3.6)
ILLINOIS	1429	3.4	0.6	(2.3, 4.4)
INDIANA	2324	3.0	0.5	(2.1, 3.9)
IOWA	3561	3.5	0.4	(2.7, 4.2)
KANSAS	1969	2.3	0.4	(1.5, 3.1)
KENTUCKY	3568	2.7	0.3	(2.1, 3.4)
LOUISIANA	1606	4.6	0.7	(3.2, 5.9)
MAINE	1664	2.7	0.5	(1.8, 3.5)
MARYLAND	4530	1.2	0.2	(0.8, 1.7)
MASSACHUSETTS	1731	3.6	0.6	(2.4, 4.8)
MICHIGAN	2524	3.8	0.4	(2.9, 4.6)
MINNESOTA	4752	2.7	0.3	(2.2, 3.3)
MISSISSIPPI	1571	2.5	0.5	(1.5, 3.4)
MISSOURI	1816	3.2	0.6	(2.1, 4.4)
MONTANA	1753	2.6	0.4	(1.8, 3.4)
NEBRASKA	2641	2.3	0.4	(1.6, 3.0)
NEVADA	2443	4.7	0.8	(3.2, 6.2)
NEW HAMPSHIRE	1489	2.5	0.5	(1.6, 3.5)
NEW JERSEY	2604	2.0	0.3	(1.4, 2.7)
NEW MEXICO	1790	3.6	0.5	(2.6, 4.5)
NEW YORK	3307	2.4	0.3	(1.8, 3.1)
NORTH CAROLINA	3537	2.7	0.3	(2.0, 3.4)
NORTH DAKOTA	1781	2.5	0.4	(1.7, 3.2)
OHIO	3142	2.4	0.4	(1.7, 3.1)
OKLAHOMA	1867	2.2	0.5	(1.1, 3.2)
OREGON	3277	3.5	0.4	(2.8, 4.3)
PENNSYLVANIA	3521	3.1	0.4	(2.4, 3.8)
RHODE ISLAND	1792	4.6	0.7	(3.3, 5.9)
SOUTH CAROLINA	2116	3.3	0.5	(2.3, 4.3)
SOUTH DAKOTA	2163	3.2	0.4	(2.3, 4.1)
TENNESSEE	2922	1.8	0.3	(1.2, 2.4)
TEXAS	2448	4.9	0.5	(3.9, 5.8)
UTAH	2845	1.7	0.3	(1.1, 2.3)
VERMONT	3115	4.1	0.5	(3.1, 5.1)
VIRGINIA	3436	3.3	0.5	(2.3, 4.3)
WASHINGTON	3528	3.6	0.4	(2.8, 4.4)
WEST VIRGINIA	2404	2.1	0.4	(1.4, 2.8)
WISCONSIN	2223	5.0	0.6	(3.9, 6.0)
WYOMING	2385	2.8	0.4	(2.0, 3.5)
PUERTO RICO	2179	3.7	0.5	(2.7, 4.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 3.0 RANGE = 1.2- 5.1

* 60 OR MORE DRINKS DURING THE PAST MONTH

August 14, 1998

TABLE 21.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHRONIC DRINKERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES-----*				FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	783	5.7	0.9	(3.9, 7.5)	*	1331	0.6	(0.2, 0.9)
ALASKA	680	6.6	1.5	(3.8, 9.5)	*	813	1.1	(-0.0, 2.3)
ARIZONA	887	4.1	1.1	(2.0, 6.3)	*	1005	1.1	(0.3, 2.0)
ARKANSAS	666	4.4	1.0	(2.4, 6.4)	*	1102	0.4	(0.0, 0.7)
COLORADO	801	5.0	0.8	(3.4, 6.6)	*	997	0.6	(-0.0, 1.2)
CONNECTICUT	864	5.1	0.9	(3.4, 6.8)	*	1335	1.5	(0.3, 2.6)
DELAWARE	997	6.0	1.0	(4.1, 8.0)	*	1533	1.0	(0.4, 1.6)
DISTRICT OF CO	567	5.3	1.1	(3.2, 7.4)	*	925	1.0	(0.3, 1.8)
FLORIDA	1425	6.8	0.8	(5.2, 8.4)	*	2035	1.4	(0.8, 1.9)
GEORGIA	940	3.0	0.5	(1.9, 4.0)	*	1332	0.7	(0.1, 1.3)
HAWAII	936	8.6	1.1	(6.5, 10.6)	*	1205	1.4	(0.6, 2.3)
IDAHO	2032	5.4	0.7	(4.1, 6.7)	*	2838	0.7	(0.4, 1.0)
ILLINOIS	571	5.8	1.1	(3.7, 7.9)	*	858	1.2	(0.4, 2.1)
INDIANA	1048	5.4	0.9	(3.7, 7.2)	*	1276	0.7	(0.2, 1.2)
IOWA	1456	6.0	0.7	(4.6, 7.4)	*	2105	1.2	(0.6, 1.8)
KANSAS	766	4.3	0.8	(2.7, 5.9)	*	1203	0.5	(0.1, 1.0)
KENTUCKY	1298	4.9	0.7	(3.6, 6.2)	*	2270	0.8	(0.4, 1.2)
LOUISIANA	609	8.9	1.4	(6.2, 11.5)	*	997	0.7	(0.2, 1.3)
MAINE	732	4.5	0.9	(2.8, 6.2)	*	932	1.0	(0.3, 1.6)
MARYLAND	1933	2.3	0.4	(1.4, 3.1)	*	2597	0.3	(0.0, 0.6)
MASSACHUSETTS	708	6.7	1.2	(4.4, 9.0)	*	1023	0.8	(0.2, 1.5)
MICHIGAN	1056	7.0	0.9	(5.3, 8.6)	*	1468	0.9	(0.4, 1.3)
MINNESOTA	2170	5.0	0.5	(3.9, 6.0)	*	2582	0.7	(0.3, 1.0)
MISSISSIPPI	571	4.4	1.0	(2.5, 6.4)	*	1000	0.8	(0.0, 1.5)
MISSOURI	683	6.2	1.2	(4.0, 8.5)	*	1133	0.6	(0.2, 1.0)
MONTANA	726	5.0	0.9	(3.3, 6.7)	*	1027	0.3	(-0.1, 0.8)
NEBRASKA	1016	4.6	0.8	(3.1, 6.1)	*	1625	0.2	(0.0, 0.4)
NEVADA	1125	7.2	1.4	(4.5, 9.9)	*	1318	2.3	(0.9, 3.7)
NEW HAMPSHIRE	594	4.9	1.0	(3.0, 6.8)	*	895	0.3	(-0.0, 0.7)
NEW JERSEY	1035	4.0	0.7	(2.6, 5.3)	*	1569	0.3	(0.0, 0.6)
NEW MEXICO	744	6.9	1.0	(5.0, 8.8)	*	1046	0.5	(0.0, 0.9)
NEW YORK	1308	3.9	0.6	(2.7, 5.1)	*	1999	1.1	(0.6, 1.6)
NORTH CAROLINA	1387	4.9	0.7	(3.5, 6.2)	*	2150	0.8	(0.3, 1.2)
NORTH DAKOTA	790	4.2	0.7	(2.8, 5.6)	*	991	0.8	(0.2, 1.3)
OHIO	1217	4.1	0.7	(2.7, 5.5)	*	1925	0.9	(0.3, 1.5)
OKLAHOMA	820	3.9	1.0	(1.9, 5.9)	*	1047	0.6	(-0.2, 1.4)
OREGON	1380	6.3	0.8	(4.8, 7.8)	*	1897	1.0	(0.5, 1.4)
PENNSYLVANIA	1405	5.7	0.7	(4.3, 7.1)	*	2116	0.8	(0.4, 1.3)
RHODE ISLAND	729	7.8	1.1	(5.6, 10.0)	*	1063	1.9	(0.4, 3.3)
SOUTH CAROLINA	802	6.3	1.0	(4.3, 8.2)	*	1314	0.6	(0.1, 1.2)
SOUTH DAKOTA	936	5.9	0.9	(4.1, 7.6)	*	1227	0.7	(0.2, 1.2)
TENNESSEE	1014	3.5	0.6	(2.2, 4.7)	*	1908	0.3	(0.1, 0.6)
TEXAS	1000	8.2	0.9	(6.4, 10.1)	*	1448	1.7	(0.9, 2.5)
UTAH	1269	3.1	0.6	(1.9, 4.2)	*	1576	0.4	(0.1, 0.7)
VERMONT	1362	6.6	0.8	(4.9, 8.2)	*	1753	1.8	(0.5, 3.1)
VIRGINIA	1370	6.1	1.0	(4.1, 8.1)	*	2066	0.7	(0.3, 1.1)
WASHINGTON	1512	6.5	0.8	(5.0, 8.1)	*	2016	0.9	(0.5, 1.3)
WEST VIRGINIA	945	3.9	0.7	(2.6, 5.3)	*	1459	0.5	(0.1, 0.9)
WISCONSIN	980	8.7	1.0	(6.6, 10.7)	*	1243	1.6	(0.7, 2.4)
WYOMING	1003	5.0	0.8	(3.5, 6.5)	*	1382	0.5	(0.2, 0.9)
PUERTO RICO	877	7.6	1.0	(5.6, 9.5)	*	1302	0.4	(0.0, 0.8)

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 51
 MEDIAN = 5.4
 RANGE = 2.3- 8.9

NUMBER OF PARTICIPANTS = 51
 MEDIAN = 0.8
 RANGE = 0.2- 2.3

TABLE 21.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 CHRONIC DRINKERS*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	2.6	3.0	0.4
ALASKA	3.0	4.0	1.0
ARIZONA	2.5	2.6	0.1
ARKANSAS	2.1	2.3	0.2
COLORADO	4.8	2.7	-2.1
CONNECTICUT	4.5	3.2	-1.3
DELAWARE	2.6	3.4	0.8
DISTRICT OF CO	N/A	3.0	N/A
FLORIDA	4.3	4.0	-0.4
GEORGIA	2.8	1.8	-1.0
HAWAII	2.9	5.1	2.2
IDAHO	2.8	3.0	0.2
ILLINOIS	2.9	3.4	0.5
INDIANA	3.7	3.0	-0.7
IOWA	3.8	3.5	-0.3
KANSAS	3.6	2.3	-1.2
KENTUCKY	2.7	2.7	0.0
LOUISIANA	2.9	4.6	1.7
MAINE	1.6	2.7	1.1
MARYLAND	2.3	1.2	-1.1
MASSACHUSETTS	3.3	3.6	0.3
MICHIGAN	2.9	3.8	0.9
MINNESOTA	4.7	2.7	-2.0
MISSISSIPPI	2.7	2.5	-0.2
MISSOURI	3.2	3.2	0.1
MONTANA	1.9	2.6	0.7
NEBRASKA	2.3	2.3	-0.0
NEVADA	5.7	4.7	-0.9
NEW HAMPSHIRE	3.2	2.5	-0.7
NEW JERSEY	2.7	2.0	-0.7
NEW MEXICO	3.0	3.6	0.6
NEW YORK	2.4	2.4	0.1
NORTH CAROLINA	2.0	2.7	0.7
NORTH DAKOTA	2.4	2.5	0.0
OHIO	1.7	2.4	0.7
OKLAHOMA	1.2	2.2	1.0
OREGON	3.8	3.5	-0.2
PENNSYLVANIA	4.3	3.1	-1.2
RHODE ISLAND	5.3	4.6	-0.7
SOUTH CAROLINA	2.2	3.3	1.1
SOUTH DAKOTA	2.0	3.2	1.2
TENNESSEE	1.5	1.8	0.3
TEXAS	3.9	4.9	1.0
UTAH	1.5	1.7	0.2
VERMONT	4.7	4.1	-0.5

VIRGINIA	3.4	3.3	-0.1
WASHINGTON	2.8	3.6	0.8
WEST VIRGINIA	1.6	2.1	0.6
WISCONSIN	4.5	5.0	0.5
WYOMING	2.4	2.8	0.3
PUERTO RICO	N/A	3.7	N/A

* 60 OR MORE DRINKS DURING THE PAST MONTH

N/A - NOT AVAILABLE

August 14, 1998

TABLE 22.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DRINKING AND DRIVING*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2162	1.5	0.3	(0.9, 2.2)
ALASKA	1538	2.2	0.7	(0.9, 3.5)
ARIZONA	1900	1.4	0.4	(0.6, 2.2)
ARKANSAS	1787	1.6	0.4	(0.9, 2.3)
CALIFORNIA	4055	2.5	0.3	(2.0, 3.1)
COLORADO	1820	2.5	0.4	(1.7, 3.3)
CONNECTICUT	2235	2.1	0.4	(1.2, 2.9)
DELAWARE	2552	2.0	0.4	(1.1, 2.9)
DISTRICT OF CO	1503	2.5	0.6	(1.4, 3.6)
FLORIDA	3486	1.9	0.3	(1.3, 2.4)
GEORGIA	2293	1.1	0.3	(0.6, 1.5)
HAWAII	2162	2.4	0.4	(1.6, 3.1)
IDAHO	4911	1.3	0.2	(0.9, 1.6)
ILLINOIS	1438	2.8	0.5	(1.9, 3.8)
INDIANA	2360	1.9	0.3	(1.3, 2.5)
IOWA	3583	3.8	0.4	(3.1, 4.5)
KANSAS	1991	2.7	0.4	(1.9, 3.5)
KENTUCKY	3598	0.6	0.1	(0.3, 0.8)
LOUISIANA	1645	3.2	0.5	(2.2, 4.3)
MAINE	1692	0.9	0.2	(0.4, 1.3)
MARYLAND	4562	0.9	0.2	(0.5, 1.3)
MASSACHUSETTS	1760	1.8	0.4	(1.1, 2.4)
MICHIGAN	2545	3.5	0.4	(2.7, 4.2)
MINNESOTA	4800	3.8	0.3	(3.2, 4.4)
MISSISSIPPI	1597	1.2	0.3	(0.5, 1.9)
MISSOURI	1846	3.0	0.5	(1.9, 4.1)
MONTANA	1786	2.6	0.4	(1.8, 3.5)
NEBRASKA	2678	3.8	0.7	(2.5, 5.1)
NEVADA	2501	3.1	0.9	(1.4, 4.8)
NEW HAMPSHIRE	1509	1.9	0.4	(1.1, 2.7)
NEW JERSEY	2645	1.5	0.3	(0.9, 2.1)
NEW MEXICO	1808	1.7	0.4	(1.0, 2.5)
NEW YORK	3393	0.9	0.2	(0.5, 1.2)
NORTH CAROLINA	3611	1.1	0.2	(0.7, 1.5)
NORTH DAKOTA	1795	3.8	0.5	(2.8, 4.7)
OHIO	3162	1.0	0.2	(0.6, 1.5)
OKLAHOMA	1884	1.5	0.4	(0.7, 2.2)

OREGON	3316	1.6	0.2	(1.1, 2.1)
PENNSYLVANIA	3581	1.7	0.2	(1.2, 2.2)
RHODE ISLAND	1835	1.6	0.3	(1.0, 2.2)
SOUTH CAROLINA	2147	0.9	0.2	(0.4, 1.3)
SOUTH DAKOTA	2190	3.7	0.5	(2.7, 4.6)
TENNESSEE	2969	1.0	0.2	(0.5, 1.4)
TEXAS	2491	4.0	0.4	(3.1, 4.9)
UTAH	2859	0.8	0.2	(0.4, 1.2)
VERMONT	3180	3.0	0.6	(1.8, 4.1)
VIRGINIA	3511	2.4	0.4	(1.7, 3.0)
WASHINGTON	3590	1.7	0.2	(1.2, 2.2)
WEST VIRGINIA	2428	0.8	0.2	(0.5, 1.2)
WISCONSIN	2236	5.3	0.6	(4.1, 6.4)
WYOMING	2408	2.6	0.4	(1.8, 3.5)
PUERTO RICO	2252	3.2	0.5	(2.3, 4.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 1.9 RANGE = 0.6- 5.3

* DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES

August 14, 1998

TABLE 22.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DRINKING AND DRIVING* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD	95% CONFIDENCE	SAMPLE SIZE	PERCENT	STANDARD	95% CONFIDENCE
			ERROR	INTERVAL			ERROR	INTERVAL
ALABAMA	806	2.4	0.6	(1.2, 3.6)	*	1356	0.7	0.3 (0.2, 1.2)
ALASKA	711	3.3	1.2	(0.9, 5.7)	*	827	0.9	0.5 (-0.0, 1.9)
ARIZONA	889	2.5	0.8	(0.9, 4.1)	*	1011	0.4	0.2 (-0.1, 0.8)
ARKANSAS	673	2.7	0.7	(1.3, 4.1)	*	1114	0.5	0.2 (0.1, 1.0)
CALIFORNIA	1755	4.1	0.6	(3.1, 5.2)	*	2300	1.0	0.2 (0.6, 1.3)
COLORADO	814	4.2	0.8	(2.7, 5.7)	*	1006	0.9	0.3 (0.2, 1.5)
CONNECTICUT	880	2.7	0.6	(1.5, 3.9)	*	1355	1.5	0.6 (0.3, 2.7)
DELAWARE	1010	3.9	0.9	(2.1, 5.7)	*	1542	0.3	0.1 (0.0, 0.5)
DISTRICT OF CO	572	3.7	1.1	(1.6, 5.8)	*	931	1.6	0.5 (0.6, 2.5)
FLORIDA	1433	2.7	0.5	(1.7, 3.7)	*	2053	1.1	0.2 (0.7, 1.6)
GEORGIA	952	1.5	0.4	(0.6, 2.3)	*	1341	0.7	0.3 (0.1, 1.2)
HAWAII	945	3.4	0.6	(2.1, 4.6)	*	1217	1.4	0.5 (0.5, 2.3)
IDAHO	2052	2.2	0.4	(1.5, 2.9)	*	2859	0.4	0.1 (0.2, 0.7)
ILLINOIS	574	5.1	1.0	(3.2, 7.0)	*	864	0.9	0.3 (0.3, 1.4)
INDIANA	1060	3.4	0.6	(2.3, 4.5)	*	1300	0.6	0.2 (0.2, 0.9)
IOWA	1469	6.0	0.7	(4.7, 7.3)	*	2114	1.9	0.3 (1.2, 2.5)
KANSAS	779	4.2	0.7	(2.7, 5.6)	*	1212	1.3	0.4 (0.6, 2.0)
KENTUCKY	1315	0.8	0.2	(0.3, 1.2)	*	2283	0.4	0.2 (0.1, 0.7)
LOUISIANA	624	5.8	1.1	(3.8, 7.9)	*	1021	0.9	0.3 (0.3, 1.5)
MAINE	747	1.3	0.4	(0.5, 2.0)	*	945	0.5	0.2 (0.0, 1.0)
MARYLAND	1953	1.7	0.4	(1.0, 2.5)	*	2609	0.2	0.1 (-0.0, 0.4)
MASSACHUSETTS	715	2.3	0.6	(1.0, 3.5)	*	1045	1.3	0.4 (0.6, 2.0)
MICHIGAN	1062	5.6	0.7	(4.1, 7.0)	*	1483	1.5	0.4 (0.8, 2.2)
MINNESOTA	2190	5.9	0.6	(4.8, 7.0)	*	2610	1.8	0.3 (1.2, 2.3)
MISSISSIPPI	591	1.4	0.5	(0.5, 2.3)	*	1006	1.0	0.5 (0.0, 2.0)
MISSOURI	695	5.2	1.1	(3.1, 7.3)	*	1151	1.0	0.4 (0.3, 1.8)
MONTANA	748	4.5	0.8	(2.9, 6.1)	*	1038	0.9	0.3 (0.3, 1.5)
NEBRASKA	1033	6.1	1.3	(3.5, 8.7)	*	1645	1.7	0.4 (1.0, 2.4)

NEVADA	1161	4.2	1.6	(1.1, 7.2)	*	1340	2.0	0.8	(0.4, 3.5)
NEW HAMPSHIRE	601	2.6	0.7	(1.2, 4.0)	*	908	1.3	0.4	(0.4, 2.1)
NEW JERSEY	1052	1.8	0.5	(0.9, 2.8)	*	1593	1.2	0.4	(0.5, 1.9)
NEW MEXICO	751	2.6	0.7	(1.2, 4.0)	*	1057	0.9	0.3	(0.3, 1.5)
NEW YORK	1347	1.6	0.4	(0.8, 2.3)	*	2046	0.2	0.1	(0.0, 0.4)
NORTH CAROLINA	1430	2.0	0.4	(1.1, 2.8)	*	2181	0.3	0.1	(0.1, 0.6)
NORTH DAKOTA	794	5.1	0.8	(3.5, 6.7)	*	1001	2.4	0.5	(1.4, 3.5)
OHIO	1228	1.6	0.4	(0.8, 2.5)	*	1934	0.5	0.2	(0.1, 0.9)
OKLAHOMA	829	2.3	0.8	(0.8, 3.8)	*	1055	0.7	0.3	(0.2, 1.2)
OREGON	1398	2.5	0.4	(1.7, 3.4)	*	1918	0.7	0.2	(0.2, 1.1)
PENNSYLVANIA	1436	3.2	0.5	(2.2, 4.2)	*	2145	0.4	0.1	(0.2, 0.6)
RHODE ISLAND	751	2.6	0.6	(1.5, 3.8)	*	1084	0.7	0.3	(0.2, 1.2)
SOUTH CAROLINA	815	1.5	0.5	(0.6, 2.4)	*	1332	0.3	0.2	(0.0, 0.6)
SOUTH DAKOTA	951	5.5	0.9	(3.8, 7.2)	*	1239	2.0	0.4	(1.2, 2.8)
TENNESSEE	1040	1.7	0.5	(0.8, 2.6)	*	1929	0.4	0.1	(0.1, 0.6)
TEXAS	1018	6.7	0.8	(5.0, 8.3)	*	1473	1.5	0.3	(0.9, 2.1)
UTAH	1275	1.4	0.4	(0.6, 2.1)	*	1584	0.3	0.2	(-0.0, 0.7)
VERMONT	1388	5.4	1.1	(3.1, 7.6)	*	1792	0.7	0.2	(0.3, 1.2)
VIRGINIA	1412	3.7	0.6	(2.5, 5.0)	*	2099	1.1	0.3	(0.4, 1.7)
WASHINGTON	1544	2.3	0.4	(1.6, 3.1)	*	2046	1.1	0.3	(0.6, 1.6)
WEST VIRGINIA	960	1.4	0.4	(0.7, 2.1)	*	1468	0.3	0.2	(0.0, 0.6)
WISCONSIN	987	8.9	1.1	(6.8, 11.0)	*	1249	1.9	0.5	(0.9, 2.8)
WYOMING	1017	4.0	0.8	(2.4, 5.6)	*	1391	1.3	0.4	(0.6, 2.0)
PUERTO RICO	929	6.6	0.9	(4.8, 8.5)	*	1323	0.2	0.1	(-0.0, 0.5)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.0
 RANGE = 0.8- 8.9

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 0.9
 RANGE = 0.2- 2.4

* DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES

August 14, 1998

TABLE 22.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 DRINKING AND DRIVING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	2.6	1.5	-1.1
ALASKA	1.3	2.2	0.9
ARIZONA	2.7	1.4	-1.3
ARKANSAS	1.5	1.6	0.1
CALIFORNIA	1.9	2.5	0.6
COLORADO	3.1	2.5	-0.6
CONNECTICUT	2.5	2.1	-0.4
DELAWARE	1.4	2.0	0.6
DISTRICT OF CO	N/A	2.5	N/A
FLORIDA	2.6	1.9	-0.7
GEORGIA	2.2	1.1	-1.1
HAWAII	2.1	2.4	0.3
IDAHO	2.0	1.3	-0.8
ILLINOIS	1.8	2.8	1.1
INDIANA	2.6	1.9	-0.7
IOWA	3.3	3.8	0.5
KANSAS	3.2	2.7	-0.5
KENTUCKY	0.6	0.6	-0.0

LOUISIANA	2.8	3.2	0.4
MAINE	0.9	0.9	-0.1
MARYLAND	1.2	0.9	-0.2
MASSACHUSETTS	3.5	1.8	-1.8
MICHIGAN	3.3	3.5	0.1
MINNESOTA	4.9	3.8	-1.2
MISSISSIPPI	1.1	1.2	0.1
MISSOURI	2.1	3.0	0.9
MONTANA	3.4	2.6	-0.8
NEBRASKA	2.8	3.8	1.0
NEVADA	3.8	3.1	-0.7
NEW HAMPSHIRE	1.6	1.9	0.3
NEW JERSEY	2.0	1.5	-0.5
NEW MEXICO	3.3	1.7	-1.5
NEW YORK	0.9	0.9	-0.1
NORTH CAROLINA	1.1	1.1	-0.0
NORTH DAKOTA	4.2	3.8	-0.5
OHIO	1.7	1.0	-0.6
OKLAHOMA	1.2	1.5	0.3
OREGON	1.8	1.6	-0.2
PENNSYLVANIA	3.6	1.7	-1.9
RHODE ISLAND	3.7	1.6	-2.1
SOUTH CAROLINA	1.4	0.9	-0.5
SOUTH DAKOTA	5.3	3.7	-1.6
TENNESSEE	1.0	1.0	-0.1
TEXAS	3.7	4.0	0.3
UTAH	1.2	0.8	-0.4
VERMONT	2.4	3.0	0.5
VIRGINIA	2.6	2.4	-0.2
WASHINGTON	2.1	1.7	-0.4
WEST VIRGINIA	0.9	0.8	-0.0
WISCONSIN	4.6	5.3	0.7
WYOMING	3.2	2.6	-0.6
PUERTO RICO	N/A	3.2	N/A

* DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES

N/A - NOT AVAILABLE

August 14, 1998

TABLE 23.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FLU SHOT*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	440	62.6	2.5	(57.6, 67.5)
ALASKA	148	58.3	5.8	(46.9, 69.7)
ARIZONA	468	72.9	2.8	(67.5, 78.3)
ARKANSAS	394	61.1	2.7	(55.8, 66.3)
CALIFORNIA	737	65.5	1.9	(61.7, 69.3)
COLORADO	321	74.4	2.8	(69.0, 79.9)
CONNECTICUT	422	67.2	2.7	(61.9, 72.6)
DELAWARE	564	68.6	2.2	(64.4, 72.8)
DISTRICT OF CO	242	54.3	3.6	(47.2, 61.4)
FLORIDA	957	62.3	1.8	(58.9, 65.8)

GEORGIA	345	58.5	2.9	(52.7, 64.3)
HAWAII	392	71.1	2.7	(65.9, 76.3)
IDAHO	934	66.4	1.8	(62.9, 70.0)
ILLINOIS	254	67.8	3.3	(61.3, 74.3)
INDIANA	425	62.5	2.7	(57.3, 67.8)
IOWA	839	69.7	1.7	(66.3, 73.1)
KANSAS	417	61.5	2.7	(56.3, 66.7)
KENTUCKY	835	61.2	1.9	(57.5, 65.0)
LOUISIANA	302	58.4	3.1	(52.3, 64.5)
MAINE	311	72.1	2.7	(66.7, 77.4)
MARYLAND	822	63.4	2.3	(59.0, 67.9)
MASSACHUSETTS	287	66.0	3.1	(60.1, 72.0)
MICHIGAN	414	63.6	2.6	(58.6, 68.6)
MINNESOTA	918	69.0	1.7	(65.7, 72.2)
MISSISSIPPI	326	61.1	2.8	(55.6, 66.6)
MISSOURI	378	70.3	2.6	(65.3, 75.3)
MONTANA	388	68.4	2.7	(63.0, 73.7)
NEBRASKA	661	65.8	2.1	(61.7, 69.9)
NEVADA	360	56.5	5.2	(46.3, 66.8)
NEW HAMPSHIRE	266	64.6	3.2	(58.4, 70.8)
NEW JERSEY	484	60.7	2.5	(55.9, 65.5)
NEW MEXICO	347	72.8	2.5	(67.9, 77.7)
NEW YORK	589	64.5	2.1	(60.3, 68.6)
NORTH CAROLINA	736	64.6	1.9	(60.8, 68.4)
NORTH DAKOTA	445	64.8	2.5	(59.9, 69.7)
OHIO	825	65.4	2.1	(61.3, 69.5)
OKLAHOMA	613	69.3	2.2	(65.1, 73.5)
OREGON	674	69.8	2.1	(65.8, 73.9)
PENNSYLVANIA	712	65.8	1.9	(62.0, 69.6)
RHODE ISLAND	368	67.7	2.7	(62.4, 73.0)
SOUTH CAROLINA	477	74.3	2.1	(70.1, 78.5)
SOUTH DAKOTA	551	65.6	2.3	(61.0, 70.1)
TENNESSEE	571	69.1	2.1	(65.0, 73.2)
TEXAS	368	68.0	2.6	(62.9, 73.2)
UTAH	427	66.2	2.9	(60.5, 71.8)
VERMONT	571	69.5	2.1	(65.4, 73.5)
VIRGINIA	558	67.7	2.3	(63.3, 72.1)
WASHINGTON	587	70.3	2.0	(66.3, 74.2)
WEST VIRGINIA	583	58.2	2.3	(53.7, 62.6)
WISCONSIN	409	66.1	2.7	(60.7, 71.4)
WYOMING	404	72.4	2.5	(67.6, 77.3)
PUERTO RICO	409	41.5	2.7	(36.2, 46.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 65.9 RANGE = 41.5-74.4 YEAR 2000 OBJ. 20.11
 * HAD FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OVER August 14, 1998

TABLE 23.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FLU SHOT* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-MALES-				-FEMALES-			
	SAMPLE SIZE	STANDARD PERCENT	95% CONFIDENCE ERROR	INTERVAL	*	SAMPLE SIZE	STANDARD PERCENT	95% CONFIDENCE ERROR
					*			
ALABAMA	133	66.4	4.5	(57.6, 75.3)	*	307	60.1	3.0

ALASKA	68	45.8	8.7	(28.7, 63.0)	*	80	69.9	7.4	(55.4, 84.3)
ARIZONA	192	72.0	4.2	(63.8, 80.2)	*	276	73.5	3.5	(66.7, 80.4)
ARKANSAS	124	69.0	4.4	(60.3, 77.6)	*	270	55.7	3.3	(49.2, 62.1)
CALIFORNIA	286	68.6	3.0	(62.7, 74.5)	*	451	63.3	2.5	(58.4, 68.2)
COLORADO	121	75.4	4.8	(66.0, 84.7)	*	200	73.7	3.4	(67.2, 80.3)
CONNECTICUT	174	70.2	3.7	(62.9, 77.5)	*	248	65.3	3.5	(58.4, 72.2)
DELAWARE	190	69.9	3.6	(62.9, 76.9)	*	374	67.7	2.7	(62.3, 73.1)
DISTRICT OF CO	72	53.1	6.5	(40.3, 65.9)	*	170	55.0	4.3	(46.6, 63.4)
FLORIDA	374	61.6	2.7	(56.2, 67.0)	*	583	62.9	2.2	(58.5, 67.3)
GEORGIA	133	56.3	4.6	(47.3, 65.3)	*	212	59.9	3.7	(52.6, 67.2)
HAWAII	153	69.7	4.2	(61.5, 78.0)	*	239	72.3	3.4	(65.7, 78.9)
IDAHO	309	67.5	3.0	(61.7, 73.4)	*	625	65.6	2.2	(61.3, 69.9)
ILLINOIS	79	70.4	5.9	(59.0, 81.9)	*	175	66.1	4.0	(58.2, 74.0)
INDIANA	157	59.5	4.1	(51.5, 67.5)	*	268	64.3	3.5	(57.5, 71.2)
IOWA	254	72.7	3.0	(66.9, 78.6)	*	585	67.7	2.1	(63.6, 71.7)
KANSAS	123	58.4	4.8	(49.0, 67.8)	*	294	63.6	3.0	(57.7, 69.5)
KENTUCKY	247	62.9	3.4	(56.3, 69.5)	*	588	60.1	2.2	(55.8, 64.4)
LOUISIANA	104	54.8	5.2	(44.5, 65.0)	*	198	60.7	3.9	(53.0, 68.4)
MAINE	124	78.8	3.9	(71.2, 86.4)	*	187	67.4	3.8	(60.0, 74.8)
MARYLAND	312	67.6	3.4	(60.9, 74.4)	*	510	60.4	2.9	(54.8, 66.1)
MASSACHUSETTS	103	70.6	4.8	(61.2, 80.0)	*	184	62.9	3.9	(55.2, 70.5)
MICHIGAN	146	59.7	4.4	(51.1, 68.2)	*	268	66.3	3.1	(60.2, 72.4)
MINNESOTA	309	67.2	2.8	(61.8, 72.6)	*	609	70.2	2.1	(66.2, 74.2)
MISSISSIPPI	91	59.5	5.4	(48.8, 70.1)	*	235	62.2	3.2	(55.8, 68.5)
MISSOURI	128	71.4	4.3	(62.9, 79.9)	*	250	69.5	3.1	(63.3, 75.6)
MONTANA	143	69.1	4.3	(60.7, 77.6)	*	245	67.8	3.4	(61.2, 74.4)
NEBRASKA	219	68.1	3.4	(61.5, 74.7)	*	442	64.2	2.7	(59.0, 69.4)
NEVADA	144	56.4	8.7	(39.4, 73.4)	*	216	56.6	6.4	(44.2, 69.1)
NEW HAMPSHIRE	107	65.1	4.8	(55.7, 74.6)	*	159	64.3	4.2	(56.1, 72.5)
NEW JERSEY	174	63.8	4.0	(55.9, 71.6)	*	310	58.7	3.1	(52.5, 64.8)
NEW MEXICO	128	74.2	4.2	(66.0, 82.3)	*	219	71.8	3.1	(65.8, 77.9)
NEW YORK	198	72.9	3.5	(66.1, 79.7)	*	391	59.0	2.6	(54.0, 64.0)
NORTH CAROLINA	232	65.7	3.3	(59.3, 72.2)	*	504	63.8	2.4	(59.1, 68.5)
NORTH DAKOTA	170	67.2	3.8	(59.8, 74.6)	*	275	63.1	3.3	(56.7, 69.4)
OHIO	299	64.8	3.4	(58.2, 71.4)	*	526	65.8	2.7	(60.6, 70.9)
OKLAHOMA	311	73.0	2.9	(67.4, 78.5)	*	302	66.9	3.0	(61.0, 72.7)
OREGON	235	72.0	3.5	(65.2, 78.8)	*	439	68.3	2.6	(63.2, 73.3)
PENNSYLVANIA	254	66.8	3.2	(60.4, 73.1)	*	458	65.1	2.4	(60.3, 69.8)
RHODE ISLAND	130	70.5	4.3	(62.1, 78.9)	*	238	65.9	3.4	(59.2, 72.6)
SOUTH CAROLINA	149	78.1	3.5	(71.3, 85.0)	*	328	71.9	2.7	(66.6, 77.1)
SOUTH DAKOTA	199	60.1	3.7	(52.8, 67.4)	*	352	69.5	2.8	(64.0, 75.0)
TENNESSEE	157	69.0	3.9	(61.5, 76.6)	*	414	69.2	2.4	(64.5, 73.8)
TEXAS	123	72.2	4.2	(64.0, 80.5)	*	245	65.1	3.4	(58.6, 71.7)
UTAH	169	69.0	4.2	(60.7, 77.3)	*	258	63.9	3.9	(56.4, 71.4)
VERMONT	209	71.4	3.4	(64.8, 78.0)	*	362	68.1	2.6	(63.0, 73.3)
VIRGINIA	180	69.3	4.0	(61.4, 77.1)	*	378	66.5	2.9	(60.7, 72.2)
WASHINGTON	202	66.7	3.5	(59.9, 73.5)	*	385	72.9	2.4	(68.1, 77.6)
WEST VIRGINIA	179	57.7	3.9	(50.0, 65.4)	*	404	58.5	2.8	(53.1, 63.9)
WISCONSIN	130	62.8	4.8	(53.4, 72.1)	*	279	68.4	3.2	(62.1, 74.7)
WYOMING	135	72.9	4.1	(64.9, 81.0)	*	269	72.1	3.0	(66.2, 78.0)
PUERTO RICO	159	36.6	4.1	(28.6, 44.6)	*	250	45.4	3.5	(38.5, 52.3)

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 67.9
 RANGE = 36.6-78.8

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 65.5
 RANGE = 45.4-73.7

* HAD FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OVER

August 14, 1998

TABLE 23.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 FLU SHOT*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	45.1	62.6	17.5
ALASKA	49.8	58.3	8.5
ARIZONA	65.3	72.9	7.6
ARKANSAS	61.0	61.1	0.1
CALIFORNIA	60.0	65.5	5.5
COLORADO	66.7	74.4	7.7
CONNECTICUT	62.5	67.2	4.7
DELAWARE	57.2	68.6	11.4
DISTRICT OF CO	N/A	54.3	N/A
FLORIDA	61.7	62.3	0.7
GEORGIA	47.0	58.5	11.5
HAWAII	62.3	71.1	8.8
IDAHO	64.2	66.4	2.3
ILLINOIS	57.9	67.8	9.9
INDIANA	59.2	62.5	3.3
IOWA	63.7	69.7	6.1
KANSAS	62.2	61.5	-0.7
KENTUCKY	53.5	61.2	7.8
LOUISIANA	52.2	58.4	6.2
MAINE	64.5	72.1	7.5
MARYLAND	58.3	63.4	5.2
MASSACHUSETTS	59.3	66.0	6.7
MICHIGAN	56.7	63.6	6.8
MINNESOTA	63.2	69.0	5.7
MISSISSIPPI	57.0	61.1	4.1
MISSOURI	66.6	70.3	3.7
MONTANA	64.0	68.4	4.4
NEBRASKA	64.4	65.8	1.4
NEVADA	52.5	56.5	4.0
NEW HAMPSHIRE	56.0	64.6	8.7
NEW JERSEY	48.0	60.7	12.7
NEW MEXICO	69.0	72.8	3.8
NEW YORK	55.9	64.5	8.5
NORTH CAROLINA	52.5	64.6	12.0
NORTH DAKOTA	57.4	64.8	7.4
OHIO	63.0	65.4	2.4
OKLAHOMA	61.1	69.3	8.2
OREGON	67.0	69.8	2.9
PENNSYLVANIA	58.6	65.8	7.2
RHODE ISLAND	66.6	67.7	1.1
SOUTH CAROLINA	51.1	74.3	23.2
SOUTH DAKOTA	60.1	65.6	5.5
TENNESSEE	63.2	69.1	5.9
TEXAS	56.8	68.0	11.3
UTAH	70.2	66.2	-4.1
VERMONT	64.1	69.5	5.4
VIRGINIA	52.5	67.7	15.2
WASHINGTON	66.7	70.3	3.6
WEST VIRGINIA	53.2	58.2	5.0
WISCONSIN	56.9	66.1	9.1
WYOMING	66.8	72.4	5.6
PUERTO RICO	N/A	41.5	N/A

* HAD FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OVER
 N/A - NOT AVAILABLE

August 14, 1998

TABLE 24.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PNEUMONIA VACCINATION*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	430	47.5	2.6	(42.3, 52.6)
ALASKA	146	39.2	5.4	(28.6, 49.9)
ARIZONA	463	59.4	3.1	(53.4, 65.5)
ARKANSAS	378	39.1	2.8	(33.7, 44.4)
CALIFORNIA	724	49.8	2.1	(45.8, 53.9)
COLORADO	309	53.3	3.1	(47.2, 59.5)
CONNECTICUT	405	43.0	2.9	(37.4, 48.7)
DELAWARE	556	52.6	2.4	(47.9, 57.3)
DISTRICT OF CO	241	32.3	3.4	(25.6, 39.0)
FLORIDA	932	45.5	1.8	(42.0, 49.0)
GEORGIA	338	48.5	2.9	(42.7, 54.3)
HAWAII	364	51.7	3.0	(45.8, 57.7)
IDAHO	914	50.2	1.9	(46.5, 54.0)
ILLINOIS	252	44.7	3.4	(38.1, 51.3)
INDIANA	419	38.0	2.7	(32.7, 43.4)
IOWA	820	51.5	2.0	(47.6, 55.3)
KANSAS	414	43.7	2.7	(38.4, 49.0)
KENTUCKY	812	38.6	1.9	(34.8, 42.3)
LOUISIANA	294	32.2	3.0	(26.4, 38.1)
MAINE	302	50.0	2.9	(44.3, 55.7)
MARYLAND	810	41.0	2.2	(36.6, 45.4)
MASSACHUSETTS	283	52.7	3.2	(46.4, 59.0)
MICHIGAN	407	45.6	2.7	(40.4, 50.8)
MINNESOTA	896	48.3	1.8	(44.8, 51.8)
MISSISSIPPI	317	45.9	3.1	(39.9, 51.9)
MISSOURI	368	44.3	2.9	(38.6, 50.0)
MONTANA	376	50.8	3.0	(45.0, 56.7)
NEBRASKA	646	49.8	2.3	(45.4, 54.2)
NEVADA	354	53.5	5.4	(43.0, 64.0)
NEW HAMPSHIRE	255	49.6	3.3	(43.1, 56.1)
NEW JERSEY	467	34.0	2.4	(29.3, 38.6)
NEW MEXICO	341	50.1	2.9	(44.4, 55.8)
NEW YORK	569	38.9	2.3	(34.5, 43.4)
NORTH CAROLINA	723	50.7	2.0	(46.7, 54.6)
NORTH DAKOTA	431	40.8	2.5	(36.0, 45.7)
OHIO	809	38.6	2.3	(34.0, 43.1)
OKLAHOMA	582	40.4	2.2	(36.1, 44.7)
OREGON	656	55.9	2.2	(51.5, 60.2)
PENNSYLVANIA	687	47.1	2.1	(42.9, 51.2)
RHODE ISLAND	355	43.0	2.9	(37.4, 48.6)
SOUTH CAROLINA	467	41.6	2.5	(36.8, 46.4)
SOUTH DAKOTA	544	40.7	2.3	(36.1, 45.2)
TENNESSEE	561	45.0	2.3	(40.5, 49.5)
TEXAS	358	44.4	2.9	(38.8, 50.0)
UTAH	419	48.5	3.2	(42.3, 54.8)
VERMONT	559	51.6	2.3	(47.1, 56.1)
VIRGINIA	549	53.6	2.6	(48.5, 58.7)
WASHINGTON	571	51.6	2.3	(47.1, 56.1)
WEST VIRGINIA	572	41.3	2.2	(36.9, 45.6)
WISCONSIN	398	42.6	2.9	(36.9, 48.3)
WYOMING	398	50.9	2.8	(45.5, 56.2)
PUERTO RICO	395	33.7	2.6	(28.5, 38.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 45.8 RANGE = 32.2-59.4 YEAR 2000 OBJ. 20.11
 * EVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER August 14, 1998

TABLE 24.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PNEUMONIA VACCINATION* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	129	51.1	4.7	(41.9, 60.3)	*	301	45.2	3.2	(38.8, 51.5)	*
ALASKA	67	36.5	8.2	(20.4, 52.5)	*	79	41.8	7.8	(26.6, 57.0)	*
ARIZONA	189	60.4	4.6	(51.4, 69.4)	*	274	58.8	4.0	(50.9, 66.6)	*
ARKANSAS	112	40.5	5.0	(30.7, 50.3)	*	266	38.1	3.2	(31.9, 44.3)	*
CALIFORNIA	280	50.1	3.2	(43.7, 56.4)	*	444	49.7	2.7	(44.4, 54.9)	*
COLORADO	114	55.9	5.2	(45.6, 66.2)	*	195	51.6	3.9	(44.0, 59.1)	*
CONNECTICUT	165	40.2	4.3	(31.8, 48.6)	*	240	44.9	3.6	(37.8, 51.9)	*
DELAWARE	183	52.1	3.9	(44.5, 59.7)	*	373	53.0	3.0	(47.2, 58.8)	*
DISTRICT OF CO	73	32.2	5.9	(20.7, 43.7)	*	168	32.3	4.2	(24.1, 40.6)	*
FLORIDA	359	43.7	2.8	(38.1, 49.2)	*	573	46.8	2.3	(42.4, 51.2)	*
GEORGIA	129	44.8	4.5	(35.9, 53.6)	*	209	50.9	3.8	(43.5, 58.3)	*
HAWAII	143	49.0	4.7	(39.8, 58.2)	*	221	54.1	4.0	(46.3, 61.9)	*
IDAHO	298	49.7	3.2	(43.4, 56.0)	*	616	50.6	2.3	(46.0, 55.2)	*
ILLINOIS	78	43.7	6.2	(31.7, 55.8)	*	174	45.3	4.1	(37.2, 53.3)	*
INDIANA	155	41.7	4.2	(33.5, 49.8)	*	264	35.9	3.5	(29.0, 42.7)	*
IOWA	243	50.7	3.6	(43.6, 57.8)	*	577	52.0	2.3	(47.4, 56.5)	*
KANSAS	122	40.3	4.7	(31.0, 49.6)	*	292	46.0	3.1	(39.9, 52.2)	*
KENTUCKY	238	41.6	3.4	(34.9, 48.2)	*	574	36.6	2.2	(32.3, 40.9)	*
LOUISIANA	99	30.9	4.8	(21.5, 40.2)	*	195	33.1	3.8	(25.7, 40.5)	*
MAINE	120	47.1	4.7	(37.8, 56.4)	*	182	52.0	3.7	(44.7, 59.4)	*
MARYLAND	305	43.7	3.5	(36.8, 50.5)	*	505	39.1	2.8	(33.6, 44.6)	*
MASSACHUSETTS	102	58.2	5.2	(48.0, 68.4)	*	181	48.9	4.0	(41.1, 56.8)	*
MICHIGAN	144	42.1	4.4	(33.6, 50.6)	*	263	48.0	3.3	(41.5, 54.4)	*
MINNESOTA	297	47.3	3.1	(41.3, 53.3)	*	599	49.0	2.2	(44.7, 53.3)	*
MISSISSIPPI	86	44.9	5.7	(33.8, 56.0)	*	231	46.6	3.5	(39.8, 53.4)	*
MISSOURI	123	41.6	4.7	(32.4, 50.9)	*	245	46.0	3.7	(38.8, 53.2)	*
MONTANA	138	41.1	4.7	(31.9, 50.3)	*	238	58.2	3.6	(51.2, 65.1)	*
NEBRASKA	214	51.5	3.8	(44.1, 58.9)	*	432	48.7	2.8	(43.2, 54.1)	*
NEVADA	141	54.1	9.0	(36.4, 71.7)	*	213	53.1	6.4	(40.5, 65.7)	*
NEW HAMPSHIRE	103	53.5	5.1	(43.5, 63.5)	*	152	46.9	4.3	(38.5, 55.4)	*
NEW JERSEY	164	38.7	4.1	(30.7, 46.6)	*	303	30.9	2.9	(25.3, 36.5)	*
NEW MEXICO	122	49.1	4.8	(39.7, 58.4)	*	219	50.9	3.6	(43.9, 57.9)	*
NEW YORK	184	38.4	3.9	(30.8, 46.1)	*	385	39.2	2.8	(33.8, 44.7)	*
NORTH CAROLINA	224	50.3	3.6	(43.3, 57.3)	*	499	50.9	2.4	(46.1, 55.7)	*
NORTH DAKOTA	160	40.9	4.1	(32.7, 49.0)	*	271	40.8	3.2	(34.5, 47.0)	*
OHIO	290	37.5	3.7	(30.2, 44.7)	*	519	39.3	2.9	(33.6, 44.9)	*
OKLAHOMA	287	41.3	3.1	(35.3, 47.3)	*	295	39.8	3.0	(34.0, 45.7)	*
OREGON	227	57.8	3.7	(50.5, 65.0)	*	429	54.5	2.7	(49.2, 59.8)	*
PENNSYLVANIA	240	42.8	3.5	(35.9, 49.6)	*	447	49.7	2.6	(44.5, 54.9)	*
RHODE ISLAND	120	43.0	4.9	(33.5, 52.6)	*	235	43.1	3.5	(36.2, 49.9)	*
SOUTH CAROLINA	142	43.8	4.4	(35.2, 52.4)	*	325	40.2	3.0	(34.3, 46.1)	*
SOUTH DAKOTA	196	38.9	3.7	(31.6, 46.2)	*	348	41.9	3.0	(36.0, 47.8)	*
TENNESSEE	152	44.9	4.2	(36.6, 53.2)	*	409	45.1	2.6	(40.0, 50.2)	*
TEXAS	118	47.4	4.9	(37.8, 56.9)	*	240	42.5	3.5	(35.7, 49.2)	*
UTAH	163	47.4	5.1	(37.3, 57.4)	*	256	49.4	3.9	(41.7, 57.0)	*
VERMONT	199	48.1	3.8	(40.7, 55.5)	*	360	53.9	2.9	(48.3, 59.5)	*
VIRGINIA	177	53.6	4.8	(44.1, 63.0)	*	372	53.6	3.2	(47.2, 60.0)	*
WASHINGTON	193	45.2	3.8	(37.8, 52.6)	*	378	56.1	2.8	(50.5, 61.6)	*

WEST VIRGINIA	174	39.8	4.1	(31.7, 47.8)	*	398	42.3	2.6	(37.2, 47.3)
WISCONSIN	127	36.0	4.8	(26.7, 45.3)	*	271	47.1	3.6	(40.1, 54.2)
WYOMING	131	45.1	4.6	(36.0, 54.1)	*	267	55.1	3.3	(48.6, 61.6)
PUERTO RICO	151	31.6	4.0	(23.8, 39.4)	*	244	35.3	3.5	(28.5, 42.1)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 44.3	MEDIAN	= 46.9
	RANGE	= 30.9-60.4	RANGE	= 30.9-58.8

* EVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER

August 14, 1998

TABLE 24.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1995 TO 1997
 PNEUMONIA VACCINATION*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	1995 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	33.2	47.5	14.3
ALASKA	45.9	39.2	-6.7
ARIZONA	49.2	59.4	10.2
ARKANSAS	37.1	39.1	1.9
CALIFORNIA	44.3	49.8	5.5
COLORADO	46.6	53.3	6.7
CONNECTICUT	38.3	43.0	4.8
DELAWARE	41.5	52.6	11.1
DISTRICT OF CO	N/A	32.3	N/A
FLORIDA	39.5	45.5	6.0
GEORGIA	39.6	48.5	8.9
HAWAII	43.0	51.7	8.8
IDAHO	40.3	50.2	10.0
ILLINOIS	28.9	44.7	15.8
INDIANA	34.1	38.0	3.9
IOWA	44.9	51.5	6.6
KANSAS	44.7	43.7	-1.0
KENTUCKY	25.3	38.6	13.3
LOUISIANA	26.0	32.2	6.3
MAINE	35.6	50.0	14.5
MARYLAND	33.6	41.0	7.4
MASSACHUSETTS	32.4	52.7	20.3
MICHIGAN	39.9	45.6	5.7
MINNESOTA	40.1	48.3	8.2
MISSISSIPPI	39.4	45.9	6.5
MISSOURI	32.2	44.3	12.1
MONTANA	34.9	50.8	15.9
NEBRASKA	36.1	49.8	13.8
NEVADA	40.3	53.5	13.3
NEW HAMPSHIRE	40.5	49.6	9.1
NEW JERSEY	13.0	34.0	20.9
NEW MEXICO	39.6	50.1	10.6
NEW YORK	26.2	38.9	12.8
NORTH CAROLINA	31.3	50.7	19.4
NORTH DAKOTA	33.3	40.8	7.6
OHIO	40.7	38.6	-2.2
OKLAHOMA	37.2	40.4	3.2
OREGON	45.9	55.9	10.0

PENNSYLVANIA	38.5	47.1	8.6
RHODE ISLAND	30.8	43.0	12.2
SOUTH CAROLINA	26.5	41.6	15.1
SOUTH DAKOTA	31.5	40.7	9.1
TENNESSEE	29.9	45.0	15.1
TEXAS	44.8	44.4	-0.3
UTAH	42.7	48.5	5.8
VERMONT	36.0	51.6	15.7
VIRGINIA	39.5	53.6	14.0
WASHINGTON	46.0	51.6	5.5
WEST VIRGINIA	37.0	41.3	4.3
WISCONSIN	35.8	42.6	6.8
WYOMING	43.8	50.9	7.0
PUERTO RICO	N/A	33.7	N/A

* EVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER

N/A - NOT AVAILABLE

August 14, 1998

TABLE 25.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SIGMOIDOSCOPY*
DENOMINATOR IS PERSON AGE 50 AND OVER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	892	43.0	1.8	(39.5, 46.6)
ALASKA	439	41.0	3.6	(34.0, 48.0)
ARIZONA	862	39.2	2.4	(34.6, 43.9)
ARKANSAS	740	34.4	1.9	(30.7, 38.1)
CALIFORNIA	1493	46.6	1.5	(43.7, 49.5)
COLORADO	667	43.5	2.1	(39.4, 47.6)
CONNECTICUT	811	45.0	2.1	(41.0, 49.0)
DELAWARE	1064	47.8	1.8	(44.4, 51.3)
DISTRICT OF CO	497	49.2	2.6	(44.0, 54.3)
FLORIDA	1579	45.8	1.4	(43.0, 48.5)
GEORGIA	737	48.1	2.1	(43.9, 52.2)
HAWAII	783	51.6	2.2	(47.4, 55.9)
IDAHO	1895	38.4	1.4	(35.7, 41.1)
ILLINOIS	516	40.3	2.4	(35.6, 44.9)
INDIANA	881	38.5	1.8	(34.9, 42.1)
IOWA	1530	40.7	1.4	(38.0, 43.4)
KANSAS	772	37.5	1.9	(33.8, 41.2)
KENTUCKY	1555	34.2	1.4	(31.6, 36.8)
LOUISIANA	616	37.0	2.1	(32.8, 41.2)
MAINE	669	42.4	2.1	(38.3, 46.4)
MARYLAND	1688	37.1	1.6	(34.0, 40.2)
MASSACHUSETTS	560	41.3	2.4	(36.7, 45.9)
MICHIGAN	876	47.8	1.8	(44.2, 51.3)
MINNESOTA	1796	50.9	1.3	(48.4, 53.5)
MISSISSIPPI	660	35.8	2.0	(31.8, 39.8)
MISSOURI	740	41.1	2.1	(36.9, 45.2)
MONTANA	738	39.7	1.9	(36.0, 43.3)
NEBRASKA	1141	38.2	1.7	(34.8, 41.5)
NEVADA	903	42.3	3.3	(35.8, 48.7)

NEW HAMPSHIRE	511	42.8	2.3	(38.2, 47.3)
NEW JERSEY	941	39.6	1.8	(35.9, 43.2)
NEW MEXICO	684	39.4	2.0	(35.5, 43.4)
NEW YORK	1187	43.7	1.6	(40.5, 46.8)
NORTH CAROLINA	1382	39.2	1.5	(36.3, 42.1)
NORTH DAKOTA	759	44.9	2.0	(40.9, 48.8)
OHIO	1468	37.5	1.6	(34.3, 40.7)
OKLAHOMA	940	22.3	1.6	(19.1, 25.4)
OREGON	1299	46.1	1.6	(43.1, 49.2)
PENNSYLVANIA	1370	40.2	1.5	(37.3, 43.1)
RHODE ISLAND	697	42.6	2.1	(38.6, 46.7)
SOUTH CAROLINA	889	29.2	1.7	(26.0, 32.4)
SOUTH DAKOTA	886	37.6	1.9	(33.9, 41.3)
TENNESSEE	1171	34.2	1.6	(31.1, 37.3)
TEXAS	803	38.2	1.9	(34.5, 42.0)
UTAH	913	45.1	2.3	(40.6, 49.5)
VERMONT	1161	40.2	1.6	(37.1, 43.3)
VIRGINIA	1281	47.6	2.0	(43.7, 51.5)
WASHINGTON	1273	44.0	1.5	(41.0, 47.0)
WEST VIRGINIA	1106	34.4	1.5	(31.5, 37.4)
WISCONSIN	814	50.3	2.1	(46.2, 54.3)
WYOMING	884	45.3	1.8	(41.7, 48.8)
PUERTO RICO	928	28.4	1.6	(25.2, 31.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 40.8 RANGE = 22.3-51.6 YEAR 2000 OBJ 16.13
 * EVER HAD A SIGMOIDOSCOPY AGE 50 AND OVER August 17, 1998

TABLE 25.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SIGMOIDOSCOPY* -BY SEX
 DENOMINATOR IS PERSON AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	308	45.8	2.9	(40.0, 51.6)	*	584	41.0	2.2	(36.6, 45.4)
ALASKA	214	42.4	5.1	(32.3, 52.5)	*	225	39.6	4.9	(30.0, 49.2)
ARIZONA	380	41.7	3.6	(34.6, 48.8)	*	482	37.3	3.3	(30.9, 43.7)
ARKANSAS	255	38.5	3.2	(32.1, 44.8)	*	485	31.3	2.2	(26.9, 35.6)
CALIFORNIA	615	49.6	2.3	(45.1, 54.0)	*	878	44.3	1.9	(40.5, 48.1)
COLORADO	277	48.1	3.3	(41.7, 54.6)	*	390	39.5	2.7	(34.2, 44.9)
CONNECTICUT	326	48.0	3.4	(41.3, 54.7)	*	485	42.6	2.5	(37.7, 47.5)
DELAWARE	378	52.2	3.0	(46.5, 58.0)	*	686	44.2	2.2	(39.8, 48.6)
DISTRICT OF CO	178	52.5	4.1	(44.4, 60.6)	*	319	46.7	3.1	(40.6, 52.8)
FLORIDA	634	49.9	2.2	(45.6, 54.2)	*	945	42.5	1.8	(39.0, 45.9)
GEORGIA	295	51.6	3.4	(45.0, 58.2)	*	442	45.3	2.6	(40.1, 50.4)
HAWAII	333	58.1	3.1	(52.0, 64.1)	*	450	45.7	2.9	(40.0, 51.5)
IDAHO	728	41.0	2.2	(36.8, 45.2)	*	1167	36.2	1.8	(32.6, 39.7)
ILLINOIS	187	45.0	3.9	(37.3, 52.7)	*	329	36.8	2.9	(31.2, 42.4)
INDIANA	386	46.1	3.0	(40.3, 51.9)	*	495	32.3	2.2	(27.9, 36.7)
IOWA	538	44.9	2.3	(40.4, 49.5)	*	992	37.4	1.7	(34.1, 40.7)
KANSAS	284	41.0	3.1	(35.0, 47.1)	*	488	34.5	2.3	(29.9, 39.0)
KENTUCKY	515	36.8	2.3	(32.3, 41.3)	*	1040	32.2	1.6	(29.0, 35.3)
LOUISIANA	227	38.6	3.5	(31.7, 45.5)	*	389	35.9	2.7	(30.5, 41.2)
MAINE	285	44.3	3.1	(38.3, 50.3)	*	384	40.8	2.8	(35.3, 46.3)

MARYLAND	703	42.9	2.5	(38.1, 47.7)	*	985	32.2	2.0	(28.4, 36.1)
MASSACHUSETTS	213	49.8	3.8	(42.3, 57.3)	*	347	34.6	2.8	(29.0, 40.2)
MICHIGAN	347	46.3	2.9	(40.6, 51.9)	*	529	49.0	2.4	(44.4, 53.6)
MINNESOTA	746	55.0	1.9	(51.2, 58.8)	*	1050	47.6	1.7	(44.3, 50.8)
MISSISSIPPI	220	35.0	3.5	(28.0, 41.9)	*	440	36.5	2.4	(31.7, 41.2)
MISSOURI	275	41.2	3.5	(34.4, 47.9)	*	465	41.0	2.6	(36.0, 46.0)
MONTANA	287	38.1	3.0	(32.2, 44.0)	*	451	41.0	2.4	(36.2, 45.8)
NEBRASKA	397	41.9	2.8	(36.3, 47.5)	*	744	35.1	2.0	(31.2, 39.0)
NEVADA	399	53.8	5.0	(44.1, 63.5)	*	504	32.2	3.7	(24.9, 39.5)
NEW HAMPSHIRE	203	46.8	3.6	(39.7, 53.9)	*	308	39.3	2.9	(33.5, 45.0)
NEW JERSEY	358	46.0	3.1	(40.0, 52.0)	*	583	34.5	2.2	(30.2, 38.8)
NEW MEXICO	274	44.5	3.3	(38.0, 50.9)	*	410	35.3	2.6	(30.3, 40.3)
NEW YORK	442	50.8	2.7	(45.5, 56.1)	*	745	38.3	1.9	(34.5, 42.0)
NORTH CAROLINA	518	40.8	2.4	(36.1, 45.5)	*	864	37.9	1.8	(34.4, 41.5)
NORTH DAKOTA	301	48.5	3.2	(42.3, 54.7)	*	458	41.8	2.5	(36.8, 46.7)
OHIO	552	43.0	2.6	(37.9, 48.1)	*	916	33.2	2.2	(29.0, 37.4)
OKLAHOMA	457	19.0	2.4	(14.4, 23.6)	*	483	24.9	2.3	(20.5, 29.3)
OREGON	499	48.3	2.5	(43.4, 53.1)	*	800	44.3	2.0	(40.4, 48.2)
PENNSYLVANIA	499	47.0	2.5	(42.1, 51.8)	*	871	35.1	1.8	(31.6, 38.5)
RHODE ISLAND	278	47.9	3.3	(41.5, 54.3)	*	419	38.6	2.6	(33.5, 43.7)
SOUTH CAROLINA	317	32.9	2.8	(27.4, 38.3)	*	572	26.3	1.9	(22.6, 30.1)
SOUTH DAKOTA	343	42.9	3.0	(37.1, 48.8)	*	543	33.5	2.3	(28.9, 38.0)
TENNESSEE	373	33.0	2.6	(27.9, 38.0)	*	798	35.2	1.9	(31.6, 38.8)
TEXAS	301	39.8	3.1	(33.8, 45.8)	*	502	37.0	2.4	(32.3, 41.6)
UTAH	397	50.4	3.4	(43.8, 56.9)	*	516	40.5	3.0	(34.7, 46.4)
VERMONT	492	45.0	2.4	(40.2, 49.7)	*	669	36.3	2.1	(32.2, 40.3)
VIRGINIA	450	52.9	3.2	(46.6, 59.2)	*	831	43.4	2.5	(38.5, 48.3)
WASHINGTON	510	45.7	2.4	(41.0, 50.3)	*	763	42.6	2.0	(38.8, 46.5)
WEST VIRGINIA	397	36.2	2.5	(31.3, 41.1)	*	709	33.1	1.9	(29.3, 36.8)
WISCONSIN	323	54.3	3.2	(48.0, 60.6)	*	491	46.9	2.6	(41.8, 52.0)
WYOMING	334	45.8	2.9	(40.1, 51.5)	*	550	44.9	2.3	(40.4, 49.3)
PUERTO RICO	367	33.1	2.6	(28.0, 38.2)	*	561	24.6	2.0	(20.8, 28.5)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 45.3
 RANGE = 19.0-58.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 37.7
 RANGE = 24.6-49.0

* EVER HAD A SIGMOIDOSCOPY AGE 50 AND OVER

August 17, 1998

TABLE 26.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HOME BLOOD STOOL TEST*
 DENOMINATOR IS PERSON AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	889	80.8	1.5	(77.8, 83.7)
ALASKA	440	75.0	3.2	(68.8, 81.2)
ARIZONA	860	70.7	2.2	(66.3, 75.1)
ARKANSAS	741	78.7	1.7	(75.4, 81.9)
CALIFORNIA	1502	75.2	1.2	(72.8, 77.6)
COLORADO	670	67.8	2.0	(63.9, 71.6)
CONNECTICUT	806	68.9	1.8	(65.4, 72.5)
DELAWARE	1060	69.5	1.7	(66.2, 72.9)
DISTRICT OF CO	503	62.2	2.6	(57.2, 67.2)
FLORIDA	1575	69.9	1.3	(67.4, 72.4)
GEORGIA	744	72.2	1.8	(68.6, 75.8)

HAWAII	784	67.8	2.0	(63.8, 71.8)
IDAHO	1905	74.6	1.3	(72.2, 77.1)
ILLINOIS	514	75.1	2.0	(71.1, 79.1)
INDIANA	884	77.8	1.7	(74.5, 81.1)
IOWA	1536	72.0	1.4	(69.3, 74.7)
KANSAS	756	68.5	1.9	(64.9, 72.2)
KENTUCKY	1544	73.9	1.3	(71.4, 76.4)
LOUISIANA	616	76.7	1.9	(73.0, 80.4)
MAINE	668	65.0	1.9	(61.2, 68.8)
MARYLAND	1674	67.9	1.5	(65.0, 70.8)
MASSACHUSETTS	559	65.8	2.3	(61.4, 70.2)
MICHIGAN	870	69.0	1.7	(65.6, 72.4)
MINNESOTA	1783	67.7	1.2	(65.3, 70.1)
MISSISSIPPI	667	86.5	1.4	(83.8, 89.3)
MISSOURI	748	74.9	1.8	(71.5, 78.4)
MONTANA	730	76.0	1.8	(72.5, 79.4)
NEBRASKA	1139	75.6	1.5	(72.7, 78.5)
NEVADA	905	84.0	2.5	(79.1, 89.0)
NEW HAMPSHIRE	525	66.1	2.4	(61.5, 70.7)
NEW JERSEY	935	71.5	1.7	(68.2, 74.8)
NEW MEXICO	682	78.3	1.8	(74.8, 81.7)
NEW YORK	1178	68.1	1.5	(65.1, 71.1)
NORTH CAROLINA	1375	63.5	1.5	(60.6, 66.3)
NORTH DAKOTA	759	77.3	1.7	(74.0, 80.6)
OHIO	1463	70.6	1.6	(67.5, 73.7)
OKLAHOMA	939	85.7	1.3	(83.1, 88.3)
OREGON	1312	65.4	1.5	(62.5, 68.3)
PENNSYLVANIA	1373	71.2	1.4	(68.4, 73.9)
RHODE ISLAND	696	73.5	1.8	(70.0, 77.1)
SOUTH CAROLINA	882	76.0	1.5	(73.0, 78.9)
SOUTH DAKOTA	889	78.3	1.5	(75.4, 81.3)
TENNESSEE	1189	77.7	1.4	(75.0, 80.4)
TEXAS	805	75.0	1.7	(71.7, 78.3)
UTAH	920	78.0	1.9	(74.2, 81.8)
VERMONT	1161	64.9	1.6	(61.8, 67.9)
VIRGINIA	1255	74.1	1.9	(70.4, 77.8)
WASHINGTON	1273	64.1	1.5	(61.2, 66.9)
WEST VIRGINIA	1099	82.0	1.3	(79.5, 84.5)
WISCONSIN	820	74.6	1.8	(71.0, 78.1)
WYOMING	892	80.7	1.4	(77.9, 83.5)
PUERTO RICO	932	77.1	1.5	(74.1, 80.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 74.0 RANGE = 62.2-86.5 YEAR 2000 OBJ 16.13
 * DID NOT HAVE A BLOOD STOOL TEST IN PAST 2 YEARS, AGE 50 AND OVER August 17, 1998

TABLE 26.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HOME BLOOD STOOL TEST* -BY SEX
 DENOMINATOR IS PERSON AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-MALES-				-FEMALES-				
	SAMPLE SIZE	PERCENT	STANDARD	95% CONFIDENCE	* SAMPLE SIZE	PERCENT	STANDARD	95% CONFIDENCE	
			ERROR	INTERVAL			*	INTERVAL	
ALABAMA	308	81.5	2.5	(76.6, 86.3)	*	581	80.2	1.8	(76.7, 83.7)
ALASKA	214	76.7	4.6	(67.7, 85.6)	*	226	73.3	4.4	(64.7, 81.9)

ARIZONA	378	67.9	3.2	(61.6, 74.2)	*	482	73.0	3.0	(67.2, 78.8)
ARKANSAS	258	76.6	2.8	(71.1, 82.0)	*	483	80.3	2.0	(76.4, 84.2)
CALIFORNIA	619	80.5	1.7	(77.2, 83.8)	*	883	70.9	1.7	(67.6, 74.2)
COLORADO	278	68.6	3.1	(62.6, 74.7)	*	392	67.0	2.6	(61.8, 72.2)
CONNECTICUT	322	70.6	2.9	(64.9, 76.3)	*	484	67.6	2.4	(62.9, 72.2)
DELAWARE	380	67.7	2.8	(62.1, 73.2)	*	680	71.1	2.0	(67.1, 75.1)
DISTRICT OF CO	180	66.8	4.1	(58.8, 74.9)	*	323	58.9	3.2	(52.5, 65.2)
FLORIDA	638	73.3	1.9	(69.7, 77.0)	*	937	67.2	1.7	(63.9, 70.5)
GEORGIA	294	73.7	2.9	(67.9, 79.4)	*	450	71.0	2.3	(66.4, 75.6)
HAWAII	333	72.0	2.9	(66.3, 77.7)	*	451	63.9	2.8	(58.4, 69.5)
IDAHO	736	79.7	1.9	(76.0, 83.3)	*	1169	70.2	1.7	(66.9, 73.6)
ILLINOIS	187	77.0	3.3	(70.7, 83.4)	*	327	73.7	2.6	(68.7, 78.7)
INDIANA	386	80.3	2.6	(75.1, 85.5)	*	498	75.8	2.2	(71.5, 80.1)
IOWA	542	73.1	2.2	(68.9, 77.4)	*	994	71.2	1.7	(67.9, 74.4)
KANSAS	279	68.6	3.0	(62.7, 74.4)	*	477	68.5	2.3	(64.0, 73.0)
KENTUCKY	509	77.0	2.0	(73.0, 80.9)	*	1035	71.5	1.6	(68.4, 74.6)
LOUISIANA	228	76.9	3.0	(71.0, 82.7)	*	388	76.6	2.3	(72.1, 81.1)
MAINE	285	69.7	2.8	(64.3, 75.2)	*	383	61.0	2.7	(55.7, 66.2)
MARYLAND	696	68.5	2.2	(64.1, 72.9)	*	978	67.4	1.9	(63.6, 71.2)
MASSACHUSETTS	214	72.8	3.4	(66.2, 79.3)	*	345	60.1	3.0	(54.3, 66.0)
MICHIGAN	345	72.9	2.6	(67.7, 78.0)	*	525	65.8	2.3	(61.4, 70.2)
MINNESOTA	740	69.6	1.9	(66.0, 73.3)	*	1043	66.1	1.6	(62.9, 69.2)
MISSISSIPPI	226	92.1	1.7	(88.8, 95.4)	*	441	82.1	2.1	(78.1, 86.2)
MISSOURI	282	78.1	2.7	(72.7, 83.5)	*	466	72.5	2.4	(67.8, 77.1)
MONTANA	288	80.0	2.5	(75.1, 84.9)	*	442	72.4	2.4	(67.8, 77.0)
NEBRASKA	399	78.4	2.4	(73.8, 83.0)	*	740	73.3	1.9	(69.6, 77.0)
NEVADA	401	83.6	4.5	(74.9, 92.4)	*	504	84.4	2.6	(79.3, 89.5)
NEW HAMPSHIRE	210	71.4	3.8	(64.0, 78.7)	*	315	61.6	2.9	(55.8, 67.3)
NEW JERSEY	358	73.0	2.6	(67.9, 78.1)	*	577	70.3	2.2	(66.0, 74.6)
NEW MEXICO	273	82.1	2.6	(77.0, 87.1)	*	409	75.1	2.4	(70.4, 79.9)
NEW YORK	438	67.5	2.5	(62.6, 72.5)	*	740	68.6	1.9	(64.8, 72.4)
NORTH CAROLINA	514	67.2	2.3	(62.8, 71.7)	*	861	60.5	1.9	(56.8, 64.1)
NORTH DAKOTA	303	83.4	2.3	(78.8, 88.0)	*	456	72.1	2.4	(67.4, 76.7)
OHIO	550	72.4	2.5	(67.5, 77.3)	*	913	69.2	2.0	(65.2, 73.1)
OKLAHOMA	456	89.3	1.8	(85.8, 92.9)	*	483	82.8	1.9	(79.1, 86.5)
OREGON	504	74.6	2.2	(70.3, 78.9)	*	808	57.6	2.0	(53.7, 61.5)
PENNSYLVANIA	495	72.2	2.3	(67.7, 76.7)	*	878	70.4	1.7	(67.1, 73.7)
RHODE ISLAND	277	76.0	2.8	(70.5, 81.4)	*	419	71.7	2.4	(67.0, 76.3)
SOUTH CAROLINA	314	78.8	2.4	(74.1, 83.6)	*	568	73.8	1.9	(70.0, 77.6)
SOUTH DAKOTA	347	81.7	2.3	(77.2, 86.1)	*	542	75.7	2.1	(71.6, 79.8)
TENNESSEE	376	81.8	2.2	(77.5, 86.2)	*	813	74.4	1.7	(71.1, 77.7)
TEXAS	297	75.4	2.7	(70.0, 80.8)	*	508	74.6	2.1	(70.5, 78.8)
UTAH	403	79.9	2.8	(74.5, 85.3)	*	517	76.4	2.6	(71.3, 81.5)
VERMONT	495	69.1	2.2	(64.7, 73.5)	*	666	61.4	2.1	(57.2, 65.5)
VIRGINIA	442	74.5	3.3	(68.1, 80.9)	*	813	73.8	2.1	(69.6, 78.0)
WASHINGTON	513	68.2	2.2	(63.8, 72.5)	*	760	60.6	2.0	(56.7, 64.4)
WEST VIRGINIA	399	83.9	2.0	(80.0, 87.7)	*	700	80.5	1.7	(77.2, 83.7)
WISCONSIN	325	79.2	2.5	(74.3, 84.2)	*	495	70.8	2.5	(65.9, 75.6)
WYOMING	337	83.1	2.2	(78.8, 87.4)	*	555	78.7	1.9	(75.0, 82.4)
PUERTO RICO	370	80.3	2.2	(76.1, 84.6)	*	562	74.5	2.1	(70.3, 78.6)

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 75.7
 RANGE = 66.8-92.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 71.3
 RANGE = 57.6-84.4

* DID NOT HAVE A BLOOD STOOL TEST IN PAST 2 YEARS, AGE 50 AND OVER

August 17, 1998

TABLE 27.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	SAMPLE PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2092	32.7	1.2	(30.4, 35.0)
ALASKA	1483	34.1	1.8	(30.6, 37.7)
ARIZONA	1861	25.4	1.4	(22.7, 28.1)
ARKANSAS	1702	32.1	1.4	(29.3, 34.9)
CALIFORNIA	4001	29.1	0.8	(27.5, 30.8)
COLORADO	1783	25.1	1.2	(22.9, 27.4)
CONNECTICUT	2110	29.1	1.3	(26.6, 31.6)
DELAWARE	2461	32.5	1.1	(30.4, 34.6)
DISTRICT OF CO	1489	28.0	1.4	(25.3, 30.7)
FLORIDA	3407	30.4	0.9	(28.6, 32.1)
GEORGIA	2246	30.8	1.2	(28.5, 33.0)
HAWAII	2143	26.6	1.2	(24.3, 28.8)
IDAHO	4802	29.6	0.8	(28.0, 31.2)
ILLINOIS	2781	32.0	1.0	(30.0, 33.9)
INDIANA	2310	34.6	1.1	(32.4, 36.9)
IOWA	3530	33.3	0.9	(31.6, 35.1)
KANSAS	1852	31.5	1.2	(29.2, 33.9)
KENTUCKY	3506	35.6	0.9	(33.8, 37.5)
LOUISIANA	1594	33.5	1.4	(30.8, 36.1)
MAINE	1634	30.2	1.3	(27.8, 32.7)
MARYLAND	4325	31.2	0.9	(29.4, 33.0)
MASSACHUSETTS	1652	25.5	1.3	(23.0, 27.9)
MICHIGAN	2499	34.9	1.1	(32.8, 36.9)
MINNESOTA	4723	29.7	0.7	(28.3, 31.2)
MISSISSIPPI	1539	35.1	1.5	(32.3, 38.0)
MISSOURI	1810	31.6	1.3	(29.0, 34.1)
MONTANA	1743	27.9	1.2	(25.5, 30.3)
NEBRASKA	2604	31.1	1.1	(28.9, 33.2)
NEVADA	2441	26.2	1.8	(22.7, 29.6)
NEW HAMPSHIRE	1464	27.5	1.4	(24.9, 30.2)
NEW JERSEY	2511	28.2	1.1	(26.1, 30.3)
NEW MEXICO	1755	29.8	1.2	(27.4, 32.1)
NEW YORK	3237	28.4	0.9	(26.6, 30.1)
NORTH CAROLINA	3393	32.4	0.9	(30.6, 34.1)
NORTH DAKOTA	1771	32.9	1.2	(30.6, 35.2)
OHIO	3011	32.6	1.1	(30.4, 34.8)
OKLAHOMA	1813	28.4	1.2	(26.0, 30.8)
OREGON	3237	32.4	1.0	(30.4, 34.3)
PENNSYLVANIA	3400	32.2	0.9	(30.4, 34.0)
RHODE ISLAND	1770	28.1	1.2	(25.7, 30.4)
SOUTH CAROLINA	2096	31.8	1.2	(29.5, 34.1)
SOUTH DAKOTA	2132	31.1	1.1	(28.9, 33.3)
TENNESSEE	2857	31.6	1.0	(29.6, 33.6)
TEXAS	2394	32.8	1.1	(30.7, 34.9)
UTAH	2808	27.0	1.1	(24.8, 29.1)
VERMONT	3032	29.9	1.0	(27.9, 31.8)
VIRGINIA	3380	30.0	1.1	(27.9, 32.2)
WASHINGTON	3405	29.1	0.9	(27.3, 30.9)
WEST VIRGINIA	2347	36.3	1.1	(34.2, 38.3)
WISCONSIN	2204	32.8	1.2	(30.5, 35.1)
WYOMING	2360	27.9	1.0	(26.0, 29.9)
PUERTO RICO	2183	34.9	1.1	(32.7, 37.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 31.1 RANGE = 25.1-36.3 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES

August 17, 1998

TABLE 27.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-MALES-					-FEMALES-				
	SAMPLE	STANDARD	95% CONFIDENCE	*	SAMPLE	STANDARD	95% CONFIDENCE	*		
	SIZE	PERCENT	ERROR	INTERVAL	SIZE	PERCENT	ERROR	INTERVAL		
ALABAMA	803	31.7	1.8	(28.2, 35.1)	*	1289	33.7	1.4	(30.8, 36.5)	
ALASKA	708	30.8	2.5	(25.9, 35.8)	*	775	38.0	2.6	(32.9, 43.0)	
ARIZONA	874	26.1	2.1	(22.0, 30.1)	*	987	24.7	1.9	(21.1, 28.4)	
ARKANSAS	659	31.5	2.3	(27.1, 35.9)	*	1043	32.7	1.8	(29.2, 36.2)	
CALIFORNIA	1746	29.5	1.3	(27.0, 32.0)	*	2255	28.8	1.1	(26.6, 30.9)	
COLORADO	809	26.2	1.7	(22.8, 29.5)	*	974	24.1	1.6	(21.0, 27.3)	
CONNECTICUT	879	32.1	2.1	(28.1, 36.2)	*	1231	26.0	1.5	(23.1, 28.9)	
DELAWARE	997	33.6	1.7	(30.3, 36.9)	*	1464	31.4	1.4	(28.7, 34.1)	
DISTRICT OF CO	570	25.0	2.1	(21.0, 29.1)	*	919	30.6	1.8	(27.1, 34.0)	
FLORIDA	1425	33.7	1.4	(30.9, 36.5)	*	1982	27.2	1.1	(25.1, 29.4)	
GEORGIA	951	33.2	1.8	(29.7, 36.7)	*	1295	28.5	1.5	(25.6, 31.3)	
HAWAII	945	28.4	1.7	(25.1, 31.8)	*	1198	24.7	1.5	(21.7, 27.7)	
IDAHO	2042	30.4	1.2	(28.0, 32.8)	*	2760	28.7	1.1	(26.5, 30.9)	
ILLINOIS	1145	32.2	1.5	(29.2, 35.2)	*	1636	31.8	1.3	(29.2, 34.4)	
INDIANA	1060	36.2	1.7	(32.8, 39.5)	*	1250	33.2	1.5	(30.3, 36.2)	
IOWA	1465	34.5	1.4	(31.7, 37.2)	*	2065	32.3	1.2	(30.0, 34.6)	
KANSAS	765	38.3	1.9	(34.5, 42.1)	*	1087	24.6	1.4	(21.9, 27.4)	
KENTUCKY	1310	37.8	1.5	(34.9, 40.8)	*	2196	33.6	1.1	(31.4, 35.8)	
LOUISIANA	626	33.4	2.2	(29.2, 37.7)	*	968	33.5	1.7	(30.2, 36.8)	
MAINE	740	33.2	2.0	(29.3, 37.0)	*	894	27.4	1.6	(24.3, 30.5)	
MARYLAND	1909	30.1	1.3	(27.4, 32.7)	*	2416	32.4	1.3	(29.9, 34.8)	
MASSACHUSETTS	708	30.3	1.9	(26.5, 34.1)	*	944	20.8	1.6	(17.7, 23.8)	
MICHIGAN	1065	36.3	1.6	(33.1, 39.4)	*	1434	33.5	1.4	(30.8, 36.2)	
MINNESOTA	2192	31.3	1.1	(29.1, 33.4)	*	2531	28.3	1.0	(26.4, 30.2)	
MISSISSIPPI	586	34.7	2.4	(30.1, 39.3)	*	953	35.5	1.8	(32.1, 38.9)	
MISSOURI	694	31.2	2.1	(27.2, 35.2)	*	1116	31.9	1.7	(28.6, 35.2)	
MONTANA	750	28.9	1.8	(25.3, 32.4)	*	993	26.9	1.6	(23.9, 30.0)	
NEBRASKA	1034	35.0	1.8	(31.5, 38.5)	*	1570	27.3	1.3	(24.8, 29.8)	
NEVADA	1160	28.6	2.8	(23.1, 34.0)	*	1281	23.6	2.1	(19.4, 27.8)	
NEW HAMPSHIRE	602	34.1	2.2	(29.8, 38.5)	*	862	21.0	1.5	(18.1, 24.0)	
NEW JERSEY	1030	29.1	1.6	(25.9, 32.3)	*	1481	27.5	1.4	(24.7, 30.2)	
NEW MEXICO	741	30.4	1.8	(26.8, 33.9)	*	1014	29.1	1.6	(26.1, 32.2)	
NEW YORK	1326	29.4	1.4	(26.6, 32.2)	*	1911	27.4	1.1	(25.1, 29.6)	
NORTH CAROLINA	1394	34.6	1.4	(31.8, 37.4)	*	1999	30.2	1.2	(28.0, 32.5)	
NORTH DAKOTA	799	34.9	1.8	(31.4, 38.4)	*	972	30.9	1.6	(27.8, 34.0)	
OHIO	1215	32.1	1.7	(28.7, 35.4)	*	1796	33.1	1.5	(30.3, 36.0)	
OKLAHOMA	816	28.0	1.9	(24.3, 31.7)	*	997	28.7	1.6	(25.5, 31.9)	
OREGON	1392	32.2	1.5	(29.2, 35.2)	*	1845	32.5	1.3	(30.1, 35.0)	
PENNSYLVANIA	1416	35.0	1.4	(32.2, 37.8)	*	1984	29.6	1.1	(27.3, 31.8)	
RHODE ISLAND	746	30.4	1.9	(26.7, 34.0)	*	1024	25.9	1.5	(22.9, 29.0)	
SOUTH CAROLINA	814	32.1	1.8	(28.6, 35.6)	*	1282	31.4	1.4	(28.6, 34.3)	
SOUTH DAKOTA	945	30.4	1.6	(27.2, 33.6)	*	1187	31.8	1.5	(28.8, 34.8)	
TENNESSEE	1028	32.4	1.6	(29.2, 35.6)	*	1829	30.9	1.2	(28.4, 33.3)	
TEXAS	1010	32.9	1.6	(29.8, 36.1)	*	1384	32.7	1.4	(29.9, 35.4)	
UTAH	1270	29.2	1.6	(26.1, 32.3)	*	1538	24.8	1.4	(22.0, 27.6)	
VERMONT	1377	33.0	1.5	(30.0, 36.0)	*	1655	26.8	1.2	(24.3, 29.2)	
VIRGINIA	1403	32.1	1.7	(28.8, 35.4)	*	1977	28.0	1.4	(25.2, 30.7)	
WASHINGTON	1522	30.1	1.3	(27.5, 32.7)	*	1883	28.1	1.2	(25.6, 30.5)	
WEST VIRGINIA	955	37.2	1.6	(34.0, 40.4)	*	1392	35.4	1.4	(32.7, 38.1)	
WISCONSIN	990	35.7	1.9	(32.1, 39.3)	*	1214	30.0	1.5	(27.0, 33.0)	
WYOMING	1014	27.9	1.6	(24.9, 31.0)	*	1346	27.9	1.3	(25.3, 30.5)	
PUERTO RICO	921	32.9	1.7	(29.7, 36.2)	*	1262	36.7	1.5	(33.7, 39.6)	

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 32.1
 RANGE = 25.0-38.3

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 29.4
 RANGE = 20.8-38.0

* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES

August 17, 1998

TABLE 27.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 OVERWEIGHT: BY BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	33.9	32.7	-1.2
ALASKA	30.0	34.1	4.1
ARIZONA	26.3	25.4	-0.9
ARKANSAS	31.8	32.1	0.3
CALIFORNIA	27.7	29.1	1.4
COLORADO	22.3	25.1	2.8
CONNECTICUT	27.2	29.1	1.9
DELAWARE	31.8	32.5	0.7
DISTRICT OF CO	29.2	28.0	-1.1
FLORIDA	30.5	30.4	-0.2
GEORGIA	24.1	30.8	6.7
HAWAII	24.0	26.6	2.6
IDAHO	29.3	29.6	0.3
ILLINOIS	29.8	32.0	2.1
INDIANA	32.7	34.6	1.9
IOWA	32.7	33.3	0.6
KANSAS	26.1	31.5	5.5
KENTUCKY	33.0	35.6	2.7
LOUISIANA	33.4	33.5	0.1
MAINE	28.1	30.2	2.1
MARYLAND	30.8	31.2	0.5
MASSACHUSETTS	26.7	25.5	-1.2
MICHIGAN	33.0	34.9	1.8
MINNESOTA	27.9	29.7	1.8
MISSISSIPPI	34.5	35.1	0.6
MISSOURI	33.1	31.6	-1.6
MONTANA	28.8	27.9	-0.9
NEBRASKA	29.0	31.1	2.1
NEVADA	27.7	26.2	-1.5
NEW HAMPSHIRE	26.0	27.5	1.5
NEW JERSEY	27.5	28.2	0.7
NEW MEXICO	26.8	29.8	2.9
NEW YORK	29.0	28.4	-0.6
NORTH CAROLINA	30.7	32.4	1.7
NORTH DAKOTA	32.9	32.9	0.0
OHIO	33.4	32.6	-0.8
OKLAHOMA	29.1	28.4	-0.7
OREGON	29.3	32.4	3.1
PENNSYLVANIA	31.8	32.2	0.4
RHODE ISLAND	28.5	28.1	-0.4
SOUTH CAROLINA	34.7	31.8	-2.9
SOUTH DAKOTA	30.1	31.1	1.0
TENNESSEE	30.8	31.6	0.8
TEXAS	31.3	32.8	1.4
UTAH	25.9	27.0	1.0
VERMONT	27.1	29.9	2.8
VIRGINIA	28.3	30.0	1.7
WASHINGTON	27.3	29.1	1.8
WEST VIRGINIA	33.5	36.3	2.8

WISCONSIN	30.5	32.8	2.3
WYOMING	27.9	27.9	0.0
PUERTO RICO	30.8	34.9	4.1

* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES
N/A - NOT AVAILABLE

August 17, 1998

TABLE 28.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY MEDIAN*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2089	38.3	1.2	(35.9, 40.8)
ALASKA	1478	36.6	1.8	(33.0, 40.2)
ARIZONA	1854	29.4	1.5	(26.5, 32.2)
ARKANSAS	1700	36.7	1.5	(33.8, 39.5)
CALIFORNIA	3995	33.6	0.9	(31.9, 35.3)
COLORADO	1782	29.4	1.2	(27.0, 31.8)
CONNECTICUT	2109	34.1	1.3	(31.5, 36.6)
DELAWARE	2459	37.4	1.1	(35.2, 39.6)
DISTRICT OF CO	1488	32.8	1.5	(29.9, 35.6)
FLORIDA	3404	35.4	0.9	(33.6, 37.1)
GEORGIA	2245	37.8	1.2	(35.4, 40.2)
HAWAII	2141	31.3	1.2	(28.9, 33.6)
IDAHO	4796	34.8	0.9	(33.1, 36.6)
ILLINOIS	2776	36.4	1.0	(34.4, 38.4)
INDIANA	2302	39.2	1.2	(36.9, 41.5)
IOWA	3522	38.0	0.9	(36.2, 39.8)
KANSAS	1848	36.0	1.2	(33.5, 38.4)
KENTUCKY	3502	40.8	1.0	(38.9, 42.7)
LOUISIANA	1592	38.4	1.4	(35.6, 41.1)
MAINE	1633	36.0	1.3	(33.5, 38.6)
MARYLAND	4323	35.0	1.0	(33.1, 36.9)
MASSACHUSETTS	1651	30.3	1.3	(27.7, 32.8)
MICHIGAN	2494	39.9	1.1	(37.8, 42.0)
MINNESOTA	4722	35.3	0.8	(33.8, 36.8)
MISSISSIPPI	1540	40.4	1.5	(37.4, 43.3)
MISSOURI	1811	36.6	1.4	(33.9, 39.3)
MONTANA	1743	33.2	1.3	(30.7, 35.7)
NEBRASKA	2604	35.9	1.1	(33.7, 38.0)
NEVADA	2436	31.6	1.9	(28.0, 35.3)
NEW HAMPSHIRE	1463	32.8	1.4	(30.1, 35.6)
NEW JERSEY	2508	33.0	1.1	(30.8, 35.2)
NEW MEXICO	1753	34.6	1.2	(32.2, 36.9)
NEW YORK	3235	32.5	1.0	(30.6, 34.4)
NORTH CAROLINA	3390	36.6	0.9	(34.7, 38.4)
NORTH DAKOTA	1770	38.7	1.2	(36.2, 41.1)
OHIO	3008	37.9	1.2	(35.5, 40.2)
OKLAHOMA	1813	32.6	1.3	(30.1, 35.1)
OREGON	3231	37.2	1.0	(35.2, 39.2)
PENNSYLVANIA	3400	36.7	0.9	(34.8, 38.5)
RHODE ISLAND	1768	33.4	1.3	(30.9, 35.8)
SOUTH CAROLINA	2093	37.7	1.2	(35.4, 40.1)
SOUTH DAKOTA	2127	36.2	1.2	(33.9, 38.5)

TENNESSEE	2853	36.7	1.0	(34.7, 38.8)
TEXAS	2389	37.6	1.1	(35.4, 39.7)
UTAH	2803	31.2	1.1	(28.9, 33.4)
VERMONT	3029	34.6	1.0	(32.6, 36.6)
VIRGINIA	3374	34.6	1.2	(32.3, 36.9)
WASHINGTON	3403	34.1	1.0	(32.2, 36.0)
WEST VIRGINIA	2343	41.2	1.1	(39.1, 43.4)
WISCONSIN	2201	37.1	1.2	(34.7, 39.5)
WYOMING	2356	32.7	1.0	(30.7, 34.7)
PUERTO RICO	2172	40.4	1.2	(38.1, 42.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 36.0 RANGE = 29.4-41.2

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

August 17, 1998

TABLE 28.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY MEDIAN* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*				FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
	*	*	*	*	*	*	*	*	
ALABAMA	802	36.5	1.9	(32.8, 40.2)	*	1287	40.0	1.5	(37.0, 43.0)
ALASKA	704	32.7	2.5	(27.8, 37.7)	*	774	41.0	2.6	(35.9, 46.1)
ARIZONA	873	28.4	2.1	(24.2, 32.5)	*	981	30.3	2.0	(26.3, 34.3)
ARKANSAS	658	36.0	2.3	(31.5, 40.5)	*	1042	37.3	1.8	(33.7, 40.8)
CALIFORNIA	1742	34.6	1.3	(32.0, 37.1)	*	2253	32.6	1.1	(30.3, 34.8)
COLORADO	808	30.8	1.8	(27.3, 34.2)	*	974	28.1	1.7	(24.8, 31.4)
CONNECTICUT	878	37.1	2.1	(33.0, 41.3)	*	1231	31.0	1.6	(28.0, 34.0)
DELAWARE	995	37.1	1.8	(33.6, 40.6)	*	1464	37.7	1.5	(34.8, 40.6)
DISTRICT OF CO	569	27.9	2.1	(23.8, 32.1)	*	919	37.0	1.9	(33.3, 40.6)
FLORIDA	1423	38.6	1.4	(35.8, 41.3)	*	1981	32.4	1.2	(30.1, 34.6)
GEORGIA	951	39.6	1.9	(35.9, 43.4)	*	1294	36.0	1.6	(33.0, 39.0)
HAWAII	944	33.9	1.8	(30.4, 37.4)	*	1197	28.6	1.6	(25.4, 31.7)
IDAHO	2041	36.5	1.3	(33.9, 39.1)	*	2755	33.2	1.2	(30.9, 35.4)
ILLINOIS	1143	35.9	1.6	(32.8, 38.9)	*	1633	36.9	1.4	(34.2, 39.5)
INDIANA	1054	40.5	1.7	(37.1, 43.9)	*	1248	37.9	1.6	(34.8, 40.9)
IOWA	1460	39.3	1.4	(36.4, 42.1)	*	2062	36.8	1.2	(34.5, 39.2)
KANSAS	764	42.7	2.0	(38.9, 46.6)	*	1084	29.1	1.5	(26.2, 32.0)
KENTUCKY	1308	43.1	1.5	(40.1, 46.1)	*	2194	38.6	1.2	(36.4, 40.9)
LOUISIANA	625	37.9	2.2	(33.6, 42.2)	*	967	38.7	1.7	(35.3, 42.2)
MAINE	740	39.4	2.0	(35.4, 43.3)	*	893	32.8	1.6	(29.6, 36.0)
MARYLAND	1907	33.6	1.4	(30.9, 36.3)	*	2416	36.4	1.3	(33.8, 38.9)
MASSACHUSETTS	707	34.4	2.0	(30.4, 38.3)	*	944	26.3	1.7	(23.0, 29.7)
MICHIGAN	1064	41.7	1.7	(38.5, 44.9)	*	1430	38.2	1.4	(35.4, 41.0)
MINNESOTA	2191	37.3	1.1	(35.1, 39.5)	*	2531	33.4	1.0	(31.4, 35.4)
MISSISSIPPI	587	39.8	2.4	(35.0, 44.5)	*	953	41.0	1.9	(37.3, 44.6)
MISSOURI	695	36.7	2.1	(32.6, 40.9)	*	1116	36.5	1.7	(33.0, 39.9)
MONTANA	750	34.2	1.9	(30.4, 38.0)	*	993	32.3	1.7	(29.0, 35.5)
NEBRASKA	1033	39.8	1.9	(36.2, 43.5)	*	1571	32.1	1.4	(29.4, 34.7)
NEVADA	1156	34.5	2.9	(28.7, 40.2)	*	1280	28.6	2.3	(24.1, 33.1)
NEW HAMPSHIRE	601	37.6	2.3	(33.1, 42.1)	*	862	28.2	1.7	(24.9, 31.4)
NEW JERSEY	1029	34.5	1.7	(31.2, 37.8)	*	1479	31.6	1.4	(28.8, 34.4)
NEW MEXICO	741	34.1	1.8	(30.6, 37.7)	*	1012	34.9	1.7	(31.7, 38.2)
NEW YORK	1325	33.5	1.5	(30.6, 36.3)	*	1910	31.6	1.2	(29.3, 33.9)
NORTH CAROLINA	1394	39.0	1.5	(36.1, 41.9)	*	1996	34.2	1.2	(31.9, 36.5)

NORTH DAKOTA	798	40.6	1.8	(37.1, 44.2)	*	972	36.7	1.7	(33.5, 40.0)
OHIO	1215	37.1	1.8	(33.7, 40.5)	*	1793	38.6	1.5	(35.6, 41.6)
OKLAHOMA	817	32.5	2.0	(28.7, 36.4)	*	996	32.6	1.7	(29.3, 35.8)
OREGON	1386	37.1	1.6	(34.0, 40.1)	*	1845	37.3	1.3	(34.7, 39.8)
PENNSYLVANIA	1415	39.7	1.5	(36.8, 42.6)	*	1985	33.8	1.2	(31.5, 36.2)
RHODE ISLAND	745	35.6	2.0	(31.7, 39.4)	*	1023	31.3	1.6	(28.1, 34.5)
SOUTH CAROLINA	813	38.7	1.9	(35.0, 42.3)	*	1280	36.9	1.5	(34.0, 39.8)
SOUTH DAKOTA	943	36.3	1.8	(32.8, 39.7)	*	1184	36.2	1.6	(33.0, 39.3)
TENNESSEE	1027	37.3	1.7	(33.9, 40.6)	*	1826	36.2	1.3	(33.7, 38.7)
TEXAS	1008	37.3	1.7	(34.0, 40.5)	*	1381	37.9	1.5	(35.0, 40.7)
UTAH	1268	33.2	1.7	(29.9, 36.5)	*	1535	29.2	1.5	(26.2, 32.1)
VERMONT	1374	37.4	1.6	(34.3, 40.4)	*	1655	31.8	1.3	(29.3, 34.4)
VIRGINIA	1400	37.4	1.8	(33.8, 40.9)	*	1974	31.9	1.5	(29.0, 34.8)
WASHINGTON	1520	35.4	1.5	(32.5, 38.2)	*	1883	32.8	1.3	(30.2, 35.3)
WEST VIRGINIA	953	41.6	1.7	(38.3, 44.8)	*	1390	40.9	1.4	(38.2, 43.7)
WISCONSIN	989	39.8	1.9	(36.1, 43.5)	*	1212	34.5	1.6	(31.4, 37.6)
WYOMING	1013	32.6	1.7	(29.4, 35.8)	*	1343	32.8	1.4	(30.1, 35.6)
PUERTO RICO	916	38.1	1.7	(34.7, 41.4)	*	1256	42.5	1.6	(39.4, 45.5)

SUMMARY STATISTICS:
 NUMBER OF PARTICIPANTS = 52
 MEDIAN = 37.1
 RANGE = 27.9-43.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 34.4
 RANGE = 26.3-42.5

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

August 17, 1998

TABLE 28.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 OVERWEIGHT: BY MEDIAN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	38.5	38.3	-0.1
ALASKA	34.3	36.6	2.3
ARIZONA	30.3	29.4	-0.9
ARKANSAS	36.4	36.7	0.3
CALIFORNIA	32.8	33.6	0.8
COLORADO	26.2	29.4	3.2
CONNECTICUT	32.0	34.1	2.0
DELAWARE	35.7	37.4	1.8
DISTRICT OF CO	33.7	32.8	-0.9
FLORIDA	35.4	35.4	-0.0
GEORGIA	29.8	37.8	7.9
HAWAII	27.8	31.3	3.4
IDAHO	34.0	34.8	0.9
ILLINOIS	34.3	36.4	2.1
INDIANA	37.8	39.2	1.4
IOWA	37.3	38.0	0.7
KANSAS	31.1	36.0	4.9
KENTUCKY	37.8	40.8	3.0
LOUISIANA	38.7	38.4	-0.3
MAINE	34.2	36.0	1.9
MARYLAND	34.9	35.0	0.1
MASSACHUSETTS	32.0	30.3	-1.7
MICHIGAN	38.4	39.9	1.5
MINNESOTA	33.0	35.3	2.3

MISSISSIPPI	38.3	40.4	2.1
MISSOURI	37.2	36.6	-0.6
MONTANA	32.9	33.2	0.3
NEBRASKA	34.4	35.9	1.5
NEVADA	31.3	31.6	0.3
NEW HAMPSHIRE	30.2	32.8	2.7
NEW JERSEY	31.4	33.0	1.6
NEW MEXICO	32.1	34.6	2.4
NEW YORK	34.1	32.5	-1.6
NORTH CAROLINA	35.6	36.6	1.0
NORTH DAKOTA	37.7	38.7	1.0
OHIO	37.9	37.9	-0.0
OKLAHOMA	33.9	32.6	-1.4
OREGON	33.9	37.2	3.2
PENNSYLVANIA	36.4	36.7	0.2
RHODE ISLAND	32.5	33.4	0.9
SOUTH CAROLINA	39.8	37.7	-2.0
SOUTH DAKOTA	34.4	36.2	1.8
TENNESSEE	34.7	36.7	2.1
TEXAS	35.6	37.6	2.0
UTAH	29.9	31.2	1.3
VERMONT	31.4	34.6	3.2
VIRGINIA	33.2	34.6	1.4
WASHINGTON	31.8	34.1	2.3
WEST VIRGINIA	38.3	41.2	2.9
WISCONSIN	35.4	37.1	1.7
WYOMING	32.1	32.7	0.6
PUERTO RICO	35.1	40.4	5.3

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN
 N/A - NOT AVAILABLE

August 17, 1998

TABLE 29.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV:ENCOURAGE TEEN CONDOM USE*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1652	91.6	0.7	(90.1, 93.0)
ALASKA	1327	89.8	1.3	(87.3, 92.4)
ARIZONA	1385	95.0	0.7	(93.6, 96.4)
ARKANSAS	1309	83.7	1.2	(81.3, 86.1)
COLORADO	1451	93.2	0.8	(91.7, 94.6)
CONNECTICUT	1694	93.2	0.7	(91.8, 94.5)
DELAWARE	1935	92.9	0.7	(91.5, 94.2)
DISTRICT OF CO	1248	90.7	1.0	(88.7, 92.6)
FLORIDA	2488	89.3	0.7	(88.0, 90.7)
GEORGIA	1911	90.6	0.8	(89.2, 92.1)
HAWAII	1722	89.9	0.9	(88.1, 91.6)
IDaho	3895	88.5	0.7	(87.2, 89.8)
ILLINOIS	2362	90.7	0.6	(89.4, 91.9)
INDIANA	1902	90.3	0.8	(88.9, 91.8)
IOWA	2663	91.6	0.6	(90.4, 92.7)
KANSAS	1483	86.9	1.0	(85.0, 88.8)

KENTUCKY	2675	89.5	0.6	(88.2, 90.8)
LOUISIANA	1332	88.5	1.0	(86.6, 90.5)
MAINE	1348	92.3	0.8	(90.8, 93.8)
MARYLAND	3597	86.7	0.8	(85.1, 88.2)
MASSACHUSETTS	1324	96.0	0.6	(94.9, 97.0)
MICHIGAN	2101	92.9	0.6	(91.7, 94.1)
MINNESOTA	3828	78.5	0.7	(77.0, 79.9)
MISSISSIPPI	1235	86.1	1.1	(84.0, 88.2)
MISSOURI	1420	89.1	1.0	(87.2, 91.1)
MONTANA	1386	84.0	1.1	(81.9, 86.1)
NEBRASKA	1928	88.4	0.9	(86.5, 90.2)
NEVADA	2112	91.8	1.2	(89.4, 94.2)
NEW HAMPSHIRE	1166	94.2	0.7	(92.8, 95.7)
NEW JERSEY	1975	92.6	0.7	(91.2, 93.9)
NEW MEXICO	1417	90.8	0.9	(89.1, 92.5)
NEW YORK	2717	89.1	0.7	(87.7, 90.4)
NORTH CAROLINA	2739	88.7	0.7	(87.4, 90.1)
NORTH DAKOTA	1302	89.7	0.9	(87.9, 91.6)
OHIO	2284	90.1	0.8	(88.6, 91.6)
OKLAHOMA	1235	89.3	1.0	(87.3, 91.3)
OREGON	2581	91.6	0.6	(90.3, 92.8)
PENNSYLVANIA	2715	91.3	0.6	(90.2, 92.5)
RHODE ISLAND	1343	91.9	0.8	(90.4, 93.5)
SOUTH CAROLINA	1624	82.5	1.1	(80.4, 84.5)
SOUTH DAKOTA	1594	86.8	0.9	(85.0, 88.7)
TENNESSEE	2195	79.8	1.0	(78.0, 81.7)
TEXAS	2066	86.1	0.9	(84.4, 87.8)
UTAH	2354	88.4	0.9	(86.6, 90.2)
VERMONT	2480	94.3	0.6	(93.2, 95.4)
VIRGINIA	2862	89.6	1.0	(87.7, 91.5)
WASHINGTON	2924	92.4	0.5	(91.4, 93.5)
WEST VIRGINIA	1803	87.8	0.8	(86.2, 89.4)
WISCONSIN	1790	93.2	0.7	(91.9, 94.6)
WYOMING	1964	93.5	0.7	(92.1, 94.9)
PUERTO RICO	1780	87.9	0.9	(86.2, 89.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 89.9 RANGE = 78.5-96.0

* WOULD ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

August 17, 1998

TABLE 29.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV:ENCOURAGE TEEN CONDOM USE* -BY SEX
DENOMINATOR IS PERSONS AGE 18-64 YEARS
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				*
	SAMPLE SIZE	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	*
				*				*	*
ALABAMA	651	92.0	1.1	(89.9, 94.1)	*	1001	91.2	1.0	(89.3, 93.1)
ALASKA	612	88.1	2.0	(84.1, 92.1)	*	715	91.7	1.5	(88.8, 94.7)
ARIZONA	670	93.7	1.3	(91.3, 96.2)	*	715	96.2	0.8	(94.8, 97.7)
ARKANSAS	529	83.7	1.8	(80.1, 87.3)	*	780	83.8	1.6	(80.7, 87.0)
COLORADO	679	91.2	1.3	(88.7, 93.6)	*	772	95.2	0.8	(93.7, 96.7)
CONNECTICUT	677	93.1	1.0	(91.1, 95.1)	*	1017	93.2	0.9	(91.5, 94.9)
DELAWARE	792	92.5	1.1	(90.3, 94.7)	*	1143	93.1	0.8	(91.6, 94.7)

DISTRICT OF CO	498	89.8	1.6	(86.6, 93.0)	*	750	91.5	1.2	(89.2, 93.8)
FLORIDA	1039	88.2	1.1	(86.0, 90.4)	*	1449	90.5	0.8	(88.9, 92.1)
GEORGIA	805	90.4	1.2	(88.1, 92.7)	*	1106	90.9	1.0	(89.0, 92.7)
HAWAII	773	89.4	1.4	(86.8, 92.1)	*	949	90.3	1.1	(88.1, 92.5)
IDAHO	1712	88.0	1.0	(86.1, 89.9)	*	2183	89.1	0.9	(87.3, 90.9)
ILLINOIS	999	91.4	0.9	(89.5, 93.2)	*	1363	90.0	0.9	(88.3, 91.8)
INDIANA	889	89.9	1.1	(87.7, 92.0)	*	1013	90.8	1.0	(88.8, 92.8)
IOWA	1179	91.0	0.9	(89.2, 92.8)	*	1484	92.2	0.8	(90.6, 93.7)
KANSAS	626	87.2	1.4	(84.4, 89.9)	*	857	86.6	1.3	(84.2, 89.1)
KENTUCKY	1030	88.8	1.0	(86.8, 90.7)	*	1645	90.2	0.8	(88.6, 91.7)
LOUISIANA	515	87.5	1.6	(84.3, 90.6)	*	817	89.6	1.2	(87.3, 91.8)
MAINE	612	91.3	1.2	(89.0, 93.6)	*	736	93.3	1.0	(91.3, 95.3)
MARYLAND	1590	85.3	1.1	(83.1, 87.6)	*	2007	88.0	0.9	(86.2, 89.8)
MASSACHUSETTS	569	95.8	0.8	(94.2, 97.4)	*	755	96.2	0.7	(94.7, 97.6)
MICHIGAN	904	91.6	1.1	(89.5, 93.6)	*	1197	94.2	0.7	(92.8, 95.6)
MINNESOTA	1853	77.1	1.1	(75.0, 79.2)	*	1975	79.8	1.0	(77.9, 81.7)
MISSISSIPPI	490	86.5	1.6	(83.3, 89.6)	*	745	85.8	1.4	(83.1, 88.5)
MISSOURI	549	88.0	1.6	(84.9, 91.1)	*	871	90.3	1.2	(87.9, 92.6)
MONTANA	603	82.9	1.6	(79.8, 86.1)	*	783	85.1	1.4	(82.4, 87.8)
NEBRASKA	780	86.5	1.6	(83.5, 89.6)	*	1148	90.2	1.0	(88.2, 92.2)
NEVADA	1007	90.5	2.1	(86.4, 94.6)	*	1105	93.3	1.2	(90.9, 95.6)
NEW HAMPSHIRE	470	93.9	1.2	(91.6, 96.1)	*	696	94.6	0.9	(92.9, 96.3)
NEW JERSEY	812	92.2	1.1	(90.1, 94.2)	*	1163	92.9	0.9	(91.2, 94.7)
NEW MEXICO	605	90.9	1.3	(88.4, 93.5)	*	812	90.6	1.2	(88.3, 93.0)
NEW YORK	1113	89.6	1.0	(87.6, 91.6)	*	1604	88.6	0.9	(86.8, 90.4)
NORTH CAROLINA	1142	88.6	1.0	(86.6, 90.6)	*	1597	88.9	0.9	(87.2, 90.6)
NORTH DAKOTA	612	89.1	1.3	(86.4, 91.7)	*	690	90.5	1.3	(88.0, 93.0)
OHIO	908	88.6	1.3	(86.0, 91.2)	*	1376	91.5	0.9	(89.7, 93.3)
OKLAHOMA	503	88.9	1.5	(86.0, 91.8)	*	732	89.7	1.3	(87.2, 92.2)
OREGON	1140	91.3	1.0	(89.4, 93.1)	*	1441	91.9	0.8	(90.2, 93.5)
PENNSYLVANIA	1135	91.6	0.9	(89.8, 93.3)	*	1580	91.1	0.8	(89.6, 92.6)
RHODE ISLAND	569	91.7	1.2	(89.3, 94.1)	*	774	92.2	1.0	(90.2, 94.2)
SOUTH CAROLINA	651	83.5	1.6	(80.3, 86.6)	*	973	81.5	1.4	(78.8, 84.2)
SOUTH DAKOTA	731	87.3	1.4	(84.6, 90.0)	*	863	86.4	1.2	(84.0, 88.8)
TENNESSEE	820	78.9	1.5	(75.9, 81.9)	*	1375	80.7	1.1	(78.5, 82.9)
TEXAS	875	84.7	1.3	(82.1, 87.3)	*	1191	87.5	1.1	(85.4, 89.5)
UTAH	1075	87.7	1.4	(84.9, 90.4)	*	1279	89.1	1.2	(86.8, 91.4)
VERMONT	1133	94.1	0.7	(92.7, 95.6)	*	1347	94.5	0.8	(92.9, 96.0)
VIRGINIA	1199	88.8	1.7	(85.5, 92.0)	*	1663	90.4	1.0	(88.4, 92.4)
WASHINGTON	1299	92.1	0.8	(90.5, 93.7)	*	1625	92.8	0.7	(91.3, 94.2)
WEST VIRGINIA	770	87.2	1.3	(84.7, 89.8)	*	1033	88.4	1.1	(86.3, 90.4)
WISCONSIN	844	93.4	1.0	(91.4, 95.4)	*	946	93.1	1.0	(91.2, 95.0)
WYOMING	865	92.1	1.0	(90.2, 94.1)	*	1099	95.0	1.0	(93.0, 97.0)
PUERTO RICO	756	89.2	1.2	(86.7, 91.6)	*	1024	86.7	1.2	(84.4, 89.0)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 51
MEDIAN = 89.4
RANGE = 77.1-95.8

NUMBER OF PARTICIPANTS = 51
MEDIAN = 90.5
RANGE = 79.8-96.2

* WOULD ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

August 17, 1998

TABLE 29.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1996 TO 1997
HIV:ENCOURAGE TEEN CONDOM USE*
DENOMINATOR IS PERSONS AGE 18-64 YEARS
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

1996 1997 PERCENT

PARTICIPANT	PERCENT	PERCENT	DIFFERENCE
ALABAMA	91.0	91.6	0.5
ALASKA	91.4	89.8	-1.6
ARIZONA	91.7	95.0	3.3
ARKANSAS	78.2	83.7	5.6
COLORADO	90.2	93.2	3.0
CONNECTICUT	92.1	93.2	1.0
DELAWARE	90.1	92.9	2.8
DISTRICT OF CO	93.8	90.7	-3.2
FLORIDA	89.1	89.3	0.2
GEORGIA	92.6	90.6	-1.9
HAWAII	87.9	89.9	2.0
IDAHO	87.4	88.5	1.1
ILLINOIS	91.0	90.7	-0.3
INDIANA	87.8	90.3	2.5
IOWA	92.6	91.6	-1.0
KANSAS	87.0	86.9	-0.1
KENTUCKY	87.6	89.5	1.9
LOUISIANA	87.3	88.5	1.3
MAINE	92.4	92.3	-0.1
MARYLAND	87.2	86.7	-0.5
MASSACHUSETTS	94.0	96.0	2.0
MICHIGAN	88.8	92.9	4.1
MINNESOTA	80.0	78.5	-1.6
MISSISSIPPI	88.4	86.1	-2.2
MISSOURI	88.5	89.1	0.6
MONTANA	84.9	84.0	-0.8
NEBRASKA	90.1	88.4	-1.7
NEVADA	90.2	91.8	1.6
NEW HAMPSHIRE	93.1	94.2	1.1
NEW JERSEY	91.5	92.6	1.0
NEW MEXICO	88.8	90.8	2.0
NEW YORK	89.1	89.1	-0.0
NORTH CAROLINA	85.3	88.7	3.5
NORTH DAKOTA	90.0	89.7	-0.2
OHIO	89.2	90.1	0.9
OKLAHOMA	86.3	89.3	3.0
OREGON	91.4	91.6	0.2
PENNSYLVANIA	89.7	91.3	1.7
RHODE ISLAND	93.0	91.9	-1.1
SOUTH CAROLINA	88.1	82.5	-5.6
SOUTH DAKOTA	82.9	86.8	3.9
TENNESSEE	81.8	79.8	-2.0
TEXAS	89.7	86.1	-3.6
UTAH	87.0	88.4	1.4
VERMONT	93.4	94.3	0.9
VIRGINIA	89.9	89.6	-0.2
WASHINGTON	90.2	92.4	2.2
WEST VIRGINIA	87.7	87.8	0.1
WISCONSIN	91.3	93.2	1.9
WYOMING	91.6	93.5	1.9
PUERTO RICO	84.7	87.9	3.2

* WOULD ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

N/A - NOT AVAILABLE

August 17, 1998

TABLE 30.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV BLOOD TEST*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1629	49.5	1.4	(46.7, 52.2)
ALASKA	1326	54.1	2.0	(50.2, 58.1)
ARIZONA	1395	31.7	1.7	(28.4, 35.0)
ARKANSAS	1287	41.7	1.8	(38.2, 45.1)
COLORADO	1434	47.1	1.5	(44.1, 50.0)
CONNECTICUT	1668	43.0	1.6	(39.9, 46.1)
DELAWARE	1911	52.1	1.4	(49.5, 54.8)
DISTRICT OF CO	1247	60.2	1.7	(57.0, 63.5)
FLORIDA	2477	55.8	1.1	(53.6, 58.0)
GEORGIA	1898	41.2	1.4	(38.6, 43.9)
HAWAII	1720	45.5	1.4	(42.7, 48.2)
IDAHO	3863	44.1	1.1	(41.9, 46.2)
ILLINOIS	2341	42.3	1.2	(40.0, 44.5)
INDIANA	1851	38.2	1.3	(35.6, 40.8)
IOWA	2641	31.9	1.0	(29.8, 33.9)
KANSAS	1485	24.4	1.2	(22.0, 26.8)
KENTUCKY	2602	37.8	1.1	(35.6, 39.9)
LOUISIANA	1303	50.1	1.6	(47.0, 53.2)
MAINE	1330	36.3	1.5	(33.4, 39.3)
MARYLAND	3558	52.6	1.1	(50.4, 54.8)
MASSACHUSETTS	1295	44.9	1.6	(41.8, 48.1)
MICHIGAN	2069	44.5	1.2	(42.2, 46.9)
MINNESOTA	3808	32.0	0.8	(30.4, 33.6)
MISSISSIPPI	1220	49.2	1.7	(45.9, 52.4)
MISSOURI	1379	41.1	1.6	(37.9, 44.2)
MONTANA	1375	38.2	1.5	(35.4, 41.1)
NEBRASKA	1924	34.8	1.5	(31.9, 37.7)
NEVADA	2054	60.5	2.1	(56.3, 64.7)
NEW HAMPSHIRE	1165	37.4	1.7	(34.2, 40.7)
NEW JERSEY	1942	48.5	1.4	(45.8, 51.1)
NEW MEXICO	1400	45.5	1.5	(42.5, 48.4)
NEW YORK	2710	43.3	1.1	(41.2, 45.5)
NORTH CAROLINA	2719	40.5	1.1	(38.3, 42.6)
NORTH DAKOTA	1327	31.0	1.4	(28.1, 33.8)
OHIO	2272	33.7	1.3	(31.2, 36.3)
OKLAHOMA	1243	36.3	1.5	(33.3, 39.3)
OREGON	2554	43.4	1.2	(41.1, 45.7)
PENNSYLVANIA	2676	42.0	1.1	(39.7, 44.2)
RHODE ISLAND	1329	46.0	1.6	(42.9, 49.2)
SOUTH CAROLINA	1581	47.9	1.5	(45.1, 50.8)
SOUTH DAKOTA	1595	37.6	1.4	(34.9, 40.3)
TENNESSEE	2230	45.5	1.2	(43.1, 47.9)
TEXAS	2049	51.2	1.2	(48.8, 53.6)
UTAH	2335	38.8	1.3	(36.2, 41.4)
VERMONT	2456	38.7	1.2	(36.4, 41.0)
VIRGINIA	2795	57.1	1.4	(54.4, 59.9)
WASHINGTON	2871	48.3	1.1	(46.1, 50.6)
WEST VIRGINIA	1784	33.1	1.2	(30.7, 35.4)
WISCONSIN	1757	40.3	1.4	(37.6, 43.0)
WYOMING	1954	38.7	1.4	(35.9, 41.4)
PUERTO RICO	1828	40.3	1.3	(37.8, 42.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 42.3 RANGE = 24.4-60.5

* EVER HAD BLOOD TESTED FOR HIV

August 17, 1998

TABLE 30.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV BLOOD TEST* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES-----*				FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	644	47.4	2.2	(43.1, 51.7)	*	985	51.4	1.8 (48.0, 54.9)
ALASKA	612	54.4	3.0	(48.5, 60.2)	*	714	53.9	2.7 (48.5, 59.3)
ARIZONA	678	31.8	2.6	(26.8, 36.9)	*	717	31.6	2.2 (27.3, 35.9)
ARKANSAS	509	41.5	2.8	(36.0, 46.9)	*	778	41.9	2.2 (37.6, 46.1)
COLORADO	675	47.0	2.1	(42.8, 51.2)	*	759	47.1	2.1 (43.0, 51.2)
CONNECTICUT	669	46.2	2.5	(41.3, 51.0)	*	999	39.8	1.9 (36.1, 43.5)
DELAWARE	779	54.2	2.1	(50.0, 58.3)	*	1132	50.2	1.7 (46.8, 53.6)
DISTRICT OF CO	496	63.6	2.5	(58.7, 68.6)	*	751	57.1	2.2 (52.9, 61.3)
FLORIDA	1040	57.0	1.7	(53.7, 60.3)	*	1437	54.6	1.5 (51.7, 57.6)
GEORGIA	810	41.0	2.1	(36.9, 45.0)	*	1088	41.5	1.8 (38.0, 45.0)
HAWAII	765	49.2	2.1	(45.2, 53.3)	*	955	41.6	1.9 (37.8, 45.3)
IDAHO	1701	45.0	1.6	(41.9, 48.2)	*	2162	43.1	1.5 (40.2, 46.1)
ILLINOIS	989	44.3	1.8	(40.9, 47.7)	*	1352	40.3	1.5 (37.4, 43.2)
INDIANA	864	38.6	1.9	(34.8, 42.4)	*	987	37.8	1.8 (34.3, 41.3)
IOWA	1179	34.6	1.6	(31.5, 37.6)	*	1462	29.1	1.3 (26.5, 31.7)
KANSAS	630	23.7	1.9	(20.0, 27.3)	*	855	25.1	1.6 (22.1, 28.2)
KENTUCKY	996	37.9	1.7	(34.6, 41.2)	*	1606	37.6	1.4 (35.0, 40.3)
LOUISIANA	506	49.7	2.5	(44.8, 54.6)	*	797	50.5	2.0 (46.7, 54.3)
MAINE	599	37.6	2.3	(33.1, 42.1)	*	731	35.0	1.9 (31.3, 38.8)
MARYLAND	1568	51.4	1.7	(48.1, 54.7)	*	1990	53.7	1.4 (50.9, 56.5)
MASSACHUSETTS	557	50.9	2.5	(46.0, 55.7)	*	738	39.1	2.1 (35.0, 43.1)
MICHIGAN	894	44.4	1.8	(40.8, 48.0)	*	1175	44.7	1.6 (41.6, 47.8)
MINNESOTA	1849	32.3	1.2	(30.0, 34.7)	*	1959	31.7	1.1 (29.5, 33.9)
MISSISSIPPI	481	49.2	2.7	(43.9, 54.4)	*	739	49.2	2.1 (45.1, 53.3)
MISSOURI	528	40.7	2.5	(35.8, 45.6)	*	851	41.5	2.0 (37.5, 45.4)
MONTANA	596	39.8	2.1	(35.6, 44.0)	*	779	36.7	1.9 (33.0, 40.3)
NEBRASKA	781	35.8	2.3	(31.4, 40.3)	*	1143	33.7	1.9 (30.0, 37.4)
NEVADA	973	61.9	3.3	(55.5, 68.3)	*	1081	59.1	2.7 (53.8, 64.4)
NEW HAMPSHIRE	474	39.0	2.6	(34.0, 44.0)	*	691	35.8	2.1 (31.8, 39.9)
NEW JERSEY	803	49.5	2.1	(45.5, 53.5)	*	1139	47.5	1.8 (44.0, 50.9)
NEW MEXICO	604	46.4	2.2	(42.0, 50.8)	*	796	44.5	1.9 (40.7, 48.3)
NEW YORK	1109	44.2	1.7	(41.0, 47.5)	*	1601	42.5	1.4 (39.7, 45.2)
NORTH CAROLINA	1144	38.9	1.6	(35.8, 42.1)	*	1575	42.0	1.4 (39.2, 44.7)
NORTH DAKOTA	613	33.5	2.0	(29.5, 37.4)	*	714	28.4	1.9 (24.6, 32.1)
OHIO	906	34.0	2.1	(29.9, 38.0)	*	1366	33.5	1.7 (30.2, 36.8)
OKLAHOMA	512	33.7	2.2	(29.4, 37.9)	*	731	39.0	2.0 (35.0, 42.9)
OREGON	1127	46.0	1.8	(42.5, 49.6)	*	1427	40.8	1.5 (37.7, 43.8)
PENNSYLVANIA	1125	44.5	1.7	(41.2, 47.8)	*	1551	39.4	1.5 (36.4, 42.4)
RHODE ISLAND	569	52.1	2.3	(47.5, 56.7)	*	760	40.1	2.1 (36.0, 44.1)
SOUTH CAROLINA	632	46.8	2.3	(42.4, 51.2)	*	949	49.1	1.9 (45.4, 52.7)
SOUTH DAKOTA	734	39.8	2.0	(35.8, 43.7)	*	861	35.4	1.9 (31.7, 39.1)
TENNESSEE	827	44.9	1.9	(41.1, 48.6)	*	1403	46.1	1.5 (43.2, 49.1)
TEXAS	872	52.1	1.9	(48.4, 55.7)	*	1177	50.3	1.6 (47.1, 53.5)
UTAH	1069	43.0	1.9	(39.2, 46.8)	*	1266	34.7	1.8 (31.2, 38.2)
VERMONT	1129	39.8	1.8	(36.4, 43.3)	*	1327	37.6	1.6 (34.5, 40.6)
VIRGINIA	1178	57.6	2.1	(53.4, 61.7)	*	1617	56.7	1.9 (53.0, 60.4)
WASHINGTON	1287	51.7	1.7	(48.4, 54.9)	*	1584	45.0	1.5 (42.0, 47.9)
WEST VIRGINIA	760	33.6	1.8	(30.1, 37.1)	*	1024	32.6	1.6 (29.4, 35.7)
WISCONSIN	829	41.1	2.1	(37.0, 45.2)	*	928	39.5	1.8 (35.9, 43.1)
WYOMING	862	39.5	2.1	(35.5, 43.5)	*	1092	37.8	1.9 (34.2, 41.4)
PUERTO RICO	769	40.3	1.9	(36.5, 44.1)	*	1059	40.3	1.7 (37.0, 43.5)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS = 51	NUMBER OF PARTICIPANTS = 51
	MEDIAN = 44.3	MEDIAN = 40.8
	RANGE = 23.7-63.6	RANGE = 25.1-59.1

TABLE 30.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 HIV BLOOD TEST*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	46.5	49.5	3.0
ALASKA	48.6	54.1	5.6
ARIZONA	44.4	31.7	-12.7
ARKANSAS	44.6	41.7	-3.0
COLORADO	45.9	47.1	1.2
CONNECTICUT	41.6	43.0	1.4
DELAWARE	46.1	52.1	6.1
DISTRICT OF CO	61.3	60.2	-1.1
FLORIDA	53.6	55.8	2.2
GEORGIA	33.6	41.2	7.6
HAWAII	43.1	45.5	2.4
IDAHO	42.6	44.1	1.5
ILLINOIS	41.2	42.3	1.1
INDIANA	39.4	38.2	-1.2
IOWA	32.9	31.9	-1.1
KANSAS	30.1	24.4	-5.8
KENTUCKY	34.6	37.8	3.2
LOUISIANA	49.0	50.1	1.1
MAINE	34.4	36.3	1.9
MARYLAND	50.4	52.6	2.1
MASSACHUSETTS	38.1	44.9	6.9
MICHIGAN	43.0	44.5	1.5
MINNESOTA	31.4	32.0	0.6
MISSISSIPPI	47.7	49.2	1.5
MISSOURI	39.7	41.1	1.4
MONTANA	35.4	38.2	2.8
NEBRASKA	35.0	34.8	-0.2
NEVADA	54.7	60.5	5.9
NEW HAMPSHIRE	36.9	37.4	0.6
NEW JERSEY	45.7	48.5	2.8
NEW MEXICO	50.2	45.5	-4.8
NEW YORK	41.1	43.3	2.3
NORTH CAROLINA	38.7	40.5	1.8
NORTH DAKOTA	29.0	31.0	2.0
OHIO	28.1	33.7	5.6
OKLAHOMA	38.6	36.3	-2.3
OREGON	42.5	43.4	0.9
PENNSYLVANIA	41.6	42.0	0.3
RHODE ISLAND	47.6	46.0	-1.6
SOUTH CAROLINA	43.0	47.9	4.9
SOUTH DAKOTA	27.3	37.6	10.3
TENNESSEE	40.0	45.5	5.5
TEXAS	51.1	51.2	0.1
UTAH	39.0	38.8	-0.2
VERMONT	39.8	38.7	-1.1
VIRGINIA	53.2	57.1	3.9

WASHINGTON	44.1	48.3	4.2
WEST VIRGINIA	30.7	33.1	2.4
WISCONSIN	41.3	40.3	-1.0
WYOMING	39.4	38.7	-0.7
PUERTO RICO	39.0	40.3	1.3

* EVER HAD BLOOD TESTED FOR HIV
N/A - NOT AVAILABLE

August 17, 1998

TABLE 31.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING HIV MEDIUM/HIGH*
DENOMINATOR IS PERSONS AGE 18-64 YEARS
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1660	5.8	0.7	(4.5, 7.2)
ALASKA	1339	5.7	0.9	(3.9, 7.5)
ARIZONA	1394	3.3	0.8	(1.8, 4.8)
ARKANSAS	1338	7.2	1.1	(5.0, 9.4)
COLORADO	1458	4.6	0.6	(3.4, 5.7)
CONNECTICUT	1692	6.1	0.6	(4.8, 7.3)
DELAWARE	1941	7.0	0.8	(5.4, 8.5)
DISTRICT OF CO	1246	7.2	1.0	(5.3, 9.1)
FLORIDA	2472	7.9	0.7	(6.6, 9.2)
GEORGIA	1908	4.9	0.6	(3.7, 6.1)
HAWAII	1732	7.0	0.8	(5.5, 8.5)
IDAH0	3941	5.2	0.4	(4.4, 6.0)
ILLINOIS	2346	7.6	0.6	(6.3, 8.8)
INDIANA	1911	6.1	0.7	(4.8, 7.4)
IOWA	2683	5.5	0.5	(4.6, 6.5)
KANSAS	1491	8.7	0.8	(7.0, 10.3)
KENTUCKY	2679	5.8	0.5	(4.8, 6.9)
LOUISIANA	1323	4.9	0.7	(3.6, 6.2)
MAINE	1359	4.9	0.7	(3.6, 6.2)
MARYLAND	3629	5.0	0.5	(4.0, 5.9)
MASSACHUSETTS	1305	6.1	0.8	(4.5, 7.6)
MICHIGAN	2108	5.4	0.5	(4.4, 6.4)
MINNESOTA	3852	5.7	0.4	(4.9, 6.5)
MISSISSIPPI	1231	6.8	1.1	(4.7, 8.9)
MISSOURI	1434	8.0	1.1	(5.8, 10.1)
MONTANA	1396	4.3	0.7	(3.0, 5.6)
NEBRASKA	1959	8.3	0.8	(6.7, 9.8)
NEVADA	2110	6.3	1.0	(4.4, 8.2)
NEW HAMPSHIRE	1175	4.2	0.8	(2.7, 5.7)
NEW JERSEY	1966	6.4	0.7	(5.1, 7.7)
NEW MEXICO	1418	7.9	0.8	(6.2, 9.5)
NEW YORK	2699	7.3	0.6	(6.2, 8.5)
NORTH CAROLINA	2760	6.7	0.6	(5.6, 7.9)
NORTH DAKOTA	1332	4.9	0.7	(3.5, 6.3)
OHIO	2298	6.2	0.6	(5.0, 7.4)
OKLAHOMA	1253	4.9	0.7	(3.5, 6.3)
OREGON	2614	5.5	0.6	(4.4, 6.6)
PENNSYLVANIA	2720	6.8	0.7	(5.5, 8.1)

RHODE ISLAND	1345	7.3	0.8	(5.6, 8.9)
SOUTH CAROLINA	1626	5.5	0.7	(4.2, 6.9)
SOUTH DAKOTA	1618	8.9	0.8	(7.4, 10.4)
TENNESSEE	2242	7.0	0.6	(5.8, 8.3)
TEXAS	2053	12.1	0.8	(10.4, 13.7)
UTAH	2405	5.2	0.6	(4.1, 6.3)
VERMONT	2489	6.6	0.8	(5.0, 8.3)
VIRGINIA	2856	5.0	0.5	(4.0, 6.0)
WASHINGTON	2961	4.5	0.4	(3.7, 5.3)
WEST VIRGINIA	1803	9.1	0.7	(7.7, 10.6)
WISCONSIN	1798	5.6	0.7	(4.3, 7.0)
WYOMING	1990	7.9	1.2	(5.5, 10.2)
PUERTO RICO	1771	10.2	0.8	(8.6, 11.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 6.1 RANGE = 3.3-12.1

* CHANCES OF GETTING HIV MEDIUM/HIGH

August 17, 1998

TABLE 31.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING HIV MEDIUM/HIGH* -BY SEX
DENOMINATOR IS PERSONS AGE 18-64 YEARS
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
					*					*
ALABAMA	661	6.5	1.2	(4.2, 8.7)	*	999	5.3	0.8	(3.7, 6.8)	*
ALASKA	618	6.1	1.3	(3.4, 8.7)	*	721	5.3	1.2	(3.0, 7.7)	*
ARIZONA	675	5.0	1.4	(2.3, 7.7)	*	719	1.6	0.6	(0.4, 2.8)	*
ARKANSAS	537	8.3	2.1	(4.3, 12.3)	*	801	6.1	0.9	(4.3, 7.9)	*
COLORADO	678	5.8	1.0	(3.9, 7.8)	*	780	3.3	0.6	(2.1, 4.6)	*
CONNECTICUT	674	7.1	1.0	(5.1, 9.2)	*	1018	5.0	0.8	(3.5, 6.5)	*
DELAWARE	797	8.4	1.3	(5.9, 11.0)	*	1144	5.6	0.9	(3.7, 7.4)	*
DISTRICT OF CO	495	8.8	1.6	(5.6, 11.9)	*	751	5.7	1.0	(3.8, 7.7)	*
FLORIDA	1039	9.9	1.1	(7.7, 12.1)	*	1433	5.9	0.7	(4.6, 7.2)	*
GEORGIA	805	5.6	1.1	(3.5, 7.6)	*	1103	4.2	0.7	(2.8, 5.7)	*
HAWAII	779	6.9	1.1	(4.7, 9.0)	*	953	7.2	1.1	(5.0, 9.3)	*
IDAHO	1736	5.9	0.6	(4.7, 7.0)	*	2205	4.5	0.5	(3.5, 5.5)	*
ILLINOIS	995	8.6	1.1	(6.6, 10.7)	*	1351	6.5	0.7	(5.1, 7.8)	*
INDIANA	895	6.1	1.0	(4.1, 8.0)	*	1016	6.1	0.8	(4.5, 7.8)	*
IOWA	1192	5.6	0.7	(4.2, 7.1)	*	1491	5.4	0.6	(4.2, 6.6)	*
KANSAS	628	9.8	1.4	(7.1, 12.5)	*	863	7.5	1.0	(5.6, 9.4)	*
KENTUCKY	1032	6.7	0.9	(4.9, 8.5)	*	1647	5.0	0.6	(3.8, 6.1)	*
LOUISIANA	513	5.2	1.1	(3.1, 7.3)	*	810	4.6	0.8	(3.0, 6.2)	*
MAINE	618	5.0	1.0	(3.1, 6.9)	*	741	4.8	0.8	(3.2, 6.4)	*
MARYLAND	1601	5.4	0.8	(3.8, 6.9)	*	2028	4.6	0.6	(3.5, 5.6)	*
MASSACHUSETTS	562	5.5	1.2	(3.1, 7.8)	*	743	6.7	1.1	(4.6, 8.8)	*
MICHIGAN	913	5.1	0.8	(3.6, 6.7)	*	1195	5.6	0.7	(4.3, 6.9)	*
MINNESOTA	1868	6.3	0.6	(5.1, 7.6)	*	1984	5.1	0.5	(4.1, 6.1)	*
MISSISSIPPI	488	9.3	2.0	(5.4, 13.3)	*	743	4.3	0.8	(2.9, 5.8)	*
MISSOURI	551	10.3	2.0	(6.4, 14.2)	*	883	5.7	1.0	(3.8, 7.7)	*
MONTANA	608	3.4	0.7	(2.1, 4.8)	*	788	5.2	1.2	(2.9, 7.5)	*
NEBRASKA	797	8.2	1.2	(5.9, 10.4)	*	1162	8.3	1.0	(6.4, 10.3)	*
NEVADA	1007	8.5	1.7	(5.2, 11.7)	*	1103	4.0	0.9	(2.2, 5.7)	*
NEW HAMPSHIRE	475	4.1	1.2	(1.7, 6.5)	*	700	4.3	0.9	(2.5, 6.1)	*

NEW JERSEY	816	7.9	1.1	(5.7, 10.1)	*	1150	4.8	0.7	(3.4, 6.3)
NEW MEXICO	609	8.1	1.3	(5.5, 10.6)	*	809	7.7	1.0	(5.6, 9.7)
NEW YORK	1104	8.1	0.9	(6.3, 9.9)	*	1595	6.6	0.7	(5.2, 8.0)
NORTH CAROLINA	1152	6.8	0.9	(5.0, 8.5)	*	1608	6.7	0.7	(5.3, 8.1)
NORTH DAKOTA	619	4.5	1.0	(2.5, 6.5)	*	713	5.3	0.9	(3.5, 7.1)
OHIO	917	6.8	0.9	(5.1, 8.6)	*	1381	5.5	0.8	(4.0, 7.1)
OKLAHOMA	513	5.6	1.2	(3.3, 7.9)	*	740	4.2	0.8	(2.7, 5.7)
OREGON	1155	6.5	0.9	(4.7, 8.3)	*	1459	4.5	0.6	(3.4, 5.6)
PENNSYLVANIA	1132	6.7	0.9	(5.0, 8.4)	*	1588	6.9	1.0	(5.0, 8.9)
RHODE ISLAND	576	9.0	1.4	(6.3, 11.7)	*	769	5.5	0.9	(3.7, 7.3)
SOUTH CAROLINA	646	7.0	1.2	(4.7, 9.3)	*	980	4.2	0.7	(2.8, 5.6)
SOUTH DAKOTA	743	10.3	1.2	(7.9, 12.6)	*	875	7.6	1.0	(5.7, 9.5)
TENNESSEE	828	6.9	1.0	(5.0, 8.9)	*	1414	7.1	0.8	(5.6, 8.6)
TEXAS	872	12.9	1.3	(10.4, 15.4)	*	1181	11.2	1.0	(9.2, 13.2)
UTAH	1097	5.4	0.8	(3.8, 7.0)	*	1308	5.0	0.8	(3.5, 6.5)
VERMONT	1137	6.5	1.3	(3.9, 9.0)	*	1352	6.7	1.1	(4.6, 8.9)
VIRGINIA	1196	5.5	0.9	(3.7, 7.3)	*	1660	4.5	0.6	(3.4, 5.6)
WASHINGTON	1318	5.7	0.7	(4.3, 7.1)	*	1643	3.3	0.5	(2.4, 4.2)
WEST VIRGINIA	769	9.1	1.1	(7.0, 11.2)	*	1034	9.2	1.0	(7.2, 11.2)
WISCONSIN	851	6.1	1.1	(4.0, 8.3)	*	947	5.2	0.8	(3.5, 6.8)
WYOMING	873	8.7	2.2	(4.5, 12.9)	*	1117	7.0	1.1	(4.8, 9.2)
PUERTO RICO	749	11.5	1.4	(8.8, 14.2)	*	1022	9.0	1.0	(7.0, 10.9)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 51
 MEDIAN = 6.7
 RANGE = 3.4-12.9

NUMBER OF PARTICIPANTS = 51
 MEDIAN = 5.4
 RANGE = 1.6-11.2

* CHANCES OF GETTING HIV MEDIUM/HIGH

August 17, 1998

STATE SURVEY DATA

TABLE 31.3 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 CHANCE OF GETTING HIV MEDIUM/HIGH*
 DENOMINATOR IS PERSONS AGE 18-64 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	6.2	5.8	-0.4
ALASKA	5.2	5.7	0.6
ARIZONA	6.7	3.3	-3.5
ARKANSAS	7.9	7.2	-0.7
COLORADO	6.1	4.6	-1.6
CONNECTICUT	6.9	6.1	-0.9
DELAWARE	5.5	7.0	1.4
DISTRICT OF CO	11.1	7.2	-3.9
FLORIDA	7.5	7.9	0.4
GEORGIA	3.9	4.9	1.0
HAWAII	6.3	7.0	0.7
IDAHO	5.1	5.2	0.1
ILLINOIS	7.6	7.6	-0.1
INDIANA	5.8	6.1	0.4
IOWA	7.3	5.5	-1.7
KANSAS	7.6	8.7	1.1
KENTUCKY	5.5	5.8	0.3
LOUISIANA	6.7	4.9	-1.8
MAINE	6.5	4.9	-1.6

MARYLAND	4.6	5.0	0.4
MASSACHUSETTS	5.4	6.1	0.7
MICHIGAN	6.1	5.4	-0.8
MINNESOTA	7.5	5.7	-1.8
MISSISSIPPI	6.9	6.8	-0.2
MISSOURI	6.7	8.0	1.2
MONTANA	4.7	4.3	-0.4
NEBRASKA	10.0	8.3	-1.8
NEVADA	8.4	6.3	-2.1
NEW HAMPSHIRE	5.5	4.2	-1.3
NEW JERSEY	7.6	6.4	-1.3
NEW MEXICO	6.2	7.9	1.7
NEW YORK	6.9	7.3	0.5
NORTH CAROLINA	7.2	6.7	-0.5
NORTH DAKOTA	6.7	4.9	-1.8
OHIO	5.0	6.2	1.1
OKLAHOMA	6.6	4.9	-1.7
OREGON	5.3	5.5	0.2
PENNSYLVANIA	7.2	6.8	-0.4
RHODE ISLAND	5.8	7.3	1.4
SOUTH CAROLINA	5.7	5.5	-0.1
SOUTH DAKOTA	4.4	8.9	4.5
TENNESSEE	9.9	7.0	-2.8
TEXAS	9.2	12.1	2.8
UTAH	5.5	5.2	-0.3
VERMONT	6.2	6.6	0.4
VIRGINIA	6.3	5.0	-1.3
WASHINGTON	5.4	4.5	-0.9
WEST VIRGINIA	14.0	9.1	-4.9
WISCONSIN	6.5	5.6	-0.8
WYOMING	6.4	7.9	1.5
PUERTO RICO	12.7	10.2	-2.5

* CHANCES OF GETTING HIV MEDIUM/HIGH

N/A - NOT AVAILABLE

August 17, 1998

TABLE 32.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2138	19.7	1.0	(17.7, 21.6)
ALASKA	1543	26.0	1.7	(22.7, 29.2)
ARIZONA	1876	5.8	0.9	(4.1, 7.5)
GEORGIA	2275	14.2	0.9	(12.4, 16.0)
INDIANA	2359	15.0	0.9	(13.3, 16.7)
KANSAS	1913	13.1	0.9	(11.3, 15.0)
KENTUCKY	1700	15.0	1.1	(12.9, 17.0)
LOUISIANA	1645	13.8	1.0	(11.8, 15.8)
MONTANA	1800	21.5	1.1	(19.3, 23.6)
OHIO	3105	11.9	0.9	(10.2, 13.7)
OKLAHOMA	1883	13.0	1.0	(11.0, 15.0)
PENNSYLVANIA	3359	13.3	0.7	(11.9, 14.8)
SOUTH CAROLINA	2152	13.8	0.9	(12.1, 15.6)

VIRGINIA	3518	15.4	0.9	(13.7, 17.2)
WASHINGTON	3594	17.8	0.8	(16.3, 19.4)
WEST VIRGINIA	2421	21.6	1.0	(19.7, 23.5)
WYOMING	2409	24.0	1.1	(21.8, 26.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 17 MEDIAN = 15.0 RANGE = 5.8-26.0

* EVER USED OR TRIED SMOKELESS TOBACCO

August 3, 1998

TABLE 32.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
					*					*
ALABAMA	803	36.7	1.8	(33.1, 40.3)	*	1335	4.5	0.6	(3.4, 5.6)	
ALASKA	715	42.3	2.7	(37.0, 47.6)	*	828	8.0	1.5	(5.1, 10.9)	
ARIZONA	877	10.7	1.7	(7.4, 14.0)	*	999	1.2	0.3	(0.6, 1.8)	
GEORGIA	950	25.1	1.7	(21.8, 28.4)	*	1325	4.2	0.6	(2.9, 5.4)	

INDIANA	1062	30.3	1.6	(27.1, 33.5)	*	1297	1.1	0.3	(0.5, 1.6)
KANSAS	752	24.7	1.8	(21.2, 28.1)	*	1161	2.4	0.5	(1.5, 3.4)
KENTUCKY	626	28.8	1.9	(25.0, 32.5)	*	1074	2.4	0.5	(1.3, 3.5)
LOUISIANA	625	26.0	1.9	(22.3, 29.7)	*	1020	2.9	0.7	(1.5, 4.3)
MONTANA	759	39.0	1.9	(35.2, 42.7)	*	1041	4.8	0.8	(3.3, 6.3)
OHIO	1209	23.6	1.7	(20.2, 26.9)	*	1896	1.4	0.4	(0.7, 2.2)
OKLAHOMA	830	25.0	2.0	(21.1, 28.8)	*	1053	1.9	0.5	(1.0, 2.8)
PENNSYLVANIA	1368	26.2	1.4	(23.5, 28.9)	*	1991	1.5	0.3	(0.9, 2.1)
SOUTH CAROLINA	819	25.2	1.7	(21.9, 28.5)	*	1333	3.5	0.6	(2.4, 4.7)
VIRGINIA	1413	29.3	1.7	(26.0, 32.6)	*	2105	2.5	0.4	(1.7, 3.2)
WASHINGTON	1540	32.7	1.4	(29.9, 35.4)	*	2054	3.7	0.6	(2.4, 4.9)
WEST VIRGINIA	960	43.7	1.7	(40.3, 47.0)	*	1461	1.9	0.4	(1.1, 2.7)
WYOMING	1017	43.7	1.9	(40.0, 47.3)	*	1392	4.8	1.0	(2.9, 6.7)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	17	NUMBER OF PARTICIPANTS =	17
	MEDIAN	= 28.8		= 2.5
	RANGE	= 10.7-43.7		= 1.1- 8.0

* EVER USED OR TRIED SMOKELESS TOBACCO

August 17, 1998

TABLE 33.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2136	5.4	0.6	(4.3, 6.5)
ALASKA	1543	5.6	0.9	(3.8, 7.3)
ARIZONA	1876	1.4	0.3	(0.7, 2.1)
GEORGIA	2274	4.0	0.5	(3.0, 4.9)
INDIANA	2359	3.3	0.4	(2.4, 4.1)
KANSAS	1912	5.0	0.6	(3.8, 6.3)
KENTUCKY	1700	6.2	0.8	(4.7, 7.7)
LOUISIANA	1645	3.7	0.5	(2.7, 4.8)
MONTANA	1800	5.3	0.6	(4.0, 6.5)
OHIO	3104	2.4	0.4	(1.7, 3.2)
OKLAHOMA	1883	3.8	0.6	(2.7, 4.9)
PENNSYLVANIA	3359	3.8	0.4	(2.9, 4.6)
SOUTH CAROLINA	2152	2.8	0.4	(2.0, 3.7)
VIRGINIA	3517	3.0	0.3	(2.3, 3.6)
WASHINGTON	3594	2.9	0.4	(2.1, 3.6)
WEST VIRGINIA	2421	8.7	0.7	(7.4, 10.1)
WYOMING	2409	7.6	0.6	(6.4, 8.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 17 MEDIAN = 3.8 RANGE = 1.4- 8.7

* CURRENT USERS OF SMOKELESS TOBACCO

August 17, 1998

TABLE 33.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-MALES-				-FEMALES-			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
	*	*	*	*	*	*	*	*
ALABAMA	802	9.9	1.1	(7.6, 12.1)	*	1334	1.4	0.4 (0.7, 2.1)
ALASKA	715	9.2	1.6	(6.1, 12.4)	*	828	1.6	0.5 (0.6, 2.5)
ARIZONA	877	2.6	0.7	(1.2, 3.9)	*	999	0.3	0.2 (-0.0, 0.6)
GEORGIA	950	6.4	0.9	(4.6, 8.2)	*	1324	1.7	0.5 (0.8, 2.6)
INDIANA	1062	6.8	0.9	(5.1, 8.5)	*	1297	0.0	0.0 (0.0, 0.0)
KANSAS	751	10.2	1.3	(7.8, 12.7)	*	1161	0.2	0.2 (-0.1, 0.5)
KENTUCKY	626	12.3	1.5	(9.3, 15.3)	*	1074	0.6	0.2 (0.1, 1.1)
LOUISIANA	625	7.5	1.1	(5.4, 9.7)	*	1020	0.3	0.2 (-0.1, 0.8)
MONTANA	759	10.5	1.3	(8.1, 13.0)	*	1041	0.2	0.2 (-0.1, 0.6)
OHIO	1209	5.1	0.8	(3.5, 6.7)	*	1895	0.0	0.0 (-0.0, 0.1)
OKLAHOMA	830	7.7	1.1	(5.5, 9.9)	*	1053	0.3	0.1 (-0.0, 0.5)
PENNSYLVANIA	1368	7.5	0.9	(5.7, 9.2)	*	1991	0.4	0.2 (0.1, 0.7)
SOUTH CAROLINA	819	4.8	0.9	(3.1, 6.5)	*	1333	1.0	0.3 (0.4, 1.6)
VIRGINIA	1412	6.1	0.7	(4.7, 7.4)	*	2105	0.1	0.1 (-0.0, 0.2)
WASHINGTON	1540	5.6	0.7	(4.2, 7.0)	*	2054	0.2	0.1 (-0.0, 0.4)
WEST VIRGINIA	960	18.4	1.4	(15.7, 21.0)	*	1461	0.2	0.1 (-0.0, 0.3)
WYOMING	1017	14.7	1.2	(12.3, 17.0)	*	1392	0.7	0.2 (0.2, 1.1)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 17
 MEDIAN = 7.5
 RANGE = 2.6-18.4

NUMBER OF PARTICIPANTS = 17
 MEDIAN = 0.3
 RANGE = 0.0- 1.7

* CURRENT USERS OF SMOKELESS TOBACCO

August 17, 1998

TABLE 34.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET*
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	126	50	1.5	(47.4, 53.3)
ALASKA	53	46	2.8	(41.0, 51.8)
ARIZONA	60	46	3.2	(39.6, 52.1)
ARKANSAS	83	49	2.1	(44.5, 52.6)
CALIFORNIA	210	46	1.2	(43.8, 48.5)
COLORADO	71	43	3.3	(36.5, 49.4)
DISTRICT OF CO	64	46	1.6	(43.0, 49.2)
FLORIDA	203	50	1.1	(47.9, 52.2)
GEORGIA	89	49	1.6	(45.6, 51.7)
HAWAII	107	49	1.7	(45.5, 52.1)
IDAHO	203	51	1.5	(47.7, 53.5)
INDIANA	117	48	1.4	(45.4, 51.0)
IOWA	172	54	1.3	(51.2, 56.2)
KANSAS	61	48	2.6	(42.4, 52.6)
KENTUCKY	212	48	1.2	(45.3, 49.8)
LOUISIANA	89	50	1.9	(46.6, 54.0)
MAINE	73	51	2.0	(47.3, 55.2)
MASSACHUSETTS	69	48	2.1	(44.2, 52.3)
MICHIGAN	147	46	1.5	(43.2, 49.2)
MINNESOTA	195	50	1.2	(48.1, 52.8)
MISSISSIPPI	101	48	1.8	(44.7, 51.6)
MISSOURI	93	50	1.9	(46.2, 53.5)
MONTANA	63	46	2.3	(41.8, 50.8)
NEBRASKA	128	50	1.9	(46.3, 53.9)
NEVADA	107	48	3.8	(40.4, 55.2)
NEW HAMPSHIRE	60	50	2.1	(45.9, 54.1)
NEW JERSEY	127	49	1.7	(45.7, 52.4)
NEW MEXICO	89	49	1.9	(45.0, 52.3)
NORTH CAROLINA	193	49	1.2	(46.8, 51.3)
NORTH DAKOTA	73	49	2.5	(44.2, 54.0)
OHIO	183	49	1.8	(45.1, 52.0)
OREGON	162	48	1.3	(45.4, 50.5)
PENNSYLVANIA	171	50	1.3	(47.3, 52.4)
SOUTH CAROLINA	118	46	1.4	(43.7, 49.1)
SOUTH DAKOTA	86	49	2.1	(44.5, 52.6)
TENNESSEE	131	44	1.5	(41.3, 47.2)
TEXAS	134	48	1.4	(45.3, 50.9)
UTAH	106	48	1.9	(43.8, 51.4)
VERMONT	135	44	4.0	(36.5, 52.2)
VIRGINIA	176	47	1.5	(44.3, 50.0)
WEST VIRGINIA	159	50	1.5	(46.7, 52.7)
WISCONSIN	97	47	2.8	(41.9, 52.7)
WYOMING	79	49	2.3	(44.4, 53.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 43 MEDIAN = 48.6 RANGE = 43.0-53.7

* MEAN AGE AT ONSET OF DIABETES

August 17, 1998

TABLE 34.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET* -BY SEX
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	43	50	2.7	(44.2, 54.9)	*	83	51	1.8	(47.5, 54.4)	*
ALASKA	21	49	4.4	(40.7, 57.9)	*	32	44	2.8	(38.4, 49.5)	*
ARIZONA	30	50	3.7	(43.2, 57.6)	*	30	43	4.6	(34.3, 52.2)	*
ARKANSAS	35	50	2.8	(44.7, 55.7)	*	48	47	2.9	(41.1, 52.5)	*
CALIFORNIA	95	47	1.8	(43.0, 50.1)	*	115	46	1.6	(42.6, 49.0)	*
COLORADO	34	47	2.9	(41.2, 52.6)	*	37	39	5.4	(28.6, 49.7)	*
DISTRICT OF CO	22	46	2.3	(41.8, 50.6)	*	42	46	2.1	(41.9, 50.3)	*
FLORIDA	100	50	1.5	(46.6, 52.6)	*	103	51	1.5	(47.6, 53.6)	*
GEORGIA	28	51	2.5	(46.3, 56.2)	*	61	47	1.9	(43.6, 51.2)	*
HAWAII	41	48	2.6	(42.8, 52.9)	*	66	49	2.1	(45.3, 53.7)	*
IDAHO	87	50	2.3	(45.7, 54.5)	*	116	51	1.8	(47.7, 54.6)	*
INDIANA	46	46	2.1	(42.2, 50.4)	*	71	50	1.9	(46.0, 53.4)	*
IOWA	62	54	1.6	(50.4, 56.9)	*	110	54	1.9	(50.0, 57.6)	*
KANSAS	21	47	4.5	(37.9, 55.7)	*	40	48	3.0	(42.1, 54.0)	*
KENTUCKY	85	46	1.7	(42.8, 49.5)	*	127	49	1.5	(46.2, 52.0)	*
LOUISIANA	36	50	2.8	(44.7, 55.6)	*	53	50	2.4	(45.7, 55.1)	*
MAINE	38	53	2.5	(48.4, 58.1)	*	35	49	3.1	(42.9, 55.1)	*
MASSACHUSETTS	30	45	2.7	(39.4, 50.2)	*	39	52	3.0	(46.0, 57.6)	*
MICHIGAN	56	49	2.2	(44.7, 53.4)	*	91	44	2.0	(40.1, 48.0)	*
MINNESOTA	86	50	1.8	(46.6, 53.5)	*	109	51	1.7	(47.5, 54.3)	*
MISSISSIPPI	29	48	3.5	(40.9, 54.6)	*	72	48	1.6	(45.3, 51.6)	*
MISSOURI	42	51	2.3	(46.7, 55.8)	*	51	48	2.7	(43.0, 53.6)	*
MONTANA	28	45	3.0	(38.9, 50.7)	*	35	48	3.5	(41.3, 54.9)	*
NEBRASKA	46	51	2.8	(45.9, 56.7)	*	82	49	2.7	(43.9, 54.4)	*
NEVADA	53	48	5.3	(37.6, 58.2)	*	54	48	3.8	(40.1, 55.1)	*
NEW HAMPSHIRE	24	50	1.9	(46.3, 53.9)	*	36	50	3.3	(43.4, 56.4)	*
NEW JERSEY	61	49	2.2	(44.8, 53.3)	*	66	49	2.6	(43.9, 54.1)	*
NEW MEXICO	44	50	2.6	(45.3, 55.3)	*	45	47	2.8	(41.4, 52.1)	*
NORTH CAROLINA	73	51	1.8	(47.3, 54.3)	*	120	48	1.5	(44.7, 50.5)	*
NORTH DAKOTA	30	44	4.1	(35.7, 51.7)	*	43	54	2.3	(49.6, 58.7)	*
OHIO	65	49	2.1	(44.9, 53.1)	*	118	48	2.5	(43.3, 53.2)	*
OREGON	68	49	1.9	(45.4, 52.6)	*	94	47	1.8	(43.5, 50.5)	*
PENNSYLVANIA	73	48	1.8	(44.8, 51.7)	*	98	51	1.9	(47.7, 55.0)	*
SOUTH CAROLINA	41	45	2.1	(41.0, 49.2)	*	77	47	1.8	(43.9, 50.8)	*
SOUTH DAKOTA	32	48	3.2	(41.8, 54.5)	*	54	49	2.6	(43.8, 53.9)	*
TENNESSEE	44	44	2.5	(39.1, 48.8)	*	87	44	1.9	(40.7, 48.1)	*
TEXAS	47	48	2.2	(44.1, 52.6)	*	87	48	1.9	(44.3, 51.7)	*
UTAH	48	48	2.9	(42.0, 53.2)	*	58	48	2.4	(42.8, 52.4)	*
VERMONT	62	40	6.1	(28.4, 52.4)	*	73	48	2.7	(43.2, 53.7)	*
VIRGINIA	64	46	2.0	(41.6, 49.6)	*	112	48	2.0	(44.5, 52.3)	*

WEST VIRGINIA	71	49	2.4	(44.5, 53.9)	*	88	50	1.8	(46.6, 53.8)
WISCONSIN	40	45	4.3	(36.6, 53.6)	*	57	50	2.7	(44.7, 55.4)
WYOMING	26	50	3.5	(42.9, 56.8)	*	53	48	3.1	(42.4, 54.3)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	43	NUMBER OF PARTICIPANTS =	43
	MEDIAN	= 49.0	MEDIAN	= 48.4
	RANGE	= 40.4-53.7	RANGE	= 39.2-54.2

* MEAN AGE AT ONSET OF DIABETES

August 17, 1998

TABLE 35.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TAKING INSULIN*
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	150	41.3	4.7	(32.2, 50.5)
ALASKA	54	38.9	10.0	(19.3, 58.6)
ARIZONA	61	28.6	8.3	(12.3, 44.9)
ARKANSAS	93	40.8	5.8	(29.5, 52.1)
CALIFORNIA	212	23.1	3.4	(16.5, 29.6)
COLORADO	72	39.3	7.4	(24.8, 53.8)
DISTRICT OF CO	69	44.5	7.0	(30.8, 58.2)
FLORIDA	205	35.5	3.7	(28.2, 42.7)
GEORGIA	95	34.1	5.2	(23.9, 44.3)
HAWAII	112	19.4	4.4	(10.8, 28.0)
IDAHO	205	32.6	3.9	(24.9, 40.3)
INDIANA	123	33.7	4.8	(24.4, 43.1)
IOWA	181	39.9	4.1	(31.8, 48.0)
KANSAS	66	30.5	6.4	(17.9, 43.1)
KENTUCKY	219	39.5	3.7	(32.3, 46.7)
LOUISIANA	92	30.0	5.6	(19.1, 40.9)
MAINE	80	29.1	5.6	(18.0, 40.1)
MASSACHUSETTS	73	34.3	6.2	(22.2, 46.4)
MICHIGAN	157	35.6	4.1	(27.6, 43.6)
MINNESOTA	200	39.8	3.8	(32.4, 47.1)
MISSISSIPPI	108	36.3	5.2	(26.1, 46.5)
MISSOURI	98	28.5	5.3	(18.1, 38.9)
MONTANA	65	44.3	6.8	(31.0, 57.5)
NEBRASKA	135	42.9	5.3	(32.4, 53.3)
NEVADA	109	33.6	10.0	(14.0, 53.1)
NEW HAMPSHIRE	64	39.4	7.0	(25.7, 53.0)
NEW JERSEY	137	38.6	4.8	(29.2, 48.0)
NEW MEXICO	93	39.0	5.6	(27.9, 50.0)
NORTH CAROLINA	208	31.5	3.5	(24.7, 38.3)
NORTH DAKOTA	76	39.3	5.9	(27.7, 50.9)
OHIO	193	34.4	4.6	(25.3, 43.4)

OREGON	168	29.8	4.3	(21.3, 38.3)
PENNSYLVANIA	184	33.3	3.8	(25.7, 40.8)
SOUTH CAROLINA	123	41.3	5.0	(31.5, 51.2)
SOUTH DAKOTA	88	34.0	5.4	(23.5, 44.5)
TENNESSEE	150	41.8	4.5	(33.0, 50.6)
TEXAS	149	30.8	4.2	(22.6, 38.9)
UTAH	109	34.7	5.9	(23.1, 46.3)
VERMONT	144	43.0	7.4	(28.4, 57.5)
VIRGINIA	194	32.3	4.4	(23.7, 41.0)
WEST VIRGINIA	163	33.6	4.1	(25.6, 41.5)
WISCONSIN	99	35.1	7.1	(21.1, 49.0)
WYOMING	82	27.4	5.6	(16.6, 38.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 43 MEDIAN = 34.7 RANGE = 19.4-44.5

* NOW TAKING INSULIN

August 17, 1998

TABLE 35.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TAKING INSULIN* -BY SEX
AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	-MALES-					-FEMALES-				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
					*				*	
ALABAMA	46	36.7	8.1	(20.9, 52.5)	*	104	44.2	5.5	(33.4, 55.0)	
ALASKA	21	23.6	13.2	(-2.2, 49.4)	*	33	52.0	13.1	(26.3, 77.7)	
ARIZONA	30	16.4	6.8	(3.2, 29.7)	*	31	35.4	12.2	(11.5, 59.3)	
ARKANSAS	39	44.9	8.6	(28.1, 61.8)	*	54	36.2	7.4	(21.7, 50.6)	
CALIFORNIA	95	19.5	4.8	(10.1, 29.0)	*	117	26.3	4.7	(17.1, 35.4)	
COLORADO	34	31.7	9.1	(13.9, 49.4)	*	38	46.6	10.9	(25.3, 67.9)	
DISTRICT OF CO	23	41.5	11.9	(18.2, 64.8)	*	46	46.4	8.5	(29.8, 63.0)	
FLORIDA	101	34.4	5.2	(24.2, 44.7)	*	104	36.8	5.1	(26.9, 46.8)	
GEORGIA	29	31.1	9.3	(12.9, 49.2)	*	66	35.5	6.2	(23.3, 47.7)	
HAWAII	42	13.4	6.6	(0.5, 26.4)	*	70	24.2	5.8	(12.8, 35.5)	
IDAHO	88	35.7	5.9	(24.1, 47.3)	*	117	28.8	4.7	(19.5, 38.0)	
INDIANA	49	28.0	7.3	(13.7, 42.3)	*	74	38.4	6.1	(26.4, 50.4)	
IOWA	64	30.7	6.1	(18.7, 42.7)	*	117	47.3	5.1	(37.3, 57.2)	
KANSAS	23	25.8	10.7	(4.9, 46.7)	*	43	33.9	8.0	(18.3, 49.5)	
KENTUCKY	86	35.4	5.5	(24.7, 46.1)	*	133	43.7	4.8	(34.3, 53.1)	
LOUISIANA	38	26.7	8.1	(10.8, 42.6)	*	54	33.0	7.2	(18.8, 47.1)	
MAINE	38	22.1	7.4	(7.6, 36.5)	*	42	35.5	8.3	(19.3, 51.7)	
MASSACHUSETTS	31	47.5	9.7	(28.4, 66.6)	*	42	21.3	6.5	(8.6, 33.9)	
MICHIGAN	59	22.7	5.6	(11.7, 33.7)	*	98	45.1	5.5	(34.3, 55.8)	
MINNESOTA	91	38.7	5.6	(27.9, 49.6)	*	109	40.7	5.1	(30.7, 50.8)	
MISSISSIPPI	29	24.3	8.7	(7.2, 41.3)	*	79	44.1	6.2	(32.0, 56.2)	
MISSOURI	43	27.9	7.9	(12.4, 43.4)	*	55	29.3	6.2	(17.2, 41.4)	

MONTANA	30	40.1	9.5	(21.6, 58.7)	*	35	49.3	9.3	(31.0, 67.5)
NEBRASKA	46	45.6	8.9	(28.2, 63.0)	*	89	40.8	6.5	(28.1, 53.5)
NEVADA	52	28.9	12.4	(4.7, 53.1)	*	57	43.4	13.9	(16.2, 70.7)
NEW HAMPSHIRE	25	35.3	11.0	(13.7, 56.9)	*	39	42.9	9.0	(25.4, 60.5)
NEW JERSEY	67	38.2	6.5	(25.6, 50.9)	*	70	39.0	7.0	(25.2, 52.7)
NEW MEXICO	45	36.4	7.8	(21.2, 51.6)	*	48	41.8	7.8	(26.6, 57.0)
NORTH CAROLINA	74	26.7	5.3	(16.3, 37.0)	*	134	35.1	4.5	(26.3, 44.0)
NORTH DAKOTA	32	36.0	8.7	(18.9, 53.1)	*	44	42.6	7.8	(27.4, 57.8)
OHIO	68	32.7	6.6	(19.7, 45.7)	*	125	35.4	6.2	(23.2, 47.6)
OREGON	68	25.5	5.7	(14.3, 36.6)	*	100	33.6	6.2	(21.3, 45.8)
PENNSYLVANIA	78	27.8	5.5	(16.9, 38.6)	*	106	38.1	5.3	(27.8, 48.4)
SOUTH CAROLINA	43	30.7	7.3	(16.3, 45.1)	*	80	50.1	6.2	(38.1, 62.2)
SOUTH DAKOTA	33	27.8	8.4	(11.5, 44.2)	*	55	38.8	6.9	(25.4, 52.3)
TENNESSEE	46	36.9	7.8	(21.6, 52.1)	*	104	44.7	5.4	(34.1, 55.2)
TEXAS	52	34.7	7.0	(20.9, 48.5)	*	97	28.0	5.1	(18.1, 38.0)
UTAH	50	29.0	8.1	(13.1, 44.9)	*	59	41.4	9.2	(23.4, 59.4)
VERMONT	64	48.4	11.0	(26.8, 69.9)	*	80	37.5	6.1	(25.5, 49.5)
VIRGINIA	67	38.5	8.0	(22.9, 54.1)	*	127	27.5	4.6	(18.6, 36.4)
WEST VIRGINIA	74	34.7	6.2	(22.5, 46.9)	*	89	32.3	5.3	(21.9, 42.7)
WISCONSIN	39	32.8	11.7	(9.9, 55.8)	*	60	37.7	7.6	(22.8, 52.5)
WYOMING	27	23.4	8.8	(6.1, 40.7)	*	55	30.2	7.1	(16.3, 44.1)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 43
 MEDIAN = 31.7
 RANGE = 13.4-48.4

NUMBER OF PARTICIPANTS = 43
 MEDIAN = 38.1
 RANGE = 21.3-52.0

* NOW TAKING INSULIN

August 17, 1998

TABLE 36.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR HEALTH PROF VISITS FOR DIABETES*
 AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	132	5	0.7	(4.0, 6.7)
ALASKA	49	4	0.8	(2.7, 5.7)
ARIZONA	60	3	0.5	(2.2, 4.1)
ARKANSAS	88	6	1.0	(4.3, 8.2)
CALIFORNIA	212	6	0.7	(4.4, 7.2)
COLORADO	70	4	0.8	(2.7, 5.8)
DISTRICT OF CO	65	4	0.4	(3.5, 5.0)
FLORIDA	197	6	0.5	(4.9, 7.0)
GEORGIA	87	4	0.4	(3.3, 5.0)
HAWAII	107	5	0.6	(4.2, 6.4)
IDAHO	203	4	0.4	(3.4, 5.0)
INDIANA	120	5	0.7	(3.5, 6.4)

IOWA	176	4	0.4	(3.6, 5.2)
KANSAS	64	3	0.6	(2.1, 4.5)
KENTUCKY	201	6	0.6	(4.5, 6.9)
LOUISIANA	88	4	0.4	(3.1, 4.6)
MAINE	76	3	0.4	(2.5, 4.1)
MASSACHUSETTS	72	4	0.4	(3.3, 4.8)
MICHIGAN	150	4	0.4	(3.4, 4.9)
MINNESOTA	197	4	0.3	(3.1, 4.3)
MISSISSIPPI	99	6	0.8	(4.2, 7.3)
MISSOURI	94	4	0.8	(2.7, 5.7)
MONTANA	61	6	1.0	(4.5, 8.4)
NEBRASKA	133	6	0.7	(4.2, 6.9)
NEVADA	105	6	1.6	(2.6, 8.8)
NEW HAMPSHIRE	61	4	0.7	(2.7, 5.3)
NEW JERSEY	129	5	0.4	(3.8, 5.3)
NEW MEXICO	88	5	0.8	(3.5, 6.6)
NORTH CAROLINA	191	5	0.5	(3.6, 5.4)
NORTH DAKOTA	75	4	0.5	(3.0, 4.8)
OHIO	191	4	0.3	(4.0, 5.0)
OREGON	162	4	0.5	(3.5, 5.3)
PENNSYLVANIA	177	5	0.6	(3.6, 5.8)
SOUTH CAROLINA	115	5	0.5	(4.2, 6.2)
SOUTH DAKOTA	86	4	0.4	(2.8, 4.4)
TENNESSEE	134	5	0.4	(4.3, 6.1)
TEXAS	138	5	0.5	(3.6, 5.4)
UTAH	108	5	0.9	(3.5, 6.9)
VERMONT	133	5	0.5	(3.5, 5.5)
VIRGINIA	178	4	0.4	(3.6, 5.3)
WEST VIRGINIA	160	4	0.3	(3.7, 4.9)
WISCONSIN	97	4	0.5	(2.5, 4.5)
WYOMING	76	3	0.4	(2.4, 4.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 43 MEDIAN = 4.4 RANGE = 3.1- 6.5

* MEAN TIMES SEEN HEALTH PROF FOR DIABETES IN PAST YEAR

August 17, 1998

TABLE 36.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR HEALTH PROF VISITS FOR DIABETES* -BY SEX
AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
					*					*
ALABAMA	43	4	0.8	(2.4, 5.4)	*	89	6	1.0	(4.4, 8.3)	*
ALASKA	20	3	0.5	(2.3, 4.3)	*	29	5	1.3	(2.4, 7.3)	*
ARIZONA	29	2	0.4	(1.3, 2.8)	*	31	4	0.7	(2.4, 5.1)	*
ARKANSAS	38	6	1.3	(3.1, 8.1)	*	50	7	1.6	(4.0, 10.1)	*

CALIFORNIA	95	5	0.8	(3.2, 6.4)	*	117	7	1.1	(4.5, 8.9)
COLORADO	34	4	1.0	(1.7, 5.8)	*	36	5	1.2	(2.4, 7.1)
DISTRICT OF CO	23	4	0.6	(2.9, 5.4)	*	42	4	0.4	(3.5, 5.2)
FLORIDA	98	6	0.8	(4.8, 7.9)	*	99	5	0.6	(4.3, 6.7)
GEORGIA	26	4	0.5	(2.6, 4.7)	*	61	4	0.6	(3.2, 5.5)
HAWAII	38	5	0.7	(3.6, 6.3)	*	69	6	0.9	(3.8, 7.3)
IDAHO	88	4	0.7	(2.9, 5.6)	*	115	4	0.4	(3.4, 4.8)
INDIANA	49	5	1.3	(2.1, 7.0)	*	71	5	0.8	(3.7, 7.0)
IOWA	62	4	0.6	(3.2, 5.5)	*	114	4	0.5	(3.5, 5.4)
KANSAS	23	3	0.7	(1.6, 4.3)	*	41	4	1.0	(1.7, 5.5)
KENTUCKY	81	6	1.1	(3.4, 7.6)	*	120	6	0.6	(4.7, 6.9)
LOUISIANA	37	4	0.6	(2.5, 4.9)	*	51	4	0.5	(3.0, 4.9)
MAINE	38	3	0.4	(2.1, 3.8)	*	38	4	0.7	(2.2, 5.0)
MASSACHUSETTS	31	4	0.6	(3.2, 5.4)	*	41	4	0.5	(2.9, 4.9)
MICHIGAN	58	3	0.5	(2.4, 4.2)	*	92	5	0.5	(3.8, 5.8)
MINNESOTA	90	3	0.4	(2.5, 4.1)	*	107	4	0.5	(3.2, 5.0)
MISSISSIPPI	28	4	1.0	(2.5, 6.2)	*	71	7	1.1	(4.5, 8.9)
MISSOURI	41	5	1.4	(2.1, 7.5)	*	53	4	0.3	(2.9, 4.2)
MONTANA	29	6	1.3	(3.7, 8.6)	*	32	7	1.5	(3.8, 9.8)
NEBRASKA	45	6	1.3	(3.0, 8.2)	*	88	6	0.7	(4.1, 6.9)
NEVADA	51	6	2.4	(1.4, 10.7)	*	54	5	1.0	(3.0, 7.0)
NEW HAMPSHIRE	25	2	0.4	(1.7, 3.1)	*	36	5	1.2	(3.2, 7.7)
NEW JERSEY	62	4	0.6	(3.2, 5.4)	*	67	5	0.5	(3.8, 5.8)
NEW MEXICO	42	5	1.4	(2.7, 8.0)	*	46	5	0.7	(3.4, 6.2)
NORTH CAROLINA	68	4	0.9	(2.7, 6.3)	*	123	5	0.4	(3.8, 5.3)
NORTH DAKOTA	32	3	0.6	(2.3, 4.6)	*	43	4	0.7	(2.9, 5.7)
OHIO	68	4	0.3	(2.9, 4.1)	*	123	5	0.4	(4.4, 5.8)
OREGON	67	4	0.7	(3.2, 5.8)	*	95	4	0.6	(3.1, 5.5)
PENNSYLVANIA	76	4	0.6	(2.8, 5.0)	*	101	5	0.9	(3.7, 7.2)
SOUTH CAROLINA	41	5	0.8	(3.1, 6.4)	*	74	6	0.7	(4.3, 6.9)
SOUTH DAKOTA	32	4	0.7	(2.4, 5.3)	*	54	3	0.4	(2.6, 4.2)
TENNESSEE	40	5	0.7	(3.4, 6.0)	*	94	5	0.6	(4.4, 6.6)
TEXAS	49	3	0.6	(2.1, 4.5)	*	89	5	0.7	(4.1, 6.6)
UTAH	50	6	1.4	(3.4, 8.8)	*	58	4	0.8	(2.5, 5.7)
VERMONT	62	4	0.7	(2.6, 5.1)	*	71	5	0.8	(3.7, 6.8)
VIRGINIA	63	4	0.7	(3.1, 5.8)	*	115	4	0.5	(3.4, 5.4)
WEST VIRGINIA	74	5	0.5	(3.6, 5.5)	*	86	4	0.4	(3.4, 4.8)
WISCONSIN	40	3	0.4	(2.0, 3.7)	*	57	4	0.9	(2.4, 6.1)
WYOMING	26	2	0.4	(1.6, 3.1)	*	50	4	0.7	(2.6, 5.2)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 43
 MEDIAN = 4.3
 RANGE = 2.1- 6.3

NUMBER OF PARTICIPANTS = 43
 MEDIAN = 4.8
 RANGE = 3.4- 7.1

* MEAN TIMES SEEN HEALTH PROF FOR DIABETES IN PAST YEAR

August 17, 1998

TABLE 37.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR RECENT DENTAL VISIT*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2107	66.6	1.2	(64.2, 68.9)
CALIFORNIA	4058	63.3	0.9	(61.5, 65.0)
COLORADO	1765	66.6	1.3	(64.1, 69.1)
FLORIDA	3426	68.5	0.9	(66.7, 70.3)
IDAHO	4897	64.8	0.9	(63.1, 66.6)
INDIANA	2352	66.8	1.1	(64.6, 69.0)
MARYLAND	2165	70.3	1.2	(67.9, 72.7)
MISSISSIPPI	1559	60.3	1.5	(57.4, 63.2)
MISSOURI	1832	61.5	1.4	(58.8, 64.3)
MONTANA	1777	67.0	1.3	(64.5, 69.5)
NEVADA	2498	64.5	2.0	(60.5, 68.4)
NEW JERSEY	2465	74.9	1.1	(72.8, 77.0)
NEW MEXICO	1776	68.1	1.2	(65.7, 70.5)
NEW YORK	3352	71.3	0.9	(69.5, 73.0)
OHIO	3133	71.5	1.1	(69.3, 73.6)
TENNESSEE	2878	66.4	1.0	(64.5, 68.3)
TEXAS	2277	61.2	1.2	(58.9, 63.5)
UTAH	2844	71.7	1.1	(69.5, 73.9)
VIRGINIA	3416	70.6	1.1	(68.3, 72.8)
WEST VIRGINIA	2402	58.4	1.1	(56.2, 60.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 20 MEDIAN = 66.7 RANGE = 58.4-74.9

* PERSONS VISITING A DENTIST OR DENTAL CLINIC WITHIN THE PAST YEAR

August 17, 1998

TABLE 37.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR RECENT DENTAL VISIT* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	797	64.9	2.0	(61.1, 68.8)	*	1310	68.0	1.5 (65.2, 70.9)
CALIFORNIA	1757	60.0	1.4	(57.3, 62.6)	*	2301	66.5	1.1 (64.3, 68.8)
COLORADO	795	64.1	1.9	(60.3, 67.9)	*	970	69.0	1.7 (65.6, 72.4)
FLORIDA	1408	67.0	1.4	(64.2, 69.8)	*	2018	69.9	1.1 (67.6, 72.1)
IDAHO	2052	62.5	1.4	(59.8, 65.2)	*	2845	67.1	1.2 (64.8, 69.4)
INDIANA	1060	63.8	1.7	(60.4, 67.1)	*	1292	69.5	1.5 (66.6, 72.4)
MARYLAND	904	68.4	1.9	(64.6, 72.2)	*	1261	72.0	1.5 (69.0, 74.9)
MISSISSIPPI	578	58.7	2.5	(53.8, 63.6)	*	981	61.7	1.7 (58.4, 65.0)
MISSOURI	693	58.0	2.3	(53.6, 62.4)	*	1139	64.7	1.7 (61.5, 68.0)
MONTANA	746	65.9	1.9	(62.1, 69.6)	*	1031	68.1	1.6 (64.9, 71.3)
NEVADA	1164	59.9	3.2	(53.5, 66.2)	*	1334	69.2	2.3 (64.8, 73.6)
NEW JERSEY	990	71.8	1.7	(68.4, 75.1)	*	1475	77.8	1.3 (75.3, 80.2)
NEW MEXICO	741	65.8	1.9	(62.1, 69.5)	*	1035	70.3	1.5 (67.2, 73.3)
NEW YORK	1342	69.9	1.4	(67.2, 72.7)	*	2010	72.5	1.1 (70.3, 74.6)
OHIO	1220	69.1	1.7	(65.7, 72.5)	*	1913	73.6	1.3 (71.0, 76.1)
TENNESSEE	1006	64.9	1.7	(61.7, 68.2)	*	1872	67.7	1.2 (65.4, 69.9)
TEXAS	939	59.4	1.8	(55.9, 62.9)	*	1338	63.0	1.5 (60.1, 65.8)
UTAH	1268	68.6	1.6	(65.4, 71.8)	*	1576	74.7	1.4 (71.9, 77.4)
VIRGINIA	1384	67.7	1.8	(64.1, 71.3)	*	2032	73.3	1.4 (70.7, 76.0)
WEST VIRGINIA	951	57.8	1.7	(54.4, 61.2)	*	1451	59.0	1.4 (56.2, 61.8)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 20
 MEDIAN = 64.9
 RANGE = 57.8-71.8

NUMBER OF PARTICIPANTS = 20
 MEDIAN = 69.1
 RANGE = 59.0-77.8

* PERSONS VISITING A DENTIST OR DENTAL CLINIC WITHIN THE PAST YEAR

August 17, 1998

TABLE 38.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOOTH LOSS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2103	28.1	1.2	(25.8, 30.3)
CALIFORNIA	4027	14.5	0.6	(13.3, 15.7)
COLORADO	1745	15.7	0.9	(13.8, 17.5)
FLORIDA	3399	23.5	0.8	(21.9, 25.2)
IDAHO	4861	18.3	0.7	(17.0, 19.6)
INDIANA	2337	22.5	1.0	(20.6, 24.4)
MARYLAND	2160	18.7	1.0	(16.7, 20.7)
MISSISSIPPI	1558	29.8	1.3	(27.3, 32.4)
MISSOURI	1827	27.1	1.2	(24.8, 29.5)
MONTANA	1785	21.8	1.1	(19.7, 23.8)
NEVADA	2467	21.4	1.6	(18.3, 24.6)
NEW JERSEY	2426	21.3	1.0	(19.3, 23.2)
NEW MEXICO	1768	15.7	0.9	(13.9, 17.5)
NEW YORK	3352	20.0	0.8	(18.5, 21.6)
OHIO	3109	25.1	1.0	(23.1, 27.1)
TENNESSEE	2874	23.5	0.9	(21.8, 25.3)
TEXAS	2270	17.0	0.9	(15.2, 18.7)
UTAH	2828	12.1	0.8	(10.7, 13.6)
VIRGINIA	3400	16.7	0.8	(15.1, 18.3)
WEST VIRGINIA	2406	37.3	1.1	(35.2, 39.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 20 MEDIAN = 21.3 RANGE = 12.1-37.3

* PERSONS WITH LOSS OF 6 OR MORE TEETH FROM DECAY OR GUM DISEASE

August 17, 1998

TABLE 38.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TOOTH LOSS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	793	26.6	1.8	(23.1, 30.1)	*	1310	29.3	1.5	(26.4, 32.2)	
CALIFORNIA	1745	13.8	0.9	(12.1, 15.6)	*	2282	15.2	0.9	(13.6, 16.9)	
COLORADO	787	15.3	1.4	(12.5, 18.0)	*	958	16.0	1.3	(13.5, 18.5)	
FLORIDA	1393	22.0	1.2	(19.6, 24.3)	*	2006	25.0	1.1	(22.8, 27.1)	
IDAHO	2040	17.2	1.0	(15.2, 19.2)	*	2821	19.4	0.9	(17.7, 21.1)	
INDIANA	1052	21.1	1.4	(18.4, 23.9)	*	1285	23.8	1.3	(21.2, 26.4)	
MARYLAND	905	19.3	1.6	(16.1, 22.4)	*	1255	18.3	1.3	(15.7, 20.8)	
MISSISSIPPI	577	27.3	2.0	(23.4, 31.2)	*	981	32.0	1.6	(28.8, 35.2)	
MISSOURI	690	25.4	1.8	(21.9, 29.0)	*	1137	28.7	1.6	(25.6, 31.7)	
MONTANA	754	20.3	1.5	(17.3, 23.2)	*	1031	23.2	1.4	(20.4, 26.0)	
NEVADA	1150	19.4	2.4	(14.8, 24.0)	*	1317	23.5	2.2	(19.3, 27.7)	
NEW JERSEY	978	20.0	1.5	(17.0, 22.9)	*	1448	22.4	1.3	(19.9, 24.9)	
NEW MEXICO	736	14.4	1.3	(11.8, 17.0)	*	1032	16.9	1.2	(14.6, 19.3)	
NEW YORK	1338	18.7	1.2	(16.3, 21.1)	*	2014	21.2	1.1	(19.2, 23.3)	
OHIO	1216	22.8	1.5	(19.9, 25.7)	*	1893	27.2	1.4	(24.6, 29.9)	
TENNESSEE	1008	20.9	1.4	(18.2, 23.6)	*	1866	25.8	1.1	(23.6, 28.0)	
TEXAS	935	15.2	1.3	(12.7, 17.8)	*	1335	18.6	1.2	(16.3, 21.0)	
UTAH	1264	10.4	1.0	(8.4, 12.3)	*	1564	13.8	1.1	(11.7, 16.0)	
VIRGINIA	1378	14.3	1.2	(12.0, 16.6)	*	2022	19.0	1.2	(16.6, 21.3)	
WEST VIRGINIA	954	34.3	1.7	(31.1, 37.5)	*	1452	40.0	1.4	(37.2, 42.8)	

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 20
 MEDIAN = 19.7
 RANGE = 10.4-34.3

NUMBER OF PARTICIPANTS = 20
 MEDIAN = 22.8
 RANGE = 13.8-40.0

* PERSONS WITH LOSS OF 6 OR MORE TEETH FROM DECAY OR GUM DISEASE

August 17, 1998

TABLE 39.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV: CONDOM EFFECTIVENESS*
 DENOMINATOR IS PERSONS AGE 18-49 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	903	44.4	2.5	(39.6, 49.2)
DELAWARE	1280	47.6	1.7	(44.3, 50.8)
DISTRICT OF CO	823	61.6	2.1	(57.4, 65.8)
FLORIDA	1641	45.0	1.4	(42.3, 47.8)
ILLINOIS	812	41.0	1.9	(37.2, 44.8)
IOWA	1748	41.1	1.3	(38.6, 43.7)
MAINE	885	46.3	1.9	(42.6, 50.0)
MARYLAND	1001	30.7	1.9	(27.0, 34.3)
MASSACHUSETTS	913	52.3	1.9	(48.6, 56.0)
MINNESOTA	2433	36.4	1.0	(34.4, 38.4)
MISSISSIPPI	799	34.3	2.0	(30.4, 38.3)
MONTANA	891	37.0	1.6	(33.8, 40.2)
NEBRASKA	1267	43.7	1.9	(40.0, 47.4)
NEVADA	1399	44.1	2.7	(38.9, 49.3)
NEW HAMPSHIRE	813	48.6	1.9	(44.8, 52.4)
NEW JERSEY	1330	48.8	1.7	(45.5, 52.0)
NEW MEXICO	912	39.1	1.8	(35.6, 42.6)
NEW YORK	1803	48.7	1.3	(46.1, 51.3)
NORTH DAKOTA	882	36.0	1.8	(32.5, 39.4)
OHIO	1349	45.1	1.8	(41.5, 48.6)
RHODE ISLAND	875	48.0	1.9	(44.2, 51.7)
TENNESSEE	388	37.6	2.6	(32.4, 42.7)
VERMONT	1688	51.5	1.4	(48.7, 54.3)
WISCONSIN	1221	41.3	1.7	(38.0, 44.5)
PUERTO RICO	946	41.3	1.7	(37.9, 44.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 25 MEDIAN = 44.1 RANGE = 30.7-61.6

* BELIEVES CONDOMS VERY EFFECTIVE IN PREVENTING HIV

August 17, 1998

TABLE 39.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV: CONDOM EFFECTIVENESS* -BY SEX
 DENOMINATOR IS PERSONS AGE 18-49 YEARS
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALASKA	412	48.4	3.6	(41.4, 55.4)	*	491	40.1	3.3	(33.7, 46.5)	*
DELAWARE	564	46.7	2.5	(41.7, 51.6)	*	716	48.5	2.2	(44.3, 52.7)	*
DISTRICT OF CO	344	60.8	3.3	(54.4, 67.2)	*	479	62.4	2.6	(57.4, 67.4)	*
FLORIDA	696	45.9	2.1	(41.8, 50.0)	*	945	44.2	1.8	(40.6, 47.8)	*
ILLINOIS	356	46.7	2.9	(41.1, 52.3)	*	456	34.8	2.5	(30.0, 39.7)	*
IOWA	794	44.3	1.9	(40.5, 48.1)	*	954	37.9	1.7	(34.5, 41.3)	*
MAINE	407	45.6	2.8	(40.0, 51.1)	*	478	47.1	2.5	(42.1, 52.0)	*
MARYLAND	444	29.8	2.7	(24.5, 35.1)	*	557	31.5	2.5	(26.6, 36.4)	*
MASSACHUSETTS	413	54.8	2.8	(49.3, 60.3)	*	500	49.6	2.5	(44.7, 54.5)	*
MINNESOTA	1192	38.9	1.5	(36.0, 41.8)	*	1241	33.8	1.4	(31.0, 36.7)	*
MISSISSIPPI	325	38.4	3.2	(32.1, 44.7)	*	474	30.3	2.3	(25.8, 34.8)	*
MONTANA	404	38.8	2.5	(33.9, 43.7)	*	487	35.1	2.2	(30.8, 39.4)	*
NEBRASKA	533	47.8	2.8	(42.4, 53.2)	*	734	39.7	2.5	(34.8, 44.5)	*
NEVADA	683	46.7	4.1	(38.6, 54.8)	*	716	41.1	3.3	(34.7, 47.5)	*
NEW HAMPSHIRE	338	50.4	3.1	(44.4, 56.5)	*	475	46.7	2.4	(42.0, 51.4)	*
NEW JERSEY	561	52.1	2.5	(47.3, 56.9)	*	769	45.3	2.2	(41.0, 49.6)	*
NEW MEXICO	403	38.3	2.6	(33.1, 43.4)	*	509	40.0	2.5	(35.2, 44.8)	*
NEW YORK	773	50.2	2.0	(46.3, 54.1)	*	1030	47.1	1.8	(43.7, 50.5)	*
NORTH DAKOTA	420	36.0	2.5	(31.2, 40.9)	*	462	35.9	2.4	(31.2, 40.7)	*
OHIO	550	45.1	2.7	(39.9, 50.4)	*	799	45.0	2.2	(40.6, 49.4)	*
RHODE ISLAND	382	50.8	2.8	(45.3, 56.4)	*	493	45.0	2.6	(40.0, 50.1)	*
TENNESSEE	169	39.6	4.1	(31.6, 47.5)	*	219	35.3	3.3	(28.8, 41.7)	*
VERMONT	791	54.6	2.1	(50.5, 58.7)	*	897	48.1	1.9	(44.3, 51.9)	*
WISCONSIN	585	42.8	2.5	(38.0, 47.7)	*	636	39.7	2.2	(35.3, 44.0)	*
PUERTO RICO	446	47.1	2.5	(42.2, 52.1)	*	500	34.8	2.3	(30.3, 39.2)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 25
 MEDIAN = 46.7

NUMBER OF PARTICIPANTS = 25
 MEDIAN = 40.1

RANGE

= 29.8-60.8

* BELIEVES CONDOMS VERY EFFECTIVE IN PREVENTING HIV

RANGE

= 30.3-62.4

August 17, 1998

TABLE 40.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM*
 DENOMINATOR IS FEMALES AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1349	65.0	1.5	(62.1, 67.9)
ALASKA	821	50.2	2.6	(45.1, 55.2)
ARIZONA	1002	54.6	2.1	(50.5, 58.7)
ARKANSAS	1106	56.0	1.7	(52.6, 59.4)
COLORADO	1001	57.3	1.8	(53.7, 60.8)
CONNECTICUT	1335	61.7	1.6	(58.6, 64.8)
DELAWARE	1537	63.1	1.5	(60.1, 66.1)
DISTRICT OF CO	931	59.9	1.9	(56.3, 63.6)
FLORIDA	2054	66.7	1.2	(64.3, 69.1)
GEORGIA	1342	59.0	1.6	(55.9, 62.1)
HAWAII	1218	62.3	1.7	(59.0, 65.6)
IDAHO	2863	55.9	1.3	(53.4, 58.4)
ILLINOIS	1721	57.6	1.3	(55.0, 60.2)
INDIANA	1301	57.1	1.6	(53.9, 60.3)
IOWA	2099	58.3	1.2	(55.9, 60.7)
KANSAS	1203	54.6	1.6	(51.5, 57.6)
KENTUCKY	2266	58.0	1.2	(55.7, 60.4)
LOUISIANA	1026	57.7	1.7	(54.3, 61.1)
MAINE	942	59.3	1.9	(55.7, 63.0)

MARYLAND	2598	59.7	1.3	(57.2, 62.2)
MASSACHUSETTS	1009	60.6	1.8	(57.2, 64.1)
MICHIGAN	1487	62.3	1.4	(59.6, 64.9)
MINNESOTA	2610	56.1	1.1	(54.0, 58.2)
MISSISSIPPI	1004	54.1	1.8	(50.5, 57.7)
MISSOURI	1144	58.7	1.7	(55.4, 62.1)
MONTANA	1042	59.6	1.7	(56.2, 62.9)
NEBRASKA	1634	55.6	1.6	(52.4, 58.7)
NEVADA	1340	60.3	2.5	(55.4, 65.1)
NEW HAMPSHIRE	873	59.1	1.9	(55.4, 62.8)
NEW JERSEY	1553	62.8	1.5	(59.9, 65.6)
NEW MEXICO	1048	54.2	1.8	(50.7, 57.7)
NEW YORK	2045	60.6	1.2	(58.1, 63.0)
NORTH CAROLINA	2162	60.9	1.2	(58.6, 63.3)
NORTH DAKOTA	1000	58.4	1.8	(54.9, 62.0)
OHIO	1931	60.4	1.5	(57.6, 63.3)
OKLAHOMA	1052	52.7	1.8	(49.2, 56.2)
OREGON	1919	61.1	1.4	(58.5, 63.8)
PENNSYLVANIA	2108	61.2	1.3	(58.7, 63.6)
RHODE ISLAND	1062	62.2	1.8	(58.8, 65.7)
SOUTH CAROLINA	1333	60.9	1.6	(57.8, 64.0)
SOUTH DAKOTA	1237	59.1	1.6	(56.0, 62.1)
TENNESSEE	1915	58.4	1.4	(55.8, 61.1)
TEXAS	1471	55.7	1.4	(52.9, 58.5)
UTAH	1581	51.3	1.7	(47.9, 54.7)
VERMONT	1765	54.5	1.4	(51.7, 57.2)
VIRGINIA	2098	57.4	1.7	(54.0, 60.8)
WASHINGTON	2051	60.1	1.3	(57.6, 62.6)
WEST VIRGINIA	1459	58.9	1.5	(55.9, 61.8)
WISCONSIN	1250	56.7	1.7	(53.4, 59.9)
WYOMING	1392	59.7	1.6	(56.5, 62.9)
PUERTO RICO	1324	55.1	1.6	(52.0, 58.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 58.9 RANGE = 50.2-66.7

* EVER HAD A MAMMOGRAM, AGE 18 AND OVER

August 17, 1998

TABLE 40.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 MAMMOGRAM *

DENOMINATOR IS FEMALES AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	60.2	65.0	4.8
ALASKA	56.9	50.2	-6.7
ARIZONA	60.9	54.6	-6.3
ARKANSAS	54.0	56.0	2.0
COLORADO	58.5	57.3	-1.2
CONNECTICUT	65.6	61.7	-3.9
DELAWARE	59.3	63.1	3.8
DISTRICT OF CO	58.6	59.9	1.3

FLORIDA	64.6	66.7	2.0
GEORGIA	60.1	59.0	-1.2
HAWAII	62.1	62.3	0.2
IDAHO	53.8	55.9	2.1
ILLINOIS	57.0	57.6	0.6
INDIANA	55.7	57.1	1.4
IOWA	57.0	58.3	1.3
KANSAS	56.1	54.6	-1.6
KENTUCKY	57.2	58.0	0.8
LOUISIANA	55.8	57.7	2.0
MAINE	59.8	59.3	-0.5
MARYLAND	61.9	59.7	-2.2
MASSACHUSETTS	59.6	60.6	1.1
MICHIGAN	61.7	62.3	0.6
MINNESOTA	55.3	56.1	0.8
MISSISSIPPI	54.4	54.1	-0.3
MISSOURI	58.2	58.7	0.6
MONTANA	60.5	59.6	-0.9
NEBRASKA	53.7	55.6	1.8
NEVADA	59.7	60.3	0.6
NEW HAMPSHIRE	57.1	59.1	2.0
NEW JERSEY	58.6	62.8	4.1
NEW MEXICO	53.3	54.2	0.9
NEW YORK	60.2	60.6	0.3
NORTH CAROLINA	57.6	60.9	3.4
NORTH DAKOTA	56.3	58.4	2.2
OHIO	57.4	60.4	3.1
OKLAHOMA	52.7	52.7	-0.0
OREGON	62.0	61.1	-0.8
PENNSYLVANIA	59.0	61.2	2.2
RHODE ISLAND	62.7	62.2	-0.4
SOUTH CAROLINA	57.8	60.9	3.1
SOUTH DAKOTA	55.8	59.1	3.3
TENNESSEE	56.6	58.4	1.8
TEXAS	52.3	55.7	3.4
UTAH	49.8	51.3	1.4
VERMONT	55.7	54.5	-1.3
VIRGINIA	59.0	57.4	-1.6
WASHINGTON	58.3	60.1	1.8
WEST VIRGINIA	58.6	58.9	0.3
WISCONSIN	62.2	56.7	-5.6
WYOMING	54.6	59.7	5.1
PUERTO RICO	51.7	55.1	3.4

* EVER HAD A MAMMOGRAM, AGE 18 AND OLDER
N/A - NOT AVAILABLE

August 17, 1998

TABLE 41.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM *
DENOMINATOR IS FEMALES AGE 40 AND OVER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
-------------	-------------	---------	----------------	-------------------------

ALABAMA	843	87.0	1.3	(84.5, 89.5)
ALASKA	460	84.2	2.8	(78.8, 89.6)
ARIZONA	637	78.4	2.5	(73.5, 83.2)
ARKANSAS	699	76.6	1.8	(72.9, 80.2)
COLORADO	607	85.0	1.7	(81.8, 88.3)
CONNECTICUT	775	87.5	1.4	(84.7, 90.3)
DELAWARE	986	89.3	1.1	(87.1, 91.5)
DISTRICT OF CO	506	87.8	1.5	(84.8, 90.7)
FLORIDA	1334	87.1	1.0	(85.1, 89.1)
GEORGIA	724	87.3	1.4	(84.6, 90.0)
HAWAII	751	88.4	1.5	(85.5, 91.2)
IDAHO	1749	81.1	1.3	(78.6, 83.6)
ILLINOIS	994	84.5	1.3	(82.1, 87.0)
INDIANA	786	80.8	1.7	(77.6, 84.0)
IOWA	1392	82.2	1.2	(79.9, 84.4)
KANSAS	710	79.5	1.7	(76.2, 82.7)
KENTUCKY	1449	80.3	1.2	(77.9, 82.6)
LOUISIANA	575	80.4	1.8	(76.8, 84.0)
MAINE	594	88.3	1.4	(85.6, 91.0)
MARYLAND	1530	87.8	1.1	(85.6, 90.0)
MASSACHUSETTS	541	88.6	1.5	(85.6, 91.6)
MICHIGAN	862	88.8	1.1	(86.6, 91.0)
MINNESOTA	1603	83.3	1.0	(81.3, 85.2)
MISSISSIPPI	625	75.2	1.9	(71.5, 79.0)
MISSOURI	699	82.8	1.7	(79.5, 86.0)
MONTANA	685	82.7	1.5	(79.7, 85.6)
NEBRASKA	1031	80.1	1.9	(76.3, 83.9)
NEVADA	797	85.2	2.2	(80.8, 89.5)
NEW HAMPSHIRE	532	88.6	1.5	(85.8, 91.5)
NEW JERSEY	911	86.0	1.2	(83.5, 88.4)
NEW MEXICO	635	81.4	1.7	(78.1, 84.7)
NEW YORK	1170	86.7	1.1	(84.5, 88.9)
NORTH CAROLINA	1294	86.5	1.0	(84.5, 88.6)
NORTH DAKOTA	648	84.5	1.5	(81.6, 87.4)
OHIO	1243	83.8	1.4	(81.1, 86.5)
OKLAHOMA	671	74.4	1.9	(70.8, 78.1)
OREGON	1228	87.4	1.1	(85.3, 89.4)
PENNSYLVANIA	1319	83.9	1.2	(81.5, 86.2)
RHODE ISLAND	629	89.0	1.3	(86.4, 91.6)
SOUTH CAROLINA	814	89.2	1.2	(86.8, 91.6)
SOUTH DAKOTA	785	84.8	1.5	(81.9, 87.7)
TENNESSEE	1183	81.0	1.3	(78.4, 83.6)
TEXAS	800	82.0	1.5	(79.0, 85.0)
UTAH	858	81.2	1.8	(77.7, 84.7)
VERMONT	1067	83.9	1.3	(81.4, 86.3)
VIRGINIA	1286	85.2	1.4	(82.5, 87.9)
WASHINGTON	1238	87.6	1.0	(85.6, 89.6)
WEST VIRGINIA	973	81.1	1.4	(78.3, 83.8)
WISCONSIN	734	82.6	1.6	(79.4, 85.8)
WYOMING	884	84.4	1.3	(81.8, 87.0)
PUERTO RICO	793	76.9	1.6	(73.7, 80.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 84.4 RANGE = 74.4-89.3

* EVER HAD A MAMMOGRAM, FEMALES AGE 40 AND OVER

August 17, 1998

TABLE 41.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

PREVALENCE DIFFERENCES FROM 1996 TO 1997

MAMMOGRAM *

DENOMINATOR IS FEMALES AGE 40 AND OVER

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	82.9	87.0	4.1
ALASKA	91.7	84.2	-7.5
ARIZONA	84.4	78.4	-6.0
ARKANSAS	75.7	76.6	0.9
COLORADO	87.8	85.0	-2.8
CONNECTICUT	91.2	87.5	-3.7
DELAWARE	86.6	89.3	2.7
DISTRICT OF CO	87.6	87.8	0.2
FLORIDA	87.0	87.1	0.1
GEORGIA	88.2	87.3	-0.9
HAWAII	90.0	88.4	-1.6
IDAHO	79.1	81.1	2.0
ILLINOIS	83.6	84.5	0.9
INDIANA	79.3	80.8	1.6
IOWA	80.7	82.2	1.5
KANSAS	79.0	79.5	0.5
KENTUCKY	79.3	80.3	1.0
LOUISIANA	76.1	80.4	4.3
MAINE	87.3	88.3	1.0
MARYLAND	89.1	87.8	-1.3
MASSACHUSETTS	89.5	88.6	-0.9
MICHIGAN	88.5	88.8	0.3
MINNESOTA	84.3	83.3	-1.0
MISSISSIPPI	76.3	75.2	-1.1
MISSOURI	81.0	82.8	1.8
MONTANA	85.9	82.7	-3.3
NEBRASKA	78.4	80.1	1.7
NEVADA	84.0	85.2	1.2
NEW HAMPSHIRE	85.9	88.6	2.7
NEW JERSEY	81.9	86.0	4.1
NEW MEXICO	81.3	81.4	0.1
NEW YORK	85.6	86.7	1.1
NORTH CAROLINA	82.0	86.5	4.5
NORTH DAKOTA	84.3	84.5	0.3
OHIO	81.3	83.8	2.5
OKLAHOMA	71.8	74.4	2.6
OREGON	87.2	87.4	0.1
PENNSYLVANIA	82.2	83.9	1.7
RHODE ISLAND	88.6	89.0	0.4
SOUTH CAROLINA	84.1	89.2	5.1
SOUTH DAKOTA	81.5	84.8	3.3
TENNESSEE	80.1	81.0	0.9
TEXAS	77.3	82.0	4.8
UTAH	82.8	81.2	-1.5
VERMONT	85.8	83.9	-1.9
VIRGINIA	85.6	85.2	-0.4
WASHINGTON	86.4	87.6	1.3
WEST VIRGINIA	80.6	81.1	0.5
WISCONSIN	89.2	82.6	-6.6
WYOMING	78.9	84.4	5.4
PUERTO RICO	72.4	76.9	4.5

* EVER HAD A MAMMOGRAM, FEMALES AGE 40 AND OLDER

N/A - NOT AVAILABLE

August 17, 1998

TABLE 42.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM *

DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	585	75.2	2.0	(71.3, 79.0)
ALASKA	230	78.2	4.3	(69.7, 86.6)
ARIZONA	482	73.8	2.9	(68.1, 79.5)
ARKANSAS	484	55.9	2.5	(51.0, 60.7)
COLORADO	394	76.1	2.3	(71.6, 80.6)
CONNECTICUT	488	77.7	2.2	(73.4, 82.0)
DELAWARE	688	79.7	1.8	(76.3, 83.2)
DISTRICT OF CO	322	83.6	2.1	(79.5, 87.8)
FLORIDA	959	78.9	1.4	(76.1, 81.8)
GEORGIA	456	75.4	2.3	(70.9, 79.9)
HAWAII	460	82.4	2.1	(78.4, 86.5)
IDAHO	1175	64.9	1.8	(61.3, 68.5)
ILLINOIS	642	73.1	1.9	(69.3, 76.8)
INDIANA	506	65.5	2.4	(60.9, 70.2)
IOWA	998	67.9	1.6	(64.6, 71.1)
KANSAS	492	70.0	2.3	(65.6, 74.5)
KENTUCKY	1041	69.7	1.5	(66.7, 72.6)
LOUISIANA	391	69.9	2.7	(64.6, 75.1)
MAINE	387	77.6	2.3	(73.0, 82.2)
MARYLAND	985	80.5	1.7	(77.2, 83.8)
MASSACHUSETTS	361	81.0	2.3	(76.5, 85.4)
MICHIGAN	539	79.9	1.8	(76.3, 83.5)
MINNESOTA	1050	73.7	1.4	(70.9, 76.5)
MISSISSIPPI	445	64.7	2.5	(59.9, 69.6)
MISSOURI	474	69.0	2.5	(64.1, 74.0)
MONTANA	450	70.4	2.2	(66.0, 74.8)
NEBRASKA	743	67.5	1.9	(63.7, 71.3)
NEVADA	507	67.7	3.9	(60.0, 75.3)
NEW HAMPSHIRE	323	79.7	2.4	(74.9, 84.4)
NEW JERSEY	590	73.7	2.0	(69.8, 77.6)
NEW MEXICO	413	67.7	2.6	(62.7, 72.7)
NEW YORK	747	78.1	1.7	(74.9, 81.4)
NORTH CAROLINA	871	74.4	1.6	(71.2, 77.6)
NORTH DAKOTA	460	72.6	2.2	(68.3, 76.9)
OHIO	917	74.9	2.0	(71.0, 78.8)
OKLAHOMA	486	65.2	2.4	(60.5, 69.9)
OREGON	813	79.3	1.6	(76.2, 82.3)
PENNSYLVANIA	888	73.8	1.7	(70.4, 77.1)
RHODE ISLAND	430	83.1	1.9	(79.4, 86.8)
SOUTH CAROLINA	573	75.6	1.9	(71.7, 79.4)
SOUTH DAKOTA	544	70.8	2.2	(66.4, 75.2)
TENNESSEE	822	74.0	1.7	(70.7, 77.3)
TEXAS	508	68.1	2.3	(63.6, 72.5)
UTAH	517	70.2	2.7	(65.0, 75.5)
VERMONT	676	73.6	1.9	(69.9, 77.2)
VIRGINIA	832	76.3	1.8	(72.8, 79.9)
WASHINGTON	773	73.4	1.8	(69.9, 76.9)
WEST VIRGINIA	708	69.3	1.9	(65.6, 73.1)
WISCONSIN	499	70.9	2.3	(66.4, 75.4)
WYOMING	553	69.3	2.1	(65.1, 73.4)
PUERTO RICO	553	61.9	2.3	(57.4, 66.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 73.7 RANGE = 55.9-83.6

TABLE 42.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 MAMMOGRAM *

DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	70.2	75.2	5.0
ALASKA	80.1	78.2	-1.9
ARIZONA	79.3	73.8	-5.5
ARKANSAS	59.9	55.9	-4.0
COLORADO	78.5	76.1	-2.4
CONNECTICUT	79.9	77.7	-2.2
DELAWARE	79.2	79.7	0.6
DISTRICT OF CO	82.6	83.6	1.1
FLORIDA	80.4	78.9	-1.5
GEORGIA	79.8	75.4	-4.4
HAWAII	78.6	82.4	3.8
IDAHO	66.4	64.9	-1.4
ILLINOIS	69.0	73.1	4.0
INDIANA	64.0	65.5	1.5
IOWA	62.1	67.9	5.7
KANSAS	71.5	70.0	-1.5
KENTUCKY	64.0	69.7	5.7
LOUISIANA	63.7	69.9	6.2
MAINE	76.8	77.6	0.8
MARYLAND	82.1	80.5	-1.7
MASSACHUSETTS	81.0	81.0	0.0
MICHIGAN	75.3	79.9	4.6
MINNESOTA	69.9	73.7	3.8
MISSISSIPPI	60.9	64.7	3.9
MISSOURI	65.5	69.0	3.5
MONTANA	68.8	70.4	1.5
NEBRASKA	66.1	67.5	1.4
NEVADA	68.1	67.7	-0.4
NEW HAMPSHIRE	74.9	79.7	4.8
NEW JERSEY	69.4	73.7	4.3
NEW MEXICO	67.8	67.7	-0.1
NEW YORK	76.7	78.1	1.5
NORTH CAROLINA	72.5	74.4	1.9
NORTH DAKOTA	69.9	72.6	2.8
OHIO	70.8	74.9	4.1
OKLAHOMA	60.5	65.2	4.6
OREGON	78.5	79.3	0.8
PENNSYLVANIA	70.0	73.8	3.8
RHODE ISLAND	77.3	83.1	5.9
SOUTH CAROLINA	75.9	75.6	-0.4
SOUTH DAKOTA	67.4	70.8	3.3
TENNESSEE	68.9	74.0	5.1
TEXAS	67.9	68.1	0.2

UTAH	69.7	70.2	0.5
VERMONT	74.7	73.6	-1.1
VIRGINIA	72.5	76.3	3.8
WASHINGTON	79.2	73.4	-5.8
WEST VIRGINIA	70.4	69.3	-1.0
WISCONSIN	74.0	70.9	-3.1
WYOMING	64.0	69.3	5.2
PUERTO RICO	61.4	61.9	0.4

* HAD A MAMMOGRAM WITHIN TWO YEARS AGE 50 AND OVER
N/A - NOT AVAILABLE

August 17, 1998

TABLE 43.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BREAST EXAM *
DENOMINATOR IS FEMALES AGE 18 AND OVER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	SAMPLE PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1348	88.7	1.0	(86.8, 90.7)
ALASKA	820	91.5	1.4	(88.6, 94.3)
ARIZONA	999	80.2	1.9	(76.4, 84.0)
ARKANSAS	1104	82.9	1.4	(80.1, 85.7)
CALIFORNIA	2300	85.6	0.9	(83.7, 87.4)
COLORADO	997	92.3	1.0	(90.4, 94.2)
CONNECTICUT	1331	87.7	1.1	(85.5, 89.9)
DELAWARE	1534	93.6	0.8	(92.0, 95.1)
DISTRICT OF CO	931	92.9	1.0	(91.1, 94.8)
FLORIDA	2044	88.5	0.8	(86.9, 90.1)
GEORGIA	1339	93.3	0.8	(91.7, 94.9)
HAWAII	1214	89.8	1.1	(87.6, 92.0)
IDAHO	2856	92.4	0.7	(91.1, 93.6)
ILLINOIS	1719	86.3	1.0	(84.4, 88.2)
INDIANA	1298	88.9	1.0	(86.9, 90.9)
IOWA	2097	89.9	0.8	(88.3, 91.5)
KANSAS	1193	91.3	0.9	(89.6, 93.0)
KENTUCKY	2263	88.1	0.8	(86.4, 89.7)
LOUISIANA	1024	84.9	1.3	(82.3, 87.5)
MAINE	939	93.4	0.9	(91.7, 95.1)
MARYLAND	2575	94.8	0.6	(93.5, 96.0)
MASSACHUSETTS	1001	89.9	1.2	(87.6, 92.1)
MICHIGAN	1483	89.3	1.0	(87.4, 91.1)
MINNESOTA	2602	91.8	0.7	(90.5, 93.0)
MISSISSIPPI	998	88.5	1.2	(86.2, 90.9)
MISSOURI	1141	92.6	0.9	(90.9, 94.3)
MONTANA	1041	93.1	1.0	(91.1, 95.2)
NEBRASKA	1629	90.4	0.9	(88.7, 92.1)
NEVADA	1338	92.0	1.2	(89.6, 94.4)
NEW HAMPSHIRE	873	93.3	0.9	(91.4, 95.1)
NEW JERSEY	1540	86.3	1.1	(84.3, 88.4)
NEW MEXICO	1047	86.8	1.3	(84.2, 89.4)
NEW YORK	2038	92.8	0.7	(91.5, 94.1)
NORTH CAROLINA	2162	93.3	0.6	(92.1, 94.4)

NORTH DAKOTA	999	91.4	1.0	(89.3, 93.4)
OHIO	1927	90.7	1.0	(88.9, 92.6)
OKLAHOMA	1052	92.5	1.0	(90.5, 94.5)
OREGON	1911	94.2	0.6	(93.0, 95.4)
PENNSYLVANIA	2094	84.6	1.0	(82.5, 86.6)
RHODE ISLAND	1057	86.8	1.3	(84.3, 89.4)
SOUTH CAROLINA	1330	93.1	0.8	(91.6, 94.7)
SOUTH DAKOTA	1229	91.2	1.0	(89.3, 93.2)
TENNESSEE	1904	89.2	0.8	(87.7, 90.7)
TEXAS	1471	83.4	1.1	(81.2, 85.6)
UTAH	1577	92.2	1.0	(90.2, 94.1)
VERMONT	1757	88.4	1.0	(86.4, 90.4)
VIRGINIA	2082	87.8	1.6	(84.7, 90.8)
WASHINGTON	2043	91.3	1.0	(89.4, 93.2)
WEST VIRGINIA	1458	90.1	0.9	(88.3, 91.9)
WISCONSIN	1243	93.5	0.8	(91.9, 95.1)
WYOMING	1389	91.2	1.0	(89.3, 93.2)
PUERTO RICO	1322	81.4	1.3	(78.8, 84.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 90.6 RANGE = 80.2-94.8

* EVER HAD A BREAST EXAM BY A DOCTOR, AGE 18 AND OLDER

August 17, 1998

TABLE 43.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1996 TO 1997
BREAST EXAM *
DENOMINATOR IS FEMALES AGE 18 AND OVER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	86.7	88.7	2.0
ALASKA	93.8	91.5	-2.4
ARIZONA	87.4	80.2	-7.2
ARKANSAS	84.9	82.9	-2.0
CALIFORNIA	86.0	85.6	-0.5
COLORADO	92.1	92.3	0.2
CONNECTICUT	87.6	87.7	0.2
DELAWARE	93.0	93.6	0.6
DISTRICT OF CO	90.6	92.9	2.4
FLORIDA	88.4	88.5	0.1
GEORGIA	92.5	93.3	0.8
HAWAII	89.9	89.8	-0.1
IDAHO	91.3	92.4	1.0
ILLINOIS	89.0	86.3	-2.7
INDIANA	89.5	88.9	-0.6
IOWA	91.7	89.9	-1.8
KANSAS	91.2	91.3	0.1
KENTUCKY	87.0	88.1	1.1
LOUISIANA	85.7	84.9	-0.8
MAINE	92.0	93.4	1.4
MARYLAND	94.2	94.8	0.6
MASSACHUSETTS	92.8	89.9	-2.9
MICHIGAN	89.9	89.3	-0.6
MINNESOTA	92.0	91.8	-0.3

MISSISSIPPI	87.7	88.5	0.8
MISSOURI	89.4	92.6	3.2
MONTANA	94.4	93.1	-1.3
NEBRASKA	92.3	90.4	-1.9
NEVADA	89.7	92.0	2.3
NEW HAMPSHIRE	90.7	93.3	2.6
NEW JERSEY	83.9	86.3	2.5
NEW MEXICO	89.0	86.8	-2.2
NEW YORK	90.2	92.8	2.6
NORTH CAROLINA	92.5	93.3	0.8
NORTH DAKOTA	90.5	91.4	0.9
OHIO	90.5	90.7	0.3
OKLAHOMA	92.3	92.5	0.2
OREGON	93.3	94.2	0.9
PENNSYLVANIA	84.1	84.6	0.5
RHODE ISLAND	90.0	86.8	-3.2
SOUTH CAROLINA	90.0	93.1	3.1
SOUTH DAKOTA	90.1	91.2	1.2
TENNESSEE	86.9	89.2	2.3
TEXAS	82.2	83.4	1.1
UTAH	89.4	92.2	2.8
VERMONT	88.3	88.4	0.1
VIRGINIA	90.3	87.8	-2.5
WASHINGTON	91.4	91.3	-0.1
WEST VIRGINIA	90.5	90.1	-0.4
WISCONSIN	94.6	93.5	-1.1
WYOMING	91.6	91.2	-0.4
PUERTO RICO	71.4	81.4	10.0

* EVER HAD A BREAST EXAM BY A DOCTOR, AGE 18 AND OLDER
 N/A - NOT AVAILABLE

August 17, 1998

TABLE 44.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BREAST EXAM *
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	841	87.8	1.2	(85.3, 90.2)
ALASKA	459	92.7	1.9	(89.0, 96.4)
ARIZONA	635	81.2	2.3	(76.7, 85.8)
ARKANSAS	697	83.3	1.5	(80.4, 86.2)
CALIFORNIA	1343	89.0	1.1	(86.8, 91.3)
COLORADO	606	93.1	1.1	(90.8, 95.3)
CONNECTICUT	772	87.7	1.5	(84.8, 90.6)
DELAWARE	983	93.9	0.8	(92.3, 95.6)
DISTRICT OF CO	506	93.0	1.3	(90.4, 95.6)
FLORIDA	1324	90.3	0.9	(88.5, 92.1)
GEORGIA	722	92.7	1.1	(90.5, 94.8)
HAWAII	747	91.4	1.3	(88.8, 94.0)
IDAHO	1741	92.8	0.7	(91.5, 94.2)
ILLINOIS	993	88.3	1.1	(86.2, 90.5)
INDIANA	784	87.2	1.4	(84.5, 89.9)

IOWA	1390	90.6	0.8	(88.9, 92.2)
KANSAS	702	90.5	1.2	(88.2, 92.9)
KENTUCKY	1446	89.1	0.9	(87.3, 90.9)
LOUISIANA	573	83.3	1.8	(79.8, 86.8)
MAINE	591	93.2	1.1	(91.0, 95.3)
MARYLAND	1513	95.8	0.6	(94.7, 97.0)
MASSACHUSETTS	537	90.6	1.5	(87.7, 93.5)
MICHIGAN	858	91.2	1.0	(89.2, 93.3)
MINNESOTA	1596	93.9	0.6	(92.7, 95.1)
MISSISSIPPI	622	90.2	1.3	(87.7, 92.7)
MISSOURI	695	91.8	1.1	(89.6, 94.0)
MONTANA	685	94.1	0.9	(92.3, 95.8)
NEBRASKA	1027	90.2	1.0	(88.2, 92.3)
NEVADA	796	91.7	1.7	(88.4, 95.1)
NEW HAMPSHIRE	532	94.3	1.1	(92.2, 96.4)
NEW JERSEY	905	86.5	1.3	(83.9, 89.1)
NEW MEXICO	634	88.2	1.5	(85.3, 91.0)
NEW YORK	1165	94.5	0.8	(93.0, 96.0)
NORTH CAROLINA	1296	94.3	0.7	(93.0, 95.6)
NORTH DAKOTA	647	92.9	1.1	(90.8, 95.0)
OHIO	1241	92.6	0.9	(90.8, 94.4)
OKLAHOMA	670	94.3	1.0	(92.4, 96.3)
OREGON	1221	95.0	0.7	(93.6, 96.3)
PENNSYLVANIA	1309	84.2	1.2	(81.9, 86.5)
RHODE ISLAND	624	88.2	1.4	(85.4, 91.0)
SOUTH CAROLINA	812	95.0	0.8	(93.4, 96.6)
SOUTH DAKOTA	777	91.1	1.2	(88.8, 93.3)
TENNESSEE	1173	89.3	1.0	(87.3, 91.2)
TEXAS	799	88.4	1.3	(85.9, 90.9)
UTAH	855	95.8	1.0	(93.8, 97.8)
VERMONT	1061	89.8	1.1	(87.7, 91.9)
VIRGINIA	1274	89.4	1.4	(86.6, 92.2)
WASHINGTON	1230	93.8	0.8	(92.3, 95.3)
WEST VIRGINIA	971	90.6	1.0	(88.6, 92.6)
WISCONSIN	726	93.6	1.1	(91.6, 95.7)
WYOMING	880	92.6	0.9	(90.7, 94.4)
PUERTO RICO	793	88.8	1.2	(86.4, 91.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 91.1 RANGE = 81.2-95.8

* EVER HAD A BREAST EXAM BY A DOCTOR, FEMALES AGE 40 AND OLDER

August 17, 1998

TABLE 44.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1996 TO 1997
BREAST EXAM *

DENOMINATOR IS FEMALES AGE 40 AND OVER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	84.8	87.8	2.9
ALASKA	95.8	92.7	-3.1
ARIZONA	92.5	81.2	-11.2
ARKANSAS	84.2	83.3	-0.9
CALIFORNIA	89.9	89.0	-0.8

COLORADO	91.7	93.1	1.3
CONNECTICUT	88.3	87.7	-0.6
DELAWARE	94.3	93.9	-0.4
DISTRICT OF CO	93.5	93.0	-0.5
FLORIDA	90.4	90.3	-0.1
GEORGIA	92.7	92.7	-0.0
HAWAII	92.1	91.4	-0.7
IDAHO	92.1	92.8	0.7
ILLINOIS	89.1	88.3	-0.8
INDIANA	89.7	87.2	-2.5
IOWA	90.7	90.6	-0.2
KANSAS	90.9	90.5	-0.4
KENTUCKY	87.6	89.1	1.5
LOUISIANA	84.5	83.3	-1.2
MAINE	92.5	93.2	0.7
MARYLAND	94.1	95.8	1.7
MASSACHUSETTS	93.9	90.6	-3.3
MICHIGAN	89.9	91.2	1.3
MINNESOTA	92.9	93.9	1.0
MISSISSIPPI	87.7	90.2	2.5
MISSOURI	89.1	91.8	2.7
MONTANA	96.0	94.1	-2.0
NEBRASKA	91.6	90.2	-1.4
NEVADA	90.7	91.7	1.0
NEW HAMPSHIRE	89.7	94.3	4.7
NEW JERSEY	83.0	86.5	3.5
NEW MEXICO	88.7	88.2	-0.6
NEW YORK	92.4	94.5	2.1
NORTH CAROLINA	93.5	94.3	0.9
NORTH DAKOTA	90.1	92.9	2.8
OHIO	93.1	92.6	-0.5
OKLAHOMA	92.7	94.3	1.7
OREGON	93.6	95.0	1.4
PENNSYLVANIA	83.0	84.2	1.2
RHODE ISLAND	89.5	88.2	-1.3
SOUTH CAROLINA	91.7	95.0	3.3
SOUTH DAKOTA	89.9	91.1	1.2
TENNESSEE	86.3	89.3	3.0
TEXAS	83.7	88.4	4.7
UTAH	91.3	95.8	4.5
VERMONT	89.0	89.8	0.9
VIRGINIA	91.1	89.4	-1.7
WASHINGTON	90.9	93.8	2.9
WEST VIRGINIA	90.4	90.6	0.3
WISCONSIN	95.6	93.6	-2.0
WYOMING	90.6	92.6	2.0
PUERTO RICO	77.6	88.8	11.2

* EVER HAD A BREAST EXAM BY A DOCTOR, FEMALES AGE 40 AND OLDER

N/A - NOT AVAILABLE

August 17, 1998

TABLE 45.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BREAST EXAM *

DENOMINATOR IS FEMALES AGE 50 AND OLDER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	580	74.0	2.0	(70.2, 77.8)
ALASKA	228	79.7	4.4	(71.1, 88.2)
ARIZONA	481	75.6	3.1	(69.4, 81.7)
ARKANSAS	481	63.4	2.3	(58.8, 67.9)
CALIFORNIA	878	77.2	1.7	(73.8, 80.6)
COLORADO	388	78.5	2.2	(74.1, 82.9)
CONNECTICUT	486	76.4	2.3	(71.9, 80.8)
DELAWARE	678	81.3	1.6	(78.1, 84.5)
DISTRICT OF CO	324	83.3	2.3	(78.8, 87.8)
FLORIDA	942	78.2	1.5	(75.4, 81.1)
GEORGIA	448	82.8	2.0	(78.9, 86.7)
HAWAII	454	80.9	2.4	(76.3, 85.5)
IDAHO	1152	71.7	1.6	(68.7, 74.8)
ILLINOIS	638	72.4	1.9	(68.7, 76.1)
INDIANA	500	69.4	2.4	(64.8, 74.1)
IOWA	989	73.7	1.5	(70.7, 76.8)
KANSAS	479	76.4	2.2	(72.1, 80.6)
KENTUCKY	1030	73.3	1.5	(70.3, 76.2)
LOUISIANA	390	64.8	2.7	(59.5, 70.1)
MAINE	384	83.0	2.2	(78.8, 87.2)
MARYLAND	968	84.5	1.5	(81.6, 87.3)
MASSACHUSETTS	355	83.8	2.1	(79.7, 87.8)
MICHIGAN	529	77.6	1.9	(73.8, 81.4)
MINNESOTA	1026	78.7	1.3	(76.1, 81.3)
MISSISSIPPI	440	73.3	2.4	(68.6, 78.0)
MISSOURI	462	74.5	2.5	(69.7, 79.4)
MONTANA	449	78.4	2.0	(74.5, 82.2)
NEBRASKA	734	71.1	1.9	(67.4, 74.8)
NEVADA	504	69.8	3.8	(62.3, 77.2)
NEW HAMPSHIRE	320	85.3	2.1	(81.1, 89.4)
NEW JERSEY	585	73.2	2.1	(69.2, 77.3)
NEW MEXICO	408	72.5	2.5	(67.6, 77.4)
NEW YORK	737	86.4	1.4	(83.7, 89.1)
NORTH CAROLINA	869	82.4	1.4	(79.6, 85.2)
NORTH DAKOTA	457	75.6	2.2	(71.3, 79.8)
OHIO	915	81.7	1.7	(78.4, 85.0)
OKLAHOMA	487	81.7	1.9	(77.9, 85.5)
OREGON	801	80.1	1.6	(77.0, 83.2)
PENNSYLVANIA	880	70.3	1.8	(66.7, 73.9)
RHODE ISLAND	426	79.7	2.1	(75.5, 83.8)
SOUTH CAROLINA	571	83.0	1.7	(79.7, 86.3)
SOUTH DAKOTA	534	75.9	2.0	(72.0, 79.9)
TENNESSEE	808	79.8	1.5	(76.8, 82.7)
TEXAS	500	71.9	2.2	(67.6, 76.2)
UTAH	517	77.3	2.4	(72.6, 81.9)
VERMONT	665	75.5	1.9	(71.8, 79.2)
VIRGINIA	814	78.5	1.8	(75.1, 82.0)
WASHINGTON	757	77.7	1.7	(74.5, 80.9)
WEST VIRGINIA	705	72.7	1.9	(69.1, 76.3)
WISCONSIN	481	75.3	2.5	(70.4, 80.1)
WYOMING	536	70.8	2.2	(66.6, 75.1)
PUERTO RICO	553	76.8	2.0	(73.0, 80.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 77.0 RANGE = 63.4-86.4

* HAD A BREAST EXAM WITHIN TWO YEARS, AGE 50 AND OVER

August 17, 1998

TABLE 45.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

PREVALENCE DIFFERENCES FROM 1996 TO 1997

BREAST EXAM *

DENOMINATOR IS FEMALES AGE 50 AND OVER

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	68.4	74.0	5.6
ALASKA	82.6	79.7	-2.9
ARIZONA	82.7	75.6	-7.1
ARKANSAS	65.4	63.4	-2.0
CALIFORNIA	77.8	77.2	-0.5
COLORADO	77.2	78.5	1.3
CONNECTICUT	74.7	76.4	1.7
DELAWARE	80.2	81.3	1.1
DISTRICT OF CO	83.4	83.3	-0.1
FLORIDA	79.8	78.2	-1.5
GEORGIA	87.1	82.8	-4.3
HAWAII	77.3	80.9	3.5
IDAHO	69.3	71.7	2.4
ILLINOIS	71.0	72.4	1.4
INDIANA	65.2	69.4	4.3
IOWA	64.3	73.7	9.5
KANSAS	75.7	76.4	0.6
KENTUCKY	70.2	73.3	3.1
LOUISIANA	67.0	64.8	-2.2
MAINE	78.7	83.0	4.3
MARYLAND	83.7	84.5	0.7
MASSACHUSETTS	82.7	83.8	1.0
MICHIGAN	73.9	77.6	3.7
MINNESOTA	75.4	78.7	3.3
MISSISSIPPI	66.2	73.3	7.1
MISSOURI	67.3	74.5	7.2
MONTANA	78.6	78.4	-0.2
NEBRASKA	74.4	71.1	-3.3
NEVADA	71.9	69.8	-2.2
NEW HAMPSHIRE	78.0	85.3	7.3
NEW JERSEY	69.3	73.2	4.0
NEW MEXICO	70.6	72.5	1.9
NEW YORK	81.4	86.4	5.1
NORTH CAROLINA	80.2	82.4	2.1
NORTH DAKOTA	74.7	75.6	0.8
OHIO	79.9	81.7	1.8
OKLAHOMA	77.6	81.7	4.1
OREGON	79.5	80.1	0.6
PENNSYLVANIA	68.8	70.3	1.5
RHODE ISLAND	78.5	79.7	1.2
SOUTH CAROLINA	82.1	83.0	0.9
SOUTH DAKOTA	70.3	75.9	5.6
TENNESSEE	71.8	79.8	8.0
TEXAS	69.5	71.9	2.4
UTAH	74.8	77.3	2.5
VERMONT	74.6	75.5	0.9
VIRGINIA	77.7	78.5	0.9
WASHINGTON	77.4	77.7	0.3
WEST VIRGINIA	74.2	72.7	-1.5
WISCONSIN	77.1	75.3	-1.9
WYOMING	69.6	70.8	1.3
PUERTO RICO	64.4	76.8	12.4

* HAD A BREAST EXAM WITHIN TWO YEARS, AGE 50 AND OVER

N/A - NOT AVAILABLE

August 17, 1998

TABLE 46.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM AND BREAST EXAM *
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	840	77.8	1.6	(74.8, 80.9)
ALASKA	458	80.1	3.0	(74.2, 86.0)
ARIZONA	637	73.1	2.7	(67.9, 78.3)
ARKANSAS	697	67.7	2.0	(63.7, 71.7)
COLORADO	606	82.1	1.8	(78.6, 85.5)
CONNECTICUT	772	80.4	1.7	(77.1, 83.7)
DELAWARE	984	85.3	1.3	(82.7, 87.9)
DISTRICT OF CO	506	83.0	1.8	(79.5, 86.4)
FLORIDA	1326	81.4	1.2	(79.1, 83.7)
GEORGIA	722	83.8	1.5	(80.8, 86.7)
HAWAII	747	82.5	1.8	(79.0, 86.0)
IDAHO	1742	77.0	1.3	(74.4, 79.6)
ILLINOIS	993	77.1	1.4	(74.2, 79.9)
INDIANA	785	75.4	1.8	(72.0, 78.9)
IOWA	1388	77.6	1.3	(75.1, 80.0)
KANSAS	704	74.7	1.8	(71.2, 78.2)
KENTUCKY	1447	77.6	1.3	(75.2, 80.1)
LOUISIANA	574	70.8	2.1	(66.7, 74.9)
MAINE	592	84.4	1.6	(81.3, 87.5)
MARYLAND	1518	85.6	1.2	(83.2, 87.9)
MASSACHUSETTS	538	82.9	1.9	(79.2, 86.5)
MICHIGAN	858	83.2	1.4	(80.6, 85.9)
MINNESOTA	1599	80.6	1.1	(78.6, 82.7)
MISSISSIPPI	623	71.7	2.0	(67.7, 75.6)
MISSOURI	696	79.0	1.8	(75.5, 82.5)
MONTANA	685	79.4	1.7	(76.0, 82.7)
NEBRASKA	1025	75.0	2.0	(71.0, 79.0)
NEVADA	797	81.0	2.4	(76.3, 85.7)
NEW HAMPSHIRE	532	85.5	1.7	(82.2, 88.8)
NEW JERSEY	907	76.7	1.6	(73.5, 79.8)
NEW MEXICO	634	75.9	1.9	(72.1, 79.6)
NEW YORK	1165	84.3	1.2	(81.9, 86.6)
NORTH CAROLINA	1294	84.1	1.1	(82.0, 86.3)
NORTH DAKOTA	647	82.0	1.6	(78.9, 85.0)
OHIO	1243	81.6	1.4	(78.8, 84.4)
OKLAHOMA	670	72.6	1.9	(68.8, 76.3)
OREGON	1221	85.0	1.2	(82.8, 87.3)
PENNSYLVANIA	1308	74.0	1.4	(71.2, 76.8)
RHODE ISLAND	624	80.8	1.7	(77.3, 84.2)
SOUTH CAROLINA	814	86.3	1.3	(83.7, 88.8)
SOUTH DAKOTA	779	79.6	1.7	(76.4, 82.8)
TENNESSEE	1176	74.7	1.5	(71.8, 77.6)
TEXAS	798	76.9	1.7	(73.6, 80.2)
UTAH	855	79.5	1.8	(75.9, 83.1)
VERMONT	1063	78.4	1.4	(75.6, 81.2)
VIRGINIA	1273	76.8	1.8	(73.3, 80.4)
WASHINGTON	1228	83.3	1.2	(81.0, 85.6)
WEST VIRGINIA	972	77.9	1.5	(74.9, 80.8)
WISCONSIN	728	80.0	1.7	(76.6, 83.3)
WYOMING	879	80.0	1.5	(77.2, 82.9)
PUERTO RICO	793	73.2	1.7	(69.8, 76.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 79.6 RANGE = 67.7-86.3 YEAR 2000 OBJ 16.11
 * HAD A MAMMOGRAM AND A BREAST EXAM, AGE 40 AND OLDER August 17, 1998

TABLE 46.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 MAMMOGRAM AND BREAST EXAM *
 DENOMINATOR IS FEMALES AGE 40 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	74.5	77.8	3.3
ALASKA	88.0	80.1	-7.9
ARIZONA	80.5	73.1	-7.4
ARKANSAS	69.6	67.7	-1.9
COLORADO	82.0	82.1	0.1
CONNECTICUT	83.1	80.4	-2.7
DELAWARE	84.0	85.3	1.3
DISTRICT OF CO	83.8	83.0	-0.9
FLORIDA	81.0	81.4	0.3
GEORGIA	84.4	83.8	-0.7
HAWAII	84.1	82.5	-1.6
IDAHO	74.7	77.0	2.3
ILLINOIS	78.0	77.1	-0.9
INDIANA	74.5	75.4	1.0
IOWA	75.4	77.6	2.2
KANSAS	73.6	74.7	1.1
KENTUCKY	76.0	77.6	1.6
LOUISIANA	69.1	70.8	1.7
MAINE	83.0	84.4	1.4
MARYLAND	85.2	85.6	0.4
MASSACHUSETTS	85.8	82.9	-3.0
MICHIGAN	81.5	83.2	1.7
MINNESOTA	82.0	80.6	-1.3
MISSISSIPPI	71.4	71.7	0.2
MISSOURI	75.0	79.0	4.0
MONTANA	84.1	79.4	-4.8
NEBRASKA	74.3	75.0	0.7
NEVADA	78.9	81.0	2.1
NEW HAMPSHIRE	81.5	85.5	4.0
NEW JERSEY	72.8	76.7	3.8
NEW MEXICO	76.1	75.9	-0.3
NEW YORK	81.9	84.3	2.4
NORTH CAROLINA	79.8	84.1	4.3
NORTH DAKOTA	79.8	82.0	2.2
OHIO	78.6	81.6	3.0
OKLAHOMA	69.9	72.6	2.6
OREGON	83.5	85.0	1.6
PENNSYLVANIA	73.3	74.0	0.7
RHODE ISLAND	81.8	80.8	-1.1
SOUTH CAROLINA	79.9	86.3	6.3
SOUTH DAKOTA	76.2	79.6	3.4
TENNESSEE	72.6	74.7	2.0
TEXAS	68.6	76.9	8.3
UTAH	78.1	79.5	1.4
VERMONT	78.8	78.4	-0.4
VIRGINIA	80.2	76.8	-3.3

WASHINGTON	81.0	83.3	2.3
WEST VIRGINIA	76.8	77.9	1.0
WISCONSIN	87.1	80.0	-7.1
WYOMING	74.6	80.0	5.5
PUERTO RICO	64.4	73.2	8.8

* HAD A MAMMOGRAM AND A BREAST EXAM, AGE 40 AND OLDER
N/A - NOT AVAILABLE

August 17, 1998

TABLE 47.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR MAMMOGRAM AND BREAST EXAM *
DENOMINATOR IS FEMALES AGE 50 AND OVER
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	577	66.4	2.1	(62.2, 70.5)
ALASKA	228	70.9	4.8	(61.6, 80.3)
ARIZONA	481	69.5	3.2	(63.2, 75.7)
ARKANSAS	480	48.3	2.5	(43.5, 53.2)
COLORADO	390	68.8	2.5	(63.9, 73.7)
CONNECTICUT	484	69.0	2.4	(64.3, 73.7)
DELAWARE	681	73.3	2.0	(69.5, 77.1)
DISTRICT OF CO	321	74.6	2.6	(69.6, 79.7)
FLORIDA	943	70.2	1.6	(67.0, 73.3)
GEORGIA	447	70.7	2.5	(65.8, 75.5)
HAWAII	452	75.2	2.5	(70.2, 80.1)
IDAHO	1156	58.3	1.9	(54.6, 62.0)
ILLINOIS	636	64.2	2.0	(60.3, 68.1)
INDIANA	501	57.2	2.5	(52.4, 62.1)
IOWA	985	62.2	1.7	(58.9, 65.5)
KANSAS	483	63.1	2.4	(58.3, 67.8)
KENTUCKY	1032	66.4	1.6	(63.3, 69.5)
LOUISIANA	388	58.1	2.7	(52.8, 63.4)
MAINE	384	73.2	2.5	(68.3, 78.0)
MARYLAND	970	76.0	1.8	(72.4, 79.5)
MASSACHUSETTS	356	74.9	2.5	(70.0, 79.8)
MICHIGAN	529	71.6	2.1	(67.5, 75.7)
MINNESOTA	1032	69.0	1.5	(66.0, 72.0)
MISSISSIPPI	442	60.4	2.6	(55.3, 65.5)
MISSOURI	464	63.9	2.6	(58.7, 69.0)
MONTANA	446	67.0	2.3	(62.4, 71.5)
NEBRASKA	731	60.0	2.1	(56.0, 64.1)
NEVADA	504	59.0	4.1	(51.0, 67.0)
NEW HAMPSHIRE	321	75.6	2.6	(70.5, 80.7)
NEW JERSEY	587	63.9	2.3	(59.4, 68.3)
NEW MEXICO	409	61.7	2.7	(56.3, 67.1)
NEW YORK	740	74.6	1.8	(71.1, 78.1)
NORTH CAROLINA	865	70.7	1.7	(67.3, 74.1)
NORTH DAKOTA	457	66.3	2.3	(61.7, 70.8)
OHIO	917	71.5	2.1	(67.4, 75.5)
OKLAHOMA	487	61.7	2.5	(56.8, 66.5)
OREGON	800	73.2	1.7	(69.8, 76.6)
PENNSYLVANIA	878	63.5	1.9	(59.8, 67.2)
RHODE ISLAND	426	73.0	2.3	(68.5, 77.6)

SOUTH CAROLINA	571	71.5	2.0	(67.5, 75.5)
SOUTH DAKOTA	533	63.3	2.4	(58.7, 68.0)
TENNESSEE	811	67.5	1.8	(64.0, 71.1)
TEXAS	500	62.4	2.4	(57.7, 67.0)
UTAH	515	63.7	2.8	(58.2, 69.3)
VERMONT	666	65.8	2.1	(61.7, 69.8)
VIRGINIA	811	68.8	2.1	(64.7, 72.8)
WASHINGTON	756	67.1	1.9	(63.4, 70.7)
WEST VIRGINIA	705	64.0	2.0	(60.0, 68.0)
WISCONSIN	483	64.8	2.5	(59.8, 69.8)
WYOMING	535	61.8	2.3	(57.3, 66.2)
PUERTO RICO	553	58.8	2.3	(54.3, 63.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 66.4 RANGE = 48.3-76.0 YEAR 2000 OBJ 16.11
 * HAD A MAMMOGRAM AND A BREAST EXAM, AGE 50 AND OLDER August 17, 1998

TABLE 47.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 MAMMOGRAM AND BREAST EXAM *

DENOMINATOR IS FEMALES AGE 50 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	59.5	66.4	6.9
ALASKA	71.8	70.9	-0.8
ARIZONA	73.8	69.5	-4.4
ARKANSAS	53.5	48.3	-5.2
COLORADO	71.5	68.8	-2.7
CONNECTICUT	69.1	69.0	-0.1
DELAWARE	73.9	73.3	-0.6
DISTRICT OF CO	76.6	74.6	-2.0
FLORIDA	72.7	70.2	-2.5
GEORGIA	75.2	70.7	-4.5
HAWAII	71.0	75.2	4.2
IDAHO	57.9	58.3	0.4
ILLINOIS	61.3	64.2	3.0
INDIANA	54.7	57.2	2.5
IOWA	54.0	62.2	8.2
KANSAS	64.7	63.1	-1.7
KENTUCKY	61.0	66.4	5.5
LOUISIANA	54.0	58.1	4.1
MAINE	70.4	73.2	2.7
MARYLAND	75.8	76.0	0.2
MASSACHUSETTS	75.9	74.9	-1.0
MICHIGAN	66.7	71.6	4.9
MINNESOTA	66.1	69.0	2.9
MISSISSIPPI	53.9	60.4	6.5
MISSOURI	57.1	63.9	6.8
MONTANA	65.0	67.0	2.0
NEBRASKA	60.4	60.0	-0.3
NEVADA	60.7	59.0	-1.7
NEW HAMPSHIRE	70.1	75.6	5.5
NEW JERSEY	60.3	63.9	3.6

NEW MEXICO	58.6	61.7	3.1
NEW YORK	72.9	74.6	1.7
NORTH CAROLINA	69.5	70.7	1.2
NORTH DAKOTA	66.4	66.3	-0.2
OHIO	68.3	71.5	3.2
OKLAHOMA	57.3	61.7	4.4
OREGON	71.3	73.2	1.9
PENNSYLVANIA	60.2	63.5	3.3
RHODE ISLAND	69.7	73.0	3.3
SOUTH CAROLINA	70.4	71.5	1.0
SOUTH DAKOTA	61.0	63.3	2.4
TENNESSEE	60.2	67.5	7.4
TEXAS	57.7	62.4	4.7
UTAH	62.3	63.7	1.5
VERMONT	66.2	65.8	-0.5
VIRGINIA	65.3	68.8	3.5
WASHINGTON	70.0	67.1	-3.0
WEST VIRGINIA	64.5	64.0	-0.5
WISCONSIN	68.2	64.8	-3.4
WYOMING	56.8	61.8	5.0
PUERTO RICO	53.4	58.8	5.4

* HAD A MAMMOGRAM AND A BREAST EXAM, AGE 50 AND OLDER
 N/A - NOT AVAILABLE

August 17, 1998

TABLE 48.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PAP SMEAR*
 DENOMINATOR IS FEMALES AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1349	96.5	0.6	(95.4, 97.6)
ALASKA	819	96.2	1.0	(94.3, 98.1)
ARIZONA	999	84.7	1.7	(81.4, 88.0)
ARKANSAS	1101	92.0	1.2	(89.7, 94.4)
COLORADO	997	96.5	0.7	(95.1, 97.9)
CONNECTICUT	1323	91.8	1.1	(89.7, 93.8)
DELAWARE	1534	96.3	0.6	(95.1, 97.5)
DISTRICT OF CO	927	96.0	0.7	(94.6, 97.4)
FLORIDA	2047	93.6	0.6	(92.4, 94.9)
GEORGIA	1338	96.8	0.6	(95.7, 97.9)
HAWAII	1213	95.9	0.7	(94.5, 97.3)
IDAHO	2859	96.2	0.4	(95.3, 97.0)
ILLINOIS	1717	93.5	0.7	(92.1, 94.9)
INDIANA	1298	95.9	0.7	(94.6, 97.2)
IOWA	2092	94.3	0.7	(93.0, 95.6)
KANSAS	1193	94.1	0.8	(92.7, 95.6)
KENTUCKY	2262	92.8	0.7	(91.5, 94.1)
LOUISIANA	1023	93.4	0.9	(91.6, 95.3)
MAINE	938	95.7	0.8	(94.2, 97.2)
MARYLAND	2580	94.6	0.6	(93.3, 95.9)
MASSACHUSETTS	997	93.5	0.9	(91.6, 95.3)
MICHIGAN	1482	94.7	0.7	(93.3, 96.0)
MINNESOTA	2600	92.9	0.6	(91.6, 94.1)

MISSISSIPPI	1000	95.9	0.7	(94.5, 97.2)
MISSOURI	1142	96.5	0.8	(95.0, 98.0)
MONTANA	1040	96.2	0.9	(94.3, 98.0)
NEBRASKA	1627	95.2	0.6	(93.9, 96.4)
NEVADA	1337	95.5	1.1	(93.4, 97.5)
NEW HAMPSHIRE	870	94.5	0.9	(92.6, 96.3)
NEW JERSEY	1535	91.4	1.0	(89.5, 93.2)
NEW MEXICO	1043	93.6	1.1	(91.4, 95.7)
NEW YORK	2037	93.1	0.7	(91.7, 94.4)
NORTH CAROLINA	2162	95.3	0.6	(94.2, 96.4)
NORTH DAKOTA	1000	94.3	0.9	(92.6, 96.0)
OHIO	1930	94.8	0.7	(93.5, 96.2)
OKLAHOMA	1053	95.2	0.9	(93.6, 96.9)
OREGON	1913	96.3	0.6	(95.2, 97.3)
PENNSYLVANIA	2093	91.8	0.9	(90.1, 93.4)
RHODE ISLAND	1055	93.7	0.9	(92.0, 95.3)
SOUTH CAROLINA	1331	96.5	0.6	(95.3, 97.7)
SOUTH DAKOTA	1227	95.6	0.7	(94.3, 97.0)
TENNESSEE	1907	94.7	0.6	(93.5, 95.9)
TEXAS	1474	93.1	0.8	(91.5, 94.6)
UTAH	1576	92.5	1.1	(90.3, 94.6)
VERMONT	1751	93.1	0.9	(91.3, 94.9)
VIRGINIA	2091	95.1	1.4	(92.4, 97.9)
WASHINGTON	2048	96.4	0.8	(94.9, 97.9)
WEST VIRGINIA	1459	93.7	0.8	(92.2, 95.2)
WISCONSIN	1244	95.6	0.7	(94.3, 97.0)
WYOMING	1393	95.7	1.1	(93.6, 97.8)
PUERTO RICO	1320	84.5	1.3	(81.8, 87.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 94.7 RANGE = 84.5-96.8 YEAR 2000 OBJ 16.12
 * EVER HAD A PAP SMEAR, AGE 18 AND OLDER August 17, 1998

TABLE 48.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 PAP SMEAR*
 DENOMINATOR IS FEMALES AGE 18 AND OLDER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	95.7	96.5	0.8
ALASKA	97.2	96.2	-1.0
ARIZONA	93.9	84.7	-9.2
ARKANSAS	92.9	92.0	-0.8
COLORADO	96.5	96.5	0.0
CONNECTICUT	93.6	91.8	-1.8
DELAWARE	94.1	96.3	2.2
DISTRICT OF CO	95.8	96.0	0.3
FLORIDA	94.4	93.6	-0.8
GEORGIA	96.0	96.8	0.8
HAWAII	94.1	95.9	1.8
IDAHO	95.9	96.2	0.3
ILLINOIS	94.0	93.5	-0.5

INDIANA	95.3	95.9	0.5
IOWA	95.6	94.3	-1.3
KANSAS	95.1	94.1	-1.0
KENTUCKY	92.9	92.8	-0.1
LOUISIANA	93.0	93.4	0.4
MAINE	95.7	95.7	0.0
MARYLAND	95.4	94.6	-0.8
MASSACHUSETTS	92.9	93.5	0.5
MICHIGAN	96.0	94.7	-1.3
MINNESOTA	93.4	92.9	-0.5
MISSISSIPPI	92.7	95.9	3.2
MISSOURI	95.7	96.5	0.9
MONTANA	96.3	96.2	-0.1
NEBRASKA	94.5	95.2	0.7
NEVADA	96.2	95.5	-0.7
NEW HAMPSHIRE	93.8	94.5	0.7
NEW JERSEY	91.0	91.4	0.3
NEW MEXICO	90.2	93.6	3.3
NEW YORK	92.3	93.1	0.8
NORTH CAROLINA	95.4	95.3	-0.1
NORTH DAKOTA	95.3	94.3	-1.0
OHIO	94.8	94.8	0.0
OKLAHOMA	95.0	95.2	0.3
OREGON	96.2	96.3	0.1
PENNSYLVANIA	93.2	91.8	-1.5
RHODE ISLAND	91.8	93.7	1.9
SOUTH CAROLINA	94.1	96.5	2.4
SOUTH DAKOTA	94.8	95.6	0.8
TENNESSEE	93.4	94.7	1.3
TEXAS	90.8	93.1	2.3
UTAH	90.7	92.5	1.8
VERMONT	94.7	93.1	-1.5
VIRGINIA	94.9	95.1	0.3
WASHINGTON	96.3	96.4	0.1
WEST VIRGINIA	94.1	93.7	-0.4
WISCONSIN	96.5	95.6	-0.8
WYOMING	96.3	95.7	-0.6
PUERTO RICO	81.9	84.5	2.5

* EVER HAD A PAP SMEAR, AGE 18 AND OLDER
 N/A - NOT AVAILABLE

August 17, 1998

TABLE 49.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER HAD A PAP SMEAR*
 DENOMINATOR IS FEMALES AGE 18 AND OVER WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	906	96.4	0.7	(95.0, 97.8)
ALASKA	661	95.9	1.1	(93.6, 98.1)
ARIZONA	793	82.9	2.0	(79.0, 86.8)
ARKANSAS	763	90.8	1.6	(87.7, 93.9)
COLORADO	751	95.8	0.9	(94.1, 97.6)
CONNECTICUT	1096	92.2	1.2	(89.8, 94.5)

DELAWARE	1194	96.5	0.7	(95.1, 97.9)
DISTRICT OF CO	809	96.3	0.8	(94.9, 97.8)
FLORIDA	1522	92.6	0.8	(91.0, 94.1)
GEORGIA	1046	96.5	0.7	(95.1, 97.8)
HAWAII	995	95.3	0.8	(93.7, 96.9)
IDAHO	2067	95.7	0.6	(94.6, 96.8)
ILLINOIS	1417	93.4	0.8	(91.8, 95.0)
INDIANA	995	96.3	0.7	(94.9, 97.6)
IOWA	1653	94.7	0.7	(93.2, 96.1)
KANSAS	955	94.1	0.9	(92.5, 95.8)
KENTUCKY	1615	92.5	0.8	(90.9, 94.1)
LOUISIANA	714	93.0	1.2	(90.7, 95.4)
MAINE	735	95.7	0.9	(93.9, 97.5)
MARYLAND	2073	94.4	0.7	(93.0, 95.8)
MASSACHUSETTS	848	93.7	1.0	(91.7, 95.7)
MICHIGAN	1167	94.5	0.8	(92.9, 96.1)
MINNESOTA	2156	92.4	0.7	(91.0, 93.9)
MISSISSIPPI	678	95.9	0.8	(94.2, 97.5)
MISSOURI	882	95.9	0.9	(94.1, 97.7)
MONTANA	778	96.4	1.1	(94.2, 98.7)
NEBRASKA	1237	94.9	0.7	(93.4, 96.4)
NEVADA	1006	95.1	1.3	(92.5, 97.6)
NEW HAMPSHIRE	710	94.9	1.0	(92.8, 96.9)
NEW JERSEY	1321	91.5	1.0	(89.5, 93.6)
NEW MEXICO	793	92.6	1.4	(89.9, 95.2)
NEW YORK	1765	93.1	0.7	(91.7, 94.6)
NORTH CAROLINA	1594	94.9	0.7	(93.5, 96.2)
NORTH DAKOTA	777	93.9	1.0	(91.8, 95.9)
OHIO	1441	95.2	0.8	(93.6, 96.7)
OKLAHOMA	702	95.1	1.0	(93.1, 97.2)
OREGON	1399	95.4	0.7	(94.0, 96.8)
PENNSYLVANIA	1660	92.5	1.0	(90.5, 94.4)
RHODE ISLAND	837	94.9	0.9	(93.1, 96.7)
SOUTH CAROLINA	978	96.3	0.7	(95.0, 97.7)
SOUTH DAKOTA	977	95.1	0.8	(93.5, 96.7)
TENNESSEE	1367	94.8	0.7	(93.4, 96.2)
TEXAS	1091	92.0	1.0	(90.1, 93.9)
UTAH	1205	91.0	1.4	(88.3, 93.7)
VERMONT	1462	93.7	1.0	(91.7, 95.7)
VIRGINIA	1622	94.7	1.7	(91.4, 98.0)
WASHINGTON	1594	96.2	1.0	(94.3, 98.0)
WEST VIRGINIA	1081	93.4	0.9	(91.5, 95.2)
WISCONSIN	1016	95.1	0.8	(93.5, 96.7)
WYOMING	991	95.0	1.4	(92.2, 97.8)
PUERTO RICO	1076	81.8	1.6	(78.8, 84.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 94.9 RANGE = 81.8-96.5

* EVER HAD A PAP SMEAR, AGE 18 AND OLDER

August 17, 1998

TABLE 49.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1996 TO 1997

EVER HAD A PAP SMEAR*

DENOMINATOR IS FEMALES AGE 18 AND OVER WITH UTERINE CERVIX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	95.7	96.4	0.7
ALASKA	97.2	95.9	-1.3
ARIZONA	93.9	82.9	-10.9
ARKANSAS	92.9	90.8	-2.1
COLORADO	96.5	95.8	-0.7
CONNECTICUT	93.6	92.2	-1.4
DELAWARE	94.1	96.5	2.4
DISTRICT OF CO	95.8	96.3	0.6
FLORIDA	94.4	92.6	-1.9
GEORGIA	96.0	96.5	0.5
HAWAII	94.1	95.3	1.1
IDAHO	95.9	95.7	-0.2
ILLINOIS	94.0	93.4	-0.6
INDIANA	95.3	96.3	0.9
IOWA	95.6	94.7	-0.9
KANSAS	95.1	94.1	-1.0
KENTUCKY	92.9	92.5	-0.4
LOUISIANA	93.0	93.0	0.0
MAINE	95.7	95.7	0.1
MARYLAND	95.4	94.4	-0.9
MASSACHUSETTS	92.9	93.7	0.7
MICHIGAN	96.0	94.5	-1.5
MINNESOTA	93.4	92.4	-1.0
MISSISSIPPI	92.7	95.9	3.2
MISSOURI	95.7	95.9	0.2
MONTANA	96.3	96.4	0.2
NEBRASKA	94.5	94.9	0.4
NEVADA	96.2	95.1	-1.1
NEW HAMPSHIRE	93.8	94.9	1.1
NEW JERSEY	91.0	91.5	0.5
NEW MEXICO	90.2	92.6	2.3
NEW YORK	92.3	93.1	0.9
NORTH CAROLINA	95.4	94.9	-0.5
NORTH DAKOTA	95.3	93.9	-1.4
OHIO	94.8	95.2	0.3
OKLAHOMA	95.0	95.1	0.1
OREGON	96.2	95.4	-0.7
PENNSYLVANIA	93.2	92.5	-0.8
RHODE ISLAND	91.8	94.9	3.1
SOUTH CAROLINA	94.1	96.3	2.3
SOUTH DAKOTA	94.8	95.1	0.3
TENNESSEE	93.4	94.8	1.4
TEXAS	90.8	92.0	1.2
UTAH	90.7	91.0	0.3
VERMONT	94.7	93.7	-1.0
VIRGINIA	94.9	94.7	-0.1
WASHINGTON	96.3	96.2	-0.1
WEST VIRGINIA	94.1	93.4	-0.7
WISCONSIN	96.5	95.1	-1.4
WYOMING	96.3	95.0	-1.3
PUERTO RICO	81.9	81.8	-0.1

* EVER HAD A PAP SMEAR, AGE 18 AND OLDER

N/A - NOT AVAILABLE

August 17, 1998

TABLE 50.1 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LAST PAP SMEAR*
 DENOMINATOR IS FEMALES AGE 18 AND OVER WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	891	86.0	1.3	(83.5, 88.4)
ALASKA	657	90.3	1.8	(86.7, 93.8)
ARIZONA	793	79.3	2.1	(75.1, 83.5)
ARKANSAS	749	78.2	1.9	(74.4, 82.0)
COLORADO	749	87.6	1.4	(85.0, 90.3)
CONNECTICUT	1091	83.0	1.5	(80.1, 85.8)
DELAWARE	1186	89.8	1.0	(87.9, 91.7)
DISTRICT OF CO	808	92.1	1.1	(89.9, 94.2)
FLORIDA	1515	84.7	1.1	(82.6, 86.8)
GEORGIA	1043	92.3	0.9	(90.4, 94.1)
HAWAII	987	86.9	1.3	(84.3, 89.4)
IDAHO	2060	81.6	1.0	(79.6, 83.6)
ILLINOIS	1415	82.7	1.1	(80.5, 85.0)
INDIANA	983	84.5	1.3	(81.9, 87.1)
IOWA	1644	80.5	1.2	(78.2, 82.8)
KANSAS	943	85.9	1.2	(83.5, 88.3)
KENTUCKY	1607	81.7	1.1	(79.6, 83.8)
LOUISIANA	709	83.8	1.6	(80.7, 86.8)
MAINE	733	88.1	1.4	(85.3, 90.9)
MARYLAND	2060	88.7	0.9	(86.9, 90.5)
MASSACHUSETTS	845	88.4	1.3	(85.9, 90.9)
MICHIGAN	1154	86.3	1.1	(84.1, 88.6)
MINNESOTA	2138	83.4	0.9	(81.6, 85.2)
MISSISSIPPI	676	85.9	1.4	(83.1, 88.7)
MISSOURI	877	83.9	1.6	(80.8, 86.9)
MONTANA	767	84.6	1.6	(81.5, 87.8)
NEBRASKA	1227	84.1	1.2	(81.8, 86.4)
NEVADA	998	85.6	2.0	(81.7, 89.5)
NEW HAMPSHIRE	709	87.0	1.5	(84.1, 89.9)
NEW JERSEY	1317	81.3	1.3	(78.7, 83.9)
NEW MEXICO	792	80.6	1.7	(77.2, 84.0)
NEW YORK	1747	87.0	0.9	(85.1, 88.8)
NORTH CAROLINA	1580	87.4	1.0	(85.4, 89.3)
NORTH DAKOTA	773	82.4	1.5	(79.4, 85.3)
OHIO	1434	87.3	1.2	(85.0, 89.5)
OKLAHOMA	699	84.4	1.6	(81.2, 87.6)
OREGON	1391	86.0	1.1	(83.9, 88.2)
PENNSYLVANIA	1643	82.3	1.3	(79.8, 84.8)
RHODE ISLAND	826	87.4	1.2	(85.0, 89.8)
SOUTH CAROLINA	974	89.7	1.1	(87.5, 91.9)
SOUTH DAKOTA	972	85.0	1.3	(82.5, 87.4)
TENNESSEE	1359	87.8	1.0	(85.7, 89.8)
TEXAS	1085	80.9	1.4	(78.2, 83.6)
UTAH	1204	78.5	1.7	(75.1, 81.9)
VERMONT	1444	84.1	1.3	(81.6, 86.5)
VIRGINIA	1605	87.9	1.8	(84.4, 91.3)
WASHINGTON	1583	87.0	1.2	(84.7, 89.3)
WEST VIRGINIA	1069	78.9	1.4	(76.1, 81.7)
WISCONSIN	1009	83.8	1.5	(80.9, 86.7)
WYOMING	982	83.7	1.7	(80.3, 87.1)
PUERTO RICO	1067	71.8	1.7	(68.6, 75.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 84.7 RANGE = 71.8-92.3

* LAST PAP SMEAR WITHIN LAST 3 YEARS, AGE 18 AND OLDER

August 25, 1998

TABLE 50.2 1997 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1997
 LAST PAP SMEAR*

DENOMINATOR IS FEMALES AGE 18 AND OVER WITH UTERINE CERVIX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1996 PERCENT	1997 PERCENT	PERCENT DIFFERENCE
ALABAMA	83.0	86.0	2.9
ALASKA	90.6	90.3	-0.3
ARIZONA	84.0	79.3	-4.7
ARKANSAS	79.9	78.2	-1.7
COLORADO	86.1	87.6	1.5
CONNECTICUT	81.2	83.0	1.8
DELAWARE	84.2	89.8	5.7
DISTRICT OF CO	90.0	92.1	2.1
FLORIDA	81.5	84.7	3.3
GEORGIA	90.8	92.3	1.5
HAWAII	86.1	86.9	0.8
IDAHO	78.0	81.6	3.6
ILLINOIS	81.3	82.7	1.5
INDIANA	78.1	84.5	6.4
IOWA	76.9	80.5	3.6
KANSAS	84.4	85.9	1.5
KENTUCKY	76.6	81.7	5.2
LOUISIANA	79.6	83.8	4.2
MAINE	81.7	88.1	6.4
MARYLAND	85.8	88.7	2.9
MASSACHUSETTS	81.4	88.4	7.0
MICHIGAN	82.7	86.3	3.7
MINNESOTA	83.8	83.4	-0.4
MISSISSIPPI	79.1	85.9	6.8
MISSOURI	82.5	83.9	1.4
MONTANA	83.1	84.6	1.5
NEBRASKA	78.7	84.1	5.3
NEVADA	82.2	85.6	3.4
NEW HAMPSHIRE	81.1	87.0	5.9
NEW JERSEY	79.3	81.3	2.0
NEW MEXICO	80.2	80.6	0.3
NEW YORK	82.2	87.0	4.8
NORTH CAROLINA	85.6	87.4	1.8
NORTH DAKOTA	78.3	82.4	4.0
OHIO	82.8	87.3	4.5
OKLAHOMA	80.2	84.4	4.1
OREGON	81.5	86.0	4.5
PENNSYLVANIA	78.2	82.3	4.1
RHODE ISLAND	77.8	87.4	9.6
SOUTH CAROLINA	85.0	89.7	4.7
SOUTH DAKOTA	82.0	85.0	2.9
TENNESSEE	77.3	87.8	10.5
TEXAS	78.8	80.9	2.1
UTAH	77.4	78.5	1.1
VERMONT	83.8	84.1	0.2
VIRGINIA	84.9	87.9	3.0
WASHINGTON	83.4	87.0	3.6
WEST VIRGINIA	78.0	78.9	0.9
WISCONSIN	83.3	83.8	0.6
WYOMING	78.9	83.7	4.8
PUERTO RICO	71.0	71.8	0.8

* LAST PAP SMEAR WITHIN LAST 3 YEARS, AGE 18 AND OLDER
N/A - NOT AVAILABLE

August 25, 1998