

Table 1.96A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	46,558	14,680	6,210
GEOGRAPHIC DIVISION			
Northeast	7,917	2,343	921
New England	2,506	834	294
Middle Atlantic	5,411	1,509	627
Midwest	9,928	3,079	1,178
East North Central	6,865	2,233	812
West North Central	3,063	846	366
South	16,186	5,458	2,450
South Atlantic	8,345	2,803	1,175
East South Central	2,667	981	546
West South Central	5,174	1,674	729
West	12,527	3,799	1,661
Mountain	3,652	1,105	526
Pacific	8,875	2,695	1,135
COUNTY TYPE			
Large Metro	23,610	7,032	2,940
Small Metro	14,805	5,104	2,187
250K – 1 Mil. Pop.	10,876	3,603	1,625
<250K Pop.	3,928	1,502	562
Nonmetro	8,144	2,544	1,083
Urbanized	2,843	805	317
Less Urbanized	4,504	1,469	599
Completely Rural	797	271	168

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.96B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	19.8	6.2	2.6
GEOGRAPHIC DIVISION			
Northeast	17.6	5.2	2.0
New England	21.2	7.1	2.5
Middle Atlantic	16.3	4.5	1.9
Midwest	18.6	5.8	2.2
East North Central	18.3	6.0	2.2
West North Central	19.1	5.3	2.3
South	19.3	6.5	2.9
South Atlantic	19.1	6.4	2.7
East South Central	18.8	6.9	3.9
West South Central	20.0	6.5	2.8
West	23.7	7.2	3.1
Mountain	23.8	7.2	3.4
Pacific	23.6	7.2	3.0
COUNTY TYPE			
Large Metro	20.1	6.0	2.5
Small Metro	20.9	7.2	3.1
250K – 1 Mil. Pop.	21.0	7.0	3.1
<250K Pop.	20.5	7.9	2.9
Nonmetro	17.3	5.4	2.3
Urbanized	19.1	5.4	2.1
Less Urbanized	16.7	5.4	2.2
Completely Rural	15.3	5.2	3.2

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.97A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	3,389	2,283	979
GEOGRAPHIC DIVISION			
Northeast	523	331	152
New England	177	105	53
Middle Atlantic	347	226	99
Midwest	811	548	204
East North Central	569	393	148
West North Central	242	155	57
South	1,212	843	393
South Atlantic	599	428	198
East South Central	202	143	67
West South Central	411	272	128
West	842	561	230
Mountain	256	171	67
Pacific	587	390	163
COUNTY TYPE			
Large Metro	1,649	1,079	455
Small Metro	1,014	721	312
250K – 1 Mil. Pop.	742	518	240
<250K Pop.	272	203	72
Nonmetro	726	483	211
Urbanized	218	141	68
Less Urbanized	429	287	121
Completely Rural	79	55	22

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.97B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	13.7	9.2	4.0
GEOGRAPHIC DIVISION			
Northeast	11.7	7.4	3.4
New England	15.3	9.1	4.5
Middle Atlantic	10.5	6.8	3.0
Midwest	14.3	9.6	3.6
East North Central	14.3	9.8	3.7
West North Central	14.2	9.1	3.3
South	13.8	9.6	4.5
South Atlantic	13.6	9.7	4.5
East South Central	14.2	10.1	4.7
West South Central	13.9	9.2	4.3
West	14.4	9.6	3.9
Mountain	15.0	10.1	3.9
Pacific	14.2	9.4	4.0
COUNTY TYPE			
Large Metro	13.3	8.7	3.7
Small Metro	13.8	9.8	4.2
250K – 1 Mil. Pop.	13.6	9.5	4.4
<250K Pop.	14.4	10.7	3.8
Nonmetro	14.6	9.7	4.3
Urbanized	14.1	9.1	4.4
Less Urbanized	15.0	10.0	4.2
Completely Rural	14.1	9.9	4.0

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.98A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	8,587	4,411	1,683
GEOGRAPHIC DIVISION			
Northeast	1,370	725	281
New England	466	271	99
Middle Atlantic	904	454	182
Midwest	1,990	995	360
East North Central	1,423	740	262
West North Central	567	255	98
South	3,095	1,727	694
South Atlantic	1,461	804	288
East South Central	586	340	150
West South Central	1,048	583	256
West	2,132	964	347
Mountain	717	354	126
Pacific	1,415	610	221
COUNTY TYPE			
Large Metro	3,813	1,829	698
Small Metro	3,199	1,763	672
250K – 1 Mil. Pop.	2,242	1,215	479
<250K Pop.	957	548	194
Nonmetro	1,575	819	312
Urbanized	567	295	119
Less Urbanized	864	452	163
Completely Rural	144	72	30

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.98B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	27.7	14.2	5.4
GEOGRAPHIC DIVISION			
Northeast	24.9	13.2	5.1
New England	32.5	18.9	6.9
Middle Atlantic	22.2	11.2	4.5
Midwest	27.6	13.8	5.0
East North Central	28.6	14.9	5.3
West North Central	25.5	11.5	4.4
South	28.1	15.7	6.3
South Atlantic	27.1	14.9	5.3
East South Central	30.7	17.8	7.8
West South Central	28.2	15.7	6.9
West	29.2	13.2	4.8
Mountain	32.8	16.2	5.8
Pacific	27.7	11.9	4.3
COUNTY TYPE			
Large Metro	25.8	12.4	4.7
Small Metro	30.6	16.9	6.4
250K – 1 Mil. Pop.	30.2	16.4	6.5
<250K Pop.	31.5	18.1	6.4
Nonmetro	27.1	14.1	5.4
Urbanized	26.7	13.9	5.6
Less Urbanized	27.3	14.3	5.2
Completely Rural	27.6	13.8	5.7

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.99A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	34,582	7,986	3,549
GEOGRAPHIC DIVISION			
Northeast	6,024	1,287	488
New England	1,864	458	142
Middle Atlantic	4,160	828	346
Midwest	7,127	1,537	614
East North Central	4,874	1,101	402
West North Central	2,254	435	212
South	11,879	2,888	1,363
South Atlantic	6,286	1,571	689
East South Central	1,878	497	329
West South Central	3,715	820	345
West	9,552	2,275	1,083
Mountain	2,679	579	333
Pacific	6,873	1,696	751
COUNTY TYPE			
Large Metro	18,148	4,124	1,786
Small Metro	10,591	2,620	1,203
250K – 1 Mil. Pop.	7,892	1,870	906
<250K Pop.	2,699	750	296
Nonmetro	5,843	1,242	560
Urbanized	2,058	369	129
Less Urbanized	3,211	730	315
Completely Rural	574	143	115

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.99B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	19.3	4.5	2.0
GEOGRAPHIC DIVISION			
Northeast	17.2	3.7	1.4
New England	20.2	5.0	1.5
Middle Atlantic	16.1	3.2	1.3
Midwest	17.6	3.8	1.5
East North Central	17.1	3.9	1.4
West North Central	18.7	3.6	1.8
South	18.6	4.5	2.1
South Atlantic	18.5	4.6	2.0
East South Central	17.4	4.6	3.0
West South Central	19.4	4.3	1.8
West	24.0	5.7	2.7
Mountain	23.5	5.1	2.9
Pacific	24.3	6.0	2.6
COUNTY TYPE			
Large Metro	20.1	4.6	2.0
Small Metro	20.0	4.9	2.3
250K – 1 Mil. Pop.	20.3	4.8	2.3
<250K Pop.	19.0	5.3	2.1
Nonmetro	16.1	3.4	1.5
Urbanized	18.4	3.3	1.2
Less Urbanized	15.3	3.5	1.5
Completely Rural	13.9	3.5	2.8

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.100A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	43,169	12,397	5,231
GEOGRAPHIC DIVISION			
Northeast	7,394	2,012	770
New England	2,329	729	241
Middle Atlantic	5,064	1,283	528
Midwest	9,117	2,531	974
East North Central	6,296	1,841	664
West North Central	2,820	691	309
South	14,974	4,615	2,057
South Atlantic	7,747	2,375	977
East South Central	2,465	838	479
West South Central	4,763	1,402	601
West	11,684	3,238	1,431
Mountain	3,396	933	459
Pacific	8,288	2,305	972
COUNTY TYPE			
Large Metro	21,961	5,953	2,484
Small Metro	13,790	4,383	1,875
250K – 1 Mil. Pop.	10,134	3,085	1,385
<250K Pop.	3,656	1,298	490
Nonmetro	7,418	2,061	872
Urbanized	2,625	664	248
Less Urbanized	4,075	1,182	478
Completely Rural	718	215	145

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.100B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2002

Geographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
TOTAL	20.5	5.9	2.5
GEOGRAPHIC DIVISION			
Northeast	18.2	5.0	1.9
New England	21.9	6.8	2.3
Middle Atlantic	16.9	4.3	1.8
Midwest	19.1	5.3	2.0
East North Central	18.8	5.5	2.0
West North Central	19.7	4.8	2.2
South	20.0	6.2	2.7
South Atlantic	19.7	6.0	2.5
East South Central	19.4	6.6	3.8
West South Central	20.8	6.1	2.6
West	24.8	6.9	3.0
Mountain	25.0	6.9	3.4
Pacific	24.8	6.9	2.9
COUNTY TYPE			
Large Metro	20.9	5.7	2.4
Small Metro	21.7	6.9	3.0
250K – 1 Mil. Pop.	21.9	6.7	3.0
<250K Pop.	21.2	7.5	2.8
Nonmetro	17.6	4.9	2.1
Urbanized	19.7	5.0	1.9
Less Urbanized	16.9	4.9	2.0
Completely Rural	15.4	4.6	3.1

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.