

Table 1.106A Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Numbers in Thousands, 2002

Drug or Method of Administration	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hallucinogens	34,314	1,419	7,517	25,378
PCP (Angel Dust, Phencyclidine)	7,418	233	834	6,352
LSD (Acid)	24,516	670	4,928	18,917
Peyote	5,738	99	533	5,105
Mescaline	9,116	106	776	8,233
Psilocybin (Mushrooms)	18,322	564	4,167	13,592
"Ecstasy" (MDMA)	10,150	820	4,674	4,656
Inhalants	22,870	2,605	4,867	15,398
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	8,915	431	899	7,585
Correction Fluid, Degreaser, or Cleaning Fluid	2,248	550	520	1,178
Gasoline or Lighter Fluid	4,023	862	910	2,250
Glue, Shoe Polish, or Toluene	4,551	1,105	805	2,641
Halothane, Ether, or Other Anesthetics	1,114	96	256	762
Lacquer Thinner or Other Paint Solvents	2,006	427	487	1,093
Lighter Gases (Butane, Propane)	1,041	270	314	458
Nitrous Oxide or "Whippets"	11,698	526	3,308	7,863
Spray Paints	2,012	631	479	903
Other Aerosol Sprays	2,011	427	613	971
Any Needle Use^{1,2}	3,982	81	375	3,526
Heroin Needle Use ^{1,2}	1,674	27	151	1,497
Cocaine Needle Use ^{1,2}	2,499	18	135	2,345
Stimulant Needle Use ^{1,2}	1,805	37	169	1,600
Methamphetamine Needle Use ^{1,2}	1,314	20	129	1,165
Heroin	3,668	94	487	3,087
Smoke Heroin ²	1,379	40	194	1,145
Sniff or Snort Heroin ²	2,554	60	336	2,159

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.106B Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Percentages, 2002

Drug or Method of Administration	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Hallucinogens	14.6	5.7	24.2	14.1
PCP (Angel Dust, Phencyclidine)	3.2	0.9	2.7	3.5
LSD (Acid)	10.4	2.7	15.9	10.5
Peyote	2.4	0.4	1.7	2.8
Mescaline	3.9	0.4	2.5	4.6
Psilocybin (Mushrooms)	7.8	2.3	13.4	7.6
"Ecstasy" (MDMA)	4.3	3.3	15.1	2.6
Inhalants	9.7	10.5	15.7	8.6
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	3.8	1.7	2.9	4.2
Correction Fluid, Degreaser, or Cleaning Fluid	1.0	2.2	1.7	0.7
Gasoline or Lighter Fluid	1.7	3.5	2.9	1.3
Glue, Shoe Polish, or Toluene	1.9	4.5	2.6	1.5
Halothane, Ether, or Other Anesthetics	0.5	0.4	0.8	0.4
Lacquer Thinner or Other Paint Solvents	0.9	1.7	1.6	0.6
Lighter Gases (Butane, Propane)	0.4	1.1	1.0	0.3
Nitrous Oxide or "Whippets"	5.0	2.1	10.7	4.4
Spray Paints	0.9	2.5	1.5	0.5
Other Aerosol Sprays	0.9	1.7	2.0	0.5
Any Needle Use^{1,2}	1.7	0.3	1.2	2.0
Heroin Needle Use ^{1,2}	0.7	0.1	0.5	0.8
Cocaine Needle Use ^{1,2}	1.1	0.1	0.4	1.3
Stimulant Needle Use ^{1,2}	0.8	0.1	0.5	0.9
Methamphetamine Needle Use ^{1,2}	0.6	0.1	0.4	0.6
Heroin	1.6	0.4	1.6	1.7
Smoke Heroin ²	0.6	0.2	0.6	0.6
Sniff or Snort Heroin ²	1.1	0.2	1.1	1.2

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

30408

Table 1.107A Specific Pain Reliever Use in Lifetime, by Age Group: Numbers in Thousands, 2002

Pain Reliever	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Darvocet®, Darvon®, or Tylenol® with Codeine	18,912	1,508	3,800	13,603
Percocet®, Percodan®, or Tylox®	9,659	407	2,285	6,967
Vicodin®, Lortab®, or Lorcet®	13,093	1,116	4,126	7,852
Codeine	6,897	513	1,878	4,506
Demerol®	2,917	116	657	2,144
Dilaudid®	1,116	7	101	1,008
Fioricet®	445	13	64	368
Fiorinal®	579	24	40	515
Hydrocodone	4,529	333	1,540	2,656
Methadone	928	116	306	506
Morphine	2,073	256	648	1,170
Oxycontin®	1,924	224	820	881
Phenaphen® with Codeine	873	69	194	610
Propoxyphene	282	31	45	206
SK-65®	42	13	10	19
Stadol®	133	6	23	103
Talacen®	55	14	13	28
Talwin®	643	10	20	613
Talwin NX®	105	10	9	86
Tramadol	52	11	20	21
Ultram®	962	98	300	564

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.107B Specific Pain Reliever Use in Lifetime, by Age Group: Percentages, 2002

Pain Reliever	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Darvocet®, Darvon®, or Tylenol® with Codeine	8.0	6.1	12.2	7.6
Percocet®, Percodan®, or Tylox®	4.1	1.6	7.4	3.9
Vicodin®, Lortab®, or Lorcet®	5.6	4.5	13.3	4.4
Codeine	2.9	2.1	6.1	2.5
Demerol®	1.2	0.5	2.1	1.2
Dilaudid®	0.5	0.0	0.3	0.6
Fioricet®	0.2	0.1	0.2	0.2
Fiorinal®	0.2	0.1	0.1	0.3
Hydrocodone	1.9	1.3	5.0	1.5
Methadone	0.4	0.5	1.0	0.3
Morphine	0.9	1.0	2.1	0.7
Oxycontin®	0.8	0.9	2.6	0.5
Phenaphen® with Codeine	0.4	0.3	0.6	0.3
Propoxyphene	0.1	0.1	0.1	0.1
SK-65®	0.0	0.1	0.0	0.0
Stadol®	0.1	0.0	0.1	0.1
Talacen®	0.0	0.1	0.0	0.0
Talwin®	0.3	0.0	0.1	0.3
Talwin NX®	0.0	0.0	0.0	0.0
Tramadol	0.0	0.0	0.1	0.0
Ultram®	0.4	0.4	1.0	0.3

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.108A Specific Tranquilizer Use in Lifetime, by Age Group: Numbers in Thousands, 2002

Tranquilizer	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Klonopin® or Clonazepam	2,374	180	724	1,470
Xanax®, Alprazolam, Ativan®, or Lorazepam	8,208	429	2,091	5,688
Valium® or Diazepam	14,386	427	2,312	11,648
Atarax®	371	26	45	301
BuSpar®	693	51	220	421
Equanil®	75	14	12	49
Flexeril®	1,781	47	295	1,439
Librium®	1,454	17	79	1,358
Limbitrol®	115	9	32	73
Meprobamate	161	9	21	131
Miltown®	169	12	6	151
Rohypnol®	375	40	160	175
Serax®	215	17	29	169
Soma®	2,276	142	729	1,405
Tranxene®	214	10	16	188
Vistaril®	356	21	64	271

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.108B Specific Tranquilizer Use in Lifetime, by Age Group: Percentages, 2002

Tranquilizer	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Klonopin® or Clonazepam	1.0	0.7	2.3	0.8
Xanax®, Alprazolam, Ativan®, or Lorazepam	3.5	1.7	6.7	3.2
Valium® or Diazepam	6.1	1.7	7.5	6.5
Atarax®	0.2	0.1	0.1	0.2
BuSpar®	0.3	0.2	0.7	0.2
Equanil®	0.0	0.1	0.0	0.0
Flexeril®	0.8	0.2	0.9	0.8
Librium®	0.6	0.1	0.3	0.8
Limbitrol®	0.0	0.0	0.1	0.0
Meprobamate	0.1	0.0	0.1	0.1
Miltown®	0.1	0.0	0.0	0.1
Rohypnol®	0.2	0.2	0.5	0.1
Serax®	0.1	0.1	0.1	0.1
Soma®	1.0	0.6	2.3	0.8
Tranxene®	0.1	0.0	0.1	0.1
Vistaril®	0.2	0.1	0.2	0.2

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.109A Specific Stimulant Use in Lifetime, by Age Group: Numbers in Thousands, 2002

Stimulant	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Methamphetamine, ¹ Desoxyn®, or Methedrine	12,383	366	1,756	10,261
Prescription Diet Pills ²	8,980	267	692	8,021
Ritalin® or Methylphenidate	4,528	587	1,662	2,280
Cylert®	196	18	39	139
Dexedrine®	3,274	70	318	2,887
Dextroamphetamine	553	22	78	453
Didrex®	177	21	31	124
Eskatrol®	302	10	15	276
Ionamin®	491	19	28	445
Mazanor®	65	16	3	45
Obedrin-LA®	79	6	6	68
Plegine®	68	12	8	48
Preludin®	856	10	35	811
Sanorex®	137	12	23	102
Tenuate®	275	10	10	255

*Low precision; no estimate reported.

¹ Crank, crystal, ice, or speed.

² For example, Amphetamines, Benzedrine®, Biphedamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.109B Specific Stimulant Use in Lifetime, by Age Group: Percentages, 2002

Stimulant	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Methamphetamine, ¹ Desoxyn®, or Methedrine	5.3	1.5	5.7	5.7
Prescription Diet Pills ²	3.8	1.1	2.2	4.5
Ritalin® or Methylphenidate	1.9	2.4	5.4	1.3
Cylert®	0.1	0.1	0.1	0.1
Dexedrine®	1.4	0.3	1.0	1.6
Dextroamphetamine	0.2	0.1	0.3	0.3
Didrex®	0.1	0.1	0.1	0.1
Eskatrol®	0.1	0.0	0.0	0.2
Ionamin®	0.2	0.1	0.1	0.2
Mazanor®	0.0	0.1	0.0	0.0
Obedrin-LA®	0.0	0.0	0.0	0.0
Plegine®	0.0	0.0	0.0	0.0
Preludin®	0.4	0.0	0.1	0.5
Sanorex®	0.1	0.0	0.1	0.1
Tenuate®	0.1	0.0	0.0	0.1

*Low precision; no estimate reported.

¹ Crank, crystal, ice, or speed.

² For example, Amphetamines, Benzedrine®, Biphedamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

30408

Table 1.110A Specific Sedative Use in Lifetime, by Age Group: Numbers in Thousands, 2002

Sedative	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Methaqualone, Sopor®, or Quaalude®	7,320	60	305	6,955
Barbiturates ¹	3,375	29	121	3,226
Restoril® or Temazepam	1,003	48	108	847
Amytal®	274	4	21	249
Butisol®	109	8	14	87
Chloral Hydrate	214	7	9	198
Dalmane®	638	22	16	600
Halcion®	939	17	64	857
Phenobarbital	1,568	37	105	1,426
Placidyl®	1,045	12	34	998
Tuinal®	1,393	8	17	1,368

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 1.110B Specific Sedative Use in Lifetime, by Age Group: Percentages, 2002

Sedative	Total	AGE GROUP (Years)		
		12-17	18-25	26 or Older
Methaqualone, Sopor®, or Quaalude®	3.1	0.2	1.0	3.9
Barbiturates ¹	1.4	0.1	0.4	1.8
Restoril® or Temazepam	0.4	0.2	0.3	0.5
Amytal®	0.1	0.0	0.1	0.1
Butisol®	0.0	0.0	0.0	0.0
Chloral Hydrate	0.1	0.0	0.0	0.1
Dalmane®	0.3	0.1	0.1	0.3
Halcion®	0.4	0.1	0.2	0.5
Phenobarbital	0.7	0.2	0.3	0.8
Placidyl®	0.4	0.1	0.1	0.6
Tuinal®	0.6	0.0	0.1	0.8

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.