

Table 1.1A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 12 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	87,734	86,931	25,402	24,535	13,829	14,027
Marijuana and Hashish	76,428	76,321	19,102	18,589	10,458	10,714
Cocaine	25,406	24,896	3,742	3,328	1,552	1,213
Crack	5,910	5,307	1,045 ^a	721	418	265
Heroin	3,054	2,779	353	308	154	130
Hallucinogens	25,061	26,125	3,191	3,483	922	971
LSD	19,215	19,642	1,905	1,749	482	403
PCP	5,693	5,804	245	264	55	54
Inhalants	17,138	16,702	1,937	1,918	570	622
Nonmedical Use of Any Psychotherapeutic ²	34,076	32,443	9,220	8,761	3,952	3,849
Pain Relievers	19,888	19,210	6,582	6,466	2,621	2,782
Tranquilizers	13,860	13,007	2,728	2,731	1,097	1,000
Stimulants	15,922 ^a	14,661	2,291	2,112	950	788
Methamphetamine	9,442	8,843	1,140	1,031	435	387
Sedatives	7,747	7,142	631	611	229	175
Any Illicit Drug Other Than Marijuana ¹	53,268	52,605	13,444	13,052	5,981	5,711

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.1B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	39.7	38.9	11.5	11.0	6.3	6.3
Marijuana and Hashish	34.6	34.2	8.6	8.3	4.7	4.8
Cocaine	11.5	11.2	1.7	1.5	0.7	0.5
Crack	2.7	2.4	0.5 ^a	0.3	0.2	0.1
Heroin	1.4	1.2	0.2	0.1	0.1	0.1
Hallucinogens	11.3	11.7	1.4	1.6	0.4	0.4
LSD	8.7	8.8	0.9	0.8	0.2	0.2
PCP	2.6	2.6	0.1	0.1	0.0	0.0
Inhalants	7.8	7.5	0.9	0.9	0.3	0.3
Nonmedical Use of Any Psychotherapeutic ²	15.4 ^a	14.5	4.2	3.9	1.8	1.7
Pain Relievers	9.0	8.6	3.0	2.9	1.2	1.2
Tranquilizers	6.3	5.8	1.2	1.2	0.5	0.4
Stimulants	7.2 ^a	6.6	1.0	0.9	0.4	0.4
Methamphetamine	4.3	4.0	0.5	0.5	0.2	0.2
Sedatives	3.5	3.2	0.3	0.3	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	24.1	23.6	6.1	5.8	2.7	2.6

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.2A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	6,415	6,288	4,584	4,357	2,265	2,264
Marijuana and Hashish	4,345	4,283	3,284	3,136	1,676	1,678
Cocaine	551	550	373	389	124	132
Crack	143	143	90	87	18	20
Heroin	102	84	65	49	38 ^a	15
Hallucinogens	1,317	1,347	889	908	254	270
LSD	877	840	554	511	150	120
PCP	217	246	99	112	26	18
Inhalants	2,118	2,079	908	826	245	223
Nonmedical Use of Any Psychotherapeutic ²	2,524	2,543	1,653	1,649	664	690
Pain Relievers	1,894	1,964	1,266	1,260	496	538
Tranquilizers	574	595	362	373	119	110
Stimulants	915	927	492	552	151	181
Methamphetamine	317	305	163	186	53	68
Sedatives	190	188	111	110	44	36
Any Illicit Drug Other Than Marijuana ¹	4,248	4,228	2,698	2,631	1,045	1,074

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.2B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	27.6	26.9	19.8 ^a	18.6	9.8	9.7
Marijuana and Hashish	18.7	18.3	14.2	13.4	7.2	7.2
Cocaine	2.4	2.4	1.6	1.7	0.5	0.6
Crack	0.6	0.6	0.4	0.4	0.1	0.1
Heroin	0.4	0.4	0.3	0.2	0.2 ^a	0.1
Hallucinogens	5.7	5.8	3.8	3.9	1.1	1.2
LSD	3.8	3.6	2.4	2.2	0.6	0.5
PCP	0.9	1.1	0.4	0.5	0.1	0.1
Inhalants	9.1	8.9	3.9	3.5	1.1	1.0
Nonmedical Use of Any Psychotherapeutic ²	10.9	10.9	7.1	7.1	2.9	3.0
Pain Relievers	8.2	8.4	5.5	5.4	2.1	2.3
Tranquilizers	2.5	2.5	1.6	1.6	0.5	0.5
Stimulants	3.9	4.0	2.1	2.4	0.7	0.8
Methamphetamine	1.4	1.3	0.7	0.8	0.2	0.3
Sedatives	0.8	0.8	0.5	0.5	0.2	0.2
Any Illicit Drug Other Than Marijuana ¹	18.3	18.1	11.6	11.3	4.5	4.6

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.3A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	14,975	14,846	8,287	8,088	4,658	4,599
Marijuana and Hashish	13,326	13,256	6,962	6,860	4,049	3,950
Cocaine	3,383	3,148	1,493 ^b	1,274	476	395
Crack	936	822	271 ^a	190	74 ^a	43
Heroin	498	403	129	108	36	26
Hallucinogens	5,490	5,592	1,928	1,959	534	532
LSD	4,185	4,072	1,132	998	276	235
PCP	691	670	124	99	29	23
Inhalants	4,014 ^a	3,701	747	696	182	162
Nonmedical Use of Any Psychotherapeutic ²	5,957	5,650	2,830	2,696	1,062	1,050
Pain Relievers	4,316	4,221	2,160	2,108	743	780
Tranquilizers	2,239	2,158	894	883	303	292
Stimulants	2,560 ^b	2,200	841 ^a	703	299 ^a	221
Methamphetamine	1,483 ^b	1,185	425	353	137	94
Sedatives	561	477	174	169	57	41
Any Illicit Drug Other Than Marijuana ¹	9,476	9,247	4,460	4,295	1,729	1,711

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.3B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	52.6	51.2	29.1	27.9	16.4	15.9
Marijuana and Hashish	46.8	45.7	24.5	23.7	14.2	13.6
Cocaine	11.9 ^a	10.9	5.2 ^b	4.4	1.7	1.4
Crack	3.3 ^a	2.8	1.0 ^b	0.7	0.3 ^a	0.1
Heroin	1.8 ^a	1.4	0.5	0.4	0.1	0.1
Hallucinogens	19.3	19.3	6.8	6.8	1.9	1.8
LSD	14.7	14.0	4.0 ^a	3.4	1.0	0.8
PCP	2.4	2.3	0.4	0.3	0.1	0.1
Inhalants	14.1 ^b	12.8	2.6	2.4	0.6	0.6
Nonmedical Use of Any Psychotherapeutic ²	20.9 ^b	19.5	9.9	9.3	3.7	3.6
Pain Relievers	15.2	14.6	7.6	7.3	2.6	2.7
Tranquilizers	7.9	7.4	3.1	3.0	1.1	1.0
Stimulants	9.0 ^b	7.6	3.0 ^a	2.4	1.1 ^a	0.8
Methamphetamine	5.2 ^b	4.1	1.5	1.2	0.5	0.3
Sedatives	2.0	1.6	0.6	0.6	0.2	0.1
Any Illicit Drug Other Than Marijuana ¹	33.3 ^a	31.9	15.7	14.8	6.1	5.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.4A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 26 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	66,344	65,797	12,531	12,089	6,906	7,164
Marijuana and Hashish	58,757	58,782	8,856	8,593	4,734	5,085
Cocaine	21,471	21,198	1,876	1,666	952	687
Crack	4,831	4,341	683	443	325	202
Heroin	2,454	2,292	159	151	80	88
Hallucinogens	18,254	19,186	374 ^a	616	134	169
LSD	14,152	14,729	219	239	56	49
PCP	4,785	4,888	23	53	*	12
Inhalants	11,006	10,922	282	396	144	236
Nonmedical Use of Any Psychotropic ²	25,596	24,250	4,737	4,415	2,225	2,108
Pain Relievers	13,678	13,025	3,156	3,099	1,382	1,464
Tranquilizers	11,047	10,254	1,472	1,475	675	599
Stimulants	12,447	11,535	958	857	500	387
Methamphetamine	7,643	7,353	551	493	246	225
Sedatives	6,995	6,477	345	332	128	98
Any Illicit Drug Other Than Marijuana ¹	39,545	39,130	6,285	6,125	3,208	2,926

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotropic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.4B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 26 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	39.2	38.5	7.4	7.1	4.1	4.2
Marijuana and Hashish	34.7	34.4	5.2	5.0	2.8	3.0
Cocaine	12.7	12.4	1.1	1.0	0.6	0.4
Crack	2.9	2.5	0.4	0.3	0.2	0.1
Heroin	1.4	1.3	0.1	0.1	0.0	0.1
Hallucinogens	10.8	11.2	0.2 ^a	0.4	0.1	0.1
LSD	8.4	8.6	0.1	0.1	0.0	0.0
PCP	2.8	2.9	0.0	0.0	*	0.0
Inhalants	6.5	6.4	0.2	0.2	0.1	0.1
Nonmedical Use of Any Psychotherapeutic ²	15.1	14.2	2.8	2.6	1.3	1.2
Pain Relievers	8.1	7.6	1.9	1.8	0.8	0.9
Tranquilizers	6.5	6.0	0.9	0.9	0.4	0.4
Stimulants	7.3	6.7	0.6	0.5	0.3	0.2
Methamphetamine	4.5	4.3	0.3	0.3	0.1	0.1
Sedatives	4.1	3.8	0.2	0.2	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	23.3	22.9	3.7	3.6	1.9	1.7

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.5A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 12 or 13: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	1,008	976	656 ^a	562	299 ^a	235
Marijuana and Hashish	306	307	242	207	110	88
Cocaine	36	30	32	24	13	11
Crack	5	8	4	8	3	2
Heroin	7	7	5	6	2	1
Hallucinogens	107	98	61	65	22	17
LSD	44	41	27	31	7	5
PCP	20	24	11	13	3	1
Inhalants	515	527	265	216	95 ^b	54
Nonmedical Use of Any Psychotherapeutic ²	436	438	297	268	134	124
Pain Relievers	320	338	217	195	103	97
Tranquilizers	62	65	41	37	14	11
Stimulants	110	130	65	83	24	27
Methamphetamine	21	24	13	19	8	9
Sedatives	40	37	28	25	11	9
Any Illicit Drug Other Than Marijuana ¹	854	837	522	456	222	180

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.5B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 or 13: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	13.3	12.6	8.6 ^b	7.2	3.9 ^b	3.0
Marijuana and Hashish	4.0	4.0	3.2	2.7	1.5	1.1
Cocaine	0.5	0.4	0.4	0.3	0.2	0.1
Crack	0.1	0.1	0.0	0.1	0.0	0.0
Heroin	0.1	0.1	0.1	0.1	0.0	0.0
Hallucinogens	1.4	1.3	0.8	0.8	0.3	0.2
LSD	0.6	0.5	0.4	0.4	0.1	0.1
PCP	0.3	0.3	0.1	0.2	0.0	0.0
Inhalants	6.8	6.8	3.5 ^a	2.8	1.3 ^b	0.7
Nonmedical Use of Any Psychotherapeutic ²	5.7	5.6	3.9	3.5	1.8	1.6
Pain Relievers	4.2	4.4	2.9	2.5	1.4	1.3
Tranquilizers	0.8	0.8	0.5	0.5	0.2	0.1
Stimulants	1.4	1.7	0.9	1.1	0.3	0.3
Methamphetamine	0.3	0.3	0.2	0.2	0.1	0.1
Sedatives	0.5	0.5	0.4	0.3	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	11.3	10.8	6.9 ^a	5.9	2.9	2.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.6A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 14 or 15: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	2,218	2,192	1,569	1,545	770	788
Marijuana and Hashish	1,383	1,407	1,062	1,074	546	556
Cocaine	143	149	101	125	38	38
Crack	51	42	39	36	9	9
Heroin	18	29	16	19	14	3
Hallucinogens	357	396	261	294	70	76
LSD	227	235	151	171	42	29
PCP	75	86	39	49	12	8
Inhalants	782	795	332	330	82	95
Nonmedical Use of Any Psychotherapeutic ²	903	883	635	592	265	244
Pain Relievers	686	686	487	445	195	192
Tranquilizers	196	183	136	120	50	35
Stimulants	304	324	188	203	61	63
Methamphetamine	98	96	66	55	19	14
Sedatives	65	60	49	38	19	11
Any Illicit Drug Other Than Marijuana ¹	1,500	1,494	952	947	385	374

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.6B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 14 or 15: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	28.2	27.2	19.9	19.2	9.8	9.8
Marijuana and Hashish	17.6	17.5	13.5	13.3	6.9	6.9
Cocaine	1.8	1.9	1.3	1.6	0.5	0.5
Crack	0.6	0.5	0.5	0.5	0.1	0.1
Heroin	0.2	0.4	0.2	0.2	0.2	0.0
Hallucinogens	4.5	4.9	3.3	3.7	0.9	0.9
LSD	2.9	2.9	1.9	2.1	0.5	0.4
PCP	1.0	1.1	0.5	0.6	0.2	0.1
Inhalants	9.9	9.9	4.2	4.1	1.0	1.2
Nonmedical Use of Any Psychotherapeutic ²	11.5	11.0	8.1	7.4	3.4	3.0
Pain Relievers	8.7	8.5	6.2	5.5	2.5	2.4
Tranquilizers	2.5	2.3	1.7	1.5	0.6	0.4
Stimulants	3.9	4.0	2.4	2.5	0.8	0.8
Methamphetamine	1.3	1.2	0.8	0.7	0.2	0.2
Sedatives	0.8	0.7	0.6	0.5	0.2	0.1
Any Illicit Drug Other Than Marijuana ¹	19.1	18.6	12.1	11.8	4.9	4.6

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.7A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 16 or 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	3,189	3,121	2,359	2,250	1,197	1,242
Marijuana and Hashish	2,656	2,569	1,980	1,855	1,020	1,034
Cocaine	372	371	240	239	73	83
Crack	86	92	48	43	7	8
Heroin	77 ^a	48	45	25	23	11
Hallucinogens	852	854	567	549	163	177
LSD	606	565	376 ^a	309	101	86
PCP	123	135	49	50	11	10
Inhalants	821	756	311	279	67	75
Nonmedical Use of Any Psychotherapeutic ²	1,185	1,222	720	790	266	323
Pain Relievers	887	940	562	619	199	248
Tranquilizers	316	348	185	216	55	64
Stimulants	502	472	240	266	65	91
Methamphetamine	197	185	84	112	25 ^a	45
Sedatives	85	91	34	46	14	17
Any Illicit Drug Other Than Marijuana ¹	1,894	1,897	1,224	1,229	437 ^a	520

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.7B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 16 or 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	41.1	41.3	30.4	29.8	15.4	16.4
Marijuana and Hashish	34.3	34.0	25.5	24.5	13.2	13.7
Cocaine	4.8	4.9	3.1	3.2	0.9	1.1
Crack	1.1	1.2	0.6	0.6	0.1	0.1
Heroin	1.0 ^a	0.6	0.6	0.3	0.3	0.1
Hallucinogens	11.0	11.3	7.3	7.3	2.1	2.3
LSD	7.8	7.5	4.8	4.1	1.3	1.1
PCP	1.6	1.8	0.6	0.7	0.1	0.1
Inhalants	10.6	10.0	4.0	3.7	0.9	1.0
Nonmedical Use of Any Psychotherapeutic ²	15.3	16.2	9.3 ^a	10.4	3.4 ^a	4.3
Pain Relievers	11.4	12.4	7.2	8.2	2.6 ^a	3.3
Tranquilizers	4.1	4.6	2.4	2.9	0.7	0.8
Stimulants	6.5	6.2	3.1	3.5	0.8 ^a	1.2
Methamphetamine	2.5	2.5	1.1	1.5	0.3 ^a	0.6
Sedatives	1.1	1.2	0.4	0.6	0.2	0.2
Any Illicit Drug Other Than Marijuana ¹	24.4	25.1	15.8	16.3	5.6 ^a	6.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.8A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 18 to 20: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	6,255	6,228	4,044	4,020	2,349	2,372
Marijuana and Hashish	5,541	5,543	3,481	3,470	2,080	2,080
Cocaine	1,271	1,182	687 ^a	578	216	209
Crack	360	324	127	103	42	31
Heroin	186	157	70	47	19	12
Hallucinogens	2,166	2,296	1,099	1,154	309	322
LSD	1,637	1,664	704	641	184	163
PCP	293	306	80	68	17	14
Inhalants	1,605	1,569	444	456	111	101
Nonmedical Use of Any Psychotherapeutic ²	2,494	2,446	1,348	1,362	536	542
Pain Relievers	1,827	1,905	1,039	1,084	383	397
Tranquilizers	894	853	423	410	139	144
Stimulants	1,112 ^a	965	434	390	160	136
Methamphetamine	626 ^b	463	220	193	71	49
Sedatives	237	203	88	103	28	20
Any Illicit Drug Other Than Marijuana ¹	3,850	3,874	2,151	2,191	887	912

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.8B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 to 20: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	51.9	51.5	33.6	33.3	19.5	19.6
Marijuana and Hashish	46.0	45.9	28.9	28.7	17.3	17.2
Cocaine	10.6	9.8	5.7 ^a	4.8	1.8	1.7
Crack	3.0	2.7	1.1	0.9	0.3	0.3
Heroin	1.5	1.3	0.6	0.4	0.2	0.1
Hallucinogens	18.0	19.0	9.1	9.6	2.6	2.7
LSD	13.6	13.8	5.8	5.3	1.5	1.4
PCP	2.4	2.5	0.7	0.6	0.1	0.1
Inhalants	13.3	13.0	3.7	3.8	0.9	0.8
Nonmedical Use of Any Psychotherapeutic ²	20.7	20.2	11.2	11.3	4.5	4.5
Pain Relievers	15.2	15.8	8.6	9.0	3.2	3.3
Tranquilizers	7.4	7.1	3.5	3.4	1.2	1.2
Stimulants	9.2 ^a	8.0	3.6	3.2	1.3	1.1
Methamphetamine	5.2 ^b	3.8	1.8	1.6	0.6	0.4
Sedatives	2.0	1.7	0.7	0.8	0.2	0.2
Any Illicit Drug Other Than Marijuana ¹	32.0	32.1	17.9	18.1	7.4	7.5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.9A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 21 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	8,720	8,618	4,243	4,068	2,310	2,227
Marijuana and Hashish	7,786	7,713	3,482	3,390	1,969	1,870
Cocaine	2,112	1,967	805	697	260 ^a	186
Crack	576	498	144 ^a	88	32 ^a	12
Heroin	312	246	58	61	17	14
Hallucinogens	3,324	3,296	829	805	225	210
LSD	2,548	2,408	428	357	93	72
PCP	398	364	44	31	12	9
Inhalants	2,409 ^a	2,132	303	240	71	61
Nonmedical Use of Any Psychotherapeutic ²	3,462	3,204	1,482	1,334	526	508
Pain Relievers	2,489	2,317	1,120	1,023	360	384
Tranquilizers	1,346	1,305	471	473	164	148
Stimulants	1,448 ^a	1,235	407 ^a	313	139 ^b	85
Methamphetamine	857 ^a	722	205	160	65	45
Sedatives	325	274	86	67	29	21
Any Illicit Drug Other Than Marijuana ¹	5,626	5,374	2,309	2,104	842	799

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.9B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 21 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	53.1 ^a	51.0	25.8 ^a	24.1	14.1	13.2
Marijuana and Hashish	47.4	45.6	21.2	20.1	12.0	11.1
Cocaine	12.9 ^a	11.6	4.9 ^a	4.1	1.6 ^a	1.1
Crack	3.5	2.9	0.9 ^b	0.5	0.2 ^a	0.1
Heroin	1.9	1.5	0.4	0.4	0.1	0.1
Hallucinogens	20.2	19.5	5.0	4.8	1.4	1.2
LSD	15.5	14.2	2.6	2.1	0.6	0.4
PCP	2.4	2.2	0.3	0.2	0.1	0.1
Inhalants	14.7 ^b	12.6	1.8	1.4	0.4	0.4
Nonmedical Use of Any Psychotherapeutic ²	21.1 ^b	19.0	9.0 ^a	7.9	3.2	3.0
Pain Relievers	15.1 ^a	13.7	6.8	6.1	2.2	2.3
Tranquilizers	8.2	7.7	2.9	2.8	1.0	0.9
Stimulants	8.8 ^b	7.3	2.5 ^a	1.9	0.8 ^b	0.5
Methamphetamine	5.2 ^a	4.3	1.2	0.9	0.4	0.3
Sedatives	2.0	1.6	0.5	0.4	0.2	0.1
Any Illicit Drug Other Than Marijuana ¹	34.2 ^b	31.8	14.1 ^b	12.4	5.1	4.7

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.10A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 26 to 34: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	17,896 ^a	16,791	4,551	4,435	2,292	2,570
Marijuana and Hashish	16,049	15,170	3,465	3,393	1,802	1,944
Cocaine	6,008 ^b	4,992	824	677	401 ^a	263
Crack	1,726 ^b	1,260	234 ^a	124	129 ^b	42
Heroin	431	361	37	15	29	4
Hallucinogens	5,528	5,200	214 ^b	391	44	126
LSD	4,184	3,908	109	130	16	44
PCP	742	590	*	22	*	4
Inhalants	3,826	3,642	101	163	64	80
Nonmedical Use of Any Psychotherapeutic ²	6,254 ^a	5,572	1,540	1,459	494 ^a	704
Pain Relievers	3,714	3,441	1,043	1,047	310 ^b	516
Tranquilizers	2,769 ^b	2,271	489	444	180	155
Stimulants	2,764 ^b	2,234	283	278	93	142
Methamphetamine	1,823	1,600	160	177	57	80
Sedatives	835	691	60	85	29	21
Any Illicit Drug Other Than Marijuana ¹	11,033 ^b	10,038	2,206	2,169	899	1,018

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.10B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 26 to 34: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	53.2 ^a	50.9	13.5	13.4	6.8	7.8
Marijuana and Hashish	47.7	46.0	10.3	10.3	5.4	5.9
Cocaine	17.8 ^b	15.1	2.4	2.1	1.2	0.8
Crack	5.1 ^b	3.8	0.7 ^a	0.4	0.4 ^b	0.1
Heroin	1.3	1.1	0.1	0.0	0.1	0.0
Hallucinogens	16.4	15.8	0.6 ^b	1.2	0.1 ^a	0.4
LSD	12.4	11.8	0.3	0.4	0.0	0.1
PCP	2.2	1.8	*	0.1	*	0.0
Inhalants	11.4	11.0	0.3	0.5	0.2	0.2
Nonmedical Use of Any Psychotherapeutic ²	18.6 ^a	16.9	4.6	4.4	1.5 ^b	2.1
Pain Relievers	11.0	10.4	3.1	3.2	0.9 ^b	1.6
Tranquilizers	8.2 ^b	6.9	1.5	1.3	0.5	0.5
Stimulants	8.2 ^b	6.8	0.8	0.8	0.3	0.4
Methamphetamine	5.4	4.8	0.5	0.5	0.2	0.2
Sedatives	2.5	2.1	0.2	0.3	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	32.8 ^b	30.4	6.6	6.6	2.7	3.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.11A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 35 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	48,448	49,006	7,980	7,654	4,613	4,594
Marijuana and Hashish	42,708	43,612	5,390	5,200	2,932	3,141
Cocaine	15,463	16,206	1,052	989	551	424
Crack	3,105	3,081	449	319	197	159
Heroin	2,023	1,931	123	136	51	84
Hallucinogens	12,726	13,986	160	225	90	43
LSD	9,969	10,822	110	109	40	*
PCP	4,043	4,298	23	31	*	9
Inhalants	7,180	7,280	182	233	80	156
Nonmedical Use of Any Psychotherapeutic ²	19,342	18,678	3,198	2,956	1,731	1,404
Pain Relievers	9,964	9,584	2,113	2,051	1,072	949
Tranquilizers	8,278	7,983	983	1,031	495	444
Stimulants	9,683	9,301	675	579	407	245
Methamphetamine	5,820	5,753	391	316	190	145
Sedatives	6,160	5,786	285	247	99	77
Any Illicit Drug Other Than Marijuana ¹	28,512	29,092	4,078	3,956	2,309	1,908

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.11B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 35 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	35.7	35.5	5.9	5.5	3.4	3.3
Marijuana and Hashish	31.5	31.6	4.0	3.8	2.2	2.3
Cocaine	11.4	11.8	0.8	0.7	0.4	0.3
Crack	2.3	2.2	0.3	0.2	0.1	0.1
Heroin	1.5	1.4	0.1	0.1	0.0	0.1
Hallucinogens	9.4	10.1	0.1	0.2	0.1	0.0
LSD	7.3	7.8	0.1	0.1	0.0	*
PCP	3.0	3.1	0.0	0.0	*	0.0
Inhalants	5.3	5.3	0.1	0.2	0.1	0.1
Nonmedical Use of Any Psychotherapeutic ²	14.2	13.5	2.4	2.1	1.3	1.0
Pain Relievers	7.3	6.9	1.6	1.5	0.8	0.7
Tranquilizers	6.1	5.8	0.7	0.7	0.4	0.3
Stimulants	7.1	6.7	0.5	0.4	0.3	0.2
Methamphetamine	4.3	4.2	0.3	0.2	0.1	0.1
Sedatives	4.5	4.2	0.2	0.2	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	21.0	21.1	3.0	2.9	1.7	1.4

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.12A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Males: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	46,359	46,703	14,687	13,880	8,593	8,272
Marijuana and Hashish	41,129	41,589	11,817	11,154	6,921	6,624
Cocaine	14,890	14,690	2,255	2,037	973	760
Crack	3,936 ^a	3,236	663	449	279	188
Heroin	2,119	1,724	169	175	94	64
Hallucinogens	14,753	15,506	1,941	2,138	566	584
LSD	11,514	11,790	1,198	1,149	291	282
PCP	3,579	3,512	160	166	34	32
Inhalants	11,223	10,480	1,235	1,216	386	410
Nonmedical Use of Any Psychotherapeutic ²	17,764	17,008	4,539	4,217	2,034	1,926
Pain Relievers	10,709	10,218	3,298	3,280	1,370	1,443
Tranquilizers	7,354	6,623	1,325	1,176	546	458
Stimulants	8,860	7,996	1,131	1,057	442	419
Methamphetamine	5,815	5,135	651	564	236	193
Sedatives	4,497	3,955	319	310	141	111
Any Illicit Drug Other Than Marijuana ¹	28,729	29,021	7,240	6,926	3,325	3,112

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.12B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	43.6	43.5	13.8	12.9	8.1	7.7
Marijuana and Hashish	38.7	38.7	11.1	10.4	6.5	6.2
Cocaine	14.0	13.7	2.1	1.9	0.9	0.7
Crack	3.7 ^a	3.0	0.6	0.4	0.3	0.2
Heroin	2.0	1.6	0.2	0.2	0.1	0.1
Hallucinogens	13.9	14.4	1.8	2.0	0.5	0.5
LSD	10.8	11.0	1.1	1.1	0.3	0.3
PCP	3.4	3.3	0.2	0.2	0.0	0.0
Inhalants	10.6	9.8	1.2	1.1	0.4	0.4
Nonmedical Use of Any Psychotherapeutic ²	16.7	15.8	4.3	3.9	1.9	1.8
Pain Relievers	10.1	9.5	3.1	3.1	1.3	1.3
Tranquilizers	6.9	6.2	1.2	1.1	0.5	0.4
Stimulants	8.3 ^a	7.4	1.1	1.0	0.4	0.4
Methamphetamine	5.5	4.8	0.6	0.5	0.2	0.2
Sedatives	4.2	3.7	0.3	0.3	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	27.0	27.0	6.8	6.5	3.1	2.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.13A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Females: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	41,375	40,228	10,715	10,654	5,236	5,755
Marijuana and Hashish	35,299	34,732	7,285	7,435	3,538 ^a	4,089
Cocaine	10,516	10,206	1,486	1,291	579	454
Crack	1,974	2,071	382	272	138	76
Heroin	935	1,055	184	133	60	65
Hallucinogens	10,308	10,619	1,250	1,345	356	387
LSD	7,701	7,852	708 ^a	600	191 ^a	121
PCP	2,114	2,292	85	98	21	22
Inhalants	5,915	6,222	702	701	184	212
Nonmedical Use of Any Psychotherapeutic ²	16,312	15,435	4,682	4,544	1,918	1,923
Pain Relievers	9,178	8,993	3,284	3,186	1,251	1,339
Tranquilizers	6,506	6,384	1,403	1,555	551	542
Stimulants	7,062	6,666	1,161	1,055	508	369
Methamphetamine	3,628	3,707	489	467	199	194
Sedatives	3,249	3,187	312	301	88	64
Any Illicit Drug Other Than Marijuana ¹	24,538	23,584	6,204	6,126	2,656	2,599

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.13B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	36.0	34.7	9.3	9.2	4.6	5.0
Marijuana and Hashish	30.7	30.0	6.3	6.4	3.1 ^a	3.5
Cocaine	9.2	8.8	1.3	1.1	0.5	0.4
Crack	1.7	1.8	0.3	0.2	0.1	0.1
Heroin	0.8	0.9	0.2	0.1	0.1	0.1
Hallucinogens	9.0	9.2	1.1	1.2	0.3	0.3
LSD	6.7	6.8	0.6 ^a	0.5	0.2 ^a	0.1
PCP	1.8	2.0	0.1	0.1	0.0	0.0
Inhalants	5.1	5.4	0.6	0.6	0.2	0.2
Nonmedical Use of Any Psychotherapeutic ²	14.2	13.3	4.1	3.9	1.7	1.7
Pain Relievers	8.0	7.8	2.9	2.7	1.1	1.2
Tranquilizers	5.7	5.5	1.2	1.3	0.5	0.5
Stimulants	6.1	5.7	1.0	0.9	0.4	0.3
Methamphetamine	3.2	3.2	0.4	0.4	0.2	0.2
Sedatives	2.8	2.7	0.3	0.3	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	21.4	20.3	5.4	5.3	2.3	2.2

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.14A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Males Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	3,364	3,250	2,365	2,206	1,195	1,178
Marijuana and Hashish	2,339	2,237	1,773 ^a	1,629	921	925
Cocaine	247	263	159	186	58	70
Crack	63	62	38	37	6	11
Heroin	54	38	34	21	20	7
Hallucinogens	680	712	457	477	114	145
LSD	452	472	295	298	66	79
PCP	116	133	55	68	11	12
Inhalants	1,111	1,100	456	430	118	109
Nonmedical Use of Any Psychotherapeutic ²	1,196	1,238	733	778	308	317
Pain Relievers	890	984	596	623	243	254
Tranquilizers	257	265	150	150	53	46
Stimulants	402	396	185	226	48 ^a	74
Methamphetamine	138	142	63	88	16	30
Sedatives	94	76	47	45	19	17
Any Illicit Drug Other Than Marijuana ¹	2,142	2,158	1,299	1,278	485	509

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.14B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	28.3	27.2	19.9 ^a	18.4	10.1	9.8
Marijuana and Hashish	19.7	18.7	14.9 ^a	13.6	7.8	7.7
Cocaine	2.1	2.2	1.3	1.6	0.5	0.6
Crack	0.5	0.5	0.3	0.3	0.0	0.1
Heroin	0.5	0.3	0.3	0.2	0.2	0.1
Hallucinogens	5.7	6.0	3.9	4.0	1.0	1.2
LSD	3.8	3.9	2.5	2.5	0.6	0.7
PCP	1.0	1.1	0.5	0.6	0.1	0.1
Inhalants	9.4	9.2	3.8	3.6	1.0	0.9
Nonmedical Use of Any Psychotherapeutic ²	10.1	10.4	6.2	6.5	2.6	2.7
Pain Relievers	7.5	8.2	5.0	5.2	2.0	2.1
Tranquilizers	2.2	2.2	1.3	1.3	0.4	0.4
Stimulants	3.4	3.3	1.6	1.9	0.4 ^a	0.6
Methamphetamine	1.2	1.2	0.5	0.7	0.1	0.3
Sedatives	0.8	0.6	0.4	0.4	0.2	0.1
Any Illicit Drug Other Than Marijuana ¹	18.0	18.0	10.9	10.7	4.1	4.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.15A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Females Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	3,051	3,038	2,219	2,151	1,070	1,086
Marijuana and Hashish	2,006	2,046	1,511	1,507	755	753
Cocaine	304	287	214	202	66	62
Crack	80	81	52	50	13	9
Heroin	48	46	31	28	19	8
Hallucinogens	637	636	432	431	140	125
LSD	425	369	258	213	83 ^b	40
PCP	101	113	43	44	14	6
Inhalants	1,007	978	452	395	127	114
Nonmedical Use of Any Psychotherapeutic ²	1,328	1,305	920	871	356	373
Pain Relievers	1,004	980	670	637	254	284
Tranquilizers	318	330	212	223	66	64
Stimulants	513	530	307	326	103	107
Methamphetamine	179	162	101	98	37	38
Sedatives	96	111	64	65	25	19
Any Illicit Drug Other Than Marijuana ¹	2,106	2,070	1,399	1,353	559	565

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.15B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	26.9	26.6	19.6	18.9	9.4	9.5
Marijuana and Hashish	17.7	17.9	13.3	13.2	6.7	6.6
Cocaine	2.7	2.5	1.9	1.8	0.6	0.5
Crack	0.7	0.7	0.5	0.4	0.1	0.1
Heroin	0.4	0.4	0.3	0.2	0.2	0.1
Hallucinogens	5.6	5.6	3.8	3.8	1.2	1.1
LSD	3.8	3.2	2.3	1.9	0.7 ^b	0.4
PCP	0.9	1.0	0.4	0.4	0.1	0.1
Inhalants	8.9	8.6	4.0	3.5	1.1	1.0
Nonmedical Use of Any Psychotherapeutic ²	11.7	11.4	8.1	7.6	3.1	3.3
Pain Relievers	8.9	8.6	5.9	5.6	2.2	2.5
Tranquilizers	2.8	2.9	1.9	2.0	0.6	0.6
Stimulants	4.5	4.7	2.7	2.9	0.9	0.9
Methamphetamine	1.6	1.4	0.9	0.9	0.3	0.3
Sedatives	0.8	1.0	0.6	0.6	0.2	0.2
Any Illicit Drug Other Than Marijuana ¹	18.6	18.1	12.4	11.9	4.9	4.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.16A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Males Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	7,954	7,888	4,736	4,599	2,869	2,745
Marijuana and Hashish	7,163	7,076	4,108	3,980	2,538	2,408
Cocaine	2,046	1,870	949 ^a	780	298	238
Crack	595	505	184	133	56 ^a	27
Heroin	312	247	82	67	24	21
Hallucinogens	3,148	3,238	1,204	1,189	336	332
LSD	2,435	2,383	746	640	184	162
PCP	408	406	84	58	22	16
Inhalants	2,482	2,267	513	444	130	103
Nonmedical Use of Any Psychotherapeutic ²	3,186	3,095	1,517	1,494	643	608
Pain Relievers	2,363	2,386	1,154	1,208	454	467
Tranquilizers	1,248	1,222	500	506	175	167
Stimulants	1,410 ^b	1,180	487 ^a	390	186 ^a	132
Methamphetamine	840 ^b	650	270 ^a	192	92 ^a	50
Sedatives	316	291	97	110	35	31
Any Illicit Drug Other Than Marijuana ¹	5,199	5,128	2,555	2,455	1,064	999

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.16B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	56.2	54.7	33.5	31.9	20.3	19.0
Marijuana and Hashish	50.7	49.1	29.1	27.6	17.9	16.7
Cocaine	14.5 ^a	13.0	6.7 ^b	5.4	2.1	1.7
Crack	4.2 ^a	3.5	1.3 ^a	0.9	0.4 ^a	0.2
Heroin	2.2	1.7	0.6	0.5	0.2	0.1
Hallucinogens	22.3	22.5	8.5	8.2	2.4	2.3
LSD	17.2	16.5	5.3 ^a	4.4	1.3	1.1
PCP	2.9	2.8	0.6	0.4	0.2	0.1
Inhalants	17.6 ^b	15.7	3.6	3.1	0.9	0.7
Nonmedical Use of Any Psychotherapeutic ²	22.5	21.5	10.7	10.4	4.5	4.2
Pain Relievers	16.7	16.6	8.2	8.4	3.2	3.2
Tranquilizers	8.8	8.5	3.5	3.5	1.2	1.2
Stimulants	10.0 ^b	8.2	3.4 ^a	2.7	1.3 ^a	0.9
Methamphetamine	5.9 ^b	4.5	1.9 ^a	1.3	0.6 ^a	0.3
Sedatives	2.2	2.0	0.7	0.8	0.2	0.2
Any Illicit Drug Other Than Marijuana ¹	36.8	35.6	18.1	17.0	7.5	6.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.17A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Females Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	7,021	6,958	3,551	3,489	1,789	1,854
Marijuana and Hashish	6,163	6,180	2,855	2,880	1,511	1,542
Cocaine	1,337	1,279	544	494	178	156
Crack	341	317	87	57	17	16
Heroin	187	156	46	40	12	5
Hallucinogens	2,342	2,355	725	770	198	200
LSD	1,751	1,689	386	358	92	72
PCP	283	265	41	41	7	7
Inhalants	1,532	1,435	234	251	52	59
Nonmedical Use of Any Psychotherapeutic ²	2,770 ^a	2,555	1,313	1,202	419	442
Pain Relievers	1,953	1,836	1,006	900	289	313
Tranquilizers	991	937	394	377	128	124
Stimulants	1,150	1,020	354	312	113	89
Methamphetamine	643 ^a	535	155	160	45	45
Sedatives	245	186	77	59	22	10
Any Illicit Drug Other Than Marijuana ¹	4,277	4,120	1,906	1,841	665	712

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.17B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	49.0	47.8	24.8	23.9	12.5	12.7
Marijuana and Hashish	43.0	42.4	19.9	19.8	10.5	10.6
Cocaine	9.3	8.8	3.8	3.4	1.2	1.1
Crack	2.4	2.2	0.6	0.4	0.1	0.1
Heroin	1.3	1.1	0.3	0.3	0.1	0.0
Hallucinogens	16.3	16.2	5.1	5.3	1.4	1.4
LSD	12.2	11.6	2.7	2.5	0.6	0.5
PCP	2.0	1.8	0.3	0.3	0.0	0.0
Inhalants	10.7	9.8	1.6	1.7	0.4	0.4
Nonmedical Use of Any Psychotherapeutic ²	19.3 ^b	17.5	9.2	8.2	2.9	3.0
Pain Relievers	13.6	12.6	7.0 ^a	6.2	2.0	2.1
Tranquilizers	6.9	6.4	2.7	2.6	0.9	0.9
Stimulants	8.0 ^a	7.0	2.5	2.1	0.8	0.6
Methamphetamine	4.5 ^a	3.7	1.1	1.1	0.3	0.3
Sedatives	1.7 ^a	1.3	0.5	0.4	0.2	0.1
Any Illicit Drug Other Than Marijuana ¹	29.9 ^a	28.3	13.3	12.6	4.6	4.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.18A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 18 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	81,319	80,643	20,818	20,177	11,564	11,763
Marijuana and Hashish	72,083	72,038	15,818	15,453	8,783	9,035
Cocaine	24,854	24,346	3,369	2,940	1,428	1,081
Crack	5,767	5,164	955 ^a	634	399	245
Heroin	2,953	2,695	288	259	116	114
Hallucinogens	23,744	24,778	2,302	2,575	668	701
LSD	18,338	18,802	1,352	1,237	332	283
PCP	5,476	5,558	147	152	29	35
Inhalants	15,020	14,623	1,029	1,092	326	398
Nonmedical Use of Any Psychotherapeutic ²	31,552	29,900	7,567	7,112	3,287	3,158
Pain Relievers	17,994	17,246	5,316	5,206	2,125	2,245
Tranquilizers	13,286	12,412	2,366	2,358	978	890
Stimulants	15,007 ^a	13,735	1,799	1,560	799	607
Methamphetamine	9,126	8,538	976	845	383	319
Sedatives	7,556	6,955	519	501	185	139
Any Illicit Drug Other Than Marijuana ¹	49,020	48,377	10,745	10,421	4,937	4,637

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.18B Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	41.1	40.3	10.5	10.1	5.8	5.9
Marijuana and Hashish	36.4	36.0	8.0	7.7	4.4	4.5
Cocaine	12.6	12.2	1.7	1.5	0.7 ^a	0.5
Crack	2.9	2.6	0.5 ^a	0.3	0.2	0.1
Heroin	1.5	1.3	0.1	0.1	0.1	0.1
Hallucinogens	12.0	12.4	1.2	1.3	0.3	0.4
LSD	9.3	9.4	0.7	0.6	0.2	0.1
PCP	2.8	2.8	0.1	0.1	0.0	0.0
Inhalants	7.6	7.3	0.5	0.5	0.2	0.2
Nonmedical Use of Any Psychotherapeutic ²	15.9 ^a	15.0	3.8	3.6	1.7	1.6
Pain Relievers	9.1	8.6	2.7	2.6	1.1	1.1
Tranquilizers	6.7	6.2	1.2	1.2	0.5	0.4
Stimulants	7.6 ^a	6.9	0.9	0.8	0.4	0.3
Methamphetamine	4.6	4.3	0.5	0.4	0.2	0.2
Sedatives	3.8	3.5	0.3	0.3	0.1	0.1
Any Illicit Drug Other Than Marijuana ¹	24.8	24.2	5.4	5.2	2.5	2.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.19A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	87,734	86,931	25,402	24,535	13,829	14,027
12	369	339	247	202	116	95
13	639	637	409	360	183 ^a	140
14	1,018 ^b	881	683 ^a	596	334	316
15	1,200	1,311	885	949	436	472
16	1,565	1,543	1,164	1,121	592	591
17	1,624	1,577	1,195	1,129	604	651
18	2,084	2,100	1,439	1,452	820	876
19	2,109	2,030	1,343	1,319	782	765
20	2,061	2,098	1,262	1,248	747	731
21	1,941	1,909	1,073	1,053	604	625
22	1,797	1,776	967	902	545	473
23	1,741	1,740	872	834	473	435
24	1,718	1,608	717	700	369	382
25	1,523	1,585	614	581	318	313
26-29	7,527	7,115	2,252	2,218	1,169	1,248
30-34	10,369 ^a	9,676	2,298	2,217	1,123	1,322
35-39	13,521 ^a	11,992	2,687 ^a	2,147	1,414	1,141
40-44	12,475	12,889	2,651	2,307	1,697	1,451
45-49	9,758	10,040	1,340	1,386	688	971
50-54	5,945	6,761	624	848	425	463
55-59	3,026 ^a	3,854	265 ^a	563	177	311
60-64	1,410	1,469	107	180	34	154
65 or Older	2,312	2,002	306	223	178	103

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.19B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	39.7	38.9	11.5	11.0	6.3	6.3
12	10.3	9.1	6.9 ^a	5.4	3.2	2.5
13	15.9	15.8	10.2	9.0	4.6 ^a	3.5
14	25.0 ^b	21.8	16.8 ^a	14.7	8.2	7.8
15	31.6	32.8	23.3	23.7	11.5	11.8
16	38.8	38.9	28.9	28.3	14.7	14.9
17	43.7	43.9	32.1	31.4	16.2	18.1
18	48.7	48.7	33.6	33.7	19.2	20.3
19	53.0	52.2	33.8	33.9	19.6	19.7
20	54.5	54.0	33.3	32.2	19.8	18.8
21	54.9	54.4	30.3	30.0	17.1	17.8
22	53.3	52.4	28.7	26.6	16.2	13.9
23	54.9 ^b	50.3	27.5 ^a	24.1	14.9	12.6
24	53.4	50.0	22.3	21.8	11.5	11.9
25	48.6	47.6	19.6	17.4	10.1	9.4
26-29	50.8	50.1	15.2	15.6	7.9	8.8
30-34	55.0 ^b	51.5	12.2	11.8	6.0	7.0
35-39	59.5 ^a	55.5	11.8	9.9	6.2	5.3
40-44	58.0	58.1	12.3	10.4	7.9	6.5
45-49	51.1	49.9	7.0	6.9	3.6	4.8
50-54	35.0	35.5	3.7	4.5	2.5	2.4
55-59	23.3	28.6	2.0 ^a	4.2	1.4	2.3
60-64	13.9	14.4	1.1	1.8	0.3	1.5
65 or Older	7.1	6.4	0.9	0.7	0.5	0.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.20A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	76,428	76,321	19,102	18,589	10,458	10,714
12	68	74	52	58	16	22
13	238	233	190	149	95	67
14	546	474	411	356	220	186
15	837 ^a	934	651	717	326	370
16	1,279	1,224	972	901	499	481
17	1,377	1,345	1,008	955	520	553
18	1,806	1,875	1,247	1,258	721	755
19	1,888	1,755	1,168	1,132	698	684
20	1,847	1,913	1,066	1,080	661	641
21	1,742	1,738	882	916	526	528
22	1,581	1,598	825	740	471	401
23	1,601	1,559	723	686	410	369
24	1,515	1,412	564	574	296	320
25	1,346	1,405	487	473	266	253
26-29	6,769	6,379	1,812	1,752	967	949
30-34	9,280	8,791	1,654	1,641	835	995
35-39	12,550 ^b	10,966	1,937	1,597	952	893
40-44	11,724	11,854	1,996	1,753	1,224	1,053
45-49	8,957	9,363	999	1,039	471	706
50-54	4,802 ^a	6,064	316	508	222	269
55-59	2,523	3,079	125	209	49	160
60-64	1,029	1,095	13	85	11	*
65 or Older	1,122	1,190	5	10	2	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.20B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	34.6	34.2	8.6	8.3	4.7	4.8
12	1.9	2.0	1.5	1.5	0.4	0.6
13	5.9	5.8	4.7	3.7	2.4	1.7
14	13.4	11.7	10.1	8.8	5.4	4.6
15	22.0	23.3	17.1	17.9	8.6	9.2
16	31.7	30.9	24.1	22.7	12.4	12.1
17	37.0	37.4	27.1	26.5	14.0	15.4
18	42.2	43.5	29.1	29.2	16.8	17.5
19	47.5	45.1	29.4	29.1	17.6	17.6
20	48.8	49.3	28.2	27.8	17.5	16.5
21	49.3	49.5	24.9	26.1	14.9	15.0
22	46.9	47.2	24.5	21.8	14.0	11.8
23	50.5 ^b	45.0	22.8 ^a	19.8	12.9	10.6
24	47.1	43.9	17.5	17.9	9.2	10.0
25	42.9	42.2	15.5	14.2	8.5	7.6
26-29	45.7	44.9	12.2	12.3	6.5	6.7
30-34	49.2 ^a	46.7	8.8	8.7	4.4	5.3
35-39	55.2 ^a	50.8	8.5	7.4	4.2	4.1
40-44	54.6	53.4	9.3	7.9	5.7	4.7
45-49	46.9	46.6	5.2	5.2	2.5	3.5
50-54	28.3	31.9	1.9	2.7	1.3	1.4
55-59	19.5	22.9	1.0	1.5	0.4	1.2
60-64	10.1	10.8	0.1	0.8	0.1	*
65 or Older	3.5	3.8	0.0	0.0	0.0	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.21A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	25,406	24,896	3,742	3,328	1,552	1,213
12	11	11	10	6	1	4
13	25	19	22	18	11	7
14	49	49	32	39	12	10
15	94	100	69	86	26	28
16	164	167	103	118	37	40
17	207	204	137	121	36	43
18	375	343	211	182	68	64
19	442	381	252 ^a	183	81	63
20	455	457	225	213	66	82
21	494	460	206	191	64	39
22	418	388	195	157	76	44
23	443	393	169	149	51	47
24	390	360	126	93	38	29
25	367	366	110	107	32	28
26-29	2,285 ^b	1,795	380	321	164	98
30-34	3,723 ^a	3,197	444	356	237	165
35-39	5,395	5,000	391	288	187	124
40-44	5,074	5,124	462	367	240	154
45-49	3,366	3,326	122	169	56	47
50-54	969 ^b	1,683	76	150	69	94
55-59	438	695	*	15	*	5
60-64	74	235	*	*	*	*
65 or Older	147	144	*	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.21B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	11.5	11.2	1.7	1.5	0.7	0.5
12	0.3	0.3	0.3	0.2	0.0	0.1
13	0.6	0.5	0.6	0.4	0.3	0.2
14	1.2	1.2	0.8	1.0	0.3	0.2
15	2.5	2.5	1.8	2.2	0.7	0.7
16	4.1	4.2	2.6	3.0	0.9	1.0
17	5.6	5.7	3.7	3.4	1.0	1.2
18	8.8	8.0	4.9	4.2	1.6	1.5
19	11.1	9.8	6.3 ^a	4.7	2.0	1.6
20	12.0	11.8	5.9	5.5	1.7	2.1
21	14.0	13.1	5.8	5.5	1.8	1.1
22	12.4	11.4	5.8	4.6	2.3	1.3
23	14.0 ^a	11.4	5.3	4.3	1.6	1.3
24	12.1	11.2	3.9	2.9	1.2	0.9
25	11.7	11.0	3.5	3.2	1.0	0.8
26-29	15.4 ^b	12.6	2.6	2.3	1.1	0.7
30-34	19.8 ^b	17.0	2.4	1.9	1.3	0.9
35-39	23.7	23.2	1.7	1.3	0.8	0.6
40-44	23.6	23.1	2.1	1.7	1.1	0.7
45-49	17.6	16.5	0.6	0.8	0.3	0.2
50-54	5.7 ^a	8.8	0.4	0.8	0.4	0.5
55-59	3.4	5.2	*	0.1	*	0.0
60-64	0.7	2.3	*	*	*	*
65 or Older	0.5	0.5	*	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.22A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	25,061	26,125	3,191	3,483	922	971
12	31	22	16	14	4	3
13	76	76	44	51	17	14
14	145	121	101	90	29	22
15	212 ^a	275	160 ^a	204	41	53
16	374	382	264	250	68	82
17	478	472	303	298	95	96
18	672	699	363	425	118	129
19	756	745	422	386	100	132
20	738	852	315	343	91	61
21	780	810	255	291	54	79
22	670	638	187	182	60	36
23	735	672	200	173	61	46
24	603	620	98	97	28	23
25	536	557	90	62	22	26
26-29	2,678	2,321	150	233	34	56
30-34	2,850	2,879	64 ^a	158	10 ^a	70
35-39	3,589	3,923	58	165	34	36
40-44	4,319	4,302	67	22	40	*
45-49	3,276	3,231	31	21	16	7
50-54	1,079 ^a	1,795	3	*	*	*
55-59	361	392	*	17	*	*
60-64	60	206	*	*	*	*
65 or Older	41	136	*	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.22B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	11.3	11.7	1.4	1.6	0.4	0.4
12	0.9	0.6	0.5	0.4	0.1	0.1
13	1.9	1.9	1.1	1.3	0.4	0.3
14	3.6	3.0	2.5	2.2	0.7	0.6
15	5.6 ^a	6.9	4.2	5.1	1.1	1.3
16	9.3	9.6	6.6	6.3	1.7	2.1
17	12.8	13.1	8.1	8.3	2.6	2.7
18	15.7	16.2	8.5	9.9	2.8	3.0
19	19.0	19.2	10.6	9.9	2.5	3.4
20	19.5	21.9	8.3	8.8	2.4	1.6
21	22.1	23.1	7.2	8.3	1.5	2.3
22	19.9	18.8	5.6	5.4	1.8	1.1
23	23.2 ^b	19.4	6.3	5.0	1.9	1.3
24	18.7	19.3	3.0	3.0	0.9	0.7
25	17.1	16.7	2.9	1.9	0.7	0.8
26-29	18.1	16.3	1.0 ^a	1.6	0.2	0.4
30-34	15.1	15.3	0.3 ^a	0.8	0.1 ^a	0.4
35-39	15.8	18.2	0.3 ^a	0.8	0.2	0.2
40-44	20.1	19.4	0.3	0.1	0.2	*
45-49	17.2	16.1	0.2	0.1	0.1	0.0
50-54	6.4 ^a	9.4	0.0	*	*	*
55-59	2.8	2.9	*	0.1	*	*
60-64	0.6	2.0	*	*	*	*
65 or Older	0.1	0.4	*	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.23A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	17,138	16,702	1,937	1,918	570	622
12	197	182	120	86	45	29
13	318	346	145	130	50 ^b	24
14	421	393	182	165	52	58
15	362	401	150	165	30	37
16	368	402	144	149	32	47
17	453 ^b	354	166	130	35	28
18	543	475	158	166	40	44
19	536	566	171	160	59	29
20	526	528	116	130	12	28
21	549	463	111	77	25	13
22	482	441	81 ^a	45	12	10
23	496	472	39	53	2 ^b	19
24	469	382	28	31	10	8
25	413	375	44	34	23	11
26-29	1,812	1,699	38	68	17	12
30-34	2,015	1,943	63	95	47	68
35-39	2,321	2,333	62	37	34	27
40-44	2,180	2,028	55	51	20	29
45-49	1,599	1,467	51	56	20	47
50-54	690	782	7	29	*	15
55-59	244	429	*	22	*	22
60-64	103	84	3	*	3	*
65 or Older	42	157	3	38	3	16

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.23B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	7.8	7.5	0.9	0.9	0.3	0.3
12	5.5	4.9	3.4 ^a	2.3	1.3	0.8
13	7.9	8.6	3.6	3.2	1.2 ^b	0.6
14	10.3	9.7	4.5	4.1	1.3	1.4
15	9.5	10.0	3.9	4.1	0.8	0.9
16	9.1	10.1	3.6	3.8	0.8	1.2
17	12.2 ^b	9.9	4.5	3.6	0.9	0.8
18	12.7	11.0	3.7	3.8	0.9	1.0
19	13.5	14.6	4.3	4.1	1.5	0.7
20	13.9	13.6	3.1	3.4	0.3	0.7
21	15.5	13.2	3.1	2.2	0.7	0.4
22	14.3	13.0	2.4 ^a	1.3	0.3	0.3
23	15.7	13.6	1.2	1.5	0.1 ^b	0.6
24	14.6 ^a	11.9	0.9	1.0	0.3	0.2
25	13.2	11.3	1.4	1.0	0.7	0.3
26-29	12.2	12.0	0.3	0.5	0.1	0.1
30-34	10.7	10.3	0.3	0.5	0.2	0.4
35-39	10.2	10.8	0.3	0.2	0.1	0.1
40-44	10.1	9.1	0.3	0.2	0.1	0.1
45-49	8.4	7.3	0.3	0.3	0.1	0.2
50-54	4.1	4.1	0.0	0.2	*	0.1
55-59	1.9	3.2	*	0.2	*	0.2
60-64	1.0	0.8	0.0	*	0.0	*
65 or Older	0.1	0.5	0.0	0.1	0.0	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.24A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	34,076	32,443	9,220	8,761	3,952	3,849
12	175	152	118	89	60	48
13	261	285	179	179	74	76
14	415	361	287	235	119	113
15	488	521	348	357	146	131
16	604	601	379	400	144	161
17	581	622	341	390	121	162
18	832	784	442	459	183	198
19	819	867	476	481	174	185
20	843	795	430	422	179	159
21	768	731	366	344	140	139
22	736	623	336	307	116	107
23	707	696	319	284	114	111
24	617	596	217	217	76	88
25	634	557	243	182	80	63
26-29	2,638	2,418	746	702	254	350
30-34	3,616 ^a	3,154	793	757	240 ^a	354
35-39	4,840	4,458	933	783	474	279
40-44	5,334	5,244	1,046	805	577	379
45-49	3,870	3,806	403	413	205	278
50-54	2,314	2,429	281	353	155	168
55-59	1,112	1,338	140	331	128	120
60-64	631	519	93	96	20	94
65 or Older	1,240	883	303	175	172	87

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.24B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	15.4 ^a	14.5	4.2	3.9	1.8	1.7
12	4.9	4.1	3.3 ^a	2.4	1.7	1.3
13	6.5	7.1	4.5	4.5	1.8	1.9
14	10.2	8.9	7.0	5.8	2.9	2.8
15	12.9	13.0	9.2	8.9	3.8	3.3
16	15.0	15.2	9.4	10.1	3.6	4.1
17	15.6	17.3	9.2 ^a	10.8	3.3 ^a	4.5
18	19.4	18.2	10.3	10.7	4.3	4.6
19	20.6	22.3	12.0	12.4	4.4	4.7
20	22.3	20.5	11.4	10.9	4.7	4.1
21	21.7	20.8	10.3	9.8	4.0	4.0
22	21.8 ^a	18.4	10.0	9.1	3.4	3.2
23	22.3	20.1	10.1	8.2	3.6	3.2
24	19.2	18.6	6.7	6.7	2.4	2.7
25	20.2 ^a	16.7	7.8 ^a	5.5	2.6	1.9
26-29	17.8	17.0	5.0	4.9	1.7	2.5
30-34	19.2 ^a	16.8	4.2	4.0	1.3 ^a	1.9
35-39	21.3	20.6	4.1	3.6	2.1	1.3
40-44	24.8	23.6	4.9	3.6	2.7	1.7
45-49	20.3	18.9	2.1	2.1	1.1	1.4
50-54	13.6	12.8	1.7	1.9	0.9	0.9
55-59	8.6	9.9	1.1	2.5	1.0	0.9
60-64	6.2	5.1	0.9	0.9	0.2	0.9
65 or Older	3.8	2.8	0.9	0.6	0.5	0.3

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.25A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	53,268	52,605	13,444	13,052	5,981	5,711
12	338	306	225 ^a	172	106	79
13	516	532	297	283	116	101
14	729	652	445	391	177	172
15	770	842	507	555	208	202
16	920	948	620	616	222	256
17	973	949	604	612	215	263
18	1,301	1,268	721	774	306	340
19	1,274	1,299	747	736	314	288
20	1,275	1,308	682	681	267	284
21	1,240	1,200	608	571	207	226
22	1,154	1,077	531	497	205	173
23	1,129	1,119	481	433	180	175
24	1,122	1,028	362	327	134	118
25	981	950	327	276	117	107
26-29	4,648	4,297	1,061	1,063	422	458
30-34	6,385 ^a	5,741	1,146	1,106	477	560
35-39	8,048	7,764	1,279	1,084	690 ^a	425
40-44	7,787	7,861	1,340	1,100	764	509
45-49	5,922	5,826	553	636	305	372
50-54	3,090	3,813	364	443	224	249
55-59	1,503	1,948	140 ^a	385	128	156
60-64	709	711	96	96	23	94
65 or Older	1,453	1,170	306	213	176	103

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.25B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	24.1	23.6	6.1	5.8	2.7	2.6
12	9.5	8.2	6.3 ^b	4.6	3.0	2.1
13	12.8	13.2	7.4	7.1	2.9	2.5
14	17.9	16.1	10.9	9.7	4.4	4.3
15	20.3	21.0	13.4	13.9	5.5	5.0
16	22.8	23.9	15.4	15.5	5.5	6.5
17	26.2	26.4	16.2	17.0	5.8 ^a	7.3
18	30.4	29.4	16.9	18.0	7.1	7.9
19	32.0	33.4	18.8	18.9	7.9	7.4
20	33.7	33.7	18.0	17.5	7.0	7.3
21	35.1	34.2	17.2	16.3	5.8	6.4
22	34.2	31.8	15.8	14.6	6.1	5.1
23	35.6	32.3	15.2 ^a	12.5	5.7	5.1
24	34.9	32.0	11.2	10.2	4.2	3.7
25	31.3	28.6	10.4	8.3	3.7	3.2
26-29	31.4	30.2	7.2	7.5	2.8	3.2
30-34	33.9 ^b	30.5	6.1	5.9	2.5	3.0
35-39	35.4	36.0	5.6	5.0	3.0	2.0
40-44	36.2	35.4	6.2	5.0	3.6	2.3
45-49	31.0	29.0	2.9	3.2	1.6	1.8
50-54	18.2	20.0	2.1	2.3	1.3	1.3
55-59	11.6	14.5	1.1 ^a	2.9	1.0	1.2
60-64	7.0	7.0	0.9	0.9	0.2	0.9
65 or Older	4.5	3.7	0.9	0.7	0.5	0.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.26A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	87,734	86,931	25,402	24,535	13,829	14,027
AGE						
12-17	6,415	6,288	4,584	4,357	2,265	2,264
18-25	14,975	14,846	8,287	8,088	4,658	4,599
26 or Older	66,344	65,797	12,531	12,089	6,906	7,164
GENDER						
Male	46,359	46,703	14,687	13,880	8,593	8,272
Female	41,375	40,228	10,715	10,654	5,236	5,755
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	68,047	67,567	18,538	18,284	9,974	10,488
Black Only	9,359	9,013	3,278 ^a	2,777	1,854	1,613
American Indian or Alaska Native Only	587	588	211	216	119	138
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	34
Asian Only	1,507	1,440	444	397	229	207
More Than One Race	822	915	302	384	201	275
Hispanic	7,182	7,142	2,538	2,412	1,405	1,272

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.26B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	39.7	38.9	11.5	11.0	6.3	6.3
AGE						
12-17	27.6	26.9	19.8 ^a	18.6	9.8	9.7
18-25	52.6	51.2	29.1	27.9	16.4	15.9
26 or Older	39.2	38.5	7.4	7.1	4.1	4.2
GENDER						
Male	43.6	43.5	13.8	12.9	8.1	7.7
Female	36.0	34.7	9.3	9.2	4.6	5.0
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	42.0	41.5	11.4	11.2	6.2	6.4
Black Only	37.7	35.5	13.2 ^b	10.9	7.5	6.4
American Indian or Alaska Native Only	51.0	53.9	18.3	19.8	10.4	12.6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	6.2
Asian Only	20.8	18.9	6.1	5.2	3.2	2.7
More Than One Race	42.2	49.2	15.5	20.6	10.3	14.8
Hispanic	31.2	29.9	11.0	10.1	6.1	5.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.27A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	6,415	6,288	4,584	4,357	2,265	2,264
GENDER						
Male	3,364	3,250	2,365	2,206	1,195	1,178
Female	3,051	3,038	2,219	2,151	1,070	1,086
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	4,264	4,227	3,077	3,021	1,496	1,546
Black Only	833	818	588	510	301	279
American Indian or Alaska Native Only	61	*	40	*	26	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	172	144	117	97	64	48
More Than One Race	109	102	79	61	37	37
Hispanic	961	904	672	601	334	313
GENDER/RACE/HISPANIC ORIGIN						
Male - White	2,204	2,140	1,547	1,502	748	795
Female - White	2,061	2,087	1,530	1,519	748	751
Male - Black	461	434	325	277	179	149
Female - Black	372	383	263	233	122	131
Male - Hispanic	506	496	358	320	191	173
Female - Hispanic	455	408	314	282	143	140

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.27B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	27.6	26.9	19.8 ^a	18.6	9.8	9.7
GENDER						
Male	28.3	27.2	19.9 ^a	18.4	10.1	9.8
Female	26.9	26.6	19.6	18.9	9.4	9.5
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	27.8	27.6	20.1	19.7	9.8	10.1
Black Only	25.9	24.5	18.3 ^a	15.3	9.4	8.4
American Indian or Alaska Native Only	46.5	*	30.5	*	20.0	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	19.1	17.3	13.0	11.6	7.1	5.8
More Than One Race	29.7	27.9	21.6	16.8	10.2	10.1
Hispanic	29.8	27.3	20.9 ^a	18.2	10.4	9.5
GENDER/RACE/HISPANIC ORIGIN						
Male - White	28.0	27.2	19.7	19.1	9.5	10.1
Female - White	27.7	28.0	20.5	20.4	10.1	10.1
Male - Black	28.2	25.5	19.9 ^a	16.3	10.9	8.7
Female - Black	23.5	23.5	16.6	14.2	7.7	8.0
Male - Hispanic	30.7	28.9	21.8	18.6	11.6	10.1
Female - Hispanic	28.9	25.6	19.9	17.7	9.1	8.8

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.28A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	14,975	14,846	8,287	8,088	4,658	4,599
GENDER						
Male	7,954	7,888	4,736	4,599	2,869	2,745
Female	7,021	6,958	3,551	3,489	1,789	1,854
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	10,781	10,732	5,980	5,882	3,296	3,368
Black Only	1,756	1,746	1,035	970	601	567
American Indian or Alaska Native Only	79 ^b	135	42	*	23	30
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	335	280	196	140	87	72
More Than One Race	232	200	123	130	80	81
Hispanic	1,748	1,700	889	876	556	467
ADULT EDUCATION						
< High School	3,118	3,195	1,822	1,852	1,119	1,147
High School Graduate	5,346	5,318	2,917	2,808	1,676	1,593
Some College	4,848	4,732	2,722	2,642	1,515	1,506
College Graduate	1,663	1,602	827	786	348	353
CURRENT EMPLOYMENT						
Full-Time	8,115	8,235	4,125	4,158	2,262	2,283
Part-Time	3,525	3,353	2,176 ^a	1,941	1,252	1,106
Unemployed	914	946	595	621	375	407
Other ¹	2,421	2,312	1,391	1,368	769	804

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.28B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	52.6	51.2	29.1	27.9	16.4	15.9
GENDER						
Male	56.2	54.7	33.5	31.9	20.3	19.0
Female	49.0	47.8	24.8	23.9	12.5	12.7
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	57.5	56.1	31.9	30.7	17.6	17.6
Black Only	45.6	44.5	26.9	24.7	15.6	14.5
American Indian or Alaska Native Only	57.9 ^b	76.3	30.8	*	16.9	17.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	30.8	27.9	18.0	14.0	8.0	7.2
More Than One Race	67.1	66.4	35.5	43.3	23.0	27.0
Hispanic	41.5	39.2	21.1	20.2	13.2 ^a	10.8
ADULT EDUCATION						
< High School	53.3	52.3	31.1	30.3	19.1	18.8
High School Graduate	52.2	51.1	28.5	27.0	16.4	15.3
Some College	53.2	52.2	29.9	29.2	16.6	16.6
College Graduate	50.9	46.9	25.3	23.0	10.6	10.3
CURRENT EMPLOYMENT						
Full-Time	55.2	53.7	28.1	27.1	15.4	14.9
Part-Time	51.7	50.2	31.9 ^a	29.0	18.4	16.5
Unemployed	55.0	58.3	35.8	38.3	22.6	25.1
Other ¹	45.7	43.3	26.2	25.6	14.5	15.1

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.29A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	66,344	65,797	12,531	12,089	6,906	7,164
GENDER						
Male	35,041	35,565	7,585	7,075	4,528	4,349
Female	31,303	30,232	4,945	5,014	2,378	2,815
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	53,001	52,608	9,482	9,381	5,181	5,574
Black Only	6,770	6,450	1,655	1,297	951	767
American Indian or Alaska Native Only	*	*	*	92	70	73
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1,000	1,016	132	160	79	86
More Than One Race	481	613	100	*	84	*
Hispanic	4,473	4,537	978	935	515	491
ADULT EDUCATION						
< High School	7,814	7,023	2,249	1,864	1,350	1,079
High School Graduate	21,102	21,158	3,767	3,978	2,105 ^a	2,714
Some College	18,259	17,336	3,337	2,872	1,778	1,636
College Graduate	19,169	20,280	3,178	3,375	1,672	1,734
CURRENT EMPLOYMENT						
Full-Time	48,010	48,244	8,799	8,621	4,659	4,978
Part-Time	6,526	6,481	1,177	1,095	690	712
Unemployed	1,914	1,602	661	439	462	313
Other ¹	9,894	9,470	1,893	1,934	1,095	1,161

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.29B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	39.2	38.5	7.4	7.1	4.1	4.2
GENDER						
Male	43.7	43.9	9.5	8.7	5.6	5.4
Female	35.1	33.6	5.5	5.6	2.7	3.1
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	41.4	41.0	7.4	7.3	4.0	4.3
Black Only	38.1	35.6	9.3 ^a	7.1	5.4	4.2
American Indian or Alaska Native Only	*	*	*	12.0	7.9	9.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	19.0	17.6	2.5	2.8	1.5	1.5
More Than One Race	38.9	51.3	8.1	*	6.8	*
Hispanic	28.7	28.0	6.3	5.8	3.3	3.0
ADULT EDUCATION						
< High School	25.5	24.0	7.3	6.4	4.4	3.7
High School Graduate	36.6	37.3	6.5	7.0	3.7 ^b	4.8
Some College	47.0	44.2	8.6	7.3	4.6	4.2
College Graduate	45.2	44.4	7.5	7.4	3.9	3.8
CURRENT EMPLOYMENT						
Full-Time	48.8	47.9	9.0	8.6	4.7	4.9
Part-Time	38.5	38.8	6.9	6.6	4.1	4.3
Unemployed	54.6	52.5	18.8	14.4	13.2	10.3
Other ¹	19.5	18.8	3.7	3.8	2.2	2.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.30A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	81,319	80,643	20,818	20,177	11,564	11,763
GENDER						
Male	42,995	43,453	12,322	11,674	7,397	7,094
Female	38,324	37,190	8,496	8,503	4,167	4,669
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	63,782	63,340	15,462	15,263	8,477	8,942
Black Only	8,526	8,195	2,690	2,267	1,553	1,334
American Indian or Alaska Native Only	*	508	171	158	93	103
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	29
Asian Only	1,335	1,296	328	300	165	159
More Than One Race	713	813	223	323	164	*
Hispanic	6,221	6,237	1,866	1,811	1,071	958
ADULT EDUCATION						
< High School	10,933	10,218	4,070	3,716	2,470	2,227
High School Graduate	26,448	26,476	6,684	6,786	3,781 ^a	4,308
Some College	23,107	22,068	6,059	5,514	3,293	3,142
College Graduate	20,832	21,881	4,005	4,161	2,020	2,086
CURRENT EMPLOYMENT						
Full-Time	56,125	56,479	12,924	12,779	6,921	7,261
Part-Time	10,050	9,833	3,353	3,036	1,942	1,817
Unemployed	2,828	2,548	1,256	1,060	837	720
Other ¹	12,316	11,783	3,284	3,302	1,864	1,965

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.30B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	41.1	40.3	10.5	10.1	5.8	5.9
GENDER						
Male	45.6	45.6	13.1	12.2	7.8	7.4
Female	37.0	35.6	8.2	8.1	4.0	4.5
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	43.5	42.9	10.5	10.3	5.8	6.1
Black Only	39.4	37.1	12.4 ^a	10.3	7.2	6.0
American Indian or Alaska Native Only	*	53.7	16.7	16.7	9.1	10.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	6.0
Asian Only	21.0	19.1	5.2	4.4	2.6	2.3
More Than One Race	45.1	54.3	14.1	21.6	10.3	*
Hispanic	31.4	30.4	9.4	8.8	5.4	4.7
ADULT EDUCATION						
< High School	30.0	28.9	11.2	10.5	6.8	6.3
High School Graduate	39.0	39.4	9.9	10.1	5.6 ^a	6.4
Some College	48.2 ^a	45.7	12.6	11.4	6.9	6.5
College Graduate	45.6	44.6	8.8	8.5	4.4	4.2
CURRENT EMPLOYMENT						
Full-Time	49.7	48.7	11.4	11.0	6.1	6.3
Part-Time	42.3	42.1	14.1	13.0	8.2	7.8
Unemployed	54.7	54.5	24.3	22.7	16.2	15.4
Other ¹	22.0	21.1	5.9	5.9	3.3	3.5

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.31A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	76,428	76,321	19,102	18,589	10,458	10,714
AGE						
12-17	4,345	4,283	3,284	3,136	1,676	1,678
18-25	13,326	13,256	6,962	6,860	4,049	3,950
26 or Older	58,757	58,782	8,856	8,593	4,734	5,085
GENDER						
Male	41,129	41,589	11,817	11,154	6,921	6,624
Female	35,299	34,732	7,285	7,435	3,538 ^a	4,089
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	60,132	60,279	14,043	14,067	7,579	8,058
Black Only	7,989	7,834	2,477	2,187	1,457	1,329
American Indian or Alaska Native Only	465	477	142	166	80	110
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	13
Asian Only	1,074	925	325 ^a	210	170	110
More Than One Race	719	801	260	333	166	233
Hispanic	5,832	5,770	1,772	1,570	962	861

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.31B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	34.6	34.2	8.6	8.3	4.7	4.8
AGE						
12-17	18.7	18.3	14.2	13.4	7.2	7.2
18-25	46.8	45.7	24.5	23.7	14.2	13.6
26 or Older	34.7	34.4	5.2	5.0	2.8	3.0
GENDER						
Male	38.7	38.7	11.1	10.4	6.5	6.2
Female	30.7	30.0	6.3	6.4	3.1 ^a	3.5
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	37.1	37.0	8.7	8.6	4.7	4.9
Black Only	32.1	30.8	10.0	8.6	5.9	5.2
American Indian or Alaska Native Only	40.4	43.7	12.3	15.3	6.9	10.1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	2.5
Asian Only	14.8	12.1	4.5 ^b	2.8	2.3	1.4
More Than One Race	36.9	43.1	13.4	17.9	8.5	12.5
Hispanic	25.3	24.2	7.7	6.6	4.2	3.6

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.32A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	4,345	4,283	3,284	3,136	1,676	1,678
GENDER						
Male	2,339	2,237	1,773 ^a	1,629	921	925
Female	2,006	2,046	1,511	1,507	755	753
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	2,924	2,967	2,234	2,238	1,132	1,194
Black Only	516	482	390 ^a	311	213	181
American Indian or Alaska Native Only	47	*	30	*	19	29
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	108 ^a	64	80 ^a	49	52 ^a	21
More Than One Race	79	68	63	51	31	30
Hispanic	662	626	480	431	223	220
GENDER/RACE/HISPANIC ORIGIN						
Male - White	1,515	1,503	1,147	1,142	585	648
Female - White	1,410	1,464	1,087	1,097	548	546
Male - Black	307	278	236	187	126	109
Female - Black	209	204	154	124	87	71
Male - Hispanic	370	347	279	228	143	125
Female - Hispanic	292	279	201	204	80	95

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.32B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	18.7	18.3	14.2	13.4	7.2	7.2
GENDER						
Male	19.7	18.7	14.9 ^a	13.6	7.8	7.7
Female	17.7	17.9	13.3	13.2	6.7	6.6
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	19.1	19.4	14.6	14.6	7.4	7.8
Black Only	16.0	14.4	12.1 ^b	9.3	6.6	5.4
American Indian or Alaska Native Only	35.6	*	22.9	*	14.4	19.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	12.1 ^a	7.6	8.9	5.9	5.8 ^a	2.5
More Than One Race	21.5	18.8	17.1	14.0	8.6	8.3
Hispanic	20.6	18.9	14.9	13.0	6.9	6.6
GENDER/RACE/HISPANIC ORIGIN						
Male - White	19.2	19.1	14.6	14.5	7.4	8.3
Female - White	18.9	19.6	14.6	14.7	7.4	7.3
Male - Black	18.8	16.3	14.4 ^a	11.0	7.7	6.4
Female - Black	13.2	12.5	9.7	7.6	5.5	4.4
Male - Hispanic	22.5	20.2	17.0 ^a	13.2	8.7	7.3
Female - Hispanic	18.6	17.5	12.8	12.8	5.1	6.0

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.33A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	13,326	13,256	6,962	6,860	4,049	3,950
GENDER						
Male	7,163	7,076	4,108	3,980	2,538	2,408
Female	6,163	6,180	2,855	2,880	1,511	1,542
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	9,723	9,694	5,055	5,052	2,877	2,917
Black Only	1,557	1,543	919	856	545	515
American Indian or Alaska Native Only	68 ^b	123	32	*	21	27
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	7
Asian Only	270	208	150 ^a	97	68	57
More Than One Race	212	188	105	110	68	69
Hispanic	1,459	1,453	680	674	456 ^a	359
ADULT EDUCATION						
< High School	2,759	2,827	1,502	1,551	934	985
High School Graduate	4,760	4,712	2,419	2,353	1,442	1,367
Some College	4,289	4,288	2,316	2,269	1,357	1,287
College Graduate	1,519	1,429	726	687	317	311
CURRENT EMPLOYMENT						
Full-Time	7,262	7,388	3,418	3,488	1,949	1,933
Part-Time	3,135	2,979	1,887	1,687	1,113	979
Unemployed	819	851	502	532	312	353
Other ¹	2,111	2,038	1,155	1,152	676	685

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.33B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	46.8	45.7	24.5	23.7	14.2	13.6
GENDER						
Male	50.7	49.1	29.1	27.6	17.9	16.7
Female	43.0	42.4	19.9	19.8	10.5	10.6
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	51.8	50.6	27.0	26.4	15.3	15.2
Black Only	40.4	39.3	23.8	21.8	14.1	13.1
American Indian or Alaska Native Only	49.8 ^b	69.5	23.8	*	15.1	15.1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	6.8
Asian Only	24.9	20.8	13.8 ^a	9.7	6.3	5.7
More Than One Race	61.3	62.5	30.2	36.4	19.5	22.8
Hispanic	34.6	33.5	16.2	15.5	10.8 ^a	8.3
ADULT EDUCATION						
< High School	47.2	46.3	25.7	25.4	16.0	16.1
High School Graduate	46.5	45.3	23.6	22.6	14.1	13.1
Some College	47.1	47.3	25.4	25.0	14.9	14.2
College Graduate	46.5 ^a	41.8	22.2	20.1	9.7	9.1
CURRENT EMPLOYMENT						
Full-Time	49.4	48.2	23.3	22.7	13.3	12.6
Part-Time	46.0	44.6	27.7 ^a	25.2	16.3	14.6
Unemployed	49.3	52.4	30.2	32.8	18.8	21.8
Other ¹	39.8	38.2	21.8	21.6	12.8	12.8

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.34A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	58,757	58,782	8,856	8,593	4,734	5,085
GENDER						
Male	31,627	32,275	5,936	5,545	3,462	3,290
Female	27,130	26,507	2,920	3,049	1,271 ^b	1,794
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	47,485	47,618	6,754	6,777	3,570	3,948
Black Only	5,916	5,809	1,168	1,021	698	633
American Indian or Alaska Native Only	351	*	80	64	40	54
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	695	653	96	64	50	32
More Than One Race	*	545	93	*	67	*
Hispanic	3,711	3,690	611	464	283	282
ADULT EDUCATION						
< High School	6,462	6,043	1,640	1,267	976	757
High School Graduate	18,517	18,623	2,773	2,742	1,452 ^a	1,867
Some College	16,613	15,831	2,364	2,103	1,262	1,227
College Graduate	17,165	18,284	2,079	2,481	1,044	1,234
CURRENT EMPLOYMENT						
Full-Time	43,819	43,942	6,449	6,358	3,389	3,617
Part-Time	5,625	5,777	766	773	448	490
Unemployed	1,705	1,386	545	346	314	267
Other ¹	7,608	7,678	1,096	1,116	583	712

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.34B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	34.7	34.4	5.2	5.0	2.8	3.0
GENDER						
Male	39.4	39.9	7.4	6.8	4.3	4.1
Female	30.4	29.5	3.3	3.4	1.4 ^b	2.0
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	37.1	37.1	5.3	5.3	2.8	3.1
Black Only	33.3	32.0	6.6	5.6	3.9	3.5
American Indian or Alaska Native Only	39.7	*	9.0	8.3	4.6	7.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	13.2	11.3	1.8	1.1	0.9	0.5
More Than One Race	*	45.6	7.5	*	5.4	*
Hispanic	23.8	22.8	3.9	2.9	1.8	1.7
ADULT EDUCATION						
< High School	21.1	20.7	5.4	4.3	3.2	2.6
High School Graduate	32.2	32.8	4.8	4.8	2.5 ^a	3.3
Some College	42.8	40.3	6.1	5.4	3.3	3.1
College Graduate	40.5	40.0	4.9	5.4	2.5	2.7
CURRENT EMPLOYMENT						
Full-Time	44.6	43.6	6.6	6.3	3.4	3.6
Part-Time	33.2	34.6	4.5	4.6	2.6	2.9
Unemployed	48.6	45.4	15.5	11.3	8.9	8.7
Other ¹	15.0	15.2	2.2	2.2	1.1	1.4

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.35A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	72,083	72,038	15,818	15,453	8,783	9,035
GENDER						
Male	38,790	39,352	10,044	9,524	6,000	5,699
Female	33,293	32,686	5,775	5,929	2,783 ^b	3,337
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	57,208	57,313	11,809	11,829	6,447	6,864
Black Only	7,473	7,352	2,087	1,877	1,244	1,148
American Indian or Alaska Native Only	418	409	112	116	61	81
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	10
Asian Only	965	861	245	161	118	88
More Than One Race	640	733	197	282	135	*
Hispanic	5,170	5,143	1,292	1,139	739	641
ADULT EDUCATION						
< High School	9,221	8,870	3,142	2,817	1,910	1,742
High School Graduate	23,277	23,335	5,192	5,095	2,893	3,234
Some College	20,902	20,119	4,680	4,372	2,619	2,514
College Graduate	18,683	19,714	2,804	3,169	1,361	1,545
CURRENT EMPLOYMENT						
Full-Time	51,080	51,330	9,867	9,846	5,338	5,550
Part-Time	8,760	8,756	2,653	2,461	1,560	1,469
Unemployed	2,524	2,237	1,048	879	626	620
Other ¹	9,719	9,715	2,250	2,268	1,259	1,397

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.35B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	36.4	36.0	8.0	7.7	4.4	4.5
GENDER						
Male	41.1	41.3	10.6	10.0	6.4	6.0
Female	32.1	31.3	5.6	5.7	2.7 ^a	3.2
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	39.0	38.8	8.0	8.0	4.4	4.7
Black Only	34.5	33.3	9.6	8.5	5.7	5.2
American Indian or Alaska Native Only	41.0	43.2	11.0	12.3	6.0	8.6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	2.1
Asian Only	15.2	12.7	3.9 ^a	2.4	1.9	1.3
More Than One Race	40.5	49.0	12.5	18.8	8.5	*
Hispanic	26.1	25.0	6.5	5.5	3.7	3.1
ADULT EDUCATION						
< High School	25.3	25.1	8.6	8.0	5.2	4.9
High School Graduate	34.3	34.8	7.7	7.6	4.3	4.8
Some College	43.6	41.7	9.8	9.1	5.5	5.2
College Graduate	40.9	40.1	6.1	6.5	3.0	3.1
CURRENT EMPLOYMENT						
Full-Time	45.2	44.2	8.7	8.5	4.7	4.8
Part-Time	36.9	37.5	11.2	10.5	6.6	6.3
Unemployed	48.8	47.8	20.3	18.8	12.1	13.3
Other ¹	17.4	17.4	4.0	4.1	2.2	2.5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.36A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	25,406	24,896	3,742	3,328	1,552	1,213
AGE						
12-17	551	550	373	389	124	132
18-25	3,383	3,148	1,493 ^b	1,274	476	395
26 or Older	21,471	21,198	1,876	1,666	952	687
GENDER						
Male	14,890	14,690	2,255	2,037	973	760
Female	10,516	10,206	1,486	1,291	579	454
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	20,202	20,200	2,699	2,459	1,095 ^a	779
Black Only	2,368 ^a	1,875	368	340	232	190
American Indian or Alaska Native Only	215	179	41	48	12	15
Native Hawaiian or Other Pacific Islander	*	*	4	*	1	*
Asian Only	171	161	36	15	12	7
More Than One Race	238	306	65	46	15	23
Hispanic	2,122	2,090	528	406	185	191

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.36B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11.5	11.2	1.7	1.5	0.7	0.5
AGE						
12-17	2.4	2.4	1.6	1.7	0.5	0.6
18-25	11.9 ^a	10.9	5.2 ^b	4.4	1.7	1.4
26 or Older	12.7	12.4	1.1	1.0	0.6	0.4
GENDER						
Male	14.0	13.7	2.1	1.9	0.9	0.7
Female	9.2	8.8	1.3	1.1	0.5	0.4
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	12.5	12.4	1.7	1.5	0.7 ^a	0.5
Black Only	9.5 ^a	7.4	1.5	1.3	0.9	0.7
American Indian or Alaska Native Only	18.7	16.4	3.5	4.4	1.0	1.4
Native Hawaiian or Other Pacific Islander	*	*	0.6	*	0.1	*
Asian Only	2.4	2.1	0.5	0.2	0.2	0.1
More Than One Race	12.2	16.4	3.3	2.5	0.8	1.2
Hispanic	9.2	8.8	2.3	1.7	0.8	0.8

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.37A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	551	550	373	389	124	132
GENDER						
Male	247	263	159	186	58	70
Female	304	287	214	202	66	62
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	379	387	268	272	96	98
Black Only	13	16	4	12	1	3
American Indian or Alaska Native Only	*	*	*	*	2	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	4	4	2	2	1	1
More Than One Race	11	6	4	5	1	1
Hispanic	129	129	84	89	23	24
GENDER/RACE/HISPANIC ORIGIN						
Male - White	155	182	99	126	41	49
Female - White	225	205	169	146	54	49
Male - Black	9	10	4	7	1	2
Female - Black	3	6	0	5	*	1
Male - Hispanic	75	68	52	50	15	17
Female - Hispanic	54	61	32	38	9	7

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.37B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	2.4	2.4	1.6	1.7	0.5	0.6
GENDER						
Male	2.1	2.2	1.3	1.6	0.5	0.6
Female	2.7	2.5	1.9	1.8	0.6	0.5
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	2.5	2.5	1.7	1.8	0.6	0.6
Black Only	0.4	0.5	0.1	0.4	0.0	0.1
American Indian or Alaska Native Only	*	*	*	*	1.5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.4	0.4	0.3	0.2	0.1	0.1
More Than One Race	2.9	1.8	1.0	1.5	0.2	0.3
Hispanic	4.0	3.9	2.6	2.7	0.7	0.7
GENDER/RACE/HISPANIC ORIGIN						
Male - White	2.0	2.3	1.3	1.6	0.5	0.6
Female - White	3.0	2.8	2.3	2.0	0.7	0.7
Male - Black	0.6	0.6	0.2	0.4	0.1	0.1
Female - Black	0.2	0.4	0.0	0.3	*	0.1
Male - Hispanic	4.5	4.0	3.2	2.9	0.9	1.0
Female - Hispanic	3.4	3.8	2.0	2.4	0.6	0.4

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.38A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	3,383	3,148	1,493 ^b	1,274	476	395
GENDER						
Male	2,046	1,870	949 ^a	780	298	238
Female	1,337	1,279	544	494	178	156
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	2,515	2,448	1,104	1,001	342	275
Black Only	147	124	72	52	31	16
American Indian or Alaska Native Only	24	*	7	*	1	1
Native Hawaiian or Other Pacific Islander	*	9	*	2	*	*
Asian Only	57	32	29	12	12	5
More Than One Race	66	49	39	19	14	5
Hispanic	566 ^a	453	241 ^a	176	75	93
ADULT EDUCATION						
< High School	909	876	384	339	141	133
High School Graduate	1,258	1,139	561 ^a	453	175	135
Some College	1,008	933	439	384	126	96
College Graduate	208	201	108	98	34	30
CURRENT EMPLOYMENT						
Full-Time	1,995	1,878	823	706	247	210
Part-Time	675	600	334	285	101	91
Unemployed	236	244	119	123	54	50
Other ¹	478	427	216	160	75	44

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.38B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11.9 ^a	10.9	5.2 ^b	4.4	1.7	1.4
GENDER						
Male	14.5 ^a	13.0	6.7 ^b	5.4	2.1	1.7
Female	9.3	8.8	3.8	3.4	1.2	1.1
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	13.4	12.8	5.9	5.2	1.8	1.4
Black Only	3.8	3.2	1.9	1.3	0.8	0.4
American Indian or Alaska Native Only	17.6	*	5.1	*	0.7	0.7
Native Hawaiian or Other Pacific Islander	*	8.3	*	1.9	*	*
Asian Only	5.3	3.2	2.7	1.2	1.1	0.5
More Than One Race	19.1	16.3	11.2	6.4	4.0	1.5
Hispanic	13.4 ^b	10.5	5.7 ^a	4.1	1.8	2.1
ADULT EDUCATION						
< High School	15.5	14.4	6.6	5.6	2.4	2.2
High School Graduate	12.3	10.9	5.5 ^a	4.4	1.7	1.3
Some College	11.1	10.3	4.8	4.2	1.4	1.1
College Graduate	6.4	5.9	3.3	2.9	1.1	0.9
CURRENT EMPLOYMENT						
Full-Time	13.6 ^a	12.2	5.6 ^a	4.6	1.7	1.4
Part-Time	9.9	9.0	4.9	4.3	1.5	1.4
Unemployed	14.2	15.0	7.2	7.6	3.2	3.1
Other ¹	9.0	8.0	4.1 ^a	3.0	1.4	0.8

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.39A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	21,471	21,198	1,876	1,666	952	687
GENDER						
Male	12,597	12,557	1,148	1,070	617	451
Female	8,874	8,641	728	595	335	235
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	17,307	17,366	1,327	1,186	657	406
Black Only	2,209	1,734	292	276	200	171
American Indian or Alaska Native Only	178	*	26	*	9	9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	110	126	4	1	*	1
More Than One Race	162	250	23	21	*	17
Hispanic	1,428	1,508	204	141	86	73
ADULT EDUCATION						
< High School	2,715	2,412	503	324	306	158
High School Graduate	6,590	6,932	597	620	298	243
Some College	6,334	6,121	405	376	205	118
College Graduate	5,832	5,732	372	347	143	168
CURRENT EMPLOYMENT						
Full-Time	15,888	15,825	1,274	1,163	700 ^a	431
Part-Time	2,075	1,978	203	229	80	109
Unemployed	954	704	208	86	97	41
Other ¹	2,555	2,690	191	188	76	106

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.39B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	12.7	12.4	1.1	1.0	0.6	0.4
GENDER						
Male	15.7	15.5	1.4	1.3	0.8	0.6
Female	9.9	9.6	0.8	0.7	0.4	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	13.5	13.5	1.0	0.9	0.5	0.3
Black Only	12.4 ^a	9.6	1.6	1.5	1.1	0.9
American Indian or Alaska Native Only	20.1	*	2.9	*	1.0	1.2
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	2.1	2.2	0.1	0.0	*	0.0
More Than One Race	13.1	20.9	1.8	1.8	*	1.4
Hispanic	9.2	9.3	1.3	0.9	0.6	0.5
ADULT EDUCATION						
< High School	8.9	8.2	1.6	1.1	1.0	0.5
High School Graduate	11.4	12.2	1.0	1.1	0.5	0.4
Some College	16.3	15.6	1.0	1.0	0.5	0.3
College Graduate	13.8	12.5	0.9	0.8	0.3	0.4
CURRENT EMPLOYMENT						
Full-Time	16.2	15.7	1.3	1.2	0.7 ^a	0.4
Part-Time	12.3	11.9	1.2	1.4	0.5	0.7
Unemployed	27.2	23.1	5.9	2.8	2.8	1.3
Other ¹	5.0	5.3	0.4	0.4	0.1	0.2

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.40A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	24,854	24,346	3,369	2,940	1,428	1,081
GENDER						
Male	14,643	14,427	2,097	1,851	915	690
Female	10,212	9,919	1,272	1,089	513	392
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	19,823	19,814	2,431	2,187	999 ^a	681
Black Only	2,356 ^a	1,859	364	328	231	187
American Indian or Alaska Native Only	202	*	33	40	10	10
Native Hawaiian or Other Pacific Islander	*	*	1	*	1	*
Asian Only	167	158	34	13	12	6
More Than One Race	228	299	61	40	14	22
Hispanic	1,993	1,961	445	317	161	166
ADULT EDUCATION						
< High School	3,624	3,289	887	663	446	292
High School Graduate	7,848	8,070	1,158	1,073	473	378
Some College	7,342	7,054	843	760	331	214
College Graduate	6,040	5,933	480	444	178	198
CURRENT EMPLOYMENT						
Full-Time	17,883	17,703	2,097	1,869	947 ^a	641
Part-Time	2,750	2,579	538	514	181	200
Unemployed	1,190	948	327	209	150	91
Other ¹	3,033	3,116	407	348	151	150

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.40B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	12.6	12.2	1.7	1.5	0.7 ^a	0.5
GENDER						
Male	15.5	15.1	2.2	1.9	1.0	0.7
Female	9.9	9.5	1.2	1.0	0.5	0.4
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	13.5	13.4	1.7	1.5	0.7 ^a	0.5
Black Only	10.9 ^a	8.4	1.7	1.5	1.1	0.8
American Indian or Alaska Native Only	19.8	*	3.2	4.3	1.0	1.1
Native Hawaiian or Other Pacific Islander	*	*	0.2	*	0.1	*
Asian Only	2.6	2.3	0.5	0.2	0.2	0.1
More Than One Race	14.4	20.0	3.9	2.7	0.9	1.5
Hispanic	10.1	9.6	2.2	1.5	0.8	0.8
ADULT EDUCATION						
< High School	9.9	9.3	2.4	1.9	1.2	0.8
High School Graduate	11.6	12.0	1.7	1.6	0.7	0.6
Some College	15.3	14.6	1.8	1.6	0.7	0.4
College Graduate	13.2	12.1	1.1	0.9	0.4	0.4
CURRENT EMPLOYMENT						
Full-Time	15.8	15.3	1.9	1.6	0.8 ^a	0.6
Part-Time	11.6	11.0	2.3	2.2	0.8	0.9
Unemployed	23.0	20.3	6.3	4.5	2.9	2.0
Other ¹	5.4	5.6	0.7	0.6	0.3	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.41A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	25,061	26,125	3,191	3,483	922	971
AGE						
12-17	1,317	1,347	889	908	254	270
18-25	5,490	5,592	1,928	1,959	534	532
26 or Older	18,254	19,186	374 ^a	616	134	169
GENDER						
Male	14,753	15,506	1,941	2,138	566	584
Female	10,308	10,619	1,250	1,345	356	387
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	21,197	22,334	2,594 ^a	2,880	743	804
Black Only	1,222	1,281	96	144	29	45
American Indian or Alaska Native Only	217	265	30	38	9	7
Native Hawaiian or Other Pacific Islander	*	45	9	7	0	4
Asian Only	272	223	82	68	23	18
More Than One Race	283	343	61	58	14	15
Hispanic	1,785	1,634	320	289	104	79

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.41B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11.3	11.7	1.4	1.6	0.4	0.4
AGE						
12-17	5.7	5.8	3.8	3.9	1.1	1.2
18-25	19.3	19.3	6.8	6.8	1.9	1.8
26 or Older	10.8	11.2	0.2 ^a	0.4	0.1	0.1
GENDER						
Male	13.9	14.4	1.8	2.0	0.5	0.5
Female	9.0	9.2	1.1	1.2	0.3	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	13.1	13.7	1.6	1.8	0.5	0.5
Black Only	4.9	5.0	0.4	0.6	0.1	0.2
American Indian or Alaska Native Only	18.8	24.3	2.6	3.4	0.8	0.7
Native Hawaiian or Other Pacific Islander	*	8.2	1.1	1.2	0.0	0.8
Asian Only	3.7	2.9	1.1	0.9	0.3	0.2
More Than One Race	14.5	18.4	3.1	3.1	0.7	0.8
Hispanic	7.8	6.9	1.4	1.2	0.5	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.42A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	1,317	1,347	889	908	254	270
GENDER						
Male	680	712	457	477	114	145
Female	637	636	432	431	140	125
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	1,026	1,053	702	718	196	213
Black Only	36	53	24	35	6	14
American Indian or Alaska Native Only	21	18	15	11	7	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	23	34	14 ^a	29	5	12
More Than One Race	23	25	13	17	1	3
Hispanic	184	163	118	97	39	25
GENDER/RACE/HISPANIC ORIGIN						
Male - White	527	557	356	375	86	118
Female - White	499	497	345	344	111	95
Male - Black	21	27	17	22	4	7
Female - Black	14	26	7	13	2	7
Male - Hispanic	98	91	61	54	19	10
Female - Hispanic	86	72	57	43	21	16

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.42B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	5.7	5.8	3.8	3.9	1.1	1.2
GENDER						
Male	5.7	6.0	3.9	4.0	1.0	1.2
Female	5.6	5.6	3.8	3.8	1.2	1.1
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	6.7	6.9	4.6	4.7	1.3	1.4
Black Only	1.1	1.6	0.7	1.1	0.2	0.4
American Indian or Alaska Native Only	16.0	12.4	11.3	7.4	5.5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	2.5	4.1	1.6 ^a	3.4	0.5	1.4
More Than One Race	6.4	6.9	3.6	4.7	0.2	0.7
Hispanic	5.7	4.9	3.7	2.9	1.2	0.8
GENDER/RACE/HISPANIC ORIGIN						
Male - White	6.7	7.1	4.5	4.8	1.1	1.5
Female - White	6.7	6.7	4.6	4.6	1.5	1.3
Male - Black	1.3	1.6	1.0	1.3	0.2	0.4
Female - Black	0.9	1.6	0.4	0.8	0.1	0.4
Male - Hispanic	5.9	5.3	3.7	3.2	1.1	0.6
Female - Hispanic	5.5	4.5	3.6	2.7	1.3	1.0

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.43A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	5,490	5,592	1,928	1,959	534	532
GENDER						
Male	3,148	3,238	1,204	1,189	336	332
Female	2,342	2,355	725	770	198	200
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	4,443	4,623	1,582	1,610	442	446
Black Only	161	200	60	98	15	28
American Indian or Alaska Native Only	39	50	10	18	2	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	149 ^b	83	62	31	19	5
More Than One Race	95	91	43	37	13	12
Hispanic	584	521	166	159	44	35
ADULT EDUCATION						
< High School	1,059	1,145	383	433	108	113
High School Graduate	1,994	1,996	734	676	194	194
Some College	1,861	1,861	662	683	186	192
College Graduate	576	591	150	166	46	32
CURRENT EMPLOYMENT						
Full-Time	3,111	3,152	945	963	265	245
Part-Time	1,229	1,285	532	524	151	137
Unemployed	314 ^a	399	133	179	51	62
Other ¹	837	756	318	293	67	88

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.43B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	19.3	19.3	6.8	6.8	1.9	1.8
GENDER						
Male	22.3	22.5	8.5	8.2	2.4	2.3
Female	16.3	16.2	5.1	5.3	1.4	1.4
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	23.7	24.2	8.4	8.4	2.4	2.3
Black Only	4.2	5.1	1.6	2.5	0.4	0.7
American Indian or Alaska Native Only	29.0	28.3	7.5	10.4	1.4	0.7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	13.7 ^a	8.3	5.7	3.1	1.7	0.5
More Than One Race	27.4	30.3	12.3	12.2	3.7	4.1
Hispanic	13.9	12.0	4.0	3.7	1.0	0.8
ADULT EDUCATION						
< High School	18.1	18.7	6.5	7.1	1.8	1.9
High School Graduate	19.5	19.2	7.2	6.5	1.9	1.9
Some College	20.4	20.5	7.3	7.5	2.0	2.1
College Graduate	17.6	17.3	4.6	4.9	1.4	0.9
CURRENT EMPLOYMENT						
Full-Time	21.2	20.5	6.4	6.3	1.8	1.6
Part-Time	18.0	19.2	7.8	7.8	2.2	2.0
Unemployed	18.9 ^b	24.6	8.0 ^a	11.0	3.1	3.8
Other ¹	15.8	14.2	6.0	5.5	1.3	1.7

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.44A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	18,254	19,186	374 ^a	616	134	169
GENDER						
Male	10,924	11,557	280	472	116	107
Female	7,329	7,629	94	144	18	62
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	15,728	16,658	309 ^b	552	105	144
Black Only	1,025	1,028	13	11	8	3
American Indian or Alaska Native Only	156	*	5	9	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	100	106	6	8	*	1
More Than One Race	165	*	5	4	*	*
Hispanic	1,017	950	36	33	21	18
ADULT EDUCATION						
< High School	2,039	2,046	60	35	37	8
High School Graduate	5,294	5,764	126	128	38	24
Some College	5,649	5,592	119	178	51	66
College Graduate	5,272	5,783	68 ^b	274	7 ^b	71
CURRENT EMPLOYMENT						
Full-Time	13,679	14,529	264 ^a	451	107	101
Part-Time	1,765	1,753	52	78	9	24
Unemployed	785	677	27	43	*	18
Other ¹	2,025	2,226	30	43	18	26

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.44B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	10.8	11.2	0.2 ^a	0.4	0.1	0.1
GENDER						
Male	13.6	14.3	0.3	0.6	0.1	0.1
Female	8.2	8.5	0.1	0.2	0.0	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	12.3	13.0	0.2 ^b	0.4	0.1	0.1
Black Only	5.8	5.7	0.1	0.1	0.0	0.0
American Indian or Alaska Native Only	17.7	*	0.6	1.1	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.9	1.8	0.1	0.1	*	0.0
More Than One Race	13.3	*	0.4	0.3	*	*
Hispanic	6.5	5.9	0.2	0.2	0.1	0.1
ADULT EDUCATION						
< High School	6.7	7.0	0.2	0.1	0.1	0.0
High School Graduate	9.2	10.2	0.2	0.2	0.1	0.0
Some College	14.6	14.3	0.3	0.5	0.1	0.2
College Graduate	12.4	12.7	0.2 ^b	0.6	0.0 ^b	0.2
CURRENT EMPLOYMENT						
Full-Time	13.9	14.4	0.3 ^a	0.4	0.1	0.1
Part-Time	10.4	10.5	0.3	0.5	0.1	0.1
Unemployed	22.4	22.2	0.8	1.4	*	0.6
Other ¹	4.0	4.4	0.1	0.1	0.0	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.45A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	23,744	24,778	2,302	2,575	668	701
GENDER						
Male	14,073	14,794	1,484	1,661	452	439
Female	9,671	9,983	818	914	216	262
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	20,171	21,281	1,892 ^a	2,162	547	590
Black Only	1,186	1,228	73	108	23	31
American Indian or Alaska Native Only	196	247	15	27	2	5
Native Hawaiian or Other Pacific Islander	*	44	*	6	0	*
Asian Only	249	189	67	39	19	5
More Than One Race	260	318	48	41	13	12
Hispanic	1,602	1,471	202	192	65	53
ADULT EDUCATION						
< High School	3,098	3,191	443	469	145	121
High School Graduate	7,287	7,760	860	804	232	218
Some College	7,510	7,453	781	861	237	258
College Graduate	5,848	6,374	218 ^b	440	53	104
CURRENT EMPLOYMENT						
Full-Time	16,790	17,681	1,210	1,415	372	347
Part-Time	2,994	3,039	583	602	160	160
Unemployed	1,099	1,076	161	222	51	80
Other ¹	2,861	2,982	348	336	85	114

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.45B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	12.0	12.4	1.2	1.3	0.3	0.4
GENDER						
Male	14.9	15.5	1.6	1.7	0.5	0.5
Female	9.3	9.6	0.8	0.9	0.2	0.3
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	13.7	14.4	1.3	1.5	0.4	0.4
Black Only	5.5	5.6	0.3	0.5	0.1	0.1
American Indian or Alaska Native Only	19.2	26.1	1.5	2.8	0.2	0.5
Native Hawaiian or Other Pacific Islander	*	9.1	*	1.2	0.0	*
Asian Only	3.9	2.8	1.1	0.6	0.3	0.1
More Than One Race	16.4	21.3	3.0	2.7	0.8	0.8
Hispanic	8.1	7.2	1.0	0.9	0.3	0.3
ADULT EDUCATION						
< High School	8.5	9.0	1.2	1.3	0.4	0.3
High School Graduate	10.7	11.6	1.3	1.2	0.3	0.3
Some College	15.7	15.4	1.6	1.8	0.5	0.5
College Graduate	12.8	13.0	0.5 ^b	0.9	0.1	0.2
CURRENT EMPLOYMENT						
Full-Time	14.9	15.2	1.1	1.2	0.3	0.3
Part-Time	12.6	13.0	2.5	2.6	0.7	0.7
Unemployed	21.3	23.0	3.1 ^a	4.7	1.0	1.7
Other ¹	5.1	5.3	0.6	0.6	0.2	0.2

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.46A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	17,138	16,702	1,937	1,918	570	622
AGE						
12-17	2,118	2,079	908	826	245	223
18-25	4,014 ^a	3,701	747	696	182	162
26 or Older	11,006	10,922	282	396	144	236
GENDER						
Male	11,223	10,480	1,235	1,216	386	410
Female	5,915	6,222	702	701	184	212
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	14,458	14,155	1,491	1,524	420	458
Black Only	859	812	115	111	24	56
American Indian or Alaska Native Only	120	135	12	5	6	1
Native Hawaiian or Other Pacific Islander	*	25	5	3	0	2
Asian Only	210	191	37	39	10	18
More Than One Race	216	190	47 ^a	19	8	7
Hispanic	1,231	1,195	230	217	101	79

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.46B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	7.8	7.5	0.9	0.9	0.3	0.3
AGE						
12-17	9.1	8.9	3.9	3.5	1.1	1.0
18-25	14.1 ^b	12.8	2.6	2.4	0.6	0.6
26 or Older	6.5	6.4	0.2	0.2	0.1	0.1
GENDER						
Male	10.6	9.8	1.2	1.1	0.4	0.4
Female	5.1	5.4	0.6	0.6	0.2	0.2
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	8.9	8.7	0.9	0.9	0.3	0.3
Black Only	3.5	3.2	0.5	0.4	0.1	0.2
American Indian or Alaska Native Only	10.4	12.4	1.1	0.5	0.5	0.1
Native Hawaiian or Other Pacific Islander	*	4.5	0.6	0.5	0.0	0.4
Asian Only	2.9	2.5	0.5	0.5	0.1	0.2
More Than One Race	11.1	10.2	2.4 ^a	1.0	0.4	0.4
Hispanic	5.3	5.0	1.0	0.9	0.4	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.47A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	2,118	2,079	908	826	245	223
GENDER						
Male	1,111	1,100	456	430	118	109
Female	1,007	978	452	395	127	114
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	1,543	1,487	674	611	161	151
Black Only	157	173	49	53	17	20
American Indian or Alaska Native Only	19	14	8	*	3	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	45	49	21	24	4	7
More Than One Race	34	33	23 ^a	6	4	3
Hispanic	313	320	127	124	55	39
GENDER/RACE/HISPANIC ORIGIN						
Male - White	816	792	346	315	80	73
Female - White	727	695	328	296	81	78
Male - Black	87	81	26	26	11	9
Female - Black	70	92	23	27	6	11
Male - Hispanic	152	182	60	78	22	25
Female - Hispanic	161	138	68	46	33 ^a	14

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.47B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	9.1	8.9	3.9	3.5	1.1	1.0
GENDER						
Male	9.4	9.2	3.8	3.6	1.0	0.9
Female	8.9	8.6	4.0	3.5	1.1	1.0
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	10.1	9.7	4.4	4.0	1.1	1.0
Black Only	4.9	5.2	1.5	1.6	0.5	0.6
American Indian or Alaska Native Only	14.3	9.6	6.4	*	2.2	0.7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	5.0	5.8	2.4	2.8	0.5	0.8
More Than One Race	9.1	9.0	6.2 ^a	1.7	1.0	0.8
Hispanic	9.7	9.7	4.0	3.8	1.7	1.2
GENDER/RACE/HISPANIC ORIGIN						
Male - White	10.4	10.1	4.4	4.0	1.0	0.9
Female - White	9.8	9.3	4.4	4.0	1.1	1.0
Male - Black	5.3	4.7	1.6	1.5	0.6	0.5
Female - Black	4.4	5.6	1.4	1.6	0.4	0.7
Male - Hispanic	9.2	10.6	3.6	4.6	1.4	1.5
Female - Hispanic	10.2	8.7	4.3 ^a	2.9	2.1 ^a	0.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.48A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	4,014 ^a	3,701	747	696	182	162
GENDER						
Male	2,482	2,267	513	444	130	103
Female	1,532	1,435	234	251	52	59
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	3,305	3,128	649	592	151	134
Black Only	129	120	12	24	6	8
American Indian or Alaska Native Only	17	21	1	0	*	0
Native Hawaiian or Other Pacific Islander	*	13	*	*	*	*
Asian Only	73	45	13	6	5	2
More Than One Race	70	56	24	13	4	4
Hispanic	408 ^a	318	48	61	15	13
ADULT EDUCATION						
< High School	749	707	148	153	50	25
High School Graduate	1,433 ^a	1,256	279	231	75	56
Some College	1,344	1,291	261	255	48	65
College Graduate	487	447	59	57	8	16
CURRENT EMPLOYMENT						
Full-Time	2,246	2,079	344	313	86	75
Part-Time	946	827	243	195	60	49
Unemployed	214	263	43	55	8	13
Other ¹	608	532	117	133	28	25

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.48B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	14.1 ^b	12.8	2.6	2.4	0.6	0.6
GENDER						
Male	17.6 ^b	15.7	3.6	3.1	0.9	0.7
Female	10.7	9.8	1.6	1.7	0.4	0.4
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	17.6 ^a	16.3	3.5	3.1	0.8	0.7
Black Only	3.3	3.1	0.3	0.6	0.1	0.2
American Indian or Alaska Native Only	12.7	11.9	0.4	0.0	*	0.0
Native Hawaiian or Other Pacific Islander	*	12.3	*	*	*	*
Asian Only	6.7	4.5	1.2	0.6	0.5	0.2
More Than One Race	20.2	18.7	6.9	4.3	1.2	1.4
Hispanic	9.7 ^a	7.3	1.1	1.4	0.4	0.3
ADULT EDUCATION						
< High School	12.8	11.6	2.5	2.5	0.9	0.4
High School Graduate	14.0 ^b	12.1	2.7	2.2	0.7	0.5
Some College	14.8	14.3	2.9	2.8	0.5	0.7
College Graduate	14.9	13.1	1.8	1.7	0.2	0.5
CURRENT EMPLOYMENT						
Full-Time	15.3 ^b	13.6	2.3	2.0	0.6	0.5
Part-Time	13.9	12.4	3.6	2.9	0.9	0.7
Unemployed	12.9	16.2	2.6	3.4	0.5	0.8
Other ¹	11.5	10.0	2.2	2.5	0.5	0.5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.49A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11,006	10,922	282	396	144	236
GENDER						
Male	7,630	7,112	266	341	139	198
Female	3,376	3,809	16	55	6	38
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	9,609	9,540	168 ^a	321	108	173
Black Only	573	519	53	34	2	27
American Indian or Alaska Native Only	84	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	92	98	3	9	*	9
More Than One Race	113	100	*	*	*	*
Hispanic	510	556	55	32	31	27
ADULT EDUCATION						
< High School	1,171	1,050	13	47	7	27
High School Graduate	3,217	3,200	36	99	29	63
Some College	3,093	2,841	108	53	42	24
College Graduate	3,525	3,831	125	197	67	123
CURRENT EMPLOYMENT						
Full-Time	8,718	8,177	255	279	132	155
Part-Time	848	1,047	9	15	5	9
Unemployed	335	311	1	29	*	25
Other ¹	1,106	1,386	17	72	7	48

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.49B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	6.5	6.4	0.2	0.2	0.1	0.1
GENDER						
Male	9.5	8.8	0.3	0.4	0.2	0.2
Female	3.8	4.2	0.0	0.1	0.0	0.0
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	7.5	7.4	0.1 ^a	0.2	0.1	0.1
Black Only	3.2	2.9	0.3	0.2	0.0	0.2
American Indian or Alaska Native Only	9.5	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.7	1.7	0.0	0.2	*	0.2
More Than One Race	9.1	8.4	*	*	*	*
Hispanic	3.3	3.4	0.4	0.2	0.2	0.2
ADULT EDUCATION						
< High School	3.8	3.6	0.0	0.2	0.0	0.1
High School Graduate	5.6	5.6	0.1	0.2	0.0	0.1
Some College	8.0	7.2	0.3	0.1	0.1	0.1
College Graduate	8.3	8.4	0.3	0.4	0.2	0.3
CURRENT EMPLOYMENT						
Full-Time	8.9	8.1	0.3	0.3	0.1	0.2
Part-Time	5.0	6.3	0.1	0.1	0.0	0.1
Unemployed	9.5	10.2	0.0	1.0	*	0.8
Other ¹	2.2	2.7	0.0	0.1	0.0	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.50A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15,020	14,623	1,029	1,092	326	398
GENDER						
Male	10,112	9,379	779	786	269	301
Female	4,908	5,244	249	306	57	97
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	12,915	12,668	817	913	259	307
Black Only	702	639	66	58	8	35
American Indian or Alaska Native Only	101	*	4	*	*	*
Native Hawaiian or Other Pacific Islander	*	22	*	*	*	*
Asian Only	165	143	16	15	5	11
More Than One Race	182	157	24	13	4	4
Hispanic	918	874	103	93	46	40
ADULT EDUCATION						
< High School	1,920	1,758	161	200	57	52
High School Graduate	4,650	4,456	314	331	104	119
Some College	4,437	4,132	369	308	90	89
College Graduate	4,012	4,278	184	253	75	138
CURRENT EMPLOYMENT						
Full-Time	10,964	10,256	598	592	218	230
Part-Time	1,794	1,875	253	210	65	58
Unemployed	548	574	43	84	8	37
Other ¹	1,714	1,918	135	205	35	73

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.50B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	7.6	7.3	0.5	0.5	0.2	0.2
GENDER						
Male	10.7	9.8	0.8	0.8	0.3	0.3
Female	4.7	5.0	0.2	0.3	0.1	0.1
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	8.8	8.6	0.6	0.6	0.2	0.2
Black Only	3.2	2.9	0.3	0.3	0.0	0.2
American Indian or Alaska Native Only	9.9	*	0.4	*	*	*
Native Hawaiian or Other Pacific Islander	*	4.5	*	*	*	*
Asian Only	2.6	2.1	0.3	0.2	0.1	0.2
More Than One Race	11.5	10.5	1.5	0.9	0.3	0.3
Hispanic	4.6	4.3	0.5	0.5	0.2	0.2
ADULT EDUCATION						
< High School	5.3	5.0	0.4	0.6	0.2	0.1
High School Graduate	6.9	6.6	0.5	0.5	0.2	0.2
Some College	9.3	8.6	0.8	0.6	0.2	0.2
College Graduate	8.8	8.7	0.4	0.5	0.2	0.3
CURRENT EMPLOYMENT						
Full-Time	9.7 ^a	8.8	0.5	0.5	0.2	0.2
Part-Time	7.6	8.0	1.1	0.9	0.3	0.2
Unemployed	10.6	12.3	0.8 ^a	1.8	0.2	0.8
Other ¹	3.1	3.4	0.2	0.4	0.1	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.51A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	34,076	32,443	9,220	8,761	3,952	3,849
AGE						
12-17	2,524	2,543	1,653	1,649	664	690
18-25	5,957	5,650	2,830	2,696	1,062	1,050
26 or Older	25,596	24,250	4,737	4,415	2,225	2,108
GENDER						
Male	17,764	17,008	4,539	4,217	2,034	1,926
Female	16,312	15,435	4,682	4,544	1,918	1,923
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	27,615	26,263	7,066	6,725	3,035	2,937
Black Only	2,675	2,463	832	702	356	315
American Indian or Alaska Native Only	299	244	*	63	40	43
Native Hawaiian or Other Pacific Islander	*	66	12	22	3	19
Asian Only	538	566	139	178	66	80
More Than One Race	310	341	104	94	53	43
Hispanic	2,571	2,499	971	976	399	413

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.51B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15.4 ^a	14.5	4.2	3.9	1.8	1.7
AGE						
12-17	10.9	10.9	7.1	7.1	2.9	3.0
18-25	20.9 ^b	19.5	9.9	9.3	3.7	3.6
26 or Older	15.1	14.2	2.8	2.6	1.3	1.2
GENDER						
Male	16.7	15.8	4.3	3.9	1.9	1.8
Female	14.2	13.3	4.1	3.9	1.7	1.7
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	17.0	16.1	4.4	4.1	1.9	1.8
Black Only	10.8	9.7	3.3	2.8	1.4	1.2
American Indian or Alaska Native Only	25.9	22.3	*	5.8	3.5	3.9
Native Hawaiian or Other Pacific Islander	*	12.1	1.5	4.1	0.3	3.5
Asian Only	7.4	7.4	1.9	2.3	0.9	1.1
More Than One Race	15.9	18.3	5.4	5.1	2.7	2.3
Hispanic	11.2	10.5	4.2	4.1	1.7	1.7

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.52A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	2,524	2,543	1,653	1,649	664	690
GENDER						
Male	1,196	1,238	733	778	308	317
Female	1,328	1,305	920	871	356	373
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	1,738	1,800	1,158	1,199	456	478
Black Only	312	288	211	183	96	89
American Indian or Alaska Native Only	20	19	14	9	9	5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	55	69	26	43	5 ^a	18
More Than One Race	43	30	27	14	8	7
Hispanic	350	327	213	196	88	91
GENDER/RACE/HISPANIC ORIGIN						
Male - White	825	852	511	553	204	217
Female - White	913	949	647	647	252	260
Male - Black	151	146	96	86	56	37
Female - Black	161	142	115	97	40	52
Male - Hispanic	163	168	93	102	36	47
Female - Hispanic	187	159	120	94	51	43

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.52B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	10.9	10.9	7.1	7.1	2.9	3.0
GENDER						
Male	10.1	10.4	6.2	6.5	2.6	2.7
Female	11.7	11.4	8.1	7.6	3.1	3.3
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	11.4	11.8	7.6	7.8	3.0	3.1
Black Only	9.7	8.6	6.6	5.5	3.0	2.7
American Indian or Alaska Native Only	15.1	13.3	10.4	6.1	7.1	3.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	6.1	8.3	2.9	5.2	0.6 ^a	2.2
More Than One Race	11.6	8.3	7.3	3.8	2.3	1.9
Hispanic	10.9	9.9	6.6	5.9	2.7	2.7
GENDER/RACE/HISPANIC ORIGIN						
Male - White	10.5	10.8	6.5	7.0	2.6	2.8
Female - White	12.3	12.7	8.7	8.7	3.4	3.5
Male - Black	9.2	8.6	5.9	5.0	3.4	2.2
Female - Black	10.1	8.7	7.3	6.0	2.5	3.2
Male - Hispanic	9.9	9.8	5.6	5.9	2.2	2.8
Female - Hispanic	11.9	10.0	7.6	5.9	3.3	2.7

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.53A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	5,957	5,650	2,830	2,696	1,062	1,050
GENDER						
Male	3,186	3,095	1,517	1,494	643	608
Female	2,770 ^a	2,555	1,313	1,202	419	442
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	4,578	4,468	2,200	2,130	821	829
Black Only	409	366	182	178	62	74
American Indian or Alaska Native Only	28	*	15	*	2	*
Native Hawaiian or Other Pacific Islander	*	*	*	5	*	*
Asian Only	127	101	64	48	23	13
More Than One Race	98	89	57	49	26	23
Hispanic	699 ^b	560	309	268	127	98
ADULT EDUCATION						
< High School	1,335	1,282	670	633	313	285
High School Graduate	2,150	2,089	1,036	1,023	413	375
Some College	1,939	1,742	907	844	293	317
College Graduate	533	537	218	196	43	74
CURRENT EMPLOYMENT						
Full-Time	3,228	3,094	1,459	1,390	523	515
Part-Time	1,370	1,299	661	633	260	236
Unemployed	366	433	202	242	110	111
Other ¹	992 ^a	824	508	432	169	188

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.53B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	20.9 ^b	19.5	9.9	9.3	3.7	3.6
GENDER						
Male	22.5	21.5	10.7	10.4	4.5	4.2
Female	19.3 ^b	17.5	9.2	8.2	2.9	3.0
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	24.4	23.3	11.7	11.1	4.4	4.3
Black Only	10.6	9.3	4.7	4.5	1.6	1.9
American Indian or Alaska Native Only	20.3	*	10.9	*	1.7	*
Native Hawaiian or Other Pacific Islander	*	*	*	5.1	*	*
Asian Only	11.7	10.0	5.9	4.8	2.1	1.3
More Than One Race	28.5	29.6	16.4	16.2	7.5	7.8
Hispanic	16.6 ^b	12.9	7.3	6.2	3.0	2.3
ADULT EDUCATION						
< High School	22.8	21.0	11.4	10.4	5.4	4.7
High School Graduate	21.0	20.1	10.1	9.8	4.0	3.6
Some College	21.3 ^a	19.2	10.0	9.3	3.2	3.5
College Graduate	16.3	15.7	6.7	5.7	1.3	2.2
CURRENT EMPLOYMENT						
Full-Time	22.0 ^a	20.2	9.9	9.1	3.6	3.4
Part-Time	20.1	19.4	9.7	9.5	3.8	3.5
Unemployed	22.0 ^a	26.7	12.2	14.9	6.6	6.8
Other ¹	18.7 ^b	15.4	9.6	8.1	3.2	3.5

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.54A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	25,596	24,250	4,737	4,415	2,225	2,108
GENDER						
Male	13,383	12,675	2,289	1,945	1,083	1,000
Female	12,213	11,575	2,448	2,471	1,142	1,108
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	21,299	19,994	3,707	3,395	1,758	1,630
Black Only	1,953	1,809	439	341	198	152
American Indian or Alaska Native Only	*	*	*	36	*	29
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	356	396	48	87	38	49
More Than One Race	169	221	21	32	19	12
Hispanic	1,522	1,612	449	512	184	224
ADULT EDUCATION						
< High School	3,322 ^a	2,636	853	769	421	327
High School Graduate	8,085	8,157	1,379	1,639	646	889
Some College	7,235	6,602	1,300 ^a	918	544	436
College Graduate	6,954	6,855	1,206	1,089	614	457
CURRENT EMPLOYMENT						
Full-Time	17,675	17,061	2,895	2,928	1,260	1,355
Part-Time	2,708	2,330	524	419	262	230
Unemployed	846	830	312 ^a	136	158	53
Other ¹	4,366	4,029	1,006	933	545	470

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.54B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15.1	14.2	2.8	2.6	1.3	1.2
GENDER						
Male	16.7	15.7	2.9	2.4	1.3	1.2
Female	13.7	12.9	2.7	2.7	1.3	1.2
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	16.6	15.6	2.9	2.6	1.4	1.3
Black Only	11.0	10.0	2.5	1.9	1.1	0.8
American Indian or Alaska Native Only	*	*	*	4.6	*	3.8
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	6.8	6.9	0.9	1.5	0.7	0.8
More Than One Race	13.6	18.5	1.7	2.7	1.5	1.0
Hispanic	9.8	10.0	2.9	3.2	1.2	1.4
ADULT EDUCATION						
< High School	10.8	9.0	2.8	2.6	1.4	1.1
High School Graduate	14.0	14.4	2.4	2.9	1.1	1.6
Some College	18.6	16.8	3.3 ^a	2.3	1.4	1.1
College Graduate	16.4	15.0	2.8	2.4	1.4	1.0
CURRENT EMPLOYMENT						
Full-Time	18.0	16.9	2.9	2.9	1.3	1.3
Part-Time	16.0	14.0	3.1	2.5	1.5	1.4
Unemployed	24.1	27.2	8.9 ^a	4.4	4.5	1.7
Other ¹	8.6	8.0	2.0	1.8	1.1	0.9

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.55A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	31,552	29,900	7,567	7,112	3,287	3,158
GENDER						
Male	16,569	15,770	3,806	3,439	1,726	1,609
Female	14,983	14,130	3,762	3,673	1,562	1,550
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	25,877	24,463	5,908	5,525	2,579	2,459
Black Only	2,363	2,175	621	519	260	226
American Indian or Alaska Native Only	279	225	*	55	31	38
Native Hawaiian or Other Pacific Islander	*	58	7	18	1	*
Asian Only	483	497	113	135	61	62
More Than One Race	267	310	77	81	45	36
Hispanic	2,221	2,172	758	780	312	322
ADULT EDUCATION						
< High School	4,657 ^a	3,918	1,522	1,402	735	612
High School Graduate	10,235	10,247	2,414	2,662	1,059	1,264
Some College	9,174	8,344	2,206 ^a	1,762	837	753
College Graduate	7,486	7,392	1,424	1,285	657	530
CURRENT EMPLOYMENT						
Full-Time	20,902	20,155	4,354	4,318	1,783	1,871
Part-Time	4,078	3,629	1,185	1,051	522	465
Unemployed	1,213	1,263	515	377	268	164
Other ¹	5,359	4,853	1,514	1,366	714	658

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.55B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15.9 ^a	15.0	3.8	3.6	1.7	1.6
GENDER						
Male	17.6	16.5	4.0	3.6	1.8	1.7
Female	14.5	13.5	3.6	3.5	1.5	1.5
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	17.6 ^a	16.6	4.0	3.7	1.8	1.7
Black Only	10.9	9.9	2.9	2.4	1.2	1.0
American Indian or Alaska Native Only	27.3	23.7	*	5.8	3.0	4.0
Native Hawaiian or Other Pacific Islander	*	12.2	0.9	3.7	0.1	*
Asian Only	7.6	7.3	1.8	2.0	1.0	0.9
More Than One Race	16.9	20.7	4.9	5.4	2.8	2.4
Hispanic	11.2	10.6	3.8	3.8	1.6	1.6
ADULT EDUCATION						
< High School	12.8	11.1	4.2	4.0	2.0	1.7
High School Graduate	15.1	15.3	3.6	4.0	1.6	1.9
Some College	19.1 ^a	17.3	4.6 ^a	3.6	1.7	1.6
College Graduate	16.4	15.0	3.1	2.6	1.4	1.1
CURRENT EMPLOYMENT						
Full-Time	18.5	17.4	3.9	3.7	1.6	1.6
Part-Time	17.2	15.5	5.0	4.5	2.2	2.0
Unemployed	23.5	27.0	10.0	8.1	5.2	3.5
Other ¹	9.6	8.7	2.7	2.4	1.3	1.2

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.56A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	53,268	52,605	13,444	13,052	5,981	5,711
AGE						
12-17	4,248	4,228	2,698	2,631	1,045	1,074
18-25	9,476	9,247	4,460	4,295	1,729	1,711
26 or Older	39,545	39,130	6,285	6,125	3,208	2,926
GENDER						
Male	28,729	29,021	7,240	6,926	3,325	3,112
Female	24,538	23,584	6,204	6,126	2,656	2,599
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	42,242	42,172	10,033	9,864	4,459	4,232
Black Only	4,653	4,097	1,275	1,110	586	529
American Indian or Alaska Native Only	426	419	140	117	55	52
Native Hawaiian or Other Pacific Islander	*	*	20	33	4	23
Asian Only	837	800	223	249	88	118
More Than One Race	498	576	165	155	69	69
Hispanic	4,485	4,399	1,588	1,523	721	688

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.56B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	24.1	23.6	6.1	5.8	2.7	2.6
AGE						
12-17	18.3	18.1	11.6	11.3	4.5	4.6
18-25	33.3 ^a	31.9	15.7	14.8	6.1	5.9
26 or Older	23.3	22.9	3.7	3.6	1.9	1.7
GENDER						
Male	27.0	27.0	6.8	6.5	3.1	2.9
Female	21.4	20.3	5.4	5.3	2.3	2.2
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	26.1	25.9	6.2	6.1	2.8	2.6
Black Only	18.7 ^a	16.1	5.1	4.4	2.4	2.1
American Indian or Alaska Native Only	37.0	38.4	12.2	10.8	4.8	4.8
Native Hawaiian or Other Pacific Islander	*	*	2.5	6.0	0.5	4.3
Asian Only	11.5	10.5	3.1	3.3	1.2	1.6
More Than One Race	25.6	30.9	8.5	8.3	3.5	3.7
Hispanic	19.5	18.4	6.9	6.4	3.1	2.9

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.57A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	4,248	4,228	2,698	2,631	1,045	1,074
GENDER						
Male	2,142	2,158	1,299	1,278	485	509
Female	2,106	2,070	1,399	1,353	559	565
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	2,954	2,928	1,926	1,865	724	735
Black Only	436	450	259	252	108	115
American Indian or Alaska Native Only	43	44	27	24	13	11
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	97	115	54	74	15 ^a	36
More Than One Race	68	66	42	34	12	10
Hispanic	639	616	384	378	171	163
GENDER/RACE/HISPANIC ORIGIN						
Male - White	1,499	1,484	930	893	327	348
Female - White	1,455	1,444	996	972	396	387
Male - Black	221	222	123	120	61	47
Female - Black	215	228	135	131	47	68
Male - Hispanic	318	336	189	207	79	90
Female - Hispanic	321	280	195	171	93	73

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.57B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	18.3	18.1	11.6	11.3	4.5	4.6
GENDER						
Male	18.0	18.0	10.9	10.7	4.1	4.3
Female	18.6	18.1	12.4	11.9	4.9	4.9
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	19.3	19.1	12.6	12.2	4.7	4.8
Black Only	13.5	13.5	8.0	7.5	3.3	3.4
American Indian or Alaska Native Only	32.5	30.2	20.3	16.5	9.8	7.8
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	10.8	13.8	6.0	8.9	1.7 ^b	4.3
More Than One Race	18.5	18.2	11.6	9.3	3.3	2.7
Hispanic	19.8	18.6	11.9	11.4	5.3	4.9
GENDER/RACE/HISPANIC ORIGIN						
Male - White	19.1	18.9	11.8	11.4	4.2	4.4
Female - White	19.5	19.4	13.4	13.0	5.3	5.2
Male - Black	13.5	13.0	7.5	7.1	3.7	2.7
Female - Black	13.6	14.0	8.5	8.0	2.9	4.2
Male - Hispanic	19.3	19.5	11.5	12.0	4.8	5.2
Female - Hispanic	20.4	17.6	12.4	10.7	5.9	4.6

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.58A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	9,476	9,247	4,460	4,295	1,729	1,711
GENDER						
Male	5,199	5,128	2,555	2,455	1,064	999
Female	4,277	4,120	1,906	1,841	665	712
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	7,175	7,151	3,418	3,327	1,327	1,318
Black Only	626	601	267	296	101	114
American Indian or Alaska Native Only	59	*	25	35	4	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	229 ^a	158	113	75	35	22
More Than One Race	150	132	81	74	38	30
Hispanic	1,204	1,087	548	478	221	208
ADULT EDUCATION						
< High School	2,035	2,038	1,028	986	482	440
High School Graduate	3,379	3,309	1,608	1,530	669	594
Some College	3,100	2,940	1,461	1,419	483	554
College Graduate	962	961	364	361	96	122
CURRENT EMPLOYMENT						
Full-Time	5,236	5,136	2,279	2,216	877	848
Part-Time	2,159	2,108	1,114	1,047	417	402
Unemployed	565	644	316	350	167	171
Other ¹	1,516	1,360	752	682	268	289

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.58B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	33.3 ^a	31.9	15.7	14.8	6.1	5.9
GENDER						
Male	36.8	35.6	18.1	17.0	7.5	6.9
Female	29.9 ^a	28.3	13.3	12.6	4.6	4.9
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	38.3	37.4	18.2	17.4	7.1	6.9
Black Only	16.2	15.3	6.9	7.5	2.6	2.9
American Indian or Alaska Native Only	43.2	*	18.6	19.7	3.0	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	21.1	15.8	10.4	7.4	3.2	2.2
More Than One Race	43.3	43.9	23.4	24.6	11.0	9.9
Hispanic	28.6 ^a	25.1	13.0	11.0	5.3	4.8
ADULT EDUCATION						
< High School	34.8	33.4	17.6	16.1	8.2	7.2
High School Graduate	33.0	31.8	15.7	14.7	6.5	5.7
Some College	34.0	32.4	16.0	15.7	5.3	6.1
College Graduate	29.4	28.1	11.1	10.6	2.9	3.6
CURRENT EMPLOYMENT						
Full-Time	35.6 ^a	33.5	15.5	14.4	6.0	5.5
Part-Time	31.7	31.5	16.3	15.7	6.1	6.0
Unemployed	34.0 ^a	39.7	19.0	21.5	10.0	10.6
Other ¹	28.6 ^a	25.5	14.2	12.8	5.0	5.4

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.59A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	39,545	39,130	6,285	6,125	3,208	2,926
GENDER						
Male	21,389	21,735	3,386	3,193	1,776	1,604
Female	18,156	17,395	2,899	2,933	1,432	1,322
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	32,112	32,092	4,690	4,672	2,408	2,179
Black Only	3,591	3,046	749	562	377	299
American Indian or Alaska Native Only	*	*	*	59	38	30
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	511	527	56	101	38	60
More Than One Race	280	*	42	47	19	29
Hispanic	2,643	2,695	656	668	328	317
ADULT EDUCATION						
< High School	4,881	4,277	1,219	985	664	478
High School Graduate	12,477	12,712	1,780	2,151	973	1,131
Some College	10,984	10,620	1,727	1,350	810	609
College Graduate	11,204	11,521	1,558	1,639	761	709
CURRENT EMPLOYMENT						
Full-Time	28,063	28,464	4,051	4,150	1,987	1,878
Part-Time	4,038	3,760	687	606	334	321
Unemployed	1,351	1,118	409	242	254	122
Other ¹	6,093	5,789	1,138	1,128	633	605

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.59B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	23.3	22.9	3.7	3.6	1.9	1.7
GENDER						
Male	26.7	26.8	4.2	3.9	2.2	2.0
Female	20.3	19.3	3.2	3.3	1.6	1.5
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	25.1	25.0	3.7	3.6	1.9	1.7
Black Only	20.2 ^a	16.8	4.2	3.1	2.1	1.6
American Indian or Alaska Native Only	*	*	*	7.6	4.3	3.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	9.7	9.1	1.1	1.7	0.7	1.0
More Than One Race	22.7	*	3.4	4.0	1.5	2.4
Hispanic	17.0	16.6	4.2	4.1	2.1	2.0
ADULT EDUCATION						
< High School	15.9	14.6	4.0	3.4	2.2	1.6
High School Graduate	21.7	22.4	3.1	3.8	1.7	2.0
Some College	28.3	27.1	4.4 ^a	3.4	2.1	1.6
College Graduate	26.4	25.2	3.7	3.6	1.8	1.6
CURRENT EMPLOYMENT						
Full-Time	28.6	28.3	4.1	4.1	2.0	1.9
Part-Time	23.8	22.5	4.1	3.6	2.0	1.9
Unemployed	38.5	36.6	11.7	7.9	7.3	4.0
Other ¹	12.0	11.5	2.2	2.2	1.2	1.2

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.60A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	49,020	48,377	10,745	10,421	4,937	4,637
GENDER						
Male	26,588	26,863	5,941	5,648	2,840	2,602
Female	22,433	21,514	4,805	4,773	2,097	2,034
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	39,287	39,244	8,108	7,999	3,735	3,497
Black Only	4,217	3,647	1,016	858	478	413
American Indian or Alaska Native Only	383	375	*	94	42	41
Native Hawaiian or Other Pacific Islander	*	*	12	27	2	20
Asian Only	740	685	169	176	73	82
More Than One Race	430	509	123	121	57	59
Hispanic	3,847	3,783	1,204	1,146	550	524
ADULT EDUCATION						
< High School	6,916	6,315	2,247	1,971	1,146	918
High School Graduate	15,855	16,021	3,388	3,681	1,641	1,725
Some College	14,083	13,560	3,187	2,769	1,293	1,163
College Graduate	12,166	12,482	1,923	2,000	857	831
CURRENT EMPLOYMENT						
Full-Time	33,299	33,599	6,330	6,366	2,864	2,726
Part-Time	6,197	5,868	1,800	1,653	751	724
Unemployed	1,916	1,762	725	591	421	293
Other ¹	7,609	7,148	1,890	1,810	900	894

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.60B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	24.8	24.2	5.4	5.2	2.5	2.3
GENDER						
Male	28.2	28.2	6.3	5.9	3.0	2.7
Female	21.7	20.6	4.6	4.6	2.0	1.9
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	26.8	26.6	5.5	5.4	2.5	2.4
Black Only	19.5 ^a	16.5	4.7	3.9	2.2	1.9
American Indian or Alaska Native Only	37.6	39.7	*	9.9	4.1	4.3
Native Hawaiian or Other Pacific Islander	*	*	1.6	5.7	0.3	4.2
Asian Only	11.6	10.1	2.7	2.6	1.1	1.2
More Than One Race	27.2	34.0	7.8	8.1	3.6	3.9
Hispanic	19.4	18.4	6.1	5.6	2.8	2.6
ADULT EDUCATION						
< High School	19.0	17.9	6.2	5.6	3.1	2.6
High School Graduate	23.4	23.9	5.0	5.5	2.4	2.6
Some College	29.4	28.1	6.7 ^a	5.7	2.7	2.4
College Graduate	26.6	25.4	4.2	4.1	1.9	1.7
CURRENT EMPLOYMENT						
Full-Time	29.5	28.9	5.6	5.5	2.5	2.3
Part-Time	26.1	25.1	7.6	7.1	3.2	3.1
Unemployed	37.1	37.7	14.0	12.6	8.1	6.3
Other ¹	13.6	12.8	3.4	3.2	1.6	1.6

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.61A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	67,807	18,411	10,231
Black	9,186	3,028	1,734
American Indian or Alaska Native	587	213	129
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	207	52	14
Filipino	372	92	41
Japanese	185	52	32
Asian Indian	205	65	33
Korean	185	60	43
Vietnamese	119	45	29
HISPANIC			
Mexican	4,627	1,561	832
Puerto Rican	1,059	418	259
Central or South American	835	264	133
Cuban	270	92	42

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.61B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	41.7	11.3	6.3
Black	36.6	12.0	6.9
American Indian or Alaska Native	52.4	19.0	11.5
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	14.5	3.6	1.0
Filipino	24.7	6.1	2.7
Japanese	29.3	8.3	5.0
Asian Indian	13.3	4.2	2.1
Korean	29.2	9.5	6.9
Vietnamese	17.8	6.7	4.3
HISPANIC			
Mexican	30.5	10.3	5.5
Puerto Rican	41.3	16.3	10.1
Central or South American	25.8	8.2	4.1
Cuban	23.8	8.1	3.7

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.62A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	4,246	3,049	1,521
Black	826	549	290
American Indian or Alaska Native	70	49	31
Native Hawaiian	*	*	*
Other Pacific Islander	8	4	2
Chinese	19	15	8
Filipino	41	24	12
Japanese	6	4	1
Asian Indian	19	14	7
Korean	13	10	7
Vietnamese	23	*	8
HISPANIC			
Mexican	654	455	226
Puerto Rican	108	73	43
Central or South American	101	57	27
Cuban	25	19	12

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.62B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	27.7	19.9	9.9
Black	25.2	16.7	8.9
American Indian or Alaska Native	50.9	35.5	22.2
Native Hawaiian	*	*	*
Other Pacific Islander	17.3	9.6	5.0
Chinese	14.6	11.1	6.2
Filipino	25.0	14.7	7.0
Japanese	15.3	10.2	1.7
Asian Indian	13.5	9.3	5.2
Korean	15.4	11.4	7.5
Vietnamese	17.8	*	6.0
HISPANIC			
Mexican	29.2	20.3	10.1
Puerto Rican	27.6	18.7	10.9
Central or South American	25.8	14.7	6.9
Cuban	26.2	20.3	12.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.63A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	10,757	5,931	3,332
Black	1,751	1,002	584
American Indian or Alaska Native	107	54	27
Native Hawaiian	*	*	*
Other Pacific Islander	*	14	*
Chinese	45	30	5
Filipino	70	36	22
Japanese	*	*	9
Asian Indian	41	20	11
Korean	37	24	*
Vietnamese	25	11	9
HISPANIC			
Mexican	1,178	564	311
Puerto Rican	237	129	91
Central or South American	189	107	58
Cuban	44	27	*

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.63B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	56.8	31.3	17.6
Black	45.0	25.8	15.0
American Indian or Alaska Native	68.3	34.3	17.0
Native Hawaiian	*	*	*
Other Pacific Islander	*	19.6	*
Chinese	26.1	17.0	2.7
Filipino	38.8	20.0	12.0
Japanese	*	*	11.8
Asian Indian	19.4	9.6	5.2
Korean	42.0	27.1	*
Vietnamese	20.8	9.1	7.5
HISPANIC			
Mexican	38.9	18.7	10.3
Puerto Rican	54.2	29.4	20.7
Central or South American	36.1	20.4	11.1
Cuban	47.8	28.7	*

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.64A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	52,804	9,432	5,378
Black	6,610	1,476	859
American Indian or Alaska Native	410	111	72
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	143	7	1
Filipino	261	32	8
Japanese	148	29	22
Asian Indian	144	32	*
Korean	*	*	*
Vietnamese	*	16	12
HISPANIC			
Mexican	2,796	543	295
Puerto Rican	714	216	125
Central or South American	546	100	48
Cuban	201	46	15

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.64B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	41.2	7.4	4.2
Black	36.8	8.2	4.8
American Indian or Alaska Native	49.6	13.4	8.7
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	12.7	0.7	0.1
Filipino	22.5	2.7	0.7
Japanese	28.5	5.5	4.3
Asian Indian	12.1	2.6	*
Korean	*	*	*
Vietnamese	*	3.9	2.8
HISPANIC			
Mexican	28.3	5.5	3.0
Puerto Rican	41.1	12.4	7.2
Central or South American	23.5	4.3	2.1
Cuban	21.3	4.8	1.6

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.65A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	63,561	15,362	8,710
Black	8,361	2,479	1,443
American Indian or Alaska Native	517	164	98
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	188	37	6
Filipino	330	68	29
Japanese	179	49	31
Asian Indian	186	52	25
Korean	*	50	37
Vietnamese	96	27	21
HISPANIC			
Mexican	3,974	1,107	606
Puerto Rican	951	345	216
Central or South American	734	207	106
Cuban	245	72	30

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.65B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	43.2	10.4	5.9
Black	38.3	11.3	6.6
American Indian or Alaska Native	52.6	16.7	10.0
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	14.5	2.9	0.5
Filipino	24.7	5.1	2.2
Japanese	30.2	8.2	5.2
Asian Indian	13.2	3.7	1.8
Korean	*	9.2	6.8
Vietnamese	17.8	5.0	3.8
HISPANIC			
Mexican	30.8	8.6	4.7
Puerto Rican	43.7	15.9	9.9
Central or South American	25.8	7.3	3.7
Cuban	23.6	6.9	2.9

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

10703

Table 1.66A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	60,206	14,055	7,819
Black	7,912	2,332	1,393
American Indian or Alaska Native	471	154	95
Native Hawaiian	*	*	8
Other Pacific Islander	*	*	*
Chinese	129	35	7
Filipino	239	76	33
Japanese	163	31	18
Asian Indian	111	25	14
Korean	130	36	26
Vietnamese	88	30	25
HISPANIC			
Mexican	3,809	1,062	566
Puerto Rican	824	281	180
Central or South American	632	179	83
Cuban	213	55	31

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.66B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	37.0	8.6	4.8
Black	31.5	9.3	5.5
American Indian or Alaska Native	42.0	13.8	8.5
Native Hawaiian	*	*	6.7
Other Pacific Islander	*	*	*
Chinese	9.0	2.4	0.5
Filipino	15.9	5.1	2.2
Japanese	26.0	5.0	2.9
Asian Indian	7.2	1.6	0.9
Korean	20.5	5.7	4.2
Vietnamese	13.1	4.4	3.7
HISPANIC			
Mexican	25.1	7.0	3.7
Puerto Rican	32.1	10.9	7.0
Central or South American	19.5	5.5	2.6
Cuban	18.8	4.8	2.7

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

10703

Table 1.67A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	2,945	2,236	1,163
Black	499	350	197
American Indian or Alaska Native	57	40	24
Native Hawaiian	*	*	*
Other Pacific Islander	3	2	1
Chinese	9	9	3
Filipino	22	16	8
Japanese	3	2	1
Asian Indian	8	6	5
Korean	9	7	*
Vietnamese	*	*	*
HISPANIC			
Mexican	465	326	153
Puerto Rican	72	56	32
Central or South American	53	35	15
Cuban	19	13	8

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.67B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	19.2	14.6	7.6
Black	15.2	10.7	6.0
American Indian or Alaska Native	41.7	29.1	17.3
Native Hawaiian	*	*	*
Other Pacific Islander	6.4	4.8	3.0
Chinese	7.2	6.8	2.3
Filipino	13.6	9.8	5.0
Japanese	8.6	4.3	1.4
Asian Indian	5.7	4.4	3.1
Korean	10.8	8.3	*
Vietnamese	*	*	*
HISPANIC			
Mexican	20.8	14.6	6.8
Puerto Rican	18.4	14.4	8.2
Central or South American	13.4	8.9	3.8
Cuban	19.3	13.9	8.7

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

10703

Table 1.68A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	9,709	5,053	2,897
Black	1,550	887	530
American Indian or Alaska Native	95	42	24
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	36	21	3
Filipino	62	30	18
Japanese	*	13	7
Asian Indian	28	14	8
Korean	31	18	*
Vietnamese	16	9	7
HISPANIC			
Mexican	982	414	242
Puerto Rican	205	107	78
Central or South American	159	87	46
Cuban	40	*	*

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.68B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	51.2	26.7	15.3
Black	39.9	22.8	13.6
American Indian or Alaska Native	60.9	27.0	15.1
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	20.6	11.9	1.9
Filipino	34.3	16.5	9.9
Japanese	*	18.0	9.8
Asian Indian	13.0	6.5	4.0
Korean	35.3	20.7	*
Vietnamese	13.2	7.1	5.7
HISPANIC			
Mexican	32.5	13.7	8.0
Puerto Rican	46.7	24.5	17.9
Central or South American	30.4	16.7	8.8
Cuban	43.0	*	*

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

10703

Table 1.69A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	47,552	6,765	3,759
Black	5,863	1,094	665
American Indian or Alaska Native	318	72	47
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	84	5	1
Filipino	155	30	8
Japanese	136	16	10
Asian Indian	76	5	1
Korean	*	*	*
Vietnamese	*	12	12
HISPANIC			
Mexican	2,362	321	170
Puerto Rican	547	117	70
Central or South American	421	57	23
Cuban	155	18	9

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.69B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	37.1	5.3	2.9
Black	32.6	6.1	3.7
American Indian or Alaska Native	38.5	8.7	5.7
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	7.5	0.5	0.1
Filipino	13.3	2.6	0.6
Japanese	26.3	3.1	2.0
Asian Indian	6.4	0.4	0.1
Korean	*	*	*
Vietnamese	*	2.8	2.8
HISPANIC			
Mexican	23.9	3.2	1.7
Puerto Rican	31.5	6.7	4.0
Central or South American	18.1	2.5	1.0
Cuban	16.4	1.9	0.9

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.70A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	57,260	11,819	6,656
Black	7,412	1,982	1,196
American Indian or Alaska Native	414	114	71
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	120	26	5
Filipino	216	60	25
Japanese	160	30	18
Asian Indian	103	19	10
Korean	*	29	21
Vietnamese	77	20	18
HISPANIC			
Mexican	3,344	735	412
Puerto Rican	752	224	148
Central or South American	579	145	68
Cuban	194	41	23

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.70B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	38.9	8.0	4.5
Black	33.9	9.1	5.5
American Indian or Alaska Native	42.1	11.6	7.2
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	9.2	2.0	0.3
Filipino	16.1	4.5	1.9
Japanese	27.1	5.0	3.0
Asian Indian	7.4	1.3	0.7
Korean	*	5.3	3.9
Vietnamese	14.2	3.7	3.4
HISPANIC			
Mexican	25.9	5.7	3.2
Puerto Rican	34.6	10.3	6.8
Central or South American	20.4	5.1	2.4
Cuban	18.7	4.0	2.2

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.71A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	87,734	86,931	25,402	24,535	13,829	14,027
GEOGRAPHIC DIVISION						
Northeast	17,564	16,604	5,068	4,956	2,951	2,820
New England	5,506	5,357	1,595	1,814	1,092	1,124
Middle Atlantic	12,058	11,247	3,473	3,142	1,858	1,696
Midwest	21,124	19,910	5,769 ^a	5,245	3,138	2,939
East North Central	14,854	14,449	4,111	3,922	2,298	2,301
West North Central	6,270 ^a	5,460	1,657 ^a	1,323	840 ^a	639
South	28,999	29,152	8,166	7,915	4,127	4,355
South Atlantic	15,714	15,674	4,422	4,409	2,279	2,381
East South Central	4,754	4,972	1,454	1,358	668	781
West South Central	8,531	8,506	2,290	2,148	1,179	1,193
West	20,047	21,266	6,399	6,418	3,614	3,913
Mountain	6,233	6,264	1,848	1,849	1,002	1,035
Pacific	13,813	15,002	4,551	4,569	2,611	2,878
COUNTY TYPE						
Large Metro	45,087	43,938	13,284	12,421	7,371	7,126
Small Metro	27,491	27,543	7,968	8,014	4,339	4,583
250K - 1 Mil. Pop.	20,826	20,650	6,036	5,884	3,307	3,281
<250K Pop.	6,665	6,893	1,932	2,130	1,031	1,303
Nonmetro	15,157	15,451	4,150	4,099	2,120	2,317
Urbanized	5,199	5,143	1,464	1,436	802	908
Less Urbanized	8,283	8,747	2,281	2,265	1,089	1,203
Completely Rural	1,674	1,560	405	398	229	206

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.71B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	39.7	38.9	11.5	11.0	6.3	6.3
GEOGRAPHIC DIVISION						
Northeast	41.2 ^a	38.8	11.9	11.6	6.9	6.6
New England	49.4	47.8	14.3	16.2	9.8	10.0
Middle Atlantic	38.2	35.6	11.0	9.9	5.9	5.4
Midwest	41.0 ^b	38.4	11.2 ^a	10.1	6.1	5.7
East North Central	41.1	39.8	11.4	10.8	6.4	6.3
West North Central	40.6 ^b	35.1	10.7 ^a	8.5	5.4 ^a	4.1
South	37.0	36.7	10.4	10.0	5.3	5.5
South Atlantic	38.8	38.2	10.9	10.7	5.6	5.8
East South Central	34.6	35.8	10.6	9.8	4.9	5.6
West South Central	35.2	34.7	9.5	8.8	4.9	4.9
West	41.4	43.3	13.2	13.1	7.5	8.0
Mountain	44.6	43.8	13.2	12.9	7.2	7.2
Pacific	40.1 ^a	43.1	13.2	13.1	7.6	8.3
COUNTY TYPE						
Large Metro	40.8	40.3	12.0	11.4	6.7	6.5
Small Metro	41.2	40.1	12.0	11.7	6.5	6.7
250K - 1 Mil. Pop.	41.7	40.5	12.1	11.5	6.6	6.4
<250K Pop.	40.0	39.1	11.6	12.1	6.2	7.4
Nonmetro	34.5	33.9	9.4	9.0	4.8	5.1
Urbanized	39.5	38.7	11.1	10.8	6.1	6.8
Less Urbanized	33.0	32.4	9.1	8.4	4.3	4.5
Completely Rural	29.2	29.6	7.1	7.5	4.0	3.9

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.72A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	6,415	6,288	4,584	4,357	2,265	2,264
GEOGRAPHIC DIVISION						
Northeast	1,095	1,123	826	810	411	403
New England	359	350	273	272	155	154
Middle Atlantic	736	773	553	537	257	249
Midwest	1,506	1,434	1,070	979	549	500
East North Central	1,066	1,008	762	693	398	351
West North Central	441	426	308	286	151	149
South	2,196	2,209	1,535	1,521	733	784
South Atlantic	1,050	1,104	735	757	348	376
East South Central	384	404	262	283	114	151
West South Central	762	701	537	481	271	256
West	1,618	1,522	1,152	1,049	572	577
Mountain	518	495	350	341	174	181
Pacific	1,100	1,028	803 ^a	707	398	396
COUNTY TYPE						
Large Metro	3,138	2,941	2,270 ^a	2,045	1,106	1,072
Small Metro	2,018	2,056	1,440	1,437	736	755
250K - 1 Mil. Pop.	1,534	1,493	1,103	1,059	576	556
<250K Pop.	483	563	337	379	160	199
Nonmetro	1,260	1,292	874	874	424	437
Urbanized	427	391	292	281	147	153
Less Urbanized	693	731	486	484	226	231
Completely Rural	140	170	96	109	51	54

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.72B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	27.6	26.9	19.8 ^a	18.6	9.8	9.7
GEOGRAPHIC DIVISION						
Northeast	26.4	26.7	19.9	19.3	9.9	9.6
New England	32.9	31.6	25.0	24.6	14.2	13.9
Middle Atlantic	24.0	25.0	18.1	17.4	8.4	8.1
Midwest	27.5	26.3	19.6	17.9	10.0	9.2
East North Central	28.2	26.8	20.2	18.4	10.5	9.3
West North Central	26.0	25.2	18.2	16.9	8.9	8.9
South	26.6	26.6	18.6	18.3	8.9	9.4
South Atlantic	26.2	27.1	18.3	18.6	8.7	9.2
East South Central	27.1	28.4	18.5	19.9	8.0	10.6
West South Central	27.1	24.8	19.1	17.0	9.6	9.1
West	30.3 ^a	28.2	21.6 ^b	19.4	10.7	10.7
Mountain	32.2	30.5	21.8	21.0	10.8	11.1
Pacific	29.5 ^a	27.2	21.5 ^b	18.8	10.7	10.5
COUNTY TYPE						
Large Metro	27.1	25.9	19.6 ^a	18.0	9.6	9.4
Small Metro	28.9	28.5	20.6	20.0	10.5	10.5
250K - 1 Mil. Pop.	29.0	27.7	20.9	19.6	10.9	10.3
<250K Pop.	28.3	31.1	19.8	20.9	9.4	11.0
Nonmetro	27.1	26.9	18.8	18.2	9.1	9.1
Urbanized	28.5	29.5	19.5	21.2	9.8	11.5
Less Urbanized	26.6	25.4	18.7	16.8	8.7	8.0
Completely Rural	25.3	28.5	17.3	18.2	9.2	9.0

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.73A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	14,975	14,846	8,287	8,088	4,658	4,599
GEOGRAPHIC DIVISION						
Northeast	2,807	2,771	1,712	1,675	990	975
New England	822	853	507	538	323	321
Middle Atlantic	1,986	1,918	1,205	1,137	667	653
Midwest	3,707	3,668	2,062	1,895	1,143	1,067
East North Central	2,601	2,641	1,483	1,360	815	794
West North Central	1,106	1,028	579	535	328	273
South	5,043	4,938	2,643	2,621	1,433	1,486
South Atlantic	2,465	2,414	1,344	1,331	737	747
East South Central	937	918	483	473	242	297
West South Central	1,642	1,605	816	817	453	443
West	3,418	3,469	1,871	1,896	1,092	1,071
Mountain	1,073	1,121	549	577	305	323
Pacific	2,345	2,348	1,322	1,319	788	748
COUNTY TYPE						
Large Metro	7,267	7,133	4,154	4,001	2,376	2,245
Small Metro	4,983	5,057	2,771	2,757	1,575	1,618
250K - 1 Mil. Pop.	3,519	3,517	1,960	1,918	1,109	1,148
<250K Pop.	1,464	1,540	811	839	466	471
Nonmetro	2,725	2,656	1,362	1,330	707	735
Urbanized	1,057	977	541	495	289	298
Less Urbanized	1,437	1,477	716	736	363	393
Completely Rural	230	203	105	99	55	44

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.73B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	52.6	51.2	29.1	27.9	16.4	15.9
GEOGRAPHIC DIVISION						
Northeast	55.6	54.2	33.9	32.8	19.6	19.1
New England	63.7	64.7	39.3	40.8	25.0	24.4
Middle Atlantic	52.9	50.6	32.1	30.0	17.8	17.2
Midwest	54.5	53.1	30.3 ^a	27.4	16.8	15.5
East North Central	54.8	55.0	31.3 ^a	28.3	17.2	16.5
West North Central	53.9	48.9	28.2	25.5	16.0	13.0
South	49.9	48.0	26.1	25.5	14.2	14.4
South Atlantic	51.0	49.0	27.8	27.0	15.2	15.2
East South Central	51.5	50.1	26.5	25.8	13.3	16.2
West South Central	47.4	45.4	23.6	23.1	13.1	12.5
West	52.5	52.0	28.7	28.4	16.8	16.0
Mountain	55.1	55.6	28.2	28.6	15.6	16.0
Pacific	51.4	50.4	29.0	28.3	17.3	16.1
COUNTY TYPE						
Large Metro	52.2	51.3	29.8	28.8	17.1	16.2
Small Metro	54.3	53.3	30.2	29.1	17.2	17.1
250K - 1 Mil. Pop.	54.7	53.0	30.5	28.9	17.2	17.3
<250K Pop.	53.4	54.0	29.6	29.4	17.0	16.5
Nonmetro	50.7 ^a	47.4	25.3	23.7	13.1	13.1
Urbanized	54.7	51.9	28.0	26.3	15.0	15.9
Less Urbanized	49.3	45.6	24.6	22.7	12.4	12.1
Completely Rural	43.3	42.0	19.7	20.5	10.3	9.1

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.74A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	66,344	65,797	12,531	12,089	6,906	7,164
GEOGRAPHIC DIVISION						
Northeast	13,662	12,710	2,530	2,472	1,549	1,442
New England	4,325	4,154	816	1,004	615	648
Middle Atlantic	9,336	8,556	1,715	1,468	934	794
Midwest	15,911	14,807	2,637	2,371	1,446	1,372
East North Central	11,187	10,800	1,867	1,868	1,085	1,156
West North Central	4,724 ^a	4,007	770 ^a	503	361	216
South	21,760	22,005	3,988	3,773	1,961	2,085
South Atlantic	12,199	12,156	2,343	2,321	1,194	1,258
East South Central	3,434	3,649	709	602	312	332
West South Central	6,126	6,200	936	850	455	495
West	15,011	16,274	3,376	3,473	1,949	2,265
Mountain	4,642	4,648	949	931	523	531
Pacific	10,369	11,626	2,427	2,543	1,426	1,734
COUNTY TYPE						
Large Metro	34,682	33,864	6,860	6,375	3,889	3,809
Small Metro	20,490	20,430	3,757	3,820	2,028	2,210
250K - 1 Mil. Pop.	15,772	15,640	2,973	2,907	1,622	1,577
<250K Pop.	4,718	4,790	784	913	405	633
Nonmetro	11,172	11,502	1,914	1,894	989	1,145
Urbanized	3,716	3,775	631	660	365	457
Less Urbanized	6,153	6,540	1,078	1,045	500	579
Completely Rural	1,304	1,187	205	190	123	108

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.74B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	39.2	38.5	7.4	7.1	4.1	4.2
GEOGRAPHIC DIVISION						
Northeast	40.8	37.9	7.6	7.4	4.6	4.3
New England	49.3	47.3	9.3	11.4	7.0	7.4
Middle Atlantic	37.8	34.6	6.9	5.9	3.8	3.2
Midwest	40.5 ^a	37.5	6.7	6.0	3.7	3.5
East North Central	40.5	38.9	6.8	6.7	3.9	4.2
West North Central	40.4 ^b	34.0	6.6 ^a	4.3	3.1	1.8
South	36.2	36.2	6.6	6.2	3.3	3.4
South Atlantic	38.5	37.9	7.4	7.2	3.8	3.9
East South Central	32.7	34.4	6.8	5.7	3.0	3.1
West South Central	34.2	34.1	5.2	4.7	2.5	2.7
West	41.0 ^a	43.9	9.2	9.4	5.3	6.1
Mountain	44.6	43.6	9.1	8.7	5.0	5.0
Pacific	39.6 ^a	44.1	9.3	9.6	5.4	6.6
COUNTY TYPE						
Large Metro	40.8	40.4	8.1	7.6	4.6	4.5
Small Metro	40.6	39.3	7.4	7.4	4.0	4.3
250K - 1 Mil. Pop.	41.2	40.1	7.8	7.5	4.2	4.0
<250K Pop.	38.6	36.9	6.4	7.0	3.3	4.9
Nonmetro	32.9	32.7	5.6	5.4	2.9	3.3
Urbanized	38.1	37.5	6.5	6.5	3.8	4.5
Less Urbanized	31.5	31.3	5.5	5.0	2.6	2.8
Completely Rural	28.1	28.3	4.4	4.5	2.7	2.6

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.75A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	81,319	80,643	20,818	20,177	11,564	11,763
GEOGRAPHIC DIVISION						
Northeast	16,469	15,481	4,242	4,147	2,539	2,416
New England	5,147	5,007	1,322	1,542	938	970
Middle Atlantic	11,322	10,474	2,920	2,605	1,601	1,447
Midwest	19,618	18,476	4,699	4,266	2,589	2,439
East North Central	13,788	13,441	3,349	3,229	1,900	1,950
West North Central	5,829 ^a	5,035	1,349 ^a	1,038	689 ^a	489
South	26,803	26,943	6,630	6,394	3,394	3,571
South Atlantic	14,664	14,570	3,686	3,652	1,931	2,004
East South Central	4,371	4,568	1,191	1,075	555	629
West South Central	7,768	7,805	1,752	1,667	908	937
West	18,429	19,743	5,247	5,370	3,042	3,337
Mountain	5,715	5,769	1,499	1,508	828	854
Pacific	12,714	13,974	3,748	3,862	2,214	2,483
COUNTY TYPE						
Large Metro	41,949	40,997	11,014	10,376	6,265	6,054
Small Metro	25,473	25,487	6,528	6,577	3,603	3,829
250K - 1 Mil. Pop.	19,292	19,157	4,933	4,825	2,732	2,725
<250K Pop.	6,182	6,330	1,595	1,751	872	1,104
Nonmetro	13,897	14,159	3,276	3,224	1,696	1,880
Urbanized	4,773	4,752	1,172	1,155	655	755
Less Urbanized	7,590	8,017	1,795	1,780	863	972
Completely Rural	1,534	1,391	309	289	178	153

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.75B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	41.1	40.3	10.5	10.1	5.8	5.9
GEOGRAPHIC DIVISION						
Northeast	42.8 ^a	40.1	11.0	10.7	6.6	6.3
New England	51.2	49.6	13.1	15.3	9.3	9.6
Middle Atlantic	39.8	36.7	10.3	9.1	5.6	5.1
Midwest	42.6 ^a	39.8	10.2	9.2	5.6	5.3
East North Central	42.6	41.3	10.4	9.9	5.9	6.0
West North Central	42.4 ^b	36.3	9.8 ^a	7.5	5.0 ^a	3.5
South	38.2	37.9	9.4	9.0	4.8	5.0
South Atlantic	40.2	39.4	10.1	9.9	5.3	5.4
East South Central	35.5	36.7	9.7	8.6	4.5	5.1
West South Central	36.3	36.0	8.2	7.7	4.2	4.3
West	42.7	45.2	12.2	12.3	7.1	7.6
Mountain	46.2	45.5	12.1	11.9	6.7	6.7
Pacific	41.3 ^a	45.0	12.2	12.4	7.2	8.0
COUNTY TYPE						
Large Metro	42.4	42.0	11.1	10.6	6.3	6.2
Small Metro	42.7	41.5	10.9	10.7	6.0	6.2
250K - 1 Mil. Pop.	43.2	42.0	11.0	10.6	6.1	6.0
<250K Pop.	41.3	40.0	10.7	11.1	5.8	7.0
Nonmetro	35.3	34.7	8.3	7.9	4.3	4.6
Urbanized	40.9	39.7	10.0	9.7	5.6	6.3
Less Urbanized	33.8	33.2	8.0	7.4	3.8	4.0
Completely Rural	29.6	29.8	6.0	6.2	3.4	3.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.76A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	76,428	76,321	19,102	18,589	10,458	10,714
GEOGRAPHIC DIVISION						
Northeast	15,417	14,634	3,892	3,882	2,170	2,307
New England	4,899	4,727	1,206 ^a	1,482	783	954
Middle Atlantic	10,518	9,907	2,686	2,400	1,387	1,353
Midwest	18,447	17,650	4,224	4,062	2,383	2,296
East North Central	13,082	12,773	3,111	3,036	1,766	1,777
West North Central	5,365	4,877	1,113	1,026	617	519
South	24,806	25,173	5,980	5,690	3,068	3,172
South Atlantic	13,373	13,777	3,344	3,306	1,857	1,836
East South Central	4,192	4,109	1,092	974	413	553
West South Central	7,241	7,287	1,544	1,409	798	784
West	17,758	18,865	5,006	4,956	2,837	2,938
Mountain	5,590	5,475	1,358	1,391	804	781
Pacific	12,168	13,389	3,648	3,564	2,033	2,157
COUNTY TYPE						
Large Metro	39,429	38,755	10,114	9,614	5,557	5,411
Small Metro	24,177	24,191	6,022	5,944	3,332	3,442
250K - 1 Mil. Pop.	18,315	18,102	4,550	4,367	2,488	2,497
<250K Pop.	5,862	6,089	1,472	1,577	843	945
Nonmetro	12,822	13,376	2,966	3,031	1,570 ^a	1,861
Urbanized	4,505	4,586	1,123	1,161	598	763
Less Urbanized	6,939	7,525	1,518	1,569	800	916
Completely Rural	1,378	1,265	325	300	172	182

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.76B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	34.6	34.2	8.6	8.3	4.7	4.8
GEOGRAPHIC DIVISION						
Northeast	36.1	34.2	9.1	9.1	5.1	5.4
New England	43.9	42.2	10.8 ^a	13.2	7.0	8.5
Middle Atlantic	33.4	31.3	8.5	7.6	4.4	4.3
Midwest	35.8	34.0	8.2	7.8	4.6	4.4
East North Central	36.2	35.2	8.6	8.4	4.9	4.9
West North Central	34.8	31.3	7.2	6.6	4.0	3.3
South	31.6	31.7	7.6	7.2	3.9	4.0
South Atlantic	33.0	33.5	8.3	8.0	4.6	4.5
East South Central	30.5	29.6	8.0	7.0	3.0	4.0
West South Central	29.9	29.7	6.4	5.7	3.3	3.2
West	36.7	38.4	10.3	10.1	5.9	6.0
Mountain	40.0	38.3	9.7	9.7	5.8	5.5
Pacific	35.3 ^a	38.5	10.6	10.2	5.9	6.2
COUNTY TYPE						
Large Metro	35.7	35.5	9.2	8.8	5.0	5.0
Small Metro	36.3	35.2	9.0	8.7	5.0	5.0
250K - 1 Mil. Pop.	36.6	35.5	9.1	8.6	5.0	4.9
<250K Pop.	35.2	34.5	8.8	8.9	5.1	5.4
Nonmetro	29.2	29.4	6.7	6.7	3.6	4.1
Urbanized	34.2	34.5	8.5	8.7	4.5	5.7
Less Urbanized	27.7	27.9	6.1	5.8	3.2	3.4
Completely Rural	24.0	24.0	5.7	5.7	3.0	3.5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.77A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	4,345	4,283	3,284	3,136	1,676	1,678
GEOGRAPHIC DIVISION						
Northeast	724	783	581	603	307	325
New England	257	265	202	222	123	134
Middle Atlantic	466	517	379	381	184	191
Midwest	1,037	979	780	731	417	392
East North Central	731	683	551	507	300	272
West North Central	306	296	229	225	117	120
South	1,447	1,436	1,068	1,003	516	530
South Atlantic	686	718	517	506	260	258
East South Central	242	267	177	189	74	108
West South Central	519	451	374	308	182	165
West	1,137	1,086	855	799	435	431
Mountain	370	366	271	270	140	136
Pacific	767	719	584	529	295	295
COUNTY TYPE						
Large Metro	2,151	2,004	1,660 ^a	1,483	834	784
Small Metro	1,396	1,427	1,029	1,037	547	559
250K - 1 Mil. Pop.	1,076	1,035	802	766	431	406
<250K Pop.	319 ^a	391	227	271	117	153
Nonmetro	798	853	594	616	295	335
Urbanized	290	289	206	219	102	124
Less Urbanized	418	458	320	315	156	163
Completely Rural	90	106	68	82	36	48

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.77B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	18.7	18.3	14.2	13.4	7.2	7.2
GEOGRAPHIC DIVISION						
Northeast	17.4	18.6	14.0	14.4	7.4	7.7
New England	23.6	24.0	18.5	20.0	11.3	12.1
Middle Atlantic	15.2	16.7	12.4	12.3	6.0	6.2
Midwest	19.0	18.0	14.3	13.4	7.6	7.2
East North Central	19.4	18.1	14.6	13.4	7.9	7.2
West North Central	18.0	17.6	13.5	13.3	6.9	7.1
South	17.6	17.3	12.9	12.1	6.3	6.4
South Atlantic	17.1	17.6	12.9	12.4	6.5	6.3
East South Central	17.0	18.7	12.5	13.3	5.2 ^a	7.6
West South Central	18.4	16.0	13.3 ^a	10.9	6.4	5.8
West	21.3	20.1	16.0	14.8	8.2	8.0
Mountain	23.0	22.6	16.8	16.7	8.7	8.4
Pacific	20.6	19.1	15.7	14.0	7.9	7.8
COUNTY TYPE						
Large Metro	18.6	17.6	14.4 ^a	13.0	7.2	6.9
Small Metro	20.0	19.8	14.7	14.4	7.8	7.8
250K - 1 Mil. Pop.	20.4	19.2	15.2	14.2	8.1	7.5
<250K Pop.	18.7	21.6	13.3	15.0	6.8	8.5
Nonmetro	17.2	17.8	12.8	12.8	6.3	7.0
Urbanized	19.4	21.8	13.8 ^a	16.5	6.8 ^a	9.3
Less Urbanized	16.1	15.9	12.3	11.0	6.0	5.7
Completely Rural	16.3	17.7	12.3	13.8	6.5	8.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.78A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	13,326	13,256	6,962	6,860	4,049	3,950
GEOGRAPHIC DIVISION						
Northeast	2,538	2,512	1,512	1,497	915	887
New England	750	795	451	492	299	298
Middle Atlantic	1,788	1,717	1,060	1,005	616	588
Midwest	3,366	3,324	1,755	1,659	993	940
East North Central	2,386	2,388	1,276	1,185	709	696
West North Central	980	935	479	474	284	244
South	4,399	4,330	2,139	2,119	1,189	1,212
South Atlantic	2,159	2,143	1,130	1,107	634	620
East South Central	817	810	367	391	186	242
West South Central	1,423	1,378	642	621	369	349
West	3,023	3,090	1,557	1,585	952	911
Mountain	947	988	449	465	267	272
Pacific	2,076	2,102	1,108	1,120	685	640
COUNTY TYPE						
Large Metro	6,484	6,424	3,537	3,412	2,090	1,937
Small Metro	4,486	4,518	2,340	2,329	1,387	1,372
250K - 1 Mil. Pop.	3,181	3,120	1,668	1,613	979	967
<250K Pop.	1,304	1,398	672	716	408	405
Nonmetro	2,357	2,315	1,086	1,119	572	641
Urbanized	946	861	450	426	239	266
Less Urbanized	1,230	1,281	557	615	302	339
Completely Rural	181	173	79	77	30	36

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.78B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	46.8	45.7	24.5	23.7	14.2	13.6
GEOGRAPHIC DIVISION						
Northeast	50.3	49.1	30.0	29.3	18.1	17.3
New England	58.2	60.3	35.0	37.3	23.2	22.6
Middle Atlantic	47.6	45.3	28.2	26.5	16.4	15.5
Midwest	49.5	48.1	25.8	24.0	14.6	13.6
East North Central	50.3	49.7	26.9	24.7	14.9	14.5
West North Central	47.8	44.5	23.3	22.6	13.9	11.6
South	43.5	42.1	21.1	20.6	11.8	11.8
South Atlantic	44.7	43.5	23.4	22.5	13.1	12.6
East South Central	44.9	44.2	20.2	21.3	10.2	13.2
West South Central	41.1	39.0	18.5	17.6	10.6	9.9
West	46.4	46.3	23.9	23.7	14.6	13.7
Mountain	48.6	49.0	23.0	23.1	13.7	13.5
Pacific	45.5	45.1	24.3	24.0	15.0	13.7
COUNTY TYPE						
Large Metro	46.6	46.2	25.4	24.6	15.0	13.9
Small Metro	48.9	47.6	25.5	24.6	15.1	14.5
250K - 1 Mil. Pop.	49.5	47.0	25.9	24.3	15.2	14.6
<250K Pop.	47.6	49.0	24.5	25.1	14.9	14.2
Nonmetro	43.8	41.3	20.2	20.0	10.6	11.5
Urbanized	48.9	45.8	23.3	22.6	12.4	14.1
Less Urbanized	42.2	39.6	19.1	19.0	10.4	10.5
Completely Rural	34.2	35.7	14.8	16.0	5.7	7.5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.79A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	58,757	58,782	8,856	8,593	4,734	5,085
GEOGRAPHIC DIVISION						
Northeast	12,155	11,339	1,799	1,782	948	1,095
New England	3,891	3,666	553	769	361	522
Middle Atlantic	8,264	7,673	1,247	1,014	586	573
Midwest	14,044	13,347	1,689	1,672	973	964
East North Central	9,964	9,701	1,283	1,344	757	810
West North Central	4,079	3,645	406	327	216	154
South	18,960	19,407	2,773	2,568	1,363	1,430
South Atlantic	10,529	10,916	1,697	1,692	963	957
East South Central	3,133	3,033	547	394	152	203
West South Central	5,298	5,458	529	481	248	270
West	13,598	14,689	2,594	2,572	1,450	1,595
Mountain	4,273	4,121	639	656	397	373
Pacific	9,325 ^a	10,568	1,955	1,916	1,053	1,223
COUNTY TYPE						
Large Metro	30,794	30,327	4,917	4,720	2,633	2,690
Small Metro	18,296	18,247	2,653	2,578	1,397	1,510
250K - 1 Mil. Pop.	14,057	13,947	2,080	1,988	1,078	1,124
<250K Pop.	4,238	4,300	573	590	319	386
Nonmetro	9,667	10,208	1,286	1,296	704	885
Urbanized	3,269	3,435	467	517	257	374
Less Urbanized	5,291	5,786	641	639	342	414
Completely Rural	1,106	986	178	141	105	97

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.79B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	34.7	34.4	5.2	5.0	2.8	3.0
GEOGRAPHIC DIVISION						
Northeast	36.3	33.8	5.4	5.3	2.8	3.3
New England	44.4	41.8	6.3	8.8	4.1	5.9
Middle Atlantic	33.4	31.0	5.0	4.1	2.4	2.3
Midwest	35.7	33.8	4.3	4.2	2.5	2.4
East North Central	36.1	35.0	4.6	4.8	2.7	2.9
West North Central	34.9	31.0	3.5	2.8	1.8	1.3
South	31.6	31.9	4.6	4.2	2.3	2.4
South Atlantic	33.3	34.0	5.4	5.3	3.0	3.0
East South Central	29.9	28.6	5.2	3.7	1.5	1.9
West South Central	29.6	30.0	2.9	2.6	1.4	1.5
West	37.1	39.6	7.1	6.9	4.0	4.3
Mountain	41.0	38.6	6.1	6.1	3.8	3.5
Pacific	35.6 ^a	40.1	7.5	7.3	4.0	4.6
COUNTY TYPE						
Large Metro	36.2	36.2	5.8	5.6	3.1	3.2
Small Metro	36.2	35.1	5.3	5.0	2.8	2.9
250K - 1 Mil. Pop.	36.7	35.8	5.4	5.1	2.8	2.9
<250K Pop.	34.7	33.1	4.7	4.5	2.6	3.0
Nonmetro	28.5	29.0	3.8	3.7	2.1	2.5
Urbanized	33.6	34.1	4.8	5.1	2.6	3.7
Less Urbanized	27.1	27.7	3.3	3.1	1.7	2.0
Completely Rural	23.8	23.5	3.8	3.4	2.3	2.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.80A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	72,083	72,038	15,818	15,453	8,783	9,035
GEOGRAPHIC DIVISION						
Northeast	14,693	13,852	3,311	3,279	1,863	1,982
New England	4,642	4,461	1,004 ^a	1,261	660	820
Middle Atlantic	10,051	9,390	2,307	2,019	1,203	1,161
Midwest	17,410	16,670	3,444	3,331	1,966	1,904
East North Central	12,351	12,090	2,560	2,530	1,466	1,506
West North Central	5,059	4,581	885	801	500	399
South	23,359	23,737	4,913	4,686	2,552	2,642
South Atlantic	12,688	13,059	2,828	2,800	1,597	1,578
East South Central	3,950	3,843	914	785	339	445
West South Central	6,722	6,835	1,170	1,102	616	619
West	16,621	17,779	4,151	4,157	2,402	2,507
Mountain	5,220	5,109	1,087	1,121	664	645
Pacific	11,401 ^a	12,670	3,064	3,036	1,738	1,862
COUNTY TYPE						
Large Metro	37,278	36,751	8,454	8,131	4,724	4,626
Small Metro	22,781	22,765	4,993	4,907	2,784	2,882
250K - 1 Mil. Pop.	17,238	17,066	3,748	3,601	2,058	2,091
<250K Pop.	5,543	5,698	1,245	1,306	726	791
Nonmetro	12,024	12,523	2,372	2,415	1,275	1,527
Urbanized	4,215	4,296	917	943	496	640
Less Urbanized	6,521	7,067	1,199	1,254	643	753
Completely Rural	1,287	1,159	256	218	135	134

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.80B Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	36.4	36.0	8.0	7.7	4.4	4.5
GEOGRAPHIC DIVISION						
Northeast	38.1	35.9	8.6	8.5	4.8	5.1
New England	46.2	44.2	10.0	12.5	6.6	8.1
Middle Atlantic	35.3	32.9	8.1	7.1	4.2	4.1
Midwest	37.8	35.9	7.5	7.2	4.3	4.1
East North Central	38.2	37.2	7.9	7.8	4.5	4.6
West North Central	36.8	33.0	6.4	5.8	3.6	2.9
South	33.3	33.4	7.0	6.6	3.6	3.7
South Atlantic	34.8	35.3	7.7	7.6	4.4	4.3
East South Central	32.1	30.9	7.4	6.3	2.8	3.6
West South Central	31.4	31.5	5.5	5.1	2.9	2.9
West	38.5	40.7	9.6	9.5	5.6	5.7
Mountain	42.2	40.3	8.8	8.8	5.4	5.1
Pacific	37.1 ^a	40.8	10.0	9.8	5.7	6.0
COUNTY TYPE						
Large Metro	37.7	37.6	8.5	8.3	4.8	4.7
Small Metro	38.2	37.1	8.4	8.0	4.7	4.7
250K - 1 Mil. Pop.	38.6	37.4	8.4	7.9	4.6	4.6
<250K Pop.	37.0	36.0	8.3	8.2	4.9	5.0
Nonmetro	30.6	30.7	6.0	5.9	3.2	3.7
Urbanized	36.1	35.9	7.9	7.9	4.3	5.3
Less Urbanized	29.0	29.3	5.3	5.2	2.9	3.1
Completely Rural	24.9	24.8	5.0	4.7	2.6	2.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.81A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	25,406	24,896	3,742	3,328	1,552	1,213
GEOGRAPHIC DIVISION						
Northeast	5,249	4,691	666	670	346	209
New England	1,696	1,695	208	256	125	73
Middle Atlantic	3,553	2,996	458	414	221	135
Midwest	4,938	4,903	731	664	287	234
East North Central	3,551	3,533	531	500	190	170
West North Central	1,386	1,370	200	165	97	63
South	7,927	7,272	1,461 ^b	1,060	586 ^a	355
South Atlantic	4,394	4,227	710	540	334 ^a	178
East South Central	1,157	1,004	284	176	85	57
West South Central	2,375	2,041	467	344	167	120
West	7,293	8,030	884	934	333	416
Mountain	1,945	2,271	263	336	105	113
Pacific	5,348	5,759	621	597	228	303
COUNTY TYPE						
Large Metro	14,074	13,368	1,877	1,648	873	688
Small Metro	7,943	7,547	1,355	1,169	519	367
250K - 1 Mil. Pop.	6,133	5,640	1,059	885	442	290
<250K Pop.	1,810	1,906	296	284	76	78
Nonmetro	3,389 ^a	3,981	510	511	160	159
Urbanized	1,144 ^a	1,537	196	157	58	42
Less Urbanized	1,802	2,074	271	295	96	99
Completely Rural	443	370	43	59	6	18

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.81B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11.5	11.2	1.7	1.5	0.7	0.5
GEOGRAPHIC DIVISION						
Northeast	12.3	11.0	1.6	1.6	0.8	0.5
New England	15.2	15.1	1.9	2.3	1.1	0.7
Middle Atlantic	11.3	9.5	1.5	1.3	0.7	0.4
Midwest	9.6	9.5	1.4	1.3	0.6	0.5
East North Central	9.8	9.7	1.5	1.4	0.5	0.5
West North Central	9.0	8.8	1.3	1.1	0.6	0.4
South	10.1	9.2	1.9 ^b	1.3	0.7 ^a	0.4
South Atlantic	10.9	10.3	1.8	1.3	0.8 ^a	0.4
East South Central	8.4	7.2	2.1	1.3	0.6	0.4
West South Central	9.8	8.3	1.9	1.4	0.7	0.5
West	15.1	16.3	1.8	1.9	0.7	0.8
Mountain	13.9	15.9	1.9	2.4	0.8	0.8
Pacific	15.5	16.5	1.8	1.7	0.7	0.9
COUNTY TYPE						
Large Metro	12.7	12.3	1.7	1.5	0.8	0.6
Small Metro	11.9	11.0	2.0	1.7	0.8	0.5
250K - 1 Mil. Pop.	12.3	11.1	2.1	1.7	0.9	0.6
<250K Pop.	10.9	10.8	1.8	1.6	0.5	0.4
Nonmetro	7.7	8.7	1.2	1.1	0.4	0.3
Urbanized	8.7 ^a	11.6	1.5	1.2	0.4	0.3
Less Urbanized	7.2	7.7	1.1	1.1	0.4	0.4
Completely Rural	7.7	7.0	0.8	1.1	0.1	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.82A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	551	550	373	389	124	132
GEOGRAPHIC DIVISION						
Northeast	68	64	54	42	19	11
New England	19	23	14	15	4	4
Middle Atlantic	49	40	40	27	14	7
Midwest	88	101	53	73	21	31
East North Central	57	73	35	50	14	21
West North Central	31	28	18	24	7	10
South	214	201	147	143	53	46
South Atlantic	87	80	64	55	20	22
East South Central	30	43	20	29	10	8
West South Central	97	78	64	58	24	17
West	181	185	118	130	30	44
Mountain	66	65	46	49	11	21
Pacific	115	119	72	82	20	23
COUNTY TYPE						
Large Metro	270	235	185	170	63	48
Small Metro	177	180	122	117	43	45
250K - 1 Mil. Pop.	131	127	86	81	35	33
<250K Pop.	46	54	36	36	9	12
Nonmetro	104	135	66	101	18	38
Urbanized	34	38	20	25	4	10
Less Urbanized	53	77	38	61	13	19
Completely Rural	16	20	7	15	0	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.82B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	2.4	2.4	1.6	1.7	0.5	0.6
GEOGRAPHIC DIVISION						
Northeast	1.6	1.5	1.3	1.0	0.4	0.3
New England	1.7	2.1	1.3	1.3	0.4	0.4
Middle Atlantic	1.6	1.3	1.3	0.9	0.5	0.2
Midwest	1.6	1.8	1.0	1.3	0.4	0.6
East North Central	1.5	1.9	0.9	1.3	0.4	0.5
West North Central	1.8	1.7	1.1	1.4	0.4	0.6
South	2.6	2.4	1.8	1.7	0.6	0.6
South Atlantic	2.2	2.0	1.6	1.4	0.5	0.5
East South Central	2.1	3.1	1.4	2.0	0.7	0.5
West South Central	3.4	2.7	2.3	2.1	0.9	0.6
West	3.4	3.4	2.2	2.4	0.6	0.8
Mountain	4.1	4.0	2.8	3.0	0.7	1.3
Pacific	3.1	3.2	1.9	2.2	0.5	0.6
COUNTY TYPE						
Large Metro	2.3	2.1	1.6	1.5	0.5	0.4
Small Metro	2.5	2.5	1.7	1.6	0.6	0.6
250K - 1 Mil. Pop.	2.5	2.3	1.6	1.5	0.7	0.6
<250K Pop.	2.7	3.0	2.1	2.0	0.5	0.7
Nonmetro	2.2	2.8	1.4	2.1	0.4	0.8
Urbanized	2.3	2.8	1.3	1.9	0.3	0.7
Less Urbanized	2.0	2.7	1.5	2.1	0.5	0.6
Completely Rural	3.0	3.3	1.3	2.5	0.1	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.83A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	3,383	3,148	1,493 ^b	1,274	476	395
GEOGRAPHIC DIVISION						
Northeast	521	544	230	234	60	67
New England	156	161	69	76	21	28
Middle Atlantic	365	383	161	157	39	39
Midwest	736	694	353	284	103	75
East North Central	505	501	256	207	73	56
West North Central	231	193	97	77	30	18
South	1,191	1,042	538 ^a	432	192 ^a	134
South Atlantic	537	487	258	204	103	65
East South Central	168	172	70	69	13	19
West South Central	487 ^a	383	210	159	76	50
West	934	869	372	325	121	119
Mountain	319	332	130	134	43	39
Pacific	615	537	242	191	78	80
COUNTY TYPE						
Large Metro	1,592	1,526	713	614	240	186
Small Metro	1,238 ^a	1,058	557 ^a	455	169	157
250K - 1 Mil. Pop.	899 ^a	744	405 ^a	312	122	114
<250K Pop.	339	314	152	142	47	42
Nonmetro	553	565	223	206	67	52
Urbanized	214	209	82	74	23	16
Less Urbanized	292	315	119	122	39	34
Completely Rural	47	41	22	10	5	1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.83B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11.9 ^a	10.9	5.2 ^b	4.4	1.7	1.4
GEOGRAPHIC DIVISION						
Northeast	10.3	10.6	4.6	4.6	1.2	1.3
New England	12.1	12.2	5.3	5.8	1.7	2.1
Middle Atlantic	9.7	10.1	4.3	4.1	1.0	1.0
Midwest	10.8	10.0	5.2 ^a	4.1	1.5	1.1
East North Central	10.6	10.4	5.4	4.3	1.5	1.2
West North Central	11.3	9.2	4.7	3.7	1.5	0.9
South	11.8 ^a	10.1	5.3 ^a	4.2	1.9 ^a	1.3
South Atlantic	11.1	9.9	5.3	4.1	2.1 ^a	1.3
East South Central	9.2	9.4	3.8	3.8	0.7	1.0
West South Central	14.1 ^a	10.8	6.1 ^a	4.5	2.2	1.4
West	14.3	13.0	5.7	4.9	1.9	1.8
Mountain	16.4	16.5	6.7	6.6	2.2	1.9
Pacific	13.5	11.5	5.3	4.1	1.7	1.7
COUNTY TYPE						
Large Metro	11.4	11.0	5.1	4.4	1.7	1.3
Small Metro	13.5 ^b	11.2	6.1 ^b	4.8	1.8	1.7
250K - 1 Mil. Pop.	14.0 ^b	11.2	6.3 ^b	4.7	1.9	1.7
<250K Pop.	12.4	11.0	5.5	5.0	1.7	1.5
Nonmetro	10.3	10.1	4.1	3.7	1.2	0.9
Urbanized	11.1	11.1	4.2	4.0	1.2	0.8
Less Urbanized	10.0	9.7	4.1	3.8	1.3	1.1
Completely Rural	8.9	8.4	4.2	2.0	1.0	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.84A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	21,471	21,198	1,876	1,666	952	687
GEOGRAPHIC DIVISION						
Northeast	4,659	4,084	382	394	268	130
New England	1,520	1,511	125	165	99	41
Middle Atlantic	3,138	2,573	257	229	168	89
Midwest	4,114	4,109	325	307	163	128
East North Central	2,989	2,960	240	243	103	93
West North Central	1,124	1,148	85	64	60	35
South	6,521	6,029	776 ^a	486	340 ^a	175
South Atlantic	3,770	3,660	388	281	212	91
East South Central	959	788	195 ^a	78	62	30
West South Central	1,792	1,580	193	127	67	53
West	6,178	6,976	394	478	182	254
Mountain	1,560	1,874	87	154	52	53
Pacific	4,618	5,103	307	324	130	201
COUNTY TYPE						
Large Metro	12,212	11,607	978	864	571	453
Small Metro	6,528	6,309	676	598	306	165
250K - 1 Mil. Pop.	5,103	4,770	568	493	286	142
<250K Pop.	1,425	1,538	108	105	20	23
Nonmetro	2,732 ^a	3,282	222	204	76	69
Urbanized	896 ^a	1,290	95	57	32	17
Less Urbanized	1,456	1,682	114	112	43	46
Completely Rural	380	309	13	34	1	6

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.84B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	12.7	12.4	1.1	1.0	0.6	0.4
GEOGRAPHIC DIVISION						
Northeast	13.9	12.2	1.1	1.2	0.8	0.4
New England	17.3	17.2	1.4	1.9	1.1	0.5
Middle Atlantic	12.7	10.4	1.0	0.9	0.7	0.4
Midwest	10.5	10.4	0.8	0.8	0.4	0.3
East North Central	10.8	10.7	0.9	0.9	0.4	0.3
West North Central	9.6	9.8	0.7	0.5	0.5	0.3
South	10.9	9.9	1.3 ^a	0.8	0.6 ^a	0.3
South Atlantic	11.9	11.4	1.2	0.9	0.7	0.3
East South Central	9.1	7.4	1.9	0.7	0.6	0.3
West South Central	10.0	8.7	1.1	0.7	0.4	0.3
West	16.9	18.8	1.1	1.3	0.5	0.7
Mountain	15.0	17.6	0.8	1.4	0.5	0.5
Pacific	17.6	19.3	1.2	1.2	0.5	0.8
COUNTY TYPE						
Large Metro	14.4	13.8	1.2	1.0	0.7	0.5
Small Metro	12.9	12.1	1.3	1.2	0.6	0.3
250K - 1 Mil. Pop.	13.3	12.2	1.5	1.3	0.7	0.4
<250K Pop.	11.7	11.9	0.9	0.8	0.2	0.2
Nonmetro	8.0	9.3	0.7	0.6	0.2	0.2
Urbanized	9.2 ^a	12.8	1.0	0.6	0.3	0.2
Less Urbanized	7.4	8.1	0.6	0.5	0.2	0.2
Completely Rural	8.2	7.4	0.3	0.8	0.0	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.85A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	24,854	24,346	3,369	2,940	1,428	1,081
GEOGRAPHIC DIVISION						
Northeast	5,180	4,628	611	628	328	197
New England	1,677	1,672	193	242	120	69
Middle Atlantic	3,503	2,956	418	387	207	128
Midwest	4,850	4,802	677	591	266	203
East North Central	3,494	3,461	495	450	176	150
West North Central	1,356	1,341	182	141	90	53
South	7,713	7,071	1,314 ^b	918	532 ^a	309
South Atlantic	4,307	4,147	646	485	315 ^a	156
East South Central	1,127	961	265	147	75	49
West South Central	2,278	1,963	403	286	143	104
West	7,112	7,846	766	803	303	372
Mountain	1,879	2,206	217	288	94	92
Pacific	5,233	5,640	549	515	209	281
COUNTY TYPE						
Large Metro	13,803	13,133	1,691	1,478	811	639
Small Metro	7,766	7,366	1,233	1,052	475	322
250K - 1 Mil. Pop.	6,002	5,514	973	805	408	256
<250K Pop.	1,764	1,853	260	248	67	66
Nonmetro	3,285 ^a	3,846	444	410	143	120
Urbanized	1,110 ^a	1,499	176	131	54	33
Less Urbanized	1,748	1,997	232	234	82	80
Completely Rural	427	350	36	44	6	7

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.85B Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	12.6	12.2	1.7	1.5	0.7 ^a	0.5
GEOGRAPHIC DIVISION						
Northeast	13.4	12.0	1.6	1.6	0.9	0.5
New England	16.7	16.6	1.9	2.4	1.2	0.7
Middle Atlantic	12.3	10.4	1.5	1.4	0.7	0.4
Midwest	10.5	10.3	1.5	1.3	0.6	0.4
East North Central	10.8	10.6	1.5	1.4	0.5	0.5
West North Central	9.9	9.7	1.3	1.0	0.7	0.4
South	11.0	9.9	1.9 ^b	1.3	0.8 ^a	0.4
South Atlantic	11.8	11.2	1.8	1.3	0.9 ^a	0.4
East South Central	9.2	7.7	2.2	1.2	0.6	0.4
West South Central	10.7	9.0	1.9	1.3	0.7	0.5
West	16.5	17.9	1.8	1.8	0.7	0.9
Mountain	15.2	17.4	1.8	2.3	0.8	0.7
Pacific	17.0	18.2	1.8	1.7	0.7	0.9
COUNTY TYPE						
Large Metro	14.0	13.4	1.7	1.5	0.8	0.7
Small Metro	13.0	12.0	2.1	1.7	0.8	0.5
250K - 1 Mil. Pop.	13.4	12.1	2.2	1.8	0.9	0.6
<250K Pop.	11.8	11.7	1.7	1.6	0.5	0.4
Nonmetro	8.4	9.4	1.1	1.0	0.4	0.3
Urbanized	9.5 ^a	12.5	1.5	1.1	0.5	0.3
Less Urbanized	7.8	8.3	1.0	1.0	0.4	0.3
Completely Rural	8.2	7.5	0.7	0.9	0.1	0.2

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.86A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	25,061	26,125	3,191	3,483	922	971
GEOGRAPHIC DIVISION						
Northeast	4,728	5,030	611 ^a	748	178	205
New England	1,656	1,959	197 ^a	281	63	92
Middle Atlantic	3,072	3,071	414	467	114	113
Midwest	5,507	5,685	776	743	224	181
East North Central	3,893	4,086	580	563	184	138
West North Central	1,614	1,599	196	180	40	43
South	7,392	7,394	996	1,080	294	334
South Atlantic	4,163	4,220	565	623	184	197
East South Central	1,019	1,061	140	150	29	56
West South Central	2,211	2,113	291	307	81	81
West	7,433	8,015	807	912	226	251
Mountain	2,105	2,317	272	286	70	92
Pacific	5,328	5,698	535	626	156	159
COUNTY TYPE						
Large Metro	13,479	13,614	1,556	1,797	478	532
Small Metro	7,877	8,397	1,086	1,169	323	309
250K - 1 Mil. Pop.	5,989	6,102	759	834	227	193
<250K Pop.	1,888	2,296	327	335	96	116
Nonmetro	3,705	4,114	549	517	121	130
Urbanized	1,316 ^a	1,642	186	215	55	63
Less Urbanized	1,981	2,104	303	258	58	61
Completely Rural	408	368	60	44	8	7

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.86B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11.3	11.7	1.4	1.6	0.4	0.4
GEOGRAPHIC DIVISION						
Northeast	11.1	11.7	1.4 ^a	1.7	0.4	0.5
New England	14.8	17.5	1.8 ^a	2.5	0.6	0.8
Middle Atlantic	9.7	9.7	1.3	1.5	0.4	0.4
Midwest	10.7	11.0	1.5	1.4	0.4	0.3
East North Central	10.8	11.3	1.6	1.6	0.5	0.4
West North Central	10.5	10.3	1.3	1.2	0.3	0.3
South	9.4	9.3	1.3	1.4	0.4	0.4
South Atlantic	10.3	10.3	1.4	1.5	0.5	0.5
East South Central	7.4	7.6	1.0	1.1	0.2	0.4
West South Central	9.1	8.6	1.2	1.3	0.3	0.3
West	15.3	16.3	1.7	1.9	0.5	0.5
Mountain	15.1	16.2	1.9	2.0	0.5	0.6
Pacific	15.5	16.4	1.6	1.8	0.5	0.5
COUNTY TYPE						
Large Metro	12.2	12.5	1.4	1.6	0.4	0.5
Small Metro	11.8	12.2	1.6	1.7	0.5	0.4
250K - 1 Mil. Pop.	12.0	12.0	1.5	1.6	0.5	0.4
<250K Pop.	11.3	13.0	2.0	1.9	0.6	0.7
Nonmetro	8.4	9.0	1.2	1.1	0.3	0.3
Urbanized	10.0 ^a	12.4	1.4	1.6	0.4	0.5
Less Urbanized	7.9	7.8	1.2	1.0	0.2	0.2
Completely Rural	7.1	7.0	1.1	0.8	0.1	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.87A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	1,317	1,347	889	908	254	270
GEOGRAPHIC DIVISION						
Northeast	253	250	192	179	56	47
New England	89	100	68	74	27	24
Middle Atlantic	163	150	125	105	30	23
Midwest	255	286	185	197	61	56
East North Central	189	208	146	144	49	47
West North Central	66	78	39	54	12	9
South	432	462	269	305	74	107
South Atlantic	206	231	128	162	33	57
East South Central	63	76	40	46	11	21
West South Central	163	155	100	97	30	29
West	377	350	244	227	63	60
Mountain	127	121	76	80	25	25
Pacific	250	229	168	146	38	35
COUNTY TYPE						
Large Metro	639	635	442	434	135	135
Small Metro	428	443	279	298	81	84
250K - 1 Mil. Pop.	338	314	214	215	63	55
<250K Pop.	90	129	65	82	18	28
Nonmetro	250	269	168	177	39	51
Urbanized	82	95	57	62	16	16
Less Urbanized	139	136	89	89	22	30
Completely Rural	29	38	22	26	1	5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.87B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	5.7	5.8	3.8	3.9	1.1	1.2
GEOGRAPHIC DIVISION						
Northeast	6.1	5.9	4.6	4.3	1.4	1.1
New England	8.2	9.0	6.2	6.7	2.4	2.2
Middle Atlantic	5.3	4.9	4.1	3.4	1.0	0.7
Midwest	4.7	5.2	3.4	3.6	1.1	1.0
East North Central	5.0	5.5	3.9	3.8	1.3	1.2
West North Central	3.9	4.6	2.3	3.2	0.7	0.5
South	5.2	5.6	3.3	3.7	0.9	1.3
South Atlantic	5.1	5.7	3.2	4.0	0.8	1.4
East South Central	4.4	5.4	2.8	3.2	0.8	1.5
West South Central	5.8	5.5	3.6	3.4	1.1	1.0
West	7.1	6.5	4.6	4.2	1.2	1.1
Mountain	7.9	7.4	4.7	4.9	1.5	1.5
Pacific	6.7	6.1	4.5	3.9	1.0	0.9
COUNTY TYPE						
Large Metro	5.5	5.6	3.8	3.8	1.2	1.2
Small Metro	6.1	6.2	4.0	4.1	1.2	1.2
250K - 1 Mil. Pop.	6.4	5.8	4.0	4.0	1.2	1.0
<250K Pop.	5.2	7.1	3.8	4.5	1.0	1.6
Nonmetro	5.4	5.6	3.6	3.7	0.8	1.1
Urbanized	5.5	7.2	3.8	4.7	1.0	1.2
Less Urbanized	5.3	4.7	3.4	3.1	0.9	1.0
Completely Rural	5.2	6.3	3.9	4.4	0.1	0.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.88A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	5,490	5,592	1,928	1,959	534	532
GEOGRAPHIC DIVISION						
Northeast	1,047	1,136	393	442	110	108
New England	334	380	125	139	33	33
Middle Atlantic	713	755	268	302	77	75
Midwest	1,313	1,325	480	423	132	108
East North Central	928	980	339	321	104	79
West North Central	386	345	142	102	28	29
South	1,683	1,682	596	636	169	204
South Atlantic	923	874	352	369	102	132
East South Central	221 ^a	300	73	90	18	29
West South Central	540	508	171	177	50	43
West	1,446	1,450	459	458	122	111
Mountain	463	519	157	159	39	51
Pacific	984	930	303	299	83	60
COUNTY TYPE						
Large Metro	2,744	2,746	916	924	270	248
Small Metro	1,875	1,940	697	741	197	208
250K - 1 Mil. Pop.	1,329	1,369	486	508	144	125
<250K Pop.	546	571	211	233	54	83
Nonmetro	871	906	316	294	66	77
Urbanized	352	365	106	142	33	44
Less Urbanized	463	481	174	134	26	31
Completely Rural	56	60	36	18	7	1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.88B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	19.3	19.3	6.8	6.8	1.9	1.8
GEOGRAPHIC DIVISION						
Northeast	20.8	22.2	7.8	8.6	2.2	2.1
New England	25.9	28.8	9.7	10.6	2.6	2.5
Middle Atlantic	19.0	19.9	7.1	8.0	2.0	2.0
Midwest	19.3	19.2	7.1	6.1	1.9	1.6
East North Central	19.6	20.4	7.1	6.7	2.2	1.6
West North Central	18.8	16.4	6.9	4.8	1.3	1.4
South	16.6	16.3	5.9	6.2	1.7	2.0
South Atlantic	19.1	17.7	7.3	7.5	2.1	2.7
East South Central	12.1 ^a	16.4	4.0	4.9	1.0	1.6
West South Central	15.6	14.4	4.9	5.0	1.5	1.2
West	22.2	21.7	7.1	6.9	1.9	1.7
Mountain	23.7	25.8	8.0	7.9	2.0	2.5
Pacific	21.6	20.0	6.6	6.4	1.8	1.3
COUNTY TYPE						
Large Metro	19.7	19.8	6.6	6.7	1.9	1.8
Small Metro	20.4	20.5	7.6	7.8	2.2	2.2
250K - 1 Mil. Pop.	20.7	20.6	7.6	7.7	2.2	1.9
<250K Pop.	19.9	20.0	7.7	8.2	2.0	2.9
Nonmetro	16.2	16.2	5.9	5.2	1.2	1.4
Urbanized	18.2	19.4	5.5 ^a	7.6	1.7	2.3
Less Urbanized	15.9	14.9	6.0 ^a	4.1	0.9	1.0
Completely Rural	10.5	12.3	6.7	3.7	1.3	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.89A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	18,254	19,186	374 ^a	616	134	169
GEOGRAPHIC DIVISION						
Northeast	3,428	3,645	26 ^a	128	11	50
New England	1,233	1,480	5	68	3	35
Middle Atlantic	2,195	2,165	21	60	8	14
Midwest	3,938	4,074	111	123	31	17
East North Central	2,776	2,898	95	98	31	12
West North Central	1,162	1,176	16	25	*	5
South	5,277	5,250	132	138	50	22
South Atlantic	3,034	3,115	86	91	50	7
East South Central	735	684	27	14	*	6
West South Central	1,509	1,451	20	33	*	8
West	5,610	6,216	104 ^a	227	42	80
Mountain	1,516	1,678	39	46	7	17
Pacific	4,094	4,538	65 ^a	181	35	63
COUNTY TYPE						
Large Metro	10,096	10,232	198 ^b	439	73	149
Small Metro	5,574	6,014	110	131	45	18
250K - 1 Mil. Pop.	4,322	4,418	59	110	21	13
<250K Pop.	1,252	1,596	51	20	25	5
Nonmetro	2,584	2,940	65	46	16	3
Urbanized	882	1,182	22	11	6	3
Less Urbanized	1,379	1,487	40	35	10	*
Completely Rural	323	271	3	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.89B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	10.8	11.2	0.2 ^a	0.4	0.1	0.1
GEOGRAPHIC DIVISION						
Northeast	10.2	10.9	0.1 ^a	0.4	0.0	0.1
New England	14.1	16.9	0.1	0.8	0.0	0.4
Middle Atlantic	8.9	8.8	0.1	0.2	0.0	0.1
Midwest	10.0	10.3	0.3	0.3	0.1	0.0
East North Central	10.1	10.5	0.3	0.4	0.1	0.0
West North Central	9.9	10.0	0.1	0.2	*	0.0
South	8.8	8.6	0.2	0.2	0.1	0.0
South Atlantic	9.6	9.7	0.3	0.3	0.2	0.0
East South Central	7.0	6.4	0.3	0.1	*	0.1
West South Central	8.4	8.0	0.1	0.2	*	0.0
West	15.3	16.8	0.3 ^a	0.6	0.1	0.2
Mountain	14.6	15.7	0.4	0.4	0.1	0.2
Pacific	15.6	17.2	0.2 ^a	0.7	0.1	0.2
COUNTY TYPE						
Large Metro	11.9	12.2	0.2 ^b	0.5	0.1	0.2
Small Metro	11.0	11.6	0.2	0.3	0.1	0.0
250K - 1 Mil. Pop.	11.3	11.3	0.2	0.3	0.1	0.0
<250K Pop.	10.2	12.3	0.4	0.2	0.2	0.0
Nonmetro	7.6	8.4	0.2	0.1	0.0	0.0
Urbanized	9.1	11.7	0.2	0.1	0.1	0.0
Less Urbanized	7.0	7.1	0.2	0.2	0.1	*
Completely Rural	7.0	6.5	0.1	*	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.90A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	23,744	24,778	2,302	2,575	668	701
GEOGRAPHIC DIVISION						
Northeast	4,475	4,781	419 ^a	569	121	158
New England	1,566	1,860	130 ^a	207	37	68
Middle Atlantic	2,909	2,921	290	362	85	90
Midwest	5,252	5,399	592	546	163	125
East North Central	3,704	3,878	434	419	135	91
West North Central	1,548	1,521	158	127	28	34
South	6,961	6,932	728	774	220	227
South Atlantic	3,957	3,989	437	460	152	140
East South Central	955	984	100	105	18	35
West South Central	2,048	1,959	191	209	50	52
West	7,057	7,666	563	685	164	191
Mountain	1,979	2,197	196	206	45	68
Pacific	5,078	5,469	367	480	118	123
COUNTY TYPE						
Large Metro	12,840	12,978	1,114 ^a	1,363	343	397
Small Metro	7,449	7,954	807	872	242	225
250K - 1 Mil. Pop.	5,651	5,787	545	619	164	137
<250K Pop.	1,798	2,167	262	253	78	88
Nonmetro	3,455	3,845	381	340	82	79
Urbanized	1,234	1,547	129	153	39	47
Less Urbanized	1,842	1,968	214	169	36	31
Completely Rural	379	330	39	18	7	1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.90B Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	12.0	12.4	1.2	1.3	0.3	0.4
GEOGRAPHIC DIVISION						
Northeast	11.6	12.4	1.1 ^a	1.5	0.3	0.4
New England	15.6	18.4	1.3 ^a	2.1	0.4	0.7
Middle Atlantic	10.2	10.2	1.0	1.3	0.3	0.3
Midwest	11.4	11.6	1.3	1.2	0.4	0.3
East North Central	11.4	11.9	1.3	1.3	0.4	0.3
West North Central	11.3	11.0	1.1	0.9	0.2	0.2
South	9.9	9.7	1.0	1.1	0.3	0.3
South Atlantic	10.8	10.8	1.2	1.2	0.4	0.4
East South Central	7.8	7.9	0.8	0.8	0.1	0.3
West South Central	9.6	9.0	0.9	1.0	0.2	0.2
West	16.4	17.5	1.3	1.6	0.4	0.4
Mountain	16.0	17.3	1.6	1.6	0.4	0.5
Pacific	16.5	17.6	1.2	1.5	0.4	0.4
COUNTY TYPE						
Large Metro	13.0	13.3	1.1 ^a	1.4	0.3	0.4
Small Metro	12.5	12.9	1.4	1.4	0.4	0.4
250K - 1 Mil. Pop.	12.6	12.7	1.2	1.4	0.4	0.3
<250K Pop.	12.0	13.7	1.8	1.6	0.5	0.6
Nonmetro	8.8	9.4	1.0	0.8	0.2	0.2
Urbanized	10.6 ^a	12.9	1.1	1.3	0.3	0.4
Less Urbanized	8.2	8.2	1.0	0.7	0.2	0.1
Completely Rural	7.3	7.1	0.7	0.4	0.1	0.0

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.91A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	17,138	16,702	1,937	1,918	570	622
GEOGRAPHIC DIVISION						
Northeast	3,309	3,300	391	437	104	160
New England	1,071	1,144	101	129	18	46
Middle Atlantic	2,238	2,157	290	308	85	114
Midwest	4,233	3,754	456	429	138	129
East North Central	3,166	2,816	361	330	115	110
West North Central	1,067	938	95	99	23	19
South	5,129	5,230	587	599	157	181
South Atlantic	2,803	3,000	322	319	85	93
East South Central	918	848	82	114	15	40
West South Central	1,408	1,382	183	165	57	48
West	4,466	4,418	502	453	172	152
Mountain	1,292	1,227	146	113	62	24
Pacific	3,174	3,191	356	340	110	128
COUNTY TYPE						
Large Metro	9,127	8,552	971	1,002	322	367
Small Metro	5,269	5,359	638	582	153	177
250K - 1 Mil. Pop.	3,963	3,929	440	428	119	134
<250K Pop.	1,306	1,430	198	154	35	43
Nonmetro	2,742	2,791	328	333	95	78
Urbanized	1,046	945	106	117	24	32
Less Urbanized	1,458	1,536	179	181	51	40
Completely Rural	238	310	43	35	19	5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.91B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	7.8	7.5	0.9	0.9	0.3	0.3
GEOGRAPHIC DIVISION						
Northeast	7.8	7.7	0.9	1.0	0.2	0.4
New England	9.6	10.2	0.9	1.2	0.2	0.4
Middle Atlantic	7.1	6.8	0.9	1.0	0.3	0.4
Midwest	8.2 ^a	7.2	0.9	0.8	0.3	0.2
East North Central	8.8	7.8	1.0	0.9	0.3	0.3
West North Central	6.9	6.0	0.6	0.6	0.1	0.1
South	6.5	6.6	0.7	0.8	0.2	0.2
South Atlantic	6.9	7.3	0.8	0.8	0.2	0.2
East South Central	6.7	6.1	0.6	0.8	0.1	0.3
West South Central	5.8	5.6	0.8	0.7	0.2	0.2
West	9.2	9.0	1.0	0.9	0.4	0.3
Mountain	9.2	8.6	1.0	0.8	0.4	0.2
Pacific	9.2	9.2	1.0	1.0	0.3	0.4
COUNTY TYPE						
Large Metro	8.3	7.8	0.9	0.9	0.3	0.3
Small Metro	7.9	7.8	1.0	0.8	0.2	0.3
250K - 1 Mil. Pop.	7.9	7.7	0.9	0.8	0.2	0.3
<250K Pop.	7.8	8.1	1.2	0.9	0.2	0.2
Nonmetro	6.2	6.1	0.7	0.7	0.2	0.2
Urbanized	7.9	7.1	0.8	0.9	0.2	0.2
Less Urbanized	5.8	5.7	0.7	0.7	0.2	0.1
Completely Rural	4.2	5.9	0.7	0.7	0.3	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.92A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	2,118	2,079	908	826	245	223
GEOGRAPHIC DIVISION						
Northeast	360	368	167	157	40	33
New England	110	111	53	57	8	13
Middle Atlantic	250	257	113	100	32	20
Midwest	486	483	215	182	61	54
East North Central	335	346	155	136	46	41
West North Central	152	137	60	46	15	13
South	713	729	292	286	71	78
South Atlantic	335	354	135	140	29	34
East South Central	128	127	51	53	11	14
West South Central	250	248	106	93	32	30
West	559	500	235	202	73	58
Mountain	189	160	69	58	23	19
Pacific	369	340	165	143	50	39
COUNTY TYPE						
Large Metro	981	947	399	383	110	108
Small Metro	660	649	301 ^a	245	78	62
250K - 1 Mil. Pop.	481	462	213	180	60	49
<250K Pop.	179	187	88	65	18	13
Nonmetro	477	483	208	198	56	53
Urbanized	152	123	56	55	11	21
Less Urbanized	266	282	125	116	39	27
Completely Rural	59	78	26	26	6	5

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.92B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	9.1	8.9	3.9	3.5	1.1	1.0
GEOGRAPHIC DIVISION						
Northeast	8.7	8.8	4.0	3.7	1.0	0.8
New England	10.1	10.0	4.9	5.1	0.7	1.1
Middle Atlantic	8.2	8.3	3.7	3.2	1.0	0.7
Midwest	8.9	8.8	3.9	3.3	1.1	1.0
East North Central	8.9	9.2	4.1	3.6	1.2	1.1
West North Central	9.0	8.1	3.5	2.7	0.9	0.8
South	8.6	8.8	3.5	3.4	0.9	0.9
South Atlantic	8.4	8.7	3.4	3.4	0.7	0.8
East South Central	9.0	8.9	3.6	3.7	0.7	1.0
West South Central	8.9	8.8	3.8	3.3	1.1	1.1
West	10.5	9.3	4.4	3.7	1.4	1.1
Mountain	11.8	9.9	4.3	3.6	1.4	1.1
Pacific	9.9	9.0	4.4	3.8	1.3	1.0
COUNTY TYPE						
Large Metro	8.5	8.3	3.5	3.4	1.0	1.0
Small Metro	9.4	9.0	4.3 ^a	3.4	1.1	0.9
250K - 1 Mil. Pop.	9.1	8.6	4.0	3.3	1.1	0.9
<250K Pop.	10.5	10.3	5.2 ^a	3.6	1.1	0.7
Nonmetro	10.3	10.1	4.5	4.1	1.2	1.1
Urbanized	10.2	9.3	3.8	4.2	0.7	1.6
Less Urbanized	10.2	9.8	4.8	4.0	1.5	0.9
Completely Rural	10.7	13.1	4.8	4.4	1.1	0.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.93A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	4,014 ^a	3,701	747	696	182	162
GEOGRAPHIC DIVISION						
Northeast	808	755	191	183	44	45
New England	213	234	42	48	8	13
Middle Atlantic	595	521	149	135	37	32
Midwest	1,017	902	189	159	49	38
East North Central	759	699	154	131	41	33
West North Central	258	203	35	29	8	6
South	1,198	1,156	218	190	46	41
South Atlantic	638	628	139	116	32	27
East South Central	203	211	27	26	4	4
West South Central	357	317	52	47	9	10
West	991	888	149	164	43	38
Mountain	295	321	34	38	3	4
Pacific	697	568	114	126	40	34
COUNTY TYPE						
Large Metro	1,946 ^a	1,703	385	344	87	83
Small Metro	1,365	1,316	262	238	66	59
250K - 1 Mil. Pop.	965	924	175	175	49	52
<250K Pop.	400	393	87	63	16	7
Nonmetro	703	683	100	114	29	20
Urbanized	282	282	42	57	11	11
Less Urbanized	368	355	45	48	9	8
Completely Rural	52	46	13	9	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.93B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	14.1 ^b	12.8	2.6	2.4	0.6	0.6
GEOGRAPHIC DIVISION						
Northeast	16.0	14.8	3.8	3.6	0.9	0.9
New England	16.5	17.7	3.3	3.6	0.6	1.0
Middle Atlantic	15.8	13.7	4.0	3.5	1.0	0.8
Midwest	15.0 ^a	13.1	2.8	2.3	0.7	0.6
East North Central	16.0	14.5	3.2	2.7	0.9	0.7
West North Central	12.6 ^a	9.7	1.7	1.4	0.4	0.3
South	11.8	11.2	2.2	1.8	0.5	0.4
South Atlantic	13.2	12.8	2.9	2.4	0.7	0.5
East South Central	11.2	11.5	1.5	1.4	0.2	0.2
West South Central	10.3	9.0	1.5	1.3	0.3	0.3
West	15.2	13.3	2.3	2.5	0.7	0.6
Mountain	15.1	15.9	1.8	1.9	0.2	0.2
Pacific	15.3 ^a	12.2	2.5	2.7	0.9	0.7
COUNTY TYPE						
Large Metro	14.0 ^a	12.3	2.8	2.5	0.6	0.6
Small Metro	14.9	13.9	2.9	2.5	0.7	0.6
250K - 1 Mil. Pop.	15.0	13.9	2.7	2.6	0.8	0.8
<250K Pop.	14.6	13.8	3.2	2.2	0.6	0.3
Nonmetro	13.1	12.2	1.9	2.0	0.5	0.3
Urbanized	14.6	15.0	2.2	3.0	0.6	0.6
Less Urbanized	12.6	11.0	1.5	1.5	0.3	0.3
Completely Rural	9.8	9.4	2.4	1.8	*	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.94A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	11,006	10,922	282	396	144	236
GEOGRAPHIC DIVISION						
Northeast	2,141	2,178	34	98	20	82
New England	749	799	6	24	3	21
Middle Atlantic	1,392	1,378	28	74	17	61
Midwest	2,730	2,369	52	87	28	36
East North Central	2,073	1,771	52	63	28	36
West North Central	658	598	*	24	*	*
South	3,218	3,345	78	124	40	62
South Atlantic	1,829	2,017	49	63	24	32
East South Central	587	511	4	35	*	21
West South Central	802	817	25	25	16	9
West	2,916	3,030	119	87	56	56
Mountain	808	746	42	17	36	2
Pacific	2,108	2,283	77	71	20	55
COUNTY TYPE						
Large Metro	6,200	5,903	187	275	125	176
Small Metro	3,244	3,393	75	99	9	56
250K - 1 Mil. Pop.	2,517	2,543	52	73	9	34
<250K Pop.	727	850	23	26	*	22
Nonmetro	1,562	1,626	20	21	10	5
Urbanized	611	540	7	5	3	*
Less Urbanized	823	899	10	16	3	5
Completely Rural	127	186	3	*	3	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.94B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	6.5	6.4	0.2	0.2	0.1	0.1
GEOGRAPHIC DIVISION						
Northeast	6.4	6.5	0.1	0.3	0.1	0.2
New England	8.5	9.1	0.1	0.3	0.0	0.2
Middle Atlantic	5.6	5.6	0.1	0.3	0.1	0.2
Midwest	6.9	6.0	0.1	0.2	0.1	0.1
East North Central	7.5	6.4	0.2	0.2	0.1	0.1
West North Central	5.6	5.1	*	0.2	*	*
South	5.4	5.5	0.1	0.2	0.1	0.1
South Atlantic	5.8	6.3	0.2	0.2	0.1	0.1
East South Central	5.6	4.8	0.0	0.3	*	0.2
West South Central	4.5	4.5	0.1	0.1	0.1	0.0
West	8.0	8.2	0.3	0.2	0.2	0.2
Mountain	7.8	7.0	0.4	0.2	0.3	0.0
Pacific	8.0	8.7	0.3	0.3	0.1	0.2
COUNTY TYPE						
Large Metro	7.3	7.0	0.2	0.3	0.1	0.2
Small Metro	6.4	6.5	0.1	0.2	0.0	0.1
250K - 1 Mil. Pop.	6.6	6.5	0.1	0.2	0.0	0.1
<250K Pop.	5.9	6.5	0.2	0.2	*	0.2
Nonmetro	4.6	4.6	0.1	0.1	0.0	0.0
Urbanized	6.3	5.4	0.1	0.1	0.0	*
Less Urbanized	4.2	4.3	0.0	0.1	0.0	0.0
Completely Rural	2.7	4.4	0.1	*	0.1	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.95A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15,020	14,623	1,029	1,092	326	398
GEOGRAPHIC DIVISION						
Northeast	2,949	2,932	225	280	64	127
New England	961	1,033	48	72	11	34
Middle Atlantic	1,988	1,899	177	208	53	93
Midwest	3,747	3,271	241	247	77	74
East North Central	2,832	2,470	206	194	69	69
West North Central	915	801	35	53	8	6
South	4,416	4,501	296	313	86	103
South Atlantic	2,467	2,646	188	179	56	59
East South Central	790	722	31	61	4	25
West South Central	1,159	1,134	77	73	25	19
West	3,908	3,918	267	251	99	95
Mountain	1,103	1,067	76	55	39	6
Pacific	2,805	2,851	191	197	60	89
COUNTY TYPE						
Large Metro	8,146	7,605	572	619	212	259
Small Metro	4,609	4,709	337	337	75	115
250K - 1 Mil. Pop.	3,482	3,467	227	249	58	85
<250K Pop.	1,127	1,243	110	89	16	29
Nonmetro	2,265	2,308	120	135	39	25
Urbanized	894	822	49	62	14	11
Less Urbanized	1,192	1,254	54	64	12	13
Completely Rural	179	232	16	9	13	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.95B Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	7.6	7.3	0.5	0.5	0.2	0.2
GEOGRAPHIC DIVISION						
Northeast	7.7	7.6	0.6	0.7	0.2	0.3
New England	9.6	10.2	0.5	0.7	0.1	0.3
Middle Atlantic	7.0	6.7	0.6	0.7	0.2	0.3
Midwest	8.1 ^a	7.0	0.5	0.5	0.2	0.2
East North Central	8.8	7.6	0.6	0.6	0.2	0.2
West North Central	6.7	5.8	0.3	0.4	0.1	0.0
South	6.3	6.3	0.4	0.4	0.1	0.1
South Atlantic	6.8	7.1	0.5	0.5	0.2	0.2
East South Central	6.4	5.8	0.3	0.5	0.0	0.2
West South Central	5.4	5.2	0.4	0.3	0.1	0.1
West	9.1	9.0	0.6	0.6	0.2	0.2
Mountain	8.9	8.4	0.6	0.4	0.3	0.0
Pacific	9.1	9.2	0.6	0.6	0.2	0.3
COUNTY TYPE						
Large Metro	8.2	7.8	0.6	0.6	0.2	0.3
Small Metro	7.7	7.7	0.6	0.5	0.1	0.2
250K - 1 Mil. Pop.	7.8	7.6	0.5	0.5	0.1	0.2
<250K Pop.	7.5	7.8	0.7	0.6	0.1	0.2
Nonmetro	5.8	5.7	0.3	0.3	0.1	0.1
Urbanized	7.7	6.9	0.4	0.5	0.1	0.1
Less Urbanized	5.3	5.2	0.2	0.3	0.1	0.1
Completely Rural	3.5	5.0	0.3	0.2	0.3	*

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.96A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	34,076	32,443	9,220	8,761	3,952	3,849
GEOGRAPHIC DIVISION						
Northeast	6,540 ^a	5,751	1,503	1,664	754	649
New England	2,206	2,098	535	580	305 ^a	196
Middle Atlantic	4,334 ^a	3,653	969	1,084	448	453
Midwest	7,646 ^b	6,640	2,154 ^b	1,661	892	733
East North Central	5,281	4,944	1,463	1,274	632	608
West North Central	2,365 ^b	1,696	691 ^b	387	260 ^b	126
South	11,601	10,987	3,348	3,116	1,388	1,404
South Atlantic	6,021	5,645	1,530	1,534	571	643
East South Central	1,998	1,979	730	593	318	299
West South Central	3,583	3,362	1,088	990	499	462
West	8,289	9,065	2,215	2,320	918	1,062
Mountain	2,461	2,458	752	657	277	288
Pacific	5,827	6,607	1,463	1,663	641	774
COUNTY TYPE						
Large Metro	17,304	16,289	4,383	4,107	1,973	1,815
Small Metro	11,034	10,332	3,096	3,103	1,269	1,395
250K - 1 Mil. Pop.	8,488	7,734	2,468	2,254	1,033	945
<250K Pop.	2,547	2,598	628 ^a	849	236 ^b	450
Nonmetro	5,738	5,822	1,741	1,552	709	639
Urbanized	1,855	1,793	523	427	224	201
Less Urbanized	3,271	3,341	1,063	977	399	376
Completely Rural	612	689	155	148	86	62

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.96B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15.4 ^a	14.5	4.2	3.9	1.8	1.7
GEOGRAPHIC DIVISION						
Northeast	15.3 ^a	13.4	3.5	3.9	1.8	1.5
New England	19.8	18.7	4.8	5.2	2.7 ^a	1.7
Middle Atlantic	13.7 ^a	11.6	3.1	3.4	1.4	1.4
Midwest	14.8 ^b	12.8	4.2 ^b	3.2	1.7	1.4
East North Central	14.6	13.6	4.0	3.5	1.7	1.7
West North Central	15.3 ^b	10.9	4.5 ^b	2.5	1.7 ^b	0.8
South	14.8	13.8	4.3	3.9	1.8	1.8
South Atlantic	14.9	13.7	3.8	3.7	1.4	1.6
East South Central	14.6	14.3	5.3	4.3	2.3	2.2
West South Central	14.8	13.7	4.5	4.0	2.1	1.9
West	17.1	18.5	4.6	4.7	1.9	2.2
Mountain	17.6	17.2	5.4	4.6	2.0	2.0
Pacific	16.9	19.0	4.2	4.8	1.9	2.2
COUNTY TYPE						
Large Metro	15.7	14.9	4.0	3.8	1.8	1.7
Small Metro	16.6 ^a	15.1	4.6	4.5	1.9	2.0
250K - 1 Mil. Pop.	17.0 ^a	15.2	4.9	4.4	2.1	1.9
<250K Pop.	15.3	14.7	3.8 ^a	4.8	1.4 ^b	2.5
Nonmetro	13.0	12.8	4.0	3.4	1.6	1.4
Urbanized	14.1	13.5	4.0	3.2	1.7	1.5
Less Urbanized	13.0	12.4	4.2	3.6	1.6	1.4
Completely Rural	10.7	13.1	2.7	2.8	1.5	1.2

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.97A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	2,524	2,543	1,653	1,649	664	690
GEOGRAPHIC DIVISION						
Northeast	398	418	270	279	99	109
New England	131	139	98	90	32	32
Middle Atlantic	266	279	172	189	67	77
Midwest	546	528	350	337	145	132
East North Central	399	403	254	266	110	101
West North Central	147	126	96	71	35	31
South	958	1,025	635	670	276	284
South Atlantic	447	488	278	310	111	133
East South Central	178	219	123	141	56	51
West South Central	333	318	235	219	109	100
West	622	571	398	363	144	167
Mountain	184	157	125	97	44	51
Pacific	438	415	273	265	100	116
COUNTY TYPE						
Large Metro	1,159	1,104	761	712	273	300
Small Metro	819	859	526	587	239	243
250K - 1 Mil. Pop.	614	614	406	429	187	182
<250K Pop.	206	245	120	158	52	62
Nonmetro	545	580	366	351	153	147
Urbanized	161	152	108	95	49	40
Less Urbanized	317	352	213	213	85	90
Completely Rural	67	76	45	43	18	16

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.97B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	10.9	10.9	7.1	7.1	2.9	3.0
GEOGRAPHIC DIVISION						
Northeast	9.6	9.9	6.5	6.6	2.4	2.6
New England	12.0	12.5	8.9	8.1	2.9	2.8
Middle Atlantic	8.7	9.0	5.6	6.1	2.2	2.5
Midwest	10.0	9.7	6.4	6.2	2.6	2.4
East North Central	10.6	10.7	6.7	7.1	2.9	2.7
West North Central	8.7	7.4	5.7	4.2	2.1	1.8
South	11.6	12.3	7.7	8.1	3.4	3.4
South Atlantic	11.2	12.0	6.9	7.6	2.8	3.3
East South Central	12.5	15.4	8.6	9.9	4.0	3.6
West South Central	11.8	11.3	8.3	7.7	3.9	3.5
West	11.7	10.6	7.5	6.7	2.7	3.1
Mountain	11.4	9.7	7.8	6.0	2.7	3.1
Pacific	11.8	11.0	7.3	7.0	2.7	3.1
COUNTY TYPE						
Large Metro	10.0	9.7	6.6	6.3	2.4	2.6
Small Metro	11.7	11.9	7.5	8.1	3.4	3.4
250K - 1 Mil. Pop.	11.6	11.4	7.7	8.0	3.5	3.4
<250K Pop.	12.1	13.5	7.0	8.7	3.1	3.4
Nonmetro	11.7	12.1	7.9	7.3	3.3	3.1
Urbanized	10.8	11.5	7.2	7.2	3.3	3.0
Less Urbanized	12.2	12.2	8.2	7.4	3.3	3.1
Completely Rural	12.1	12.7	8.2	7.2	3.3	2.8

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.98A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	5,957	5,650	2,830	2,696	1,062	1,050
GEOGRAPHIC DIVISION						
Northeast	949	988	434	482	142	198
New England	273	335	146	144	47	57
Middle Atlantic	676	653	288	338	96	140
Midwest	1,377	1,248	653 ^a	548	234	194
East North Central	930	948	460	412	171	157
West North Central	448 ^b	301	193	136	63	37
South	2,170	2,009	1,083	1,035	445	444
South Atlantic	1,017	896	510	454	189	183
East South Central	413	438	210	223	96	115
West South Central	741	675	363	358	161	146
West	1,460	1,405	660	632	241	215
Mountain	457	468	192	202	64	63
Pacific	1,002	937	468	430	177	152
COUNTY TYPE						
Large Metro	2,827 ^a	2,549	1,302	1,221	469	450
Small Metro	2,066	2,022	1,016	988	397	410
250K - 1 Mil. Pop.	1,469	1,443	723	715	289	296
<250K Pop.	597	580	293	273	108	114
Nonmetro	1,064	1,078	512	487	195	190
Urbanized	370	365	180	158	66	69
Less Urbanized	601	629	292	294	112	104
Completely Rural	93	85	40	36	17	17

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.98B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	20.9 ^b	19.5	9.9	9.3	3.7	3.6
GEOGRAPHIC DIVISION						
Northeast	18.8	19.3	8.6	9.4	2.8	3.9
New England	21.2	25.4	11.4	10.9	3.6	4.4
Middle Atlantic	18.0	17.2	7.7	8.9	2.5	3.7
Midwest	20.3 ^a	18.1	9.6 ^a	7.9	3.4	2.8
East North Central	19.6	19.7	9.7	8.6	3.6	3.3
West North Central	21.8 ^b	14.3	9.4 ^a	6.5	3.1	1.8
South	21.5 ^a	19.5	10.7	10.1	4.4	4.3
South Atlantic	21.0 ^a	18.2	10.6	9.2	3.9	3.7
East South Central	22.7	23.9	11.6	12.2	5.3	6.3
West South Central	21.4	19.1	10.5	10.1	4.6	4.1
West	22.4	21.0	10.1	9.5	3.7	3.2
Mountain	23.5	23.2	9.8	10.0	3.3	3.1
Pacific	22.0	20.1	10.3	9.2	3.9	3.3
COUNTY TYPE						
Large Metro	20.3 ^a	18.3	9.4	8.8	3.4	3.2
Small Metro	22.5	21.3	11.1	10.4	4.3	4.3
250K - 1 Mil. Pop.	22.8	21.7	11.2	10.8	4.5	4.5
<250K Pop.	21.8	20.3	10.7	9.6	3.9	4.0
Nonmetro	19.8	19.3	9.5	8.7	3.6	3.4
Urbanized	19.1	19.4	9.3	8.4	3.4	3.7
Less Urbanized	20.6	19.4	10.0	9.1	3.9	3.2
Completely Rural	17.4	17.5	7.5	7.4	3.3	3.5

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.99A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	25,596	24,250	4,737	4,415	2,225	2,108
GEOGRAPHIC DIVISION						
Northeast	5,193 ^a	4,345	799	903	513	343
New England	1,802	1,624	291	346	227 ^a	107
Middle Atlantic	3,391 ^a	2,721	509	557	286	236
Midwest	5,723 ^a	4,863	1,151 ^b	776	514	408
East North Central	3,953	3,594	749	595	351	350
West North Central	1,770 ^b	1,270	402 ^a	181	163 ^a	58
South	8,473	7,953	1,630	1,411	666	677
South Atlantic	4,557	4,261	742	770	270	328
East South Central	1,407	1,322	397	229	166	133
West South Central	2,510	2,370	491	413	229	216
West	6,207	7,089	1,157	1,325	533	681
Mountain	1,820	1,834	435	357	169	175
Pacific	4,387 ^a	5,255	723	968	364	506
COUNTY TYPE						
Large Metro	13,318	12,635	2,320	2,174	1,231	1,064
Small Metro	8,149	7,451	1,554	1,528	633	742
250K - 1 Mil. Pop.	6,405	5,677	1,339	1,110	557	467
<250K Pop.	1,744	1,774	215 ^a	418	75 ^b	274
Nonmetro	4,129	4,164	863	713	361	302
Urbanized	1,324	1,275	235	174	110	92
Less Urbanized	2,352	2,361	558	469	201	182
Completely Rural	453	528	70	70	51	29

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.99B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15.1	14.2	2.8	2.6	1.3	1.2
GEOGRAPHIC DIVISION						
Northeast	15.5 ^a	13.0	2.4	2.7	1.5	1.0
New England	20.6	18.5	3.3	3.9	2.6 ^a	1.2
Middle Atlantic	13.7 ^a	11.0	2.1	2.3	1.2	1.0
Midwest	14.6 ^b	12.3	2.9 ^b	2.0	1.3	1.0
East North Central	14.3	13.0	2.7	2.1	1.3	1.3
West North Central	15.1 ^b	10.8	3.4 ^b	1.5	1.4 ^a	0.5
South	14.1	13.1	2.7	2.3	1.1	1.1
South Atlantic	14.4	13.3	2.3	2.4	0.9	1.0
East South Central	13.4	12.5	3.8	2.2	1.6	1.3
West South Central	14.0	13.0	2.7	2.3	1.3	1.2
West	17.0	19.1	3.2	3.6	1.5	1.8
Mountain	17.5	17.2	4.2	3.3	1.6	1.6
Pacific	16.7 ^a	19.9	2.8	3.7	1.4	1.9
COUNTY TYPE						
Large Metro	15.7	15.1	2.7	2.6	1.4	1.3
Small Metro	16.1 ^a	14.3	3.1	2.9	1.3	1.4
250K - 1 Mil. Pop.	16.7 ^a	14.6	3.5	2.8	1.5	1.2
<250K Pop.	14.3	13.7	1.8 ^a	3.2	0.6 ^b	2.1
Nonmetro	12.2	11.8	2.5	2.0	1.1	0.9
Urbanized	13.6	12.7	2.4	1.7	1.1	0.9
Less Urbanized	12.0	11.3	2.9	2.2	1.0	0.9
Completely Rural	9.7	12.6	1.5	1.7	1.1	0.7

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.100A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	31,552	29,900	7,567	7,112	3,287	3,158
GEOGRAPHIC DIVISION						
Northeast	6,142 ^a	5,334	1,233	1,385	655	541
New England	2,075	1,960	437	489	273 ^a	164
Middle Atlantic	4,067 ^a	3,374	796	895	381	376
Midwest	7,100 ^b	6,112	1,804 ^b	1,324	747	602
East North Central	4,882	4,541	1,209	1,007	522	507
West North Central	2,218 ^b	1,570	595 ^b	317	225 ^b	95
South	10,643	9,961	2,713	2,446	1,111	1,121
South Atlantic	5,573	5,157	1,252	1,223	459	511
East South Central	1,820	1,760	607	451	262	248
West South Central	3,250	3,045	853	771	390	362
West	7,666	8,493	1,817	1,957	774	895
Mountain	2,277	2,301	627	559	233	237
Pacific	5,389	6,192	1,190	1,398	541	658
COUNTY TYPE						
Large Metro	16,144	15,185	3,622	3,395	1,700	1,514
Small Metro	10,215	9,473	2,570	2,516	1,030	1,152
250K - 1 Mil. Pop.	7,874	7,120	2,062	1,825	847	764
<250K Pop.	2,341	2,353	508	691	184 ^b	388
Nonmetro	5,193	5,242	1,375	1,201	557	492
Urbanized	1,694	1,640	416	332	176	161
Less Urbanized	2,953	2,989	850	763	313	286
Completely Rural	545	613	110	105	68	46

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.100B Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	15.9 ^a	15.0	3.8	3.6	1.7	1.6
GEOGRAPHIC DIVISION						
Northeast	15.9 ^a	13.8	3.2	3.6	1.7	1.4
New England	20.6	19.4	4.3	4.8	2.7 ^a	1.6
Middle Atlantic	14.3 ^a	11.8	2.8	3.1	1.3	1.3
Midwest	15.4 ^b	13.2	3.9 ^b	2.9	1.6	1.3
East North Central	15.1	14.0	3.7	3.1	1.6	1.6
West North Central	16.1 ^b	11.3	4.3 ^b	2.3	1.6 ^b	0.7
South	15.2	14.0	3.9	3.4	1.6	1.6
South Atlantic	15.3	13.9	3.4	3.3	1.3	1.4
East South Central	14.8	14.1	4.9	3.6	2.1	2.0
West South Central	15.2	14.0	4.0	3.6	1.8	1.7
West	17.8	19.4	4.2	4.5	1.8	2.0
Mountain	18.4	18.1	5.1	4.4	1.9	1.9
Pacific	17.5	19.9	3.9	4.5	1.8	2.1
COUNTY TYPE						
Large Metro	16.3	15.5	3.7	3.5	1.7	1.5
Small Metro	17.1 ^a	15.4	4.3	4.1	1.7	1.9
250K - 1 Mil. Pop.	17.6 ^a	15.6	4.6	4.0	1.9	1.7
<250K Pop.	15.6	14.9	3.4	4.4	1.2 ^b	2.5
Nonmetro	13.2	12.9	3.5	2.9	1.4	1.2
Urbanized	14.5	13.7	3.6	2.8	1.5	1.3
Less Urbanized	13.1	12.4	3.8	3.2	1.4	1.2
Completely Rural	10.5	13.1	2.1	2.3	1.3	1.0

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.101A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	53,268	52,605	13,444	13,052	5,981	5,711
GEOGRAPHIC DIVISION						
Northeast	10,524	9,871	2,403	2,625	1,183	1,066
New England	3,353	3,424	786	938	433	341
Middle Atlantic	7,171	6,447	1,618	1,686	749	725
Midwest	12,061	11,387	3,086 ^b	2,597	1,326	1,111
East North Central	8,454	8,423	2,155	1,975	946	898
West North Central	3,607 ^b	2,964	932 ^b	622	380 ^b	214
South	17,285	16,950	4,676	4,399	2,029	1,940
South Atlantic	9,445	9,045	2,285	2,250	960	928
East South Central	2,676	2,846	889	796	396	395
West South Central	5,164	5,059	1,502	1,353	673	617
West	13,398	14,397	3,278	3,432	1,444	1,593
Mountain	3,838	4,128	1,089	1,028	440	434
Pacific	9,560	10,269	2,189	2,404	1,004	1,159
COUNTY TYPE						
Large Metro	27,821	27,044	6,669	6,449	3,140	2,927
Small Metro	16,724	16,558	4,418	4,455	1,875	1,925
250K - 1 Mil. Pop.	12,787	12,375	3,388	3,260	1,522	1,345
<250K Pop.	3,937	4,183	1,031	1,195	353 ^b	581
Nonmetro	8,723	9,003	2,357	2,148	966	859
Urbanized	2,965	2,984	757	668	335	280
Less Urbanized	4,782	5,114	1,390	1,283	524	508
Completely Rural	976	906	210	198	108	70

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.101B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	24.1	23.6	6.1	5.8	2.7	2.6
GEOGRAPHIC DIVISION						
Northeast	24.7	23.1	5.6	6.1	2.8	2.5
New England	30.1	30.6	7.0	8.4	3.9	3.0
Middle Atlantic	22.7	20.4	5.1	5.3	2.4	2.3
Midwest	23.4	22.0	6.0 ^b	5.0	2.6	2.1
East North Central	23.4	23.2	6.0	5.4	2.6	2.5
West North Central	23.4 ^b	19.0	6.0 ^b	4.0	2.5 ^b	1.4
South	22.0	21.3	6.0	5.5	2.6	2.4
South Atlantic	23.3	22.0	5.6	5.5	2.4	2.3
East South Central	19.5	20.5	6.5	5.7	2.9	2.9
West South Central	21.3	20.6	6.2	5.5	2.8	2.5
West	27.7	29.3	6.8	7.0	3.0	3.2
Mountain	27.5	28.8	7.8	7.2	3.1	3.0
Pacific	27.7	29.5	6.3	6.9	2.9	3.3
COUNTY TYPE						
Large Metro	25.2	24.8	6.0	5.9	2.8	2.7
Small Metro	25.1	24.1	6.6	6.5	2.8	2.8
250K - 1 Mil. Pop.	25.6	24.3	6.8	6.4	3.0	2.6
<250K Pop.	23.6	23.7	6.2	6.8	2.1 ^b	3.3
Nonmetro	19.8	19.8	5.4	4.7	2.2	1.9
Urbanized	22.5	22.5	5.7	5.0	2.5	2.1
Less Urbanized	19.1	18.9	5.5	4.8	2.1	1.9
Completely Rural	17.0	17.2	3.7	3.8	1.9	1.3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.102A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	4,248	4,228	2,698	2,631	1,045	1,074
GEOGRAPHIC DIVISION						
Northeast	743	712	503	448	182	160
New England	231	233	164	154	58	54
Middle Atlantic	512	479	339	294	124	106
Midwest	929	930	585	542	230	211
East North Central	657	668	427	408	174	156
West North Central	273	262	158	135	56	54
South	1,480	1,564	930	1,003	378	441
South Atlantic	693	769	425	481	152 ^a	207
East South Central	263	296	159	190	68	81
West South Central	523	500	346	332	157	153
West	1,096	1,022	680	638	255	263
Mountain	355	328	209	195	77	85
Pacific	741	694	471	442	178	178
COUNTY TYPE						
Large Metro	2,026	1,910	1,281	1,178	483	478
Small Metro	1,332	1,397	850	897	343	362
250K - 1 Mil. Pop.	994	1,002	636	651	267	268
<250K Pop.	337	395	213	246	76	95
Nonmetro	890	922	567	557	218	233
Urbanized	292	257	183	165	73	71
Less Urbanized	495	539	316	327	122	138
Completely Rural	104	126	68	65	23	25

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.102B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	18.3	18.1	11.6	11.3	4.5	4.6
GEOGRAPHIC DIVISION						
Northeast	17.9	16.9	12.1	10.7	4.4	3.8
New England	21.2	21.0	15.1	13.9	5.3	4.9
Middle Atlantic	16.7	15.5	11.1	9.5	4.1	3.4
Midwest	17.0	17.0	10.7	9.9	4.2	3.9
East North Central	17.4	17.7	11.3	10.8	4.6	4.1
West North Central	16.1	15.5	9.3	8.0	3.3	3.2
South	17.9	18.8	11.3	12.1	4.6	5.3
South Atlantic	17.3	18.9	10.6	11.8	3.8 ^a	5.1
East South Central	18.6	20.8	11.2	13.4	4.8	5.7
West South Central	18.6	17.7	12.3	11.8	5.6	5.4
West	20.5	19.0	12.8	11.8	4.8	4.9
Mountain	22.1	20.2	13.0	12.0	4.8	5.2
Pacific	19.9	18.4	12.7	11.7	4.8	4.7
COUNTY TYPE						
Large Metro	17.5	16.8	11.1	10.4	4.2	4.2
Small Metro	19.0	19.4	12.2	12.4	4.9	5.0
250K - 1 Mil. Pop.	18.8	18.6	12.0	12.1	5.0	5.0
<250K Pop.	19.8	21.8	12.5	13.6	4.5	5.2
Nonmetro	19.1	19.2	12.2	11.6	4.7	4.9
Urbanized	19.5	19.4	12.2	12.5	4.9	5.3
Less Urbanized	19.0	18.7	12.2	11.4	4.7	4.8
Completely Rural	18.8	21.1	12.3	10.8	4.2	4.1

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.103A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	9,476	9,247	4,460	4,295	1,729	1,711
GEOGRAPHIC DIVISION						
Northeast	1,746	1,751	795	840	273	327
New England	517	561	246	263	89	101
Middle Atlantic	1,229	1,190	549	577	184	226
Midwest	2,251	2,145	1,055	916	408	336
East North Central	1,547	1,585	750	685	296	265
West North Central	705 ^a	561	305	231	112	72
South	3,153	3,021	1,536	1,489	638	642
South Atlantic	1,546	1,459	740	698	314	300
East South Central	541	571	271	293	110	139
West South Central	1,066	991	524	497	214	202
West	2,326	2,330	1,074	1,051	410	406
Mountain	752	780	343	346	119	134
Pacific	1,574	1,551	732	704	291	272
COUNTY TYPE						
Large Metro	4,588	4,440	2,148	2,048	816	783
Small Metro	3,184	3,180	1,535	1,526	612	662
250K - 1 Mil. Pop.	2,257	2,267	1,079	1,079	438	472
<250K Pop.	927	914	456	448	174	190
Nonmetro	1,704	1,627	777	721	300	266
Urbanized	625	589	278	272	117	106
Less Urbanized	930	909	437	397	150	142
Completely Rural	149	128	62	52	33	17

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.103B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	33.3 ^a	31.9	15.7	14.8	6.1	5.9
GEOGRAPHIC DIVISION						
Northeast	34.6	34.3	15.8	16.4	5.4	6.4
New England	40.1	42.6	19.1	20.0	6.9	7.7
Middle Atlantic	32.7	31.4	14.6	15.2	4.9	5.9
Midwest	33.1	31.1	15.5 ^a	13.3	6.0	4.9
East North Central	32.6	33.0	15.8	14.3	6.2	5.5
West North Central	34.3 ^b	26.7	14.9 ^a	11.0	5.4 ^a	3.4
South	31.2	29.3	15.2	14.5	6.3	6.2
South Atlantic	32.0	29.6	15.3	14.2	6.5	6.1
East South Central	29.7	31.1	14.9	16.0	6.1	7.6
West South Central	30.8	28.0	15.1	14.1	6.2	5.7
West	35.7	34.9	16.5	15.7	6.3	6.1
Mountain	38.6	38.7	17.6	17.2	6.1	6.6
Pacific	34.5	33.3	16.0	15.1	6.4	5.8
COUNTY TYPE						
Large Metro	33.0	32.0	15.4	14.7	5.9	5.6
Small Metro	34.7	33.5	16.7	16.1	6.7	7.0
250K - 1 Mil. Pop.	35.1	34.2	16.8	16.3	6.8	7.1
<250K Pop.	33.8	32.0	16.6	15.7	6.3	6.7
Nonmetro	31.7 ^a	29.0	14.5	12.9	5.6	4.7
Urbanized	32.3	31.3	14.4	14.5	6.1	5.7
Less Urbanized	31.9 ^a	28.1	15.0 ^a	12.3	5.1	4.4
Completely Rural	28.0	26.5	11.6	10.7	6.2	3.6

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.104A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	39,545	39,130	6,285	6,125	3,208	2,926
GEOGRAPHIC DIVISION						
Northeast	8,036	7,409	1,105	1,336	728	579
New England	2,606	2,630	376	521	286	186
Middle Atlantic	5,430	4,778	729	815	442	393
Midwest	8,880	8,312	1,446	1,139	688	564
East North Central	6,250	6,171	977	882	475	477
West North Central	2,630 ^a	2,141	469 ^a	256	212 ^a	88
South	12,652	12,365	2,210	1,907	1,014	858
South Atlantic	7,206	6,817	1,119	1,071	494	421
East South Central	1,871	1,979	459	313	218	175
West South Central	3,575	3,568	631	523	302	262
West	9,976	11,044	1,524	1,743	779	925
Mountain	2,731	3,020	538	486	244	216
Pacific	7,245	8,024	986	1,257	534	709
COUNTY TYPE						
Large Metro	21,207	20,694	3,239	3,223	1,841	1,665
Small Metro	12,208	11,981	2,033	2,032	920	901
250K - 1 Mil. Pop.	9,536	9,106	1,672	1,531	816	605
<250K Pop.	2,673	2,875	362	501	103 ^b	296
Nonmetro	6,129	6,455	1,012	871	448	359
Urbanized	2,049	2,138	296	231	145	103
Less Urbanized	3,357	3,666	637	559	252	228
Completely Rural	724	652	80	81	51	29

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.104B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	23.3	22.9	3.7	3.6	1.9	1.7
GEOGRAPHIC DIVISION						
Northeast	24.0	22.1	3.3	4.0	2.2	1.7
New England	29.7	30.0	4.3	5.9	3.3	2.1
Middle Atlantic	22.0	19.3	3.0	3.3	1.8	1.6
Midwest	22.6	21.0	3.7 ^a	2.9	1.7	1.4
East North Central	22.6	22.3	3.5	3.2	1.7	1.7
West North Central	22.5 ^b	18.2	4.0 ^a	2.2	1.8 ^a	0.7
South	21.1	20.3	3.7	3.1	1.7	1.4
South Atlantic	22.8	21.2	3.5	3.3	1.6	1.3
East South Central	17.8	18.7	4.4	2.9	2.1	1.6
West South Central	19.9	19.6	3.5	2.9	1.7	1.4
West	27.3	29.8	4.2	4.7	2.1	2.5
Mountain	26.2	28.3	5.2	4.6	2.3	2.0
Pacific	27.7	30.4	3.8	4.8	2.0	2.7
COUNTY TYPE						
Large Metro	24.9	24.7	3.8	3.8	2.2	2.0
Small Metro	24.2	23.1	4.0	3.9	1.8	1.7
250K - 1 Mil. Pop.	24.9	23.4	4.4	3.9	2.1	1.6
<250K Pop.	21.9	22.2	3.0	3.9	0.8 ^b	2.3
Nonmetro	18.1	18.4	3.0	2.5	1.3	1.0
Urbanized	21.0	21.2	3.0	2.3	1.5	1.0
Less Urbanized	17.2	17.6	3.3	2.7	1.3	1.1
Completely Rural	15.6	15.6	1.7	1.9	1.1	0.7

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.105A Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	49,020	48,377	10,745	10,421	4,937	4,637
GEOGRAPHIC DIVISION						
Northeast	9,782	9,159	1,900	2,177	1,001	906
New England	3,122	3,192	621	784	376	287
Middle Atlantic	6,659	5,968	1,279	1,393	625	619
Midwest	11,132	10,457	2,502 ^a	2,055	1,095	901
East North Central	7,797	7,755	1,728	1,567	771	741
West North Central	3,335 ^b	2,702	774 ^b	487	324 ^a	160
South	15,805	15,386	3,745	3,395	1,652	1,500
South Atlantic	8,752	8,277	1,859	1,769	808	721
East South Central	2,412	2,550	730	606	328	314
West South Central	4,641	4,559	1,155	1,021	516	464
West	12,302	13,375	2,598	2,794	1,188	1,330
Mountain	3,483	3,800	880	832	363	350
Pacific	8,819	9,575	1,718	1,962	826	981
COUNTY TYPE						
Large Metro	25,795	25,135	5,387	5,271	2,657	2,448
Small Metro	15,393	15,161	3,569	3,558	1,532	1,563
250K - 1 Mil. Pop.	11,793	11,373	2,751	2,610	1,255	1,077
<250K Pop.	3,600	3,789	817	949	277 ^b	486
Nonmetro	7,833	8,082	1,790	1,592	748	625
Urbanized	2,674	2,727	574	502	262	209
Less Urbanized	4,287	4,575	1,074	956	402	370
Completely Rural	872	780	142	133	84	46

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.105B Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	24.8	24.2	5.4	5.2	2.5	2.3
GEOGRAPHIC DIVISION						
Northeast	25.4	23.7	4.9	5.6	2.6	2.3
New England	31.0	31.6	6.2	7.8	3.7	2.8
Middle Atlantic	23.4	20.9	4.5	4.9	2.2	2.2
Midwest	24.1 ^a	22.5	5.4 ^b	4.4	2.4	1.9
East North Central	24.1	23.8	5.3	4.8	2.4	2.3
West North Central	24.3 ^b	19.5	5.6 ^b	3.5	2.4 ^b	1.2
South	22.5	21.6	5.3	4.8	2.4	2.1
South Atlantic	24.0	22.4	5.1	4.8	2.2	1.9
East South Central	19.6	20.5	5.9	4.9	2.7	2.5
West South Central	21.7	21.0	5.4	4.7	2.4	2.1
West	28.5	30.6	6.0	6.4	2.8	3.0
Mountain	28.2	30.0	7.1	6.6	2.9	2.8
Pacific	28.7	30.8	5.6	6.3	2.7	3.2
COUNTY TYPE						
Large Metro	26.1	25.7	5.4	5.4	2.7	2.5
Small Metro	25.8	24.7	6.0	5.8	2.6	2.5
250K - 1 Mil. Pop.	26.4	24.9	6.2	5.7	2.8	2.4
<250K Pop.	24.1	23.9	5.5	6.0	1.9 ^b	3.1
Nonmetro	19.9	19.8	4.6	3.9	1.9	1.5
Urbanized	22.9	22.8	4.9	4.2	2.2	1.8
Less Urbanized	19.1	19.0	4.8	4.0	1.8	1.5
Completely Rural	16.9	16.7	2.7	2.8	1.6	1.0

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.106A Estimated Numbers (in Thousands) of Lifetime Users of Specific Hallucinogens, Inhalants, Needles, and Heroin, by Age Group: 1999 and 2000

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Hallucinogens	25,061	26,125	1,317	1,347	5,490	5,592	18,254	19,186
PCP	5,693	5,804	217	246	691	670	4,785	4,888
LSD	19,215	19,642	877	840	4,185	4,072	14,152	14,729
Peyote	5,013	4,947	104	113	483	498	4,427	4,337
Mescaline	7,947	7,728	93	103	696	700	7,158	6,924
Psilocybin (Mushrooms)	14,006	14,494	573	613	3,257	3,191	10,176	10,690
"Ecstasy" (MDMA)	5,097 ^b	6,482	424 ^b	615	2,172 ^b	2,800	2,501 ^a	3,067
Inhalants	17,138	16,702	2,118	2,079	4,014 ^a	3,701	11,006	10,922
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	7,488 ^a	6,669	417	397	932 ^a	782	6,139	5,490
Correction Fluid, Degreaser, or Cleaning Fluid	2,016 ^a	1,639	498	477	529 ^b	397	990	766
Gasoline or Lighter Fluid	3,039	3,014	781	763	765	722	1,493	1,530
Glue, Shoe Polish, or Toluene	3,195	3,132	878	900	671 ^a	560	1,646	1,672
Halothane, Ether, or Other Anesthetics	944	1,015	117	99	262 ^a	192	564	724
Lacquer Thinner or Other Paint Solvents	1,682	1,419	405	367	424	345	852	708
Lighter Gases (Butane, Propane)	828	779	251	243	304	244	273	292
Nitrous Oxide or "Whippets"	7,875	8,270	457	456	2,702	2,653	4,716	5,161
Spray Paints	1,524	1,530	578	510	368	333	578	686
Other Aerosol Sprays	1,586	1,578	428	400	409	384	750	794
Any Needle Use^{1,2}	3,065	2,643	76	62	305	270	2,684	2,311
Heroin Needle Use ^{1,2}	1,441	1,170	28	23	139	141	1,274	1,007
Cocaine Needle Use ^{1,2}	1,930	1,555	23	14	133	110	1,774	1,431
Stimulant Needle Use ^{1,2}	1,076	1,054	23	20	115	77	937	957
Heroin	3,054	2,779	102	84	498	403	2,454	2,292
Smoke Heroin ²	981	990	38	33	179	156	764	801
Sniff or Snort Heroin ²	1,701	1,817	31	30	316 ^a	236	1,354	1,551

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "non-core" module of the NHSDA questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NHSDA data are not edited for consistency across the "core" and "non-core" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.106B Percentages Reporting Lifetime Use of Specific Hallucinogens, Inhalants, Needles, and Heroin, by Age Group: 1999 and 2000

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Hallucinogens	11.3	11.7	5.7	5.8	19.3	19.3	10.8	11.2
PCP	2.6	2.6	0.9	1.1	2.4	2.3	2.8	2.9
LSD	8.7	8.8	3.8	3.6	14.7	14.0	8.4	8.6
Peyote	2.3	2.2	0.4	0.5	1.7	1.7	2.6	2.5
Mescaline	3.6	3.5	0.4	0.4	2.4	2.4	4.2	4.1
Psilocybin (Mushrooms)	6.3	6.5	2.5	2.6	11.4	11.0	6.0	6.3
"Ecstasy" (MDMA)	2.3 ^b	2.9	1.8 ^b	2.6	7.6 ^b	9.7	1.5 ^a	1.8
Inhalants	7.8	7.5	9.1	8.9	14.1 ^b	12.8	6.5	6.4
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	3.4 ^a	3.0	1.8	1.7	3.3 ^a	2.7	3.6	3.2
Correction Fluid, Degreaser, or Cleaning Fluid	0.9 ^a	0.7	2.1	2.0	1.9 ^b	1.4	0.6	0.4
Gasoline or Lighter Fluid	1.4	1.3	3.4	3.3	2.7	2.5	0.9	0.9
Glue, Shoe Polish, or Toluene	1.4	1.4	3.8	3.9	2.4 ^a	1.9	1.0	1.0
Halothane, Ether, or Other Anesthetics	0.4	0.5	0.5	0.4	0.9 ^a	0.7	0.3	0.4
Lacquer Thinner or Other Paint Solvents	0.8	0.6	1.7	1.6	1.5 ^a	1.2	0.5	0.4
Lighter Gases (Butane, Propane)	0.4	0.3	1.1	1.0	1.1	0.8	0.2	0.2
Nitrous Oxide or "Whippets"	3.6	3.7	2.0	2.0	9.5	9.2	2.8	3.0
Spray Paints	0.7	0.7	2.5	2.2	1.3	1.1	0.3	0.4
Other Aerosol Sprays	0.7	0.7	1.8	1.7	1.4	1.3	0.4	0.5
Any Needle Use^{1,2}	1.4	1.2	0.3	0.3	1.1	0.9	1.6	1.4
Heroin Needle Use ^{1,2}	0.7	0.5	0.1	0.1	0.5	0.5	0.8	0.6
Cocaine Needle Use ^{1,2}	0.9	0.7	0.1	0.1	0.5	0.4	1.0	0.8
Stimulant Needle Use ^{1,2}	0.5	0.5	0.1	0.1	0.4	0.3	0.6	0.6
Heroin	1.4	1.2	0.4	0.4	1.8 ^a	1.4	1.4	1.3
Smoke Heroin ²	0.4	0.4	0.2	0.1	0.6	0.5	0.5	0.5
Sniff or Snort Heroin ²	0.8	0.8	0.1	0.1	1.1 ^a	0.8	0.8	0.9

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "non-core" module of the NHSDA questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NHSDA data are not edited for consistency across the "core" and "non-core" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.107A Estimated Numbers (in Thousands) of Lifetime Users of Specific Pain Relievers, by Age Group: 1999 and 2000

Pain Relievers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Darvocet®, Darvon®, or Tylenol® with Codeine	14,389	13,549	1,170	1,214	2,881 ^a	2,613	10,337	9,722
Percocet®, Percodan®, or Tylox®	6,420	6,402	284	292	1,376	1,372	4,759	4,738
Vicodin®, Lortab®, or Lorcet®	6,067	6,708	380 ^b	520	1,864 ^a	2,114	3,823	4,075
Codeine	4,423	4,442	319	385	1,189	1,230	2,914	2,828
Demerol®	2,294	2,151	95	125	481	465	1,718	1,561
Dilaudid®	680	757	12	15	73	71	595	670
Fioricet®	311	286	12	14	54	42	244	229
Fiorinal®	390	309	18	12	35	37	338	261
Hydrocodone	1,622	1,788	104 ^a	147	524 ^b	729	994	912
Methadone	464 ^b	819	72	76	156	201	236 ^a	542
Morphine	1,234	1,536	147	167	341 ^a	425	746	944
Oxycontin®	221 ^b	399	20 ^a	41	71 ^b	178	130	179
Phenaphen® with Codeine	502	711	50	49	140	163	312	499
Propoxyphene	219	153	22	17	39	34	158	101
SK-65®	39	73	15	15	19	5	*	53
Stadol®	81	61	9	*	24	27	48	33
Talacen®	65	67	17 ^a	5	10	20	39	42
Talwin®	391	278	10	10	27	22	354	246
Talwin NX®	76	76	10	4	9	3	57	69
Tramadol	42	83	9	12	27	29	*	41
Ultram®	395	456	40	53	118	170	236	234

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.107B Percentages Reporting Lifetime Use of Specific Pain Relievers, by Age Group: 1999 and 2000

Pain Relievers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Darvocet®, Darvon®, or Tylenol® with Codeine	6.5	6.1	5.0	5.2	10.1 ^b	9.0	6.1	5.7
Percocet®, Percodan®, or Tylox®	2.9	2.9	1.2	1.2	4.8	4.7	2.8	2.8
Vicodin®, Lortab®, or Lorcet®	2.7	3.0	1.6 ^b	2.2	6.5 ^a	7.3	2.3	2.4
Codeine	2.0	2.0	1.4	1.6	4.2	4.2	1.7	1.7
Demerol®	1.0	1.0	0.4	0.5	1.7	1.6	1.0	0.9
Dilaudid®	0.3	0.3	0.1	0.1	0.3	0.2	0.4	0.4
Fioricet®	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1
Fiorinal®	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.2
Hydrocodone	0.7	0.8	0.4 ^a	0.6	1.8 ^b	2.5	0.6	0.5
Methadone	0.2 ^b	0.4	0.3	0.3	0.5	0.7	0.1 ^a	0.3
Morphine	0.6	0.7	0.6	0.7	1.2	1.5	0.4	0.6
Oxycontin®	0.1 ^b	0.2	0.1 ^a	0.2	0.2 ^b	0.6	0.1	0.1
Phenaphen® with Codeine	0.2	0.3	0.2	0.2	0.5	0.6	0.2	0.3
Propoxyphene	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
SK-65®	0.0	0.0	0.1	0.1	0.1	0.0	*	0.0
Stadol®	0.0	0.0	0.0	*	0.1	0.1	0.0	0.0
Talacen®	0.0	0.0	0.1 ^a	0.0	0.0	0.1	0.0	0.0
Talwin®	0.2	0.1	0.0	0.0	0.1	0.1	0.2	0.1
Talwin NX®	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Tramadol	0.0	0.0	0.0	0.1	0.1	0.1	*	0.0
Ultram®	0.2	0.2	0.2	0.2	0.4	0.6	0.1	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.108A Estimated Numbers (in Thousands) of Lifetime Users of Specific Tranquilizers, by Age Group: 1999 and 2000

Tranquilizers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Klonopin® or Clonazepam	1,254	1,223	76	96	353 ^a	445	824	682
Xanax®, Alprazolam, Ativan®, or Lorazepam	5,042	4,666	250	266	1,206	1,164	3,586	3,236
Valium® or Diazepam	10,982	10,235	325	348	1,638	1,552	9,019	8,335
Atarax®	334	335	22	20	36	53	275	263
BuSpar®	333	415	20 ^b	50	90 ^a	138	223	228
Equanil®	57	119	14	17	18	16	26	86
Flexeril®	816	1,035	27	30	169	207	619	797
Librium®	891	841	24	21	55	77	812	743
Limbitrol®	42	80	13	9	9	22	21	49
Meprobamate	106	136	14	10	17	21	75	104
Miltown®	79	145	9	11	19	8	52	127
Rohypnol®	275	292	29	40	146	147	100	106
Serax®	132	97	13	15	26	36	94	46
Soma®	1,042	1,057	61	68	330	382	651	606
Tranxene®	251	143	18	17	39	30	195	96
Vistaril®	251	198	21	16	28 ^a	50	202	131

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.108B Percentages Reporting Lifetime Use of Specific Tranquilizers, by Age Group: 1999 and 2000

Tranquilizers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Klonopin® or Clonazepam	0.6	0.5	0.3	0.4	1.2	1.5	0.5	0.4
Xanax®, Alprazolam, Ativan®, or Lorazepam	2.3	2.1	1.1	1.1	4.2	4.0	2.1	1.9
Valium® or Diazepam	5.0	4.6	1.4	1.5	5.8	5.4	5.3	4.9
Atarax®	0.2	0.2	0.1	0.1	0.1	0.2	0.2	0.2
BuSpar®	0.2	0.2	0.1 ^b	0.2	0.3 ^a	0.5	0.1	0.1
Equanil®	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.1
Flexeril®	0.4	0.5	0.1	0.1	0.6	0.7	0.4	0.5
Librium®	0.4	0.4	0.1	0.1	0.2	0.3	0.5	0.4
Limbitrol®	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0
Meprobamate	0.0	0.1	0.1	0.0	0.1	0.1	0.0	0.1
Miltown®	0.0	0.1	0.0	0.0	0.1	0.0	0.0	0.1
Rohypnol®	0.1	0.1	0.1	0.2	0.5	0.5	0.1	0.1
Serax®	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.0
Soma®	0.5	0.5	0.3	0.3	1.2	1.3	0.4	0.4
Tranxene®	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Vistaril®	0.1	0.1	0.1	0.1	0.1 ^a	0.2	0.1	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.109A Estimated Numbers (in Thousands) of Lifetime Users of Specific Stimulants, by Age Group: 1999 and 2000

Stimulants	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methamphetamine, Desoxyn®, or Methedrine	9,442	8,843	317	305	1,483 ^b	1,185	7,643	7,353
Prescription Diet Pills ¹	7,293	6,605	240	282	633	548	6,420	5,776
Ritalin® or Methylphenidate	2,831	2,761	518	506	1,021	1,055	1,292	1,201
Cylert®	293	246	22	24	49	52	221	169
Dexedrine®	1,855	1,982	44	63	156	191	1,654	1,729
Dextroamphetamine	459	634	17	28	71	77	372	528
Didrex®	200	182	6	16	18	20	176	145
Eskatrol®	127	152	8	11	12	22	106	118
Ionamin®	403	333	9	14	24	41	370	279
Mazanor®	79	93	8 ^a	23	17	19	54	51
Obedrin-LA®	45	91	3	5	5	13	37	73
Plegine®	29	84	6	12	2	4	20	67
Preludin®	521	671	14	10	53	47	455	614
Sanorex®	147	106	20	18	28	26	99	63
Tenuate®	97	81	8	9	16	18	74	54

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ For example, Amphetamines, Benzedrine®, Biphedamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.109B Percentages Reporting Lifetime Use of Specific Stimulants, by Age Group: 1999 and 2000

Stimulants	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methamphetamine, Desoxyn®, or Methedrine	4.3	4.0	1.4	1.3	5.2 ^b	4.1	4.5	4.3
Prescription Diet Pills ¹	3.3	3.0	1.0	1.2	2.2	1.9	3.8	3.4
Ritalin® or Methylphenidate	1.3	1.2	2.2	2.2	3.6	3.6	0.8	0.7
Cylert®	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.1
Dexedrine®	0.8	0.9	0.2	0.3	0.5	0.7	1.0	1.0
Dextroamphetamine	0.2	0.3	0.1	0.1	0.2	0.3	0.2	0.3
Didrex®	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Eskatrol®	0.1	0.1	0.0	0.0	0.0	0.1	0.1	0.1
Ionamin®	0.2	0.1	0.0	0.1	0.1	0.1	0.2	0.2
Mazanor®	0.0	0.0	0.0 ^a	0.1	0.1	0.1	0.0	0.0
Obedrin-LA®	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Plegine®	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
Preludin®	0.2	0.3	0.1	0.0	0.2	0.2	0.3	0.4
Sanorex®	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.0
Tenuate®	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ For example, Amphetamines, Benzedrine®, Biphedamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.110A Estimated Numbers (in Thousands) of Lifetime Users of Specific Sedatives, by Age Group: 1999 and 2000

Sedatives	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methaqualone, Sopor®, or Quaalude®	5,964	5,416	70	45	296	259	5,599	5,111
Barbiturates ¹	2,981	2,847	41	50	158	129	2,782	2,668
Restoril® or Temazepam	639	622	38	34	93	70	508	518
Amytal®	203	221	19	19	12	20	171	182
Butisol®	91	134	10	10	17	17	64	107
Chloral Hydrate	170	163	7	13	17	18	146	132
Dalmane®	273	477	17	16	19	17	238	443
Halcion®	502	441	19	17	58	35	425	389
Phenobarbital	891	991	23	23	85	71	782	897
Placidyl®	696	640	8	9	37	22	651	609
Tuinal®	842	1,030	8	8	13	20	821	1,002

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.110B Percentages Reporting Lifetime Use of Specific Sedatives, by Age Group: 1999 and 2000

Sedatives	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methaqualone, Sopor®, or Quaalude®	2.7	2.4	0.3	0.2	1.0	0.9	3.3	3.0
Barbiturates ¹	1.3	1.3	0.2	0.2	0.6	0.4	1.6	1.6
Restoril® or Temazepam	0.3	0.3	0.2	0.1	0.3	0.2	0.3	0.3
Amytal®	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1
Butisol®	0.0	0.1	0.0	0.0	0.1	0.1	0.0	0.1
Chloral Hydrate	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Dalmane®	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.3
Halcion®	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.2
Phenobarbital	0.4	0.4	0.1	0.1	0.3	0.2	0.5	0.5
Placidyl®	0.3	0.3	0.0	0.0	0.1	0.1	0.4	0.4
Tuinal®	0.4	0.5	0.0	0.0	0.0	0.1	0.5	0.6

*Low precision; no estimate reported.

^aDifference between estimate and 2000 estimate is statistically significant at the .05 level.

^bDifference between estimate and 2000 estimate is statistically significant at the .01 level.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.