

Table 1.113A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	46,558	47,882	14,680	14,986	6,210	6,336
GEOGRAPHIC DIVISION						
Northeast	7,917	8,156	2,343	2,342	921	944
New England	2,506	2,539	834	875	294	300
Middle Atlantic	5,411	5,617	1,509	1,467	627	643
Midwest	9,928	10,405	3,079	3,194	1,178	1,297
East North Central	6,865	7,195	2,233	2,169	812	894
West North Central	3,063	3,209	846	1,025	366	402
South	16,186	16,418	5,458	5,346	2,450	2,255
South Atlantic	8,345	8,478	2,803	2,697	1,175	1,076
East South Central	2,667	2,794	981	856	546 ^a	362
West South Central	5,174	5,145	1,674	1,793	729	817
West	12,527	12,903	3,799	4,104	1,661	1,840
Mountain	3,652	3,911	1,105	1,266	526	518
Pacific	8,875	8,992	2,695	2,838	1,135	1,322
COUNTY TYPE¹						
Large Metro	25,296	25,933	7,552	7,749	3,215	3,283
Small Metro	14,267	14,595	4,829	4,915	1,995	2,030
250K – 1 Mil. Pop.	9,426	9,531	3,072	3,129	1,307	1,224
< 250K Pop.	4,841	5,064	1,757	1,786	688	806
Nonmetro	6,995	7,353	2,299	2,321	1,000	1,023
Urbanized	3,101	3,350	983	1,142	389	522
Less Urbanized	3,242	3,335	1,048	1,047	428	468
Completely Rural	653	669	267	133	182 ^a	33

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.113B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	19.8	20.1	6.2	6.3	2.6	2.7
GEOGRAPHIC DIVISION						
Northeast	17.6	18.0	5.2	5.2	2.0	2.1
New England	21.2	21.4	7.1	7.4	2.5	2.5
Middle Atlantic	16.3	16.8	4.5	4.4	1.9	1.9
Midwest	18.6	19.4	5.8	5.9	2.2	2.4
East North Central	18.3	19.1	6.0	5.8	2.2	2.4
West North Central	19.1	19.9	5.3	6.4	2.3	2.5
South	19.3	19.4	6.5	6.3	2.9	2.7
South Atlantic	19.1	19.1	6.4	6.1	2.7	2.4
East South Central	18.8	19.7	6.9	6.0	3.9 ^a	2.5
West South Central	20.0	19.7	6.5	6.9	2.8	3.1
West	23.7	24.0	7.2	7.6	3.1	3.4
Mountain	23.8	25.1	7.2	8.1	3.4	3.3
Pacific	23.6	23.5	7.2	7.4	3.0	3.5
COUNTY TYPE¹						
Large Metro	20.1	20.6	6.0	6.1	2.6	2.6
Small Metro	20.6	20.4	7.0	6.9	2.9	2.8
250K – 1 Mil. Pop.	20.7	20.3	6.8	6.7	2.9	2.6
< 250K Pop.	20.3	20.7	7.4	7.3	2.9	3.3
Nonmetro	17.4	18.3	5.7	5.8	2.5	2.5
Urbanized	19.3	18.7	6.1	6.4	2.4	2.9
Less Urbanized	16.5	18.3	5.3	5.7	2.2	2.6
Completely Rural	14.7	16.3	6.0	3.2	4.1 ^a	0.8

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.114A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	3,389	3,351	2,283	2,289	979	997
GEOGRAPHIC DIVISION						
Northeast	523	520	331	352	152	160
New England	177	147	105	105	53	40
Middle Atlantic	347	373	226	247	99	120
Midwest	811 ^a	714	548	485	204	209
East North Central	569	504	393	345	148	159
West North Central	242	211	155	140	57	50
South	1,212	1,254	843	866	393	379
South Atlantic	599	590	428	406	198	161
East South Central	202	234	143	175	67	74
West South Central	411	431	272	284	128	144
West	842	862	561	587	230	249
Mountain	256	284	171	190	67	82
Pacific	587	579	390	396	163	167
COUNTY TYPE¹						
Large Metro	1,735	1,612	1,134	1,094	471	462
Small Metro	1,016	1,104	722	736	321	328
250K – 1 Mil. Pop.	693	738	495	479	237	214
< 250K Pop.	323	366	228	257	83	113
Nonmetro	637	635	426	459	188	207
Urbanized	239	290	157 ^a	220	76	101
Less Urbanized	335	294	222	202	92	87
Completely Rural	63	51	48	37	19	18

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.114B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	13.7	13.4	9.2	9.2	4.0	4.0
GEOGRAPHIC DIVISION						
Northeast	11.7	11.6	7.4	7.8	3.4	3.6
New England	15.3	12.5	9.1	9.0	4.5	3.4
Middle Atlantic	10.5	11.3	6.8	7.4	3.0	3.6
Midwest	14.3 ^b	12.5	9.6 ^a	8.5	3.6	3.7
East North Central	14.3 ^a	12.6	9.8	8.6	3.7	4.0
West North Central	14.2	12.4	9.1	8.2	3.3	3.0
South	13.8	14.0	9.6	9.7	4.5	4.2
South Atlantic	13.6	13.1	9.7	9.0	4.5	3.6
East South Central	14.2	16.1	10.1	12.1	4.7	5.1
West South Central	13.9	14.5	9.2	9.6	4.3	4.8
West	14.4	14.7	9.6	10.0	3.9	4.2
Mountain	15.0	16.4	10.1	11.0	3.9	4.8
Pacific	14.2	13.9	9.4	9.5	4.0	4.0
COUNTY TYPE¹						
Large Metro	13.1	12.1	8.6	8.2	3.6	3.5
Small Metro	14.0	14.9	10.0	9.9	4.4	4.4
250K – 1 Mil. Pop.	14.1	14.8	10.1	9.6	4.8	4.3
< 250K Pop.	13.8	15.1	9.7	10.6	3.6	4.7
Nonmetro	14.8	14.8	9.9	10.7	4.4	4.8
Urbanized	13.9	15.6	9.1 ^a	11.8	4.4	5.4
Less Urbanized	15.9	14.8	10.5	10.2	4.4	4.4
Completely Rural	13.2	11.6	10.0	8.5	4.1	4.2

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.115A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	8,587 ^b	9,211	4,411	4,598	1,683 ^a	1,913
GEOGRAPHIC DIVISION						
Northeast	1,370 ^a	1,555	725	787	281	331
New England	466	501	271	246	99	106
Middle Atlantic	904 ^a	1,055	454	541	182	225
Midwest	1,990	2,091	995	1,049	360	412
East North Central	1,423	1,445	740	719	262	279
West North Central	567	646	255 ^a	330	98	133
South	3,095	3,283	1,727	1,664	694	724
South Atlantic	1,461	1,562	804	791	288	362
East South Central	586	567	340	279	150	122
West South Central	1,048	1,154	583	595	256	240
West	2,132	2,282	964	1,097	347 ^a	445
Mountain	717	743	354	366	126	140
Pacific	1,415	1,538	610	731	221 ^a	305
COUNTY TYPE¹						
Large Metro	4,204 ^b	4,666	2,102	2,281	819 ^a	965
Small Metro	3,009	3,054	1,582	1,537	602	614
250K – 1 Mil. Pop.	1,853	1,978	961	961	371	373
< 250K Pop.	1,156	1,076	621	575	230	241
Nonmetro	1,373	1,491	727	780	262	334
Urbanized	624	735	350	388	121 ^a	174
Less Urbanized	629	668	313	358	114	151
Completely Rural	119	88	65 ^a	34	28 ^b	9

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.115B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	27.7 ^a	29.0	14.2	14.5	5.4 ^a	6.0
GEOGRAPHIC DIVISION						
Northeast	24.9 ^a	27.6	13.2	14.0	5.1	5.9
New England	32.5	34.1	18.9	16.7	6.9	7.2
Middle Atlantic	22.2 ^a	25.3	11.2	13.0	4.5	5.4
Midwest	27.6	28.4	13.8	14.3	5.0	5.6
East North Central	28.6	28.4	14.9	14.2	5.3	5.5
West North Central	25.5	28.4	11.5	14.5	4.4	5.9
South	28.1	29.0	15.7	14.7	6.3	6.4
South Atlantic	27.1	28.0	14.9	14.2	5.3	6.5
East South Central	30.7	29.3	17.8	14.4	7.8	6.3
West South Central	28.2	30.3	15.7	15.6	6.9	6.3
West	29.2	30.8	13.2	14.8	4.8 ^a	6.0
Mountain	32.8	33.2	16.2	16.4	5.8	6.3
Pacific	27.7	29.7	11.9 ^a	14.1	4.3 ^a	5.9
COUNTY TYPE¹						
Large Metro	26.5	28.1	13.2	13.7	5.2	5.8
Small Metro	29.6	29.9	15.6	15.1	5.9	6.0
250K – 1 Mil. Pop.	28.8	29.7	14.9	14.5	5.8	5.6
< 250K Pop.	31.0	30.3	16.7	16.2	6.2	6.8
Nonmetro	27.6 ^a	30.3	14.6	15.9	5.3 ^a	6.8
Urbanized	27.8	31.1	15.6	16.4	5.4 ^a	7.4
Less Urbanized	27.8	30.5	13.8	16.3	5.0	6.9
Completely Rural	25.6	24.2	14.0	9.4	6.0 ^b	2.4

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.116A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	34,582	35,320	7,986	8,099	3,549	3,426
GEOGRAPHIC DIVISION						
Northeast	6,024	6,081	1,287	1,203	488	453
New England	1,864	1,892	458	524	142	154
Middle Atlantic	4,160	4,189	828	679	346	298
Midwest	7,127	7,600	1,537	1,660	614	676
East North Central	4,874	5,247	1,101	1,104	402	457
West North Central	2,254	2,352	435	556	212	219
South	11,879	11,881	2,888	2,816	1,363	1,152
South Atlantic	6,286	6,327	1,571	1,500	689	553
East South Central	1,878	1,993	497	402	329	166
West South Central	3,715	3,560	820	914	345	433
West	9,552	9,759	2,275	2,420	1,083	1,146
Mountain	2,679	2,885	579	709	333	295
Pacific	6,873	6,874	1,696	1,711	751	851
COUNTY TYPE¹						
Large Metro	19,356	19,655	4,315	4,374	1,926	1,855
Small Metro	10,241	10,437	2,525	2,643	1,073	1,089
250K – 1 Mil. Pop.	6,879	6,815	1,617	1,689	698	637
< 250K Pop.	3,362	3,622	908	954	375	452
Nonmetro	4,985	5,228	1,145	1,082	550	482
Urbanized	2,237	2,325	477	534	193	246
Less Urbanized	2,277	2,373	514	487	222	230
Completely Rural	471	530	155	62	135 ^a	6

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.116B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	19.3	19.5	4.5	4.5	2.0	1.9
GEOGRAPHIC DIVISION						
Northeast	17.2	17.3	3.7	3.4	1.4	1.3
New England	20.2	20.5	5.0	5.7	1.5	1.7
Middle Atlantic	16.1	16.1	3.2	2.6	1.3	1.1
Midwest	17.6	18.7	3.8	4.1	1.5	1.7
East North Central	17.1	18.4	3.9	3.9	1.4	1.6
West North Central	18.7	19.4	3.6	4.6	1.8	1.8
South	18.6	18.4	4.5	4.4	2.1	1.8
South Atlantic	18.5	18.4	4.6	4.4	2.0	1.6
East South Central	17.4	18.4	4.6	3.7	3.0 ^a	1.5
West South Central	19.4	18.4	4.3	4.7	1.8	2.2
West	24.0	24.1	5.7	6.0	2.7	2.8
Mountain	23.5	24.8	5.1	6.1	2.9	2.5
Pacific	24.3	23.8	6.0	5.9	2.6	2.9
COUNTY TYPE¹						
Large Metro	20.0	20.4	4.5	4.5	2.0	1.9
Small Metro	19.7	19.4	4.9	4.9	2.1	2.0
250K – 1 Mil. Pop.	20.1	19.3	4.7	4.8	2.0	1.8
< 250K Pop.	19.0	19.6	5.1	5.2	2.1	2.4
Nonmetro	16.2	16.9	3.7	3.5	1.8	1.6
Urbanized	18.6	17.0	4.0	3.9	1.6	1.8
Less Urbanized	14.9	16.9	3.4	3.5	1.5	1.6
Completely Rural	13.5	16.1	4.4	1.9	3.9 ^a	0.2

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.117A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	43,169	44,530	12,397	12,697	5,231	5,339
GEOGRAPHIC DIVISION						
Northeast	7,394	7,636	2,012	1,990	770	784
New England	2,329	2,392	729	770	241	260
Middle Atlantic	5,064	5,243	1,283	1,220	528	523
Midwest	9,117	9,691	2,531	2,709	974	1,088
East North Central	6,296	6,692	1,841	1,824	664	736
West North Central	2,820	2,999	691	885	309	352
South	14,974	15,163	4,615	4,480	2,057	1,877
South Atlantic	7,747	7,889	2,375	2,291	977	916
East South Central	2,465	2,560	838	680	479 ^a	288
West South Central	4,763	4,714	1,402	1,509	601	673
West	11,684	12,041	3,238	3,517	1,431	1,591
Mountain	3,396	3,628	933	1,075	459	436
Pacific	8,288	8,413	2,305	2,442	972	1,155
COUNTY TYPE¹						
Large Metro	23,561	24,321	6,418	6,655	2,744	2,821
Small Metro	13,251	13,491	4,107	4,180	1,675	1,702
250K – 1 Mil. Pop.	8,733	8,793	2,578	2,650	1,070	1,009
< 250K Pop.	4,518	4,698	1,529	1,529	605	693
Nonmetro	6,358	6,718	1,873	1,862	812	816
Urbanized	2,861	3,060	826	922	313	421
Less Urbanized	2,906	3,041	827	844	336	381
Completely Rural	590	618	220	96	163 ^a	15

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.117B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	20.5	20.9	5.9	6.0	2.5	2.5
GEOGRAPHIC DIVISION						
Northeast	18.2	18.7	5.0	4.9	1.9	1.9
New England	21.9	22.3	6.8	7.2	2.3	2.4
Middle Atlantic	16.9	17.4	4.3	4.1	1.8	1.7
Midwest	19.1	20.2	5.3	5.6	2.0	2.3
East North Central	18.8	19.9	5.5	5.4	2.0	2.2
West North Central	19.7	20.8	4.8	6.1	2.2	2.4
South	20.0	20.0	6.2	5.9	2.7	2.5
South Atlantic	19.7	19.8	6.0	5.7	2.5	2.3
East South Central	19.4	20.1	6.6	5.3	3.8 ^a	2.3
West South Central	20.8	20.4	6.1	6.5	2.6	2.9
West	24.8	25.1	6.9	7.3	3.0	3.3
Mountain	25.0	26.1	6.9	7.7	3.4	3.1
Pacific	24.8	24.7	6.9	7.2	2.9	3.4
COUNTY TYPE¹						
Large Metro	20.9	21.6	5.7	5.9	2.4	2.5
Small Metro	21.4	21.1	6.6	6.5	2.7	2.7
250K – 1 Mil. Pop.	21.5	21.0	6.4	6.3	2.6	2.4
< 250K Pop.	21.1	21.3	7.1	6.9	2.8	3.1
Nonmetro	17.8	18.7	5.2	5.2	2.3	2.3
Urbanized	20.0	19.1	5.8	5.8	2.2	2.6
Less Urbanized	16.6	18.7	4.7	5.2	1.9	2.3
Completely Rural	14.9	16.9	5.6	2.6	4.1 ^a	0.4

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.