

**Table 1.128A Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Numbers in Thousands, 2002 and 2003**

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>HALLUCINOGENS</b>	34,314	34,363	1,419 <sup>a</sup>	1,250	7,517	7,377	25,378	25,736
PCP (Angel Dust, Phencyclidine)	7,418	7,107	233	204	834	948	6,352	5,955
LSD (Acid)	24,516	24,424	670 <sup>b</sup>	397	4,928 <sup>b</sup>	4,450	18,917	19,577
Peyote	5,738	5,783	99	101	533	555	5,105	5,127
Mescaline	9,116	9,455	106 <sup>b</sup>	61	776	671	8,233	8,724
Psilocybin (Mushrooms)	18,322	19,507	564	566	4,167	4,238	13,592	14,703
"Ecstasy" (MDMA)	10,150	10,904	820 <sup>b</sup>	603	4,674	4,683	4,656 <sup>b</sup>	5,619
<b>INHALANTS</b>	22,870	22,995	2,605	2,670	4,867	4,729	15,398	15,596
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	8,915	8,957	431	450	899	851	7,585	7,656
Correction Fluid, Degreaser, or Cleaning Fluid	2,248	2,262	550	578	520	463	1,178	1,221
Gasoline or Lighter Fluid	4,023	3,856	862	890	910	853	2,250	2,113
Glue, Shoe Polish, or Toluene	4,551	4,269	1,105	1,112	805	788	2,641	2,369
Halothane, Ether, or Other Anesthetics	1,114	1,366	96	116	256	236	762	1,014
Lacquer Thinner or Other Paint Solvents	2,006	1,968	427	417	487	450	1,093	1,101
Lighter Gases (Butane, Propane)	1,041	951	270	294	314	278	458	380
Nitrous Oxide or "Whippets"	11,698	11,619	526	457	3,308	3,231	7,863	7,930
Spray Paints	2,012	2,086	631 <sup>a</sup>	734	479	438	903	913
Other Aerosol Sprays	2,011	2,046	427 <sup>a</sup>	516	613	616	971	914
<b>ANY NEEDLE USE<sup>1,2</sup></b>	3,982	3,683	81	77	375	409	3,526	3,197
Heroin Needle Use <sup>1,2</sup>	1,674	1,661	27	30	151	162	1,497	1,469
Cocaine Needle Use <sup>1,2</sup>	2,499	2,166	18	12	135	166	2,345	1,988
Stimulant Needle Use <sup>1,2</sup>	1,805	1,655	37	32	169	186	1,600	1,437
Methamphetamine Needle Use <sup>1,2</sup>	1,314	1,258	20	22	129	135	1,165	1,101
<b>HEROIN</b>	3,668	3,744	94	80	487	516	3,087	3,148
Smoke Heroin <sup>2</sup>	1,379	1,268	40	44	194	199	1,145	1,025
Sniff or Snort Heroin <sup>2</sup>	2,554	2,580	60	43	336	350	2,159	2,187

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

<sup>2</sup> These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.128B Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Percentages, 2002 and 2003**

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>HALLUCINOGENS</b>	14.6	14.5	5.7 <sup>b</sup>	5.0	24.2	23.3	14.1	14.2
PCP (Angel Dust, Phencyclidine)	3.2	3.0	0.9	0.8	2.7	3.0	3.5	3.3
LSD (Acid)	10.4	10.3	2.7 <sup>b</sup>	1.6	15.9 <sup>b</sup>	14.0	10.5	10.8
Peyote	2.4	2.4	0.4	0.4	1.7	1.7	2.8	2.8
Mescaline	3.9	4.0	0.4 <sup>b</sup>	0.2	2.5 <sup>a</sup>	2.1	4.6	4.8
Psilocybin (Mushrooms)	7.8	8.2	2.3	2.3	13.4	13.4	7.6	8.1
"Ecstasy" (MDMA)	4.3	4.6	3.3 <sup>b</sup>	2.4	15.1	14.8	2.6 <sup>b</sup>	3.1
<b>INHALANTS</b>	9.7	9.7	10.5	10.7	15.7	14.9	8.6	8.6
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	3.8	3.8	1.7	1.8	2.9	2.7	4.2	4.2
Correction Fluid, Degreaser, or Cleaning Fluid	1.0	1.0	2.2	2.3	1.7	1.5	0.7	0.7
Gasoline or Lighter Fluid	1.7	1.6	3.5	3.6	2.9	2.7	1.3	1.2
Glue, Shoe Polish, or Toluene	1.9	1.8	4.5	4.4	2.6	2.5	1.5	1.3
Halothane, Ether, or Other Anesthetics	0.5	0.6	0.4	0.5	0.8	0.7	0.4	0.6
Lacquer Thinner or Other Paint Solvents	0.9	0.8	1.7	1.7	1.6	1.4	0.6	0.6
Lighter Gases (Butane, Propane)	0.4	0.4	1.1	1.2	1.0	0.9	0.3	0.2
Nitrous Oxide or "Whippets"	5.0	4.9	2.1	1.8	10.7	10.2	4.4	4.4
Spray Paints	0.9	0.9	2.5	2.9	1.5	1.4	0.5	0.5
Other Aerosol Sprays	0.9	0.9	1.7 <sup>a</sup>	2.1	2.0	1.9	0.5	0.5
<b>ANY NEEDLE USE<sup>1,2</sup></b>	1.7	1.5	0.3	0.3	1.2	1.3	2.0	1.8
Heroin Needle Use <sup>1,2</sup>	0.7	0.7	0.1	0.1	0.5	0.5	0.8	0.8
Cocaine Needle Use <sup>1,2</sup>	1.1	0.9	0.1	0.0	0.4	0.5	1.3	1.1
Stimulant Needle Use <sup>1,2</sup>	0.8	0.7	0.1	0.1	0.5	0.6	0.9	0.8
Methamphetamine Needle Use <sup>1,2</sup>	0.6	0.5	0.1	0.1	0.4	0.4	0.6	0.6
<b>HEROIN</b>	1.6	1.6	0.4	0.3	1.6	1.6	1.7	1.7
Smoke Heroin <sup>2</sup>	0.6	0.5	0.2	0.2	0.6	0.6	0.6	0.6
Sniff or Snort Heroin <sup>2</sup>	1.1	1.1	0.2	0.2	1.1	1.1	1.2	1.2

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

<sup>2</sup> These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.129A Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Numbers in Thousands, 2002 and 2003**

Pain Reliever	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Darvocet®, Darvon®, or Tylenol® with Codeine	18,912	19,672	1,508	1,458	3,800	4,020	13,603	14,193
Percocet®, Percodan®, or Tylox®	9,659 <sup>b</sup>	10,811	407	473	2,285	2,471	6,967 <sup>a</sup>	7,867
Vicodin®, Lortab®, or Lorcet®	13,093 <sup>b</sup>	15,717	1,116	1,129	4,126 <sup>b</sup>	4,773	7,852 <sup>b</sup>	9,815
Codeine	6,897	6,985	513	526	1,878	2,074	4,506	4,385
Demerol®	2,917	2,986	116	109	657	686	2,144	2,191
Dilaudid®	1,116	914	7	13	101	80	1,008	821
Fioricet®	445	466	13	17	64	49	368	401
Fiorinal®	579	525	24	18	40	43	515	465
Hydrocodone	4,529 <sup>b</sup>	5,716	333 <sup>a</sup>	410	1,540 <sup>b</sup>	2,088	2,656 <sup>a</sup>	3,218
Methadone	928 <sup>a</sup>	1,159	116	109	306 <sup>a</sup>	396	506	655
Morphine	2,073	2,060	256	215	648	737	1,170	1,107
Oxycontin®	1,924 <sup>b</sup>	2,832	224	254	820 <sup>b</sup>	1,145	881 <sup>b</sup>	1,433
Phenaphen® with Codeine	873	857	69	67	194	236	610	554
Propoxyphene	282	290	31	33	45	68	206	189
SK-65®	42	48	13	16	10	14	19	18
Stadol®	133	157	6	5	23	16	103	136
Talacen®	55	43	14	7	13	13	28	23
Talwin®	643 <sup>a</sup>	425	10	6	20	22	613 <sup>a</sup>	397
Talwin NX®	105	53	10	8	9	12	86	33
Tramadol	52 <sup>b</sup>	186	11	14	20 <sup>b</sup>	68	21 <sup>a</sup>	103
Ultram®	962	1,105	98	66	300	330	564	709

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.129B Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Percentages, 2002 and 2003**

Pain Reliever	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Darvocet®, Darvon®, or Tylenol® with Codeine	8.0	8.3	6.1	5.8	12.2	12.7	7.6	7.8
Percocet®, Percodan®, or Tylox®	4.1 <sup>a</sup>	4.5	1.6	1.9	7.4	7.8	3.9 <sup>a</sup>	4.3
Vicodin®, Lortab®, or Lorcet®	5.6 <sup>b</sup>	6.6	4.5	4.5	13.3 <sup>b</sup>	15.0	4.4 <sup>b</sup>	5.4
Codeine	2.9	2.9	2.1	2.1	6.1	6.5	2.5	2.4
Demerol®	1.2	1.3	0.5	0.4	2.1	2.2	1.2	1.2
Dilaudid®	0.5	0.4	0.0	0.1	0.3	0.3	0.6	0.5
Fioricet®	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.2
Fiorinal®	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.3
Hydrocodone	1.9 <sup>b</sup>	2.4	1.3	1.6	5.0 <sup>b</sup>	6.6	1.5 <sup>a</sup>	1.8
Methadone	0.4	0.5	0.5	0.4	1.0 <sup>a</sup>	1.2	0.3	0.4
Morphine	0.9	0.9	1.0	0.9	2.1	2.3	0.7	0.6
Oxycontin®	0.8 <sup>b</sup>	1.2	0.9	1.0	2.6 <sup>b</sup>	3.6	0.5 <sup>b</sup>	0.8
Phenaphen® with Codeine	0.4	0.4	0.3	0.3	0.6	0.7	0.3	0.3
Propoxyphene	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
SK-65®	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0
Stadol®	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1
Talacen®	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Talwin®	0.3 <sup>a</sup>	0.2	0.0	0.0	0.1	0.1	0.3 <sup>a</sup>	0.2
Talwin NX®	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Tramadol	0.0 <sup>b</sup>	0.1	0.0	0.1	0.1 <sup>b</sup>	0.2	0.0 <sup>a</sup>	0.1
Ultram®	0.4	0.5	0.4	0.3	1.0	1.0	0.3	0.4

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.130A Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Numbers in Thousands, 2002 and 2003**

Tranquilizer	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Klonopin® or Clonazepam	2,374 <sup>a</sup>	2,851	180	160	724 <sup>b</sup>	964	1,470	1,727
Xanax®, Alprazolam, Ativan®, or Lorazepam	8,208 <sup>b</sup>	9,500	429	437	2,091 <sup>b</sup>	2,379	5,688 <sup>b</sup>	6,684
Valium® or Diazepam	14,386	14,671	427	424	2,312	2,473	11,648	11,773
Atarax®	371	292	26	17	45	42	301	233
BuSpar®	693	587	51	53	220	205	421	329
Equanil®	75	128	14	13	12	14	49	101
Flexeril®	1,781	1,994	47	49	295 <sup>a</sup>	377	1,439	1,569
Librium®	1,454	1,121	17	20	79	84	1,358	1,017
Limbitrol®	115 <sup>b</sup>	41	9	7	32 <sup>a</sup>	11	73	23
Meproamate	161	170	9	12	21	11	131	148
Miltown®	169	98	12	8	6	8	151	82
Rohypnol®	375	408	40	46	160	189	175	173
Serax®	215	170	17	9	29	17	169	144
Soma®	2,276	2,572	142	152	729	793	1,405	1,626
Tranxene®	214	189	10	13	16	23	188	153
Vistaril®	356	258	21	11	64 <sup>a</sup>	35	271	212

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.130B Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Percentages, 2002 and 2003**

Tranquilizer	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Klonopin® or Clonazepam	1.0 <sup>a</sup>	1.2	0.7	0.6	2.3 <sup>b</sup>	3.0	0.8	1.0
Xanax®, Alprazolam, Ativan®, or Lorazepam	3.5 <sup>b</sup>	4.0	1.7	1.7	6.7 <sup>a</sup>	7.5	3.2 <sup>b</sup>	3.7
Valium® or Diazepam	6.1	6.2	1.7	1.7	7.5	7.8	6.5	6.5
Atarax®	0.2	0.1	0.1	0.1	0.1	0.1	0.2	0.1
BuSpar®	0.3	0.2	0.2	0.2	0.7	0.6	0.2	0.2
Equanil®	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.1
Flexeril®	0.8	0.8	0.2	0.2	0.9	1.2	0.8	0.9
Librium®	0.6	0.5	0.1	0.1	0.3	0.3	0.8	0.6
Limbitrol®	0.0 <sup>b</sup>	0.0	0.0	0.0	0.1 <sup>a</sup>	0.0	0.0 <sup>a</sup>	0.0
Meprobamate	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1
Miltown®	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0
Rohypnol®	0.2	0.2	0.2	0.2	0.5	0.6	0.1	0.1
Serax®	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1
Soma®	1.0	1.1	0.6	0.6	2.3	2.5	0.8	0.9
Tranxene®	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1
Vistaril®	0.2	0.1	0.1	0.0	0.2 <sup>a</sup>	0.1	0.2	0.1

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.131A Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Numbers in Thousands, 2002 and 2003**

Stimulant	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methamphetamine, <sup>1</sup> Desoxyn®, or Methedrine	12,383	12,303	366	328	1,756	1,650	10,261	10,325
Prescription Diet Pills <sup>2</sup>	8,980	8,671	267	251	692	743	8,021	7,677
Ritalin® or Methylphenidate	4,528	4,168	587	547	1,662	1,806	2,280 <sup>a</sup>	1,816
Cylert®	196	198	18	23	39	36	139	140
Dexedrine®	3,274 <sup>b</sup>	2,553	70	69	318	291	2,887 <sup>b</sup>	2,193
Dextroamphetamine	553	498	22	31	78	80	453	387
Didrex®	177	196	21	19	31	25	124	152
Eskatrol®	302 <sup>a</sup>	89	10	4	15	4	276 <sup>a</sup>	81
Ionamin®	491	583	19	26	28	23	445	535
Mazanor®	65	30	16	10	3	8	45	13
Obedrin-LA®	79	18	6	13	6	5	68	*
Plegine®	68	45	12	5	8	2	48	37
Preludin®	856	705	10	13	35	52	811	640
Sanorex®	137	105	12	20	23	17	102	68
Tenuate®	275	146	10	13	10	17	255	116

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Crank, crystal, ice, or speed.

<sup>2</sup> For example, Amphetamines, Benzedrine®, Biphedamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.131B Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Percentages, 2002 and 2003**

Stimulant	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methamphetamine, <sup>1</sup> Desoxy <sup>®</sup> , or Methedrine	5.3	5.2	1.5	1.3	5.7	5.2	5.7	5.7
Prescription Diet Pills <sup>2</sup>	3.8	3.6	1.1	1.0	2.2	2.3	4.5	4.2
Ritalin <sup>®</sup> or Methylphenidate	1.9	1.8	2.4	2.2	5.4	5.7	1.3 <sup>a</sup>	1.0
Cylert <sup>®</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Dexedrine <sup>®</sup>	1.4 <sup>b</sup>	1.1	0.3	0.3	1.0	0.9	1.6 <sup>b</sup>	1.2
Dextroamphetamine	0.2	0.2	0.1	0.1	0.3	0.3	0.3	0.2
Didrex <sup>®</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Eskatrol <sup>®</sup>	0.1 <sup>a</sup>	0.0	0.0	0.0	0.0	0.0	0.2 <sup>a</sup>	0.0
Ionamin <sup>®</sup>	0.2	0.2	0.1	0.1	0.1	0.1	0.2	0.3
Mazanor <sup>®</sup>	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Obedrin-LA <sup>®</sup>	0.0	0.0	0.0	0.1	0.0	0.0	0.0	*
Plegine <sup>®</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Preludin <sup>®</sup>	0.4	0.3	0.0	0.1	0.1	0.2	0.5	0.4
Sanorex <sup>®</sup>	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.0
Tenuate <sup>®</sup>	0.1	0.1	0.0	0.1	0.0	0.1	0.1	0.1

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Crank, crystal, ice, or speed.

<sup>2</sup> For example, Amphetamines, Benzedrine<sup>®</sup>, Biphedamine<sup>®</sup>, Fastin<sup>®</sup>, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 1.132A Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Numbers in Thousands, 2002 and 2003**

Sedative	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methaqualone, Sopor®, or Quaalude®	7,320	6,997	60	64	305	258	6,955	6,676
Barbiturates <sup>1</sup>	3,375	3,301	29	44	121	101	3,226	3,156
Restoril® or Temazepam	1,003	923	48	50	108	90	847	783
Amytal®	274	209	4 <sup>b</sup>	26	21	19	249	164
Butisol®	109	103	8	15	14	12	87	76
Chloral Hydrate	214	149	7	11	9 <sup>a</sup>	25	198	114
Dalmane®	638	435	22	23	16	19	600	393
Halcion®	939 <sup>a</sup>	668	17	23	64	53	857 <sup>a</sup>	592
Phenobarbital	1,568	1,300	37	29	105	91	1,426	1,180
Placidyl®	1,045	783	12	12	34 <sup>a</sup>	14	998	758
Tuinal®	1,393	1,077	8	14	17	10	1,368	1,053

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.132B Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Percentages, 2002 and 2003**

Sedative	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Methaqualone, Sopor®, or Quaalude®	3.1	2.9	0.2	0.3	1.0	0.8	3.9	3.7
Barbiturates <sup>1</sup>	1.4	1.4	0.1	0.2	0.4	0.3	1.8	1.7
Restoril® or Temazepam	0.4	0.4	0.2	0.2	0.3	0.3	0.5	0.4
Amytal®	0.1	0.1	0.0 <sup>b</sup>	0.1	0.1	0.1	0.1	0.1
Butisol®	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
Chloral Hydrate	0.1	0.1	0.0	0.0	0.0 <sup>a</sup>	0.1	0.1	0.1
Dalmane®	0.3	0.2	0.1	0.1	0.1	0.1	0.3	0.2
Halcion®	0.4 <sup>a</sup>	0.3	0.1	0.1	0.2	0.2	0.5 <sup>a</sup>	0.3
Phenobarbital	0.7	0.5	0.2	0.1	0.3	0.3	0.8	0.7
Placidyl®	0.4	0.3	0.1	0.0	0.1 <sup>a</sup>	0.0	0.6	0.4
Tuinal®	0.6	0.5	0.0	0.1	0.1	0.0	0.8	0.6

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.