

**Table 1.108C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	474	441	95	83	64	45
<b>GEOGRAPHIC DIVISION</b>						
Northeast	223	220	36	31	21	12
New England	149	107	28	15	18	7
Middle Atlantic	165	193	22	27	11	10
Midwest	177	168	50	33	38	23
East North Central	146	139	39	26	36	16
West North Central	100	94	31	21	12	16
South	257	248	43	51	23	23
South Atlantic	201	201	31	45	21	18
East South Central	91	72	25	15	3	6
West South Central	136	126	18	21	10	13
West	275	235	57	47	41	28
Mountain	92	134	18	19	12	12
Pacific	259	194	54	43	39	25
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	373	363	70	67	50	37
Small Metro	248	212	58	40	39	23
250K – 1 Mil. Pop.	175	175	51	31	38	18
< 250K Pop.	182	127	29	23	10	13
Nonmetro	149	144	25	27	17	11
Urbanized	115	101	20	15	15	7
Less Urbanized	81	96	13	21	7	6
Completely Rural	49	31	9	7	3	5

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.108D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.20	0.19	0.04	0.03	0.03	0.02
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.50	0.49	0.08	0.07	0.05	0.03
New England	1.26	0.90	0.24	0.13	0.15	0.06
Middle Atlantic	0.50	0.58	0.07	0.08	0.03	0.03
Midwest	0.33	0.31	0.09	0.06	0.07	0.04
East North Central	0.39	0.37	0.10	0.07	0.10	0.04
West North Central	0.63	0.58	0.20	0.13	0.07	0.10
South	0.31	0.29	0.05	0.06	0.03	0.03
South Atlantic	0.46	0.45	0.07	0.10	0.05	0.04
East South Central	0.64	0.50	0.17	0.10	0.02	0.04
West South Central	0.52	0.48	0.07	0.08	0.04	0.05
West	0.52	0.44	0.11	0.09	0.08	0.05
Mountain	0.60	0.86	0.12	0.12	0.08	0.08
Pacific	0.69	0.51	0.14	0.11	0.10	0.07
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.30	0.29	0.06	0.05	0.04	0.03
Small Metro	0.36	0.30	0.08	0.06	0.06	0.03
250K – 1 Mil. Pop.	0.38	0.37	0.11	0.07	0.08	0.04
< 250K Pop.	0.77	0.52	0.12	0.10	0.04	0.05
Nonmetro	0.37	0.36	0.06	0.07	0.04	0.03
Urbanized	0.72	0.56	0.13	0.08	0.09	0.04
Less Urbanized	0.41	0.53	0.07	0.12	0.04	0.04
Completely Rural	1.10	0.77	0.21	0.16	0.07	0.12

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.108P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.8484		0.8088		0.3622	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.5401		0.8165		0.7973	
New England	0.7902		0.2198		0.4350	
Middle Atlantic	0.3430		0.1437		0.1614	
Midwest	0.2377		0.6845		0.7028	
East North Central	0.8079		0.7998		0.3961	
West North Central	0.0894		0.7406		0.4200	
South	0.3543		0.4865		0.4113	
South Atlantic	0.5271		0.8233		0.1463	
East South Central	0.4546		0.2820		0.7107	
West South Central	0.7121		0.8506		0.5718	
West	0.4933		0.6489		0.4911	
Mountain	0.8283		0.5800		0.9399	
Pacific	0.5130		0.7798		0.4562	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.6938		0.8956		0.4985	
Small Metro	0.2311		0.7621		0.8927	
250K – 1 Mil. Pop.	0.9710		0.8614		0.6309	
< 250K Pop.	0.0698		0.7721		0.4488	
Nonmetro	0.7179		0.9173		0.2469	
Urbanized	0.3834		0.1719		0.3271	
Less Urbanized	0.1087		0.0480		0.2127	
Completely Rural	0.3766		0.1765		0.5754	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.109C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	67	68	44	48	22	25
<b>GEOGRAPHIC DIVISION</b>						
Northeast	29	28	17	18	7	9
New England	10	14	7	10	4	4
Middle Atlantic	27	24	15	15	6	9
Midwest	25	28	16	17	8	9
East North Central	17	25	11	15	6	7
West North Central	18	13	11	10	4	5
South	42	41	28	29	15	14
South Atlantic	31	31	23	23	13	9
East South Central	18	13	9	6	3	3
West South Central	22	23	12	16	6	11
West	35	38	26	29	12	16
Mountain	12	19	9	13	4	7
Pacific	32	33	24	25	12	15
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	52	50	32	36	17	18
Small Metro	38	36	28	23	14	15
250K – 1 Mil. Pop.	32	28	23	18	11	12
< 250K Pop.	21	22	16	14	8	9
Nonmetro	24	26	15	18	10	9
Urbanized	14	16	10	11	6	6
Less Urbanized	18	18	12	13	6	6
Completely Rural	9	9	7	5	3	2

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.109D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.27	0.27	0.18	0.19	0.09	0.10
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.64	0.62	0.38	0.40	0.16	0.21
New England	0.85	1.20	0.61	0.89	0.35	0.31
Middle Atlantic	0.81	0.72	0.47	0.45	0.18	0.26
Midwest	0.44	0.49	0.28	0.31	0.14	0.15
East North Central	0.43	0.62	0.28	0.36	0.16	0.18
West North Central	1.07	0.74	0.65	0.56	0.25	0.31
South	0.48	0.45	0.31	0.32	0.17	0.16
South Atlantic	0.71	0.68	0.53	0.50	0.30	0.20
East South Central	1.23	0.88	0.62	0.43	0.23	0.24
West South Central	0.74	0.78	0.40	0.54	0.22	0.36
West	0.59	0.65	0.44	0.49	0.21	0.27
Mountain	0.73	1.12	0.52	0.78	0.25	0.38
Pacific	0.78	0.79	0.58	0.61	0.28	0.35
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.39	0.37	0.24	0.27	0.13	0.13
Small Metro	0.53	0.49	0.39	0.32	0.19	0.20
250K – 1 Mil. Pop.	0.65	0.56	0.46	0.37	0.22	0.23
< 250K Pop.	0.90	0.93	0.66	0.59	0.35	0.38
Nonmetro	0.55	0.61	0.36	0.42	0.22	0.21
Urbanized	0.80	0.86	0.59	0.59	0.38	0.30
Less Urbanized	0.85	0.89	0.57	0.65	0.29	0.32
Completely Rural	1.89	2.05	1.46	1.13	0.55	0.53

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.109P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.6767		0.6422		0.6467	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.6921		0.6327		0.3477	
New England	0.5682		0.5634		0.1779	
Middle Atlantic	0.8407		0.3658		0.0954	
Midwest	0.5518		0.2905		0.3473	
East North Central	0.1917		0.2590		0.9208	
West North Central	0.4713		0.7684		0.1049	
South	0.3687		0.3734		0.3474	
South Atlantic	0.0757		0.3032		0.0098	
East South Central	0.7949		0.2600		0.7689	
West South Central	0.3044		0.5490		0.0678	
West	0.0820		0.4178		0.5692	
Mountain	0.8817		0.8404		0.7193	
Pacific	0.0646		0.4161		0.6388	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.9325		0.7886		0.6800	
Small Metro	0.4108		0.7721		0.7096	
250K – 1 Mil. Pop.	0.4732		0.7017		0.7882	
< 250K Pop.	0.7055		0.9621		0.7889	
Nonmetro	0.9258		0.7683		0.8235	
Urbanized	0.3631		0.5757		0.7965	
Less Urbanized	0.4034		0.6573		0.4657	
Completely Rural	0.7551		0.2906		0.6656	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.110C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	111	104	41	39	18	19
<b>GEOGRAPHIC DIVISION</b>						
Northeast	47	46	22	19	5	9
New England	28	17	17	8	2	3
Middle Atlantic	37	43	13	17	5	9
Midwest	45	41	19	16	7	7
East North Central	38	35	14	12	5	6
West North Central	23	21	12	10	5	3
South	61	62	22	21	11	11
South Atlantic	47	47	17	16	10	9
East South Central	23	23	9	5	*	*
West South Central	30	33	10	12	5	7
West	66	56	19	22	11	11
Mountain	25	27	7	10	3	4
Pacific	61	49	17	20	11	10
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	85	81	33	30	15	14
Small Metro	58	52	21	20	8	10
250K – 1 Mil. Pop.	45	43	16	15	6	9
< 250K Pop.	35	30	14	13	6	6
Nonmetro	37	37	11	16	5	8
Urbanized	28	29	8	11	4	8
Less Urbanized	21	23	5	12	3	1
Completely Rural	9	7	5	0	2	0

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.110D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.36	0.33	0.13	0.12	0.06	0.06
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.85	0.81	0.39	0.34	0.09	0.16
New England	1.95	1.19	1.16	0.54	0.14	0.19
Middle Atlantic	0.90	1.02	0.33	0.42	0.12	0.21
Midwest	0.63	0.56	0.26	0.21	0.10	0.09
East North Central	0.77	0.69	0.29	0.24	0.10	0.12
West North Central	1.04	0.94	0.53	0.44	0.24	0.12
South	0.55	0.55	0.20	0.19	0.10	0.10
South Atlantic	0.87	0.84	0.32	0.29	0.19	0.16
East South Central	1.18	1.19	0.45	0.27	*	*
West South Central	0.82	0.88	0.26	0.32	0.14	0.17
West	0.90	0.76	0.26	0.30	0.15	0.15
Mountain	1.13	1.23	0.33	0.45	0.13	0.18
Pacific	1.19	0.95	0.34	0.39	0.21	0.19
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.54	0.49	0.21	0.18	0.10	0.08
Small Metro	0.57	0.51	0.21	0.19	0.08	0.10
250K – 1 Mil. Pop.	0.70	0.65	0.24	0.23	0.09	0.13
< 250K Pop.	0.95	0.85	0.37	0.35	0.15	0.16
Nonmetro	0.74	0.76	0.23	0.33	0.11	0.16
Urbanized	1.24	1.23	0.38	0.47	0.16	0.32
Less Urbanized	0.95	1.06	0.24	0.55	0.14	0.06
Completely Rural	2.02	1.89	1.05	0.10	0.48	0.06

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 1.110P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.1087		0.5097		0.4771	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4457		0.9755		0.1971	
New England	0.1519		0.2723		0.3986	
Middle Atlantic	0.9038		0.2933		0.2849	
Midwest	0.0723		0.5283		0.4254	
East North Central	0.0432		0.7995		0.9635	
West North Central	0.7742		0.5061		0.2169	
South	0.3936		0.0852		0.3489	
South Atlantic	0.3582		0.1371		0.4802	
East South Central	0.8793		0.1372		*	
West South Central	0.8647		0.7927		0.5063	
West	0.9245		0.3311		0.6089	
Mountain	0.9412		0.6162		0.6026	
Pacific	0.8781		0.4048		0.7061	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.1632		0.2100		0.1342	
Small Metro	0.1105		0.8317		0.3165	
250K – 1 Mil. Pop.	0.8112		0.5251		0.2582	
< 250K Pop.	0.0140		0.7086		0.8743	
Nonmetro	0.3427		0.5503		0.9431	
Urbanized	0.6159		0.6183		0.2771	
Less Urbanized	0.3311		0.0407		0.0917	
Completely Rural	0.4376		0.0200		0.2242	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.111C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	457	407	66	54	56	32
<b>GEOGRAPHIC DIVISION</b>						
Northeast	217	213	23	16	20	7
New England	144	103	20	6	18	4
Middle Atlantic	161	187	11	14	9	5
Midwest	169	158	41	22	38	20
East North Central	139	128	37	16	36	13
West North Central	96	92	18	15	9	15
South	239	221	28	36	12	15
South Atlantic	182	175	14	34	11	14
East South Central	86	60	23	13	*	4
West South Central	132	120	7	*	6	*
West	273	215	38	29	35	20
Mountain	95	123	13	15	11	13
Pacific	256	176	35	25	33	15
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	361	337	53	47	44	29
Small Metro	241	197	37	25	32	14
250K – 1 Mil. Pop.	166	166	35	22	32	11
< 250K Pop.	180	113	12	12	*	7
Nonmetro	140	128	14	11	12	4
Urbanized	109	90	14	4	12	*
Less Urbanized	72	85	1	9	1	*
Completely Rural	43	30	*	4	*	4

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.111D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.25	0.22	0.04	0.03	0.03	0.02
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.62	0.61	0.06	0.04	0.06	0.02
New England	1.57	1.12	0.21	0.06	0.19	0.05
Middle Atlantic	0.62	0.72	0.04	0.06	0.03	0.02
Midwest	0.42	0.39	0.10	0.05	0.09	0.05
East North Central	0.49	0.45	0.13	0.06	0.13	0.05
West North Central	0.80	0.76	0.15	0.12	0.08	0.12
South	0.37	0.34	0.04	0.06	0.02	0.02
South Atlantic	0.54	0.51	0.04	0.10	0.03	0.04
East South Central	0.80	0.55	0.22	0.12	*	0.04
West South Central	0.69	0.62	0.04	*	0.03	*
West	0.69	0.53	0.09	0.07	0.09	0.05
Mountain	0.83	1.06	0.12	0.13	0.10	0.11
Pacific	0.90	0.61	0.12	0.09	0.12	0.05
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.37	0.35	0.05	0.05	0.05	0.03
Small Metro	0.47	0.37	0.07	0.05	0.06	0.03
250K – 1 Mil. Pop.	0.49	0.47	0.10	0.06	0.09	0.03
< 250K Pop.	1.02	0.61	0.07	0.06	*	0.04
Nonmetro	0.45	0.41	0.05	0.04	0.04	0.01
Urbanized	0.90	0.66	0.12	0.03	0.10	*
Less Urbanized	0.47	0.61	0.01	0.07	0.01	*
Completely Rural	1.24	0.90	*	0.13	*	0.13

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.111P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.9175		0.7414		0.2953	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.4017		0.8799		0.3841	
New England	0.9190		0.5694		0.5454	
Middle Atlantic	0.3404		0.6766		0.4769	
Midwest	0.0996		0.4079		0.6377	
East North Central	0.5605		0.4467		0.3637	
West North Central	0.0480		0.7400		0.3884	
South	0.5111		0.4021		0.7410	
South Atlantic	0.8401		0.1210		0.5885	
East South Central	0.4897		0.8603		*	
West South Central	0.5866		*		*	
West	0.3433		0.6017		0.3644	
Mountain	0.7940		0.8723		0.9495	
Pacific	0.3539		0.5149		0.3027	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.4798		0.8185		0.6929	
Small Metro	0.3412		0.4774		0.4623	
250K – 1 Mil. Pop.	0.9189		0.4235		0.3051	
< 250K Pop.	0.1868		0.9317		*	
Nonmetro	0.8804		0.4707		0.1014	
Urbanized	0.2658		0.0509		*	
Less Urbanized	0.1178		0.2535		*	
Completely Rural	0.4038		*		*	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.112C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	472	435	80	66	59	37
<b>GEOGRAPHIC DIVISION</b>						
Northeast	227	220	32	25	20	11
New England	150	108	27	10	18	5
Middle Atlantic	168	192	17	23	10	10
Midwest	175	165	48	28	38	21
East North Central	145	136	40	22	37	14
West North Central	99	93	27	18	11	15
South	258	242	36	42	17	18
South Atlantic	202	197	23	38	15	17
East South Central	91	68	24	13	*	4
West South Central	136	122	12	13	8	7
West	269	233	43	35	37	22
Mountain	92	132	15	18	12	13
Pacific	252	193	40	30	35	18
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	372	362	63	54	47	32
Small Metro	248	206	45	33	33	17
250K – 1 Mil. Pop.	172	174	39	27	33	14
< 250K Pop.	182	120	23	19	6	9
Nonmetro	148	141	19	21	13	9
Urbanized	115	99	18	12	12	8
Less Urbanized	77	93	5	16	3	1
Completely Rural	45	30	5	4	2	4

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.112D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.22	0.20	0.04	0.03	0.03	0.02
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.56	0.54	0.08	0.06	0.05	0.03
New England	1.41	1.01	0.25	0.09	0.17	0.05
Middle Atlantic	0.56	0.64	0.06	0.08	0.03	0.03
Midwest	0.37	0.34	0.10	0.06	0.08	0.04
East North Central	0.43	0.40	0.12	0.06	0.11	0.04
West North Central	0.69	0.65	0.19	0.13	0.08	0.10
South	0.34	0.32	0.05	0.06	0.02	0.02
South Atlantic	0.51	0.49	0.06	0.09	0.04	0.04
East South Central	0.72	0.54	0.19	0.10	*	0.03
West South Central	0.60	0.53	0.05	0.05	0.04	0.03
West	0.57	0.49	0.09	0.07	0.08	0.05
Mountain	0.67	0.95	0.11	0.13	0.09	0.09
Pacific	0.75	0.57	0.12	0.09	0.10	0.05
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.33	0.32	0.06	0.05	0.04	0.03
Small Metro	0.40	0.32	0.07	0.05	0.05	0.03
250K – 1 Mil. Pop.	0.42	0.42	0.10	0.06	0.08	0.03
< 250K Pop.	0.85	0.55	0.11	0.08	0.03	0.04
Nonmetro	0.41	0.39	0.05	0.06	0.04	0.03
Urbanized	0.81	0.62	0.12	0.08	0.09	0.05
Less Urbanized	0.44	0.57	0.03	0.10	0.02	0.01
Completely Rural	1.13	0.83	0.13	0.12	0.06	0.12

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.112P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.7991		0.5630		0.2223	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.5037		0.9872		0.8263	
New England	0.8287		0.2576		0.6583	
Middle Atlantic	0.3251		0.2261		0.8203	
Midwest	0.2673		0.3742		0.5255	
East North Central	0.9671		0.4720		0.3808	
West North Central	0.0667		0.6021		0.7337	
South	0.4371		0.8244		0.7494	
South Atlantic	0.7127		0.6166		0.9787	
East South Central	0.4871		0.5402		*	
West South Central	0.5874		0.3420		0.2298	
West	0.3472		0.9269		0.2968	
Mountain	0.8110		0.6658		0.9329	
Pacific	0.3553		0.9022		0.2697	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.6842		0.7347		0.3674	
Small Metro	0.1772		0.5892		0.7133	
250K – 1 Mil. Pop.	0.8763		0.6808		0.5054	
< 250K Pop.	0.0548		0.7225		0.3896	
Nonmetro	0.7259		0.9468		0.1935	
Urbanized	0.3234		0.0680		0.2155	
Less Urbanized	0.0756		0.0189		0.0786	
Completely Rural	0.2844		0.3110		0.7145	

\*Low precision; no estimate reported.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.