

**Table 1.113C Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	737	641	377	354	229	250
<b>GEOGRAPHIC DIVISION</b>						
Northeast	283	261	157	117	83	83
New England	164	154	116	85	39	46
Middle Atlantic	229	213	104	82	73	69
Midwest	256	243	127	129	80	80
East North Central	199	179	104	112	59	70
West North Central	161	165	73	66	53	38
South	476	345	235	190	161	137
South Atlantic	386	238	194	140	126	101
East South Central	170	143	87	70	69	52
West South Central	219	204	102	107	73	76
West	403	399	215	240	115	174
Mountain	167	148	60	89	38	60
Pacific	367	370	206	223	109	163
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	582	508	279	279	168	202
Small Metro	387	330	209	185	122	121
250K – 1 Mil. Pop.	308	274	152	156	99	95
< 250K Pop.	228	190	136	105	74	77
Nonmetro	265	241	135	125	91	82
Urbanized	155	152	78	82	41	54
Less Urbanized	190	153	79	87	46	61
Completely Rural	102	86	76	25	70	9

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.113D Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.31	0.27	0.16	0.15	0.10	0.11
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.63	0.58	0.35	0.26	0.18	0.18
New England	1.39	1.29	0.98	0.72	0.33	0.39
Middle Atlantic	0.69	0.64	0.31	0.24	0.22	0.21
Midwest	0.48	0.45	0.24	0.24	0.15	0.15
East North Central	0.53	0.47	0.28	0.30	0.16	0.19
West North Central	1.00	1.03	0.46	0.41	0.33	0.24
South	0.57	0.41	0.28	0.22	0.19	0.16
South Atlantic	0.88	0.54	0.44	0.32	0.29	0.23
East South Central	1.20	1.01	0.61	0.49	0.49	0.37
West South Central	0.85	0.78	0.39	0.41	0.28	0.29
West	0.76	0.74	0.41	0.45	0.22	0.32
Mountain	1.09	0.95	0.39	0.57	0.24	0.39
Pacific	0.98	0.97	0.55	0.58	0.29	0.43
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.46	0.40	0.22	0.22	0.13	0.16
Small Metro	0.56	0.46	0.30	0.26	0.18	0.17
250K – 1 Mil. Pop.	0.68	0.58	0.33	0.33	0.22	0.20
< 250K Pop.	0.96	0.78	0.57	0.43	0.31	0.31
Nonmetro	0.66	0.60	0.34	0.31	0.23	0.20
Urbanized	0.97	0.85	0.48	0.46	0.26	0.30
Less Urbanized	0.97	0.84	0.40	0.48	0.23	0.34
Completely Rural	2.29	2.11	1.72	0.60	1.58	0.23

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.113P Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.3887		0.7744		0.8646	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.6161		0.9412		0.8739	
New England	0.9396		0.8005		0.9418	
Middle Atlantic	0.5797		0.6878		0.8885	
Midwest	0.2479		0.6052		0.3257	
East North Central	0.2983		0.6071		0.3857	
West North Central	0.5759		0.0815		0.6246	
South	0.9450		0.5436		0.2780	
South Atlantic	0.9910		0.5297		0.4531	
East South Central	0.5308		0.2440		0.0301	
West South Central	0.7456		0.4382		0.4519	
West	0.7926		0.4725		0.5008	
Mountain	0.4061		0.1886		0.7987	
Pacific	0.9451		0.7550		0.4265	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.4444		0.6432		0.8249	
Small Metro	0.8359		0.8339		0.8803	
250K – 1 Mil. Pop.	0.6535		0.8630		0.3517	
< 250K Pop.	0.7654		0.9074		0.3905	
Nonmetro	0.3429		0.9254		0.8667	
Urbanized	0.6292		0.7063		0.2378	
Less Urbanized	0.1690		0.4892		0.3347	
Completely Rural	0.6324		0.1335		0.0325	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.114C Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	75	76	64	64	42	42
<b>GEOGRAPHIC DIVISION</b>						
Northeast	26	31	23	24	14	16
New England	16	14	11	14	6	8
Middle Atlantic	21	27	20	19	13	13
Midwest	29	29	24	25	15	16
East North Central	24	25	18	22	12	14
West North Central	16	14	16	12	9	8
South	46	47	41	41	27	26
South Atlantic	33	36	31	31	21	18
East South Central	18	20	15	16	10	11
West South Central	27	24	23	21	14	15
West	44	41	36	36	25	24
Mountain	18	21	15	21	12	13
Pacific	40	35	33	29	22	21
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	58	52	48	45	31	32
Small Metro	38	41	34	33	24	19
250K – 1 Mil. Pop.	31	34	29	29	21	16
< 250K Pop.	24	22	19	17	11	11
Nonmetro	27	31	26	27	17	19
Urbanized	17	18	15	17	11	12
Less Urbanized	22	23	18	19	11	13
Completely Rural	11	7	11	7	7	6

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.114D Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.30	0.30	0.26	0.26	0.17	0.17
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.59	0.69	0.51	0.53	0.32	0.35
New England	1.35	1.24	0.94	1.18	0.53	0.69
Middle Atlantic	0.63	0.82	0.60	0.59	0.39	0.41
Midwest	0.51	0.50	0.42	0.44	0.26	0.29
East North Central	0.60	0.62	0.45	0.54	0.30	0.36
West North Central	0.96	0.85	0.92	0.73	0.52	0.47
South	0.53	0.53	0.47	0.46	0.31	0.29
South Atlantic	0.75	0.81	0.71	0.70	0.48	0.41
East South Central	1.28	1.37	1.06	1.10	0.72	0.77
West South Central	0.92	0.80	0.77	0.70	0.47	0.51
West	0.75	0.70	0.62	0.61	0.42	0.41
Mountain	1.03	1.21	0.88	1.24	0.68	0.73
Pacific	0.98	0.84	0.80	0.69	0.53	0.50
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.44	0.39	0.36	0.34	0.23	0.24
Small Metro	0.53	0.55	0.47	0.45	0.33	0.26
250K – 1 Mil. Pop.	0.64	0.69	0.60	0.58	0.44	0.33
< 250K Pop.	1.01	0.90	0.83	0.70	0.48	0.45
Nonmetro	0.64	0.73	0.60	0.64	0.39	0.44
Urbanized	0.99	0.99	0.86	0.92	0.66	0.66
Less Urbanized	1.06	1.16	0.87	0.97	0.51	0.68
Completely Rural	2.41	1.69	2.39	1.65	1.54	1.40

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.114P Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.5012		0.8584		0.8951	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.8755		0.5924		0.7289	
New England	0.1951		0.9567		0.2120	
Middle Atlantic	0.4341		0.4977		0.2619	
Midwest	0.0071		0.0400		0.8490	
East North Central	0.0292		0.0595		0.5678	
West North Central	0.1118		0.3760		0.6069	
South	0.7513		0.8950		0.5407	
South Atlantic	0.6253		0.4547		0.0824	
East South Central	0.2730		0.1498		0.7109	
West South Central	0.6587		0.7596		0.4798	
West	0.8402		0.6958		0.6398	
Mountain	0.3538		0.5211		0.3937	
Pacific	0.8221		0.9342		0.9377	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.0798		0.4603		0.7803	
Small Metro	0.2344		0.9568		0.9952	
250K – 1 Mil. Pop.	0.4708		0.5753		0.3089	
< 250K Pop.	0.3285		0.4186		0.0732	
Nonmetro	0.9949		0.3311		0.4158	
Urbanized	0.2158		0.0267		0.2604	
Less Urbanized	0.4488		0.7708		0.9864	
Completely Rural	0.5902		0.5897		0.9677	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.115C Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	135	135	96	103	60	70
<b>GEOGRAPHIC DIVISION</b>						
Northeast	54	47	38	41	22	26
New England	30	21	23	26	12	14
Middle Atlantic	44	43	30	33	18	22
Midwest	54	56	42	40	25	28
East North Central	44	47	35	32	21	22
West North Central	31	30	22	24	14	17
South	83	88	61	65	41	43
South Atlantic	60	63	43	48	29	35
East South Central	37	34	29	25	17	15
West South Central	45	51	33	37	23	19
West	75	72	47	55	28	41
Mountain	24	34	18	29	14	18
Pacific	69	63	42	47	24	37
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	105	105	73	77	42	54
Small Metro	73	70	52	56	38	36
250K – 1 Mil. Pop.	58	60	42	47	30	26
< 250K Pop.	45	41	31	32	23	25
Nonmetro	44	43	36	38	20	25
Urbanized	32	30	27	24	13	17
Less Urbanized	27	29	21	25	14	17
Completely Rural	15	10	10	7	7	4

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.115D Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.43	0.43	0.31	0.33	0.19	0.22
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.98	0.83	0.68	0.73	0.39	0.45
New England	2.08	1.44	1.61	1.80	0.82	0.92
Middle Atlantic	1.07	1.02	0.74	0.79	0.45	0.53
Midwest	0.76	0.76	0.58	0.55	0.35	0.38
East North Central	0.88	0.93	0.71	0.64	0.43	0.44
West North Central	1.39	1.32	0.97	1.06	0.61	0.76
South	0.76	0.78	0.56	0.58	0.37	0.38
South Atlantic	1.12	1.13	0.79	0.85	0.54	0.64
East South Central	1.91	1.75	1.51	1.28	0.90	0.75
West South Central	1.20	1.35	0.89	0.97	0.63	0.50
West	1.03	0.97	0.64	0.75	0.38	0.56
Mountain	1.11	1.51	0.83	1.29	0.65	0.81
Pacific	1.36	1.22	0.81	0.90	0.46	0.71
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.66	0.63	0.46	0.46	0.27	0.32
Small Metro	0.72	0.69	0.51	0.55	0.37	0.36
250K – 1 Mil. Pop.	0.91	0.90	0.64	0.70	0.47	0.40
< 250K Pop.	1.21	1.14	0.84	0.89	0.62	0.70
Nonmetro	0.89	0.87	0.73	0.78	0.41	0.51
Urbanized	1.41	1.27	1.19	1.03	0.59	0.71
Less Urbanized	1.19	1.30	0.93	1.13	0.63	0.79
Completely Rural	3.13	2.73	2.15	1.96	1.57	1.07

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 1.115P Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0228		0.5291		0.0300	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0246		0.4294		0.1957	
New England	0.5230		0.3651		0.7610	
Middle Atlantic	0.0270		0.0911		0.2025	
Midwest	0.4769		0.5996		0.2033	
East North Central	0.8867		0.4646		0.6581	
West North Central	0.1361		0.0609		0.1582	
South	0.3854		0.2161		0.8469	
South Atlantic	0.5821		0.4924		0.1389	
East South Central	0.5941		0.1116		0.2431	
West South Central	0.1789		0.9803		0.4599	
West	0.2529		0.0597		0.0420	
Mountain	0.8290		0.9110		0.6294	
Pacific	0.2431		0.0318		0.0378	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.0648		0.4359		0.0953	
Small Metro	0.7526		0.4769		0.8517	
250K – 1 Mil. Pop.	0.4706		0.5927		0.7847	
< 250K Pop.	0.6339		0.7015		0.5400	
Nonmetro	0.0259		0.2670		0.0265	
Urbanized	0.0862		0.6091		0.0406	
Less Urbanized	0.0878		0.0890		0.0618	
Completely Rural	0.7201		0.0730		0.0052	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.116C Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	707	617	345	331	215	237
<b>GEOGRAPHIC DIVISION</b>						
Northeast	270	257	146	104	80	72
New England	152	155	108	81	39	44
Middle Atlantic	221	207	98	67	70	57
Midwest	238	240	112	120	70	72
East North Central	184	174	90	102	50	62
West North Central	152	165	68	63	49	36
South	462	332	213	172	151	127
South Atlantic	378	238	175	126	116	91
East South Central	157	128	78	64	67	52
West South Central	214	193	93	97	70	72
West	380	374	197	232	110	172
Mountain	154	140	57	82	40	58
Pacific	347	347	188	216	103	161
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	555	484	259	267	161	195
Small Metro	371	318	190	174	108	112
250K – 1 Mil. Pop.	299	257	140	144	88	87
< 250K Pop.	213	178	123	97	65	72
Nonmetro	252	233	115	104	85	74
Urbanized	146	144	66	69	36	46
Less Urbanized	181	147	65	73	41	58
Completely Rural	86	89	61	25	61	4

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.116D Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.39	0.34	0.19	0.18	0.12	0.13
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.77	0.73	0.42	0.30	0.23	0.20
New England	1.64	1.68	1.17	0.88	0.42	0.48
Middle Atlantic	0.85	0.80	0.38	0.26	0.27	0.22
Midwest	0.59	0.59	0.28	0.30	0.17	0.18
East North Central	0.65	0.61	0.32	0.36	0.18	0.22
West North Central	1.25	1.36	0.56	0.52	0.40	0.30
South	0.72	0.51	0.33	0.27	0.24	0.20
South Atlantic	1.11	0.69	0.51	0.37	0.34	0.27
East South Central	1.45	1.18	0.72	0.59	0.62	0.48
West South Central	1.12	1.00	0.49	0.50	0.37	0.37
West	0.96	0.92	0.50	0.57	0.28	0.42
Mountain	1.35	1.20	0.50	0.71	0.35	0.50
Pacific	1.22	1.20	0.66	0.75	0.36	0.56
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.57	0.50	0.27	0.28	0.17	0.20
Small Metro	0.72	0.59	0.37	0.32	0.21	0.21
250K – 1 Mil. Pop.	0.88	0.73	0.41	0.41	0.26	0.25
< 250K Pop.	1.20	0.96	0.69	0.52	0.37	0.39
Nonmetro	0.82	0.75	0.37	0.34	0.27	0.24
Urbanized	1.21	1.05	0.55	0.51	0.30	0.34
Less Urbanized	1.18	1.05	0.43	0.52	0.27	0.41
Completely Rural	2.47	2.70	1.76	0.76	1.75	0.14

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.116P Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.6417		0.9305		0.6434	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.9117		0.6155		0.7207	
New England	0.9157		0.6222		0.8430	
Middle Atlantic	0.9507		0.1875		0.5642	
Midwest	0.2014		0.4882		0.5729	
East North Central	0.1730		0.9887		0.5103	
West North Central	0.6941		0.2237		0.9211	
South	0.8518		0.7116		0.2661	
South Atlantic	0.9493		0.6737		0.3327	
East South Central	0.5287		0.3379		0.0489	
West South Central	0.4430		0.4801		0.4252	
West	0.9715		0.7478		0.8506	
Mountain	0.4874		0.2512		0.5442	
Pacific	0.7880		0.9524		0.6803	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.5745		0.8274		0.8166	
Small Metro	0.7151		0.9240		0.8798	
250K – 1 Mil. Pop.	0.4703		0.9312		0.4898	
< 250K Pop.	0.6793		0.9677		0.5397	
Nonmetro	0.5415		0.6519		0.5433	
Urbanized	0.3378		0.9575		0.6633	
Less Urbanized	0.2256		0.8801		0.7137	
Completely Rural	0.5149		0.1961		0.0373	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.117C Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	730	635	367	351	226	249
<b>GEOGRAPHIC DIVISION</b>						
Northeast	280	263	154	115	81	83
New England	159	153	114	85	38	49
Middle Atlantic	229	216	101	80	71	67
Midwest	249	240	123	128	77	78
East North Central	195	178	101	110	57	68
West North Central	156	161	71	66	51	36
South	471	345	228	190	160	135
South Atlantic	380	240	186	138	124	99
East South Central	166	140	85	73	67	55
West South Central	220	204	100	106	75	74
West	401	391	209	239	115	175
Mountain	163	142	58	87	38	60
Pacific	367	364	201	223	108	164
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	581	504	275	282	169	202
Small Metro	380	328	203	183	118	120
250K – 1 Mil. Pop.	304	270	145	154	94	95
< 250K Pop.	225	187	136	101	73	74
Nonmetro	259	237	126	120	87	80
Urbanized	153	149	75	80	39	51
Less Urbanized	185	151	72	81	43	60
Completely Rural	94	88	67	24	64	6

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.117D Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.35	0.30	0.17	0.17	0.11	0.12
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.69	0.64	0.38	0.28	0.20	0.20
New England	1.49	1.43	1.07	0.79	0.36	0.46
Middle Atlantic	0.76	0.72	0.34	0.26	0.24	0.22
Midwest	0.52	0.50	0.26	0.27	0.16	0.16
East North Central	0.58	0.53	0.30	0.33	0.17	0.20
West North Central	1.09	1.12	0.49	0.46	0.36	0.25
South	0.63	0.46	0.30	0.25	0.21	0.18
South Atlantic	0.97	0.60	0.47	0.35	0.32	0.25
East South Central	1.31	1.10	0.66	0.57	0.53	0.43
West South Central	0.96	0.88	0.44	0.46	0.33	0.32
West	0.85	0.82	0.44	0.50	0.24	0.36
Mountain	1.20	1.02	0.42	0.62	0.28	0.43
Pacific	1.10	1.07	0.60	0.65	0.32	0.48
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.52	0.45	0.24	0.25	0.15	0.18
Small Metro	0.61	0.51	0.33	0.29	0.19	0.19
250K – 1 Mil. Pop.	0.75	0.64	0.36	0.37	0.23	0.23
< 250K Pop.	1.05	0.85	0.63	0.46	0.34	0.34
Nonmetro	0.72	0.66	0.35	0.33	0.24	0.22
Urbanized	1.07	0.93	0.53	0.50	0.27	0.32
Less Urbanized	1.05	0.93	0.41	0.50	0.25	0.37
Completely Rural	2.38	2.41	1.70	0.67	1.62	0.16

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.117P Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.3485		0.7429		0.8811	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.6035		0.8556		0.9273	
New England	0.8196		0.7865		0.7703	
Middle Atlantic	0.6367		0.5684		0.9311	
Midwest	0.1505		0.3766		0.3284	
East North Central	0.1955		0.8429		0.4421	
West North Central	0.4861		0.0547		0.5443	
South	0.9717		0.5144		0.3242	
South Atlantic	0.9446		0.6054		0.6285	
East South Central	0.6267		0.1581		0.0221	
West South Central	0.6893		0.4860		0.5405	
West	0.8145		0.4959		0.5489	
Mountain	0.4694		0.2429		0.6519	
Pacific	0.9568		0.7539		0.4311	
<b>COUNTY TYPE<sup>1</sup></b>						
Large Metro	0.3386		0.5561		0.7910	
Small Metro	0.7281		0.8429		0.8828	
250K – 1 Mil. Pop.	0.5852		0.9523		0.4600	
< 250K Pop.	0.8466		0.8071		0.5150	
Nonmetro	0.3356		0.9266		0.9909	
Urbanized	0.5107		0.9835		0.3308	
Less Urbanized	0.1321		0.4365		0.3345	
Completely Rural	0.5901		0.1125		0.0174	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.