

Table 1.98C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	306	264	124	124	68	91
GEOGRAPHIC DIVISION						
Northeast	126	117	67	43	27	26
New England	76	59	53	11	8	6
Middle Atlantic	100	100	42	41	26	25
Midwest	117	105	61	52	31	29
East North Central	88	77	46	38	28	28
West North Central	77	71	40	35	14	6
South	181	165	64	97	43	76
South Atlantic	145	132	49	86	36	67
East South Central	76	53	32	28	22	25
West South Central	80	82	24	36	8	26
West	178	133	56	37	32	32
Mountain	49	63	24	27	15	24
Pacific	171	118	50	26	28	21
COUNTY TYPE¹						
Large Metro	238	200	94	91	57	63
Small Metro	159	140	73	52	30	34
250K – 1 Mil. Pop.	129	103	45	35	24	24
< 250K Pop.	94	83	58	37	18	24
Nonmetro	88	101	39	65	19	55
Urbanized	52	61	20	27	15	8
Less Urbanized	66	75	33	57	11	54
Completely Rural	25	30	8	9	2	8

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.98D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.13	0.11	0.05	0.05	0.03	0.04
GEOGRAPHIC DIVISION						
Northeast	0.28	0.26	0.15	0.09	0.06	0.06
New England	0.65	0.50	0.45	0.09	0.07	0.05
Middle Atlantic	0.30	0.30	0.12	0.12	0.08	0.07
Midwest	0.22	0.20	0.11	0.10	0.06	0.05
East North Central	0.23	0.21	0.12	0.10	0.07	0.07
West North Central	0.48	0.44	0.25	0.22	0.09	0.04
South	0.22	0.19	0.08	0.11	0.05	0.09
South Atlantic	0.33	0.30	0.11	0.19	0.08	0.15
East South Central	0.54	0.37	0.23	0.20	0.16	0.18
West South Central	0.31	0.31	0.09	0.14	0.03	0.10
West	0.34	0.25	0.11	0.07	0.06	0.06
Mountain	0.32	0.40	0.16	0.17	0.10	0.15
Pacific	0.46	0.31	0.13	0.07	0.08	0.06
COUNTY TYPE¹						
Large Metro	0.19	0.16	0.07	0.07	0.05	0.05
Small Metro	0.23	0.20	0.11	0.07	0.04	0.05
250K – 1 Mil. Pop.	0.28	0.22	0.10	0.07	0.05	0.05
< 250K Pop.	0.40	0.34	0.24	0.15	0.08	0.10
Nonmetro	0.22	0.25	0.10	0.16	0.05	0.14
Urbanized	0.33	0.34	0.13	0.15	0.09	0.05
Less Urbanized	0.34	0.41	0.17	0.31	0.06	0.30
Completely Rural	0.57	0.73	0.18	0.22	0.04	0.20

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.98P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.1637		0.3425		0.7889	
GEOGRAPHIC DIVISION						
Northeast	0.8895		0.7020		0.6313	
New England	0.6143		0.2812		0.5589	
Middle Atlantic	0.9146		0.7404		0.7556	
Midwest	0.7119		0.0877		0.0896	
East North Central	0.8987		0.2928		0.6215	
West North Central	0.6808		0.1702		0.0006	
South	0.6863		0.3064		0.1164	
South Atlantic	0.4611		0.9120		0.3598	
East South Central	0.1875		0.5379		0.7392	
West South Central	0.1978		0.0881		0.0570	
West	0.0527		0.0869		0.7645	
Mountain	0.5253		0.8570		0.7829	
Pacific	0.0183		0.0629		0.5592	
COUNTY TYPE¹						
Large Metro	0.3532		0.4479		0.6751	
Small Metro	0.0565		0.3124		0.4953	
250K – 1 Mil. Pop.	0.1338		0.1566		0.9190	
< 250K Pop.	0.2040		0.9203		0.3591	
Nonmetro	0.2588		0.7365		0.5194	
Urbanized	0.1593		0.7140		0.5588	
Less Urbanized	0.6802		0.7235		0.4591	
Completely Rural	0.8289		0.5113		0.4317	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.99C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	17	17	14	13	9	8
GEOGRAPHIC DIVISION						
Northeast	5	5	5	3	2	1
New England	3	1	2	1	2	0
Middle Atlantic	4	4	4	3	1	1
Midwest	5	7	5	6	1	4
East North Central	4	5	4	4	0	2
West North Central	4	4	4	4	0	3
South	10	10	10	8	8	5
South Atlantic	8	8	8	7	7	4
East South Central	4	2	3	2	2	1
West South Central	5	6	5	4	2	2
West	11	10	7	7	3	4
Mountain	5	6	4	5	1	4
Pacific	10	9	5	5	3	*
COUNTY TYPE¹						
Large Metro	10	13	7	10	2	7
Small Metro	10	7	8	5	5	2
250K – 1 Mil. Pop.	9	6	5	3	3	2
< 250K Pop.	6	5	5	4	3	2
Nonmetro	10	8	9	6	7	2
Urbanized	7	3	6	3	6	1
Less Urbanized	7	5	5	4	3	2
Completely Rural	3	5	3	3	*	0

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.99D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.07	0.07	0.06	0.05	0.03	0.03
GEOGRAPHIC DIVISION						
Northeast	0.12	0.10	0.11	0.07	0.05	0.02
New England	0.28	0.09	0.21	0.06	0.19	0.03
Middle Atlantic	0.13	0.13	0.12	0.10	0.02	0.02
Midwest	0.10	0.12	0.09	0.10	0.01	0.07
East North Central	0.10	0.13	0.09	0.10	0.01	0.06
West North Central	0.21	0.26	0.21	0.24	0.01	0.20
South	0.12	0.11	0.11	0.09	0.09	0.06
South Atlantic	0.19	0.17	0.19	0.15	0.17	0.10
East South Central	0.25	0.12	0.18	0.10	0.15	0.08
West South Central	0.17	0.21	0.16	0.15	0.06	0.08
West	0.20	0.18	0.12	0.12	0.05	0.06
Mountain	0.30	0.34	0.23	0.30	0.03	0.21
Pacific	0.25	0.21	0.13	0.12	0.07	*
COUNTY TYPE¹						
Large Metro	0.07	0.10	0.05	0.08	0.01	0.05
Small Metro	0.14	0.10	0.10	0.07	0.07	0.03
250K – 1 Mil. Pop.	0.18	0.11	0.11	0.06	0.07	0.04
< 250K Pop.	0.27	0.20	0.22	0.15	0.15	0.06
Nonmetro	0.24	0.18	0.21	0.13	0.16	0.04
Urbanized	0.38	0.19	0.37	0.17	0.36	0.06
Less Urbanized	0.34	0.24	0.25	0.19	0.14	0.08
Completely Rural	0.68	1.07	0.68	0.57	*	0.04

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.99P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.7236		0.7667		0.6129	
GEOGRAPHIC DIVISION						
Northeast	0.6201		0.6339		0.3046	
New England	0.0433		0.1282		0.2379	
Middle Atlantic	0.5686		0.8446		0.8927	
Midwest	0.0567		0.2914		0.0019	
East North Central	0.2418		0.4075		0.0402	
West North Central	0.1044		0.4998		0.0157	
South	0.5724		0.4811		0.5116	
South Atlantic	0.3147		0.5889		0.4566	
East South Central	0.0954		0.2881		0.4404	
West South Central	0.2078		0.9952		0.4314	
West	0.3959		0.8859		0.7935	
Mountain	0.8242		0.8135		0.2597	
Pacific	0.3832		0.7005		*	
COUNTY TYPE¹						
Large Metro	0.0468		0.0854		0.0085	
Small Metro	0.0439		0.1445		0.5601	
250K – 1 Mil. Pop.	0.0753		0.1100		0.7904	
< 250K Pop.	0.5420		0.6269		0.5655	
Nonmetro	0.1994		0.1908		0.1804	
Urbanized	0.4623		0.4202		0.3172	
Less Urbanized	0.0943		0.3089		0.2952	
Completely Rural	0.5078		0.7580		*	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.100C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	51	54	23	23	9	13
GEOGRAPHIC DIVISION						
Northeast	18	17	7	9	3	2
New England	8	7	5	4	2	1
Middle Atlantic	16	15	5	8	2	2
Midwest	22	21	12	10	5	4
East North Central	15	15	6	8	4	3
West North Central	16	15	11	7	3	3
South	31	30	14	13	6	4
South Atlantic	21	21	11	9	4	4
East South Central	15	14	5	5	2	2
West South Central	16	16	8	7	3	1
West	30	36	12	14	4	12
Mountain	14	14	6	8	2	5
Pacific	26	34	10	12	4	11
COUNTY TYPE¹						
Large Metro	36	40	16	18	7	12
Small Metro	31	30	13	13	5	5
250K – 1 Mil. Pop.	22	26	10	11	4	4
< 250K Pop.	21	16	8	7	2	3
Nonmetro	19	20	12	9	4	3
Urbanized	11	13	5	6	1	2
Less Urbanized	13	13	7	5	3	1
Completely Rural	6	5	6	3	*	*

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.100D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.16	0.17	0.08	0.07	0.03	0.04
GEOGRAPHIC DIVISION						
Northeast	0.32	0.30	0.13	0.16	0.05	0.04
New England	0.54	0.48	0.33	0.27	0.13	0.06
Middle Atlantic	0.39	0.36	0.13	0.19	0.04	0.05
Midwest	0.30	0.28	0.17	0.14	0.07	0.06
East North Central	0.29	0.29	0.13	0.16	0.09	0.06
West North Central	0.73	0.65	0.47	0.29	0.14	0.12
South	0.28	0.26	0.13	0.11	0.05	0.04
South Atlantic	0.39	0.38	0.20	0.16	0.08	0.07
East South Central	0.80	0.73	0.28	0.25	0.11	0.10
West South Central	0.43	0.41	0.21	0.19	0.09	0.02
West	0.41	0.49	0.16	0.19	0.06	0.16
Mountain	0.62	0.62	0.29	0.36	0.10	0.21
Pacific	0.51	0.65	0.20	0.22	0.08	0.20
COUNTY TYPE¹						
Large Metro	0.22	0.24	0.10	0.11	0.04	0.07
Small Metro	0.30	0.29	0.13	0.13	0.05	0.05
250K – 1 Mil. Pop.	0.34	0.38	0.16	0.17	0.07	0.06
< 250K Pop.	0.57	0.44	0.20	0.21	0.06	0.09
Nonmetro	0.39	0.40	0.24	0.17	0.08	0.05
Urbanized	0.48	0.55	0.23	0.24	0.03	0.10
Less Urbanized	0.57	0.59	0.33	0.24	0.14	0.06
Completely Rural	1.30	1.50	1.23	0.90	*	*

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.100P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.8333		0.9486		0.9523	
GEOGRAPHIC DIVISION						
Northeast	0.1984		0.3262		0.5772	
New England	0.5534		0.9021		0.0689	
Middle Atlantic	0.0930		0.2308		0.4943	
Midwest	0.7987		0.5564		0.6635	
East North Central	0.1433		0.0591		0.6700	
West North Central	0.4103		0.5779		0.8747	
South	0.3846		0.2675		0.2465	
South Atlantic	0.2627		0.1849		0.5549	
East South Central	0.7696		0.6057		0.8643	
West South Central	0.9371		0.5966		0.1866	
West	0.7849		0.9149		0.2437	
Mountain	0.0220		0.7451		0.4634	
Pacific	0.4031		0.7315		0.3436	
COUNTY TYPE¹						
Large Metro	0.7483		0.8087		0.9528	
Small Metro	0.7194		0.5029		0.7957	
250K – 1 Mil. Pop.	0.6826		0.3434		0.9347	
< 250K Pop.	0.3130		0.8804		0.5392	
Nonmetro	0.5081		0.8458		0.7030	
Urbanized	0.0734		0.2384		0.1259	
Less Urbanized	0.7861		0.7695		0.3006	
Completely Rural	0.5367		0.2505		*	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.101C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	303	257	121	120	66	90
GEOGRAPHIC DIVISION						
Northeast	126	117	67	43	27	25
New England	73	60	53	10	7	6
Middle Atlantic	102	101	41	41	26	25
Midwest	116	101	60	49	30	27
East North Central	86	76	44	36	26	27
West North Central	78	66	40	33	14	3
South	175	157	61	94	42	76
South Atlantic	142	127	47	82	35	67
East South Central	73	48	32	29	22	24
West South Central	77	78	23	35	7	26
West	179	133	55	36	31	31
Mountain	47	61	21	26	15	23
Pacific	174	118	51	25	28	21
COUNTY TYPE¹						
Large Metro	237	199	92	89	56	61
Small Metro	159	135	72	50	29	35
250K – 1 Mil. Pop.	127	98	44	34	23	26
< 250K Pop.	95	83	57	36	18	23
Nonmetro	84	98	38	62	17	55
Urbanized	50	60	19	26	14	8
Less Urbanized	64	75	32	56	10	54
Completely Rural	21	29	6	8	*	8

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.101D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.17	0.14	0.07	0.07	0.04	0.05
GEOGRAPHIC DIVISION						
Northeast	0.36	0.33	0.19	0.12	0.08	0.07
New England	0.80	0.65	0.57	0.10	0.08	0.06
Middle Atlantic	0.39	0.39	0.16	0.16	0.10	0.10
Midwest	0.29	0.25	0.15	0.12	0.07	0.07
East North Central	0.30	0.27	0.16	0.13	0.09	0.10
West North Central	0.65	0.55	0.33	0.28	0.11	0.03
South	0.27	0.24	0.10	0.15	0.07	0.12
South Atlantic	0.42	0.37	0.14	0.24	0.10	0.19
East South Central	0.67	0.44	0.29	0.26	0.20	0.23
West South Central	0.40	0.40	0.12	0.18	0.04	0.14
West	0.45	0.33	0.14	0.09	0.08	0.08
Mountain	0.41	0.52	0.19	0.22	0.13	0.20
Pacific	0.61	0.41	0.18	0.09	0.10	0.07
COUNTY TYPE¹						
Large Metro	0.25	0.21	0.10	0.09	0.06	0.06
Small Metro	0.31	0.25	0.14	0.09	0.06	0.06
250K – 1 Mil. Pop.	0.37	0.28	0.13	0.10	0.07	0.07
< 250K Pop.	0.54	0.45	0.32	0.20	0.10	0.13
Nonmetro	0.27	0.32	0.12	0.20	0.05	0.18
Urbanized	0.42	0.44	0.16	0.19	0.11	0.06
Less Urbanized	0.42	0.53	0.21	0.40	0.07	0.38
Completely Rural	0.59	0.87	0.16	0.25	*	0.25

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.101P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.1751		0.3408		0.8281	
GEOGRAPHIC DIVISION						
Northeast	0.9845		0.6283		0.7074	
New England	0.5148		0.3170		0.9270	
Middle Atlantic	0.7501		0.8816		0.7153	
Midwest	0.5847		0.0488		0.0444	
East North Central	0.6526		0.1395		0.5689	
West North Central	0.7527		0.1937		0.0000	
South	0.8369		0.1891		0.0792	
South Atlantic	0.6292		0.6985		0.2898	
East South Central	0.2301		0.5460		0.6782	
West South Central	0.2128		0.0637		0.0480	
West	0.0636		0.0790		0.5063	
Mountain	0.2332		0.9063		1.0000	
Pacific	0.0117		0.0532		0.3971	
COUNTY TYPE¹						
Large Metro	0.3357		0.3691		0.5170	
Small Metro	0.0760		0.3105		0.4793	
250K – 1 Mil. Pop.	0.1280		0.1227		0.8878	
< 250K Pop.	0.3066		0.9888		0.3531	
Nonmetro	0.2513		0.5625		0.3958	
Urbanized	0.2820		0.7465		0.6525	
Less Urbanized	0.5331		0.6304		0.3876	
Completely Rural	0.8907		0.7981		*	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.102C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	307	263	124	122	67	91
GEOGRAPHIC DIVISION						
Northeast	125	116	67	43	27	26
New England	76	59	53	11	8	6
Middle Atlantic	100	100	41	41	26	25
Midwest	118	105	61	51	31	28
East North Central	89	78	46	37	28	28
West North Central	78	70	40	35	14	4
South	181	163	63	95	43	76
South Atlantic	145	132	49	83	35	67
East South Central	76	51	32	28	22	25
West South Central	79	82	23	35	8	26
West	180	134	56	37	32	32
Mountain	49	62	24	27	15	24
Pacific	173	119	50	26	28	21
COUNTY TYPE¹						
Large Metro	239	200	93	90	57	62
Small Metro	160	140	73	51	30	34
250K – 1 Mil. Pop.	129	102	45	34	24	24
< 250K Pop.	94	83	57	37	18	24
Nonmetro	87	99	39	62	17	55
Urbanized	52	61	19	27	14	8
Less Urbanized	65	75	32	56	10	54
Completely Rural	24	29	8	9	2	8

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.102D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.15	0.12	0.06	0.06	0.03	0.04
GEOGRAPHIC DIVISION						
Northeast	0.31	0.28	0.17	0.10	0.07	0.06
New England	0.71	0.55	0.50	0.10	0.07	0.06
Middle Atlantic	0.33	0.33	0.14	0.14	0.09	0.08
Midwest	0.25	0.22	0.13	0.11	0.06	0.06
East North Central	0.27	0.23	0.14	0.11	0.08	0.08
West North Central	0.54	0.49	0.28	0.24	0.10	0.03
South	0.24	0.22	0.08	0.12	0.06	0.10
South Atlantic	0.37	0.33	0.12	0.21	0.09	0.17
East South Central	0.60	0.40	0.26	0.22	0.18	0.19
West South Central	0.35	0.35	0.10	0.15	0.04	0.11
West	0.38	0.28	0.12	0.08	0.07	0.07
Mountain	0.36	0.45	0.18	0.19	0.11	0.17
Pacific	0.52	0.35	0.15	0.08	0.08	0.06
COUNTY TYPE¹						
Large Metro	0.21	0.18	0.08	0.08	0.05	0.06
Small Metro	0.26	0.22	0.12	0.08	0.05	0.05
250K – 1 Mil. Pop.	0.32	0.24	0.11	0.08	0.06	0.06
< 250K Pop.	0.44	0.38	0.27	0.17	0.08	0.11
Nonmetro	0.24	0.28	0.11	0.17	0.05	0.15
Urbanized	0.36	0.38	0.14	0.17	0.10	0.05
Less Urbanized	0.37	0.46	0.18	0.35	0.06	0.33
Completely Rural	0.61	0.79	0.20	0.24	0.05	0.23

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.102P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.1706		0.3562		0.8271	
GEOGRAPHIC DIVISION						
Northeast	0.8722		0.7265		0.6744	
New England	0.5517		0.3157		0.6992	
Middle Atlantic	0.8959		0.7515		0.7526	
Midwest	0.6322		0.0708		0.0424	
East North Central	0.8433		0.2590		0.5373	
West North Central	0.6168		0.1514		0.0000	
South	0.7171		0.2628		0.1001	
South Atlantic	0.5082		0.8612		0.3158	
East South Central	0.2112		0.4883		0.6946	
West South Central	0.2275		0.0763		0.0655	
West	0.0629		0.0947		0.7426	
Mountain	0.5217		0.8261		0.8951	
Pacific	0.0228		0.0729		0.6094	
COUNTY TYPE¹						
Large Metro	0.3084		0.3575		0.5274	
Small Metro	0.0721		0.3868		0.4502	
250K – 1 Mil. Pop.	0.1609		0.2096		0.8970	
< 250K Pop.	0.2214		0.9574		0.3171	
Nonmetro	0.1998		0.5986		0.4144	
Urbanized	0.1402		0.5768		0.8197	
Less Urbanized	0.5659		0.6550		0.4217	
Completely Rural	0.7421		0.5692		0.4434	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.