

Table 3.1C Perceived Risk and Availability of Drugs, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

Risk/Availability ¹	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	750	829	108	104	138	138	732	795
Marijuana								
Smoke Once a Month	912	897	97	101	130	124	872	861
Smoke Once or Twice a Week	881	860	105	109	148	150	828	817
Cocaine								
Use Once a Month	757	759	108	112	140	133	726	721
Use Once or Twice a Week	484	539	86	88	86	98	477	507
Heroin								
Try Once or Twice	563	595	108	108	111	118	536	565
Use Once or Twice a Week	363	378	82	85	73	74	342	353
LSD								
Try Once or Twice	696	732	109	111	140	136	670	700
Use Once or Twice a Week	474	497	93	99	92	97	464	465
Alcohol								
Four or Five Drinks Nearly Every Day	775	770	118	112	139	144	764	753
Five or More Drinks Once or Twice a Week	863	831	106	103	145	134	834	818
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	898	890	109	106	131	119	873	861
Cocaine	816	768	86	90	136	144	792	736
Crack	821	750	90	91	133	142	793	733
Heroin	669	651	73	79	110	109	649	639
LSD	661	612	79	84	117	126	638	598
Approached by Someone Selling Drugs in the Past Month	359	376	82	83	110	116	330	328

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.1D Perceived Risk and Availability of Drugs, by Age Group: Standard Errors of Percentages, 2002 and 2003

Risk/Availability ¹	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	0.32	0.35	0.44	0.42	0.45	0.44	0.41	0.44
Marijuana								
Smoke Once a Month	0.39	0.38	0.40	0.41	0.42	0.39	0.50	0.48
Smoke Once or Twice a Week	0.38	0.37	0.43	0.44	0.48	0.48	0.47	0.46
Cocaine								
Use Once a Month	0.33	0.32	0.44	0.46	0.45	0.42	0.41	0.40
Use Once or Twice a Week	0.21	0.23	0.35	0.36	0.28	0.31	0.27	0.28
Heroin								
Try Once or Twice	0.24	0.25	0.45	0.44	0.36	0.38	0.30	0.32
Use Once or Twice a Week	0.16	0.16	0.34	0.35	0.24	0.24	0.19	0.20
LSD								
Try Once or Twice	0.30	0.31	0.46	0.46	0.46	0.44	0.38	0.39
Use Once or Twice a Week	0.21	0.21	0.39	0.41	0.30	0.31	0.26	0.26
Alcohol								
Four or Five Drinks Nearly Every Day	0.33	0.33	0.48	0.45	0.45	0.45	0.43	0.42
Five or More Drinks Once or Twice a Week	0.37	0.35	0.43	0.42	0.47	0.42	0.47	0.45
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	0.40	0.39	0.45	0.44	0.43	0.38	0.51	0.50
Cocaine	0.37	0.34	0.36	0.38	0.45	0.47	0.47	0.43
Crack	0.37	0.33	0.37	0.38	0.44	0.47	0.47	0.43
Heroin	0.30	0.29	0.31	0.33	0.37	0.36	0.39	0.38
LSD	0.30	0.27	0.34	0.36	0.39	0.41	0.38	0.35
Approached by Someone Selling Drugs in the Past Month	0.15	0.16	0.33	0.33	0.35	0.37	0.18	0.18

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.1P Perceived Risk and Availability of Drugs, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Risk/Availability ¹	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
PERCEPTIONS OF GREAT RISK								
Cigarettes								
Smoke One or More Packs Per Day	0.5454		0.0748		0.3932		0.8126	
Marijuana								
Smoke Once a Month	0.0145		0.0000		0.0259		0.0825	
Smoke Once or Twice a Week	0.0057		0.0000		0.0483		0.0445	
Cocaine								
Use Once a Month	0.3103		0.1375		0.4129		0.2606	
Use Once or Twice a Week	0.2835		0.0868		0.2007		0.2449	
Heroin								
Try Once or Twice	0.4826		0.6598		0.2807		0.5517	
Use Once or Twice a Week	0.7993		0.9426		0.8247		0.7943	
LSD								
Try Once or Twice	0.4889		0.2293		0.3597		0.2847	
Use Once or Twice a Week	0.8013		0.2351		0.1919		0.4431	
Alcohol								
Four or Five Drinks Nearly Every Day	0.2372		0.3437		0.0950		0.4663	
Five or More Drinks Once or Twice a Week	0.2480		0.6807		0.0487		0.3758	
PERCEIVED AVAILABILITY								
Fairly or Very Easy to Obtain								
Marijuana	0.1380		0.0231		0.3576		0.1095	
Cocaine	0.9737		0.9463		0.2586		0.7912	
Crack	0.3518		0.5767		0.7384		0.2890	
Heroin	0.7316		0.2090		0.7914		0.5922	
LSD	0.2167		0.0001		0.0001		0.9896	
Approached by Someone Selling Drugs in the Past Month	0.3770		0.1751		0.2241		0.4470	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.2C Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	108	104	108	104
(I) Demographic Characteristic				
AGE				
12-13	62	65	62	65
14-15	62	59	62	59
16-17	53	58	53	58
GENDER				
Male	76	72	76	72
Female	69	69	69	69
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	97	95	97	95
White	80	79	80	79
Black or African American	35	40	35	40
American Indian or Alaska Native	7	8	7	8
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	26	27	26	27
Two or More Races	11	11	11	11
Hispanic or Latino	49	46	49	46
COUNTY TYPE²				
Large Metro	85	78	85	78
Small Metro	53	51	53	51
Nonmetro	36	39	36	39
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	52	51	46	43
Marijuana	43	46	39	38
Any Illicit Drug Other Than Marijuana ³	38	35	34	32
Cigarettes	51	51	50	46
Alcohol	64	63	51	51
Binge Alcohol ⁴	47	52	46	40
PAST YEAR				
Any Illicit Drug ³	69	69	59	57
Marijuana	58	60	53	49
Any Illicit Drug Other Than Marijuana ³	56	56	48	48
Cigarettes	64	64	57	53
Alcohol	83	79	62	63
LIFETIME				
Any Illicit Drug ³	76	81	66	61
Marijuana	65	69	59	55
Any Illicit Drug Other Than Marijuana ³	66	71	57	56
Cigarettes	84	82	61	64
Alcohol	82	88	65	64

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.2D Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.44	0.42	0.44	0.42
(I) Demographic Characteristic				
AGE				
12-13	0.76	0.78	0.76	0.78
14-15	0.74	0.71	0.74	0.71
16-17	0.67	0.71	0.67	0.71
GENDER				
Male	0.61	0.57	0.61	0.57
Female	0.57	0.57	0.57	0.57
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.47	0.46	0.47	0.46
White	0.52	0.51	0.52	0.51
Black or African American	0.99	1.08	0.99	1.08
American Indian or Alaska Native	3.94	4.71	3.94	4.71
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.64	2.78	2.64	2.78
Two or More Races	2.94	2.77	2.94	2.77
Hispanic or Latino	1.28	1.15	1.28	1.15
COUNTY TYPE²				
Large Metro	0.65	0.59	0.65	0.59
Small Metro	0.73	0.70	0.73	0.70
Nonmetro	0.84	0.92	0.84	0.92
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.34	0.32	0.51	0.49
Marijuana	0.28	0.29	0.43	0.43
Any Illicit Drug Other Than Marijuana ³	0.25	0.22	0.38	0.36
Cigarettes	0.33	0.32	0.55	0.52
Alcohol	0.41	0.39	0.56	0.57
Binge Alcohol ⁴	0.30	0.32	0.51	0.46
PAST YEAR				
Any Illicit Drug ³	0.45	0.43	0.65	0.64
Marijuana	0.37	0.38	0.58	0.55
Any Illicit Drug Other Than Marijuana ³	0.36	0.35	0.53	0.54
Cigarettes	0.42	0.40	0.63	0.60
Alcohol	0.53	0.49	0.68	0.71
LIFETIME				
Any Illicit Drug ³	0.49	0.51	0.73	0.69
Marijuana	0.42	0.43	0.65	0.62
Any Illicit Drug Other Than Marijuana ³	0.42	0.45	0.63	0.63
Cigarettes	0.54	0.51	0.68	0.72
Alcohol	0.53	0.55	0.71	0.72

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.2P Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking One or More Packs of Cigarettes Per Day (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Smoke One or More Packs of Cigarettes Per Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.0748		0.0748	
(I) Demographic Characteristic				
AGE				
12-13	0.2439		0.2439	
14-15	0.0359		0.0359	
16-17	0.7074		0.7074	
GENDER				
Male	0.1635		0.1635	
Female	0.2444		0.2444	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0899		0.0899	
White	0.2606		0.2606	
Black or African American	0.0921		0.0921	
American Indian or Alaska Native	0.4905		0.4905	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.7607		0.7607	
Two or More Races	0.7052		0.7052	
Hispanic or Latino	0.5567		0.5567	
COUNTY TYPE²				
Large Metro	0.0783		0.0783	
Small Metro	0.9327		0.9327	
Nonmetro	0.2056		0.2056	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.5459		0.4167	
Marijuana	0.3596		0.7622	
Any Illicit Drug Other Than Marijuana ³	0.8599		0.5969	
Cigarettes	0.0926		0.4224	
Alcohol	0.9499		0.5218	
Binge Alcohol ⁴	0.9156		0.8720	
PAST YEAR				
Any Illicit Drug ³	0.7612		0.4794	
Marijuana	0.0848		0.4264	
Any Illicit Drug Other Than Marijuana ³	0.7903		0.6193	
Cigarettes	0.0068		0.3261	
Alcohol	0.4625		0.9611	
LIFETIME				
Any Illicit Drug ³	0.7313		0.6819	
Marijuana	0.0205		0.5949	
Any Illicit Drug Other Than Marijuana ³	0.7683		0.6473	
Cigarettes	0.0009		0.0696	
Alcohol	0.1562		0.7604	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.3C Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	97	101	97	101
(I) Demographic Characteristic				
AGE				
12-13	58	58	58	58
14-15	58	57	58	57
16-17	53	55	53	55
GENDER				
Male	64	70	64	70
Female	70	75	70	75
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	89	93	89	93
White	77	77	77	77
Black or African American	37	39	37	39
American Indian or Alaska Native	7	6	7	6
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	25	23	25	23
Two or More Races	11	12	11	12
Hispanic or Latino	41	43	41	43
COUNTY TYPE²				
Large Metro	74	77	74	77
Small Metro	49	50	49	50
Nonmetro	34	40	34	40
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	24	27	66	59
Marijuana	16	16	55	55
Any Illicit Drug Other Than Marijuana ³	21	22	46	43
Cigarettes	29	26	66	63
Alcohol	34	36	72	69
Binge Alcohol ⁴	24	26	61	60
PAST YEAR				
Any Illicit Drug ³	34	39	85	77
Marijuana	24	26	73	70
Any Illicit Drug Other Than Marijuana ³	29	34	68	67
Cigarettes	38	37	77	75
Alcohol	48	50	89	81
LIFETIME				
Any Illicit Drug ³	44	51	92	85
Marijuana	30	32	79	78
Any Illicit Drug Other Than Marijuana ³	39	49	79	78
Cigarettes	51	53	88	87
Alcohol	56	58	89	87

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.3D Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.40	0.41	0.40	0.41
(I) Demographic Characteristic				
AGE				
12-13	0.72	0.71	0.72	0.71
14-15	0.69	0.70	0.69	0.70
16-17	0.67	0.67	0.67	0.67
GENDER				
Male	0.51	0.56	0.51	0.56
Female	0.59	0.62	0.59	0.62
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.43	0.45	0.43	0.45
White	0.50	0.50	0.50	0.50
Black or African American	1.03	1.06	1.03	1.06
American Indian or Alaska Native	3.92	3.77	3.92	3.77
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.54	2.46	2.54	2.46
Two or More Races	2.79	3.18	2.79	3.18
Hispanic or Latino	1.07	1.08	1.07	1.08
COUNTY TYPE²				
Large Metro	0.57	0.59	0.57	0.59
Small Metro	0.69	0.69	0.69	0.69
Nonmetro	0.79	0.95	0.79	0.95
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.30	0.32	0.40	0.37
Marijuana	0.20	0.19	0.34	0.34
Any Illicit Drug Other Than Marijuana ³	0.26	0.26	0.28	0.27
Cigarettes	0.37	0.30	0.40	0.39
Alcohol	0.43	0.41	0.44	0.43
Binge Alcohol ⁴	0.31	0.31	0.37	0.37
PAST YEAR				
Any Illicit Drug ³	0.42	0.45	0.52	0.48
Marijuana	0.30	0.30	0.44	0.43
Any Illicit Drug Other Than Marijuana ³	0.37	0.40	0.41	0.42
Cigarettes	0.47	0.44	0.46	0.47
Alcohol	0.60	0.58	0.54	0.51
LIFETIME				
Any Illicit Drug ³	0.56	0.59	0.56	0.53
Marijuana	0.38	0.37	0.48	0.48
Any Illicit Drug Other Than Marijuana ³	0.49	0.57	0.48	0.49
Cigarettes	0.64	0.61	0.53	0.54
Alcohol	0.71	0.67	0.54	0.54

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.3P Risk Perceptions of Smoking Marijuana Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once a Month (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Smoke Marijuana Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.0000		0.0000	
(I) Demographic Characteristic				
AGE				
12-13	0.0605		0.0605	
14-15	0.0011		0.0011	
16-17	0.0044		0.0044	
GENDER				
Male	0.0010		0.0010	
Female	0.0015		0.0015	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0000		0.0000	
White	0.0000		0.0000	
Black or African American	0.7649		0.7649	
American Indian or Alaska Native	0.7661		0.7661	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.0601		0.0601	
Two or More Races	0.7919		0.7919	
Hispanic or Latino	0.4618		0.4618	
COUNTY TYPE²				
Large Metro	0.0000		0.0000	
Small Metro	0.0079		0.0079	
Nonmetro	0.3573		0.3573	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.5987		0.8376	
Marijuana	0.8230		0.8491	
Any Illicit Drug Other Than Marijuana ³	0.5063		0.6769	
Cigarettes	0.2151		0.3800	
Alcohol	0.8061		0.2150	
Binge Alcohol ⁴	0.4777		0.3369	
PAST YEAR				
Any Illicit Drug ³	0.5624		0.8402	
Marijuana	0.7824		0.3602	
Any Illicit Drug Other Than Marijuana ³	0.7384		0.8344	
Cigarettes	0.0961		0.1231	
Alcohol	0.4752		0.4166	
LIFETIME				
Any Illicit Drug ³	0.9454		0.8245	
Marijuana	0.1760		0.4249	
Any Illicit Drug Other Than Marijuana ³	0.6393		0.8184	
Cigarettes	0.0409		0.0088	
Alcohol	0.8241		0.9689	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.4C Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	105	109	105	109
(I) Demographic Characteristic				
AGE				
12-13	63	58	63	58
14-15	60	59	60	59
16-17	59	64	59	64
GENDER				
Male	72	74	72	74
Female	72	79	72	79
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	94	99	94	99
White	81	87	81	87
Black or African American	38	38	38	38
American Indian or Alaska Native	8	9	8	9
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	26	29	26	29
Two or More Races	11	12	11	12
Hispanic or Latino	48	49	48	49
COUNTY TYPE²				
Large Metro	80	86	80	86
Small Metro	56	55	56	55
Nonmetro	36	40	36	40
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	30	28	59	56
Marijuana	19	19	53	53
Any Illicit Drug Other Than Marijuana ³	26	24	42	40
Cigarettes	36	32	60	58
Alcohol	42	45	64	66
Binge Alcohol ⁴	30	32	58	55
PAST YEAR				
Any Illicit Drug ³	46	48	75	67
Marijuana	34	34	67	64
Any Illicit Drug Other Than Marijuana ³	38	42	61	62
Cigarettes	47	46	69	67
Alcohol	63	64	72	74
LIFETIME				
Any Illicit Drug ³	57	61	77	73
Marijuana	42	42	71	70
Any Illicit Drug Other Than Marijuana ³	50	56	69	69
Cigarettes	67	64	76	75
Alcohol	72	73	70	73

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.4D Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.43	0.44	0.43	0.44
(I) Demographic Characteristic				
AGE				
12-13	0.78	0.70	0.78	0.70
14-15	0.71	0.72	0.71	0.72
16-17	0.74	0.77	0.74	0.77
GENDER				
Male	0.58	0.58	0.58	0.58
Female	0.60	0.65	0.60	0.65
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.46	0.48	0.46	0.48
White	0.53	0.56	0.53	0.56
Black or African American	1.06	1.02	1.06	1.02
American Indian or Alaska Native	4.52	4.89	4.52	4.89
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.68	2.99	2.68	2.99
Two or More Races	2.97	2.97	2.97	2.97
Hispanic or Latino	1.24	1.24	1.24	1.24
COUNTY TYPE²				
Large Metro	0.62	0.65	0.62	0.65
Small Metro	0.79	0.76	0.79	0.76
Nonmetro	0.85	0.94	0.85	0.94
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.24	0.21	0.50	0.50
Marijuana	0.15	0.14	0.45	0.47
Any Illicit Drug Other Than Marijuana ³	0.21	0.18	0.35	0.36
Cigarettes	0.29	0.24	0.51	0.51
Alcohol	0.33	0.34	0.54	0.59
Binge Alcohol ⁴	0.24	0.24	0.49	0.49
PAST YEAR				
Any Illicit Drug ³	0.37	0.36	0.63	0.60
Marijuana	0.27	0.25	0.57	0.57
Any Illicit Drug Other Than Marijuana ³	0.30	0.31	0.52	0.55
Cigarettes	0.37	0.34	0.58	0.60
Alcohol	0.50	0.48	0.61	0.65
LIFETIME				
Any Illicit Drug ³	0.45	0.46	0.65	0.64
Marijuana	0.33	0.31	0.59	0.62
Any Illicit Drug Other Than Marijuana ³	0.40	0.42	0.58	0.61
Cigarettes	0.53	0.47	0.64	0.66
Alcohol	0.57	0.54	0.59	0.65

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.4P Risk Perceptions of Smoking Marijuana Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Smoking Marijuana Once or Twice a Week (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Smoke Marijuana Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.0000		0.0000	
(I) Demographic Characteristic				
AGE				
12-13	0.0273		0.0273	
14-15	0.0002		0.0002	
16-17	0.0127		0.0127	
GENDER				
Male	0.0000		0.0000	
Female	0.1766		0.1766	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0000		0.0000	
White	0.0000		0.0000	
Black or African American	0.9042		0.9042	
American Indian or Alaska Native	0.5758		0.5758	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.2988		0.2988	
Two or More Races	0.5184		0.5184	
Hispanic or Latino	0.6857		0.6857	
COUNTY TYPE²				
Large Metro	0.0000		0.0000	
Small Metro	0.0039		0.0039	
Nonmetro	0.8307		0.8307	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.3691		0.4757	
Marijuana	0.7916		0.6442	
Any Illicit Drug Other Than Marijuana ³	0.2831		0.2363	
Cigarettes	0.0061		0.4601	
Alcohol	0.4869		0.1745	
Binge Alcohol ⁴	0.9899		0.2154	
PAST YEAR				
Any Illicit Drug ³	0.9903		0.4101	
Marijuana	0.4734		0.9896	
Any Illicit Drug Other Than Marijuana ³	0.4967		0.6738	
Cigarettes	0.0111		0.8078	
Alcohol	0.7492		0.4081	
LIFETIME				
Any Illicit Drug ³	0.9415		0.2227	
Marijuana	0.0605		0.6706	
Any Illicit Drug Other Than Marijuana ³	0.2296		0.8525	
Cigarettes	0.0164		0.1432	
Alcohol	0.9127		0.6166	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.5C Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	108	112	108	112
(I) Demographic Characteristic				
AGE				
12-13	60	60	60	60
14-15	62	63	62	63
16-17	59	62	59	62
GENDER				
Male	75	74	75	74
Female	73	83	73	83
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	99	101	99	101
White	80	82	80	82
Black or African American	39	43	39	43
American Indian or Alaska Native	10	7	10	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	25	26	25	26
Two or More Races	12	13	12	13
Hispanic or Latino	49	48	49	48
COUNTY TYPE²				
Large Metro	82	87	82	87
Small Metro	56	55	56	55
Nonmetro	41	42	41	42
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	44	49	53	50
Marijuana	39	43	43	41
Any Illicit Drug Other Than Marijuana ³	30	32	42	38
Cigarettes	52	52	52	48
Alcohol	57	59	57	53
Binge Alcohol ⁴	46	48	47	44
PAST YEAR				
Any Illicit Drug ³	59	65	69	61
Marijuana	55	55	56	52
Any Illicit Drug Other Than Marijuana ³	46	51	58	54
Cigarettes	63	65	62	58
Alcohol	74	74	74	70
LIFETIME				
Any Illicit Drug ³	66	76	77	69
Marijuana	61	64	61	58
Any Illicit Drug Other Than Marijuana ³	58	65	68	66
Cigarettes	74	80	73	70
Alcohol	78	79	80	74

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.5D Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.44	0.46	0.44	0.46
(I) Demographic Characteristic				
AGE				
12-13	0.75	0.73	0.75	0.73
14-15	0.75	0.78	0.75	0.78
16-17	0.74	0.76	0.74	0.76
GENDER				
Male	0.61	0.59	0.61	0.59
Female	0.61	0.69	0.61	0.69
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.48	0.49	0.48	0.49
White	0.52	0.54	0.52	0.54
Black or African American	1.09	1.18	1.09	1.18
American Indian or Alaska Native	5.26	4.16	5.26	4.16
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.58	2.68	2.58	2.68
Two or More Races	3.16	3.26	3.16	3.26
Hispanic or Latino	1.26	1.20	1.26	1.20
COUNTY TYPE²				
Large Metro	0.63	0.66	0.63	0.66
Small Metro	0.78	0.76	0.78	0.76
Nonmetro	0.98	1.00	0.98	1.00
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.36	0.39	0.44	0.42
Marijuana	0.32	0.34	0.35	0.35
Any Illicit Drug Other Than Marijuana ³	0.25	0.25	0.34	0.32
Cigarettes	0.43	0.42	0.43	0.40
Alcohol	0.46	0.47	0.47	0.45
Binge Alcohol ⁴	0.37	0.38	0.39	0.37
PAST YEAR				
Any Illicit Drug ³	0.48	0.51	0.57	0.51
Marijuana	0.45	0.44	0.46	0.43
Any Illicit Drug Other Than Marijuana ³	0.38	0.40	0.48	0.46
Cigarettes	0.52	0.52	0.51	0.48
Alcohol	0.60	0.59	0.62	0.59
LIFETIME				
Any Illicit Drug ³	0.54	0.60	0.64	0.58
Marijuana	0.49	0.51	0.51	0.49
Any Illicit Drug Other Than Marijuana ³	0.47	0.51	0.56	0.55
Cigarettes	0.60	0.63	0.61	0.59
Alcohol	0.63	0.62	0.66	0.62

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.5P Risk Perceptions of Using Cocaine Once a Month, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once a Month (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Use Cocaine Once a Month ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.1375		0.1375	
(I) Demographic Characteristic				
AGE				
12-13	0.3629		0.3629	
14-15	0.0487		0.0487	
16-17	0.7022		0.7022	
GENDER				
Male	0.0350		0.0350	
Female	0.9923		0.9923	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1099		0.1099	
White	0.0798		0.0798	
Black or African American	0.7447		0.7447	
American Indian or Alaska Native	0.5692		0.5692	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.8513		0.8513	
Two or More Races	0.5984		0.5984	
Hispanic or Latino	0.8674		0.8674	
COUNTY TYPE²				
Large Metro	0.0506		0.0506	
Small Metro	0.9160		0.9160	
Nonmetro	0.8181		0.8181	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.8080		0.1046	
Marijuana	0.8785		0.1227	
Any Illicit Drug Other Than Marijuana ³	0.6106		0.9619	
Cigarettes	0.0336		0.5231	
Alcohol	0.8189		0.6951	
Binge Alcohol ⁴	0.9549		0.9829	
PAST YEAR				
Any Illicit Drug ³	0.6193		0.5087	
Marijuana	0.1890		0.1534	
Any Illicit Drug Other Than Marijuana ³	0.7830		0.9602	
Cigarettes	0.0083		0.2263	
Alcohol	0.1523		0.5995	
LIFETIME				
Any Illicit Drug ³	0.2683		0.6995	
Marijuana	0.0155		0.4775	
Any Illicit Drug Other Than Marijuana ³	0.5889		0.6100	
Cigarettes	0.0002		0.0965	
Alcohol	0.0953		0.7034	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.6C Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	86	88	86	88
(I) Demographic Characteristic				
AGE				
12-13	60	54	60	54
14-15	49	49	49	49
16-17	43	44	43	44
GENDER				
Male	60	57	60	57
Female	60	63	60	63
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	76	73	76	73
White	58	57	58	57
Black or African American	34	36	34	36
American Indian or Alaska Native	9	5	9	5
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	26	27	26	27
Two or More Races	11	8	11	8
Hispanic or Latino	39	47	39	47
COUNTY TYPE²				
Large Metro	65	68	65	68
Small Metro	44	49	44	49
Nonmetro	31	32	31	32
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	58	58	37	32
Marijuana	50	52	29	26
Any Illicit Drug Other Than Marijuana ³	40	38	30	26
Cigarettes	64	62	34	33
Alcohol	69	72	37	36
Binge Alcohol ⁴	58	59	32	28
PAST YEAR				
Any Illicit Drug ³	76	82	46	37
Marijuana	68	69	36	32
Any Illicit Drug Other Than Marijuana ³	60	67	40	35
Cigarettes	78	80	40	39
Alcohol	89	90	49	46
LIFETIME				
Any Illicit Drug ³	84	96	49	43
Marijuana	76	81	39	35
Any Illicit Drug Other Than Marijuana ³	71	84	47	42
Cigarettes	95	97	46	44
Alcohol	93	98	49	46

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.6D Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.35	0.36	0.35	0.36
(I) Demographic Characteristic				
AGE				
12-13	0.75	0.66	0.75	0.66
14-15	0.58	0.60	0.58	0.60
16-17	0.55	0.54	0.55	0.54
GENDER				
Male	0.48	0.45	0.48	0.45
Female	0.50	0.52	0.50	0.52
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.37	0.35	0.37	0.35
White	0.38	0.37	0.38	0.37
Black or African American	0.95	1.00	0.95	1.00
American Indian or Alaska Native	4.83	3.19	4.83	3.19
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.61	2.83	2.61	2.83
Two or More Races	2.78	2.11	2.78	2.11
Hispanic or Latino	1.01	1.19	1.01	1.19
COUNTY TYPE²				
Large Metro	0.50	0.52	0.50	0.52
Small Metro	0.62	0.67	0.62	0.67
Nonmetro	0.72	0.76	0.72	0.76
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.30	0.29	0.74	0.67
Marijuana	0.26	0.26	0.58	0.55
Any Illicit Drug Other Than Marijuana ³	0.21	0.19	0.61	0.55
Cigarettes	0.33	0.31	0.69	0.70
Alcohol	0.35	0.36	0.75	0.76
Binge Alcohol ⁴	0.30	0.30	0.65	0.60
PAST YEAR				
Any Illicit Drug ³	0.39	0.41	0.93	0.77
Marijuana	0.35	0.35	0.73	0.66
Any Illicit Drug Other Than Marijuana ³	0.31	0.34	0.82	0.74
Cigarettes	0.40	0.40	0.82	0.82
Alcohol	0.46	0.45	1.00	0.96
LIFETIME				
Any Illicit Drug ³	0.43	0.49	1.00	0.90
Marijuana	0.39	0.41	0.79	0.74
Any Illicit Drug Other Than Marijuana ³	0.37	0.42	0.95	0.88
Cigarettes	0.49	0.49	0.93	0.92
Alcohol	0.48	0.49	0.99	0.96

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.6P Risk Perceptions of Using Cocaine Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Cocaine Once or Twice a Week (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Use Cocaine Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.0868		0.0868	
(I) Demographic Characteristic				
AGE				
12-13	0.2311		0.2311	
14-15	0.2063		0.2063	
16-17	0.7805		0.7805	
GENDER				
Male	0.2368		0.2368	
Female	0.1851		0.1851	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.0404		0.0404	
White	0.4139		0.4139	
Black or African American	0.0420		0.0420	
American Indian or Alaska Native	0.4080		0.4080	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.3562		0.3562	
Two or More Races	0.6921		0.6921	
Hispanic or Latino	0.9617		0.9617	
COUNTY TYPE²				
Large Metro	0.0504		0.0504	
Small Metro	0.5914		0.5914	
Nonmetro	0.8657		0.8657	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.4216		0.6114	
Marijuana	0.4095		0.6667	
Any Illicit Drug Other Than Marijuana ³	0.5995		0.7235	
Cigarettes	0.0109		0.4563	
Alcohol	0.5389		0.1237	
Binge Alcohol ⁴	0.7297		0.5115	
PAST YEAR				
Any Illicit Drug ³	0.1752		0.3399	
Marijuana	0.0251		0.8056	
Any Illicit Drug Other Than Marijuana ³	0.5530		0.4528	
Cigarettes	0.0004		0.3397	
Alcohol	0.0773		0.0421	
LIFETIME				
Any Illicit Drug ³	0.3221		0.3664	
Marijuana	0.0044		0.3875	
Any Illicit Drug Other Than Marijuana ³	0.7994		0.6294	
Cigarettes	0.0000		0.4898	
Alcohol	0.0350		0.0326	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.7C Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	108	108	108	108
(I) Demographic Characteristic				
AGE				
12-13	62	61	62	61
14-15	65	60	65	60
16-17	55	59	55	59
GENDER				
Male	77	73	77	73
Female	74	79	74	79
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	98	101	98	101
White	82	80	82	80
Black or African American	38	43	38	43
American Indian or Alaska Native	7	9	7	9
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	27	27	27	27
Two or More Races	13	12	13	12
Hispanic or Latino	49	44	49	44
COUNTY TYPE²				
Large Metro	82	81	82	81
Small Metro	50	58	50	58
Nonmetro	41	41	41	41
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	55	55	38	37
Marijuana	48	51	29	30
Any Illicit Drug Other Than Marijuana ³	40	37	30	31
Cigarettes	59	59	41	37
Alcohol	66	65	45	46
Binge Alcohol ⁴	56	54	34	36
PAST YEAR				
Any Illicit Drug ³	72	72	53	54
Marijuana	66	65	40	42
Any Illicit Drug Other Than Marijuana ³	57	56	45	48
Cigarettes	69	69	49	49
Alcohol	80	80	63	60
LIFETIME				
Any Illicit Drug ³	76	82	65	63
Marijuana	71	74	48	48
Any Illicit Drug Other Than Marijuana ³	68	71	56	56
Cigarettes	83	83	59	61
Alcohol	78	82	67	65

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.7D Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.45	0.44	0.45	0.44
(I) Demographic Characteristic				
AGE				
12-13	0.76	0.74	0.76	0.74
14-15	0.78	0.74	0.78	0.74
16-17	0.69	0.72	0.69	0.72
GENDER				
Male	0.62	0.59	0.62	0.59
Female	0.63	0.66	0.63	0.66
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.48	0.49	0.48	0.49
White	0.54	0.52	0.54	0.52
Black or African American	1.09	1.19	1.09	1.19
American Indian or Alaska Native	3.93	5.28	3.93	5.28
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.80	2.85	2.80	2.85
Two or More Races	3.39	3.05	3.39	3.05
Hispanic or Latino	1.27	1.12	1.27	1.12
COUNTY TYPE²				
Large Metro	0.64	0.62	0.64	0.62
Small Metro	0.70	0.80	0.70	0.80
Nonmetro	0.96	0.99	0.96	0.99
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.39	0.38	0.38	0.37
Marijuana	0.34	0.35	0.29	0.30
Any Illicit Drug Other Than Marijuana ³	0.28	0.26	0.30	0.31
Cigarettes	0.42	0.41	0.40	0.36
Alcohol	0.46	0.45	0.45	0.46
Binge Alcohol ⁴	0.39	0.37	0.33	0.35
PAST YEAR				
Any Illicit Drug ³	0.50	0.50	0.52	0.53
Marijuana	0.47	0.45	0.40	0.41
Any Illicit Drug Other Than Marijuana ³	0.40	0.39	0.45	0.48
Cigarettes	0.49	0.48	0.49	0.48
Alcohol	0.56	0.55	0.62	0.60
LIFETIME				
Any Illicit Drug ³	0.53	0.57	0.65	0.62
Marijuana	0.50	0.51	0.48	0.47
Any Illicit Drug Other Than Marijuana ³	0.48	0.50	0.56	0.56
Cigarettes	0.59	0.58	0.59	0.61
Alcohol	0.55	0.57	0.67	0.64

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.7P Risk Perceptions of Trying Heroin Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying Heroin Once or Twice (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Try Heroin Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.6598		0.6598	
(I) Demographic Characteristic				
AGE				
12-13	0.5973		0.5973	
14-15	0.2361		0.2361	
16-17	0.2048		0.2048	
GENDER				
Male	0.6184		0.6184	
Female	0.8926		0.8926	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.5758		0.5758	
White	0.6946		0.6946	
Black or African American	0.4735		0.4735	
American Indian or Alaska Native	1.0000		1.0000	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.5352		0.5352	
Two or More Races	0.3795		0.3795	
Hispanic or Latino	0.8704		0.8704	
COUNTY TYPE²				
Large Metro	0.3951		0.3951	
Small Metro	0.6740		0.6740	
Nonmetro	0.9492		0.9492	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.2159		0.9088	
Marijuana	0.3902		0.6731	
Any Illicit Drug Other Than Marijuana ³	0.3152		0.6054	
Cigarettes	0.0376		0.6070	
Alcohol	0.3846		0.1327	
Binge Alcohol ⁴	0.2416		0.0513	
PAST YEAR				
Any Illicit Drug ³	0.1183		0.5188	
Marijuana	0.0311		0.9102	
Any Illicit Drug Other Than Marijuana ³	0.2815		0.4491	
Cigarettes	0.0040		0.3211	
Alcohol	0.1082		0.2827	
LIFETIME				
Any Illicit Drug ³	0.3006		0.6220	
Marijuana	0.0156		0.8391	
Any Illicit Drug Other Than Marijuana ³	0.3011		0.2925	
Cigarettes	0.0002		0.1249	
Alcohol	0.0319		0.2750	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.8C Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	82	85	82	85
(I) Demographic Characteristic				
AGE				
12-13	57	55	57	55
14-15	49	47	49	47
16-17	35	35	35	35
GENDER				
Male	58	57	58	57
Female	57	62	57	62
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	73	72	73	72
White	54	53	54	53
Black or African American	33	36	33	36
American Indian or Alaska Native	7	5	7	5
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	25	25	25	25
Two or More Races	9	9	9	9
Hispanic or Latino	39	42	39	42
COUNTY TYPE²				
Large Metro	67	69	67	69
Small Metro	40	40	40	40
Nonmetro	27	30	27	30
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	62	62	25	25
Marijuana	53	56	18	20
Any Illicit Drug Other Than Marijuana ³	46	43	20	22
Cigarettes	66	67	24	26
Alcohol	75	77	26	29
Binge Alcohol ⁴	62	62	21	22
PAST YEAR				
Any Illicit Drug ³	83	83	33	33
Marijuana	74	74	24	25
Any Illicit Drug Other Than Marijuana ³	66	68	29	30
Cigarettes	78	84	30	32
Alcohol	91	94	38	39
LIFETIME				
Any Illicit Drug ³	92	96	40	39
Marijuana	81	85	29	29
Any Illicit Drug Other Than Marijuana ³	79	84	36	36
Cigarettes	98	100	37	39
Alcohol	93	100	42	43

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.8D Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.34	0.35	0.34	0.35
(I) Demographic Characteristic				
AGE				
12-13	0.70	0.67	0.70	0.67
14-15	0.59	0.58	0.59	0.58
16-17	0.45	0.43	0.45	0.43
GENDER				
Male	0.47	0.45	0.47	0.45
Female	0.48	0.52	0.48	0.52
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.36	0.35	0.36	0.35
White	0.35	0.35	0.35	0.35
Black or African American	0.93	0.99	0.93	0.99
American Indian or Alaska Native	3.93	2.88	3.93	2.88
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.58	2.68	2.58	2.68
Two or More Races	2.29	2.19	2.29	2.19
Hispanic or Latino	1.02	1.05	1.02	1.05
COUNTY TYPE²				
Large Metro	0.52	0.53	0.52	0.53
Small Metro	0.56	0.56	0.56	0.56
Nonmetro	0.65	0.72	0.65	0.72
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.31	0.31	0.58	0.59
Marijuana	0.26	0.28	0.42	0.47
Any Illicit Drug Other Than Marijuana ³	0.23	0.21	0.47	0.52
Cigarettes	0.33	0.33	0.56	0.60
Alcohol	0.37	0.38	0.61	0.67
Binge Alcohol ⁴	0.31	0.31	0.49	0.52
PAST YEAR				
Any Illicit Drug ³	0.41	0.41	0.77	0.78
Marijuana	0.37	0.36	0.56	0.59
Any Illicit Drug Other Than Marijuana ³	0.33	0.34	0.67	0.71
Cigarettes	0.39	0.42	0.70	0.75
Alcohol	0.46	0.46	0.90	0.90
LIFETIME				
Any Illicit Drug ³	0.46	0.47	0.93	0.90
Marijuana	0.40	0.42	0.69	0.67
Any Illicit Drug Other Than Marijuana ³	0.40	0.42	0.85	0.84
Cigarettes	0.49	0.49	0.87	0.91
Alcohol	0.46	0.49	0.98	1.02

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.8P Risk Perceptions of Using Heroin Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using Heroin Once or Twice a Week (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Use Heroin Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.9426		0.9426	
(I) Demographic Characteristic				
AGE				
12-13	0.7750		0.7750	
14-15	0.8301		0.8301	
16-17	0.9248		0.9248	
GENDER				
Male	0.8068		0.8068	
Female	0.8985		0.8985	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.7275		0.7275	
White	0.7377		0.7377	
Black or African American	0.1663		0.1663	
American Indian or Alaska Native	0.9780		0.9780	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.8562		0.8562	
Two or More Races	0.5009		0.5009	
Hispanic or Latino	0.6810		0.6810	
COUNTY TYPE²				
Large Metro	0.3606		0.3606	
Small Metro	0.3668		0.3668	
Nonmetro	0.4755		0.4755	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.1060		0.2404	
Marijuana	0.1281		0.1261	
Any Illicit Drug Other Than Marijuana ³	0.3897		0.4638	
Cigarettes	0.0090		0.1809	
Alcohol	0.5584		0.0410	
Binge Alcohol ⁴	0.5154		0.0691	
PAST YEAR				
Any Illicit Drug ³	0.1995		0.3858	
Marijuana	0.0214		0.2354	
Any Illicit Drug Other Than Marijuana ³	0.4054		0.3135	
Cigarettes	0.0004		0.1104	
Alcohol	0.1259		0.0208	
LIFETIME				
Any Illicit Drug ³	0.4046		0.4602	
Marijuana	0.0093		0.2546	
Any Illicit Drug Other Than Marijuana ³	0.6386		0.4364	
Cigarettes	0.0000		0.4263	
Alcohol	0.0485		0.0200	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.9C Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	109	111	109	111
(I) Demographic Characteristic				
AGE				
12-13	60	54	60	54
14-15	67	60	67	60
16-17	56	68	56	68
GENDER				
Male	78	75	78	75
Female	73	76	73	76
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	98	102	98	102
White	79	81	79	81
Black or African American	39	45	39	45
American Indian or Alaska Native	9	8	9	8
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	26	27	26	27
Two or More Races	14	12	14	12
Hispanic or Latino	48	47	48	47
COUNTY TYPE²				
Large Metro	85	84	85	84
Small Metro	53	55	53	55
Nonmetro	41	42	41	42
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	44	48	51	46
Marijuana	37	43	42	40
Any Illicit Drug Other Than Marijuana ³	32	31	39	36
Cigarettes	52	52	50	48
Alcohol	57	60	57	51
Binge Alcohol ⁴	46	49	48	41
PAST YEAR				
Any Illicit Drug ³	62	66	65	59
Marijuana	56	56	55	51
Any Illicit Drug Other Than Marijuana ³	48	51	57	52
Cigarettes	62	62	59	60
Alcohol	70	73	69	68
LIFETIME				
Any Illicit Drug ³	69	76	70	65
Marijuana	63	66	60	57
Any Illicit Drug Other Than Marijuana ³	58	64	65	60
Cigarettes	77	79	68	70
Alcohol	73	77	72	73

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.9D Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.46	0.46	0.46	0.46
(I) Demographic Characteristic				
AGE				
12-13	0.77	0.69	0.77	0.69
14-15	0.83	0.76	0.83	0.76
16-17	0.71	0.83	0.71	0.83
GENDER				
Male	0.64	0.62	0.64	0.62
Female	0.63	0.64	0.63	0.64
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.49	0.51	0.49	0.51
White	0.53	0.54	0.53	0.54
Black or African American	1.13	1.25	1.13	1.25
American Indian or Alaska Native	4.92	5.00	4.92	5.00
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.79	2.93	2.79	2.93
Two or More Races	3.71	3.22	3.71	3.22
Hispanic or Latino	1.26	1.21	1.26	1.21
COUNTY TYPE²				
Large Metro	0.67	0.66	0.67	0.66
Small Metro	0.76	0.77	0.76	0.77
Nonmetro	0.98	1.03	0.98	1.03
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.35	0.38	0.45	0.41
Marijuana	0.30	0.33	0.37	0.35
Any Illicit Drug Other Than Marijuana ³	0.26	0.25	0.34	0.32
Cigarettes	0.41	0.40	0.44	0.43
Alcohol	0.45	0.47	0.50	0.46
Binge Alcohol ⁴	0.37	0.38	0.42	0.37
PAST YEAR				
Any Illicit Drug ³	0.49	0.52	0.58	0.53
Marijuana	0.45	0.44	0.49	0.46
Any Illicit Drug Other Than Marijuana ³	0.38	0.40	0.51	0.47
Cigarettes	0.50	0.48	0.52	0.54
Alcohol	0.56	0.57	0.61	0.61
LIFETIME				
Any Illicit Drug ³	0.55	0.59	0.62	0.58
Marijuana	0.50	0.52	0.53	0.51
Any Illicit Drug Other Than Marijuana ³	0.47	0.50	0.58	0.53
Cigarettes	0.62	0.62	0.60	0.62
Alcohol	0.58	0.60	0.64	0.65

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.9P Risk Perceptions of Trying LSD Once or Twice, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Trying LSD Once or Twice (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Try LSD Once or Twice ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.2293		0.2293	
(I) Demographic Characteristic				
AGE				
12-13	0.6088		0.6088	
14-15	0.1119		0.1119	
16-17	0.4217		0.4217	
GENDER				
Male	0.0208		0.0208	
Female	0.5912		0.5912	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1778		0.1778	
White	0.1118		0.1118	
Black or African American	0.3444		0.3444	
American Indian or Alaska Native	0.3203		0.3203	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.3866		0.3866	
Two or More Races	0.3301		0.3301	
Hispanic or Latino	0.9536		0.9536	
COUNTY TYPE²				
Large Metro	0.1818		0.1818	
Small Metro	0.5939		0.5939	
Nonmetro	0.8856		0.8856	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.3509		0.0278	
Marijuana	0.1519		0.0145	
Any Illicit Drug Other Than Marijuana ³	0.9974		0.6741	
Cigarettes	0.3938		0.0625	
Alcohol	0.5874		0.6598	
Binge Alcohol ⁴	0.5194		0.4460	
PAST YEAR				
Any Illicit Drug ³	0.1507		0.0195	
Marijuana	0.3653		0.0007	
Any Illicit Drug Other Than Marijuana ³	0.4557		0.3430	
Cigarettes	0.1529		0.0137	
Alcohol	0.6303		0.1636	
LIFETIME				
Any Illicit Drug ³	0.2504		0.1074	
Marijuana	0.8894		0.0065	
Any Illicit Drug Other Than Marijuana ³	0.3704		0.4333	
Cigarettes	0.0823		0.0004	
Alcohol	0.7883		0.0984	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.10C Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	93	99	93	99
(I) Demographic Characteristic				
AGE				
12-13	56	55	56	55
14-15	55	53	55	53
16-17	47	48	47	48
GENDER				
Male	63	66	63	66
Female	66	67	66	67
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	79	84	79	84
White	61	64	61	64
Black or African American	34	42	34	42
American Indian or Alaska Native	9	5	9	5
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	25	28	25	28
Two or More Races	10	8	10	8
Hispanic or Latino	45	44	45	44
COUNTY TYPE²				
Large Metro	74	76	74	76
Small Metro	48	53	48	53
Nonmetro	31	33	31	33
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	56	58	37	34
Marijuana	47	51	30	27
Any Illicit Drug Other Than Marijuana ³	41	39	29	27
Cigarettes	61	64	36	34
Alcohol	69	73	39	37
Binge Alcohol ⁴	54	59	35	29
PAST YEAR				
Any Illicit Drug ³	74	80	47	43
Marijuana	65	67	39	34
Any Illicit Drug Other Than Marijuana ³	59	65	41	38
Cigarettes	73	79	43	40
Alcohol	85	87	49	47
LIFETIME				
Any Illicit Drug ³	82	93	52	48
Marijuana	72	78	43	39
Any Illicit Drug Other Than Marijuana ³	73	83	47	45
Cigarettes	93	96	48	48
Alcohol	89	94	51	49

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.10D Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.39	0.41	0.39	0.41
(I) Demographic Characteristic				
AGE				
12-13	0.71	0.69	0.71	0.69
14-15	0.68	0.66	0.68	0.66
16-17	0.60	0.58	0.60	0.58
GENDER				
Male	0.52	0.54	0.52	0.54
Female	0.57	0.57	0.57	0.57
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.39	0.42	0.39	0.42
White	0.41	0.43	0.41	0.43
Black or African American	0.98	1.19	0.98	1.19
American Indian or Alaska Native	4.80	2.98	4.80	2.98
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.70	3.01	2.70	3.01
Two or More Races	2.75	2.16	2.75	2.16
Hispanic or Latino	1.19	1.13	1.19	1.13
COUNTY TYPE²				
Large Metro	0.58	0.59	0.58	0.59
Small Metro	0.69	0.74	0.69	0.74
Nonmetro	0.75	0.81	0.75	0.81
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.31	0.31	0.65	0.60
Marijuana	0.26	0.28	0.53	0.49
Any Illicit Drug Other Than Marijuana ³	0.22	0.21	0.51	0.49
Cigarettes	0.34	0.34	0.64	0.60
Alcohol	0.38	0.40	0.69	0.67
Binge Alcohol ⁴	0.30	0.32	0.62	0.52
PAST YEAR				
Any Illicit Drug ³	0.41	0.43	0.83	0.78
Marijuana	0.36	0.36	0.70	0.62
Any Illicit Drug Other Than Marijuana ³	0.33	0.35	0.73	0.69
Cigarettes	0.40	0.43	0.76	0.73
Alcohol	0.47	0.47	0.87	0.84
LIFETIME				
Any Illicit Drug ³	0.45	0.50	0.92	0.86
Marijuana	0.40	0.42	0.76	0.70
Any Illicit Drug Other Than Marijuana ³	0.40	0.45	0.83	0.80
Cigarettes	0.51	0.52	0.85	0.87
Alcohol	0.49	0.51	0.90	0.88

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.10P Risk Perceptions of Using LSD Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Using LSD Once or Twice a Week (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Use LSD Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.2351		0.2351	
(I) Demographic Characteristic				
AGE				
12-13	0.6317		0.6317	
14-15	0.3343		0.3343	
16-17	0.5086		0.5086	
GENDER				
Male	0.0047		0.0047	
Female	0.3188		0.3188	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.1028		0.1028	
White	0.1582		0.1582	
Black or African American	0.6751		0.6751	
American Indian or Alaska Native	0.5537		0.5537	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.5341		0.5341	
Two or More Races	0.2437		0.2437	
Hispanic or Latino	0.6400		0.6400	
COUNTY TYPE²				
Large Metro	0.1882		0.1882	
Small Metro	0.6052		0.6052	
Nonmetro	0.6921		0.6921	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.4728		0.4770	
Marijuana	0.7234		0.2736	
Any Illicit Drug Other Than Marijuana ³	0.6826		0.8870	
Cigarettes	0.0659		0.6017	
Alcohol	0.8516		0.5510	
Binge Alcohol ⁴	0.8825		0.6741	
PAST YEAR				
Any Illicit Drug ³	0.5582		0.7092	
Marijuana	0.1816		0.1813	
Any Illicit Drug Other Than Marijuana ³	0.8492		0.8160	
Cigarettes	0.0073		0.4976	
Alcohol	0.3206		0.6403	
LIFETIME				
Any Illicit Drug ³	0.5545		0.6691	
Marijuana	0.0843		0.2536	
Any Illicit Drug Other Than Marijuana ³	0.8771		0.8027	
Cigarettes	0.0012		0.1984	
Alcohol	0.1572		0.5343	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.11C Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	118	112	118	112
(I) Demographic Characteristic				
AGE				
12-13	60	63	60	63
14-15	58	59	58	59
16-17	65	61	65	61
GENDER				
Male	81	76	81	76
Female	72	75	72	75
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	102	99	102	99
White	83	85	83	85
Black or African American	39	40	39	40
American Indian or Alaska Native	9	8	9	8
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	25	24	25	24
Two or More Races	11	12	11	12
Hispanic or Latino	53	52	53	52
COUNTY TYPE²				
Large Metro	91	87	91	87
Small Metro	62	53	62	53
Nonmetro	37	43	37	43
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	45	45	47	50
Marijuana	36	39	41	43
Any Illicit Drug Other Than Marijuana ³	34	34	36	35
Cigarettes	48	45	52	51
Alcohol	55	59	57	54
Binge Alcohol ⁴	39	43	50	48
PAST YEAR				
Any Illicit Drug ³	64	65	59	64
Marijuana	54	52	53	56
Any Illicit Drug Other Than Marijuana ³	53	54	50	55
Cigarettes	58	60	60	59
Alcohol	75	80	63	67
LIFETIME				
Any Illicit Drug ³	70	77	65	69
Marijuana	60	64	59	62
Any Illicit Drug Other Than Marijuana ³	61	67	57	63
Cigarettes	74	71	64	70
Alcohol	80	88	67	67

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.11D Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.48	0.45	0.48	0.45
(I) Demographic Characteristic				
AGE				
12-13	0.73	0.76	0.73	0.76
14-15	0.68	0.72	0.68	0.72
16-17	0.82	0.74	0.82	0.74
GENDER				
Male	0.65	0.60	0.65	0.60
Female	0.60	0.61	0.60	0.61
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.49	0.48	0.49	0.48
White	0.54	0.55	0.54	0.55
Black or African American	1.10	1.07	1.10	1.07
American Indian or Alaska Native	4.93	4.34	4.93	4.34
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.49	2.48	2.49	2.48
Two or More Races	2.88	2.96	2.88	2.96
Hispanic or Latino	1.36	1.28	1.36	1.28
COUNTY TYPE²				
Large Metro	0.70	0.66	0.70	0.66
Small Metro	0.86	0.72	0.86	0.72
Nonmetro	0.87	1.02	0.87	1.02
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.29	0.30	0.51	0.52
Marijuana	0.24	0.25	0.44	0.45
Any Illicit Drug Other Than Marijuana ³	0.22	0.22	0.39	0.37
Cigarettes	0.31	0.29	0.56	0.53
Alcohol	0.36	0.39	0.61	0.57
Binge Alcohol ⁴	0.26	0.28	0.54	0.50
PAST YEAR				
Any Illicit Drug ³	0.42	0.42	0.64	0.67
Marijuana	0.35	0.34	0.57	0.58
Any Illicit Drug Other Than Marijuana ³	0.35	0.35	0.54	0.57
Cigarettes	0.38	0.39	0.64	0.62
Alcohol	0.49	0.53	0.68	0.70
LIFETIME				
Any Illicit Drug ³	0.46	0.50	0.70	0.73
Marijuana	0.40	0.42	0.64	0.65
Any Illicit Drug Other Than Marijuana ³	0.40	0.44	0.62	0.66
Cigarettes	0.49	0.47	0.69	0.74
Alcohol	0.52	0.58	0.72	0.71

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.11P Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having Four or Five Drinks of an Alcoholic Beverage Nearly Every Day (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Have Four or Five Drinks of Alcoholic Beverage Nearly Every Day ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.3437		0.3437	
(I) Demographic Characteristic				
AGE				
12-13	0.7577		0.7577	
14-15	0.0424		0.0424	
16-17	0.6261		0.6261	
GENDER				
Male	0.7052		0.7052	
Female	0.0646		0.0646	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.9455		0.9455	
White	0.7352		0.7352	
Black or African American	0.2267		0.2267	
American Indian or Alaska Native	0.9161		0.9161	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.2653		0.2653	
Two or More Races	0.3479		0.3479	
Hispanic or Latino	0.0653		0.0653	
COUNTY TYPE²				
Large Metro	0.9657		0.9657	
Small Metro	0.0483		0.0483	
Nonmetro	0.8757		0.8757	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.5698		0.2680	
Marijuana	0.4606		0.4371	
Any Illicit Drug Other Than Marijuana ³	0.8779		0.6249	
Cigarettes	0.0077		0.6055	
Alcohol	0.4945		0.5365	
Binge Alcohol ⁴	0.3693		0.2983	
PAST YEAR				
Any Illicit Drug ³	0.6029		0.3977	
Marijuana	0.0710		0.2433	
Any Illicit Drug Other Than Marijuana ³	0.7871		0.4350	
Cigarettes	0.0080		0.0781	
Alcohol	0.8713		0.4060	
LIFETIME				
Any Illicit Drug ³	0.8607		0.3008	
Marijuana	0.0640		0.1346	
Any Illicit Drug Other Than Marijuana ³	0.8148		0.7439	
Cigarettes	0.0009		0.0097	
Alcohol	0.2711		0.7280	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.12C Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	106	103	106	103
(I) Demographic Characteristic				
AGE				
12-13	62	60	62	60
14-15	62	58	62	58
16-17	57	59	57	59
GENDER				
Male	71	72	71	72
Female	73	76	73	76
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	96	92	96	92
White	78	74	78	74
Black or African American	35	44	35	44
American Indian or Alaska Native	8	7	8	7
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	28	27	28	27
Two or More Races	13	13	13	13
Hispanic or Latino	48	49	48	49
COUNTY TYPE²				
Large Metro	84	79	84	79
Small Metro	58	56	58	56
Nonmetro	38	41	38	41
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	36	33	58	57
Marijuana	29	27	49	51
Any Illicit Drug Other Than Marijuana ³	25	25	43	40
Cigarettes	38	37	64	60
Alcohol	41	44	68	69
Binge Alcohol ⁴	28	32	60	57
PAST YEAR				
Any Illicit Drug ³	50	47	77	76
Marijuana	41	39	67	66
Any Illicit Drug Other Than Marijuana ³	37	39	63	63
Cigarettes	47	49	73	71
Alcohol	59	62	82	84
LIFETIME				
Any Illicit Drug ³	59	58	84	82
Marijuana	49	48	73	73
Any Illicit Drug Other Than Marijuana ³	50	49	70	74
Cigarettes	62	62	84	84
Alcohol	62	66	87	84

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.12D Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

(I) Demographic Characteristic and (II) Drug Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.43	0.42	0.43	0.42
(I) Demographic Characteristic				
AGE				
12-13	0.76	0.73	0.76	0.73
14-15	0.74	0.70	0.74	0.70
16-17	0.72	0.71	0.72	0.71
GENDER				
Male	0.56	0.57	0.56	0.57
Female	0.61	0.63	0.61	0.63
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.46	0.44	0.46	0.44
White	0.51	0.48	0.51	0.48
Black or African American	0.98	1.19	0.98	1.19
American Indian or Alaska Native	4.23	4.02	4.23	4.02
Native Hawaiian or Other Pacific Islander	*	*	*	*
Asian	2.83	2.80	2.83	2.80
Two or More Races	3.40	3.34	3.40	3.34
Hispanic or Latino	1.24	1.22	1.24	1.22
COUNTY TYPE²				
Large Metro	0.64	0.60	0.64	0.60
Small Metro	0.81	0.77	0.81	0.77
Nonmetro	0.90	0.97	0.90	0.97
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.39	0.34	0.38	0.37
Marijuana	0.31	0.29	0.32	0.33
Any Illicit Drug Other Than Marijuana ³	0.26	0.26	0.29	0.26
Cigarettes	0.40	0.39	0.42	0.39
Alcohol	0.44	0.46	0.45	0.45
Binge Alcohol ⁴	0.30	0.34	0.39	0.38
PAST YEAR				
Any Illicit Drug ³	0.53	0.49	0.51	0.50
Marijuana	0.44	0.41	0.44	0.43
Any Illicit Drug Other Than Marijuana ³	0.40	0.41	0.41	0.42
Cigarettes	0.50	0.52	0.48	0.47
Alcohol	0.62	0.65	0.54	0.55
LIFETIME				
Any Illicit Drug ³	0.63	0.61	0.55	0.54
Marijuana	0.52	0.50	0.48	0.48
Any Illicit Drug Other Than Marijuana ³	0.54	0.51	0.46	0.48
Cigarettes	0.66	0.65	0.55	0.55
Alcohol	0.66	0.69	0.57	0.55

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 3.12P Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week, by Demographic Characteristics (I), and Selected Drug Use, by Risk Perceptions of Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week (II), among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

(I) Demographic Characteristic and (II) Drug Use	Have Five or More Drinks of Alcoholic Beverage Once or Twice a Week ¹			
	Great Risk		Moderate/Slight/No Risk	
	2002	2003	2002	2003
TOTAL	0.6807		0.6807	
(I) Demographic Characteristic				
AGE				
12-13	0.4124		0.4124	
14-15	0.7903		0.7903	
16-17	0.9055		0.9055	
GENDER				
Male	0.1127		0.1127	
Female	0.3426		0.3426	
HISPANIC ORIGIN AND RACE				
Not Hispanic or Latino	0.3069		0.3069	
White	0.4134		0.4134	
Black or African American	0.8575		0.8575	
American Indian or Alaska Native	0.9450		0.9450	
Native Hawaiian or Other Pacific Islander	*		*	
Asian	0.7635		0.7635	
Two or More Races	0.5490		0.5490	
Hispanic or Latino	0.2930		0.2930	
COUNTY TYPE²				
Large Metro	0.5807		0.5807	
Small Metro	0.9152		0.9152	
Nonmetro	0.8234		0.8234	
(II) Drug Use				
PAST MONTH				
Any Illicit Drug ³	0.0983		0.9112	
Marijuana	0.3424		0.6749	
Any Illicit Drug Other Than Marijuana ³	0.1686		0.6667	
Cigarettes	0.0285		0.4164	
Alcohol	0.2854		0.2928	
Binge Alcohol ⁴	0.8469		0.9366	
PAST YEAR				
Any Illicit Drug ³	0.6745		0.5901	
Marijuana	0.2066		0.1834	
Any Illicit Drug Other Than Marijuana ³	0.7195		0.6873	
Cigarettes	0.0055		0.1492	
Alcohol	0.3843		0.9026	
LIFETIME				
Any Illicit Drug ³	0.5178		0.8442	
Marijuana	0.1564		0.1109	
Any Illicit Drug Other Than Marijuana ³	0.6153		0.8412	
Cigarettes	0.0034		0.0106	
Alcohol	0.0247		0.4944	

*Low precision; no estimate reported.

¹ Respondents with missing data were excluded.

² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.