

Table 6.33A Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Numbers in Thousands, 2002 and 2003

Location of Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	3,136	2,983	479	507	1,956	1,710	702	766
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	8,461	8,216	1,091	1,112	4,914	5,034	2,456	2,069
Doctor's Office - Not Part of a Clinic	3,796	3,669	383	396	2,173	2,113	1,239	1,159
Outpatient Medical Clinic	950	997	167	159	400	539	383	299
Partial Day Hospital or Day Treatment Program	289	310	72	49	199	191	19	70
School or University Setting/Clinic/Center ²	107	78	70	49	37	30	*	*
Church/Religious Setting ²	82	138	25	27	57	93	*	17
Some Other Place	389	403	56	77	229	209	104	117

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.33B Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Percentages, 2002 and 2003

Location of Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	20.1	19.6	22.7	23.7	21.7	19.5	15.6	17.8
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	54.1	54.0	51.7	52.1	54.5	57.4	54.5	48.0
Doctor's Office - Not Part of a Clinic	24.3	24.1	18.2	18.6	24.1	24.1	27.5	26.9
Outpatient Medical Clinic	6.1	6.5	7.9	7.4	4.4	6.1	8.5	6.9
Partial Day Hospital or Day Treatment Program	1.9	2.0	3.4	2.3	2.2	2.2	0.4	1.6
School or University Setting/Clinic/Center ²	0.7	0.5	3.3	2.3	0.4	0.3	*	*
Church/Religious Setting ²	0.5	0.9	1.2	1.3	0.6	1.1	*	0.4
Some Other Place	2.5	2.6	2.7	3.6	2.5	2.4	2.3	2.7

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.34A Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Numbers in Thousands, 2002 and 2003

Location of Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	1,415	1,531	263	294	928	961	*	276
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	3,034	3,125	554	557	1,861	1,829	*	*
Doctor's Office - Not Part of a Clinic	1,528	1,621	194	208	1,044	990	*	*
Outpatient Medical Clinic	370	413	65	91	166	238	*	*
Partial Day Hospital or Day Treatment Program	100	171	21	30	64	104	*	37
School or University Setting/Clinic/Center ²	36	41	27	16	9	25	*	*
Church/Religious Setting ²	25	58	8	16	17	42	*	*
Some Other Place	138	188	21	38	89	105	*	*

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.34B Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Percentages, 2002 and 2003

Location of Treatment/Counseling ¹	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	24.7	25.2	26.7	27.7	25.9	26.8	*	19.3
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	52.9	51.4	56.2	52.4	51.9	51.0	*	*
Doctor's Office - Not Part of a Clinic	26.6	26.7	19.7	19.6	29.1	27.6	*	*
Outpatient Medical Clinic	6.5	6.8	6.7	8.5	4.6	6.7	*	*
Partial Day Hospital or Day Treatment Program	1.7	2.8	2.2	2.8	1.8	2.9	*	2.6
School or University Setting/Clinic/Center ²	0.6	0.7	2.7	1.5	0.3	0.7	*	*
Church/Religious Setting ²	0.4	1.0	0.8	1.5	0.5	1.2	*	*
Some Other Place	2.4	3.1	2.2	3.6	2.5	2.9	*	*

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.