

**Table 6.30C Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling:
Standard Errors of Numbers in Thousands, 2002 and 2003**

| Drug | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG³ | 587 | 574 | 236 | 240 | 531 | 490 |
| Marijuana and Hashish | 464 | 457 | 185 | 179 | 419 | 396 |
| Cocaine | 221 | 232 | 122 | 87 | 191 | 213 |
| Crack | 124 | 123 | 81 | 60 | 99 | 103 |
| Heroin | 54 | 47 | 38 | 23 | 39 | 41 |
| Hallucinogens | 145 | 141 | 62 | 51 | 125 | 129 |
| LSD | 58 | 41 | 27 | 13 | 49 | 38 |
| PCP | 21 | 20 | 9 | 10 | 19 | 17 |
| Ecstasy | 123 | 102 | 53 | 39 | 105 | 94 |
| Inhalants | 80 | 67 | 35 | 29 | 73 | 62 |
| Nonmedical Use of Any Psychotherapeutic ⁴ | 383 | 361 | 177 | 169 | 321 | 301 |
| Pain Relievers | 297 | 304 | 147 | 135 | 253 | 263 |
| Tranquilizers | 222 | 195 | 116 | 112 | 180 | 157 |
| Stimulants | 144 | 124 | 81 | 61 | 114 | 106 |
| Methamphetamine | 105 | 93 | 46 | 43 | 89 | 82 |
| Sedatives | 115 | 83 | 76 | 64 | 67 | 54 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA³ | 438 | 425 | 199 | 188 | 377 | 366 |

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.30D Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling:
Standard Errors of Percentages, 2002 and 2003**

| Drug | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG³ | 0.27 | 0.25 | 0.80 | 0.77 | 0.28 | 0.26 |
| Marijuana and Hashish | 0.22 | 0.21 | 0.66 | 0.60 | 0.23 | 0.21 |
| Cocaine | 0.10 | 0.11 | 0.44 | 0.30 | 0.10 | 0.11 |
| Crack | 0.06 | 0.06 | 0.29 | 0.21 | 0.05 | 0.06 |
| Heroin | 0.03 | 0.02 | 0.14 | 0.08 | 0.02 | 0.02 |
| Hallucinogens | 0.07 | 0.07 | 0.22 | 0.18 | 0.07 | 0.07 |
| LSD | 0.03 | 0.02 | 0.10 | 0.05 | 0.03 | 0.02 |
| PCP | 0.01 | 0.01 | 0.03 | 0.04 | 0.01 | 0.01 |
| Ecstasy | 0.06 | 0.05 | 0.19 | 0.14 | 0.06 | 0.05 |
| Inhalants | 0.04 | 0.03 | 0.13 | 0.10 | 0.04 | 0.03 |
| Nonmedical Use of Any Psychotherapeutic ⁴ | 0.17 | 0.17 | 0.59 | 0.57 | 0.17 | 0.16 |
| Pain Relievers | 0.14 | 0.14 | 0.51 | 0.47 | 0.14 | 0.14 |
| Tranquilizers | 0.10 | 0.09 | 0.41 | 0.39 | 0.10 | 0.08 |
| Stimulants | 0.07 | 0.06 | 0.29 | 0.22 | 0.06 | 0.06 |
| Methamphetamine | 0.05 | 0.04 | 0.17 | 0.15 | 0.05 | 0.04 |
| Sedatives | 0.05 | 0.04 | 0.28 | 0.23 | 0.04 | 0.03 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA³ | 0.20 | 0.19 | 0.66 | 0.62 | 0.20 | 0.20 |

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.30P Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling:
P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

| Drug | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG³ | 0.5876 | | 0.6250 | | 0.3849 | |
| Marijuana and Hashish | 0.3442 | | 0.7276 | | 0.3319 | |
| Cocaine | 0.9550 | | 0.0673 | | 0.3259 | |
| Crack | 0.3562 | | 0.4088 | | 0.5902 | |
| Heroin | 0.2823 | | 0.4333 | | 0.4550 | |
| Hallucinogens | 0.0004 | | 0.0292 | | 0.0042 | |
| LSD | 0.0001 | | 0.0125 | | 0.0010 | |
| PCP | 0.9285 | | 0.7231 | | 0.9850 | |
| Ecstasy | 0.0000 | | 0.0055 | | 0.0000 | |
| Inhalants | 0.5630 | | 0.6988 | | 0.6015 | |
| Nonmedical Use of Any Psychotherapeutic ⁴ | 0.7429 | | 0.3491 | | 0.8845 | |
| Pain Relievers | 0.1889 | | 0.5030 | | 0.2576 | |
| Tranquilizers | 0.6296 | | 0.4526 | | 0.9973 | |
| Stimulants | 0.0513 | | 0.3120 | | 0.0900 | |
| Methamphetamine | 0.1943 | | 0.4504 | | 0.2686 | |
| Sedatives | 0.2878 | | 0.6576 | | 0.2426 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA³ | 0.5770 | | 0.9161 | | 0.4715 | |

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

⁴ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.31C Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--------------------------------|--------------------|-------|---|------|--------------|-------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| PAST YEAR | | | | | | |
| Any Tobacco ³ | 1,096 | 1,042 | 344 | 356 | 1,035 | 913 |
| Cigarettes | 1,003 | 956 | 332 | 333 | 946 | 837 |
| Smokeless Tobacco | 321 | 314 | 87 | 104 | 306 | 291 |
| Cigars | 495 | 498 | 162 | 171 | 466 | 453 |
| Pipe Tobacco ⁴ | -- | -- | -- | -- | -- | -- |
| Alcohol | 1,796 | 1,725 | 516 | 524 | 1,666 | 1,537 |
| Binge Alcohol Use ⁵ | -- | -- | -- | -- | -- | -- |
| Heavy Alcohol Use ⁵ | -- | -- | -- | -- | -- | -- |
| PAST MONTH | | | | | | |
| Any Tobacco ³ | 1,002 | 972 | 320 | 339 | 956 | 857 |
| Cigarettes | 936 | 901 | 309 | 324 | 884 | 796 |
| Smokeless Tobacco | 282 | 288 | 67 | 94 | 273 | 271 |
| Cigars | 348 | 353 | 102 | 125 | 339 | 316 |
| Pipe Tobacco | 164 | 130 | 48 | 52 | 156 | 122 |
| Alcohol | 1,531 | 1,507 | 447 | 439 | 1,419 | 1,362 |
| Binge Alcohol Use ⁵ | 838 | 787 | 232 | 234 | 794 | 717 |
| Heavy Alcohol Use ⁵ | 400 | 401 | 117 | 109 | 377 | 378 |

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.31D Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2002 and 2003

| Drug | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--------------------------------|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| PAST YEAR | | | | | | |
| Any Tobacco ³ | 0.42 | 0.38 | 1.08 | 1.02 | 0.44 | 0.40 |
| Cigarettes | 0.39 | 0.38 | 1.07 | 0.99 | 0.42 | 0.39 |
| Smokeless Tobacco | 0.15 | 0.15 | 0.32 | 0.36 | 0.16 | 0.16 |
| Cigars | 0.23 | 0.21 | 0.57 | 0.57 | 0.24 | 0.23 |
| Pipe Tobacco ⁴ | -- | -- | -- | -- | -- | -- |
| Alcohol | 0.43 | 0.42 | 1.08 | 1.01 | 0.47 | 0.44 |
| Binge Alcohol Use ⁵ | -- | -- | -- | -- | -- | -- |
| Heavy Alcohol Use ⁵ | -- | -- | -- | -- | -- | -- |
| PAST MONTH | | | | | | |
| Any Tobacco ³ | 0.39 | 0.37 | 1.06 | 0.97 | 0.42 | 0.39 |
| Cigarettes | 0.38 | 0.36 | 1.03 | 0.96 | 0.40 | 0.38 |
| Smokeless Tobacco | 0.13 | 0.13 | 0.24 | 0.33 | 0.15 | 0.15 |
| Cigars | 0.16 | 0.16 | 0.37 | 0.42 | 0.18 | 0.17 |
| Pipe Tobacco | 0.08 | 0.06 | 0.18 | 0.18 | 0.08 | 0.07 |
| Alcohol | 0.46 | 0.43 | 1.14 | 1.06 | 0.50 | 0.46 |
| Binge Alcohol Use ⁵ | 0.35 | 0.32 | 0.78 | 0.73 | 0.38 | 0.34 |
| Heavy Alcohol Use ⁵ | 0.19 | 0.18 | 0.42 | 0.37 | 0.20 | 0.20 |

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.³ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.⁴ Information about past year use of pipe tobacco was not collected.⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.31P Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--------------------------------|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| PAST YEAR | | | | | | |
| Any Tobacco ³ | 0.0973 | | 0.9305 | | 0.0656 | |
| Cigarettes | 0.1001 | | 0.7721 | | 0.0800 | |
| Smokeless Tobacco | 0.4865 | | 0.9408 | | 0.4873 | |
| Cigars | 0.2509 | | 0.7449 | | 0.2485 | |
| Pipe Tobacco ⁴ | -- | | -- | | -- | |
| Alcohol | 0.0414 | | 0.3357 | | 0.0109 | |
| Binge Alcohol Use ⁵ | -- | | -- | | -- | |
| Heavy Alcohol Use ⁵ | -- | | -- | | -- | |
| PAST MONTH | | | | | | |
| Any Tobacco ³ | 0.2451 | | 0.7633 | | 0.1548 | |
| Cigarettes | 0.2930 | | 0.9339 | | 0.2151 | |
| Smokeless Tobacco | 0.6821 | | 0.6123 | | 0.5723 | |
| Cigars | 0.9369 | | 0.9200 | | 0.9488 | |
| Pipe Tobacco | 0.2325 | | 0.7681 | | 0.1677 | |
| Alcohol | 0.0971 | | 0.8387 | | 0.0639 | |
| Binge Alcohol Use ⁵ | 0.5452 | | 0.4493 | | 0.7361 | |
| Heavy Alcohol Use ⁵ | 0.8533 | | 0.7163 | | 0.7902 | |

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column represent past year and past month use for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

⁴ Information about past year use of pipe tobacco was not collected.

⁵ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.32C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Numbers in Thousands, 2002 and 2003

| Dependence/Abuse | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| DEPENDENCE OR ABUSE³ | | | | | | |
| Any Illicit Drug or Alcohol ⁴ | 471 | 428 | 215 | 196 | 403 | 368 |
| Any Illicit Drug ⁴ | 229 | 205 | 141 | 112 | 162 | 166 |
| Alcohol | 436 | 397 | 186 | 167 | 387 | 345 |
| DEPENDENCE³ | | | | | | |
| Any Illicit Drug or Alcohol ⁴ | 330 | 303 | 174 | 150 | 269 | 256 |
| Any Illicit Drug ⁴ | 179 | 159 | 109 | 97 | 135 | 124 |
| Alcohol | 288 | 269 | 151 | 122 | 241 | 231 |

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.32D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: Standard Errors of Percentages, 2002 and 2003

| Dependence/Abuse | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| DEPENDENCE OR ABUSE³ | | | | | | |
| Any Illicit Drug or Alcohol ⁴ | 0.21 | 0.19 | 0.73 | 0.65 | 0.20 | 0.20 |
| Any Illicit Drug ⁴ | 0.11 | 0.09 | 0.50 | 0.39 | 0.09 | 0.09 |
| Alcohol | 0.19 | 0.18 | 0.66 | 0.56 | 0.20 | 0.19 |
| DEPENDENCE³ | | | | | | |
| Any Illicit Drug or Alcohol ⁴ | 0.15 | 0.14 | 0.62 | 0.51 | 0.14 | 0.14 |
| Any Illicit Drug ⁴ | 0.08 | 0.07 | 0.39 | 0.34 | 0.07 | 0.07 |
| Alcohol | 0.13 | 0.12 | 0.55 | 0.43 | 0.13 | 0.12 |

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 6.32P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Dependence/Abuse | Total ¹ | | MENTAL HEALTH TREATMENT/COUNSELING ² | | | |
|--|--------------------|------|---|------|--------------|------|
| | | | Received | | Not Received | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| DEPENDENCE OR ABUSE³ | | | | | | |
| Any Illicit Drug or Alcohol ⁴ | 0.2436 | | 0.1139 | | 0.6550 | |
| Any Illicit Drug ⁴ | 0.3827 | | 0.2936 | | 0.7203 | |
| Alcohol | 0.3540 | | 0.2651 | | 0.6425 | |
| DEPENDENCE³ | | | | | | |
| Any Illicit Drug or Alcohol ⁴ | 0.0992 | | 0.0715 | | 0.3582 | |
| Any Illicit Drug ⁴ | 0.3491 | | 0.7253 | | 0.3253 | |
| Alcohol | 0.0578 | | 0.0416 | | 0.2898 | |

*Low precision; no estimate reported.

¹ Estimates in the Total column represent past year dependence/abuse for all persons aged 18 or older, including those with unknown mental health treatment/counseling information.

² Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

³ Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

⁴ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.