

**Table 7.17A Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	5,178	5,247	6,307	6,216	3,853	4,136	4,185	3,871
Marijuana and Hashish	4,327	4,266	5,079	5,185	2,912	2,999	2,266	2,189
Cocaine	904	1,096	695	646	206	316	215	222
Crack	187	219	181	133	81	123	118	129
Heroin	26	24	54	39	18	28	69	28
Hallucinogens	472	424	435	325	129	163	161	131
LSD	31 <sup>a</sup>	68	56	46	15	5	10	14
PCP	17	24	29	26	*	*	10	*
Ecstasy	268 <sup>b</sup>	158	252 <sup>a</sup>	156	88	111	69	44
Inhalants	125	125	123	132	93	97	293	216
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	1,486	1,697	1,730	1,670	1,137	1,330	1,858	1,639
Pain Relievers	1,046 <sup>a</sup>	1,281	1,258	1,228	717	874	1,357	1,310
Tranquilizers	470	563	501	517	398	454	435	296
Stimulants	481	480	323	336	163	197	251	179
Methamphetamine	236	269	162	160	80	99	120	79
Sedatives	90	98	124	66	119	84	104	47
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	2,363	2,612	2,561	2,452	1,469	1,730	2,385	2,055

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.17B Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	32.6	32.5	16.6	16.5	5.8	6.3	3.6	3.3
Marijuana and Hashish	27.3	26.4	13.4	13.8	4.4	4.6	2.0	1.8
Cocaine	5.7	6.8	1.8	1.7	0.3	0.5	0.2	0.2
Crack	1.2	1.4	0.5	0.4	0.1	0.2	0.1	0.1
Heroin	0.2	0.1	0.1	0.1	0.0	0.0	0.1	0.0
Hallucinogens	3.0	2.6	1.1	0.9	0.2	0.2	0.1	0.1
LSD	0.2 <sup>a</sup>	0.4	0.1	0.1	0.0	0.0	0.0	0.0
PCP	0.1	0.1	0.1	0.1	*	*	0.0	*
Ecstasy	1.7 <sup>b</sup>	1.0	0.7 <sup>a</sup>	0.4	0.1	0.2	0.1	0.0
Inhalants	0.8	0.8	0.3	0.4	0.1	0.1	0.3	0.2
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	9.4	10.5	4.6	4.4	1.7	2.0	1.6	1.4
Pain Relievers	6.6 <sup>a</sup>	7.9	3.3	3.3	1.1	1.3	1.2	1.1
Tranquilizers	3.0	3.5	1.3	1.4	0.6	0.7	0.4	0.2
Stimulants	3.0	3.0	0.9	0.9	0.2	0.3	0.2	0.2
Methamphetamine	1.5	1.7	0.4	0.4	0.1	0.2	0.1	0.1
Sedatives	0.6	0.6	0.3	0.2	0.2	0.1	0.1	0.0
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	14.9	16.2	6.8	6.5	2.2	2.7	2.1	1.7

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.18A Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	422	421	887	913	435	428	1,134	1,049
Marijuana and Hashish	366	366	728	763	336	311	593	531
Cocaine	44	56	63	57	15	20	23	20
Crack	17	21	4	6	*	1	4	4
Heroin	3	7	6	8	*	0	3	3
Hallucinogens	61	89	90	86	36	27	68	57
LSD	20	22	25	16	6	1	10	12
PCP	3	10	9	8	2	*	10	2
Ecstasy	24	40	50	28	15	11	34 <sup>a</sup>	13
Inhalants	38	39	55	76	36	53	172	152
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	135 <sup>a</sup>	188	287	282	122	114	436	413
Pain Relievers	112	151	236	213	87	99	360	339
Tranquilizers	44	68	68	78	25	20	50	61
Stimulants	39	66	67	74	28	23	66	58
Methamphetamine	20	29	24	12	10	10	10	18
Sedatives	4	12	17	14	8	6	18	15
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	213	242	391	396	176	181	641	598

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.18B Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	67.0	64.5	44.1	45.5	25.2	24.2	5.6	5.1
Marijuana and Hashish	58.1	56.1	36.2	38.0	19.5	17.6	2.9	2.6
Cocaine	7.1	8.5	3.1	2.9	0.9	1.2	0.1	0.1
Crack	2.8	3.2	0.2	0.3	*	0.1	0.0	0.0
Heroin	0.4	1.1	0.3	0.4	*	0.0	0.0	0.0
Hallucinogens	9.7	13.7	4.5	4.3	2.1	1.5	0.3	0.3
LSD	3.1	3.3	1.3	0.8	0.3	0.1	0.0	0.1
PCP	0.5	1.6	0.4	0.4	0.1	*	0.0	0.0
Ecstasy	3.8	6.1	2.5	1.4	0.9	0.6	0.2 <sup>a</sup>	0.1
Inhalants	6.1	6.0	2.7	3.8	2.1	3.0	0.8	0.7
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	21.5 <sup>a</sup>	28.8	14.2	14.1	7.1	6.5	2.1	2.0
Pain Relievers	17.7	23.1	11.7	10.6	5.0	5.6	1.8	1.6
Tranquilizers	7.0	10.4	3.4	3.9	1.4	1.1	0.2	0.3
Stimulants	6.2	10.1	3.3	3.7	1.6	1.3	0.3	0.3
Methamphetamine	3.1	4.5	1.2	0.6	0.6	0.6	0.0	0.1
Sedatives	0.7	1.8	0.9	0.7	0.5	0.3	0.1	0.1
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	33.8	37.2	19.4	19.7	10.2	10.2	3.1	2.9

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.19A Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	2,277	2,336	2,255	2,307	906	1,013	827	778
Marijuana and Hashish	2,020	2,015	1,968	1,990	797	872	590	522
Cocaine	346	425	211	196	39	42	35	34
Crack	21	23	17	21	10	7	5	4
Heroin	11	10	12	8	1	*	15	6
Hallucinogens	257	262	235	180	54	57	43	46
LSD	11 <sup>a</sup>	35	16	26	9	4	*	2
PCP	1	11	14	16	*	*	*	3
Ecstasy	155 <sup>b</sup>	95	140 <sup>a</sup>	81	34	28	25	26
Inhalants	70	58	50	35	5	22	25	19
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	680	766	552	623	201	229	250	295
Pain Relievers	510	585	420	480	135	175	197	246
Tranquilizers	213	252	171	165	51	68	52	40
Stimulants	211	191	101	137	41	33	33	39
Methamphetamine	80	74	44	70	20	17	16	24
Sedatives	18	26	25	19	0	3	5	10
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	1,032	1,147	841	874	256	296	330	348

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.19B Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	49.2	48.8	27.9	27.5	14.9	16.1	6.7	6.3
Marijuana and Hashish	43.7	42.1	24.4	23.7	13.2	13.9	4.8	4.3
Cocaine	7.5	8.9	2.6	2.3	0.6	0.7	0.3	0.3
Crack	0.5	0.5	0.2	0.2	0.2	0.1	0.0	0.0
Heroin	0.2	0.2	0.2	0.1	0.0	*	0.1	0.1
Hallucinogens	5.6	5.5	2.9	2.1	0.9	0.9	0.4	0.4
LSD	0.2 <sup>a</sup>	0.7	0.2	0.3	0.2	0.1	*	0.0
PCP	0.0	0.2	0.2	0.2	*	*	*	0.0
Ecstasy	3.4 <sup>b</sup>	2.0	1.7 <sup>b</sup>	1.0	0.6	0.4	0.2	0.2
Inhalants	1.5	1.2	0.6	0.4	0.1	0.3	0.2	0.2
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	14.7	16.0	6.8	7.4	3.3	3.6	2.0	2.4
Pain Relievers	11.0	12.2	5.2	5.7	2.2	2.8	1.6	2.0
Tranquilizers	4.6	5.3	2.1	2.0	0.8	1.1	0.4	0.3
Stimulants	4.6	4.0	1.3	1.6	0.7	0.5	0.3	0.3
Methamphetamine	1.7	1.5	0.5	0.8	0.3	0.3	0.1	0.2
Sedatives	0.4	0.5	0.3	0.2	0.0	0.0	0.0	0.1
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	22.3	24.0	10.4	10.4	4.2	4.7	2.7	2.8

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.20A Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	2,479	2,490	3,165	2,997	2,512	2,694	2,224	2,043
Marijuana and Hashish	1,941	1,885	2,382	2,431	1,779	1,816	1,083	1,136
Cocaine	513	616	420	393	152	254	157	168
Crack	148	175	161	107	71	115	109	120
Heroin	12	7	36	23	17	28	51	19
Hallucinogens	154	73	110	60	39	78	50	28
LSD	*	11	15	4	*	*	*	*
PCP	13	3	7	2	*	*	*	*
Ecstasy	88 <sup>a</sup>	24	61	47	38	73	9	5
Inhalants	17	28	18	21	52	23	97	44
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	671	743	891	765	814	987	1,172	931
Pain Relievers	424	545	602	535	496	600	800	726
Tranquilizers	213	243	262	274	322	366	333	195
Stimulants	231	224	155	125	94	141	151	81
Methamphetamine	137	166	95	78	50	72	93	36
Sedatives	67	61	81	33	111	75	81 <sup>a</sup>	21
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	1,118	1,222	1,329	1,181	1,037	1,254	1,413	1,110

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.20B Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	23.4	23.3	11.4	11.0	4.3	4.7	2.7	2.4
Marijuana and Hashish	18.3	17.6	8.6	8.9	3.1	3.2	1.3	1.3
Cocaine	4.8	5.8	1.5	1.4	0.3	0.4	0.2	0.2
Crack	1.4	1.6	0.6	0.4	0.1	0.2	0.1	0.1
Heroin	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0
Hallucinogens	1.4	0.7	0.4	0.2	0.1	0.1	0.1	0.0
LSD	*	0.1	0.1	0.0	*	*	*	*
PCP	0.1	0.0	0.0	0.0	*	*	*	*
Ecstasy	0.8 <sup>a</sup>	0.2	0.2	0.2	0.1	0.1	0.0	0.0
Inhalants	0.2	0.3	0.1	0.1	0.1	0.0	0.1	0.1
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	6.3	6.9	3.2	2.8	1.4	1.7	1.4	1.1
Pain Relievers	4.0	5.1	2.2	2.0	0.9	1.1	1.0	0.8
Tranquilizers	2.0	2.3	0.9	1.0	0.6	0.6	0.4	0.2
Stimulants	2.2	2.1	0.6	0.5	0.2	0.2	0.2	0.1
Methamphetamine	1.3	1.6	0.3	0.3	0.1	0.1	0.1	0.0
Sedatives	0.6	0.6	0.3	0.1	0.2	0.1	0.1 <sup>a</sup>	0.0
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	10.5	11.4	4.8	4.3	1.8	2.2	1.7	1.3

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.