

**Table 7.21A Tobacco and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	10,363	10,323
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	2,946	2,946	4,841	4,779
Cigars	7,358	7,354	5,394	5,483
Pipe Tobacco	902	837	915	782
<b>ALCOHOL</b>	40,688	39,747	79,133	79,217
Binge Alcohol Use <sup>2</sup>	26,323	26,258	27,464	27,512
Heavy Alcohol Use <sup>2</sup>	9,722	9,953	6,138	6,191

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.21B Tobacco and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	6.0	5.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.8	4.9	2.8	2.7
Cigars	12.0	12.2	3.1	3.1
Pipe Tobacco	1.5	1.4	0.5	0.4
<b>ALCOHOL</b>	66.6	65.8	45.5	44.7
Binge Alcohol Use <sup>2</sup>	43.1	43.4	15.8	15.5
Heavy Alcohol Use <sup>2</sup>	15.9	16.5	3.5	3.5

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.22A Tobacco and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	558	566
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	273	279	215	232
Cigars	773	777	344	338
Pipe Tobacco	91	120	46	40
<b>ALCOHOL</b>	1,893	1,867	2,472	2,558
Binge Alcohol Use <sup>2</sup>	1,435	1,438	1,206	1,221
Heavy Alcohol Use <sup>2</sup>	449	472	181	180

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.22B Tobacco and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	2.6	2.6
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	8.5	9.2	1.0	1.1
Cigars	24.1	25.5	1.6	1.5
Pipe Tobacco	2.8	3.9	0.2	0.2
<b>ALCOHOL</b>	59.0	61.4	11.5	11.7
Binge Alcohol Use <sup>2</sup>	44.7	47.2	5.6	5.6
Heavy Alcohol Use <sup>2</sup>	14.0	15.5	0.8	0.8

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.23A Tobacco and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	1,399	1,437
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,009	1,023	493	462
Cigars	2,479	2,618	938	1,004
Pipe Tobacco	271	222	61	71
<b>ALCOHOL</b>	9,896	10,036	8,863 <sup>a</sup>	9,430
Binge Alcohol Use <sup>2</sup>	7,776	7,960	4,923	5,228
Heavy Alcohol Use <sup>2</sup>	3,293	3,383	1,332	1,405

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.23B Tobacco and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	7.6	7.6
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	8.0	8.0	2.7	2.4
Cigars	19.6	20.5	5.1	5.3
Pipe Tobacco	2.1	1.7	0.3	0.4
<b>ALCOHOL</b>	78.2	78.6	48.2	49.7
Binge Alcohol Use <sup>2</sup>	61.5	62.4	26.8	27.6
Heavy Alcohol Use <sup>2</sup>	26.0	26.5	7.3	7.4

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.24A Tobacco and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	8,405	8,319
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,664	1,643	4,133	4,085
Cigars	4,105	3,960	4,112	4,140
Pipe Tobacco	540	495	808	671
<b>ALCOHOL</b>	28,898	27,844	67,798	67,229
Binge Alcohol Use <sup>2</sup>	17,112	16,860	21,335	21,064
Heavy Alcohol Use <sup>2</sup>	5,981	6,098	4,624	4,605

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.24B Tobacco and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2002 and 2003**

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
<b>ANY TOBACCO<sup>1</sup></b>	N/A	N/A	6.3	6.1
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3.7	3.7	3.1	3.0
Cigars	9.1	8.9	3.1	3.0
Pipe Tobacco	1.2	1.1	0.6	0.5
<b>ALCOHOL</b>	63.8	62.4	50.6	49.3
Binge Alcohol Use <sup>2</sup>	37.8	37.8	15.9	15.5
Heavy Alcohol Use <sup>2</sup>	13.2	13.7	3.4	3.4

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.