

Table 7.17C Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	162	150	199	191	203	218	190	177
Marijuana and Hashish	148	140	171	173	168	160	125	127
Cocaine	78	86	71	68	41	69	39	44
Crack	39	41	44	37	29	52	32	38
Heroin	12	9	24	15	13	17	29	13
Hallucinogens	47	37	43	37	38	39	29	24
LSD	7	13	15	12	6	3	4	5
PCP	11	10	10	8	*	*	4	*
Ecstasy	35	23	34	31	37	37	14	12
Inhalants	19	23	18	20	27	16	54	28
Nonmedical Use of Any Psychotherapeutic ³	90	83	99	98	111	148	137	128
Pain Relievers	72	76	82	82	72	115	119	116
Tranquilizers	48	48	60	53	85	76	68	49
Stimulants	59	50	46	47	28	41	40	31
Methamphetamine	48	44	37	37	20	28	33	19
Sedatives	25	30	28	16	44	40	28	15
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	118	113	123	121	129	167	152	135

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.17D Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	1.02	0.93	0.52	0.51	0.31	0.33	0.17	0.15
Marijuana and Hashish	0.93	0.87	0.45	0.46	0.25	0.24	0.11	0.11
Cocaine	0.49	0.53	0.19	0.18	0.06	0.11	0.03	0.04
Crack	0.24	0.26	0.12	0.10	0.04	0.08	0.03	0.03
Heroin	0.08	0.06	0.06	0.04	0.02	0.03	0.03	0.01
Hallucinogens	0.30	0.23	0.11	0.10	0.06	0.06	0.03	0.02
LSD	0.05	0.08	0.04	0.03	0.01	0.00	0.00	0.00
PCP	0.07	0.06	0.03	0.02	*	*	0.00	*
Ecstasy	0.22	0.14	0.09	0.08	0.06	0.06	0.01	0.01
Inhalants	0.12	0.14	0.05	0.05	0.04	0.03	0.05	0.02
Nonmedical Use of Any Psychotherapeutic ³	0.57	0.51	0.26	0.26	0.17	0.23	0.12	0.11
Pain Relievers	0.46	0.47	0.22	0.22	0.11	0.18	0.10	0.10
Tranquilizers	0.30	0.30	0.16	0.14	0.13	0.12	0.06	0.04
Stimulants	0.37	0.31	0.12	0.12	0.04	0.06	0.03	0.03
Methamphetamine	0.30	0.27	0.10	0.10	0.03	0.04	0.03	0.02
Sedatives	0.16	0.18	0.07	0.04	0.07	0.06	0.02	0.01
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.74	0.70	0.33	0.32	0.19	0.26	0.13	0.11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.17P Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	0.9168		0.8866		0.2621		0.1000	
Marijuana and Hashish	0.4972		0.5676		0.5760		0.4314	
Cocaine	0.1283		0.6592		0.1570		0.9941	
Crack	0.6148		0.4232		0.4631		0.8841	
Heroin	0.8909		0.6008		0.6052		0.1860	
Hallucinogens	0.3423		0.0634		0.5152		0.3574	
LSD	0.0119		0.5992		0.1658		0.6280	
PCP	0.6289		0.8105		*		*	
Ecstasy	0.0068		0.0409		0.6371		0.1418	
Inhalants	0.9473		0.7215		0.8605		0.1660	
Nonmedical Use of Any Psychotherapeutic ³	0.1438		0.7319		0.2555		0.1559	
Pain Relievers	0.0485		0.8594		0.2166		0.5958	
Tranquilizers	0.2031		0.7964		0.5903		0.0741	
Stimulants	0.9113		0.8213		0.4711		0.1061	
Methamphetamine	0.6653		0.9865		0.5572		0.2476	
Sedatives	0.8561		0.0780		0.5694		0.0657	
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.2111		0.6026		0.1798		0.0561	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.18C Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	14	16	28	30	24	25	48	42
Marijuana and Hashish	14	17	28	29	21	23	34	31
Cocaine	9	10	13	9	6	6	6	5
Crack	8	7	3	2	*	1	2	2
Heroin	1	4	5	4	*	0	2	2
Hallucinogens	10	11	13	12	8	6	10	11
LSD	6	6	9	6	4	1	4	5
PCP	2	5	4	3	1	*	4	1
Ecstasy	6	8	11	7	5	5	8	5
Inhalants	10	8	10	12	8	9	17	17
Nonmedical Use of Any Psychotherapeutic ³	13	15	22	20	14	15	28	28
Pain Relievers	12	13	22	18	12	15	26	25
Tranquilizers	8	11	11	12	8	6	8	12
Stimulants	8	11	11	11	6	6	12	10
Methamphetamine	6	7	8	4	4	4	4	6
Sedatives	2	5	5	6	4	2	5	6
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	15	16	25	22	17	18	34	32

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.18D Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	2.22	2.50	1.39	1.51	1.40	1.42	0.24	0.21
Marijuana and Hashish	2.26	2.55	1.41	1.45	1.25	1.32	0.16	0.15
Cocaine	1.46	1.55	0.64	0.47	0.35	0.34	0.03	0.03
Crack	1.22	1.01	0.14	0.12	*	0.04	0.01	0.01
Heroin	0.20	0.68	0.23	0.19	*	0.02	0.01	0.01
Hallucinogens	1.55	1.71	0.63	0.61	0.47	0.37	0.05	0.05
LSD	1.00	0.97	0.43	0.29	0.26	0.07	0.02	0.02
PCP	0.25	0.76	0.17	0.14	0.07	*	0.02	0.01
Ecstasy	1.00	1.16	0.54	0.36	0.30	0.26	0.04	0.02
Inhalants	1.65	1.23	0.49	0.58	0.47	0.54	0.08	0.08
Nonmedical Use of Any Psychotherapeutic ³	2.01	2.29	1.12	1.01	0.84	0.85	0.14	0.13
Pain Relievers	1.97	2.01	1.09	0.88	0.72	0.82	0.13	0.12
Tranquilizers	1.29	1.72	0.53	0.58	0.44	0.35	0.04	0.06
Stimulants	1.21	1.66	0.54	0.52	0.37	0.32	0.06	0.05
Methamphetamine	1.02	1.10	0.39	0.21	0.21	0.22	0.02	0.03
Sedatives	0.34	0.75	0.24	0.28	0.24	0.13	0.02	0.03
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	2.34	2.47	1.22	1.11	0.98	1.04	0.17	0.16

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.18P Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	0.4541		0.5030		0.6285		0.1469	
Marijuana and Hashish	0.5476		0.3602		0.3014		0.1477	
Cocaine	0.4938		0.7276		0.5862		0.6851	
Crack	0.8057		0.6215		*		0.9910	
Heroin	0.3404		0.7799		*		0.9739	
Hallucinogens	0.0805		0.8332		0.3536		0.4506	
LSD	0.8822		0.3849		0.3031		0.7698	
PCP	0.1662		0.8880		*		0.0915	
Ecstasy	0.1299		0.0948		0.4780		0.0161	
Inhalants	0.9686		0.1532		0.2212		0.3947	
Nonmedical Use of Any Psychotherapeutic ³	0.0168		0.9030		0.6210		0.5050	
Pain Relievers	0.0558		0.4418		0.6146		0.5172	
Tranquilizers	0.1237		0.5236		0.5672		0.4898	
Stimulants	0.0613		0.6382		0.4920		0.5972	
Methamphetamine	0.3767		0.2058		0.9892		0.2871	
Sedatives	0.1884		0.6810		0.6094		0.7356	
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.3204		0.8532		0.9920		0.3080	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.19C Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	51	52	63	64	41	46	44	44
Marijuana and Hashish	52	50	59	61	39	44	38	36
Cocaine	25	31	22	20	8	10	8	8
Crack	6	5	5	7	4	5	3	2
Heroin	4	5	5	3	1	*	7	4
Hallucinogens	21	24	25	21	10	11	12	11
LSD	4	9	5	9	5	2	*	1
PCP	0	7	6	7	*	*	*	2
Ecstasy	17	14	20	13	8	7	9	9
Inhalants	12	12	11	9	3	9	9	7
Nonmedical Use of Any Psychotherapeutic ³	36	39	33	38	22	23	25	26
Pain Relievers	32	34	30	32	15	20	23	25
Tranquilizers	21	26	21	18	13	12	11	9
Stimulants	21	19	14	20	10	8	8	9
Methamphetamine	15	13	9	15	7	6	6	8
Sedatives	6	7	7	6	0	2	3	6
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	43	46	41	45	24	24	30	28

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.19D Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	1.11	1.08	0.78	0.76	0.68	0.74	0.36	0.36
Marijuana and Hashish	1.12	1.03	0.73	0.72	0.64	0.70	0.31	0.29
Cocaine	0.55	0.65	0.27	0.23	0.14	0.15	0.06	0.06
Crack	0.12	0.11	0.07	0.09	0.06	0.08	0.02	0.02
Heroin	0.09	0.10	0.07	0.04	0.01	*	0.06	0.03
Hallucinogens	0.46	0.49	0.31	0.25	0.17	0.17	0.10	0.09
LSD	0.08	0.18	0.06	0.11	0.08	0.04	*	0.01
PCP	0.01	0.16	0.07	0.08	*	*	*	0.02
Ecstasy	0.38	0.30	0.25	0.15	0.14	0.11	0.07	0.08
Inhalants	0.25	0.26	0.13	0.11	0.05	0.14	0.07	0.05
Nonmedical Use of Any Psychotherapeutic ³	0.78	0.81	0.41	0.45	0.36	0.36	0.20	0.21
Pain Relievers	0.69	0.70	0.37	0.39	0.25	0.32	0.19	0.20
Tranquilizers	0.46	0.54	0.27	0.21	0.21	0.18	0.09	0.07
Stimulants	0.46	0.40	0.18	0.24	0.17	0.13	0.06	0.08
Methamphetamine	0.32	0.28	0.12	0.17	0.12	0.10	0.05	0.06
Sedatives	0.13	0.14	0.08	0.07	0.01	0.03	0.02	0.05
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.92	0.95	0.51	0.54	0.39	0.38	0.24	0.23

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.19P Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	0.7689		0.6717		0.2315		0.4348	
Marijuana and Hashish	0.2903		0.5120		0.4511		0.1948	
Cocaine	0.1065		0.4170		0.8769		0.9008	
Crack	0.9091		0.7036		0.6228		0.8167	
Heroin	0.8683		0.4836		*		0.2663	
Hallucinogens	0.8895		0.0640		0.9059		0.8602	
LSD	0.0120		0.3766		0.3136		*	
PCP	0.2050		0.8310		*		*	
Ecstasy	0.0043		0.0094		0.4406		0.9575	
Inhalants	0.3842		0.2309		0.0805		0.5988	
Nonmedical Use of Any Psychotherapeutic ³	0.2404		0.3257		0.5090		0.2107	
Pain Relievers	0.2284		0.3165		0.1379		0.1413	
Tranquilizers	0.3402		0.6392		0.3594		0.3915	
Stimulants	0.3383		0.1989		0.5041		0.6214	
Methamphetamine	0.6658		0.1721		0.6873		0.3973	
Sedatives	0.4304		0.3895		0.1914		0.4283	
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.2035		0.9898		0.3728		0.6673	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.20C Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	142	132	181	167	198	210	174	162
Marijuana and Hashish	126	122	154	149	161	148	114	115
Cocaine	70	77	66	64	40	67	38	43
Crack	38	40	44	37	29	52	31	38
Heroin	11	6	23	14	13	17	28	12
Hallucinogens	40	24	32	26	36	36	24	17
LSD	*	7	11	4	*	*	*	*
PCP	11	3	7	2	*	*	*	*
Ecstasy	29	15	25	25	36	36	8	5
Inhalants	10	17	10	13	25	10	48	22
Nonmedical Use of Any Psychotherapeutic ³	80	71	93	86	109	149	133	119
Pain Relievers	62	65	78	74	70	113	115	109
Tranquilizers	41	38	54	53	83	75	67	47
Stimulants	54	44	43	39	26	39	39	27
Methamphetamine	45	40	35	34	19	27	32	17
Sedatives	24	29	27	14	44	40	28	12
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	106	98	117	110	127	165	145	126

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.20D Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	1.34	1.23	0.65	0.61	0.34	0.37	0.21	0.19
Marijuana and Hashish	1.19	1.14	0.55	0.55	0.28	0.26	0.14	0.13
Cocaine	0.66	0.72	0.24	0.23	0.07	0.12	0.05	0.05
Crack	0.36	0.38	0.16	0.13	0.05	0.09	0.04	0.04
Heroin	0.11	0.06	0.08	0.05	0.02	0.03	0.03	0.01
Hallucinogens	0.38	0.22	0.12	0.09	0.06	0.06	0.03	0.02
LSD	*	0.06	0.04	0.01	*	*	*	*
PCP	0.10	0.03	0.03	0.01	*	*	*	*
Ecstasy	0.28	0.14	0.09	0.09	0.06	0.06	0.01	0.01
Inhalants	0.09	0.16	0.04	0.05	0.04	0.02	0.06	0.03
Nonmedical Use of Any Psychotherapeutic ³	0.75	0.66	0.34	0.32	0.19	0.26	0.16	0.14
Pain Relievers	0.58	0.60	0.28	0.27	0.12	0.20	0.14	0.13
Tranquilizers	0.39	0.35	0.19	0.19	0.14	0.13	0.08	0.06
Stimulants	0.51	0.41	0.15	0.14	0.04	0.07	0.05	0.03
Methamphetamine	0.43	0.38	0.13	0.13	0.03	0.05	0.04	0.02
Sedatives	0.23	0.27	0.10	0.05	0.07	0.07	0.03	0.01
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	1.00	0.92	0.42	0.40	0.22	0.29	0.17	0.15

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.20P Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
ANY ILLICIT DRUG²	0.9522		0.7008		0.4334		0.2763	
Marijuana and Hashish	0.6729		0.6447		0.7414		0.9507	
Cocaine	0.3476		0.8384		0.1775		0.9314	
Crack	0.6383		0.3754		0.4378		0.8845	
Heroin	0.6723		0.6497		0.5888		0.2774	
Hallucinogens	0.0715		0.2395		0.4307		0.4270	
LSD	*		0.3206		*		*	
PCP	0.3860		0.4917		*		*	
Ecstasy	0.0473		0.7056		0.4826		0.6517	
Inhalants	0.5641		0.8331		0.2931		0.3021	
Nonmedical Use of Any Psychotherapeutic ³	0.5569		0.3999		0.3088		0.1203	
Pain Relievers	0.2101		0.6152		0.3946		0.5049	
Tranquilizers	0.5992		0.8036		0.6537		0.0716	
Stimulants	0.9017		0.6419		0.3057		0.1037	
Methamphetamine	0.6459		0.7659		0.4830		0.1051	
Sedatives	0.8547		0.1218		0.5684		0.0434	
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.5285		0.4481		0.2567		0.0719	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.