

Table 7.21C Tobacco and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	336	340
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	149	144	226	229
Cigars	240	240	240	242
Pipe Tobacco	85	86	141	102
ALCOHOL	441	401	855	842
Binge Alcohol Use ²	408	388	574	511
Heavy Alcohol Use ²	266	270	278	244

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.21D Tobacco and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	0.19	0.19
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.24	0.24	0.13	0.13
Cigars	0.39	0.40	0.14	0.14
Pipe Tobacco	0.14	0.14	0.08	0.06
ALCOHOL	0.72	0.66	0.49	0.47
Binge Alcohol Use ²	0.67	0.64	0.33	0.29
Heavy Alcohol Use ²	0.44	0.45	0.16	0.14

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.21P Tobacco and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A		0.6249	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.8709		0.6466	
Cigars	0.8157		0.9733	
Pipe Tobacco	0.6598		0.3760	
ALCOHOL	0.4137		0.2468	
Binge Alcohol Use ²	0.6614		0.5461	
Heavy Alcohol Use ²	0.3259		0.8692	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.22C Tobacco and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	30	33
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	20	20	17	19
Cigars	32	30	25	25
Pipe Tobacco	12	15	7	9
ALCOHOL	35	34	62	63
Binge Alcohol Use ²	37	35	45	45
Heavy Alcohol Use ²	25	26	17	16

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.22D Tobacco and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	0.14	0.15
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.63	0.65	0.08	0.09
Cigars	0.99	0.99	0.12	0.11
Pipe Tobacco	0.38	0.51	0.03	0.04
ALCOHOL	1.10	1.10	0.29	0.29
Binge Alcohol Use ²	1.17	1.14	0.21	0.20
Heavy Alcohol Use ²	0.79	0.86	0.08	0.07

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.22P Tobacco and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A		0.9529	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.4485		0.6318	
Cigars	0.3277		0.7282	
Pipe Tobacco	0.0847		0.5676	
ALCOHOL	0.1228		0.6615	
Binge Alcohol Use ²	0.1149		0.8999	
Heavy Alcohol Use ²	0.1825		0.8518	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.23C Tobacco and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	57	57
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	47	44	33	34
Cigars	67	65	46	48
Pipe Tobacco	21	21	12	14
ALCOHOL	76	78	123	123
Binge Alcohol Use ²	91	89	115	104
Heavy Alcohol Use ²	83	83	58	62

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.23D Tobacco and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	0.31	0.30
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.37	0.34	0.18	0.18
Cigars	0.53	0.51	0.25	0.25
Pipe Tobacco	0.17	0.16	0.06	0.07
ALCOHOL	0.60	0.61	0.67	0.65
Binge Alcohol Use ²	0.72	0.69	0.62	0.55
Heavy Alcohol Use ²	0.66	0.65	0.32	0.33

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.23P Tobacco and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A		0.9301	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.9316		0.3429	
Cigars	0.2203		0.5960	
Pipe Tobacco	0.0845		0.6419	
ALCOHOL	0.6365		0.0767	
Binge Alcohol Use ²	0.3556		0.3180	
Heavy Alcohol Use ²	0.5925		0.7174	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.24C Tobacco and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	329	334
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	146	132	221	225
Cigars	224	235	233	240
Pipe Tobacco	80	81	139	100
ALCOHOL	424	389	836	809
Binge Alcohol Use ²	384	364	558	497
Heavy Alcohol Use ²	243	250	265	235

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.24D Tobacco and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2002 and 2003

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A	N/A	0.25	0.24
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.32	0.30	0.16	0.17
Cigars	0.49	0.53	0.17	0.18
Pipe Tobacco	0.18	0.18	0.10	0.07
ALCOHOL	0.94	0.87	0.62	0.59
Binge Alcohol Use ²	0.85	0.82	0.42	0.36
Heavy Alcohol Use ²	0.54	0.56	0.20	0.17

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.24P Tobacco and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Drug	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2002	2003	2002	2003
ANY TOBACCO¹	N/A		0.6282	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.9891		0.7168	
Cigars	0.7971		0.9041	
Pipe Tobacco	0.7489		0.3705	
ALCOHOL	0.2621		0.1424	
Binge Alcohol Use ²	0.9908		0.4007	
Heavy Alcohol Use ²	0.5273		0.7861	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.