

Table 7.62C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Drug	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
ANY ILLICIT DRUG²	138	13	136
Marijuana and Hashish	110	11	109
Cocaine	41	3	41
Crack	21	0	21
Heroin	8	1	8
Hallucinogens	27	3	26
LSD	7	0	7
PCP	4	1	4
Ecstasy	22	2	22
Inhalants	16	1	16
Nonmedical Use of Any Psychotherapeutic ³	83	7	83
Pain Relievers	70	7	70
Tranquilizers	44	2	44
Stimulants	37	3	37
Methamphetamine	27	1	27
Sedatives	22	1	22
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	94	8	93

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.62D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Drug	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
ANY ILLICIT DRUG²	0.22	0.54	0.23
Marijuana and Hashish	0.18	0.45	0.19
Cocaine	0.07	0.13	0.07
Crack	0.03	0.02	0.03
Heroin	0.01	0.03	0.01
Hallucinogens	0.04	0.11	0.04
LSD	0.01	0.02	0.01
PCP	0.01	0.06	0.01
Ecstasy	0.04	0.08	0.04
Inhalants	0.03	0.03	0.03
Nonmedical Use of Any Psychotherapeutic ³	0.14	0.31	0.14
Pain Relievers	0.11	0.28	0.12
Tranquilizers	0.07	0.08	0.07
Stimulants	0.06	0.13	0.06
Methamphetamine	0.04	0.04	0.05
Sedatives	0.04	0.06	0.04
ANY ILLICIT DRUG OTHER THAN MARIJUANA²	0.15	0.31	0.16

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

³ Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.63C Any Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	138	13	136
AGE			
15-17	27	3	26
18-25	50	8	49
26-44	119	9	118
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	129	12	127
White	114	10	112
Black or African American	43	6	42
American Indian or Alaska Native	9	*	9
Native Hawaiian or Other Pacific Islander	7	*	7
Asian	22	*	22
Two or More Races	14	*	13
Hispanic or Latino	43	5	43
TRIMESTER²			
First	N/A	9	N/A
Second	N/A	6	N/A
Third	N/A	6	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.63D Any Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	0.22	0.54	0.23
AGE			
15-17	0.44	3.39	0.45
18-25	0.32	0.86	0.34
26-44	0.30	0.64	0.31
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.25	0.63	0.25
White	0.29	0.70	0.29
Black or African American	0.52	2.09	0.54
American Indian or Alaska Native	2.34	*	2.44
Native Hawaiian or Other Pacific Islander	3.89	*	4.01
Asian	0.72	*	0.76
Two or More Races	2.07	*	2.17
Hispanic or Latino	0.46	0.92	0.49
TRIMESTER²			
First	N/A	1.28	N/A
Second	N/A	0.61	N/A
Third	N/A	0.82	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.64C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	110	11	109
AGE			
15-17	24	3	24
18-25	46	7	45
26-44	91	7	91
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	105	10	104
White	92	9	91
Black or African American	38	4	37
American Indian or Alaska Native	8	*	8
Native Hawaiian or Other Pacific Islander	3	*	3
Asian	14	*	13
Two or More Races	12	*	12
Hispanic or Latino	31	4	31
TRIMESTER²			
First	N/A	7	N/A
Second	N/A	5	N/A
Third	N/A	6	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.64D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	0.18	0.45	0.19
AGE			
15-17	0.40	2.91	0.40
18-25	0.30	0.76	0.31
26-44	0.23	0.52	0.24
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.20	0.52	0.21
White	0.23	0.63	0.24
Black or African American	0.46	1.38	0.47
American Indian or Alaska Native	2.02	*	2.15
Native Hawaiian or Other Pacific Islander	1.45	*	1.50
Asian	0.45	*	0.47
Two or More Races	1.82	*	1.92
Hispanic or Latino	0.34	0.81	0.35
TRIMESTER²			
First	N/A	1.05	N/A
Second	N/A	0.53	N/A
Third	N/A	0.81	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.65C Tobacco and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Drug	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
ANY TOBACCO²	232	29	227
Cigarettes	229	29	225
Smokeless Tobacco	22	3	22
Cigars	66	7	64
Pipe Tobacco	19	4	19
ALCOHOL	237	23	231
Binge Alcohol Use ³	192	14	191
Heavy Alcohol Use ³	94	5	94

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.65D Tobacco and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Drug	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
ANY TOBACCO²	0.38	1.21	0.39
Cigarettes	0.37	1.18	0.38
Smokeless Tobacco	0.04	0.14	0.04
Cigars	0.11	0.28	0.11
Pipe Tobacco	0.03	0.16	0.03
ALCOHOL	0.38	0.94	0.39
Binge Alcohol Use ³	0.31	0.58	0.32
Heavy Alcohol Use ³	0.15	0.22	0.16

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.66C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	229	29	225
AGE			
15-17	30	4	29
18-25	71	15	69
26-44	209	22	205
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	213	27	209
White	189	26	185
Black or African American	78	5	78
American Indian or Alaska Native	15	*	14
Native Hawaiian or Other Pacific Islander	10	*	10
Asian	43	*	43
Two or More Races	20	*	20
Hispanic or Latino	75	7	74
TRIMESTER²			
First	N/A	15	N/A
Second	N/A	17	N/A
Third	N/A	17	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.66D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	0.37	1.18	0.38
AGE			
15-17	0.49	4.11	0.49
18-25	0.45	1.58	0.47
26-44	0.52	1.57	0.53
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.41	1.42	0.42
White	0.47	1.80	0.48
Black or African American	0.94	1.67	0.98
American Indian or Alaska Native	3.83	*	3.97
Native Hawaiian or Other Pacific Islander	5.41	*	5.55
Asian	1.41	*	1.50
Two or More Races	3.10	*	3.21
Hispanic or Latino	0.80	1.31	0.85
TRIMESTER²			
First	N/A	2.15	N/A
Second	N/A	1.75	N/A
Third	N/A	2.23	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.67C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	237	23	231
AGE			
15-17	35	3	35
18-25	81	10	79
26-44	216	20	211
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	221	21	214
White	183	18	176
Black or African American	83	4	83
American Indian or Alaska Native	14	*	14
Native Hawaiian or Other Pacific Islander	11	*	11
Asian	66	*	65
Two or More Races	22	*	22
Hispanic or Latino	98	10	98
TRIMESTER²			
First	N/A	17	N/A
Second	N/A	12	N/A
Third	N/A	10	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.67D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	0.38	0.94	0.39
AGE			
15-17	0.58	3.53	0.59
18-25	0.52	1.05	0.54
26-44	0.54	1.43	0.55
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.42	1.07	0.43
White	0.46	1.26	0.46
Black or African American	1.01	1.39	1.05
American Indian or Alaska Native	3.64	*	3.81
Native Hawaiian or Other Pacific Islander	5.63	*	5.81
Asian	2.17	*	2.27
Two or More Races	3.37	*	3.50
Hispanic or Latino	1.05	1.88	1.12
TRIMESTER²			
First	N/A	2.36	N/A
Second	N/A	1.25	N/A
Third	N/A	1.39	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.68C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	192	14	191
AGE			
15-17	30	2	30
18-25	73	6	72
26-44	168	12	167
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	182	13	181
White	153	10	152
Black or African American	71	2	69
American Indian or Alaska Native	15	*	14
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	33	*	32
Two or More Races	16	*	16
Hispanic or Latino	70	3	70
TRIMESTER²			
First	N/A	12	N/A
Second	N/A	5	N/A
Third	N/A	4	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.68D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	0.31	0.58	0.32
AGE			
15-17	0.49	2.32	0.50
18-25	0.47	0.66	0.49
26-44	0.42	0.88	0.43
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.35	0.70	0.36
White	0.39	0.74	0.40
Black or African American	0.86	0.81	0.88
American Indian or Alaska Native	3.75	*	3.96
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	1.09	*	1.11
Two or More Races	2.47	*	2.60
Hispanic or Latino	0.75	0.66	0.80
TRIMESTER²			
First	N/A	1.75	N/A
Second	N/A	0.50	N/A
Third	N/A	0.49	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.69C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	94	5	94
AGE			
15-17	13	0	13
18-25	42	2	41
26-44	78	5	78
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	90	5	89
White	80	5	79
Black or African American	31	1	31
American Indian or Alaska Native	8	*	7
Native Hawaiian or Other Pacific Islander	4	*	4
Asian	22	*	22
Two or More Races	8	*	8
Hispanic or Latino	24	1	23
TRIMESTER²			
First	N/A	4	N/A
Second	N/A	*	N/A
Third	N/A	1	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 7.69D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002 and 2003 NSDUHs

Demographic Characteristic	Total ¹	PREGNANCY STATUS	
		Pregnant	Not Pregnant
TOTAL	0.15	0.22	0.16
AGE			
15-17	0.22	0.09	0.22
18-25	0.27	0.26	0.28
26-44	0.20	0.35	0.20
HISPANIC ORIGIN AND RACE			
Not Hispanic or Latino	0.17	0.28	0.18
White	0.20	0.34	0.21
Black or African American	0.37	0.26	0.39
American Indian or Alaska Native	2.05	*	1.95
Native Hawaiian or Other Pacific Islander	1.90	*	1.93
Asian	0.71	*	0.75
Two or More Races	1.16	*	1.22
Hispanic or Latino	0.25	0.19	0.27
TRIMESTER²			
First	N/A	0.54	N/A
Second	N/A	*	N/A
Third	N/A	0.08	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.