

REVIEWED LABELS

Drug Facts (continued)

Other information ● each caplet contains: sodium 20 mg ● store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-800-331-4536 or www.midol.com

24 CAPLETS*
capsule-shaped tablets

Cramps • Backache • Headache

MAXIMUM STRENGTH

Extended Relief
for up to 12 HOURS

NAPROXEN SODIUM TABLETS, 220 mg PAIN RELIEVER

Midol

F.P.O.

| | | | |
|---|----------------|---|--|
| Drug Facts | | Drug Facts (continued) | |
| Active ingredient (in each caplet) | Purpose | Warnings | |
| Naproxen sodium 220 mg (naproxen 200 mg)..... | Pain reliever | Allergy alert: Naproxen sodium may cause a severe allergic reaction which may include: ● hives ● facial swelling ● asthma (wheezing) ● shock | |
| Uses temporarily relieves minor aches and pains due to ● menstrual cramps ● muscle aches ● headache ● backache | | | |

| | | | | | | | |
|---|--|----------------|----------------|----------------|---|-------------|---|
| Drug Facts (continued) | Drug Facts (continued) | | | | | | |
| <p>Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take naproxen sodium or other pain relievers/fever reducers. Naproxen sodium may cause stomach bleeding.</p> <p>Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer</p> <p>Ask a doctor before use if you have ever had serious side effects from any pain reliever</p> <p>Ask a doctor or pharmacist before use if you are ● under a doctor's care for any serious condition ● taking any other drug ● taking any other product that contains naproxen sodium, or any other pain reliever/fever reducer</p> <p>Stop use and ask a doctor if ● an allergic reaction occurs. Seek medical help right away ● pain gets worse or lasts more than 10 days ● you have difficulty swallowing ● it feels like the pill is stuck in your throat ● you develop heartburn ● stomach pain occurs or lasts, even if symptoms are mild ● redness or swelling is present in the painful area ● any new symptoms appear</p> | <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ● do not take more than directed ● drink a full glass of water with each dose</p> <table border="1"> <tr> <td>under 12 years</td> <td>● ask a doctor</td> </tr> <tr> <td>12 to 65 years</td> <td>● take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. The smallest effective dose should be used. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period.</td> </tr> <tr> <td>over age 65</td> <td>● do not take more than 1 caplet every 12 hours unless directed by a doctor</td> </tr> </table> | under 12 years | ● ask a doctor | 12 to 65 years | ● take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. The smallest effective dose should be used. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. | over age 65 | ● do not take more than 1 caplet every 12 hours unless directed by a doctor |
| under 12 years | ● ask a doctor | | | | | | |
| 12 to 65 years | ● take 1 caplet every 8 to 12 hours while symptoms last. For the first dose you may take 2 caplets within the first hour. The smallest effective dose should be used. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period. | | | | | | |
| over age 65 | ● do not take more than 1 caplet every 12 hours unless directed by a doctor | | | | | | |

12

MAXIMUM STRENGTH
Midol® Extended Relief
Dist. by: Bayer HealthCare LLC

READ CARTON BEFORE USE. Each caplet contains: Naproxen sodium 220 mg 8875440

SEPARATE, PEEL PAPER, PUSH THROUGH FOIL

EPR-4/07 UOE: 287624D