

FPO 128 code

FPO 128 code

Drug Facts

Active ingredient (in each tablet) Purpose
Omeprazole magnesium delayed-release tablet 20.6 mg (equivalent to 20 mg omeprazole).....Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- digoxin (heart medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Drug Facts (continued)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not chew or crush the tablets
- do not crush tablets in food
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor

Other information

- read the directions, warnings and package insert before use

Drug Facts (continued)

- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polysorbate 80, polyvinylpyrrolidone, sodium stearyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate

Questions? 1-800-289-9181

Two 14-day courses of treatment
28 TABLETS

omeprazole delayed-release tablets
20 mg / acid reducer
OTC®

Prilosec

Treats Frequent Heartburn!

NDC 37000-455-03

Prilosec OTC

Prilosec OTC

Dist. by Preter & Gamble, Cincinnati, OH 45202
Product of Sweden 00000000

Safety Feature—
Do not use
if tablet blister
unit is open
or torn.

Prilosec OTC

100% SIZE
12 digit UPC
(non-suppressed)
For Position Only
37000 455906 7