

7905119

MIDAMOR® (Amiloride HCl)

hypokalemia has been documented and only with careful titration of the dose and close monitoring of serum electrolytes.

CONTRAINDICATIONS**Hyperkalemia**

MIDAMOR should not be used in the presence of elevated serum potassium levels (greater than 5.5 mEq per liter).

Antikaliuretic Therapy or Potassium Supplementation

MIDAMOR should not be given to patients receiving other potassium-conserving agents, such as spironolactone or triamterene. Potassium supplementation in the form of medication, potassium-containing salt substitutes or a potassium-rich diet should not be used with MIDAMOR except in severe and/or refractory cases of hypokalemia. Such concomitant therapy can be associated with rapid increases in serum potassium levels. If potassium supplementation is used, careful monitoring of the serum potassium level is necessary.

Impaired Renal Function

Anuria, acute or chronic renal insufficiency, and evidence of diabetic nephropathy are contraindications to the use of MIDAMOR. Patients with evidence of renal functional impairment (blood urea nitrogen [BUN] levels over 30 mg per 100 mL or serum creatinine levels over 1.5 mg per 100 mL) or diabetes mellitus should not receive the drug without careful, frequent and continuing monitoring of serum electrolytes, creatinine, and BUN levels. Potassium retention associated with the use of an antikaliuretic agent is accentuated in the presence of renal impairment and may result in the rapid development of hyperkalemia.

Hypersensitivity

MIDAMOR is contraindicated in patients who are hypersensitive to this product.

WARNINGS**Hyperkalemia**

Like other potassium-conserving agents, amiloride may cause hyperkalemia (serum potassium levels greater than 5.5 mEq per liter) which, if uncorrected, is potentially fatal. Hyperkalemia occurs commonly (about 10%) when amiloride is used without a kaliuretic diuretic. This incidence is greater in patients with renal impairment, diabetes mellitus (with or without recognized renal insufficiency), and in the elderly. When MIDAMOR is used concomitantly with a thiazide diuretic in patients without these complications, the risk of hyperkalemia is reduced to about 1-2 percent. It is thus essential to monitor serum potassium levels carefully in any patient receiving amiloride, particularly when it is first introduced, at the time of diuretic dosage adjustments, and during any illness that could affect renal function.

The risk of hyperkalemia may be increased when potassium-conserving agents, including MIDAMOR, are administered concomitantly with an angiotensin-converting enzyme inhibitor, an angiotensin II receptor antagonist, cyclosporine or tacrolimus. (See PRECAUTIONS, *Drug Interactions*.) Warning signs or symptoms of hyperkalemia include paresthesias, muscular weakness, fatigue, flaccid paralysis of the extremities, bradycardia, shock, and ECG abnormalities. Monitoring of the serum potassium level is essential because mild hyperkalemia is not usually associated with an abnormal ECG.

When abnormal, the ECG in hyperkalemia is characterized primarily by tall, peaked T waves or elevations from previous tracings. There may also be lowering of the R wave and increased depth of the S wave, widening and even disappearance of the P wave, progressive widening of the QRS complex, prolongation of the PR interval, and ST depression.

Treatment of hyperkalemia: If hyperkalemia occurs in patients taking MIDAMOR, the drug should be discontinued immediately. If the serum potassium level exceeds 6.5 mEq per liter, active measures should be taken to reduce it. Such measures include the intravenous administration of sodium bicarbonate solution or oral or parenteral glucose with a rapid-acting insulin preparation. If needed, a cation exchange resin such as sodium polystyrene sulfonate may be given orally or by enema. Patients with persistent hyperkalemia may require dialysis.

Diabetes Mellitus

In diabetic patients, hyperkalemia has been reported with the use of all potassium-conserving diuretics, including MIDAMOR, even in patients without evidence of diabetic nephropathy. Therefore, MIDAMOR should be avoided, if possible, in diabetic patients and, if it is used, serum electrolytes and renal function must be monitored frequently.

MIDAMOR should be discontinued at least three days before glucose tolerance testing.

Metabolic or Respiratory Acidosis

Antikaliuretic therapy should be instituted only with caution in severely ill patients in whom respiratory or metabolic acidosis may occur, such as patients with cardiopulmonary disease or poorly controlled diabetes. If MIDAMOR is given to these patients, frequent monitoring of acid-base balance is necessary. Shifts in acid-base balance alter the ratio of extracellular/intracellular potassium, and the development of acidosis may be associated with rapid increases in serum potassium levels.

MIDAMOR® (Amiloride HCl)

PRECAUTIONS**General****Electrolyte Imbalance and BUN Increases**

Hyponatremia and hypochloremia may occur when MIDAMOR is used with other diuretics and increases in BUN levels have been reported. These increases usually have accompanied vigorous fluid elimination, especially when diuretic therapy was used in seriously ill patients, such as those who had hepatic cirrhosis with ascites and metabolic alkalosis, or those with resistant edema. Therefore, when MIDAMOR is given with other diuretics to such patients, careful monitoring of serum electrolytes and BUN levels is important. In patients with pre-existing severe liver disease, hepatic encephalopathy, manifested by tremors, confusion, and coma, and increased jaundice, have been reported in association with diuretics, including amiloride HCl.

Drug Interactions

When amiloride HCl is administered concomitantly with an angiotensin-converting enzyme inhibitor, an angiotensin II receptor antagonist, cyclosporine or tacrolimus, the risk of hyperkalemia may be increased. Therefore, if concomitant use of these agents is indicated because of demonstrated hypokalemia, they should be used with caution and with frequent monitoring of serum potassium. (See WARNINGS.)

Lithium generally should not be given with diuretics because they reduce its renal clearance and add a high risk of lithium toxicity. Read circulars for lithium preparations before use of such concomitant therapy.

In some patients, the administration of a non-steroidal anti-inflammatory agent can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing and thiazide diuretics. Therefore, when MIDAMOR and non-steroidal anti-inflammatory agents are used concomitantly, the patient should be observed closely to determine if the desired effect of the diuretic is obtained. Since indomethacin and potassium-sparing diuretics, including MIDAMOR, may each be associated with increased serum potassium levels, the potential effects on potassium kinetics and renal function should be considered when these agents are administered concurrently.

Carcinogenicity, Mutagenicity, Impairment of Fertility

There was no evidence of a tumorigenic effect when amiloride HCl was administered for 92 weeks to mice at doses up to 10 mg/kg/day (25 times the maximum daily human dose). Amiloride HCl has also been administered for 104 weeks to male and female rats at doses up to 6 and 8 mg/kg/day (15 and 20 times the maximum daily dose for humans, respectively) and showed no evidence of carcinogenicity.

Amiloride HCl was devoid of mutagenic activity in various strains of *Salmonella typhimurium* with or without a mammalian liver microsomal activation system (Ames test).

Pregnancy

Pregnancy Category B. Teratogenicity studies with amiloride HCl in rabbits and mice given 20 and 25 times the maximum human dose, respectively, revealed no evidence of harm to the fetus, although studies showed that the drug crossed the placenta in modest amounts. Reproduction studies in rats at 20 times the expected maximum daily dose for humans showed no evidence of impaired fertility. At approximately 5 or more times the expected maximum daily dose for humans, some toxicity was seen in adult rats and rabbits and a decrease in rat pup growth and survival occurred.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Studies in rats have shown that amiloride is excreted in milk in concentrations higher than those found in blood, but it is not known whether MIDAMOR is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from MIDAMOR, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

Clinical studies of MIDAMOR did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function. (See CONTRAINDICATIONS, *Impaired Renal Function*.)

7905119

MIDAMOR® (Amiloride HCl)

ADVERSE REACTIONS

MIDAMOR is usually well tolerated and, except for hyperkalemia (serum potassium levels greater than 5.5 mEq per liter — see WARNINGS), significant adverse effects have been reported infrequently. Minor adverse reactions were reported relatively frequently (about 20%) but the relationship of many of the reports to amiloride HCl is uncertain and the overall frequency was similar in hydrochlorothiazide treated groups. Nausea/anorexia, abdominal pain, flatulence, and mild skin rash have been reported and probably are related to amiloride. Other adverse experiences that have been reported with amiloride are generally those known to be associated with diuresis, or with the underlying disease being treated.

The adverse reactions for MIDAMOR listed in the following table have been arranged into two groups: (1) incidence greater than one percent; and (2) incidence one percent or less. The incidence for group (1) was determined from clinical studies conducted in the United States (837 patients treated with MIDAMOR). The adverse effects listed in group (2) include reports from the same clinical studies and voluntary reports since marketing. The probability of a causal relationship exists between MIDAMOR and these adverse reactions, some of which have been reported only rarely.

<i>Incidence >1%</i>	<i>Incidence ≤1%</i>
Body as a Whole	
Headache**	Back pain
Weakness	Chest pain
Fatigability	Neck/shoulder ache
	Pain, extremities
Cardiovascular	
None	Angina pectoris
	Orthostatic hypotension
	Arrhythmia
	Palpitation
Digestive	
Nausea/anorexia**	Jaundice
Diarrhea**	GI bleeding
Vomiting**	Abdominal fullness
Abdominal pain	GI disturbance
Gas pain	Thirst
Appetite changes	Heartburn
Constipation	Flatulence
	Dyspepsia
Metabolic	
Elevated serum potassium levels (>5.5 mEq per liter)***	None
Skin	
None	Skin rash
	Itching
	Dryness of mouth
	Pruritus
	Alopecia
Musculoskeletal	
Muscle cramps	Joint pain
	Leg ache
Nervous	
Dizziness	Paresthesia
Encephalopathy	Tremors
	Vertigo
Psychiatric	
None	Nervousness
	Mental confusion
	Insomnia
	Decreased libido
	Depression
	Somnolence
Respiratory	
Cough	Shortness of breath
Dyspnea	
Special Senses	
None	Visual disturbances
	Nasal congestion
	Tinnitus
	Increased intraocular pressure
Urogenital	
Impotence	Polyuria
	Dysuria
	Urinary frequency
	Bladder spasms
	Gynecomastia

**Reactions occurring in 3% to 8% of patients treated with MIDAMOR. (Those reactions occurring in less than 3% of the patients are unmarked.)

***See WARNINGS.

Causal Relationship Unknown

Other reactions have been reported but occurred under circumstances where a causal relationship could not be established. However, in these rarely reported events, that possibility cannot be excluded. Therefore, these observations are listed to serve as alerting information to physicians.

- Activation of probable pre-existing peptic ulcer
- Aplastic anemia
- Neutropenia
- Abnormal liver function

MIDAMOR® (Amiloride HCl)

OVERDOSAGE

No data are available in regard to overdosage in humans.

The oral LD₅₀ of amiloride hydrochloride (calculated as the base) is 56 mg/kg in mice and 36 to 85 mg/kg in rats, depending on the strain.

It is not known whether the drug is dialyzable.

The most likely signs and symptoms to be expected with overdosage are dehydration and electrolyte imbalance. These can be treated by established procedures. Therapy with MIDAMOR should be discontinued and the patient observed closely. There is no specific antidote. Emesis should be induced or gastric lavage performed. Treatment is symptomatic and supportive. If hyperkalemia occurs, active measures should be taken to reduce the serum potassium levels.

DOSAGE AND ADMINISTRATION

MIDAMOR should be administered with food.

MIDAMOR, one 5 mg tablet daily, should be added to the usual antihypertensive or diuretic dosage of a kaliuretic diuretic. The dosage may be increased to 10 mg per day, if necessary. More than two 5 mg tablets of MIDAMOR daily usually are not needed, and there is little controlled experience with such doses. If persistent hypokalemia is documented with 10 mg, the dose can be increased to 15 mg, then 20 mg, with careful monitoring of electrolytes.

In treating patients with congestive heart failure after an initial diuresis has been achieved, potassium loss may also decrease and the need for MIDAMOR should be re-evaluated. Dosage adjustment may be necessary. Maintenance therapy may be on an intermittent basis.

If it is necessary to use MIDAMOR alone (see INDICATIONS), the starting dosage should be one 5 mg tablet daily. This dosage may be increased to 10 mg per day, if necessary. More than two 5 mg tablets usually are not needed, and there is little controlled experience with such doses. If persistent hypokalemia is documented with 10 mg, the dose can be increased to 15 mg, then 20 mg, with careful monitoring of electrolytes.

HOW SUPPLIED

No. 3381 — Tablets MIDAMOR, 5 mg, are yellow, diamond-shaped, compressed tablets, coded MSD 92 on one side and MIDAMOR on the other. They are supplied as follows:

NDC 0006-0092-68 bottles of 100.


Storage

Protect from moisture, freezing and excessive heat.

MERCK & CO., INC., Whitehouse Station, NJ 08889, USA

Issued November 2002

Printed in USA



7905119

