

PATIENT INFORMATION

DETROL® (DE-trol)

(tolterodine tartrate tablets)

Read the Patient Information that comes with DETROL before you start using it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your condition or your treatment. Only your doctor can determine if treatment with DETROL is right for you.

What is DETROL?

DETROL is a prescription medicine for **adults** used to treat the following symptoms due to a condition called **overactive bladder**:

- having a strong need to urinate with leaking or wetting accidents (urge urinary incontinence)
- having a strong need to urinate right away (urgency)
- having to urinate often (frequency)

DETROL LA (tolterodine tartrate extended release capsules) did not help the symptoms of overactive bladder when studied in children.

What is overactive bladder?

Overactive bladder happens when you cannot control your bladder muscle. When the muscle contracts too often or cannot be controlled, you get symptoms of overactive bladder, which are leakage of urine (urge urinary incontinence), needing to urinate right away (urgency), and needing to urinate often (frequency).

Who should not take DETROL?

Do not take DETROL if:

- you have trouble emptying your bladder (also called “urinary retention”)
- your stomach empties slowly (also called “gastric retention”)
- you have an eye problem called “uncontrolled narrow-angle glaucoma”
- you are allergic to DETROL or to any of its ingredients. See the end of this leaflet for a complete list of ingredients

What should I tell my doctor before starting DETROL?

Before starting DETROL, tell your doctor about all of your medical conditions, including if you:

- have any stomach or intestinal problems
- have trouble emptying your bladder or you have a weak urine stream
- have an eye problem called narrow-angle glaucoma
- have liver problems
- have kidney problems
- **have a condition called myasthenia gravis**
- or any family members have a rare heart condition called QT prolongation (long QT syndrome)
- are pregnant or trying to become pregnant. It is not known if DETROL could harm your unborn baby
- are breastfeeding. It is not known if DETROL passes into your milk and if it can harm your child

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins and herbal supplements. Other drugs can affect how your body handles DETROL. Your doctor may use a lower dose of DETROL if you are taking:

- Certain medicines for fungus or yeast infections such as Nizoral® (ketoconazole), Sporanox® (itraconazole), and Monistat® (miconazole)
- Certain medicines for bacteria infections such as Biaxin® (clarithromycin), and erythromycin
- Sandimmune® (cyclosporine) or Velban® (vinblastine)

Know the medicines you take. Keep a list of them with you to show your doctor or pharmacist each time you get a new medicine.

How should I take DETROL?

- Take DETROL exactly as prescribed. Your doctor will prescribe the dose that is right for you. Do not change your dose unless told to do so by your doctor.
- Take DETROL tablets two times a day.
- DETROL can be taken with or without food.
- Take DETROL the same time each day.
- If you miss a dose of DETROL, begin taking DETROL again the next day. Do not take 2 doses of DETROL in the same day.
- If you took more than your prescribed dose of DETROL, call your doctor, or poison control center, or go to the hospital emergency room.

What are possible side effects of DETROL?

The most common side effects with DETROL are:

- dry mouth
- headache
- constipation
- dizziness
- stomach pain

Medicines like DETROL can cause blurred vision or drowsiness.

Use caution while driving or doing other dangerous activities until you know how DETROL affects you.

These are not all the side effects with DETROL. For a complete list, ask your doctor or pharmacist.

How do I store DETROL?

- Store DETROL at room temperature (59 to 86° F). Keep it in a dry place.
- **Keep DETROL and all medicines out of the reach of children.**

General Information about DETROL

Medicines are sometimes prescribed for conditions that are not in the patient information leaflet. Only use DETROL the way your doctor tells you. Do not share it with other people even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about DETROL. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about DETROL that is written for health professionals. You can also visit www.DETROL.com on the Internet, or call 1-888-4-DETROL (1-888-433-8765).

What are the ingredients in DETROL?

Active ingredients: tolterodine tartrate

Inactive ingredients: colloidal anhydrous silica, calcium hydrogen phosphate dihydrate, cellulose microcrystalline, hypromellose, magnesium stearate, sodium starch glycolate (pH 3.0 to 5.0), stearic acid, and titanium dioxide.

Distributed by:



Pharmacia & Upjohn
Division of Pfizer Inc, NY, NY 10017
March 2008

Registered trademarks are the property of their respective owners.

LAB-0329-3.0