



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez

Display

Abstract

Show:

20

Sort

Send to

Text

Text Version

1: Ned Tijdschr Geneeskd. 2002 Apr 27;146(17):811-2.

Related Articles

[Adverse effects after switching to a different generic form of paroxetine paroxetine mesylate instead of paroxetine HCl hemihydrate]

[Article in Dutch]

Vergouwen AC, Bakker A.

Sint Lucas Andreas Ziekenhuis, locatie Lucas, afd. Psychiatrie, Postbus 9243, 1006 A Amsterdam. a.vergouwen@slaz.nl

Two patients, a 29-year-old man and a 32-year-old woman who were being treated with paroxetine HCl hemihydrate for depression and panic disorder with agoraphobia, respectively, unexpectedly developed complaints after switching to paroxetine mesylate. The symptoms were generalised itching with relapse of the depressive disorder in the man and nausea with diarrhoea in the woman. The symptoms disappeared after the original treatment with paroxetine HCl hemihydrate was resumed. In the first patient, a rechall with the mesylate form of paroxetine led to the same complaints which subsided again using the first product. Those treating such patients, and the patients themselves, should be informed if the delivery form of paroxetine is changed.

PMID: 12014241 [PubMed - indexed for MEDLINE]

Display

Abstract

Show:

20

Sort

Send to

Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)