

Appendix

Table A1. Concentration Ranges Used for Susceptibility Testing of *Salmonella* and *E. coli*, 2004

Antimicrobial Class	Antimicrobial Agent	Concentration Range ($\mu\text{g/ml}$)
Aminoglycosides	Amikacin	0.5 - 64
	Gentamicin	0.25 - 16
	Kanamycin	8 - 64
	Streptomycin	32 - 64
Aminopenicillins	Ampicillin	1 - 32
β-Lactam/β-Lactamase Inhibitor Combinations	Amoxicillin–Clavulanic Acid	1 / 0.5 - 32 / 16
Cephalosporins	Ceftiofur	0.12 - 8
	Ceftriaxone	0.25 - 64
Cephamycins	Cefoxitin	0.5 - 32
Folate Pathway Inhibitors	Sulfisoxazole	16 - 256
	Trimethoprim–Sulfamethoxazole	0.12 / 2.4 - 4 / 76
Phenicols	Chloramphenicol	2 - 32
Quinolones	Ciprofloxacin	0.015 - 4
	Nalidixic acid	0.5 - 32
Tetracyclines	Tetracycline	4 - 32

Table A2. Concentration Ranges Used for Susceptibility Testing¹ of *Campylobacter*, 2004

Antimicrobial Class	Antimicrobial Agent	Concentration Range (µg/ml)	
		Etest®	Broth Microdilution
Aminoglycosides	Gentamicin	0.016 - 256	0.12 - 32
Ketolides	Telithromycin	N/A	0.015 - 8
Lincosamides	Clindamycin	0.016 - 256	0.03 - 16
Macrolides	Azithromycin	0.016 - 256	0.015 - 64
	Erythromycin	0.016 - 256	0.03 - 64
Phenicols	Chloramphenicol	0.016 - 256	N/A
	Florfenicol	N/A	0.03 - 64
Quinolones	Ciprofloxacin	0.002 - 32	0.015 - 64
	Nalidixic acid	0.016 - 256	4 - 64
Tetracyclines	Tetracycline	0.016 - 256	0.06 - 64

¹ Etest® was used to test human and food animal isolates while a broth microdilution method was used to test retail meat isolates