

Wind Direction and Its Linkage with *Vibrio cholerae* Dissemination

Shlomit Paz¹ and Meir Broza²

¹Department of Geography and Environmental Studies, University of Haifa, Haifa, Israel; ²Department of Biology, Faculty of Science and Science Education, University of Haifa, Oranim, Tivon, Israel

BACKGROUND: The relevance of climatic events as causative factors for cholera epidemics is well known. However, examinations of the involvement of climatic factors in intracontinental disease distribution are still absent.

OBJECTIVES: The spreading of cholera epidemics may be related to the dominant wind direction over land.

METHODS: We examined the geographic diffusion of three cholera outbreaks through their linkage with the wind direction: *a*) the progress of *Vibrio cholerae* O1 biotype El Tor in Africa during 1970–1971 and *b*) again in 2005–2006; and *c*) the rapid spread of *Vibrio cholerae* O139 over India during 1992–1993. We also discuss the possible influence of the wind direction on windborn dissemination by flying insects, which may serve as vectors.

RESULTS: Analysis of air pressure data at sea level and at several altitudes over Africa, India, and Bangladesh show a correspondence between the dominant wind direction and the intracontinental spread of cholera.

CONCLUSIONS: We explored the hypothesis that winds have assisted the progress of cholera *Vibrios* throughout continents. The current analysis supports the hypothesis that aeroplankton (the tiny life forms that float in the air and that may be caught and carried upward by the wind, landing far from their origin) carry the cholera bacteria from one body of water to an adjacent one. This finding may improve our understanding of how climatic factors are involved in the rapid distribution of new strains throughout a vast continental area. Awareness of the aerial transfer of *Vibrio cholerae* may assist health authorities by improving the prediction of the disease's geographic dissemination.

KEY WORDS: Africa, Chironomidae, India, *Vibrio cholerae*, wind. *Environ Health Perspect* 115:195–200 (2007). doi:10.1289/ehp.9391 available via <http://dx.doi.org/> [Online 18 October 2006]

Cholera continues to be an important cause of morbidity and mortality in many areas of the world, and there is currently a high frequency of new outbreaks in Africa. The fatal effects of the disease are mainly due to the cholera toxin produced by specific strains of *Vibrio cholerae*.

In 1817, cholera spread out of its endemic distribution on the Indian subcontinent and reached many countries around the world, appearing as a global event, or pandemic. In 1961, *Vibrio cholerae* O1 biotype El Tor spread out of Indonesia toward India, replacing the O1 “classical biotype,” and creating the “seventh pandemic.” During summer 1970, it continued to spread westward through the Middle East into Africa. The seventh pandemic began in Africa in August 1970 on two fronts, simultaneously striking the north and the west, with devastating consequences. Cholera quickly spread throughout the continent. Inadequate resources and limited access to health care led to high death rates (Swerdlow and Isaäcson 1994).

In October 1992 a new serogroup, now known as O139, caused a severe outbreak of cholera in the city of Madras, in southeast India. Within 10 months, the O139 serogroup had disseminated all over the Indian subcontinent and soon thereafter spread to 11 neighboring countries in Asia and beyond, temporarily replacing the O1 serogroup as the most common (Ramamurthy et al. 1993).

What could enable such rapid distribution of the *Vibrio cholerae* O139 throughout the subcontinent? The answer is still a mystery.

While maps and reports describing the El Tor pandemic of 1970 showed its spread on a continental and global scale [e.g., Todar 2005; World Health Organization (WHO) 2000], researchers in India were able to track the replacement of the O1 by O139 serogroup in town after town, over a period of 10 months (Nair et al. 1994). However, both the spread of El Tor in 1970–1971 and O139 in 1992–1993 show a tendency for wave after wave of gradual westward movement. Although person-to-person contact plays an important role in the rapid spread of cholera, especially in crowded religious festivals and on all types of modern public transportation, these pandemics included cases that cannot be explained by any human activity or mechanical transportation. In some other cases, epidemics appeared simultaneously in remote unconnected locations.

Chironomids, the nonbiting midges (Diptera; Chironomidae), are one of the most abundant macroinvertebrate groups that exist in most freshwater aquatic habitats (Armitage et al. 1995), especially in estuarine (Benke 1998) and organic-rich water bodies typical to *Vibrio cholerae*.

The eggs, larvae, and pupae are aquatic; adults emerge into the air for mating and

dispersion. Chironomid adults are highly mobile, present in very large numbers, and found frequently in the air at various elevations, as was noted by Glick (1939) and more recently by Reynolds et al. (1999) (see also Bowden and Johnson 1976; Sparks et al. 1986).

Broza and Halpern (2001) reported that the huge number of chironomid egg masses attached to hard substrates on the water surface serve as a reservoir for the free-living *Vibrio* bacteria. Also, Broza et al. (2005) observed that both male and female chironomids, while emerging from the water, may carry cholera bacteria into the mating swarms. In experimental simulation, they demonstrated that the cholera-bearing adult midges are carried by the wind, and transmit the bacteria from one body of water to another. Broza et al. (2005) suggested that flying adult midges can act as a vector for cholera in continental and even intercontinental spread. However, this hypothesis has not yet been tested in a true cholera epidemic or pandemic.

Much information is available regarding the relevance of climatic events as causative factors for cholera epidemics. Several studies (e.g., Huq et al. 2005; Lipp et al. 2002) focused on the effect of the main climatic factors on the initial eruption of cholera, such as precipitation, air and water-surface temperature, and relative humidity. These studies did not, however, address the climatic connection to disease dissemination over vast continental areas.

Because winds are the driving force behind the long-distance flight of insects, we suggest exploring the linkage between wind direction on a local and global scale and the sequence of epidemic dissemination. In the present study, we examine three cases of cholera outbreaks

Address correspondence to S. Paz, Department of Geography and Environmental Studies, University of Haifa, Haifa, Israel 31905. Telephone: 972-4-8249617. Fax: 972-4-8249605. E-mail: shlomit@geo.haifa.ac.il

We thank B.G. Nair (Director of Laboratory Sciences Division, Center for Health and Population, Dhaka, Bangladesh) for reading the manuscript and supporting our conclusions. We also thank R. Fluss (Statistics Consulting Unit, University of Haifa) for statistical advice and N. Yoselevich for help with preparing the final figures.

This study was partially supported by the Israel Science Foundation.

The authors declare they have no competing financial interests.

Received 1 June 2006; accepted 18 October 2006.

that occurred in Africa and on the Indian subcontinent within the last 35 years, focusing on linkages between dominant wind direction and the disease dissemination.

The African monsoonal winds are ultimately caused by the differential heating of land and ocean, which inevitably leads to a pressure gradient. Atmospheric circulation transports very different types of air masses, whose presence results in the occurrence of diametrically different wet and dry seasons. The circulation over West Africa is typically monsoonal with seasonal change in wind direction. The humid Guinea monsoon blowing from the southwest moves northward behind the convergence zone, and by April it reaches the borders of Mali and Niger. In eastern Africa, the seasonal exchange of air masses flow from opposite directions, from parts of continental Africa in the northern and southern hemispheres. In April, Indian Ocean trade winds blow from the sea to the coastline and the mountain ranges. In East and West Africa and in the interior areas of the equatorial zone, the first of two annual precipitation peaks occurs in April and May, whereas

the second peak occurs in October and November with lower amounts of precipitation [Intergovernmental Panel on Climate Change (IPCC) 1996; Martyn 1992].

Climatic patterns in southern Asia depend primarily on the geographic position in tropical and equatorial latitudes. The Indian climate is dominated by the monsoon, which shows spatial, interannual, and intraseasonal variability. The monsoon is characterized by a reversal in the winds over the Indian subcontinent between summer and winter. In winter the winds blow off the continent to the southwest, and in summer (starting in May) they blow from the southeast. The wind shift is associated with heavy rains over India in the summer. After the withdrawal of the summer monsoon, the northeast monsoon sets in by November and continues through December. Due to very active weather systems in the Bay of Bengal, many cyclones occur during this season, mainly affecting the eastern coast of India [Martyn 1992; National Oceanic and Atmospheric Administration (NOAA) and Climate Diagnostic Center 2004].

We hypothesize that a potential aerial transfer of *Vibrio cholerae* over vast continental areas is related to a correlation between spatial incidence patterns and wind direction. In the present study we evaluated three cholera outbreaks within the seventh and the eighth pandemics: the disease dissemination over parts of Africa throughout 1970–1971; the disease dissemination in the same areas in 2005–2006; and the *Vibrio cholerae* O139 rapid spread over the Indian subcontinent during 1992–1993. We evaluated the dissemination of these epidemics through wind direction analysis (the dominant wind direction in a selected region computed based on monthly synoptic maps), examining the possible influence of wind direction on flying insects carrying viable cholera bacteria attached to their cuticle.

In this study, we also investigated the hypothesis of Broza et al. (2005) that aeroplankton may carry the cholera bacteria from one body of water to an adjacent one. Aeroplankton are the tiny life forms that drift through the air, are carried upward by the wind into the atmosphere, and continue to be carried by the wind for a considerable length of time, landing far from their origin. Small organisms such as flies may be part of the aeroplankton. They may be driven by strong winds in lower altitudes during their swarming behavior at a height of 5–100 m.

Methods

Data for Africa, 1970–1971. We obtained data for cholera dissemination in Africa for 1970–1971 from Swerdlow and Isaacson (1994) and for October 2004 to March 2006 from the World Health Organization (WHO 2005, 2006). To create wind direction maps for Africa [using National Centers for Environmental Prediction–National Center for Atmospheric Research (NCEP–NCAR) Reanalysis Project database and software (NOAA NCEP–NCAR 2006)], we used mean monthly sea level pressure [SLP; in millibars (mb)] and mean monthly geopotential values for three altitude levels: 1,000 mb (150 to ~200 m), 925 mb (700 to ~800 m), and 850 mb (~1,500 m) covering the whole African continent (NOAA NCEP–NCAR 2005). These values were measured for each month from July 1970, one month before the West African cholera epidemic appeared in Guinea, until December 1971, the last month of the epidemic in Africa (NOAA NCEP–NCAR 2005). The altitude levels are in accordance with those examined for aeroplankton by Glick (1939).

Data for Africa, 2005–2006. We also mapped the same SLP, 1,000 mb, 925 mb, and 850 mb levels (NOAA NCEP–NCAR 2005) for Africa for each month from October 2004, when *Vibrio cholerae* O1 El Tor was




Figure 1. Maps showing movement of cholera in Africa during 1970–1971 (adapted from Swerdlow and Isaacson 1994) (A) and the dominant wind direction in Africa in SLP (in millibars) in August 1970 (B), air pressure of 925 mb level in September 1970 (C), and SLP in November 1970 (D). Black arrows in (A) indicate other waves of cholera movement. Values in (C) indicate altitude in meters.

confirmed in Senegal, to March 2006 (cholera has been reported in southern Sudan).

Data for India. Our study of India was based on a summary of the temporal and spatial spread of O139 during 1992–1993 by the National Institute of Cholera and Enteric Diseases in Calcutta, India (Nair et al. 1994). We analyzed mean monthly SLP and mean monthly geopotential values for 1,000 mb, 925 mb, and 850 mb levels, covering the subcontinent of India and Bangladesh for each of the 10 months from October 1992 (when the *Vibrio cholerae* serogroup O139 caused an outbreak in Madras) to July 1993 (when the disease reached northwestern India). We produced wind-direction maps using the NCEP-NCAR Reanalysis Project database (NOAA NCEP-NCAR 2006). Anomalies from the monthly averages, based on the 1951–2000 period, were also mapped and analyzed.

Correlations. In the present study, the dominant wind direction refers to the prevailing winds over a selected area during a chosen month using monthly average pressure maps at different altitudes. The main direction was identified based on the principles for the Northern Hemisphere—that the wind blows counterclockwise around a low-pressure area and clockwise around a high-pressure area. Although the wind behavior is characterized by a rapid change in direction and speed, the maps present the most expectable wind direction throughout the month.

To quantify the correlation between wind direction and cholera spatial dissemination pattern for *Vibrio cholerae* dissemination in Africa during 1970–1971, we used the formula for circular correlation (ρ_{circ}) of Jammalamadaka and SenGupta (2001):

$$(\alpha_i, \beta_i), i = 1, 2, 3 \dots, n.$$

This formula denotes the sample pairs of directions (wind direction and the cholera

spatial dissemination direction). The formula for the general sample correlation is

$$\rho_{circ} = \frac{\sum_i \sin(\alpha_i - \bar{\alpha}) \sin(\beta_i - \bar{\beta})}{\sqrt{\sum_i \sin^2(\alpha_i - \bar{\alpha}) \sum_i \sin^2(\beta_i - \bar{\beta})}}, \quad [1]$$

where $\bar{\alpha}$ is the sample mean defined as

$$\bar{\alpha} = \begin{cases} \arctan(\bar{\alpha}_y / \bar{\alpha}_x) & \text{if } \bar{\alpha}_x > 0 \\ 180^\circ + \arctan(\bar{\alpha}_y / \bar{\alpha}_x) & \text{if } \bar{\alpha}_x < 0 \end{cases}$$

and

$$\bar{\alpha}_x = \sum_i \cos \alpha_i, \quad \bar{\alpha}_y = \sum_i \sin \alpha_i.$$

$\bar{\beta}$ is defined similarly. The ρ_{circ} ranges between -1 and 1 , similar to the linear correlation.

Results

Africa. Figure 1A–D shows examples of the results of mapping monthly average pressure values in the different altitudes over Africa during summer and autumn 1970. Note the impressive overlap between the dominant wind direction and the cholera progress in North Africa (August 1970), West Africa (September 1970), and East Africa (November 1970). Although we have not attempted to explain the disease spread throughout the entire area in 1970–1971 (e.g., the spreading from Mopti, the pilgrimage site in Senegal, could be a result of human dissemination), the influence of wind direction on the cholera dispersion adds a new explanation for disease dissemination in key parts of the continent.

Moreover, we obtained the sample circular correlation using 11 paired direction measurements from the *Vibrio cholerae* progress in Africa during 1970–1971 for the stages presented in Figure 1B–D. Our result

is $\rho_{circ} = 0.958$ ($p = 0.006$), indicating an impressive significantly high positive correlation. Figure 2 shows the relationship between the geographic dissemination of cholera over western and northeast Africa during March–September 2005 and February–March 2006 and the dominant wind direction over the continent in the same periods. Figure 2 shows a convincing correspondence (see selected examples in Figure 2A–C).

India. During March and April 1993, the air pressure values in the eastern part of India were significantly higher (by t -test analysis) than the average values for 1951–2000 (Figures 3 and 4A–B), and therefore caused a strengthening of the eastern winds. Table 1 presents the significant deviation of the mean SLP values in March 1993 from the mean values of 1951–2000, as well as the t -test results ($p \leq 0.05$) for selected grid points. In addition, after averaging all standardized deviations per year, we found that March 1993 was the month with the second-highest positive deviations throughout the entire 50 years, with an average t -score of 1.52.

These extreme easterly winds could carry the cholera bacteria from the east coast to Pune in the west. Indeed, there is an overlap between the wind direction and the cholera progress from eastern India to western India in March–April 1993 (Figures 3 and 4A–B).

To observe the diseases spreading in northern India, we compared the progress of cholera in May, June, and July (Figure 3) with the wind patterns at the SLP (an example is presented in Figure 4C). Geographic dissemination of *Vibrio cholerae* O139 completely corresponds to the dominant monthly wind direction, from the east (Jamshedpur) to the northwest.

Results of mapping monthly average geopotential values at the three different altitudes (1,000 mb, 925 mb, and 850 mb) together with their anomalies show the same trends as




Figure 2. Progression of cholera over Africa in March–September 2005 and February–March 2006 [A; adapted from WHO (2005) and WHO (2006)] and the dominant wind direction (shown by SLP maps in millibars) over the continent in July 2005 (B) and February 2006 (C).

during spring 1993, causing a significant strengthening of the easterly winds and therefore encouraging the cholera distribution from east to west.

- During a long monsoon period in May–July 1993, cholera's progress up the river was speeded up by the wind, all the way from Calcutta to northwest India (Jodhpur, July 1993). This last phase coincided with the usual timing and location of cholera in the Ganga Valley, which indicates a correlation between wind direction and cholera dissemination. Overall, the first O139 epidemic (sometimes called the eighth pandemic) did not follow the “seasonal” pattern of cholera occurrence in India but occurred over 10 consecutive months.

Conclusions

In a recent study, it was suggested that aeroplanktonic adult flies may carry the cholera bacteria between bodies of water (Broza et al. 2005). The aim of the present study was to further support this claim by showing linkages between wind (the vehicle of aeroplankton) and cholera-spreading patterns. Indeed, the present data connects wind direction with cholera dissemination patterns. Moreover, there is a clear relationship between the intra-continental distributions of cholera epidemics and the dominant wind direction over land. Synoptic data can be an important indication for predicting the direction in which cholera is spreading, by daily monitoring and forecasting of wind direction and intensity.

Because cholera continues to be an important cause of morbidity and mortality in Africa, Asia, and Latin America, further investigations of aerial transfer may improve the crucial ability to predict its dissemination.

Interactions between climatologists, entomologists, and physicians are generally limited. They do not exchange information that can be readily used in vector management. By raising the awareness of the relationship between cholera outbreaks and synoptic conditions, disease spread could be halted before a full-blown epidemic develops. Today, the importance of such collaboration is increasing dramatically because human activity enhances the greenhouse effect, causing climatic changes (e.g., Gartell 2001; IPCC 2001). This trend has many global and local aspects, one of which is the impact on human health. Global

Table 1. Deviation of mean monthly SLPs in March 1993 from the mean monthly values of March 1951–2000 at selected grid-points.

Selected grid-points	SD for March 1951–2000	March 1993 deviation	<i>t</i>	<i>p</i> -Value
15°00'N/75°00'E	0.94	1.67	1.773	0.041
20°00'N/75°00'E	1.12	1.99	1.769	0.041
20°00'N/87°30'E	1.11	1.85	1.671	0.050
22°30'N/75°00'E	1.15	2.21	1.916	0.030
22°30'N/80°00'E	1.32	2.23	1.682	0.049
25°00'N/75°00'E	1.26	2.38	1.897	0.032
25°00'N/87°30'E	1.45	2.73	1.887	0.033
27°30'N/80°00'E	1.63	2.76	1.692	0.049

Values are based on *t*-test results ($p \leq 0.05$).




Figure 4. Maps showing (A) mean SLP (in millibars) for 1951–2000; (B) deviations in SLP for March 1993 calculated based on the 1951–2000 averages for March; (C) mean monthly SLP for June 1993; and (D) dominant wind direction over northern India during June 1993 at the 925 mb level. Values in (D) indicate altitude in meters. Note how the geographic dissemination of *Vibrio cholerae* O139 corresponds to the dominant wind direction.




Figure 5. Photographs of (A) sunset on Lake Victoria (Kenya) showing a huge swarm of nonbiting midges moving over Rusinga Island toward the northern shore of the lake (photographed by H. Nadel), and (B) adult chironomids adhered to the engine cover of a car passing beside a sewage pond before sunrise (photographed by M. Broza).

climate change may affect human health via pathways of varying complexity, scale, directness, and timing. Similarly, the geographic impact may vary depending on the physical and environmental conditions and on the vulnerability of the local human population (Marsh and Gross 2001; McMichael 2003). Although most research deals with the influence of temperature increase on vector-borne diseases, it is important to study the impact of wind (direction and intensity) on spread of disease because wind is directly influenced by changes in atmospheric circulation. A good tool for monitoring and predicting cholera outbreaks will challenge a similar analysis of other pandemic diseases.

REFERENCES

- Armitage P, Cranston PS, Pinder CV, eds. 1995. *The Chironomidae: The Biology and Ecology of Non-biting Midges*. London: Chapman and Hall.
- Benke AC. 1998. Production dynamics of riverine chironomids: extremely high biomass turnover rates of primary consumers. *Ecology* 79:899–910.
- Bowden J, Johnson CG. 1976. Migrating and other terrestrial insects at sea. In: *Marine Insects* (Cheng L, ed). Amsterdam: North Holland Publishing, 97–118.
- Broza M, Gancz H, Halpern M, Kashi Y. 2005. Adult non-biting midges: possible windborne carriers of *Vibrio cholerae*. *Environ Microbiol* 7:576–585.
- Broza M, Halpern M. 2001. Pathogen reservoirs: chironomid egg masses and *Vibrio cholerae*. *Nature* 412:40.
- Colwell RR. 1996. Global climate and infectious disease: the cholera paradigm. *Science* 274:2025–2031.
- Faruque SM, Sack DA, Sack RB, Colwell RR, Takeda Y, Nair GB. 2003. Emergence and evolution of *Vibrio cholerae* O139. *Proc Natl Acad Sci USA* 100:1304–1309.
- Glick PA. 1939. The distribution of insects, spiders and mites in the air. *USDA Tech Bull* 673:1–150.
- Gartel AC. 2001. Health impacts of global environmental change. In: *Geographies of Health: An Introduction*. Oxford, UK: Blackwell Publishing, 234–255.
- Huq A, Sack B, Nizam A, Longini IM, Nair B, Ali A, et al. 2005. Critical factors influencing the occurrence of *Vibrio cholerae* in the environment of Bangladesh. *Appl Environ Microbiol* 71(8):4645–4654.
- IPCC (Intergovernmental Panel on Climate Change). 1996. Africa: regional climate. In: *IPCC Special Report on the Regional Impacts of Climate Change: An Assessment of Vulnerability*. Available: <http://www.grida.no/climate/ipcc/regional/010.htm> [accessed 11 April 2006].
- IPCC (Intergovernmental Panel on Climate Change). 2001. *Climate Change 2001: The Scientific Basis*. Available: http://www.grida.no/climate/ipcc_tar/wg1/ [accessed 4 December 2005].
- Jammalamadaka SR, SenGupta A. 2001. A circular correlation measure. In: *Topics in Circular Statistics*. Singapore: World Scientific Press, 176–183.
- Lipp EK, Huq A, Colwell RR. 2002. Effects of global climate on infectious disease: the cholera model. *Clin Microbiol Rev* 15:757–770.
- Marsh MM, Gross JM Jr. 2001. *Environmental Geography: Science, Land Use and Earth Systems*. 2nd ed. New York: John Wiley and Sons.
- Martyn D. 1992. *Climates of the World, Developments in Atmospheric Science*, Vol. 18, Amsterdam: Elsevier Press.
- McMichael AJ. 2003. Global climate change and health: an old story writ large. In: *Climate Change and Human Health: Risks and Responses* (McMichael AJ, Campbell-Lendrum DH, Corvalán CF, Ebi KL, Githeko A, Scheraga JD, et al. eds). Geneva: World Health Organization, 1–17.
- Nair GB, Ramamurthy T, Bhattacharya SK, Mukhopadhyay AK, Garg S, Bhattacharya MK et al. 1994. Spread of *Vibrio cholerae* O139 Bengal in India. *J Infect Dis* 169:1029–1034.
- NOAA (National Oceanic and Atmospheric Administration) and Climate Diagnostic Center. 2004. *Indian Monsoon*. Available: http://www.cdc.noaa.gov/ncep_reanalysis/examples.html [accessed 12 July 2005].
- NOAA NCEP-NCAR (National Oceanic and Atmospheric Administration, National Centers for Environmental Prediction, National Center for Atmospheric Research). 2005. *NOAA NCEP-NCAR CDAS-1: Climate Data Assimilation System I; NCEP-NCAR Reanalysis Project*. Available: ingrid.ideo.columbia.edu/SOURCES/NOAA/NCEP-NCAR/CDAS-1 [accessed 10 July 2005].
- NOAA NCEP-NCAR (National Oceanic and Atmospheric Administration, National Centers for Environmental Prediction, National Center for Atmospheric Research). 2006. *Monthly Intrinsic from NOAA NCEP-NCAR CDAS-1: Climate Data Assimilation System I; NCEP-NCAR Reanalysis Project*. Available: <http://iridl.ideo.columbia.edu/SOURCES/.NOAA/.NCEP-NCAR/.CDAS-1/.MONTHLY/Intrinsic/> [accessed 9 January 2006].
- Ramamurthy T, Garg S, Sharma R, Bhattacharya SK, Nair GB, Shimada T, et al. 1993. Emergence of a novel strain of *Vibrio cholerae* with epidemic potential in southern and eastern India [Letter]. *Lancet* 341:703–704.
- Reynolds DR, Mukhopadhyay S, Riley JR, Das BK, Nath PS, Mandal SK. 1999. Seasonal variation in the windborne movement of insect pests over northeast India. *Int J Pest Manag* 45:195–205.
- Ritchie SA, Rochester W. 2001. Wind blown mosquitoes and introduction of Japanese encephalitis into Australia. *Emerg Infect Dis* 7:900–903.
- Sellers RF, Pedgley DE, Tucker MR. 1982. Rift Valley fever, Egypt 1977: disease spread by windborne insect vectors? *Vet Rec* 110:73–77.
- Sparks AN, Jackson RD, Carpenter JE, Muller RA. 1986. Insects captured in light traps in the Gulf of Mexico. *Ann Entomol Soc Am* 79:132–139.
- Swerdlow LD, Isaacson M. 1994. The epidemiology of cholera in Africa. In: *Vibrio cholerae and Cholera: Molecular to Global Perspectives* (Wachsmuth IK, Blake PA, Olsvik Ø, eds). Washington, DC: American Society for Microbiology, 297–308.
- Tissot-Dupont H, Amadei MA, Nezri M, Raoult D. 2004. Wind in November, Q fever in December. *Emerg Infect Dis* 10:1264–1269.
- Todar K. 2005. *Vibrio cholerae* and Asiatic Cholera. Available: <http://textbookofbacteriology.net/cholera.html> [accessed 12 January 2006].
- WHO (World Health Organization). 2000. *WHO Report on Global Surveillance of Epidemic-prone Infectious Diseases*. Available: <http://www.who.int/csr/resources/publications/surveillance/en/cholera.pdf> [accessed 22 March 2006].
- WHO (World Health Organization). 2005. *Cholera in West Africa*. Available: http://www.who.int/csr/don/2005_08_26/en/index.html [accessed 10 October 2005].
- WHO (World Health Organization). 2006. *Cholera in Southern Sudan – Update*. Available: http://www.who.int/csr/don/2006_03_06/en/index.html [accessed 20 March 2006].