

Table 23. Natural Gas Home Customer-Weighted Heating Degree Days

Month/Year/Type of Data	New England	Middle Atlantic	East North Central	West North Central	South Atlantic
	CT, ME, MA, NH, RI, VT	NJ, NY, PA	IL, IN, MI, OH, WI	IA, KS, MN, MO, ND, NE, SD	DE, FL, GA, MD, DC, NC, SC, VA, WV
November					
Normal ^a	703	665	758	841	443
2007.....	751	692	764	790	458
2008.....	741	696	787	807	525
% Diff (normal to 2008)	5.41	4.66	3.83	-4.04	18.51
% Diff (2007 to 2008)	-1.33	0.58	3.01	2.15	14.63
November to November					
Normal ^a	703	665	758	841	443
2007.....	751	692	764	790	458
2008.....	741	696	787	807	525
% Diff (normal to 2008)	5.41	4.66	3.83	-4.04	18.51
% Diff (2007 to 2008)	-1.33	0.58	3.01	2.15	14.63

Month/Year/Type of Data	East South Central	West South Central	Mountain	Pacific ^b	U.S. Average ^b
	AL, KY, MS, TN	AR, LA, OK, TX	AZ, CO, ID, MT, NV, NM, UT, WY	CA, OR, WA	
November					
Normal ^a	454	305	739	366	589
2007.....	456	264	610	287	566
2008.....	541	287	599	242	578
% Diff (normal to 2008)	19.16	-5.90	-18.94	-33.88	-1.87
% Diff (2007 to 2008)	18.64	8.71	-1.80	-15.68	2.12
November to November					
Normal ^a	454	305	739	366	589
2007.....	456	264	610	287	566
2008.....	541	287	599	242	578
% Diff (normal to 2008)	19.16	-5.90	-18.94	-33.88	-1.87
% Diff (2007 to 2008)	18.64	8.71	-1.80	-15.68	2.12

^a Normal is based on calculations of data from 1971 through 2000.
^b Excludes Alaska and Hawaii.

Note: See Appendix A, Explanatory Note 10, for discussion of Heating Degree-Days computations.
Source: National Oceanic and Atmospheric Administration.