



OCRWM Office of Science and Technology and International



# THE PEÑA BLANCA NATURAL ANALOGUE MODEL

Presented to:  
**International High Level  
Radioactive Waste Management  
Conference**

Presented by:  
**George Saulnier  
AREVA NP**

**May 2, 2006  
Las Vegas, NV**


# ACKNOWLEDGEMENTS

- **William “Buddy” Statham was responsible for coding and model development**
- **The Bechtel-SAIC TSPA Department, especially Jerry McNeish and Kearn Patrick Lee**
- **Support from DOE Office of Science and Technology International**
- **The AREVA NP Federal Group, Tom Stevens and Joe Stringer**


# WHY A YUCCA MOUNTAIN ANALOGUE?

- **Analogous site: UO<sub>2</sub> uranium ore deposit = spent nuclear fuel in the repository**
- **Analogous geology: (i.e., Fractured, welded, and altered rhyolitic ash-flow tuffs)**
- **Analogous climate: Semiarid to arid**
- **Analogous setting: Volcanic tuffs overlie carbonate rocks**
- **Analogous geochemistry: Oxidizing conditions**
- **Analogous hydrogeology: Ore deposit lies in the unsaturated zone well above water table**

# Location of the Peña Blanca Mining District




# LOCATION OF THE NOPAL I URANIUM MINE





# LOCATION OF OBSERVATION WELLS



# LOOKING NORTHWEST AT NOPAL I URANIUM MINE





# VIEW SOUTHEAST FROM ABOVE +10 M LEVEL, NOPAL I





# OBJECTIVES

- **Develop a Total System Performance Assessment (TSPA) Model for the Peña Blanca Nopal I site**
- **Calibrate Peña Blanca Natural Analogue Model and compare predicted concentrations of uranium to observations at Nopal I, Peña Blanca**
- **Assist in building confidence in the modeling of total system performance of the Yucca Mountain repository**

# PROCESS



- **Develop a TSPA Model analogous to performance assessment models developed for the Yucca Mountain repository site**
- **Utilize probabilistic simulations to estimate the outcomes of hydrogeologic processes operating at the Nopal I site**
- **Perform model simulations using available data and field observations from ongoing investigations at Nopal I**

# HYDROGEOLOGIC SETTING

- **Fractured, welded rhyolitic ash-flow tuffs**
- **Arid climate**
- **Volcanic tuffs overlie carbonate rocks**
- **Nopal I ore deposit exposed to oxidizing conditions for 3.2 to 3.4 million years ago**
- **The Nopal I ore deposit lies in the unsaturated zone above the water table**



# HYDROGEOLOGIC SETTING



abq0063G337c.ai

# CONCEPTUAL MODEL

- **Nopal I ore deposit exposed to oxidation for ~3 million years**
- **Nopal I ore deposit extends from land surface to a depth of ~120 m, ~130 m above saturated zone**
- **Percolation of recharge through Nopal I ore deposit is the primary means of release of radionuclides**


# CONCEPTUAL MODEL (cont'd)

- **Simulate dissolution and release of uranium using the Yucca Mountain TSPA spent-fuel dissolution model**
- **Generate and release of three Uranium species, Technetium 99, and Thorium**
- **Assume the Nopal I source term consists of two cells analogous to two waste packages containing uranium oxide**
- **Use a defined steady-state dissolution rate**



# CONCEPTUAL MODEL (cont'd)

## CONCEPTUAL MODEL



# MODEL PARAMETERS & ASSUMPTIONS

- **Nopal I ore deposit consists of 408 Metric Tons Uraninite, an analogue to commercial spent nuclear fuel**
- **Ore deposit is idealized as a vertically oriented ovoid cylinder 18 x 30 x 100 meters**
- **Peña Blanca Natural Analogue Model for Nopal I assumes Percolation = infiltration at 6 mm/yr or 3.24 m<sup>3</sup>/yr**

# MODEL PARAMETERS & ASSUMPTIONS

- **Assume Groundwater flows generally West to East**
- **Hydraulic Parameters of ore deposit**
- **Porosity 0.075 (unaltered) to 0.30 (altered)**
- **$K_d$  Tc-99 = 0;**
- **$K_d$  U, varied between 0 and 1;**



# RADIONUCLIDE INVENTORY


- **Estimates based on 99.5% Uranium 238**
- **Extrapolations from vegetation study (Leslie et al., 1999)**
- **Technetium from Curtis et al., (1999)**
  - **Tc99 = 0.025 atoms/min/gU-238**
  - **Tc99/U238 = 1.57 E -12**
- **Radioactive decay assumed for radionuclides**
- **Radium based on site investigations**

# PEÑA BLANCA NATURAL ANALOGUE MODEL RESULTS

- **Base case simulations estimated dissolved concentrations of Tc-99 and Uranium in the saturated zone beneath the Nopal I ore deposit**
- **Results sensitive to Uranium solubility, infiltration rate, surface area available for dissolution, and  $K_d$**
- **Observed concentrations bracketed by the range of Peña Blanca Natural Analogue Model Results, within the uncertainty of the source-term dissolution parameters**


# PEÑA BLANCA NATURAL ANALOGUE MODEL RESULTS

Calculated radionuclide concentrations directly beneath the Nopal I ore deposit



# PEÑA BLANCA NATURAL ANALOGUE MODEL RESULTS


Calculated radionuclide concentrations beneath Nopal I using high Kd for all Uranium species



# NOPAL I URANIUM CONCENTRATIONS

Uranium concentrations in water samples from Nopal I observation wells

U Time Series for Wells




00318DC\_405.ai



# PENA BLANCA NATURAL ANALOGUE MODEL RESULTS

Results of the base case simulation for  $^{238}\text{U}$  at 65 m for 100 realizations of the uncertain dissolution parameters for late time samples from PB1, PB2, and PB3



# CONCLUSIONS

- **The Nopal I ore deposit is a useful analogue**
  - Nopal I ore deposit is smaller than YMP
  - Nopal I ore deposit has similar geology
- **Tc-99 may be detectable in picograms/L at 100 m**
- **Thorium 230, 232 predicted**
- **Uranium 238, 234 predicted but possibly limited due to mineral precipitation**

# Future Plans

- **Modify Peña Blanca Natural Analogue Model to reflect ongoing data collection and revised versions of Yucca Mountain process models**
- **Analyze water samples for Tc-99**
- **Vary Thorium  $K_d$**
- **Add Radium to Model Inventory**
- **Revisit Model Inventory after completion of UTEP investigations**