

Rapid Assessment Reference Condition Model

The Rapid Assessment is a component of the LANDFIRE project. Reference condition models for the Rapid Assessment were created through a series of expert workshops and a peer-review process in 2004 and 2005. For more information, please visit www.landfire.gov. Please direct questions to helpdesk@landfire.gov.

Potential Natural Vegetation Group (PNVG)

R2CRBU Creosotebush Shrublands with Grasses

General Information

Contributors (additional contributors may be listed under "Model Evolution and Comments")

Modelers

Sandy Gregory s50grego@nv.blm.gov

Reviewers

Patti Novak-Echenique patti.novak@nv.usda.gov
Tim Duck tim_duck@blm.gov
Stanley D. Smith ssmith@ccmail.nevada.edu

Vegetation Type

Shrubland

General Model Sources

- Literature
 Local Data
 Expert Estimate

Rapid Assessment Model Zones

- California Pacific Northwest
 Great Basin South Central
 Great Lakes Southeast
 Northeast S. Appalachians
 Northern Plains Southwest
 N-Cent. Rockies

Dominant Species*

AMDU
LATR2
EPNE
ATRIP

LANDFIRE Mapping Zones

12	17
13	18
16	

Geographic Range

This area extends from southern California, western Arizona, southern Nevada, and southwestern Utah.

Biophysical Site Description

Creosotebush Scrub is wide ranging across the warm deserts and is most common in the Mojave Desert. Creosotebush scrub is typically found below the blackbrush zone on well-drained alluvial flats and slopes and above the saltbush zone. Elevation zones 600 to 4200 ft. Creosote shrublands occur on several soil types from shallow to deep. Desert pavement is common. Soils are well drained, available water capacity is very low and runoff is moderate to rapid.

Vegetation Description

Creosotebush (*Larrea tridentata*) dominates shrublands. Plant community associates change from east to west Mojave Desert. Creosotebush codominates in desert shrub communities with saltbush (*Atriplex* spp.), *Ambrosia dumosa*, white bursage, and bladder sage (*Salazaria mexicana*). Joshua tree forests are included in this community type. Grass species include galleta grasses (*Hilaria* spp.). Creosotebush scrub is characterized by low cover (5-30%) of woody shrubs of various heights. With the exception of Joshua tree, creosotebush has the highest cover and is the most wide-ranging plant species in the Mojave Desert.

Disturbance Description

Not fire tolerant, because of their drought-tolerant features (thin bark, slow growth, shallow root system, small leaves). Some species resprout after fire depending on fire severity, but native plants are slow to recover or re-establish after fire.

We do not know the pre-settlement fire conditions in warm desert plant communities. However, it is thought that fires in creosotebush scrub were an infrequent events in pre-settlement desert habitats, because fine fuels from winter annual plants were probably sparse, only occurring in large amounts during the spring

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

following exceptionally wet winters.

It appears that wildfire was not historically a landscape dominating influence in creosotebush scrub habitat.

Adjacency or Identification Concerns

The issue is not as much of a concern as in other systems.

Fine fuels adjacency from alien annual grasses currently represent the most important fuelbed component in creosotebush scrub. Alien annual grasses can comprise 66-97% of the total annual biomass in this system.

Livestock grazing has contributed to the deterioration of this system.

Scale Description

Sources of Scale Data	<input checked="" type="checkbox"/> Literature	<input checked="" type="checkbox"/> Local Data	<input checked="" type="checkbox"/> Expert Estimate
------------------------------	--	--	---

Patch sizes vary to slope and aspect and precipitation.

Issues/Problems

There little information regarding fire frequency, fire severity in pre-settlement fire conditions in warm desert plan communities is not well documented. It is thought that fire was infrequent and not landscape dominating.

Model Evolution and Comments

For the Rapid Assessment, this model was also used only in the western portion of the Southwest model zone where creosote bush occurs.

As ecological condition deteriorates, creosotebush, blackbrush, black gramma, Indian ricegrass, desert needlegrass decrease. Species likely to invade this site such as Bromus rubens and Schismus barbatus.

Succession Classes**
Succession classes are the equivalent of "Vegetation Fuel Classes" as defined in the Interagency FRCC Guidebook (www.frcc.gov).

Class A 10%

Early1 Open

Description

Creosotebush scrub is characterized by low cover 5 to 10%. Little disturbance was considered in Class A, except for replacement fire every 300 yrs on average. Historical condition where invasive annual grasses are absent, the fire return interval is virtually non-existent except for areas near the base of mountains experiencing locally higher rainfall and fine fuel buildup from native annual. After 100 yrs, class A transitions to B.

Dominant Species* and Canopy Position

PPGG

AMUD

LATR2

ARIST

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	5 %	10 %
Height	no data	no data
Tree Size Class	no data	

Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Class B 90 %

Late I Closed

Description

Greater than 15% shrub cover and 20-40 percent grass and forb cover; associated with more productive soils. Less fine fuels are associated with this community, therefore the FRIs for replacement fire and mixed severity fire is 650 years (min-max: 300-1000 yrs). Wind/weather stress also affected this community on average every 80 yrs, but did not cause a transition to class A.

Dominant Species* and Canopy Position

LATR2
AMUD

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

Structure Data (for upper layer lifeform)

	Min	Max
Cover	15 %	20 %
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class C 0 %

Late I Open

Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	%	%
Height	no data	no data
Tree Size Class	no data	

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class D 0 %

Late I Open

Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	0 %
Height	no data	no data
Tree Size Class	no data	

Upper Layer Lifeform

- Herbaceous
- Shrub
- Tree

Fuel Model no data

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

Class E 0 %

Late I Closed

Description

Dominant Species* and Canopy Position

Structure Data (for upper layer lifeform)

	Min	Max
Cover	0 %	%
Height	no data	no data
Tree Size Class	no data	

- Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.

Upper Layer Lifeform Upper layer lifeform differs from dominant lifeform. Height and cover of dominant lifeform are:
 Herbaceous
 Shrub
 Tree

Fuel Model no data

Disturbances

Disturbances Modeled

- Fire
- Insects/Disease
- Wind/Weather/Stress
- Native Grazing
- Competition
- Other:
- Other

Historical Fire Size (acres)

Avg: no data
 Min: no data
 Max: no data

Fire Regime Group: 5

- I: 0-35 year frequency, low and mixed severity
- II: 0-35 year frequency, replacement severity
- III: 35-200 year frequency, low and mixed severity
- IV: 35-200 year frequency, replacement severity
- V: 200+ year frequency, replacement severity

Fire Intervals (FI)

Fire interval is expressed in years for each fire severity class and for all types of fire combined (All Fires). Average FI is central tendency modeled. Minimum and maximum show the relative range of fire intervals, if known. Probability is the inverse of fire interval in years and is used in reference condition modeling. Percent of all fires is the percent of all fires in that severity class. All values are estimates and not precise.

Sources of Fire Regime Data

- Literature
- Local Data
- Expert Estimate

	Avg FI	Min FI	Max FI	Probability	Percent of All Fires
<i>Replacement</i>	588	300	1000	0.00170	56
<i>Mixed</i>	769	300	1000	0.00130	43
<i>Surface</i>					
<i>All Fires</i>	333			0.00301	

References

Brooks, M.L., T.C. Esque, and T. Duck. 2003. Fuels and fire regimes in creosotebush, blackbrush, and interior chaparral shrublands. Report for the Southern Utah Demonstration Fuels Project, USDA, Forest Service, Rocky Mountain Research Station, Fire Science Lab, Missoula, Montana. 17pp.

Brown, D. E. and R. A. Minnich. 1986. Fire and creosote bush scrub of the western Sonoran Desert, California. *American Midland Naturalist* 116:411-422.

Brown, J. K., and J. K. Smith, eds. 2000. Willdand fire in ecosystems: effects of fire on flora. Gen. Tech. Rep RMRS-GTR-42-vol.2. Odgen, UT; US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Kuchler, A.W. 1964. Manual to accompany the map of potential natural vegetation of the conterminous United States. American Geographical Society. Spec. Publ. NO. 36. Lib. Congress Cat. Card Num. 64-15417

Marshall, K. Anna. 1995. *Larrea tridentata*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2004, November 14].

United States Department of Agriculture Natural Resources Conservation Service, Nevada Rangeland Ecological Site Description. 0300XB078NV, 030XB019NV

*Dominant Species are from the NRCS PLANTS database. To check a species code, please visit <http://plants.usda.gov>.